SapL1: A New Target Lectin for the Development of Antiadhesive Therapy Against *Scedosporium apiospermum*

Dania Martínez-Alarcón[1, 2], Jean-Philippe Bouchara[3], Roland J. Pieters[2,] Annabelle Varrot[1*].

1   Univ. Grenoble Alpes, CNRS, CERMAV, 38000 Grenoble, France. dania.martinez.alarcon@gmail.com, annabelle.varrot@cermav.cnrs.fr

2   Utrecht University, 3584 CG Utrecht, The Netherlands. R.J.Pieters@uu.nl

3   Host-Pathogen Interaction Study Group, GEIHP, EA 3142, SFR ICAT 4208, UNIV Angers, UNIV Brest, Institut de Biologie en Santé, IRIS, CHU d'Angers, Angers, France

* Corresponding author

Email: annabelle.varrot@cermav.cnrs.fr

Abstract

Scedosporium apiospermum is an emerging opportunistic fungal pathogen responsible for life-threatening infections in immunocompromised patients. This fungus exhibits limited susceptibility to all current antifungals and, due its emerging character, its pathogeny and virulence factors remain largely unknown. Carbohydrate binding proteins such as lectins are involved in host-pathogen interactions and may constitute valuable therapeutic targets to inhibit microbial adhesion to the host cells by using carbohydrate mimics. However, such lectins are still unidentified in *S. apiospermum*. Here, we present the first report of the identification and characterization of a lectin from *S. apiospermum* named SapL1. SapL1 is homologous to the conidial surface lectin FleA from *Aspergillus fumigatus* known to be involved in the adhesion to host glycoconjugates present in human lung epithelium. The present report includes a detailed strategy to achieve SapL1 soluble expression in bacteria, its biochemical characterization, an analysis of its specificity and affinity by glycan arrays and isothermal titration calorimetry (ITC), as well as the structural characterization of its binding mode by X–ray crystallography. The information gathered here contribute to the understanding of glycosylated surface recognition by *Scedosporium* species and is essential for the design and development of antiadhesive glycodrugs targeting SapL1.

Keywords: fucolectin, glycodrug, host recognition, glycans, opportunistic pathogen.

Introduction

During the last decades, an increased incidence of invasive infections, especially in immunosuppressed patients, has been caused by previously rare fungal pathogens.[1] Among those pathogenic fungi, species from the *Scedosporium* genus are particularly dangerous since their mortality rate may be over 80%.[2] *Scedosporium* infections mainly affect immunocompromised patients such as solid organ transplant (SOT) or hematopoietic stem cell transplant (HSCT) recipients, or patients with chronic granulomatous disease. Together with *Fusarium* species and some dematiaceous fungi, *Scedosporium* species are responsible for approximately 10% of the mycosis caused by filamentous fungi in hematopoietic cell transplant recipients and up to 19% in solid organ transplant patients.[3] In addition, *Scedosporium* species rank the second among the filamentous fungi colonizing the respiratory tract of people with cystic fibrosis (CF), after *Aspergillus fumigatus*, and because of the propensity of these fungi to disseminate in case of immune deficiency, this fungal colonization of the airways is considered in some centers as a contraindication to lung transplantation which remains the ultimate treatment in CF.[4] Apart from lung transplantation, this fungal colonization of the airways which usually starts in the adolescence and becomes chronic in up to 19% of the patients, may also lead to a chronic inflammation and sometimes to an allergic bronchopulmonary mycosis.[2] Furthermore, it has been reported that some species of this genus can also affect immunocompetent patients.[5]

The genus *Scedosporium* comprises more than ten species *(Scedosporium angustum, S. apiospermum, S. aurantiacum, S. boydii, S. cereisporum, S. dehoogii, S. desertorum, S. ellipsoideum, S. fusoideum, S. minutisporum, S. rarisporum and S. sanyaense)* whose taxonomy has been previously described based on molecular phylogenetic by using the sequences of four genetic loci.[2,6] However only the genome of *S. boydii, S. aurantiacum, S. dehoogii* and *S. apiospermum* have been completely sequenced to date.[7-9] Among *Scedosporium* species, *S. apiospermum* (formerly considered as the asexual form of *Pseudallescheria boydii*) is the most outstanding. This cosmopolitan microfungus is responsible for localized to severe or fatal disseminated infections in humans and is one of the most common *Scedosporium* species (with *S. boydii* and *S. aurantiacum*) capable to colonize chronically the lungs of CF patients.[3,10]

The treatment of *Scedosporium* infections is challenging because its efficacy depends on timely diagnosis, which is based on morphological detection by microscopy, histological analysis, and culture on selective media. This process is time consuming and may lead to false negative results, especially from respiratory secretions of patients with CF because of the common co-colonization by other fungi or bacteria. [2] Furthermore, *Scedosporium* species display a primary resistance to classical antifungals such as 5-flucytosine, amphotericin B, and the first-generation triazole drugs, fluconazole or itraconazole and exhibit a limited susceptibility to the newest generations of antifungal drugs, *i.e.* echinocandins, voriconazole and isavuconazole.[2,11] Nowadays, the first-line treatment for *Scedosporium* infections involves a combination therapy that includes the use of voriconazole in conjunction with other antifungals.[11,12] However, due to common recurrences, even without interruption of treatment, the recovery rates are poor and mortality remains over 65% while it is almost 100% when dissemination occurs, reason why it has aroused special interest despite its emerging character.[12,13]

Scedosporium infections begin with conidial adherence to tissues, followed by germination and hyphal elongation.[14] This adherence process allows it to avoid cleansing mechanisms aimed to eradicate the invading pathogens and represents the initial step towards infection.[14-17] Conidial adhesion is mediated by cell surface molecules (CSM), including different types of carbohydrates such as polysaccharides and glycoconjugates. Some of the most important carbohydrates CSM described to date in *Scedosporium* species include peptidorhamnomannans (PRMs),[18-20] α-glucan,[21] melanin,[22] ceramide monohexosides,[20,23] N-acetyl-D-glucosamine-containing molecules[24] and mannose/glucose-rich glycoconjugates.[20] The presence and/or abundance of these molecules on the cell surface varies according to the stage of development and is of great relevance to understand fungal pathobiology.[20] The carbohydrates binding proteins known as lectins also act as CSMs and were shown to have an essential role during pathogenesis. However, the knowledge about lectins in emerging microfungi is very limited to date.

Since some lectins from microorganisms have been demonstrated to be involved in the host recognition and adhesion process, they have been used to develop anti-adhesive therapy through the use of carbohydrates and their mimics.[15-17,25-31] This approach is particularly promising since it does not kill the pathogen nor arrest its cell cycle. Consequently, resistance frequencies are rare [16,26] and expected side effects

in non-target tissues are lower than those caused by conventional antifungal compounds. This may allow its implementation as a prophylactic therapy for immunocompromised patients.[17,27]

Due to the emerging character of *Scedosporium apiospermum*, there is very limited information on its mechanisms of recognition and anchoring to the host. Furthermore, lectins from this microorganism have not been characterized to date, what hinders the development of an anti-adhesive therapy. Conversely, closely related filamentous fungi were investigated leading to the identification and characterization of their host binding modes. For example, in *A. fumigatus,* which is a saprophytic mold also responsible for bronchopulmonary infections in receptive hosts, the lectin FleA (or AFL) was identified and revealed a role in host-pathogen interactions.[32] FleA is a six-bladed β-propeller homodimer located on the conidial surface that recognize human blood group antigens and mediates *A. fumigatus* binding to airway mucins and macrophages glycoproteins in a fucose-dependent manner.[33] In healthy individuals, this anchorage is critical for the mucociliary clearance process and the macrophagy, in fact it has been described that fleA-deficient (ΔfleA) conidia are even more pathogenic than wild type (WT) conidia both, in healthy and chemically immunocompromised mice.[33,34] However, CF patients represent a very particular scenario because the mucus in their lungs is thicker, in relation with mucin overproduction and the high content of calcium ions, which modulates the supramolecular organization of MUC5B by protein cross-linkages.[34,35] This contributes to the suboptimal transport properties of mucus and compromises the pathogen clearance mechanisms [34]. Furthermore, the aberrant glycosylation of the mucins MUC5B and MUC5B in CF lungs causes, among other things, an increase in the abundance of sialyl-Lewis X and Lewis X determinants in the non-reducing end of mucins carbohydrate chains, [36,37]which can also be translated as an increase in the fucose content. Therefore, in this context, the FleA (and homologous proteins) anchoring to the mucus layer plays an essential role in the colonization of the CF lungs by *A. fumigatus*.

Here, we have used the recently sequenced genome of *S. apiospermum*,[9] to identify a putative homologue of FleA that we have called SapL1 for *Scedosporium apiospermum* Lectin 1. The present report comprises SapL1 identification, its production in bacteria, an analysis of the fine specificity and affinity of the recombinant protein, as well as its structural characterization by X–ray crystallography.

Results

Production and purification of SapL1

The hypothetical protein XP_016640003.1 (EMBL accession number), encoded by SAPIO_CDS9261 and from now on referred as SapL1, was identified through data mining using the FleA (pdb entry 4D4U[38]) sequence as bait into the genome of the reference strain *S. apiospermum* IHEM 14462. [9] During the sequence analysis, we identified that the first 74 amino acids of the putative protein (Uniprot A0A084FYP2) are a very disordered region that is not present in any other protein of the same family. We suspected that this peptide could derived from a misplacement of the open reading frame (ORF) of SapL1 during the genome assembly and that its sequence actually starts at methionine-75 (Figure S1). The SapL1 coding sequence (from Met-75) was fused to an N-terminal 6xHis tag cleavable by the Tobacco Etch virus protease (TEV) under regulation of *trc* and *T7* promoters into pProNde and pET-TEV vectors, respectively (Figure 1A). Expression was performed in *Escherichia coli* and purification was carried out using immobilized metal affinity chromatography (IMAC). Unfortunately, the original expression yield with both vectors (~0.35 mg·L^{-1} of culture) was too low to proceed with characterization studies. Therefore, we explored new alternatives to enhance the expression. First, the thioredoxin protein (Trx) as well as 6-His tag and TEV cleavage site were fused at the N-terminus of SapL1 by subcloning into the pET32-TEV vector (Figure 1A). This strategy substantially increased the SapL1 production yield but most of the protein remained insoluble as part of inclusion bodies (Figure 1B). A wide range of expression conditions were subsequently assayed to achieve sufficient soluble expression by modification of various parameters such as growth temperature, host strain, inducer concentration, optical density of the culture at induction, culture duration, etc. Sixty-nine different sets of parameters were assayed (S1 Table) and it was possible to improve the yield up to 4 mg·L^{-1} (Figure 1C). The best set of conditions for SapL1 expression was using *E. coli* strain TRX, pProNde vector, LB media, growth at 37°C and 160 rpm until OD$_{600}$ = 0.4, before a switch of the temperature to 16°C and overnight induction at OD 0.8 with 0.05 mM IPTG and 1% L-rhamnose (Rh).

Rhamnose influence on SapL1 solubility

Interestingly, during the optimization of the expression conditions, we found that
rhamnose (Rh) plays an essential role in the solubility and stability of SapL1.
Therefore, we performed a new set of experiments to demonstrate the correlation
between the amount of protein recovered after purification and the Rh concentration
in the media. For this, we investigated SapL1 expression under the previously
described conditions modified by supplementation of the culture medium with
different concentrations of Rh (0.1%, 0.15%, 0.2%, 0.5% and 1%). Purification
parameters were set to obtain high purity product and kept identical for all
experiments. Figure 2 shows the chromatograms obtained for representative
concentrations accompanied by their respective SDS-PAGE profile. To quantify
SapL1 expression in these experiments, we integrated the area under peaks
corresponding to the protein (Figure 2, black bars). Thus we confirmed that SapL1
recovery is directly proportional to the Rh concentration in the culture medium.
Then, the experiment was repeated at three different concentrations (0.1%, 0.15%,
0.2%) using 20°C as the induction temperature, and production behavior was similar
to that found at 16°C, but with lower performance.

Additional experiments also showed that SapL1 expression can be induced at high
concentration of rhamnose in a dose dependent manner, even without addition of
IPTG when the pProNde vector is used (data not shown). For this reason, we initially
consider that the higher production rates of SapL1 could be attributed to the
dissociation of the *LacI*-DNA complex caused by an increase in cytosolic potassium
concentration since the accumulation of this ion is one of the main adaptive
responses of *E. coli* to hypertonic shocks, and its concentration has been correlated
to the loss of interactions between DNA and proteins in *in vitro* assays.[39-41] However,
previous reports have demonstrated that, *in vivo*, the effects of macromolecular
crowding caused by plasmolysis (the reduction of intracellular water content, as an
adaptive response to osmotic shock) on the activity of cytoplasmic proteins, is
sufficient to buffer the kinetics of association of DNA-protein against changes in
$[K^+]$.[41,42] Thus, it seems that the higher recovery rates of SapL1 is actually caused by
a combination of two main parameters: 1) the leak of genetic repression by
dissociation of the *LacI-DNA* complex (characteristic of *lac* promoters) and 2) a
positive effect on the solubility and folding of the protein due to the increased
concentration of organic osmoregulators, whose synthesis is induced in *E. coli* as an
adaptive response to the external hypertonicity caused, in this case, by the addition

of rhamnose to the culture medium[39-47] This hypothesis was later supported by the

finding that the addition of glycerol, instead of rhamnose, displays the same effect

on production and solubility of SapL1 (data not shown). Plasmolysis may also play

an essential role in SapL1 stabilization, since it has been shown that macromolecular

crowding has positive effects on protein folding and in some cases it can drive self-

association of improper folded proteins into functional oligomers.[41]

Biochemical characterization

We assayed the thermal stability of SapL1 in 26 different buffers in a pH range of 5

to 10 through Thermal Shift Assay (TSA). The most suitable condition for this

protein was MES buffer 100 mM pH 6.5, where a single denaturing event at T_m of

55°C was observed (Figure S2). Then, to estimate the molecular size of the native

protein, we performed size exclusion chromatography using an ENrich™ SEC 70

column (Bio-Rad) and 20 mM MES, 100 mM NaCl, pH 6.5 as mobile phase.

However, the protein displayed strong non-specific interactions with the matrix of

the column, and SapL1 could not be eluted even using 5 M of NaCl. Interestingly, it

could be recovered when the buffer was supplemented with 20 mM α-methyl

fucoside or L-rhamnose, evidencing similar effects on SapL1 elution for both sugars.

Due to the impossibility to estimate the molecular weight of SapL1 by size exclusion

chromatography on this resin, we performed measurements in solution using

Dynamic Light Scattering (DLS). We obtained a monodisperse peak corresponding

to a protein of 72 ± 29.4 kDa corroborating that SapL1 forms dimers as FleA and

other proteins of this family (monomer MW: 40 kDa, data not shown).[32] The range

of the standard deviation also suggested an ellipsoidal shape, which is characteristic

for the dimers in this lectin family[48].

Carbohydrate binding properties

A hemagglutination assay showed that recombinant SapL1 agglutinated rabbit red

blood cells at 0.97 µg·ml[-1] (Figure 3A). It confirms that the recombinant lectin is

active and that its heterologous production in *E. coli* did not alter its

hemagglutinating properties.

To identify the potential ligands of SapL1 on epithelial cell surfaces, we submitted

SapL1 to the glycan array version 5.4 of the Consortium for Functional Glycomics

(USA) consisting of 585 mammalian glycans. It was labelled with Fluorescein

Isothiocyanate (FITC) in a molar ratio of 0.426 and its binding properties were

analysed at two different concentrations (5 and 50 µg·mL^{-1}). As expected from its
homology with FleA, SapL1 recognizes fucosylated oligosaccharides independently
of the fucose linkage. The α1-2 and α1-3/4 linked fucosides displayed the highest
affinity whilst α1-6 linked ones, the lowest. The weakest interactions with
fucosylated compounds were reported for branched oligosaccharides (Figure S3).
The affinity of SapL1 for L-fucose and α-methyl-fucoside was determined by
Isothermal Titration Calorimetry (ITC) and the K_d was found to be 225 µM and 188
µM, respectively with stoichiometry fixed to 1 since the measurements were done
in the presence of an excess of ligand (Figure 3B). These values are in agreement
with the affinity constant (around 110 µM) reported for FleA for α-methyl-
fucoside.[49] No binding interaction was observed for SapL1 with rhamnose by ITC
(data not shown).

Analysis of the glycans constituting blood group determinants revealed that SapL1
binds to all epitopes with a preference for H type 2 (Fucα1-2Galβ1-4GlcNAcβ) then
Lewis^a (Galβ1-3(Fucα1-4)GlcNAcβ) and Lewis^x (Galβ1-4(Fucα1-3)GlcNAcβ).
However, most of the recognized branched oligosaccharides contained the core
fucose Fucα1-6. Epitopes with two fucose units, such as Lewis^b and fucosylated
polylactosamine, were also well recognized. Addition of a galactose or a GalNAc as
in blood group B or A antigens did not impair Fucα1-2 recognition (Figure 3C).
Spacers used to join the carbohydrates to the chip also display a strong influence on
binding. It is remarkable that 70% of the 90 positive binders contained either the
spacer Sp0 (CH$_2$CH$_2$NH$_2$) or Sp8 (CH$_2$CH$_2$CH$_2$NH$_2$). Those spacers also display a
strong influence on binding, especially for small glycans such as Galα1-3(Fucα1-
2)Galβ1-4(Fucα1-3)GlcNAcβ which was recognized when attached to Sp0 but not to
Sp8. This may be due to a steric hindrance caused by the modification of
carbohydrate presentation on the surface of the chip.

Overall structure of SapL1
SapL1 was co-crystallized with α-methyl-fucoside and the structure of the complex
was solved by molecular replacement at 2.3 Å resolution in the P2$_1$ space group using
the coordinates of FleA (PDB code 4D4U[38]) as the search model. The asymmetric unit
contained two monomers, assembled as a dimer with all 295 amino acids visible
apart of the N-terminal methionine. SapL1 folds into the canonical six-bladed β-
propeller with six-binding sites at the interface between blades typical for this family
of lectin (Figure 4). A fucose moiety was found in 5 and 4 of the six binding pockets

of chains A and B, respectively while glycerol originating from the crystallizing
solution was found in the other binding sites.

The overall fold of SapL1 and FleA as well as the overall dimer are very similar with
a rsmd of 1.2 and 1.26 Å, respectively. They share 43% of sequence identity and both
proteins present the same distribution of β-strands except for the lack of the last
blade of strand 4 in SapL1, and the external face of blade 5, in which FleA displays
an elongated β-strand (β22) that is split in two in SapL1 (β22/23). It is remarkable
that the structure of the first 3 blades is highly conserved in both proteins, while the
second half (blades 3-6) displays the largest discrepancies. Furthermore, FleA also
presents two additional small α-helices (ή3 and ή4), located in the loops between
sheets that serve as connectors for blades (2/3 and 5/6, respectively, Figure 5). Similar
conclusions can be drawn when comparing SapL1 to its homologue AOL in
Aspergillus oryzae.[50] Those proteins have similar dimerization interface and we
noticed differences on the other side of the propeller where sugar binding occurs.
The lectins present surface loops of different size and sequence leading to change in
the architecture of the related binding sites that impacts their affinity as discussed
in more details below. The first member structurally characterized in this lectin
family was AAL from the mushroom *Aleuria aurantia*.[51] It presents differences on
both sides of the β-propeller with the microfungal members of this family mainly at
the level of the surface loops. This leads to a different dimerization interface and the
dimer cannot overlay. This also affects fucose binding since AAL only has five
functional binding sites. The role of these lectins is still unknown, in particular for
the mushroom members.
289

Protein-ligand interactions
Due to divergence in the tandem repeat sequence forming each blade of the
propeller, the six binding sites of SapL1 monomer are not equivalent, but they share
important conserved features. Hydrophobic interactions are observed between the
C6 of the fucose and at least three residues of the protein (mainly isoleucine,
tryptophan/tyrosine and leucine). The O2 and O3 hydroxyls make strong hydrogen
bonds with a conserved triad of amino acid consisting of an arginine, a glutamic acid
and a tryptophan. In the cases where glycerol was found in the binding pocket, its
own hydroxyls mimic the interactions of fucose with these same residues (Figure 6).
It is to be noted that SapL1 binding sites are more conserved than FleA binding site

where a glutamine can replace the glutamic acid and a tyrosine the tryptophan in the triad (Figure S4).

The O2 hydroxyl seemed to be the most versatile position, since it established interactions with a loop adjacent in four of the six binding sites (BPs 2, 4, 5 and 6). This is particularly interesting since the binding pockets that do not contain this loop (BPs 1 and 3) were mostly occupied by glycerol instead of fucose, indicating that those interactions could be responsible for enhancing the affinity and might be explored for development of inhibitors.

Discussion

The development novel techniques for diagnosis has exposed that, until now, the incidence of fungal infections has been underestimated.[1,52] In comparison with pathogenic bacteria, which have been the focus of attention for several years, there is a lack of information about the virulence factors and host recognition systems of pathogenic fungi.[53] This hinders the development of new drugs against this microorganisms, while their resistance to the current ones is progressing quickly.[13,24,52] Therefore, there is a rush in obtaining information that may contribute to the development of new antifungal agents.

In this study, we present the identification, production and characterization of a new carbohydrate binding protein from the emerging microfungus *S. apiospermum*, whose host-anchoring mechanism is completely unknown. To the best of our knowledge, this is the first studied lectin for this opportunistic pathogen. SapL1 is homologous to the conidial surface lectin FleA from *A. fumigatus* known to be involved in adhesion to the host glycoconjugates present in mucins and human lung epithelium.[33] During our analysis, we have identified a possible shift in the ORF of SapL1 and we have relayed this information to corroborate this finding and to update the entry if necessary.

In our initial attempts to produce recombinant SapL1, we observe that the conventional parameters of expression in *E. coli*, result in the production of insoluble protein that was mainly found in inclusion bodies. Later, experimental data indicated that the presence of rhamnose or glycerol was required for SapL1 solubilization during bacterial expression and that the production yield was directly

proportional to their concentration in the culture medium. This suggests that the stabilization of SapL1 derives from an osmomodulator effect that, either directly or indirectly, enhances its proper folding. This osmomodulator effect may be due to the leak of genetic repression by dissociation of the repressor (*LacI-DNA)* complex[39-41] in the first case, or in the second case to the increased concentration of organic osmoregulators, which are known to enhance protein folding and solubility, presumably also reinforced by the macromolecular crowding produced by plasmolysis.[39-47] These findings, and the extensive research carried out to produce SapL1 in its soluble form, provide important insights for the heterologous expression of eukaryotic lectins tending to be produced as insoluble proteins in *E. coli.* Besides, our findings highlight the useful role of organic osmoregulators during heterologous expression of proteins to favour their proper folding and to avoid the formation of inclusion bodies.

We demonstrated that SapL1 is strictly specific for fucosylated carbohydrates and recognized all blood group types present in the glycan array screening. These results are particularly interesting since it has been shown that there is six times more content of fucose α1,3/4 linked to glycoproteins in the CF airways.[36] This phenomenon is mainly due to an increased expression of the α1,3-fucosyltransferase, which is involved in the synthesis of sialyl-Lewis X and Lewis X determinants attached to mucus mucins.[37] Therefore, the fact that SapL1 recognizes equally the α1-2 and α1-/3/4-linked fucosides may explain the high incidence of scedosporiosis in CF patients. Besides, it correlates with the presence of *SapL1* gene in all the pathogenic strains of *Scedosporium* whose genome has been sequenced to date. However, there is no evidence suggesting that the phenotype of blood group could have an influence in this recognition conversely to previous reports on other pathogens like *Pseudomonas aeruginosa* and *Haemophilus influenza.*[54-57]

Finally, the specificity and affinity of each one of the six binding pockets of SapL1 has been deeply analyzed to highlight the features that must be explored for the design of efficient inhibitors. Within our analysis, we have found that the binding pockets are non-equivalent but they all share the features necessary for fucose recognition. This example of divergence can also be found in the other lectins of the same family with FleA being the member with the greatest differences between pockets known to date.[38] It is remarkable that although SapL1 has a relatively low sequence identity with other members of this family, its specificity and affinity are

very similar to those reported for FleA from *A. fumigatus*,[49] AAL from the orange

peel mushroom *Aleuria aurantia*[51] and the bacterial lectins BambL and RSL, from

Burkholderia ambifaria and *Ralstonia solanacearum,* respectively.[58,59] Together with the

crystallographic data, this confirms that the structure and function of this lectin

family are highly conserved, settling the possibility for development of a broad-

spectrum therapy.

Overall, our research has revealed the first insights about the recognition of human

glycoconjugates by *S. apiospermum* lectins and contributes to the general

understanding of the host-binding process during the early stages of infection. The

detailed information exposed here, places SapL1 as a promising target for the

diagnosis and/or treatment of *Scedosporium* infections and it will be of great value to

guide the development of antiadhesive glycodrugs against this pathogen.

Material and methods

Production

The coding sequence for SapL1 was optimized for expression in *Escherichia coli* and

ordered at Eurofins Genomics (Ebersberg, Germany) before cloning into the

expression vectors pET-TEV,[60] pET32-TEV[49] and pProNde. pProNde is a homemade

vector where the NcoI restriction site of pProEx HTb (EMBL, Heidelberg) has been

replaced by a NdeI restriction site by PCR. Then, plasmids were introduced into *E.*

coli strains by thermal shock at 42°C and different expression parameters were

assayed (S1 Table) until soluble expression was achieved. The best set of conditions

for SapL1 expression were: *E. coli* TRX strain, pProNde vector, LB medium, cells

grown at 37°C and 160 rpm until 0.4 DO_{600}, then cultures transferred to 16°C until

0.8 OD_{600nm} and protein induction carried out overnight by the addition of isopropyl-

β-D-1-thiogalactoside (IPTG) to a final concentration of 0.05 mM and 1% L-

rhamnose (Rh).

Purification

Cells were subsequently harvested by centrifugation at 5000 g for 10 min and

resuspended in buffer A (50 mM Tris-HCl, 500 mM NaCl, pH 8.5). After addition of

1 µl of Denarase (C-LEcta GmbH, Leipzig, Germany) and moderate agitation during

30 minutes at room temperature, cells were disrupted at 1.9 kbar using a cell

disruptor (Constant Ltd Systems). Cell debris were removed by centrifugation at
22000 g for 30 minutes. The supernatant was filtered using 0.45 micrometer
membranes (PES, ClearLine) and the protein purification was carried out by IMAC
using 1 mL His-Trap FF columns (GE Healthcare Life Sciences) and a NGC™
chromatography system (Bio-Rad). After loading the supernatant, the column was
rinsed thoroughly with buffer A until stabilization of the baseline. Bound proteins
were eluted through the addition of buffer B (50 mM Tris-HCl, 500 mM NaCl, 500
mM Imidazole, pH 8.5) in an 0-500 mM imidazole gradient over 20 mL. Fractions
containing SapL1 were pooled and concentrated by ultrafiltration (Pall, 10kDa cut-
off) prior buffer exchange to 50 mM Tris-HCl, 100 mM NaCl, pH 8.5 using PD10
desalting columns (GE Healthcare Life Sciences). Then, the fusion was cleaved off
overnight at 19°C using the TEV protease produced in the lab in 1:50 ratio and
addition of 0.5 mM EDTA and 0.25 mM TCEP. The sample was loaded on the His-
trap column to separate the cleaved protein collected in the flow-through from the
TEV protease and potential uncleaved sample retained and eluted with imidazole.
Sapl1 containing fractions were concentrated by centrifugation to the desired
concentration.

Size exclusion chromatography
Size exclusion chromatography (SEC) was performed using a High-Resolution
ENrich™ SEC 70 column (Bio-Rad) on NGC™ chromatography system (Bio-Rad).
Column was equilibrated with 50 ml of buffer D (20 mM MES, 100 mM NaCl, pH
6.5) and 200 µl of sample at 10 mg mL^{-1} were injected into the system followed by 40
ml isocratic elution on buffer D supplemented with 20 mM α-methyl fucoside, 0.1%
L-rhamnose, 5 M NaCl, 2 M NaCl or 500 mM NaCl, according to the experiment.
Fractions were monitored by the absorbance at 280 nm and 0.5 mL fractions were
collected in the resolving region of the column.

Dynamic light scattering (DLS)
DLS analyses were performed using a Zetasizer™ NanoS (Malvern Panalytical) with
a 40 µl quartz cuvette. Measurements were performed in triplicate on protein sample
at 1 mg·ml^{-1} in buffer C (50 mM Tris-HCl, 100 mM NaCl, pH 8.5) after centrifugation.

Thermal Shift Assay (TSA)
The thermal stability of SapL1 was analyzed by TSA with the MiniOpticon real-time
PCR system (Bio-Rad). Prior assay, buffer stocks at 100 mM and a mixture

containing 70 μl of SapL1 at 1 mg ml^{-1}, 7 μl of 500x Sypro Orange (Merk Sigma-
Aldrich,) and 63 μl of ultrapure H_2O were prepared. Then, 7.5 μl of H_2O, 12.5 μl of
the corresponding buffer and 5 μl of the protein/Sypro mixture were mixed in 96-
well PCR microplates. The heat exchange test was then carried out from 20°C to
100°C with a heating rate of 1°C·min^{-1}. Fluorescence intensity was measured with
Ex/Em: 490/530 nm and the data processing was performed with the CFX Manager
software.

Isothermal Titration Calorimetry (ITC)

437   Experiments were performed using a Microcal ITC200 calorimeter (Malvern
438   Panalytical) with 40 μl of L-fucose 2.5 mM in the syringe and 200 μl of protein 0.04
439   mM in the sample cell. 2 μl injections in a range of 120 seconds while stirring at 1000
440   rpm were performed. Experimental data were adjusted to a theoretical titration
441   curve by the Origin ITC Analysis software.

Hemagglutination assay

Agglutination test was performed with fresh rabbit erythrocytes (bioMérieux, Lyon)
in U 96-well plates (Nalgene). For the test, 150 mM NaCl was used as a negative
control and 1 mg ml^{-1} LecB of *P. aeruginosa* as a positive control. 50 μl of sample was
prepared at 0.1 mg·ml^{-1} and submitted to serial double dilutions. 50 μl of rabbit
erythrocytes 3% were added to each well prior incubation of the plate at room
temperature. After 2 hours, the result of the experiment was evaluated and
agglutination activity was calculated according to the dilution of the protein.

Glycan arrays

Protein was labeled with Fluorescein Isothiocyanate (FITC) (Merk Sigma-Aldrich)
according to the supplier's instructions with slight modifications. Briefly two
milligrams of protein were dissolved in 1 ml of buffer E (100 mM Na_2CO_3, 100 mM
NaCl, pH 9); then, 40 μl of FITC at 1 mg·ml^{-1}, previously dissolved in DMSO, were
gradually added to the protein solution and the mixture was gently stirred at room
temperature overnight. Next day, the solution was supplemented with NH_4Cl to a
final concentration of 50 mM and free FITC was removed using PD10 column with
PBS as mobile phase. Protein concentration was determined at A_{280} and FITC at A_{490}
using a NanoDrop™ 200 (Thermo Scientific) and Fluorescein/Protein molar ratio
(F/P) was estimated by the following formula:

461
$$\text{Molar}\,\frac{F}{P} = \frac{MW}{389} \times \frac{\dfrac{A_{495}}{195}}{[A_{280} - (0.35 \times A_{495})]E^{0.1\%}}$$

Where MW is the molecular weight of the protein, 389 is the molecular weight of FITC, 195 is the
absorption E 0.1% of bound FITC at 490 nm at pH 13.0, (0.35 X A495) is the correction factor due to
the absorbance of FITC at 280 nm, and E 0.1% is the absorption at 280 nm of a protein at 1.0 mg·ml^{-1},
Being an ideal F/P should be 0.3 > 1.

Labeled lectin was sent to the Consortium for Functional Glycomics (CFG; Boston,
MA, USA) and its binding properties were assayed at 5 and 50 µg·ml^{-1} on a
"Mammalian Glycan Array version 5.4" which contain 585 glycans in replicates of
6. The highest and lowest signal of each set of replicates were eliminated and the
average of the remaining data was normalized to the percentages of the highest RFU
value for each analysis. Finally, the percentages for each glycan were averaged at
different lectin concentrations.

Crystallization and data collection
Crystal screening was performed using the hanging-drop vapour diffusion
technique by mixing equal volumes of pure protein at 5 mg·ml^{-1} and precipitant
solutions from commercial screenings of Molecular Dimensions (Newmarket, UK).
2 µl drops were incubated at 19°C until crystals appeared. A subsequent
optimization of positive conditions for SapL1 crystallization was carried out and
crystals suitable for X-ray diffraction analysis were obtained under solution
containing 100 mM Bicine pH 8.5, 1.5 M ammonium sulfate (NH$_4$)$_2$SO$_4$ and 12% v/v
glycerol. Crystals were soaked in mother liquor supplemented with 10% (v/v)
glycerol, prior to flash cooling in liquid nitrogen. Data collection was performed on
PX1 beamline at SOLEIL Synchrotron (Saint Aubin, FR) using a Pixel detector.

Structure determination
Data were processed using XDS [61]software and were converted to structure factors
using the CCP4 program package v.6.1,[62] with 5% of the data reserved for Rfree
calculation. The structure was determined using the molecular-replacement method
with Phaser v.2.5,[63] using the structure of FleA dimer (PDB entry 4D4U[38]) as starting
model. Model refinement was performed using REFMAC 5.8 [64] alternated with
manual model building in Coot v.0.7.[65] Sugar residues and other compounds that
were present were placed manually using Coot and validated using Privateer.[66] The

final model has been validated and deposited in the PDB Database with accession number 6TRV.

Figures

Figures were created using PyMOL Version 1.8.4 (Schrödinger), ChemDraw Version 15, ESPript Version 3.0[67] and PowerPoint 16.

Accession numbers

The structure of SapL1 was deposited as PDB ID: 6TRV.

Author Contributions: Conceptualization, A.V.; formal analysis, A.V. and D.M.A; investigation, D.M.A.; methodology A.V. and D.M.A; visualization A.V. and D.M.A; writing—original draft preparation, D.M.A.; writing—review and editing, A.V, B.J-P, R.P and D.M.A; supervision, A.V. and R.P.; project administration, A.V.; funding acquisition, A.V. All authors have read and agreed to the published version of the manuscript.

Funding: This project has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant (H2020-MSCA-ITN-2017-EJD-765581), from the French cystic fibrosis association Vaincre la Mucoviscidose. The project was also supported by Glyco@Alps (ANR-15-IDEX02).

Acknowledgments: We would like to thank to Valérie Chazalet and Emilie Gillon for their technical assistance. We also would like to thank for access to the beamlines BM30A-FIP at the European Synchrotron Radiation Facility (ESRF), Grenoble, France where initial tests were performed and Proxima 1 at SOLEIL Synchrotron, Saint Aubin, France (proposal number 20170827), where final data were collected. Thanks to our local contacts Jean-Luc Ferrer, Serena Sirigu, and Pierre Legrand for their assistance and technical support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tuite NL, Lacey K. Overview of invasive fungal infections. In: O'Connor L, Glynn B, editors. NY: Humana Press, Totowa, NJ; 2013. p. 1-23.

2. Ramirez-Garcia A, Pellon A, Rementeria A, Buldain I, Barreto-Bergter E, Rollin-
Pinheiro R, et al. (2018). *Scedosporium* and *Lomentospora*: An updated overview of
underrated opportunists. *Med Mycol.* 56:S102-S25.
3. Goldman C, Akiyama MJ, Torres J, Louie E, Meehan SA (2016). *Scedosporium*
apiospermum infections and the role of combination antifungal therapy and GM-
CSF: a case report and review of the literature. *Medical Mycol Case Rep.* 11:40-3.
4. Rammaert B, Puyade M, Cornely OA, Seidel D, Grossi P, Husain S, et al. (2019).
Perspectives on Scedosporium species and Lomentospora prolificans in lung
transplantation: Results of an international practice survey from ESCMID fungal
infection study group and study group for infections in compromised hosts, and
European Confederation. *Transpl Infect Dis.* 21:1-8.
5. Luplertlop N (2018). *Pseudallescheria/Scedosporium* complex species: from saprobic
to pathogenic fungus. *J Mycol Med.* 28:249-56.
6. Gilgado F, Cano J, Gené J, Guarro J (2005). Molecular phylogeny of the
Pseudallescheria boydii species complex: proposal of two new species. *J Clin*
Microbiol. 43:4930-42.
7. Duvaux L, Shiller J, Vandeputte P, Bernonville TDD, Thornton C, Papon N, et al.
(2017). Draft genome sequence of the human-pathogenic fungus *Scedosporium*
boydii. *Genome Announc.* 5:e00871-17.
8. Pérez-Bercoff Å, Papanicolaou A, Ramsperger M, Kaur J, Patel HR, Harun A, et
al. (2015). Draft genome of Australian environmental strain WM 09.24 of the
opportunistic human pathogen *Scedosporium aurantiacum*. *Genome Announc.*
3:e01526-14.
9. Vandeputte P, Ghamrawi S, Rechenmann M, Iltis A, Giraud S, Fleury M, et al.
(2014). Draft genome sequence of the pathogenic fungus *Scedosporium*
apiospermum. *Genome Announc.* 2:e00988-14.
10. Pereira De Mello T, Aor AC, Santiago S, De Oliveira C, Branquinha MH, Luis A
(2016). Conidial germination in *Scedosporium apiospermum, S . aurantiacum, S .*
minutisporum and *Lomentospora prolificans*; influence of growth conditions and
antifungal susceptibility profiles. *Mem Inst Oswaldo Cruz.* 111:484-94.
11. Tortorano AM, Richardson M, Roilides E, van Diepeningen A, Caira M, Munoz
P, et al. (2014). ESCMID and ECMM joint guidelines on diagnosis and
management of hyalohyphomycosis: *Fusarium spp., Scedosporium spp.* and others.
Clin Microbiol Infect. 20:27-46.
12. Troke P, Aguirrebengoa K, Arteaga C, Ellis D, Heath CH, Lutsar I, et al. (2008).
Treatment of scedosporiosis with voriconazole: clinical experience with 107
patients. *Antimicrob Agents Chemother.* 52:1743-50.
13. Rodriguez-tudela JL, Guarro J, Kantarcioglu AS, Horre R, Estrella MC, Berenguer
J, et al. (2006). *Scedosporium apiospermum:* changing clinical spectrum of a therapy-
refractory opportunist. *Med Mycol.* 44:295-327.

14. Cortez KJ, Roilides E, Quiroz-telles F, Meletiadis J, Antachopoulos C, Knudsen
T, et al. (2008). Infections Caused by Scedosporium spp. *Clin Microbiol Rev.* 21:157-
97.
15. Theuretzbacher U, Piddock LJV (2019). Non-traditional antibacterial therapeutic
options and challenges. *Cell Host Microbe.* 26:61-72.
16. Krachler AM, Orth K (2013). Targeting the bacteria-host interface strategies in
anti-adhesion therapy. *Virulence.* 4:284-94.
17. Ofek I, Hasty DL, Sharon N (2003). Anti-adhesion therapy of bacterial diseases:
prospects and problems. *FEMS Immunol Med Microbiol.* 38:181-91.
18. Figueiredo RT, Fernandez PL, Dutra FF, Gonza Y, Cristina L, Bittencourt VCB, et
al. (2010). TLR4 recognizes *Pseudallescheria boydii* conidia and purified
rhamnomannans. *J Biol Chem.* 285:40714-23.
19. Pinto MR, Gorin PAJ, Wait R, Mulloy B, Barreto-Bergter E (2005). Structures of
the O-linked oligosaccharides of a complex glycoconjugate from *Pseudallescheria
boydii. Glycobiology.* 15:895-904.
20. Pinto MR, De Sá ACM, Limongi CL, Rozental S, Santos ALS, Barreto-Bergter E
(2004). Involvement of peptidorhamnomannan in the interaction of
Pseudallescheria boydii and HEp2 cells. *Microbes Infect.* 6:1259-67.
21. Bittencourt VCB, Figueiredo RT, Silva RB, Moura DS, Fernandez PL, Sassaki GL,
et al. (2006). An a-glucan of *Pseudallescheria boydii* is involved in fungal
phagocytosis and Toll-like receptor activation. *J Biol Chem.* 281:22614-23.
22. Ghamrawi S, Renier G, Saulnier P, Cuenot S, Zykwinska A, Dutilh BE, et al.
(2014). Cell wall modifications during conidial maturation of the human
pathogenic fungus *Pseudallescheria boydii. PLoS ONE.* 9:e100290-e.
23. Rollin-Pinheiro R, Liporagi-lopes LC, Meirelles JVD, Souza LMD (2014).
Characterization of *Scedosporium apiospermum* glucosylceramides and their
involvement in fungal development and macrophage functions. *PLoS ONE.*
9:e98149-e.
24. De Mello TP, Aor AC, Gonçalves DDS, Seabra SH, Branquinha MH, Luis A
(2018). *Scedosporium apiospermum, Scedosporium aurantiacum, Scedosporium
minutisporum* and *Lomentospora prolificans*: a comparative study of surface
molecules produced by conidial and germinated conidial cells. *Mem Inst Oswaldo
Cruz.* 113:1-8.
25. Shoaf-sweeney KD, Hutkins RW. Adherence, anti-adherence, and
oligosaccharides: preventing pathogens from sticking to the host. 2009. p. 101-61.
26. Sharon N (2006). Carbohydrates as future anti-adhesion drugs for infectious
diseases. *Biochim Biophys Acta.* 1760:527-37.
27. Sharon N, Ofek I (2000). Safe as mother's milk: carbohydrates as future anti-
adhesion drugs for bacterial diseases. *Glycoconj J.* 17:659-64.

28. Tamburrini A, Colombo C, Bernardi A (2020). Design and synthesis of glycomimetics: Recent advances. *Med Res Rev.* 40:495-531.

29. Sattin S, Bernardi A (2016). Glycoconjugates and glycomimetics as microbial anti-adhesives. *Trends Biotechnol.* 34:483-95.

30. Pieters RJ (2011). Carbohydrate mediated bacterial adhesion. *Adv Exp Med Biol.* 715:227-40.

31. Visini R, Jin X, Bergmann M, Michaud G, Pertici F, Fu O, et al. (2015). Structural Insight into Multivalent Galactoside Binding to *Pseudomonas aeruginosa* Lectin LecA. *ACS Chem Biol.* 10:2455-62.

32. Houser J, Komarek J, Kostlanova N, Cioci G, Varrot A, Kerr SC, et al. (2013). A soluble fucose-specific lectin from *Aspergillus fumigatus* conidia - structure, specificity and possible role in fungal pathogenicity. *PLoS ONE.* 8:e83077-e.

33. Kerr SC, Fischer GJ, Sinha M, McCabe O, Palmer M, Choera T, et al. (2016). FleA expression in *Aspergillus fumigatus* is recognized by fucosylated structures on mucins and macrophages to prevent lung infection. *PLOS Pathog.*e100555-e.

34. Sakai K, Hiemori K, Tateno H, Hirabayashi J, Gonoi T (2019). Fucose-specific lectin of *Aspergillus fumigatus:* binding properties and effects on immune response stimulation. *Med Mycol.* 57:71-83.

35. Thornton DJ, Rousseau K, McGuckin MA (2008). Structure and function of the polymeric mucins in airways mucus. *Annu Rev Physiol.* 70:459-86.

36. Glick MC, Kothari VA, Liu A, Stoykova LI, Scanlin TF (2001). Activity of fucosyltransferases and altered glycosylation in cystic fibrosis airway epithelial cells. *Biochimie.* 83:743-7.

37. Lamblin G, Degroote S, Perini J-mM, Delmotte P, Scharfman A, Davril M, et al. (2001). Human airway mucin glycosylation: a combinatory of carbohydrate determinants which vary in cystic fibrosis. *Glycoconj J.* 18:661-84.

38. Houser J, Komarek J, Cioci G, Varrot A, Imberty A, Wimmerova M (2015). Structural insights into *Aspergillus fumigatus* lectin specificity : AFL binding sites are functionally non-equivalent. *Acta Crystallogr D Biol Crystallogr.*442-53.

39. Frank DE, Saecker RM, Bond JP, Capp MW, Tsodikov OV, Melcher SE, et al. (1997). Thermodynamics of the interactions of Lac repressor with variants of the symmetric Lac operator: effects of converting a consensus site to a non-specific site. *J Mol Biol.* 267:1186-206.

40. Capp MW, Cayley DS, Zhang W, Guttman HJ, Melcher SE, Saecker RM, et al. (1996). Compensating effects of opposing changes in putrescine (2+) and K+ concentrations on lac repressor-lac operator binding: in vitro thermodynamic analysis and in vivo relevance. *J Mol Biol.* 258:25-36.

41. Cayley S, Lewis BA, Guttman HJ, Record MT (1991). Characterization of the cytoplasm of *Escherichia coli K-12* as a function of external osmolarity implications for protein-DNA interactions in vivo. *J Mol Biol.* 222:281-300.

42. Richeys B, Cayley DS, Mossing MC, Kolka C, Anderson CF, Farrar TC, et al.
(1987). Variability of the intracellular ionic environment of *Escherichia coli.*
Differences between *in vitro* and *in vivo* effects of ion concentrations on protein-
DNA interactions and gene expression. *J Biol Chem.* 262:7157-64.

43. Harries D, Rosgen J. A practical guide on how osmolytes modulate
macromolecular properties. Elsevier; 2008. p. 679-735.

44. Massiah MA, Wright KM, Du H (2016). Obtaining soluble folded proteins from
inclusion bodies using sarkosyl, Triton X-100, and CHAPS: application to LB and
M9 minimal media. *Curr Protoc Protein Sci.* 84:1-24.

45. Pettitt BM, Bolen DW (2005). Protein folding, stability, and solvation structure in
osmolyte solutions. *Biophys J.* 89:2988-97.

46. Singh LR, Poddar NK, Dar TA, Rahman S, Kumar R, Ahmad F (2011). Forty years
of research on osmolyte-induced protein folding and stability. *J Iran Chem Soc.* 8:1-
23.

47. Ajito S, Iwase H, Takata S-i, Hirai M (2018). Sugar-mediated stabilization of
protein against chemical or thermal denaturation. *The Journal of Physical Chemistry*
B. 122:8685-97.

48. Bonnardel F, Kumar A, Wimmerova M, Lahmann M, Perez S, Varrot A, et al.
(2019). Architecture and evolution of blade assembly in β-propeller lectins.
Structure. 27:764-75.e3.

49. Lehot V, Brissonnet Y, Dussouy C, Brument S, Cabanettes A, Gillon E, et al.
(2018). Multivalent fucosides with nanomolar affinity for the *Aspergillus fumigatus*
lectin FleA prevent spore adhesion to pneumocytes. *Chem Eur J.* 24:19243-9.

50. Makyio H, Shimabukuro J, Suzuki T, Imamura A, Ishida H, Kiso M, et al. (2016).
Six independent fucose-binding sites in the crystal structure of *Aspergillus oryzae*
lectin. *Biochem Biophys Res Commun.* 477:477-82.

51. Wimmerova M, Mitchell E, Sanchez JF, Gautier C, Imberty A (2003). Crystal
structure of fungal lectin. Six-bladed β-propeller fold and novel fucose
recognition mode for *Aleuria aurantia* lectin. *J Biol Chem.* 278:27059-67.

52. Debourgogne A, Dorin J, Machouart M (2016). Emerging infections due to
filamentous fungi in humans and animals: only the tip of the iceberg? *Env*
Microbiol Rep. 8:332-42.

53. Hogan LH, Klein BS, Levitz SM (1996). Virulence factors of medically important
fungi. *Clin Microbiol Rev.* 9:469-88.

54. Kuo KC, Kuo HC, Huang LT, Lin CS, Yang SN (2013). The clinical implications
of ABO blood groups in *Pseudomonas aeruginosas* sepsis in children. *J Microbiol*
Immunol Infect. 46:109-14.

55. Gilboa-Garber N, Sudakevitz D, Sheffi M, Sela R, Levene C (1994). PA-I and PA-
II lectin interactions with the ABO(H) and P blood group glycosphingolipid

antigens may contribute to the broad spectrum adherence of *Pseudomonas*
aeruginosa to human tissues in secondary infections. *Glycoconj J.* 11:414-7.

56. Scanlin TF, Glick MC (1999). Terminal glycosylation in cystic fibrosis. *Biochim*
Biophys Acta. 1455:241-53.

57. Taylor-Cousar JL, Zariwala MA, Burch LH, Pace RG, Drumm ML, Calloway H,
et al. (2009). Histo-blood group gene polymorphisms as potential genetic
modifiers of infection and cystic fibrosis lung disease severity. *PLoS ONE.*
4:e4270-e.

58. Audfray A, Claudinon J, Abounit S, Ruvoe N, Smith DF, Pendu JL, et al. (2012).
Fucose-binding lectin from opportunistic pathogen *Burkholderia ambifaria* binds to
both plant and human oligosaccharidic epitopes. *J Biol Chem.* 287:4335-47.

59. Kostlanova N, Mitchell EP, Lortat-Jacob H, Oscarson S, Lahmann M, Gilboa-
Garber N, et al. (2005). The fucose-binding lectin from *Ralstonia solanacearum*. A
new type of β-propeller architecture formed by oligomerization and interacting
with fucoside, fucosyllactose, and plant xyloglucan. *J Biol Chem.* 280:27839-49.

60. Houben K, Marion D, Tarbouriech N, Ruigrok RWH, Blanchard L (2007).
Interaction of the C-terminal domains of sendai virus N and P proteins:
comparison of polymerase-nucleocapsid interactions within the paramyxovirus
family. *J Virol.* 81:6807-16.

61. Kabsch W (2010). XDS. *Acta Crystallographica.* 66:125-32.

62. Winn MD, Ballard CC, Cowtan KD, Dodson EJ, Emsley P, Evans PR, et al. (2011).
Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol*
Crystallogr. 67:235-42.

63. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ
(2007). Phaser crystallographic software. *J Appl Cryst.* 40:658-74.

64. Murshudov GN, Nicholls RA (2011). REFMAC 5 for the refinement of
macromolecular crystal structures. *Acta Crystallogr D Biol Crystallogr.* 67:355-67.

65. Emsley P, Lohkamp B (2010). Features and development of Coot. *Acta Crystallogr*
D Biol Crystallogr. 66:486-501.

66. Aguirre J, Iglesias-Fernandez J, Rovira C, Davies GJ, Wilson KS, Cowtan KD
(2015). Privateer: software for the conformational validation of carbohydrate
structures. *Nat Struct Mol Biol.* 22:833-4.

67. Robert X, Gouet P (2014). Deciphering key features in protein structures with the
new ENDscript server. *Nucleic Acids Res.* 42:320-4.

 Figure captions

 Figure 1. SapL1 production. A) Schematic representation of the genetic constructs
used for the expression of SapL1. B) SDS-PAGE of fractions collected from SapL1
purification using pET32-TEV vector. IF: Insoluble fraction, SF: Soluble fraction, FT:
Flow through, W: wash and elution, from right to left respectively. Red arrow
indicates inclusion bodies from insoluble fractions. C) Representative
chromatograms of SapL1 purification before and after the process with their
respective profile on 15% SDS-polyacrylamide gels (insoluble fraction, soluble
fraction, flow through, washing and elution, from left to right respectively). The
fractions containing SapL1 are delimited.

Figure 2. Rhamnose influence on SapL1 solubility. A) SapL1 purification
chromatograms at 0.1%, 0.2%, and 1% rhamnose with their corresponding SDS-
PAGE profile. IF: Insoluble fraction, SF: Soluble fraction, FT: Flow through, W: wash
and elution, from left to right respectively. Black rectangles indicate the elution
peaks and elution fractions containing SapL1 on SDS-polyacrylamide gels. B)
Numerical values derived from the integration of areas under the peaks
corresponding to SapL1 for each experiment. Black and striped bars represent the
results for experiments performed at 16°C and 20°C, respectively.

Figure 3. SapL1 carbohydrate binding properties. A) Hemagglutination assay of
SapL1 on fresh rabbit erythrocytes. Negative and positive controls consist of 150 mM
NaCl and the lectin LecB from *Pseudomonas aeruginosa*, respectively. B) Titration of
SapL1 with L-fucose and α-methyl-fucoside with the thermogram and the
integration displayed at the top and bottom, respectively. C) Analysis of the
interactions of SapL1 with blood group epitopes. The graph shows the relative
binding of SapL1 to glycans containing Lewis and ABH blood group antigens from
the 90 hits identified as binders.

Figure 4: SapL1 overall structure. A) Surface and cartoon representation of SapL1
monomer colored from blue (*N*-terminal end) to red (*C*-terminal end). BP: Binding
pockets. B) Representation of SapL1 dimer colored by monomer with ligand
depicted in sticks.

Figure 5: SapL1/FleA comparison. A) Overlay of SapL1 (blue, PDB: 6TRV)) and
FleA (green, PDB: 4D4U) structures. Red arrows indicate the main differences
between structures. B) Sequence alignment of SapL1 and FleA with display of their

secondary structure elements. The residues involved in ligand interactions of each pocket are indicated by circles for hydrophobic and stacking interactions and by triangles for hydrogen bonds colored according to the binding pocket: BP-1, blue; BP-2 cyan; BP-3 green; BP-4, orange, BP-5, brown; BP-6, red.

Figure 6: SapL1-ligand interactions. A) The sequence alignment of the 6 blades of SapL1. The residues involved in ligand binding interactions are indicated in solid boxes: BP-1, blue; BP-2 cyan; BP-3 green; BP-4, orange, BP-5, brown; BP-6, red. Striped green boxes indicate the four additional residues expected to be involved in fucose binding within BP-3 that were not visible in the structure since only glycerol was attached to this pocket. B) Zoom on the interactions of ligand with each binding site of SapL1 and their schematic representation.

763 **Table 1.** Data-collection and refinement statistics.

Data collection		
Beamline	SOLEIL Proxima 1	
Wavelength (A°)	0.97857	
Space group	$P2_1$	
Unit cell dimensions a, b, c (Å), α, β, γ (°)	76.06, 45.66, 83.48, 90, 105.05, 90	
No. of monomers in ASU	2	
Resolution (Å)	40.0 – 2.4 (2.46-2.4)	
*R*merge	0.085 (0.428)	
*R*pim	0.07 (0.356)	
Mean I/σ (I)	5.3 (1.6)	
Completeness (%)	98.2 (98.4)	
Multiplicity	2.7 (2.7)	
CC1/2	0.991(0.7)	
No. reflections /No. Unique reflections	58305/ 21570	
Refinement		
Resolution (Å)	20.0-2.3	
No. of reflections in working set / Free set	21561 / 1089	
R work/ R free	17.5 / 24.0	
R.m.s Bond lengths (Å)	0.017	
Rmsd Bond angles (°)	1.94	
Rmsd Chiral (Å³)	0.087	
No. atoms / Bfac (Å²)	Chain A	Chain B
Protein	2222/32.64	2220/40.83
Ligand and heterogen	89/31.56	66/36.914
Waters	111/31.018	69/33.206
Ramachandran Allowed / Favored/ Outliers (%)	96 / 4 / 0	95 / 4 / 1
PDB Code	6TRV	

764 *Values in parentheses are for the outer shell

767   Figure 1

771   Figure 2

SapL1 monomer

SapL1 dimer

774   Figure 4

776   Figure 5

A)

```
SapL1      1        10                  20         30              40              50
Blade_1    MSGVLQISFPAGIAAI.....RNNSSLRVYEAALD..GGVREAQYEG...RWAGGKPDNVI.
Blade_2    ....ATGKIGTPIAATS..VG.HQYTRVYYVGAD..NKAREACKDG.K.GWYTGAFV....
Blade_3    ...KDVAPYSSIGAVFL...GKNIVVRVYTQNHD.NTIQEWVWDSPSTGWTAGANFGAA.
Blade_4    .......LPGTAIAATSWGAG.PYHIRVYFQDTN.RNVIESGWDG.S.GWYTGGLKISN.
Blade_5    ......QSPRASLGATSWGESGSSLGIRLYYATQD.NLIKEKAWDGGG.GWYDGGFQQRS.
Blade_6    ........IPGSRVAAIP......LPVLRVYLQNGTEVSGITEYAWNS...GWVVGCAVLPPA
```

B)

Figure 6