Hippocampal-cortical encoding activity predicts the precision of episodic memory

Saana M. Korkki[1], Franziska R. Richter[2], Jon S. Simons[1]

[1]Department of Psychology, University of Cambridge, Cambridge CB2 3EB, United Kingdom

[2]Cognitive Psychology Unit, Institute of Psychology, University of Leiden, Leiden 2333 AK, Netherlands

Corresponding authors: Saana M. Korkki, smk62@cam.ac.uk; Jon S. Simons, jss30@cam.ac.uk

Acknowledgements

This study was funded by BBSRC grant BB/L02263X/1 and James S. McDonnell Foundation Scholar Award #220020333, and was carried out within the University of Cambridge Behavioural and Clinical Neuroscience Institute, funded by a joint award from the Medical Research Council and the Wellcome Trust. We are grateful to Paul Bays for valuable advice and to the staff of the MRC Cognition and Brain Sciences Unit MRI facility for scanning assistance.

Abstract

Our recollections of past experiences can vary both in the number of specific event details accessible from memory and the precision with which such details are reconstructed. Prior neuroimaging evidence suggests the success and precision of episodic recollection to rely on distinct neural substrates during memory *retrieval*. In contrast, the specific *encoding* mechanisms supporting later memory precision, and whether they differ from those underlying successful memory formation in general, are currently unknown. Here, we combined continuous measures of memory retrieval with model-based analyses of behavioural and neuroimaging data to tease apart the encoding correlates of successful memory formation and mnemonic precision. In the MRI scanner, participants encoded object-scene displays, and later reconstructed features of studied objects using a continuous scale. We observed overlapping encoding activity in inferior prefrontal and posterior perceptual regions to predict both which object features were later remembered versus forgotten, and the precision with which they were reconstructed from memory. In contrast, hippocampal encoding activity significantly predicted the precision, but not overall success, of subsequent memory retrieval. The current results identify a hippocampal-cortical encoding basis for episodic memory precision, and suggest a contribution of shared cortical encoding mechanisms to the formation of both accessible and precise memory representations.

Introduction

Our memories are not an exact reproduction of the past, but can range from high-fidelity, precise, reconstructions of previous experiences to less accurate, lower-resolution, representations (Brady et al., 2013; Harlow & Donaldson, 2013; Harlow & Yonelinas, 2016). Prior studies suggest such variation in mnemonic precision to be underpinned by at least partly distinct neurocognitive mechanisms from those underlying the binary success vs. failure of memory retrieval (Cooper & Ritchey, 2019; Nilakantan et al., 2017, 2018; Richter et al., 2016). Indeed, during *retrieval*, these two memory components have been shown to rely on dissociable regions of the posterior-medial memory network (Richter et al., 2016; Cooper et al., 2017). However, despite increased interest in the neural basis of mnemonic precision (e.g., Cooper & Ritchey, 2019; Montchal et al., 2019; Richter et al., 2016; Stevenson et al., 2018), the associated encoding mechanisms, and whether they differ from those supporting successful encoding in general, remains unresolved.

Successful episodic memory formation is typically associated with activity increases in a network of medial temporal, lateral prefrontal, and posterior perceptual regions (Kim, 2011; Spaniol et al., 2009). The hippocampus receives input from content-specific perceptual regions, and is thought to bind disparate event features into a coherent memory representation (Cooper & Ritchey, 2020; Davachi, 2006; Paller & Wagner, 2002; Ranganath, 2010), and allow for the storage of similar experiences in an orthogonalised, or non-overlapping, manner (Norman & O'Reilly, 2003; O'Reilly & McClelland, 1994), whereas lateral prefrontal regions are involved in the goal-driven and controlled encoding of information into memory (Blumenfeld & Ranganath, 2007; Simons & Spiers, 2003). The specific neural substrates supporting successful memory formation have been found to vary depending on informational content

(Cooper & Ritchey, 2020; Staresina et al., 2011; Uncapher et al., 2006), the depth of stimulus processing engaged in (Fletcher et al., 2003; Otten et al., 2001), and the type of retrieval process later recruited (Ranganath et al., 2004; Staresina & Davachi, 2006). Moreover, encoding correlates appear sensitive to more subtle differences in the quality of retained representations, including their amount of detail (Cooper & Ritchey, 2020; Qin et al., 2011), and subjective vividness (Kensinger et al., 2011).

It is possible that, in addition to being dissociable at *retrieval* (Richter et al., 2016), the success and precision of episodic recollection may be supported by at least partly distinct neural mechanisms during memory *encoding*. For instance, the successful retrieval of information from memory may depend on the strength of an association between a retrieval cue and the target memory, thus drawing in particular on associative encoding processes supported by the hippocampus and the prefrontal cortex (Blumenfeld & Ranganath, 2007; Davachi, 2006). In contrast, the precision with which specific mnemonic features can be reconstructed from memory may closely relate to the fidelity of stimulus encoding in posterior perceptual regions (Emrich et al., 2013), and/or to hippocampal function supporting the formation of distinct and detailed memory traces (Moscovitch et al., 2016; Yonelinas, 2013). Alternatively, it is possible that the success and precision of episodic retrieval rely on similar encoding mechanisms that perhaps act to increase the strength of the memory more generally, rendering it both accessible and precise at retrieval (Schurgin et al., 2020).

In the current study, we employed continuous measures of memory retrieval and model-based analyses of behavioural and neuroimaging data to elucidate the encoding substrates of mnemonic precision. In the MRI scanner, participants

encoded visual stimulus displays depicting an object overlaid on a scene background. The location and colour of the objects were drawn from circular spaces, and at retrieval participants recreated these attributes of the studied items using a continuous response dial. This approach allowed us to segregate encoding activity supporting later successful memory retrieval from that supporting subsequent mnemonic precision in a manner not afforded by more typical categorical measures of retrieval performance (e.g., old/new, remember/know), thus providing novel insight into the encoding mechanisms supporting the acquisition of high-fidelity episodic memories.

Methods

Participants

Twenty-one young adults (18-29 years old) participated in the current experiment. All participants were right-handed, native English-speakers, had normal or corrected-to-normal vision, no colour blindness, and no current or historical diagnosis of any neurological, psychiatric, or developmental disorder, or learning difficulty. Participants indicated no current use of any psychoactive medication, and no medical or other contradictions to MRI scanning. One participant was excluded from all analyses due to excessive movement (> 4mm) in the scanner, leaving 20 participants to contribute to the present analyses (8 male, 12 female; mean age 22.15 years, *SD*: 3.10). The participants were recruited via the University of Cambridge Psychology Department Sona volunteer recruitment system (Sona Systems, Ltd) and community advertisements, and were reimbursed with £30 for

their participation. All participants gave written informed consent in a manner approved by the Cambridge Psychology Research Ethics Committee.

Materials

Stimuli for the memory task comprised 180 images of outdoor scenes and 180 images of distinct everyday objects. The images were obtained from existing stimuli sets (scenes: Richter et al., 2016; objects: Brady et al., 2013) and Google image search. Each object image was randomly paired with a scene image to generate a total of 180 trial-unique study displays (size 750 x 750 pixels). Across the study displays, we varied the appearance of two object features: colour and location. For each display, object colour and location were randomly selected from circular parameter spaces (0-360°) (cf. Cooper et al., 2017; Richter et al., 2016) (see Figure 1). All participants viewed the same study displays in a randomized order.

Design and procedure

Prior to the scan, participants read the instructions and undertook practice trials of the memory task. The task was adapted from Richter et al. (2016) to optimize the investigation of encoding-related effects, and comprised 9 study-test blocks completed over 9 functional runs (one study and one test phase per run). At study, participants sequentially viewed 20 object-background displays (stimulus duration: 5s), and were instructed to try and memorise the appearance of each display, including the location and colour of the object. The study phase was followed by a 10s delay during which a "Get Ready" message was presented on a black screen. Following this delay, participants were asked to reconstruct *either* the location *or* the

colour of each object viewed in the preceding study phase (one feature question per each encoding trial, total of 20 retrieval trials per block). At retrieval, the test object reappeared on its associated background with a central cue word "Location" or "Colour" indicating the type of feature tested on that trial. The initial appearance of the tested feature was randomly selected from a circular parameter space (0-360°), while the appearance of the untested feature remained unchanged from study to test. In other words, for location trials, the test object reappeared in its original colour, but in a randomly selected location, whereas for colour trials the test object reappeared in its original location but in a randomly chosen colour. Participants were asked to recreate the object's original features as accurately as they could by moving a slider around a 360-degree response dial using their middle and index finger on a button box, and were able to confirm their answer by pressing a third key with their thumb. The retrieval phase was self-paced with the constraint of a minimum trial length of 7s and a maximum response time of 11s. Participants on average produced response times that were well under this limit (*M:* 5.64s, *SD:* 0.68s), and the percentage of trials where response selection was not confirmed in time was very low (*M*: 1.36%, *SD*: 1.77%). Note that if a participant failed to confirm their answer within 11s, their last position on the response wheel was recorded as their answer for that trial.

Participants completed 90 location and 90 colour trials in total (10 trials of each type per task block). The type of feature tested for each object was randomised across displays, but constant across participants. To ensure that memory was tested for feature values spanning the entire circular space, this randomisation was conducted with a constraint of roughly equal number (i.e., 20-25) of target feature values sampled from each quadrant around the circular space for both the location and

colour condition. The order of study and test displays was randomised across participants with the constraint of no more than four encoding or retrieval trials in a row for which the same type of feature was tested. Study and test trials were separated by a fixation cross with jittered duration between 0.4s and 2.4s (mean ISI duration: 1s) following an approximate Poisson distribution. After the first five of the nine task blocks, participants were given a 10-minute break from the memory task in the scanner, during which a diffusion-weighted structural scan was acquired (analysis of diffusion-weighted data not reported here).

Figure 1. Example study and test trials of the memory task. At study, participants viewed stimuli displays consisting of one object overlaid on a scene background (stimulus duration: 5s). The location and colour of the objects at study were randomly chosen from circular parameter spaces (0-360°). At test, participants

recreated *either* the location *or* the colour of each studied object using a 360-degree continuous response dial, allowing for a fine-grained assessment of memory fidelity.

Behavioural analysis

For each trial, we calculated participants' retrieval error as the angular difference between their response value and the target feature value (0 ± 180°). To distinguish the likelihood of successful memory retrieval from the precision of the retrieved information, we fitted a two-component mixture model (Bays et al., 2009; Zhang & Luck, 2008) to each participant's retrieval error data using maximum likelihood estimation (code available at: https://www.paulbays.com/code/JV10/index.php). This mixture model has previously been shown to characterize long-term memory performance in similar tasks (e.g., Brady et al., 2013; Korkki et al., 2020; Richter et al., 2016), and has been employed to gain insights about the neural basis of the precision of episodic recollection (Cooper & Ritchey, 2019; Richter et al., 2016; Stevenson et al., 2018). The model assumes that two distinct sources of error contribute to participants' retrieval performance across trials: variability, that is, noise, in successful retrieval of target features, and the presence of random guess responses where memory retrieval has failed to bring any diagnostic information about the target to mind. These two sources of error are modelled by a von Mises distribution (circular equivalent of a Gaussian distribution) centred at a mean error of zero degrees from the target value, with a concentration, K, and a circular uniform distribution with a probability, pU, respectively. Precision of memory retrieval can be estimated as the concentration parameter (K, higher values reflect higher precision) of the target von Mises distribution, and the likelihood of successful memory retrieval

(pT) as the probability of responses stemming from the target von Mises distribution ($pT = 1 - pU$). Note that a low number of guess trials for some individual subjects prevented the investigation of feature-specific fMRI effects, however, for completeness, we report participants' behavioural performance both across features and separated by feature condition.

MRI acquisition

MRI scanning took place at the University of Cambridge Medical Research Council Cognition and Brain Sciences Unit using a 3T Siemens Tim Trio scanner (Siemens, Germany) with a 32-channel head coil. For each participant, a whole brain structural image was acquired using a T1-weighted 3D magnetization prepared rapid gradient echo (MPRAGE) sequence (repetition time (TR): 2.25s, echo time (TE): 3ms, flip angle = 9°, field of view (FOV): 256 x 256 x 192mm, resolution: 1mm isotropic, GRAPPA acceleration factor 2). Functional data were acquired over 9 runs each comprising one task block (one encoding and one retrieval phase), using a single-shot echoplanar imaging (EPI) sequence (TR: 2s, TE: 30ms, flip angle° = 78, FOV: 192 x 192mm, resolution: 3mm isotropic). Each volume consisted of 32 sequential oblique-axial slices (interslice gap: 0.75mm) acquired parallel to the anterior commissure – posterior commissure transverse plane. Across the participants, the mean number of volumes acquired per functional run was 161.60 (*SD*: 9.63). The scanning protocol further included a diffusion-weighted structural scan that was acquired after the first five functional runs (not analysed here).

fMRI preprocessing

Data preprocessing and analysis was performed with Statistical Parametric Mapping (SPM) 12 (https://www.fil.ion.ucl.ac.uk/spm/) implemented in MATLAB R2016a. The first five volumes of each functional run were discarded to allow for T1 equilibration. Furthermore, any additional volumes collected after each task block had finished were discarded for each participant so that the last volume of each run corresponded to a time point of ~2s after the last fixation cross. The functional images were spatially realigned to the mean image to correct for head motion and temporally interpolated to the middle slice to correct for differences in slice acquisition time. The anatomical image was coregistered to the mean EPI image, bias-corrected and segmented into different tissue classes (grey matter, white matter, cerebrospinal fluid). These segmentations were used to create a study-specific structural template image using the DARTEL (Diffeomorphic Anatomical Registration Through Exponentiated Lie Algebra) toolbox (Ashburner, 2007). The functional data was normalized to MNI space using DARTEL and spatially smoothed with an isotropic 8mm full-width at half-maximum (FWHF) Gaussian kernel.

fMRI analysis

In order to obtain trial-specific estimates of the success and precision of memory retrieval for the fMRI analyses, we fitted the two-component mixture model (von Mises + uniform distribution) to all retrieval errors across all participants (3600 trials in total), and calculated the cut-off point at which the probability of participants' responses stemming from the target von Mises distribution was less than 5%, beyond which responses likely reflected guessing (cf., Cooper et al., 2017; Richter et

al., 2016). This cut-off point was established as +/- 51 degrees in the current data, and was then used to classify each encoding trial as successful (absolute subsequent retrieval error ≤ 51°) or unsuccessful (absolute subsequent retrieval error > 51 degrees). For successful encoding trials, we further calculated a trial-specific measure of memory precision as 180 – participant's absolute retrieval error on that trial so that higher values (smaller error) reflected higher precision (range: 129 – 180°). A measure of memory precision was not calculated for unsuccessful encoding trials as retrieval errors in this condition were approximately randomly distributed and thus did not carry meaningful information about memory quality.

For each participant, a first level General Linear Model (GLM) was constructed containing three regressors corresponding to each event of interest (successful location encoding, successful colour encoding, unsuccessful encoding), and a fourth regressor modelling the retrieval trials. For the successful encoding trials, the trial-specific estimates of memory precision were included as parametric modulators comprising two additional regressors in the model. The precision parametric modulators were rescaled to range between 0 and 1 to facilitate the direct comparison of success and precision-related effects, and mean centred for each participant. Neural activity was modelled with a boxcar function convolved with the canonical hemodynamic response function (HRF), with a duration of 5s for the encoding trials and a variable duration (7s-11s) for the retrieval trials, capturing the duration of the study and test displays, respectively. Six participant-specific movement parameters estimated during realignment (3 rigid-body translations, 3 rotations) were further included as covariates in the first level model to capture any residual movement-related artefacts. Due to the small number of guessing trials in each functional run, data from all functional runs were concatenated for each

participant, and 9 constant block regressors included as additional covariates. Autocorrelation in the data was estimated with an AR(1) model and a temporal high pass filter with a 1/128 Hz cut-off was used to eliminate low frequency noise. First level subject-specific parameter estimates were submitted to second level random effects analyses.

Contrasts

The contrasts for the fMRI analyses focused on identifying regions where encoding activity positively predicted the subsequent success and/or precision of episodic memory retrieval (i.e., increases in BOLD signal for successful encoding, or higher memory precision). To examine encoding activity associated with the subsequent success of memory retrieval, we contrasted encoding trials for which memory retrieval subsequently succeeded against trials for which memory retrieval subsequently failed (*subsequent retrieval success effects*). To identify encoding activity predicting the later precision of memory retrieval, positive correlations between BOLD signal and the precision parametric modulator were examined (i.e., linear relationship between BOLD signal and precision parametric modulator; *subsequent precision effects*). We further assessed the overlap between subsequent success and subsequent precision effects using conjunction analyses. Conjunction analyses were conducted testing against the conjunction null hypothesis to ensure that regions identified in this analysis displayed reliable encoding activity associated with each individual contrast, i.e., both subsequent success and subsequent precision of memory retrieval (see Nichols et al., 2005).

Regions of interest

The main analyses focused on a small number of a priori regions of interest (ROIs) implicated by meta-analytic evidence in supporting the successful formation of episodic memories for visual information (Kim, 2011; Spaniol et al., 2009). Specifically, the ROIs included the hippocampus (HC), the inferior frontal gyrus (IFG) and the fusiform gyrus (FFG). Given evidence for greater consistency of subsequent memory effects in the left hemisphere (Spaniol et al., 2009), left-lateralized ROIs were used, each comprising the left anatomical region as defined by the Automated Anatomical Labelling (AAL) atlas. Statistical significance within each anatomical ROI was assessed using small-volume correction with a peak-level familywise error (FWE) corrected (based on random field theory) threshold of $p < .05$, correcting for the number of voxels in each ROI. In addition to the ROI analyses, we sought to identify any additional brain regions displaying a relationship between encoding activity and the subsequent success and/or precision of memory retrieval in exploratory whole brain analyses conducted at a whole brain FWE-corrected threshold of $p < .05$, minimum extent of 5 contiguous voxels.

Results

Behavioural results

For each trial, we calculated participants' retrieval error as the angular difference between their response value and the target feature value (0±180°) (see Figure 2A). Across participants and feature conditions, overall task performance, as measured by the mean absolute retrieval error, was 30.43° (*SD*: 15.04°), with significantly higher mean absolute error in the colour (*M*: 34.48°, *SD*: 15.90°) in comparison to

the location condition (*M*: 26.37°, *SD*: 15.80°), *t*(19) = 3.63, *p* = .002, *d* = 0.81. To

further decompose the specific sources of error contributing to participants' overall

performance, we fitted the two component mixture model (von Mises + uniform

distribution) to each individual participant's retrieval error data using maximum

likelihood estimation (Bays et al., 2009). The mean model-estimated probability of

successful memory retrieval, defined as the probability of responses stemming from

a von Mises distribution centred at the target feature value (*pT*), was 0.73 (*SD*: 0.18)

across participants (see Figure 2B). The mean model-estimated precision of memory

retrieval, estimated as the concentration parameter, *K*, of the target von Mises

distribution, was 16.79 (*SD*: 7.92) across participants (see Figure 2B) (note that this

value of *K* is comparable to an *SD* of approximately 14.20°). Mean memory precision

(*K*) was significantly higher in the location (*M*: 34.65, *SD*: 27.24) in comparison to the

colour condition (*M*: 10.94, *SD*: 7.15), *t*(19) = 4.04, p = 001, *d* = 0.90, whereas mean

probability of successful memory retrieval (*pT*) did not significantly differ between the

two feature conditions (location *M*: 0.75, *SD*: 0.18; colour *M*: 0.73, *SD*: 0.20), *t*(19) =

0.65, *p* = .524. Consistent with previous results (Richter et al., 2016), we also

observed a moderate positive correlation between estimates of the probability of

successful memory retrieval and memory precision across participants, r_s = .54, *p* =

.014.

Figure 2. A) Distribution of retrieval errors (response – target) across all trials and participants. Black line illustrates response probabilities predicted by the two-component mixture model (von Mises + uniform distribution; model fitted to data across all participants for visualization). B) Mean model-estimated probability of successful memory retrieval (*pT*) and memory precision (*K*) across participants. Error bars display 95% confidence interval of the mean and data points individual participant parameter estimates.

fMRI results

Encoding activity predicting subsequent retrieval success and memory precision in a priori ROIs

Our ROI analyses focused on examining whether encoding activity in three regions typically displaying subsequent memory effects for visual information; the hippocampus, the inferior frontal gyrus and the fusiform gyrus, differentially contributes to the later success and precision of episodic memory retrieval. We first examined increases in encoding activity for trials that were subsequently successfully retrieved (absolute retrieval error ≤ 51°) in contrast to trials that were subsequently forgotten (absolute retrieval error > 51°). Within our anatomical ROIs,

we observed increased encoding activity in the inferior frontal gyrus, $t(19) = 6.75$, $p = .001$, peak: -36, 27, 18, and the fusiform gyrus, $t(19) = 8.88$, $p < .001$, peak: -30, -63, -9, to predict whether object features were later successfully retrieved from memory or forgotten (see Figure 3A and 4A). In contrast, no significant subsequent retrieval success effects were detected in the hippocampus (ps > .151).

We next examined whether encoding activity in these regions predicted the graded precision with which object features were later successfully retrieved from memory (linear relationship between BOLD signal and precision parametric modulator). In addition to predicting which trials were successfully remembered, encoding activity in the inferior frontal gyrus, $t(19) = 5.63$, $p = .011$, peak: -57, 15, 15, and the fusiform gyrus, $t(19) = 6.27$, $p = .001$, peak: -33, -75, -18, positively correlated with the precision of later memory retrieval (see Figure 3B and 4B). Furthermore, increased encoding activity in the hippocampus, $t(19) = 4.20$, $p = .029$, peak: -33, -30, -9, was associated with greater mnemonic precision for object features (see Figure 3B and 3C).

Figure 3. Mean parameter estimates for A) subsequent success and B) subsequent

precision effects in the left inferior frontal gyrus (IFG), hippocampus (HC) and

fusiform gyrus (FFG). Error bars display +/- 1 SEM. C) Encoding activity correlating

with the subsequent precision of memory retrieval in the hippocampal ROI

(visualised at an uncorrected threshold of *p* < .001).

Thus, results from the ROI analyses suggest encoding activity in the inferior frontal

and fusiform cortex to support both the later success and precision of memory

retrieval, while significant increases in BOLD signal in the hippocampus were

observed for subsequent memory precision only. We next sought to assess whether

encoding activity predicting these two aspects of later retrieval performance

overlapped in any of the ROIs. Conjunction analyses indicated significant overlap

between subsequent success and subsequent precision effects in both the inferior

frontal $t(19) = 4.86$, $p = .007$, peak: -42, 3, 27, and the fusiform gyrus, $t(19) = 6.12$, p

< .001, peak: -42, -57, -12, whereas no significant overlap was detected in the

hippocampus (ps > .778). We further directly compared the mean beta-estimates for

the subsequent success and subsequent precision effects in each anatomical ROI to

assess evidence for disproportionate involvement in one, or the other, aspect of later

retrieval performance. A within subjects ANOVA with the factors of region (HC, IFG,

FFG) and memory measure (success vs. precision) did not provide evidence for a

significant effect of memory measure, $F(1, 19) = 0.05$, $p = .825$, or a significant

interaction between memory measure and brain region, $F(2, 38) = 0.58$, $p = .568$,

however. Indeed, although no conjoint encoding activity relating to subsequent

retrieval success and precision was detected in the hippocampus as described

above, the mean subsequent success and subsequent precision effects did not

significantly differ in the hippocampus (HC: $p = .529$), or the other regions of interest

(IFG: $p = .808$, FFG: $p = .742$).

Encoding activity predicting subsequent retrieval success and memory precision
across the whole brain

To identify any additional brain regions, beyond our *a priori* ROIs, where encoding

activity predicted the later success and/or precision of memory retrieval, we further

performed complementary whole brain analyses. While for subsequent memory

precision, no significant clusters survived a whole brain corrected significance

threshold ($p < .050$ FWE-corrected, $k > 5$), activity in several regions of the dorsal

and ventral visual stream, including the middle occipital gyrus, inferior parietal gyrus,

fusiform gyrus, and inferior temporal gyrus, was found to predict which object

features were later successfully remembered vs. forgotten (see Table 1 and Figure 4A). Whole brain conjunction analyses further indicated significant overlap between subsequent success and subsequent precision effects in the middle occipital and fusiform gyri (see Table 1 and Figure 4C).

Table 1. *Encoding activity associated with the subsequent success and precision of memory retrieval in the whole brain analyses, p < .050 FWE, k > 5.*

Region	Voxels	x	y	z	t	p
Subsequent retrieval success						
L middle occipital gyrus	355	-36	-87	12	10.50	< .001
L inferior parietal gyrus	87	-33	-45	39	9.41	< .001
R middle occipital gyrus	63	33	-75	30	8.83	.001
R inferior temporal gyrus	15	45	-54	-12	8.27	.003
R middle occipital gyrus	21	42	-81	9	7.95	.005
R fusiform gyrus	8	30	-27	-21	7.33	.016
Subsequent precision						
No significant clusters.						
Conjoint activity						
L middle occipital gyrus	30	-24	-75	30	6.59	.002
R middle occipital gyrus	41	33	-78	24	6.39	.004
L fusiform gyrus	30	-42	-57	-12	6.12	.008
L middle occipital gyrus	22	-39	-84	6	6.01	.010

Note. L = left, R = right.

Figure 4. Encoding activity predicting the A) subsequent success and B) subsequent precision of memory retrieval, and C) overlap between encoding activity predicting both later success and precision of memory retrieval. Visualized at an uncorrected threshold of *p* < .001, minimum cluster of 10 voxels.

Discussion

Amid growing interest in the neural substrates underlying the precision of episodic memory, prior studies have predominantly focused on *retrieval* processes (Cooper & Ritchey, 2019; Montchal et al., 2019; Richter et al., 2016; Stevenson et al., 2018), leaving the specific *encoding* mechanisms supporting the acquisition of high-fidelity memories largely uncharacterised. Here, we employed continuous measures of memory retrieval in combination with model-based analyses of fMRI data to

segregate the encoding activity supporting the later success and precision of episodic retrieval. We observed encoding activity in overlapping cortical regions, including the inferior frontal, fusiform and middle occipital gyri, to predict both which object features were later successfully retrieved from memory versus forgotten, and the precision with which they were reconstructed. In contrast, encoding activity in the hippocampus significantly predicted the precision of later memory retrieval only. Together, these findings highlight a hippocampal-cortical basis for the formation of high-fidelity memories of perceptual information, and provide novel insight into the encoding substrates supporting the accessibility and precision of episodic memory.

The current findings demonstrating a relationship between hippocampal encoding activity and later memory precision are consistent with previous accounts emphasizing a critical role for this region in supporting detailed episodic memories (Moscovitch et al., 2016; Robin & Moscovitch, 2017). Prior neuroimaging studies have found hippocampal encoding activity to correlate with both behavioural and neural indices of later retrieval quality, including the number of event details remembered (Cooper & Ritchey, 2020; Qin et al., 2011; Staresina & Davachi, 2008), and the specificity of subsequent neural reinstatement of mnemonic content (Danker et al., 2016; Wing et al., 2015). Aligning with these accounts, we here provide evidence to demonstrate a relationship between hippocampal encoding activity and subsequent behavioural memory precision. Specifically, we observed linear increases in BOLD signal in the left hippocampus to predict the graded precision with which object features were later reconstructed from memory. Interestingly, the peak of this subsequent precision effect was located in the posterior hippocampus, consistent with emerging evidence suggesting posterior hippocampus to be

particularly important for supporting the fine-grained representation of perceptual details (Brunec et al., 2018; Poppenk et al., 2013).

In contrast, we did not observe significant subsequent retrieval success effects in the hippocampus in our current paradigm, nor any significant overlap in this region between encoding activity associated with the subsequent success and subsequent precision of episodic recollection. The lack of significant retrieval success effects in the hippocampus may seem surprising given previous evidence for hippocampal encoding increases for successful versus unsuccessful encoding of associative information (Davachi, 2006; Staresina & Davachi, 2008), but aligns with the proposed involvement of the hippocampus in supporting complex, high-resolution bindings between separate event features (Yonelinas, 2013). Specifically, hippocampal involvement is postulated to increase with greater demands for representation of both high-dimensional and high-fidelity associations (Yonelinas, 2013), a proposal consistent with our current finding of greater hippocampal encoding activity with higher precision of object-feature bindings. Our findings are further in line with patient evidence demonstrating medial temporal lesions to impair both short and long term memory for high-fidelity, but not coarse-grained, bindings (Koen et al., 2017; Nilakantan et al., 2018), and suggest a potential role for a deficient hippocampal encoding mechanism in such impairments. However, it is important to note that despite a lack of significant subsequent retrieval success effects in the hippocampus, we did not observe evidence for significantly greater involvement of hippocampal encoding activity in subsequent memory precision when compared to subsequent retrieval success either.

Interestingly, a prior study employing a similar paradigm to the one used here found hippocampal *retrieval* activity to be associated with the success, but not precision, of

episodic memory retrieval (Richter et al., 2016). This apparent difference between encoding and retrieval effects in the hippocampus may reflect differential involvement of hippocampal *pattern separation* and *pattern completion* during encoding and retrieval, respectively. Specifically, at encoding, hippocampal pattern separation is thought to facilitate the formation of distinct memory traces (Norman & O'Reilly, 2003; Yassa & Stark, 2011), thus posing a putative prerequisite for later high-fidelity memory retrieval (Moscovitch et al., 2016; Xie et al., 2020). In contrast, at retrieval, hippocampal *pattern completion* enables access to stored memory representations when presented with a noisy or partial cue (Norman, 2010; Norman & O'Reilly, 2003), while retrieval precision may be more reliant on cortical regions including the lateral parietal cortex (Richter et al., 2016; Ritchey & Cooper, 2020). Pattern separation and pattern completion are thought to be supported by different subfields of the hippocampus (Hunsaker & Kesner, 2013). An interesting question for future studies to explore is whether different hippocampal subfields also support the precision and accessibility of episodic memory during encoding and retrieval, respectively.

Beyond the hippocampus, we observed activity in overlapping cortical regions, including the inferior frontal, fusiform, and middle occipital gyrus, to predict both the later success and precision of episodic memory retrieval. Our finding of left inferior frontal involvement in subsequent retrieval success and precision is consistent with previous evidence implicating this region in cognitive control of memory encoding, supporting successful memory formation across a range of encoding tasks and mnemonic content (Blumenfeld et al., 2011; Blumenfeld & Ranganath, 2006; Murray & Ranganath, 2007; Park & Rugg, 2008, 2011). Specifically, ventrolateral regions of the prefrontal cortex have been proposed to support the attentional selection and

elaboration of goal-relevant information, leading to formation of strong and distinctive memory traces for specific item features (Blumenfeld et al., 2014; Blumenfeld & Ranganath, 2007; Simons & Spiers, 2003). Such selective encoding processes supported by this region may act to enhance the representation of goal-relevant features in posterior perceptual regions (Chun & Turk-Browne, 2007; Gilbert & Li, 2013; Sprague et al., 2015; Xue et al., 2013), and/or modulate hippocampal encoding more directly (Aly & Turk-Browne, 2017; Carr et al., 2013), aiding the formation of durable and precise memory representations.

The current results further emphasize the role of perceptual regions in supporting the formation of accessible and precise memory traces. Specifically, we observed encoding activity in the fusiform gyrus, a region typically associated with object perception and memory (Bar et al., 2001; Haxby et al., 2001; Vaidya et al., 2002), to predict both the later success and precision of object feature retrieval. This finding is consistent with previous studies that have observed memory-related activity increases in the fusiform gyrus during episodic encoding (reviewed in Kim, 2011; Spaniol et al., 2009), potentially playing an important role in formation of detailed object representations (Garoff et al., 2005; Kensinger et al., 2007), and with evidence suggesting representational specificity in the occipitotemporal cortex during encoding to predict subsequent memory performance (Gordon et al., 2014; Ward et al., 2013; Xue et al., 2010). Beyond our regions of interest, we further observed that encoding activity in a wider network of ventral and dorsal visual regions predicted the subsequent success of memory retrieval. Of these regions, conjoint subsequent retrieval success and precision effects were observed in the middle occipital gyrus. The involvement of a broader set of ventral and dorsal visual regions, implicated in processing of "what" and "where" visual information, respectively (Haxby et al., 2001;

Mishkin et al., 1983), may reflect the demands of the current task for processing both item and spatial stimulus features, such as object colour and location, respectively. This interpretation is consistent with the idea that activity in content-specific perceptual regions supports successful memory formation for distinct event attributes (Cooper & Ritchey, 2020; Gottlieb et al., 2010).

At the behavioural level, we observed the probability and precision of successful memory retrieval to correlate moderately across participants. Although both behavioural (Harlow & Donaldson, 2013; Harlow & Yonelinas, 2016, but see Schurgin et al., 2020) and neural (Richter et al., 2016) evidence suggests partial independence, these two aspects of episodic retrieval are also typically characterised by a degree of shared variance (Richter et al., 2016). Our current findings highlight common cortical encoding substrates as one possible factor contributing to this shared variance. Notably, the representation of stimulus-specific information in posterior perceptual regions, and strategic encoding operations mediated by the lateral prefrontal cortex, may be important for the formation of durable memory traces that are readily accessible from memory and can be reconstructed with a high degree of precision. The computational properties of the hippocampus, on the other hand, may render this structure important for the acquisition of differentiated, high-fidelity, mnemonic representations (LaRocque et al., 2013; Norman & O'Reilly, 2003; Wing et al., 2020).

In summary, the current study aimed to elucidate the encoding mechanisms supporting the formation of accessible and precise memory traces. We observed encoding activity in prefrontal and posterior perceptual regions to support both the later success and precision of episodic memory retrieval. In contrast, activity in the hippocampus was found to significantly predict later memory precision only. These

results highlight the importance of hippocampal-cortical encoding mechanisms for the formation of precise episodic memories, and have important implications for understanding memory deficits in distinct populations.

References

Aly, M., & Turk-Browne, N. B. (2017). How hippocampal memory shapes, and is shaped by, attention. In *The hippocampus from cells to systems: structure, connectivity, and functional contributions to memory and flexible cognition*. https://doi.org/10.1007/978-3-319-50406-3_12

Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *NeuroImage*, *38*(1), 95–113. https://doi.org/10.1016/j.neuroimage.2007.07.007

Bar, M., Tootell, R. B. H., Schacter, D. L., Greve, D. N., Fischl, B., Mendola, J. D., … Dale, A. M. (2001). Cortical mechanisms specific to explicit visual object recognition. *Neuron*, *29*(2), 529–535. https://doi.org/10.1016/S0896-6273(01)00224-0

Bays, P. M., Catalao, R. F. G., & Husain, M. (2009). The precision of visual working memory is set by allocation of a shared resource. *Journal of Vision*, *9*(10), 7–7. https://doi.org/10.1167/9.10.7

Blumenfeld, R. S., Lee, T. G., & D'Esposito, M. (2014). The effects of lateral prefrontal transcranial magnetic stimulation on item memory encoding. *Neuropsychologia*, *53*, 197–202. https://doi.org/10.1016/j.neuropsychologia.2013.11.021

Blumenfeld, R. S., Parks, C. M., Yonelinas, A. P., & Ranganath, C. (2011). Putting

the pieces together: the role of dorsolateral prefrontal cortex in relational memory encoding. *Journal of Cognitive Neuroscience*, *23*(1), 257–265. https://doi.org/10.1162/jocn.2010.21459

Blumenfeld, R. S., & Ranganath, C. (2006). Dorsolateral prefrontal cortex promotes long-term memory formation through its role in working memory organization. *The Journal of Neuroscience*, *26*(3), 916–925. https://doi.org/10.1523/JNEUROSCI.2353-05.2006

Blumenfeld, R. S., & Ranganath, C. (2007). Prefrontal cortex and long-term memory encoding: an integrative review of findings from neuropsychology and neuroimaging. *The Neuroscientist*, *13*(3), 280–291. https://doi.org/10.1177/1073858407299290

Brady, T. F., Konkle, T., Gill, J., Oliva, A., & Alvarez, G. A. (2013). Visual long-term memory has the same limit on fidelity as visual working memory. *Psychological Science*, *24*(6), 981–990. https://doi.org/10.1177/0956797612465439

Brunec, I. K., Bellana, B., Ozubko, J. D., Man, V., Robin, J., Liu, Z.-X., … Moscovitch, M. (2018). Multiple scales of representation along the hippocampal anteroposterior axis in humans. *Current Biology*, *28*(13), 2129-2135.e6. https://doi.org/10.1016/j.cub.2018.05.016

Carr, V. A., Engel, S. A., & Knowlton, B. J. (2013). Top-down modulation of hippocampal encoding activity as measured by high-resolution functional MRI. *Neuropsychologia*, *51*(10), 1829–1837. https://doi.org/10.1016/j.neuropsychologia.2013.06.026

Chun, M. M., & Turk-Browne, N. B. (2007). Interactions between attention and memory. *Current Opinion in Neurobiology*, *17*(2), 177–184.

https://doi.org/10.1016/j.conb.2007.03.005

Cooper, R. A., Richter, F. R., Bays, P. M., Plaisted-Grant, K. C., Baron-Cohen, S., & Simons, J. S. (2017). Reduced hippocampal functional connectivity during episodic memory retrieval in autism. *Cerebral Cortex*, *27*(2), 888–902. https://doi.org/10.1093/cercor/bhw417

Cooper, R. A., & Ritchey, M. (2019). Cortico-hippocampal network connections support the multidimensional quality of episodic memory. *ELife*, *8*, e45591. https://doi.org/10.7554/eLife.45591

Cooper, R. A., & Ritchey, M. (2020). Progression from feature-specific brain activity to hippocampal binding during episodic encoding. *Journal of Neuroscience*, *40*(8), 1701–1709. https://doi.org/10.1523/JNEUROSCI.1971-19.2019

Danker, J. F., Tompary, A., & Davachi, L. (2016). Trial-by-trial hippocampal encoding activation predicts the fidelity of cortical reinstatement during subsequent retrieval. *Cerebral Cortex*, *27*(7), 3515–3524. https://doi.org/10.1093/cercor/bhw146

Davachi, L. (2006). Item, context and relational episodic encoding in humans. *Current Opinion in Neurobiology*, *16*(6), 693–700. https://doi.org/10.1016/j.conb.2006.10.012

Emrich, S. M., Riggall, A. C., LaRocque, J. J., & Postle, B. R. (2013). Distributed patterns of activity in sensory cortex reflect the precision of multiple items maintained in visual short-term memory. *Journal of Neuroscience*, *33*(15), 6516–6523. https://doi.org/10.1523/JNEUROSCI.5732-12.2013

Fletcher, P., Stephenson, C., Carpenter, T., Donovan, T., & Bullmore, E. (2003).

Regional brain activations predicting subsequent memory success: an event-related fMRI study of the influence of encoding tasks. *Cortex*, *39*(4–5), 1009–1026. https://doi.org/10.1016/S0010-9452(08)70875-X

Garoff, R. J., Slotnick, S. D., & Schacter, D. L. (2005). The neural origins of specific and general memory: the role of the fusiform cortex. *Neuropsychologia*, *43*(6), 847–859. https://doi.org/10.1016/j.neuropsychologia.2004.09.014

Gilbert, C. D., & Li, W. (2013). Top-down influences on visual processing. *Nature Reviews Neuroscience*, *14*(5), 350–363. https://doi.org/10.1038/nrn3476

Gordon, A. M., Rissman, J., Kiani, R., & Wagner, A. D. (2014). Cortical reinstatement mediates the relationship between content-specific encoding activity and subsequent recollection decisions. *Cerebral Cortex*, *24*(12), 3350–3364. https://doi.org/10.1093/cercor/bht194

Gottlieb, L. J., Uncapher, M. R., & Rugg, M. D. (2010). Dissociation of the neural correlates of visual and auditory contextual encoding. *Neuropsychologia*, *48*(1), 137–144. https://doi.org/10.1016/j.neuropsychologia.2009.08.019

Harlow, I. M., & Donaldson, D. I. (2013). Source accuracy data reveal the thresholded nature of human episodic memory. *Psychonomic Bulletin & Review*, *20*(2), 318–325. https://doi.org/10.3758/s13423-012-0340-9

Harlow, I. M., & Yonelinas, A. P. (2016). Distinguishing between the success and precision of recollection. *Memory*, *24*(1), 114–127. https://doi.org/10.1080/09658211.2014.988162

Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001). Distributed and overlapping representations of faces and objects in

ventral temporal cortex. *Science*, *293*(5539), 2425–2430.

https://doi.org/10.1126/science.1063736

Hunsaker, M. R., & Kesner, R. P. (2013). The operation of pattern separation and

pattern completion processes associated with different attributes or domains of

memory. *Neuroscience & Biobehavioral Reviews*, *37*(1), 36–58.

https://doi.org/10.1016/j.neubiorev.2012.09.014

Kensinger, E. A., Addis, D. R., & Atapattu, R. K. (2011). Amygdala activity at

encoding corresponds with memory vividness and with memory for select

episodic details. *Neuropsychologia*, *49*(4), 663–673.

https://doi.org/10.1016/j.neuropsychologia.2011.01.017

Kensinger, E. A., Garoff-Eaton, R. J., & Schacter, D. L. (2007). How negative

emotion enhances the visual specificity of a memory. *Journal of Cognitive*

Neuroscience, *19*(11), 1872–1887. https://doi.org/10.1162/jocn.2007.19.11.1872

Kim, H. (2011). Neural activity that predicts subsequent memory and forgetting: A

meta-analysis of 74 fMRI studies. *NeuroImage*, *54*(3), 2446–2461.

https://doi.org/10.1016/j.neuroimage.2010.09.045

Koen, J. D., Borders, A. A., Petzold, M. T., & Yonelinas, A. P. (2017). Visual short-

term memory for high resolution associations is impaired in patients with medial

temporal lobe damage. *Hippocampus*, *27*(2), 184–193.

https://doi.org/10.1002/hipo.22682

Korkki, S. M., Richter, F. R., Jeyarathnarajah, P., & Simons, J. S. (2020). Healthy

ageing reduces the precision of episodic memory retrieval. *Psychology and*

Aging, *35*(1), 124–142. https://doi.org/10.1037/pag0000432

LaRocque, K. F., Smith, M. E., Carr, V. A., Witthoft, N., Grill-Spector, K., & Wagner, A. D. (2013). Global similarity and pattern separation in the human medial temporal lobe predict subsequent memory. *Journal of Neuroscience*, *33*(13), 5466–5474. https://doi.org/10.1523/JNEUROSCI.4293-12.2013

Mishkin, M., Ungerleider, L. G., & Macko, K. A. (1983). Object vision and spatial vision: two cortical pathways. *Trends in Neurosciences*, *6*, 414–417. https://doi.org/10.1016/0166-2236(83)90190-X

Montchal, M. E., Reagh, Z. M., & Yassa, M. A. (2019). Precise temporal memories are supported by the lateral entorhinal cortex in humans. *Nature Neuroscience*, *22*(2), 284–288. https://doi.org/10.1038/s41593-018-0303-1

Moscovitch, M., Cabeza, R., Winocur, G., & Nadel, L. (2016). Episodic memory and beyond: the hippocampus and neocortex in transformation. *Annual Review of Psychology*, *67*(1), 105–134. https://doi.org/10.1146/annurev-psych-113011-143733

Murray, L. J., & Ranganath, C. (2007). The dorsolateral prefrontal cortex contributes to successful relational memory encoding. *Journal of Neuroscience*, *27*(20), 5515–5522. https://doi.org/10.1523/JNEUROSCI.0406-07.2007

Nichols, T., Brett, M., Andersson, J., Wager, T., & Poline, J.-B. (2005). Valid conjunction inference with the minimum statistic. *NeuroImage*, *25*(3), 653–660. https://doi.org/10.1016/j.neuroimage.2004.12.005

Nilakantan, A. S., Bridge, D. J., Gagnon, E. P., VanHaerents, S. A., & Voss, J. L. (2017). Stimulation of the posterior cortical-hippocampal network enhances precision of memory recollection. *Current Biology*, *27*(3), 465–470. https://doi.org/10.1016/j.cub.2016.12.042

Nilakantan, A. S., Bridge, D. J., VanHaerents, S., & Voss, J. L. (2018). Distinguishing the precision of spatial recollection from its success: Evidence from healthy aging and unilateral mesial temporal lobe resection. *Neuropsychologia*, *119*, 101–106. https://doi.org/10.1016/j.neuropsychologia.2018.07.035

Norman, K. A. (2010). How hippocampus and cortex contribute to recognition memory: Revisiting the complementary learning systems model. *Hippocampus*, *20*(11), 1217–1227. https://doi.org/10.1002/hipo.20855

Norman, K. A., & O'Reilly, R. C. (2003). Modeling hippocampal and neocortical contributions to recognition memory: A complementary-learning-systems approach. *Psychological Review*, *110*(4), 611–646. https://doi.org/10.1037/0033-295X.110.4.611

O'Reilly, R. C., & McClelland, J. L. (1994). Hippocampal conjunctive encoding, storage, and recall: Avoiding a trade-off. *Hippocampus*, *4*(6), 661–682. https://doi.org/10.1002/hipo.450040605

Otten, L. J., Henson, R. N., & Rugg, M. D. (2001). Depth of processing effects on neural correlates of memory encoding: Relationship between findings from across- and within-task comparisons. *Brain*, *124*(2), 399–412. https://doi.org/10.1093/brain/124.2.399

Paller, K. A., & Wagner, A. D. (2002). Observing the transformation of experience into memory. *Trends in Cognitive Sciences*, *6*(2), 93–102. https://doi.org/10.1016/S1364-6613(00)01845-3

Park, H., & Rugg, M. D. (2008). Neural correlates of successful encoding of semantically and phonologically mediated inter-item associations. *NeuroImage*, *43*(1), 165–172. https://doi.org/10.1016/j.neuroimage.2008.06.044

Park, H., & Rugg, M. D. (2011). Neural correlates of encoding within- and across-domain inter-item associations. *Journal of Cognitive Neuroscience*, *23*(9), 2533–2543. https://doi.org/10.1162/jocn.2011.21611

Poppenk, J., Evensmoen, H. R., Moscovitch, M., & Nadel, L. (2013). Long-axis specialization of the human hippocampus. *Trends in Cognitive Sciences*, *17*(5), 230–240. https://doi.org/10.1016/j.tics.2013.03.005

Prince, S. E., Dennis, N. A., & Cabeza, R. (2009). Encoding and retrieving faces and places: Distinguishing process- and stimulus-specific differences in brain activity. *Neuropsychologia*, *47*(11), 2282–2289. https://doi.org/10.1016/j.neuropsychologia.2009.01.021

Qin, S., van Marle, H. J. F., Hermans, E. J., & Fernandez, G. (2011). Subjective sense of memory strength and the objective amount of information accurately remembered are related to distinct neural correlates at encoding. *Journal of Neuroscience*, *31*(24), 8920–8927. https://doi.org/10.1523/JNEUROSCI.2587-10.2011

Ranganath, C. (2010). A unified framework for the functional organization of the medial temporal lobes and the phenomenology of episodic memory. *Hippocampus*, *20*(11), 1263–1290. https://doi.org/10.1002/hipo.20852

Ranganath, C., Yonelinas, A. P., Cohen, M. X., Dy, C. J., Tom, S. M., & D'Esposito, M. (2004). Dissociable correlates of recollection and familiarity within the medial temporal lobes. *Neuropsychologia*, *42*(1), 2–13. https://doi.org/10.1016/j.neuropsychologia.2003.07.006

Richter, F. R., Cooper, R. A., Bays, P. M., & Simons, J. S. (2016). Distinct neural mechanisms underlie the success, precision, and vividness of episodic memory.

ELife, *5*, e18260. https://doi.org/10.7554/eLife.18260

Ritchey, M., & Cooper, R. A. (2020). Deconstructing the posterior medial episodic

network. *Trends in Cognitive Sciences*, *24*(6), 451–465.

https://doi.org/10.1016/j.tics.2020.03.006

Robin, J., & Moscovitch, M. (2017). Details, gist and schema: hippocampal–

neocortical interactions underlying recent and remote episodic and spatial

memory. *Current Opinion in Behavioral Sciences*, *17*, 114–123.

https://doi.org/10.1016/j.cobeha.2017.07.016

Schurgin, M. W., Wixted, J. T., & Brady, T. F. (2020). Psychophysical scaling reveals

a unified theory of visual memory strength. *Nature Human Behaviour*, 1–17.

https://doi.org/10.1038/s41562-020-00938-0

Simons, J. S., & Spiers, H. J. (2003). Prefrontal and medial temporal lobe

interactions in long-term memory. *Nature Reviews Neuroscience*, *4*(8), 637–

648. https://doi.org/10.1038/nrn1178

Spaniol, J., Davidson, P. S. R., Kim, A. S. N., Han, H., Moscovitch, M., & Grady, C.

L. (2009). Event-related fMRI studies of episodic encoding and retrieval: Meta-

analyses using activation likelihood estimation. *Neuropsychologia*, *47*(8–9),

1765–1779. https://doi.org/10.1016/j.neuropsychologia.2009.02.028

Sprague, T. C., Saproo, S., & Serences, J. T. (2015). Visual attention mitigates

information loss in small- and large-scale neural codes. *Trends in Cognitive

Sciences*, *19*(4), 215–226. https://doi.org/10.1016/j.tics.2015.02.005

Staresina, B. P., & Davachi, L. (2006). Differential encoding mechanisms for

subsequent associative recognition and free recall. *Journal of Neuroscience*,

26(36), 9162–9172. https://doi.org/10.1523/JNEUROSCI.2877-06.2006

Staresina, B. P., Duncan, K. D., & Davachi, L. (2011). Perirhinal and parahippocampal cortices differentially contribute to later recollection of object- and scene-related event details. *Journal of Neuroscience*, *31*(24), 8739–8747. https://doi.org/10.1523/JNEUROSCI.4978-10.2011

Staresina, Bernhard P., & Davachi, L. (2008). Selective and shared contributions of the hippocampus and perirhinal cortex to episodic item and associative encoding. *Journal of Cognitive Neuroscience*, *20*(8), 1478–1489. https://doi.org/10.1162/jocn.2008.20104

Stevenson, R. F., Zheng, J., Mnatsakanyan, L., Vadera, S., Knight, R. T., Lin, J. J., & Yassa, M. A. (2018). Hippocampal CA1 gamma power predicts the precision of spatial memory judgments. *Proceedings of the National Academy of Sciences*, *115*(40), 10148–10153. https://doi.org/10.1073/pnas.1805724115

Uncapher, M. R., Otten, L. J., & Rugg, M. D. (2006). Episodic encoding is more than the sum of its parts: an fMRI investigation of multifeatural contextual encoding. *Neuron*, *52*(3), 547–556. https://doi.org/10.1016/j.neuron.2006.08.011

Vaidya, C. J., Zhao, M., Desmond, J. E., & Gabrieli, J. D. E. (2002). Evidence for cortical encoding specificity in episodic memory: memory-induced re-activation of picture processing areas. *Neuropsychologia*, *40*(12), 2136–2143. https://doi.org/10.1016/S0028-3932(02)00053-2

Ward, E. J., Chun, M. M., & Kuhl, B. A. (2013). Repetition suppression and multi-voxel pattern similarity differentially track implicit and explicit visual memory. *Journal of Neuroscience*, *33*(37), 14749–14757. https://doi.org/10.1523/JNEUROSCI.4889-12.2013

Wing, E. A., Geib, B. R., Wang, W.-C., Monge, Z., Davis, S. W., & Cabeza, R.

    (2020). Cortical overlap and cortical-hippocampal interactions predict

    subsequent true and false memory. *The Journal of Neuroscience*, *40*(9), 1920–

    1930. https://doi.org/10.1523/JNEUROSCI.1766-19.2020

Wing, E. A., Ritchey, M., & Cabeza, R. (2015). Reinstatement of individual past

    events revealed by the similarity of distributed activation patterns during

    encoding and retrieval. *Journal of Cognitive Neuroscience*, *27*(4), 679–691.

    https://doi.org/10.1162/jocn_a_00740

Xie, W., Park, H.-B., Zaghloul, K. A., & Zhang, W. (2020). Correlated individual

    differences in the estimated precision of working memory and long-term

    memory: commentary on the study by Biderman, Luria, Teodorescu, Hajaj, and

    Goshen-Gottstein (2019). *Psychological Science*, *31*(3), 345–348.

    https://doi.org/10.1177/0956797620903718

Xue, G., Dong, Q., Chen, C., Lu, Z.-L., Mumford, J. A., & Poldrack, R. A. (2013).

    Complementary role of frontoparietal activity and cortical pattern similarity in

    successful episodic memory encoding. *Cerebral Cortex*, *23*(7), 1562–1571.

    https://doi.org/10.1093/cercor/bhs143

Xue, G., Dong, Q., Chen, C., Lu, Z., Mumford, J. A., & Poldrack, R. A. (2010).

    Greater neural pattern similarity across repetitions is associated with better

    memory. *Science*, *330*(6000), 97–101. https://doi.org/10.1126/science.1193125

Yassa, M. A., & Stark, C. E. L. (2011). Pattern separation in the hippocampus.

    Trends in Neurosciences, *34*(10), 515–525.

    https://doi.org/10.1016/j.tins.2011.06.006

Yonelinas, A. P. (2013). The hippocampus supports high-resolution binding in the

service of perception, working memory and long-term memory. *Behavioural Brain Research*, *254*, 34–44. https://doi.org/10.1016/j.bbr.2013.05.030

Zhang, W., & Luck, S. J. (2008). Discrete fixed-resolution representations in visual working memory. *Nature*, *453*(7192), 233–235. https://doi.org/10.1038/nature06860