STOMAGEN-like Regulates Stomatal Generation Causing Increased Water Use

Efficiency during C[4] Evolution

Yong-Yao Zhao[1,2], Mingju Amy Lyu, FenFen Miao, Genyun Chen, Xin-Guang Zhu[1,2]

1. Center of Excellence for Molecular Plant Sciences, Chinese Academy of Sciences

2. State Key Laboratory for Plant Molecular Genetics

Abstract

Compared with C_3 plants, C_4 plants maintain lower stomatal conductance (g_s),

attributed to decreased maximal stomatal conductance (g_{smax}), without losing

photosynthetic CO_2 uptake rate. Which stomatal traits caused the decreased g_{smax} and

how did the g_{smax} change along C_4 evolution as well as the molecular mechanism

underlying this change remain unknown.

Stomatal patternings were examined in *Flaveria* genus, which contains species at

different evolutionary stages from C_3 to C_4 photosynthesis. We further used

comparative transcriptomics, transgenic experiments and semi-*in-vitro* analysis to

identify the gene underlying decreased g_{smax} in C_4 species.

We found that the evolution from C_3 to C_4 species was accompanied by a

stepwise decrease in g_{smax} and stomatal density (*SD*) and the dramatic decrease

occurred between type II and C_4-like species. *SD* controled the decrease of g_{smax}

during C_4 evolution; *FSTOMAGEN* gene positively regulated stomatal development

and its changed expression underlay the decreased *SD* during C_4 evolution;

STOMAGEN-like genes expression also decreased along photosynthetic evolution in

other C_4 lineages.

Our study suggests that the lower *SD* is another major feature of C_4 evolution,

besides C_4 enzymes and Kranz anatomy. *FSTOMAGEN* can be used to engineer *SD*, a

C_4 feature required in the current effort of C_4 crop engineering.

Key words:

Water use efficiency, Stomatal density, C_4 photosynthesis, STOMAGEN, Stomatal

conductance

The total word count for the main body of the text is 6346. The word count of
Introduction, Materials and Methods, Results, Discussion and Acknowledgement is
811, 2293, 2100, 1099 and 43 respectively. This paper includes 8 colour figures. The
supporting material includes 18 figures and 5 tables.

Introduction

Compared with C_3 plants, C_4 plants have higher light use efficiency (LUE) as a
result of a CO_2 concentrating mechanism (*1*). At the same time, C_4 plants have higher
water use efficiency (WUE), as a result of both lower g_s and higher photosynthetic
CO_2 uptake rate (*A*) due to the CO_2 concentrating mechanism (*2, 3*). C_4 photosynthesis
therefore optimizes water and carbon relations by reducing transpiration without
compromising carbon assimilation (*4*). Improvements in WUE enables C_4 plants to
better adapt to saline lands, hot and drought subtropical environments and open
habitats compared with their close C_3 relatives, which raises the possibility that C_4
photosynthesis might have been selected as a water-conserving mechanism (*4*) and
that water limitation might be the primary driver for the initial ecological advantage
of C_4 over C_3 (*5-7*). Given that C_4 plants have higher photosynthetic energy
conversion efficiency and higher potential for biomass productivity, their lower g_s is
critical for the survival and fitness of C_4 plants in the field, especially in a world
where water shortage is predicted to occur more often.

C_3 plants can gain improved WUE through reducing transpiration under
long-term changed environmental conditions by decreasing g_{smax} (*8, 9*), which implies
that the decreasing g_{smax} may be an option that plants use to reduce water loss. Indeed,
phylogenetically informed comparison shows that C_4 plants have a lower g_{smax}
compared with their close C_3 relatives (*10*), indicating that the lower g_{smax} contribute
to the higher water-conserving ability of C_4 land plants (*3, 10*). Therefore, low g_{smax},
same as the C_4 metabolite shuttle and Kranz anatomy, may be a required feature for
C_4 land plants. However, so far, in comparison with the extensive studies on the C_4
metabolite shuttle and Kranz anatomy, relatively little effort has been invested into
studying the changes in g_{smax} and the mechanism underlying these changes during C_4

evolution (*11*).

In principle, g_{smax} positively correlates to operating g_s (*8, 12*) and the g_{smax} is

determined by stomatal density (*SD*) and stomatal size (*SS*). In Eucalyptus and

Arabidopsis, *SD* positively controls g_{smax} and operating g_s (*8, 13-15*). Higher *SD*

appears to have been selected as a strategy to gain higher g_{smax} to cope with low

atmospheric CO_2 levels over the last 400 million years (*16*). In contrast, when grown

under drought, potato plants severely reduce *SS* to lower operating g_s and increase

WUE (*17*). Under ABA treatment, a sufficiently small *SS* can lower g_{smax} to achieve

higher WUE in *Tradescantia virginiana (18)*. Therefore, in different species or under

different environmental conditions, either *SD* or *SS* is used as a primary determinant

for g_{smax} in plants. In the case of C_4 photosynthesis, it also remains unknown whether

SD or SS is the dominant characteristic responsible for the lower g_{smax}. For example,

lower g_{smax} of different species in one C_4 clade were attributed to either smaller *SS* or

lower *SD* (*10*), which may be a result of a compounded phylogenetic effect, i.e.

species used in the study were evolutionarily distant from each other and hence

genetic factors unrelated to photosynthetic type may influenced stomatal

patterning(*3*).

Stomatal development is controlled by a well-studied genetic program (*19, 20*),

which mainly involves a series of ligands and the transmembrane receptors, leading to

downstream changes of mitogen-activated protein kinase (MAPK) and transcription

factors (*19, 20*). TOO MANY MOUTHS (TMM) and ERECTA-family (ERfs: ER,

ERL1, ERL2) are receptor-like kinases that negatively regulate stomatal development

(*21, 22*), and the TMM interacts with ERfs (*21, 23*). The ligands that bind to ERfs and

TMM receptors are mainly EPIDERMAL PATTERNING FACTOR/EPIDERMAL

PATTERNING FACTOR-LIKE family (EPF/EPFL-family) peptides (*24*). The

EPF/*EPFL*-family genes encode cysteine-rich peptides, which can fold into a

three-dimensional structure through disulfide bond (*25*). Overexpression of *EPF1*

decreases *SD*, whose loss of function mutation increases *SD* and results in stomatal

cluster (*26*). *EPF2*, which acts earlier in stomatal development relative to *EPF1*, has a

similar genetic effect as *EPF1* (*27*). STOMAGEN can compete with EPF1 and EPF2

for ERfs (*28*), and then prevents the phosphorylation of downstream receptors, which

results in the changed expression of basic Helix-loop-Helix (bHLH) paralogs, SPCH,

MUTE and FAMA, ultimately positively regulates stomatal development (*29-32*).

To study whether *SD* or *SS* underlies the decreased g_{smax} in plants with C_4

photosynthesis, here we use the *Flaveria* genus, a representative of dicotyledons

which contains closely related species at different evolutionary stages from C_3 to C_4

photosynthesis. Using a genus containing species with intermediate photosynthetic

types enables examination of changes in g_{smax} and stomatal patterning during C_4

evolution. This study aims to answer the following questions: 1) which stomatal

patterning trait, i.e. *SD* and *SS*, underlies the decreased g_{smax} in C_4 evolution and how

did the g_{smax} and stomatal patterning traits change during C_4 evolution? 2) What's the

molecular mechanism underlying the change of the stomatal patterning trait? 3) Is this

molecular mechanism conserved across different C_4 evolutionary events?

Result

Maximum stomatal conductance and stomatal pattern during the evolution of C_4

photosynthesis

As a model of C_4 evolution, there are distinct evolutionary phases in *Flaveria*

genus (Fig 1a). The maximum value of operational g_s (marked with circles on figure)

for C_3 *Flaveria* species under saturated light intensity was higher than that for C_4

species, indicating that the C_3 *Flaveria* species tends to have a higher g_s

(Supplementary Fig 1a). The pore length of stomata was approximately equal to

0.41×stomatal length, and the guard cell width was approximately equal to

0.33×stomatal length (Fig. 1b). Through measuring stomatal density and stomatal

length, we calculated the g_{smax} for *Flaveria* species. The results show that, whether

growing under outdoor natural light or constant light in the greenhouse, the g_{smax}

decreased gradually from C_3 through intermediate species to C_4 species (Fig 1d, e). A

significant correlation was found between the g_{smax} and g_s under saturated light for the

Flaveria species with different photosynthetic types (Fig 1f). Under saturated light,

the g_s of *son* (Type I) was extremely high (marked with circle on figure), matching

with its highest g_{smax} within the *Flaveria* genus (Supplementary Fig 1a, Fig1d).

Similar to that under the saturated light intensity, the WUE (A/g_s) of *rob* (C_3)

measured under unsaturated light intensity was higher than that of *bid* (C_4), and the g_s

of *rob* was 71% higher than that of *bid* (Supplementary Fig. 1c). The observed

reductions in both g_s and g_{smax} of C_4 and C_4-like *Flaveria* species compared to C_3 and

Type I species suggest that the decrease in g_{smax} contributes to the decrease in g_s

during C_4 evolution.

In comparison to C_4 and C_4-like photosynthesis, the stomatal densities (*SD*) of C_3

and Type I photosynthesis on the abaxial leaf surface of *Flaveria* were nearly 150%

higher (Fig. 2a). With the progression from C_3 to C_4 species, there was a distinctly

gradual decrease in *SD* (Fig. 2a, b). The difference in *SD* between C_3 and Type I

species wasn't as large as that between C_3 and C_4 species; and the *SD* of Type II

species fell in between Type I and C_4-like species (Fig. 2b). The *SD* of C_4 and C_4-like

species were similar, although the *SD* of C_4-like species was slightly less than that of

C_4 (Fig. 2b). For the adaxial leaf surface, the *SD* of C_3 species was again higher than

that of the C_4 species, although the difference between *SD* in the intermediate species

was less significant (Supplementary fig. 4a, b). There was a strong positive

correlation of g_{smax} and *SD* either on abaxial leaves ($R^2=0.8955$, p<0.0001) (Fig. 2C)

or on adaxial leaves ($R^2=0.9165$, p<0.0001) (Supplementary fig. 4C) and a significant

negative correlation of g_{smax} and l on abaxial leaves ($R^2=0.2596$, *p*<0.005) (Fig. 2d),

indicating the *SD* determined the change of g_{smax}, not stomatal size (SS). *SD* and l

showed a significant negative correlation on abaxial leaves ($R^2=0.5093$, *p*<0.0001)

(Fig. 2e) or on adaxial leaves ($R^2=0.2055$, *p*<0.05) (Supplementary fig. 4e), the same

as reported previously   (*8, 13, 16*).

Flaveria species, i.e. *rob*, *ram* and *bid*, were chosen to represent C_3, intermediate

and C_4 species, respectively, to analyse the ratio between stomatal number and the

number of epidermal cells, i.e. the stomatal index (*SI*). The *SI* of *rob* was nearly 50%

higher than *bid* and *ram* in plants grown in either greenhouse or outdoors, suggesting

that the proportion of stomata occupying the epidermis, contributed to the highest *SD*

in *rob* (Fig. 2g). In addition to the analysis of *Flaveria* plants grown from cuttings, we

also planted *Flaveria* germinated from seeds. Results show that the *SD* of C_3 species

was higher than C_4 species, consistent with the results from cuttings (Supplementary

Fig. 5).

Comparison of the sequence and transcription abundance of genes related to

stomatal pattern between C_3 and C_4 *Flaveria* species

The signaling pathway for stomatal development has been studied extensively

(*19*). We identified amino acid sequence encoded by homologs of stomatal

development related genes in *Flaveria*. Amino acid sequence alignment analysis

showed that epidermal patterning factor/epidermal patterning factor-Like gene family

(*EPF*/*EPFL*-family) homologs was conserved at the C-terminal part of amino acid

sequence (Fig. 4A, Supplementary fig. 6). Members of *EPF*/*EPFL*-family genes

encode secretory proteins containing signal peptides at the N-terminus and functional

domains at the C-terminus (*26, 27, 33*). Analysis shows that the hydrophobicity of

EPFs/EPFL homologs in *Flaveria* was similar to EPF/EPFL in Arabidopsis

(Supplementary fig 7), implying the homologs are also the secretory proteins and

should have the same physiological function as EPF/EPFL. We hypothesized that

either the amino acid sequence or the expression of stomatal development related

genes changed during C_4 evolution. Analysis results reveals no differences within the

functional region of amino acid sequence for STOMAGEN, EPF1 and EPF2

homologs between C_3 and C_4 species (Fig. 3f, Supplementary Fig. 6a, b), indicating

the change of amino acid sequence is unlikely to be a cause of the changed *SD* in C_4

evolution.

As up-regulation of transcription has been a major approach to increase the

expression of many genes participating in the building of C_4 phenotype during C_3 to

C_4 evolution in *Flaveria* (*34, 35*), we compared the transcript abundance of stomatal

development related genes between C_3 and C_4 *Flaveria* species. The expression

patterns of known C_4 related genes in our data analysis were same as the previous

works, which indicates that our data collection and analysis procedure were reliable

(Supplementary Fig. 12) (*34, 35*). We found that the homologs of most stomatal

development related genes showed higher transcript abundance in developing tissues

as compared to developed tissues (Fig. 3b), which is consistent with their function of

regulating stomatal differentiation in developing leaf (*26, 30, 32, 36*). *STOMAGEN*,

EPF1, *SDD1* and *TMM* showed obviously lower expression levels in C_4 species than

in C_3 species (Fig. 3c). Among these genes, *STOMAGEN* was the one that changed

the most (Fig. 3c) and showed the most significant difference (Fig. 3d). Considering

that the *EPF1*, *SDD1* and *TMM* are negative regulators for stomatal development

(Supplementary table 4) (*24, 26-28*), the decrease in their expression should increase

SD. But actually SD in C_4 species was lower than that in C_3, indicating these genes

are not be the factors responsible for the decrease in *SD* of C_4 species. *STOMAGEN* is

a positive regulator of *SD* (Supplementary table 4) (*24, 30, 31*), whose lower

expression in C_4 species was consistent with the decreased *SD* in C_4 species (Fig. 2a,

b, Supplementary Fig. 4a, b). RT-qPCR quantification of the expression levels of

STOMAGEN, *EPF1*, *EPF2* homologs between C_3 and C_4 species grown under our

conditions also confirmed the results from RNA-SEQ (Fig. 3e, Supplementary Fig.

15). In this study, we renamed *STOMAGEN* in *Flaveria* as *FSTOMAGEN*.

There are considerable differences in amino acid sequence of signal peptide of

STOMAGEN homologs, especially between species with large evolutionary distance

(Fig. 4a). The functional domain of *STOMAGEN* is however conserved, not only

between different *Flaveria* species (Fig. 3f), but also between species that have a large

evolutionary distance (Fig. 4a). In Arabidopsis, increased expression of *STOMAGEN*

increases *SD*; experiments with exogenously applied STOMAGEN peptides shows

that the signal peptide is not required for STOMAGEN to induce increased SD (*30,*

31, 33, 36). A phylogenic tree constructed for *STOMAGEN* from different plants can

be divided into two clades, i.e. dicotyledon (blue clades) and monocotyledon (red

clades) (Fig. 4b). Previous studies showed that overexpressing rice *STOMAGEN*

homologs increases *SD* in Arabidopsis and decreased expression of rice STOMAGEN

homologs reduces *SD* in rice (*37-39*), implying conserved function of *STOMAGEN*

across monocots and dicots. When plants are grown under conditions that promote or

inhibit stomatal development, the expression of *STOMAGEN* is known to increase or

decrease, respectively (*14, 36, 40*). A similar modulation of *STOMAGEN* expression

levels was observed for *rob* when grown at different light conditions (Supplementary

Fig 10).

To confirm that *FSTOMAGEN* can induce stomatal generation, we overexpressed

the *FSTOMAGEN* in *Arabidopsis*. Specifically, we recombined the signal peptide of

STOMAGEN from *Arabidopsis* to the functional region of *FSTOMAGEN* and named

the hybrid gene as AFSTO (Fig, 4c). Overexpressing either the AFSTO or the intact

FSTOMAGEN (FSTO) from *F.rob* into *Arabidopsis* resulted in similar increased *SD*

(Fig. 4e, f, g, j, Supplementary fig 16). Identification of transcription showed

successful transgenes and the expression level of the AFSTO or FSTO increased with

the increase in SD (Fig. 4e, f, h, i). Furthermore, overexpression of *FSTOMAGEN*

also induced clustered stomata (Fig. 4d), as also reported earlier for the

over-expression of *STOMAGEN* (*30*). To further confirm that *FSTOMAGEN* can

increase *SD* in *Flaveria*, we synthesized a peptide representing the FSTOMAGEN

functional region, consisting of 45 amino acids. In order to confirm our synthesized

peptides that have been folded were biologically active, the FSTOMAGEN peptides

were applied in vitro on the leaves of Arabidopsis and resulted in increased *SD* and

cluster stomata (Fig 5b, c). Application of the peptides on the leaves of *bid* also

increased *SD* and *SI* and caused stomatal cluster (Fig 5a, d, e). These results

demonstrate that, in addition to its *in vivo* function, *in vitro* application of

FSTOMAGEN also directly positively regulate *SD* in *Flaveria*. This is consistent with

the notion that *STOMAGEN* is expressed in mesophyll cells and secreted out of cells

to influence the stomatal development in the epidermal layer in *Arabidopsis* (*30, 31*).

The transcript abundance of *FSTOMAGEN* during C$_4$ evolution in *Flaveria*

We examined the copy number of *FSTOMAGEN* in *Flaveria* using blastn with

the coding sequence of *FSTOMAGEN* as query sequence against the assembled

contigs of each *Flaveria* species from 1KP. Results show that there was only one hit

sequence with e-value less than 10^{-5} in each species, suggesting that there is a single

copy of *FSTOMAGEN* in *Flaveria* (Supplementary Fig. 13). Therefore we can ignore

the impact of other *FSTOMAGEN* paralogs on stomatal development, and there will be no quantitative errors in the gene expression levels caused by other paralogs that were not considered. Examination of the RNA-seq data of *Flaveria* showed that the expression levels of *FSTOMAGEN* in C_3 species were markedly higher than those in C_4 species along different developmental stages of leaves (Fig. 6a). During leaf growth and development, the expression of *FSTOMAGEN* in both *bid* and *rob* increased first, reaching the highest level at leaf development stage 3 (leaf length 1.5 cm) or 4 (leaf length 2.0 cm), and then declined (Fig. 6a). The similar patterns of *FSTOMAGEN* expression in these two *Flaveria* species indicates that it would be valid to compare the *FSTOMAGEN* expression levels in leaves at the same development stage between *Flaveria* species. The expression level of *FSTOMAGEN* in *bid* (C_4) was lower than that in *rob* (C_3), and that of *ram* (intermediate) lied between them (Fig. 6b). This encouraged us to test more *Flaveria* species with different photosynthetic types. Similar results were obtained with either RNA-seq or RT-qPCR (Fig. 6c, Supplementary Fig. 14), where we found the gradual decrease of *FSTOMAGEN* expression from C_3 to intermediate species to C_4 species. Together these results suggest that the gradual change in the *FSTOMAGEN* expression levels between different species underlay the stepwise change in *SD* along C_4 evolution.

The expression of *FSTOMAGEN* homologs for species with different photosynthetic type in multiple C_4 lineages.

To gain insight into whether *STOMAGEN* homolog might be a generic factor leading to the difference in *SD* between C_3 and C_4 species in higher plants, we tested its expression level in different C_4 lineages other than the *Flaveria genus*. The *SD* of abaxial leaf in C_3 rice was 240% higher than that in C_4 maize, consistent with a 574% higher g_s in rice compared to maize (Fig. 7a, b). The stomatal length (*l*) in maize was 133% longer (Fig. 7a, b). We analysed the transcript abundance of *STOMAGEN* homologs, since there are two and three paralogs of *STOMAGEN* in rice and maize respectively (*37*), only one in rice and two in maize are expressed. The *STOMAGEN* homologs in rice had much higher transcript abundance than those in maize (Fig. 7d),

while the transcript abundance of other stomatal development related genes did not

differ to the same extent between rice and maize (Fig. 4c). This result encouraged us

to test more C_4 lineages (Fig. 7e). C_4 photosynthesis related genes were indeed higher

in C_4 species, suggesting the data and analysis pipeline can be used to compare the

differences in expression of genes between C_3 and C_4 photosynthetic types (Fig 7f). It

was showed that the expression levels of *STOMAGEN* homologs for C_4 were lower

than those for C_3, while intermediate had expression between them (Fig 7g).

Method

Plant materials, sample condition and data sources

All *Flaveria* species used in this study were kindly provided by Prof. Rowan F.

Sage (University of Toronto). Plants used for stomatal phenotyping were re-grown

from cuttings. Placed in an environment humidified with humidifiers to generate roots.

After plants generated roots, they were transferred outdoors (Institute of Plant

Physiology and Ecology, Chinese Academy of Sciences, Shanghai, China), and

transplanted into 4-L pots filled with topsoil in April. In July, the number of stomata

on abaxial and adaxial surfaces of leaves were counted. Another batch of *Flaveria*,

which were also grown from cuttings, were transferred to the phytotron (Center of

Excellence for Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai,

China). Growth temperature of the phytotron was maintained at 25-28□, photoperiod

was maintained at 16h light / 8h dark, relative humidity (RH) was at 60-65% and

photosynthetic photon flux density (PPFD) was maintained at about 180 μmol $m^{-2} s^{-1}$.

Two months later, the numbers of stomata on the abaxial and adaxial surfaces of

leaves were counted. RNA was extracted from 1.5~3.0 cm long young leaves on the

axial shoot which were still expanding. Three hours before sampling, we provided

sufficient water to ensure that plants were not water limited, since drought and water

status can affect expression levels of *STOMAGEN* and *SD* (*14, 41*). Gene expression

was also examined under normal growth conditions by RT-qPCR for the *rob*, *ram* and

bid, which are representative of C_3, intermediate and C_4 species in the *Flaveria* genus.

These three species were grown in the same phytotron with the same environmental

conditions, except that we did not water plants before sampling. To test whether

plants grown from cuttings or from seeds show difference in the *SD*, we further grew

Flaveria species from seeds in phytotron and examined stomatal properties. Same

environmental conditions were maintained for phytotron as above. Two months after

the germination of seeds, we counted stomatal number.

All plants were well watered to avoid drought stress during their growth. We

supplied the same amount of commercial nutrient solution (Peters Professional,

SCOTTS, N:P:K = 20:20:20+TE, at recommended dose for application) to the

different species to avoid nutrient deficiency.

Rice (*Oryza Sativa Nipponbare*) and maize (*Zea maize* B73) were grown in the

greenhouse (Center of Excellence for Molecular Plant Sciences, Chinese Academy of

Sciences, Shanghai, China) under a PPFD of 550 umol/m^2/sec, a photoperiod of 12h

light / 12h dark, and a daytime temperature of around 31℃ and a night-time

temperature of around 22℃. The Relative humidity was maintained at around 50-55%.

The plants were regularly watered to avoid drought stress.

In addition to the expression data collected by in this study, we also used public

data for comparison of transcript abundance. Considering that stomatal development

related genes are mainly expressed in young tissues, in each of these used public data

source, samples for young leaves were included. For the *Flaveria*, the samples data

from four source containing the leaves of plants grown under normal condition were

analysed. The first data source is the One Thousand Plants (1KP) Consortium, and we

subsequently analyzed the data according to the methods described in previous work

(*42, 43*); The second and third data sources were Mallman et al (2014) (*35*) and Udo

Gowik et al (*34*); The fourth data source was Kumari Billakurthi et al (2018) (*44*).

The last three data sources are all from Heinrich-Heine-University. For the second and

third data source, the second and fourth visible leaves were sampled; the second leaf

was not completely mature, and stomatal developmental related genes were still

actively expressed. In addition to *Flaveria*, transcripts of C$_3$ and C$_4$ species from other

C$_4$ lineage sequenced from 1KP were assembled using Trinity (*45*) with default

parameters. Transcript abundances were analyzed by mapping the reads to assembled

contigs for corresponding samples using the RSEM package with mapping option of

bowtie 1 (*46*), and outliers that less than Q1 (lower quartile) - 1.5IQR (interquartile

range) and greater than Q3 (upper quartile) + 1.5IQR (interquartile range) were

removed. Expression levels for stomatal development related genes in developing

leaves for rice and maize were obtained from Wang et al. (*47*).

Conservation and hydrophobicity analysis of proteins

Amino acid sequence of the homologs of *STOMAGEN*, *EPF1* and *EPF2* were

aligned by clustalx, and graphs for the multiple alignments were drawn with ESPrit

3.0 (*48*) (http://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi). Hydrophobicity of proteins

was determined by ProtScale (*49*) (https://web.expasy.org/protscale/).

Construction of the phylogenetic tree of *STOMAGEN*

In order to construct the gene evolutionary tree of *STOMAGEN*, we searched the

orthologs of *STOMAGEN* in 28 representative species among Viridisplantae as

demonstrated in (*50*),The protein sequences of representative species were

downloaded from Phytozome (https://phytozome.jgi.doe.gov/). Orthologs of

STOMAGEN were predicted applying OrthoFinder (V 0.7.1) (*51*)using default

parameters. Proteins sequences of STOMAGEN orthologs were then aligned using

MUSCLE (*52*). We than inferred the gene phylogeny using both maximum likelihood

(ML) and Bayesian-inference (BI) methods. The ML tree was constructed applying

RAxML (V2.2.3) (*53*), where bootstrap scores were obtained by running 100,000

independent trees. BI tree was built applying MrBayes package (v 3.2.1) (*54*), where

posterior probabilities were calculated by running 2000,000 generations with the first

2000 generations being used as burn-in. In both cases, PROTGAMMAILG model

(General Time Reversible amino acid substitution model with assumption that

variations in sites follow gamma distribution) was used to infer the gene phylogeny.

The protein sequence of AT4G14723 (*CLL2*) from A. thaliana was used as outgroup

as it showed the highest identity with STOMAGEN in A. thaliana, with a protein

identity of 26.97% and E-value of 0.037 based on Blastp (*55*). Protein sequences for

the *Flaveria* species were predicted using OrfPredictor (*56*) based on de novo

assembled transcripts as described in (*43*).

Measurement of stomatal conductance (g_s)

The LI-6400 Portable Photosynthesis System (Li-Cor, Inc., Lincoln, NE, USA)

was used to measure g_s on the youngest fully expanded leaves from approximately

three-months-old *Flaveria* plants. The cuvette temperature was maintained at 25□,

the PPFD was at ~500 µmol/m²s and the CO_2 concentration was at around 400 ppm.

Chamber humidity was matched to ambient air humidity. Leaf-to-air vapor pressure

difference (VPD) was maintained around 1.1 kPa. Before the g_s measurements,

Flaveria plants were maintained under constant light with a PPFD of ~500 µmol/m^{2}s

until g_s reached a steady state. External environments during the g_s measurements for

rice and maize were controlled to be their growth conditions. In maize and rice,

the16-day-old third leaves which were the youngest mature leaves, were used for gas

exchange measurements (*47*).

Phenotyping stomatal traits and the calculation of g_{smax}

The youngest fully expanded leaves (usually the third or fourth leaves from the

apex in *Flaveria*) at the main stem were used to phenotype stomata related traits. In

Flaveria and maize, stomata related traits were observed using both nail polish

method and epidermis directly peeled from leaves (*6, 13*). For the first method, nail

polish was painted on both leaf surfaces to produce the impressions of epidermis.

After drying of the nail polish, the nail polish was peeled off the leaf and placed on a

microscope slide for examination. For the second method, leaf epidermis was peeled

off manually and transferred onto microscope slide. For the count of SD in

Arabidopsis, the rosette leaves was soaked in 9:1 (V/V) ethanol:acetic acid overnight,

washed with water and cleared in chloral hydrate solution (glycerol:chloral

hydrate:water, 1:8:1) overnight, and then stomata of Arabidopsis were stained with

Safranin-O, finally leaves were placed on a microscope slide for imaging. Light

microscope (smarte; CNOPTEC), camera and image processed software (OPTPRO;

CNOPTEC) were used to observe and image epidermis at 100x magnification in

Flaveria. Center of a leaf was chosen to count stomatal density and measure stomatal

length. Five stomata randomly chosen along a diagonal of the varnish peels or

epidermis were measured for stomatal length.

G_{smax} was calculated using the follow equation (*16*):

$$g_{smax} = \frac{d}{v} \cdot D \cdot a_{max} / (e + \frac{\pi\sqrt{a_{max}/\pi}}{2})$$

The constants *d* is the diffusivity of H_2O in air; the constant *v* is the molar

volume of air; D is the stomatal density; a_{max} is the maximum stomatal pore area

which is calculated as $a_{max} = \pi p^2/4$ in *Flaveria* species, and the *p* is the stomatal

pore length; *e* is the stomatal pore depth, which is known as equal to guard cell width.

RNA isolation, semiquantitative RT-PCR and quantitative real-time PCR

The young leaf samples taken from the main stem in *Flaveria* or the developing

rosette leaves in Arabidopsis were immediately frozen in liquid nitrogen and used

directly for RNA extraction. Total RNA was extract from the young leaves using the

PureLink RNA Mini kit (Thermo Fisher Scientifc). The first-strand cDNA was

synthesized by the TransScript One-step gDNA Removal and cDNA Synthesis

SuperMix (TRANSGEN). Semiquantitative RT-PCR was performed with Phanta®

Max Super-Fidelity DNA Polymerase (vazyme). The actin8 was chosen as the loading

control. Real-time PCR was performed on the CFX connect[TM] (Bio-Rad) with the

UNICON[TM] qPCR SYBR ®Green Master Mix (YEASEN) according to the

manufacturer's instructions. The calculation of expression according to the $2^{-\Delta\Delta CT}$.

The expression stability of internal reference genes for the quantitative real-time

PCR was assessed by the software: geNORM and NormFinder (*57, 58*). Specifically,

we first calculated the coefficient of variations (CV) of all genes in different *Flaveria*

species. After that, all genes were sorted according to the calculated CV values. 150

genes with the lowest CV were selected as potential candidate reference genes.

Among these, those with mean expression level >300 were chosen as candidate

reference genes. We also included those commonly used house-keeping genes, which

include *UBC9* (AT4G27960), *UBQ11* (AT4G05050), *ACT7* (AT5G09810), *GAPDH*

(AT1G13440), *EF1a* (AT5G60390), *CACMS* (AT5G46630) (Supplementary fig. 6a, b,

d), as the internal reference genes in this study. The specificity of the primers

(Supplementary table 2) for all these candidate reference genes were identified by

agarose electrophoresis (Supplementary fig. 6c). With Genorm (Supplementary fig. 6e)

and Normfinder (Supplementary fig. 6f), we concluded that *ACT7*, *EF1a* and *HEME2*

showed stable expression levels across *rob*, *ram* and *bid*. *EF1a* and *ACT7* are classic

reference genes, and they showed minor variations under different conditions. The

geometrical means of the expression level of *ACT7* and *EF1a* were finally used as

normalization factors. All primers used for RT-qPCR are listed in Supplementary table

2. For each gene, three technical replicates and at least three biological replicates were

performed.

Vector construction and transformation

A list of primer sequences were used for plasmid construction (Supplementary

table 2). The *STOMAGEN* and *FSTOMAGEN* DNA sequences, named ATSTO-cDNA

and FloSTO-cDNA respectively, were PCR-amplified from cDNA in Arabidopsis and

rob. To obtain the AFSTO fragment, the signal peptide region of *STOMAGEN* was

cloned from ATSTO-cDNA by PCR-amplification, and the functional region of

FSTOMAGEN was PCR-amplified from FloSTO-cDNA. The signal peptide and

functional region were combined by fusion PCR. To obtain an intact *FSTOMAGEN*

(FSTO) gene, the whole gene in *rob*, including the signal peptide, propeptide and the

STOMAGEN functional region, was amplified from the cDNA of *rob*. To over-express

AFSTO and FSTO, the AFSTO and FSTO were cloned separately to a pcambia-3300

binary vector which has a CaMV 35S promoter. This construct was introduced into *A.*

tumefaciens strain GV3101, transformed into Arabidopsis using a floral dipping

procedure. Transgenic lines were selected on soil that had been treated with a diluted

solution of phosphinothricin (diluted 3000 times) over night. About a month after the

seeds were sowed on the soil, the stomata were observed on the biggest mature rosette

leaf. The phenotype of Arabidopsis T1 and T2 generations was observed at the 400x

magnification using the light microscope. For each epidermal peel, five fields were

counted for density.

Refolding of synthetic *FSTOMAGEN*

Chemically synthesized peptide (Synthesized peptide sequence: IGSVKPTCTY

NECRGCRSRCRAEQVPVEGNDPINSAYHYRCICHR, Sangon Biotech, Shanghai,

China) was dissolved in 20 mM Tris-HCl, pH 8.8, and 50 mM NaCl and then dialysed

(Sangon Biotech,0.5 KD-1 KD) for 1 day at 4□ against 0.5 mM glutathione (reduced

and oxidized forms, Macklin) and 200 mML-arginine (meilunbio) at pH 8.0, which

were further dialysed three times against 20 mM Tris-HCl, pH 8.8, and 50 mM NaCl

for 1.5 days to remove glutathione.

Stomata induction assay

When the first true leaf or cotyledons appeared in *bid* and Arabidopsis that had

germinated in 1/2 MS sterilized solid medium, F*STOMAGEN* peptide was applied on

the plants. After the *bid* and Arabidopsis was further incubated in 1/2 MS liquid

medium at 22℃ for 5 days, stomatal number on the abaxial surfaces for the first true

leaf and cotyledons was counted under a differential interference contrast microscope

(DIC)**.**

Plotting and statistical analysis

The correlation, linear regressions and t-test (a=0.05) was performed by

GraphPad Prism 8. Differences between means of multi-groups were assessed with

one-way analysis of variance (ANOVA). Duncan's multiple range test was performed

using the agricolae package of the R (https://www.r-project.org/) to detect differences

between two groups. The A and g_s at the saturated light intensity were digitized from

figure 1 and figure 3 in (*2, 59*) using the OriginPro 2018.

Accession numbers

The *Flaveria* read data have been submitted to the National Center for Biotechnology

Information Short Read Archive under accession numbers: SRP036880, SRP036881,

SRP036883, SRP036884, SRP036885, SRP037526, SRP037527, SRP037528,

SRP037529, SRP006166, SRP169568. The *Flaveria* RNA-Seq datasets from 1KP

Consortium were available at National Center for Biotechnology Information (NCBI)

Gene Expression Omnibus (GEO) under accession number: GSE54339. The rice and

maize RNA-seq datasets under accession number: GSE54274.

Discussion

C_4 plants have higher water use efficiency and light use efficiency, as result of the

combined effect of decreased operating g_s and a CO_2 concentrating mechanism (Fig

1b, c) (*3, 4*). Previous work suggests that the g_{smax} in C_4 plants are lower than their C_3

relatives (*10*). Comparative studies using plants from different C_4 clades show that

lower g_{smax} is due to either a smaller stomatal size (*SS*) for some C_4 clades, or lower

stomatal density (*SD*) for other clades (*10*). We collected and analysed *SD* data from

literature for a large number of C_3 and C_4 species (*10, 60-66*). The average *SD* on both

sides of leaf surface of C_3 species was significantly higher than that of C_4 species

(Supplementary Fig. 1). Recent studies have shown that when the physiological and

ecological differences between C_3 and C_4 plants are compared, the phylogenic effect

must be considered (*3, 67, 68*), *i.e.* comparing either stomatal density or size without

considering the phylogeny inevitably has the compounded effect of other genetic

factors other than photosynthetic types. Here we used the *Flaveria* genus which

contains species with close evolutionary distance and intermediate photosynthetic

types to examine changes in stomatal traits during C_4 evolution. We showed that, in

addition to the grass, the g_{smax} of C_4 plants in dicots was lower than that of C_3 plants,

and there was a stepwise decrease during C_4 evolution indeed (Fig 1). Compared to

the *SD* of C_3 species, the *SD* of C_4 species decreased (Fig. 2a, b). This decrease in *SD*

is due to less stomata generation rather than augmented guard cell and epidermal cells

(Fig. 2g). In two other closely related Cleomaecae species, *Gynandropsis gynandra*

(C_4) also has a lower SD than *Tareneya hassleriana* (C_3), consistent with the

difference in their g_s (*69*).

The lower SD in phylogenetically closely related C_4 species compared to C_3

species raises the possibility that, during evolution from C_3 to C_4 species, the SD may

gradually decrease. This is indeed shown in (Fig. 2a, b) where with the gradual

transition of photosynthetic type from C_3 to type I, to type II, to C_4-like and to C_4,

there is a gradually decrease in SD. Usually stomata conductance is significantly

higher on the abaxial surface, as shown in maize, broad bean, and wild tomato species

(*70-72*). Here we found that there is a highly significant difference in SD on the

abaxial surface between the C_3 species and the C_4 species; by comparison, on the

adaxial surface, though the difference between SD in the C_3 and C_4 species was

smaller, C_4 species still had lower SD (Supplemental Fig 4,b). In contrast, the SS as

represented by stomatal length was longer in intermediate species and C_4 species

compared to C_3 species in *Flaveria*, consistent with previous reports that SD

negatively correlates with SS (*8, 16*), suggesting that the SS is not the crucial

contributor to the decreased g_{smax} and operational g_s in C_4 evolution. These evidences

show that lower SD is the determinant of the commonly observed lower g_{smax} and

correspondingly higher water use efficiency in C_4 compared to C_3 plants (*2, 73*).

Therefore, lower SD could be another major feature of C_4 evolution, besides C_4

enzymes and Kranz anatomy.

Many earlier studies suggest that water limitation might have been a primary

selection pressure for the evolution of C_4 photosynthesis (*5-7*). In *F. angustifolia*, a

type I intermediate *Flaveria* species, when grown under outdoor natural light

conditions, SD, g_{smax} and g_s were still similar to those in two C_3 *Flaveria* species, i.e.

F. cronquistii and *F. pringlei* (*2*) (Fig. 1d, 3b), as also the case in *Helliotropin* (*74*). In

contrast, the CO_2 concentration point of *F. angustifolia* was dramatically decreased

(*75*) compared to their C_3 ancestors, as a result of an operating CO_2 concentrating

mechanism of C_2 (*76*). This suggests that during the evolution from C_3 to intermediate

species, where the movement of GDC from mesophyll cell to bundle sheath cell

occurred, the primary evolutionary driving force should be to elevate CO_2 levels

around Rubisco, rather than H_2O saving (*77*). After this step, reducing water loss

might became crucial. The emergence of C_2 mechanisms can increase leaf

photosynthetic CO_2 uptake rate and biomass production, which can in principle create

a higher water demand and increased the danger of embolism in the intermediate

species, especially in the arid and semi-arid areas (*4*). This will create an even greater

challenge once the complete C_4 photosynthetic metabolism emerged where

photosynthetic CO_2 uptake rate and biomass production are further increased.

Therefore, decreased *SD* and further increased water use efficiency observed in this

study and earlier studies (*2, 59*) should be a required change during the transition

from the intermediate species and later C_4 species.

Since *SD* is controlled by a molecular signaling network composed of many

interacting components, many options can be used to alter *SD*. Out of these different

options to regulate *SD*, here we present results which indicate that a down-regulation

of the expression of *FSTOMAGEN*, a *STOMAGEN*-like gene, has been selected as a

molecular switch to reduce *SD* during C_4 evolution in *Flaveria*, which is also the case

in other C_4 lineages. When synthetic FSTOMAGEN peptide was applied on the leaves

or *FSTOMAGEN* was over-expressed in Arabidopsis, the increases in SD were

observed. Our evidences showed that the *FSTOMAGEN* positively regulates stomatal

development. The result suggests that the declining expression of *STOMAGEN*

homologs underlies the observed decrease in *SD* during the evolution of C_4

photosynthesis. In theory, Any of genes controlling stomatal patterning can be

potentially selected to alter *SD* by regulation of expression during the emergence of

C_4 photosynthesis. And in fact, these genes have innate mechanisms to alter their

expression under different environments, such as low CO_2 (*78*). However,

STOMAGEN homologs were shown to be the regulator selected to reduce *SD*. The

two specialities of *STOMAGEN* probably have accounted for the phenomenon. First,

STOMAGEN can regulates the entire molecular process of stomatal development,

rather than *EPF1*, *EPF2*, *TMM* and the members of *ERECTA,* which individually only

regulates particular phases of the stomatal development (*20, 21, 23, 24, 26, 27*).

Second, which is potentially more important, *STOMAGEN* is expressed in the

mesophyll cell, while all other regulators controlling stomatal development are
expressed in the cell of stomatal lineage (*24, 26, 27, 30, 31*). Having a genetic
regulator controlling stomatal density in the mesophyll cell, where many of the C_4
related genes are expressed, can enable an easier coordination between stomatal
development and C_4 photosynthesis. Future work is still needed to elucidate molecular
level coordinating the expression of *STOMAGEN* and C_4 photosynthetic related genes.

Reference

1. O. V. Sedelnikova, T. E. Hughes, J. A. Langdale, Understanding the Genetic Basis of C_4 Kranz
Anatomy with a View to Engineering C_3 Crops. *Annu Rev Genet* **52**, 249-270 (2018).
2. P. J. Vogan, R. F. Sage, Water-use efficiency and nitrogen-use efficiency of C_3-C_4 intermediate
species of Flaveria Juss. (Asteraceae). *Plant, cell & environment* **34**, 1415-1430 (2011).
3. S. H. H. Taylor, S. P. Rees, M Ripley, B. S. Woodward, F. I. Osborne, C. P., Ecophysiological traits
in C_3 and C_4 grasses: a phylogenetically controlled screening experiment. *New Phytol* **185**,
780-791 (2010).
4. C. P. Osborne, L. Sack, Evolution of C_4 plants: a new hypothesis for an interaction of CO_2 and
water relations mediated by plant hydraulics. *Philos Trans R Soc Lond B Biol Sci* **367**, 583-600
(2012).
5. H. Zhou, B. R. Helliker, M. Huber, A. Dicks, E. Akcay, C_4 photosynthesis and climate through
the lens of optimality. *Proc Natl Acad Sci U S A* **115**, 12057-12062 (2018).
6. G. Reeves *et al.*, Natural Variation within a Species for Traits Underpinning C_4 Photosynthesis.
Plant physiology **177**, 504-512 (2018).
7. B. P. Williams, I. G. Johnston, S. Covshoff, J. M. Hibberd, Phenotypic landscape inference
reveals multiple evolutionary paths to C_4 photosynthesis. *Elife* **2**, e00961 (2013).
8. P. J. Franks, P. L. Drake, D. J. Beerling, Plasticity in maximum stomatal conductance
constrained by negative correlation between stomatal size and density: an analysis using
Eucalyptus globulus. *Plant, cell & environment* **32**, 1737-1748 (2009).
9. T. Doheny-Adams, L. Hunt, P. J. Franks, D. J. Beerling, J. E. Gray, Genetic manipulation of
stomatal density influences stomatal size, plant growth and tolerance to restricted water
supply across a growth carbon dioxide gradient. *Philos Trans R Soc Lond B Biol Sci* **367**,
547-555 (2012).
10. S. H. Taylor *et al.*, Photosynthetic pathway and ecological adaptation explain stomatal trait
diversity amongst grasses. *New Phytol* **193**, 387-396 (2012).
11. D. A. Way, What lies between : the evolution of stomatal traits on the road to C4
photosynthesis. *New phytologist* **193**, 291-293 (2012).
12. G. J. Dow, D. C. Bergmann, J. A. Berry, An integrated model of stomatal development and leaf
physiology. *New Phytol* **201**, 1218-1226 (2014).
13. P. J. Franks, W. Doheny-Adams T, Z. J. Britton-Harper, J. E. Gray, Increasing water-use
efficiency directly through genetic manipulation of stomatal density. *New Phytol* **207**,
188-195 (2015).

14.   E. T. Hamanishi, B. R. Thomas, M. M. Campbell, Drought induces alterations in the stomatal
development program in Populus. *J Exp Bot* **63**, 4959-4971 (2012).

15.   Y. Tanaka, S. S. Sugano, T. Shimada, I. Hara-Nishimura, Enhancement of leaf photosynthetic
capacity through increased stomatal density in Arabidopsis. *New Phytologist* **198**, 757-764
(2013).

16.   P. J. F. a. D. J. Beerling, Maximum leaf conductance driven by CO_2 effects on stomatal size and
density over geologic time. *PNAS* **106**, 10343–10347 (2009).

17.   S. h. Zhang, X. f. Xu, Y. m. Sun, J. l. Zhang, C. z. Li, Influence of drought hardening on the
resistance physiology of potato seedlings under drought stress. *Journal of Integrative*
Agriculture **17**, 336-347 (2018).

18.   Peter J. Franks, G. D. Farquhar, The Effect of Exogenous Abscisic Acid on Stomatal
Development, Stomatal Mechanics, and Leaf Gas Exchange in Tradescantia virginiana. *Plant*
physiology **125**, 935–942 (2001).

19.   L. J. Pillitteri, K. U. Torii, Mechanisms of stomatal development. *Annu Rev Plant Biol* **63**,
591-614 (2012).

20.   N. Zoulias, E. L. Harrison, S. A. Casson, J. E. Gray, Molecular control of stomatal development.
Biochem J **475**, 441-454 (2018).

21.   G. Lin *et al.*, A receptor-like protein acts as a specificity switch for the regulation of stomatal
development. *Genes Dev* **31**, 927-938 (2017).

22.   E. D. Shpak, J. M. McAbee, L. J. Pillitteri, K. U. Torii, Stomatal patterning and differentiation by
synergistic interactions of receptor kinases. *Science* **309**, 290-293 (2005).

23.   J. S. Lee *et al.*, Direct interaction of ligand-receptor pairs specifying stomatal patterning.
Genes Dev **26**, 126-136 (2012).

24.   T. Shimada, S. S. Sugano, I. Hara-Nishimura, Positive and negative peptide signals control
stomatal density. *Cell Mol Life Sci* **68**, 2081-2088 (2011).

25.   A. L. Rychel, K. M. Peterson, K. U. Torii, Plant twitter: ligands under 140 amino acids enforcing
stomatal patterning. *J Plant Res* **123**, 275-280 (2010).

26.   K. Hara, R. Kajita, K. U. Torii, D. C. Bergmann, T. Kakimoto, The secretory peptide gene EPF1
enforces the stomatal one-cell-spacing rule. *Genes Dev* **21**, 1720-1725 (2007).

27.   L. Hunt, J. E. Gray, The signaling peptide EPF2 controls asymmetric cell divisions during
stomatal development. *Curr Biol* **19**, 864-869 (2009).

28.   J. S. Lee *et al.*, Competitive binding of antagonistic peptides fine-tunes stomatal patterning.
Nature **522**, 439-443 (2015).

29.   L. J. Pillitteri, K. U. Torii, Breaking the silence: three bHLH proteins direct cell-fate decisions
during stomatal development. *Bioessays* **29**, 861-870 (2007).

30.   S. S. Sugano *et al.*, Stomagen positively regulates stomatal density in Arabidopsis. *Nature* **463**,
241-244 (2010).

31.   T. Kondo *et al.*, Stomatal density is controlled by a mesophyll-derived signaling molecule.
Plant Cell Physiol **51**, 1-8 (2010).

32.   L. Hunt, K. J. Bailey, J. E. Gray, The signalling peptide EPFL9 is a positive regulator of stomatal
development. *New Phytol* **186**, 609-614 (2010).

33.   S. Ohki, M. Takeuchi, M. Mori, The NMR structure of stomagen reveals the basis of stomatal
density regulation by plant peptide hormones. *Nat Commun* **2**, 512 (2011).

34. U. Gowik, A. Bräutigam, K. L. Weber, A. P. M. Weber, P. Westhoff, Evolution of C_4
Photosynthesis in the Genus Flaveria: How Many and Which Genes Does It Take to Make C_4?
The Plant Cell **23**, 2087-2105 (2011).

35. J. Mallmann *et al.*, The role of photorespiration during the evolution of C_4 photosynthesis in
the genus Flaveria. *Elife* **3**, e02478 (2014).

36. M. Hronkova *et al.*, Light-induced STOMAGEN-mediated stomatal development in
Arabidopsis leaves. *J Exp Bot* **66**, 4621-4630 (2015).

37. J. Lu *et al.*, Homologous genes of epidermal patterning factor regulate stomatal development
in rice. *J Plant Physiol* **234-235**, 18-27 (2019).

38. L. Xuxin, The Effects for Stomata Developmental on Arabidopsis by Overexpression of Rice
EPFL Family Genes. *Molecular Plant Breeding* **15**, 2042- 2047 (2017).

39. X. Yin *et al.*, CRISPR-Cas9 and CRISPR-Cpf1 mediated targeting of a stomatal developmental
gene EPFL9 in rice. *Plant Cell Rep* **36**, 745-757 (2017).

40. J. Y. Zhang, S. B. He, L. Li, H. Q. Yang, Auxin inhibits stomatal development through
MONOPTEROS repression of a mobile peptide gene STOMAGEN in mesophyll. *Proc Natl Acad*
Sci U S A **111**, E3015-3023 (2014).

41. Z. Xu, G. Zhou, Responses of leaf stomatal density to water status and its relationship with
photosynthesis in a grass. *J Exp Bot* **59**, 3317-3325 (2008).

42. M. J. Lyu *et al.*, RNA-Seq based phylogeny recapitulates previous phylogeny of the genus
Flaveria (Asteraceae) with some modifications. *BMC Evol Biol* **15**, 116 (2015).

43. A. L. Ming-Ju *et al.*, The Coordination and Jumps along C_4 Photosynthesis Evolution in the
Genus Flaveria. *bioRxiv*, 460287 (2018).

44. K. Billakurthi *et al.*, Transcriptome dynamics in developing leaves from C_3 and C_4 Flaveria
species reveal determinants of Kranz anatomy. *bioRxiv*, (2018).

45. M. G. Grabherr *et al.*, Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat Biotechnol* **29**, 644-652 (2011).

46. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

47. L. Wang *et al.*, Comparative analyses of C_4 and C_3 photosynthesis in developing leaves of
maize and rice. *Nat Biotechnol* **32**, 1158-1165 (2014).

48. X. Robert, P. Gouet, Deciphering key features in protein structures with the new ENDscript
server. *Nucleic Acids Res* **42**, W320-324 (2014).

49. D. R. F. Kyte J, A simple method for displaying the hydropathic character of a protein. *Journal*
of Molecular Biology **157**, 105-132 (1982).

50. S. Perveen *et al.*, Overexpression of maize transcription factor mEmBP-1 increases
photosynthesis, biomass, and yield in rice. *J Exp Bot* **71**, 4944-4957 (2020).

51. D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol* **16**, 157 (2015).

52. R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res **32**, 1792-1797 (2004).

53. A. Stamatakis, RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
thousands of taxa and mixed models. *Bioinformatics* **22**, 2688-2690 (2006).

54. F. Ronquist *et al.*, MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice
across a large model space. *Syst Biol* **61**, 539-542 (2012).

55. S. A. Shiryev, J. S. Papadopoulos, A. A. Schaffer, R. Agarwala, Improved BLAST searches using
longer words for protein seeding. *Bioinformatics* **23**, 2949-2951 (2007).
56. X. J. Min, G. Butler, R. Storms, A. Tsang, OrfPredictor: predicting protein-coding regions in
EST-derived sequences. *Nucleic Acids Res* **33**, W677-680 (2005).
57. J. L. J. a. T. F. Ø. Claus Lindbjerg Andersen, Normalization of real-time quantitative reverse
transcription-PCR data a model-based variance estimation approach to identify genes suited
for normalization, applied to bladder and colon cancer data sets. *Cancer Res* **64**, 5245-5250
(2004).
58. D. P. K. Vandesompele J, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F., Accurate
normalization of real-time quantitative RT-PCR data by geometric averaging of multiple
internal control genes. *Genome Biology* **3**, research0034.0031–0034.0011 (2002).
59. D. A. Way, G. G. Katul, S. Manzoni, G. Vico, Increasing water use efficiency along the C3 to C4
evolutionary pathway: a stomatal optimization perspective. *J Exp Bot* **65**, 3683-3693 (2014).
60. V. S. R. D. a. M. Santakumari, Stomatal characteristics of some dicotyledonous plants in
relation to the C_4 and C_3 pathways of photosynthesis. *Plant and cell physiology* **18**, 935-938
(1977).
61. M. E. K. a. S. A. M. Marc D. Abrams, Relating Wet and Dry Year Ecophysiology to Leaf
Structure in Contrasting Temperate Tree Species. *Ecology* **75**, 123-133 (1994).
62. C. T. a. D. R. Murray, Effects of Elevated Atmospheric CO_2 Concentration on Leaf Anatomy and
Morphology in Panicum Species Representing Different Photosynthetic Modes. *International*
Journal of Plant Sciences **160**, 1063-1073 (1999).
63. A. S. Soares *et al.*, Adaxial/abaxial specification in the regulation of photosynthesis and
stomatal opening with respect to light orientation and growth with CO_2 enrichment in the C_4
species Paspalum dilatatum. *New Phytologist* **177**, 186-198 (2007).
64. I. Gindel, Stomatal Number and Size as Related to Soil Moisture in Tree Xerophytes in Israel
Ecological Society of America **50**, 263-267 (1968).
65. V. S. Pathare, N. Koteyeva, A. B. Cousins, Increased adaxial stomatal density is associated with
greater mesophyll surface area exposed to intercellular air spaces and mesophyll
conductance in diverse C4 grasses. *The New phytologist* **225**, 169-182 (2019).
66. J. A. Cardoso, M. Pineda, L. Jimenez Jde, M. F. Vergara, I. M. Rao, Contrasting strategies to
cope with drought conditions by two tropical forage C_4 grasses. *AoB Plants* **7**, plv107 (2015).
67. E. J. Edwards, C. J. Still, M. J. Donoghue, The relevance of phylogeny to studies of global
change. *Trends Ecol Evol* **22**, 243-249 (2007).
68. E. J. Edwards, C. J. Still, Climate, phylogeny and the ecological distribution of C4 grasses. *Ecol*
Lett **11**, 266-276 (2008).
69. S. Aubry *et al.*, A Specific Transcriptome Signature for Guard Cells from the C_4 Plant
Gynandropsis gynandra. *Plant physiology* **170**, 1345-1357 (2016).
70. S. P. Driscoll, A. Prins, E. Olmos, K. J. Kunert, C. H. Foyer, Specification of adaxial and abaxial
stomata, epidermal structure and photosynthesis to CO_2 enrichment in maize leaves. *J Exp*
Bot **57**, 381-390 (2006).
71. W.-H. W. Xi-Qing Wang, and Sarah M. Assmann, Differential Responses of Abaxial and Adaxial
Guard Cells of Broad Bean to Abscisic Acid and Calcium. *Plant physiology* **118**, 1421–1429
(1998).

72.   C. D. Muir, M. À. Conesa, J. Galmés, Independent evolution of ab- and adaxial stomatal
density enables adaptation. *biorxiv*,   (2015).
73.   R. J. Webster *et al.*, High C_3 photosynthetic capacity and high intrinsic water use efficiency
underlies the high productivity of the bioenergy grass Arundo donax. *Sci Rep* **6**, 20694 (2016).
74.   P. J. Vogan, M. W. Frohlich, R. F. Sage, The functional significance of C_3-C_4 intermediate traits
in Heliotropium L. (Boraginaceae): gas exchange perspectives. *Plant, cell & environment* **30**,
1337-1345 (2007).
75.   J. W. Maurice S. B. Ku, Ziyu Dai, Rick A. Scott, Chun Chu, and Gerald E. Edwards,
Photosynthetic and Photorespiratory Characteristics of Flaveria species. *Plant physiology* **96**,
518-528 (1991).
76.   O. Keerberg, T. Parnik, H. Ivanova, B. Bassuner, H. Bauwe, C_2 photosynthesis generates about
3-fold elevated leaf CO_2 levels in the C_3-C_4 intermediate species Flaveria pubescens. *J Exp Bot*
65, 3649-3656 (2014).
77.   R. F. Sage, T. L. Sage, F. Kocacinar, Photorespiration and the evolution of C_4 photosynthesis.
Annu Rev Plant Biol **63**, 19-47 (2012).
78.   C. B. Engineer *et al.*, Carbonic anhydrases, EPF2 and a novel protease mediate CO_2 control of
stomatal development. *Nature* **513**, 246-250 (2014).

Acknowledgements

The work was funded by Strategic Priority Research Program of the Chinese

Academy of Sciences (grant number: XDB27020105) and the general program of

National Science Foundation of China (31870214). We thank Prof. Rowan F. Sage

(University of Toronto) for providing the Flaveria cuttings.

Author contributions

YYZ and XGZ designed the study. YYZ observed and measured the stomatal

characteristic. YYZ carried out the molecular experiment and the treatment of

synthetic peptides in vivo. FFM conducted the transgene experiment. GYC guided the

treatment of peptides in vivo. Amy MJL analyzed the one kp RNA-seq data. YYZ and

XGZ interpreted results and wrote the article.

Competing interests

The authors declare no competing financial interests.

Figure Legend

Fig 1: Maximum stomatal conductance (g_{smax}) of *Flaveria* species

a, Schematic of the progresssion of C_4 evolution in *Flaveria* genus. **b**, A stomatal

graph taken with a light microscope showing stomatal parameters for *Flaveria* species

which are needed to be measured to calculate the g_{smax}. The graph of stomata is *rob*. *l*

(stomatal length), PL (pore length) and GCW (guard cell width). **c**, Correlation of the

guard cell width and stomatal length (left), correlation of the stomatal length and pore

length (right). **d**, The change of g_{smax} on the abaxial surface in *Flaveria* species with

different photosynthetic types from C_3 to C_4 grown outdoors (left) and in greenhouse

(right). **e**, The difference in g_{smax} of the abaxial surface between species with different

photosynthetic types in the *Flaveria* genus grown outdoors (left) and in greenhouse

(right). **f**, Correlation of g_{smax} and g_s for *Flaveria* species with different photosynthetic

types. Error bar indicates s.d. Different letters represent statistically significant

differences (one-way ANOVA, Duncan's multiple range test ($\alpha = 0.05$)). Pearson

correlation coefficient and the significance were noted in each panel.

Fig 2: Stomatal density (*SD*) and its patterning.

a, The difference in *SD* of abaxial surface between species with different

photosynthetic types in *Flaveria* genus grown outdoors (left) and in greenhouse

(right). **b**, The change of *SD* of abaxial surface from C_3 to C_4 in *Flaveria* species

grown outdoors (left) and in greenhouse (right). (n=4, biologically independent

replicates). **c,** Correlation of g_{smax} and *SD* for different *Flaveria* species. **d**, Correlation

of g_{smax} and *l* for different *Flaveria* species. **e**, Correlation of *l* and *SD* for different

Flaveria species. **f,** The photographs of stomatal patterning on abaxial surface in *rob*,

ram and *bid*. The photographs from the left to right are for *rob*, *ram* and *bid* grown

outdoors (top) and in greenhouse (bottom). **g**, Comparison of stomatal index (*SI*)

between *rob*, *ram*, and *bid*. Different colours represent plants grown outdoors (blue)

or in greenhouse (green). *l*, (n=20, biologically independent replicates). *SI*, (n=3,

biologically independent replicates). Error bar indicates s.d. Different letters represent

statistically significant differences (one-way ANOVA, Duncan's multiple range test (α

= 0.05)). Pearson correlation coefficient and the significance are noted in each panel.

Fig 3: Transcript abundance of stomatal developmental related genes between C_3 and

C_4 *Flaveria* species.

a, A model of the signaling pathway for the stomatal development in Arabidopsis. A

protodermal cell (PDC, white) becomes a meristemoid mother cell (MMC, orange),

which divides asymmetrically into two daughter cells, with one being a meristemoid

(M, red) and the other being a stomatal-lineage ground cell (SLGC). The M cell

develops into a guard mother cell (GMC, yellow); the SLGC develops into a

pavement cell (PC). The GMC conducts a symmetrical division to produce two

equal-sized guard cells (GC, green). EPF1, EPF2 and STOMAGEN competitively

bind to ER and TMM, which can deliver a signal to the YDA MAPK cascade. SPCH,

MUTE and FAMA ultimately are inactivated through phosphorylation by the YDA

MAPK cascade. **b**, Comparison of the expression of stomatal developmental related

genes between juvenile and mature leaves. **c**, The foldchanges of stomatal

development related genes for the C_3 and C_4 *Flaveria* species. Fold change means the

ratio between the gene expression in a C_3 *Flaveria* specie and gene expression in a C_4

Flaveria specie under the same leaf development stage and growth condition as the C_3

Flaveria species. The expression data were from (*44*), (*35*), (*34*) and One Thousand

Plants (1KP). **d,** The volcano plot for the expression of the stomatal development

related genes between C_3 and C_4 *Flaveria* species. The expression data were from (*35*)

and (*44*). **e**, Relative expression levels determined by RT-qPCR and expression level

determined by RNA-seq of FSTO (FSTOMAGEN) in *rob* and *bid* (n=4, biologically

independent replicates). **f**, Protein conservation analysis of the homologs of

FSTOMAGEN between different species in the *Flaveria* genus. The amino acid

regions marked by green line and red line represent the signal peptide and the

functional domain of STOMAGEN, respectively. Red letters represent conserved

amino acids in most species; white letters with red background represent amino acids

conserved across all species.

Fig 4: The Function of *FSTOMAGEN* in controlling the *SD*

a, Protein conservation analysis of the homologs of FSTOMAGEN in different species. The amino acid regions marked by green line and red line represent the signal peptide and the functional domain of FSTOMAGEN, respectively. Red letters represent conserved amino acids in most species, white letters with red backgrounds represent amino acid conserved across all species. **b**, Phylogenetic tree for FSTOMAGEN. blue represents dicotyledon and red represents monocotyledon. **c**, The structure of recombinant genes used to transform Arabidopsis. AFSTO represents the recombination of the signal peptide of STOMAGEN from Arabidopsis and the functional domain of FSTOMAGEN from *F.robusta*. FSTO represents the intact STOMAGEN from *F.robusta*. **d**, Photographs showing clustered stomata on a leaf for the transgenic Arabidopsis. Scar bar represents 10 um. **e**, **h**, The *SD* (**e**) and RNA-level (**h**) of the different transgenic Arabidopsis (FSTO) lines. **f**, **i**, The *SD* (**f**) and RNA-level (**i**) of the different transgenic Arabidopsis (AR) lines. **g**, Comparison of the transgenic FSTO lines and AR lines. **j**, Photographs of stomata in a leaf from Arabidopsis with FSTO and AFSTO over-expressed. Scar bar represents 20 um. Error bar indicates s.d. The n.s. represent statistically no significant differences (P>0.05, t-test, two-tailed distribution, unpaired).

Fig 5: The FSTOMAGEN peptide was applied to plants in vitro.

a, Photographs of leaf surface taken with differential interference contrast microscope showing the increased stomata density in *F. bidentis* (*bid*) leaves with FSTOMAGEN applied. Scale bar represents 20 μm. **b,** Photographs of leaf surface taken with differential interference contrast microscope showing the increased stomata density in Arabidopsis leaves with FSTOMAGEN applied. Scale bar represents 20 μm. **c**, The stomatal density for Arabidopsis leaves with FSTOMAGEN applied at different concentrations (n=8, biologically independent replicates). **d**, The stomatal density for *bid* leaves with FSTOMAGEN applied at different concentrations (n=8, biologically independent replicates). **e**, The stomatal index (SI) for *bid* leaves with FSTOMAGEN

applied at different concentrations (n=8, biologically independent replicates). Error bar indicates s.d. The asterisks represent statistically significant differences ($P<0.05$, t-test, two-tailed distribution, unpaired).

Fig 6: Expression level of *FSTOMAGEN* at different stages of C_4 evolution.

a, Expression levels of *FSTOMAGEN* at the different developmental stages of leaves in *rob* and *bid*. P denotes leaf primordia. Approximate leaf length for the different leaf stages was 0.3 cm (stage 1), 0.5 cm (stage 2), 1.5 cm (stage 3), 2 cm (stage 4), 3 cm (stage 5), 4 cm (stage 6), 6 cm (stage7) , 8 cm (stage 8) and 10 cm (stage 9). **b**, Comparison of relative expression levels of *FSTOMAGEN* between *rob*, *ram* and *bid*. (n=3, biologically independent replicates). **c,** Comparison of the expression levels of *FSTOMAGEN* between different species with different photosynthetic types in *Flaveria* genus. For RT-qPCR, C_3 species incude *pri* (n=5) and *rob* (n=13), intermediates species includes *ang* (n=3), *ram* (n=5) and *vag* (n=3), C_4 species include *tri* (n=4), *aus* (n=3), *bid* (n=4) and *koc* (n=3). For RNA-seq, C_3 species include *pri* (n=4) and *rob* (n=4), intermediates species includes *ano* (n=4), *pub* (n=4), *chl* (n=4), *bro* (n=4), C_4 speices include *tri* (n=4) and *bid* (n=4). The relative expression indicates the measurements with RT-qPCR, and the others were measured with RNA-seq. The geometric means of *EF1a* and *ACT7* were used as the reference in the calculation of expression levels for other genes. Error bars indicate s.d. The asterisks represent statistically significant differences ($P<0.05$, t-test, two-tailed distribution, unpaired). Different letters represent statistically significant differences (one-way ANOVA, Duncan's multiple range test ($\alpha = 0.05$)).

Fig 7, Comparison of stomatal pattern and expression levels of stomatal development related genes in the C_4 lineages other than *Flaveria*

a, Stomatal characteristics for rice and maize. *SD*, (n=8, biologically independent replicates); *l*, (n=25, biologically independent replicates); *SI*, (n=3, biologically independent replicates); g_s, (n=8, biologically independent replicates). Error bar

indicates s.d. The asterisks represent statistically significant differences ($P<0.05$, t-test, two-tailed distribution, unpaired). **b**, Photographs of stomatal pattern in rice and maize. Scale bars represent 50 μm. **c**, Comparison of transcript abundance of stomatal development related genes between rice and maize. The developmental stages of leaves were divided into 11 and 15 parts in rice and maize, respectively. R represents the leaves of rice, M represents the leaves of maize. **d**, Comparison of the expression levels of *STOMAGEN* homologs between rice and maize. There are one and two paralogs of *STOMAGEN* that is expressing in rice and maize, respectively. *OsSTO* represents the homologs of *STOMAGEN* in rice. *ZmSTO* represents the homologs of *STOMAGEN* in maize. **e**, The families and numbers of species with different photosynthetic type that we used to analyse. **f**, Comparison of the expression of C_4 related genes between species with different photosynthetic types. **g**, Comparison of the expression of *STOMAGEN* homologs between species with different photosynthetic types. The asterisks represent statistically significant differences ($P<0.05$, t-test, two-tailed distribution, unpaired).

Fig 8：Schematic of the role of *FSTOMAGEN* in the changes of WUE during C_4 evolution. During the C_4 evolution, *FSTOMAGEN* (*FSTO*) changes g_{smax} through regulating stomatal density, which in turn affectes gas exchange and improves the water use efficiency of C_4 plants. C_3-C_4 denotes the intermediate plants between C_3 and C_4 plants during C_4 evolution.

Legends for Supplemental Figures

Supplementary fig 1. Comparison of *SD* between C_3 and C_4 species. **a**, Comparison of *SD* on the abaxial surface of leaves between 70 C_3 and 47 C_4 species, with data collected from literature (see details in the discussion section). b, Comparison of *SD* on adaxial surface of leaves between 26 C_3 and 27 C_4 species with data collected from literature (see details in the discussion section). Error bar indicates s.d. The asterisks represent statistically significant differences ($P<0.05$, t-test, two-tailed distribution, unpaired).

Supplementary fig 2. a, Stomatal conductance (g_s) responses to net photosynthesis rate (A) under saturated light. The circles on the graph indicate larger g_s values in different photosynthetic types (Date from (*2, 59*)). **b,** The correlation of g_{smax} (abaxial + adxial) and g_s. **c,** G_s responses to A at the growth light in two *Flaveria* species (*rob* and *bid*) (n=6).

Supplementary fig 3. g_{smax} on the adxial surface of leaves in *Flaveria*. **a**, The difference in g_{smax} in species with different photosynthetic types grown outdoors (left) or in greenhouse (right)**. b,** The changes of g_{smax} on the adaxial leaf surface from C_3 to C_4 species in the *Flaveria* genus grown outdoors (left) and in greenhouse (right). (n=3, biologically independent replicates). Correlation of g_{smax} and g_s on the adaxial surface of leaves. Different letters represent statistically significant differences (one-way ANOVA, Duncan's multiple range test ($\alpha = 0.05$)).

Supplementary fig 4. *SD* on the adaxial surface of leaves in *Flaveria*. **a**, The difference in *SD* in species with different photosynthetic types grown outdoors (left) or in greenhouse (right)**. b**, The changes of *SD* on the adaxial leaf surface from C_3 to C_4 species in the *Flaveria* genus grown outdoors (left) and in greenhouse (right). (n=3, biologically independent replicates). **c**, Correlation of g_{smax} and *SD* on the adaxial surface of leaves. **d**, Correlation of g_{smax} and *l* on the adaxial surface of leaves. **e**, Correlation of *l* and *SD* on the adaxial surface of leaves. Different letters represent statistically significant differences (one-way ANOVA, Duncan's multiple range test ($\alpha = 0.05$)).

Supplementary fig 5. *SD* in *rob* (C_3), *ram* (intermediate) and *bid* (C_4) grown from seeds. The *SD* on abaxial (ab) (n=5, biologically independent replicates) and adaxial (ad) (n=4, biologically independent replicates) surfaces were observed in *rob*, *ram* and *bid* grown in greenhouse. Different colours represent different photosynthetic types. Different letters represent statistically significant differences (one-way ANOVA, Duncan's multiple range test ($\alpha = 0.05$)).

Supplementary fig 6. Conservative analysis of amino acid sequences for EPF1 and EPF2. **a**. Comparison of amino acid sequences of homologs of EPF1 between

different *Flaveria* species and Arabidopsis. **b**. Comparison of amino acid sequences of homologs of EPF2 between *Flaveria* species and Arabidopsis. White letters with red background represent amino acids conserved across all species. Red letters represent amino acids with similar biochemical property.

Supplementary fig 7. Hydrophobicity analysis of proteins. **a**, the protein of stomagen (left) and the protein of FSTOMAGEN (right). **b**, the protein of EPF1 (left) and the protein of FEPF1 (right). **c**, the protein of EPF2 (left) and the protein of FEPF2 (right).

Supplementary fig 8. Assessment of the stability of expression levels of reference genes. **a,b**, Comparison of the stability of expression levels of candidate reference genes in the *Flaveria* species by RNA-seq. **c**, Gel electrophoresis showing the specificity of the primers for candidate reference genes. From left to right, *ACT7*, *EF1a*, *J3*, *PNdO*, *CPN20*, *UBC9*, *GAPDH*, *F2N1.5*, *PSRP4*, *CACMS*, *HEME2*, and *UBQ11*. **d**, The coefficient of variations (CV) for the candidate reference genes. **e**, The expression stability for candidate reference genes calculated by Genorm. **f**, The expression stability for candidate reference genes calculated by Normfinder.

Supplementary fig 9. The expression levels of stomatal development related genes in *Flaveria* by RNA-seq. Expression levels of stomatal development related genes in species with different photosynthetic types in the *Flaveria* genus.

Supplementary fig 10. Relative expression levels of *FSTOMAGEN* and *SD* for *rob* (C$_3$) under different light conditions. (A) Relative expression levels for *FSTOMAGEN* under two different light conditions. (B) *SD* under two different light conditions. Natural light: the maximal photosynthetic photon flux density in the greenhouse was about 1300~1400 mmol m^{-2} s^{-1}. Light in the phytotron: 20~100 mmol m^{-2} s^{-1}. Reference gene used to calculate the expression level was *ACT7*. The asterisks represent statistically significant differences ($P<0.05$, t-test, two-tailed distribution, unpaired).

Supplementary fig 11. Relative expression levels of stomata development related genes determined with different reference genes and different primers. **a, b, c,** Quantification with different primers and the same *EF1a* reference gene. Relative

expressions shown in **a and b** were determined using the same primer pairs while that

in **c** used other primer pairs (See details of the primers used in Table S2). **d,** Relative

expression levels determined with the gene *J3* used as the reference gene.

Supplementary fig 12. Transcript abundance of genes in the C_4 cycle and C_4 related

transporters. J represents juvenile leaves and M represents mature leaves. Different

colours represent species with different photosynthetic types in the *Flaveria* genus.

Red colour represents C_3 species, grey colour represents C_3-C_4 species, green colour

represents C_4-like species, and blue colour represents C_4 species.

Supplementary fig 13. Alignment of the functional domain of *STOMAGEN* with

transcripts from different *Flaveria* species by Blast. From the top to bottom, *F.*

robusta, F. pringlei, F. angustifolia, F. ramosissima, F. vaginata, F. australasica, F.

bidentis.

Supplementary fig 14. The expression level of *FSTOMAGEN* in different *Flaveria*

species based on RT-qPCR (**a**) or RNA-seq (**b**). **c,** The correlation of *SD* and

FSTOMAGEN expression in *Flaveria* species.

Supplementary fig 15. Comparison of the expression level of *FEPF1* and *FEPF2*

between *rob* and *bid Flaveria* species.

Supplementary fig 16. The analysis for the *SD* of T2 generation of transgenic

Arabidopsis (AFSTO and FSTO). **a,** Comparison of *SD* between WT and transgenic

Arabidopsis AR, FSTO respectively. **b**, RNA level identification of transgenic

Arabidospsis. Two independent lines was analyzed for each trangene event.

Supplementary fig 17. The increase of *SD* in cotyledon of *F. bidentis* (*bid*) with in

vitro application of FSTOMAGEN. Photographs of the leaf surface in the untreated

(control) and treated bid plants. scale bar, 20 mm.

Supplementary fig 18. The gene tree of STOMAGEN based on bayesian-inference

method.

1004

1005

1006

1007

Supplementary table 1. Statistics for the mapping of the RNA-seq data of the

juvenile and mature leaves in different species of the *Flaveria* genus.

Supplementary table 2. Primers used in this study

Supplementary table 3. The effect of genes related to stomatal development in

Arabidopsis on stomatal phenotypes.

Supplementary table 4. The N50 of the mapping of the RNA-SEQ data taken from

1KP database for the juvenile and mature leaves

Supplementary table 5. The mapping rates of the mapping of the RNA-SEQ data

taken from 1KP database for the juvenile and mature leaves.

1017