The evolution of stomatal traits along the trajectory towards C$_4$ photosynthesis

Yong-Yao Zhao[1,2], Mingju Amy Lyu, FenFen Miao, Genyun Chen, Xin-Guang Zhu[1,2]

1. Center of Excellence for Molecular Plant Sciences, Chinese Academy of Sciences

2. State Key Laboratory for Plant Molecular Genetics

Abstract

C$_4$ photosynthesis optimizes plants carbon and water relations, allowing high photosynthetic rate with low stomatal conductance. Stomata have long been believed as a part of C$_4$ syndrome. However, it remains unclear how stomata traits evolved along the path from C$_3$ to C$_4$. Stomatal patterning was examined in *Flaveria* genus, a model for studying C$_4$ evolution. Comparative, transgenic and semi-*in-vitro* experiments were used to study molecular basis that underlies stomatal traits along C$_4$ evolution. Novel results: the evolution from C$_3$ to C$_4$ species through intermediate species is accompanied by a stepwise rather than an abrupt change in the stomatal traits. The initial change occurs near Type II and dramatic change occurs at the C$_4$-like species. On the road to C$_4$, stomata become less in number but bigger in size and changes in stomatal density dominates changes in maximum stomatal conductance (g_{smax}). The reduction of *FSTOMAGEN* expression underlies altered g_{smax} between *Flaveria* species with different photosynthetic pathways and likely occurs in other C$_4$ lineages. Our study provides insight into the pattern, mechanism and role of stomatal evolution along the road towards C$_4$. This work highlights the stomatal traits in the current C$_4$ evolutionary model and the co-evolution of photosynthetic pathway and stomata.

Key words:

Stomatal traits, *Flaveria*, C$_4$ evolution, STOMAGEN, photosynthetic pathway

Introduction

Stomata are small pores surrounded by a pair of guard cells on the leaf surface, which control the passage of water and CO_2 between leaf and atmosphere (1). The emergence of C_4 photosynthesis optimizes plant water and carbon relations, allowing low stomatal conductance (g_s) (i.e reduced water loss) without compromising carbon assimilation (2-4), which raises a possibility that C_4 photosynthesis might have been selected as a water-conserving mechanism (3) and that water limitation might be a primary driver of C_4 evolution (5-7). Comparison study on multiple independent lineages of C_4 grasses showed that the g_s is significantly lower in C_4 species relative to C_3 relatives (2, 8), and the higher intrinsic water use efficiency (iWUE) of C_4 plants compared to the C_3 plants can be attributed to both lower g_s and higher photosynthetic CO_2 uptake rate (A) (2, 9-11). The high carbon assimilation capacity of C_4 plants contributes to their high biomass, while low g_s is crucial for C_4 plants to better adapt to saline land, and hot, arid & open habitat (12, 13).

In a long-term adaptation, C_3 plants generally adjust their maximal stomatal conductance (g_{smax}) (14-16). Phylogenetically controlled comparison shows that C_4 grass have lower g_{smax} compared with their close related C_3 grass (17). In line with this, C_4 grass also have lower operating g_s (2, 17). Though low g_{smax} is regarded as an important feature in the C_4 syndrome (18), the changes of g_{smax} along the road to C_4 photosynthesis have not been studied (19). G_{smax} is determined by stomatal density (SD) and stomatal size (SS). Although higher SD of C_4 species compared to C_3 species has been observed (20); in some cases, C_4 plants can have a lower SD compared to C_3 species (21). These differential changes in stomatal traits between photosynthetic types might be related to phylogenetic effects (2, 22, 23). As a reflection of this, comparative study using plants from different C_4 clades shows that lower g_{smax} in C_4 plants is due to either a smaller stomatal size (SS) for some C_4 clades, or lower stomatal density (SD) for other clades (17), but after accounting for the phylogenetic effects, SD in C_4 is lower than C_3 species and there is no significant difference in SS between C_3 and C_4 grass (17). Earlier study suggests that the influence of

photosynthetic pathway on *SD* and *SS* is only statistically marginally significant (*17*),

which might because stomatal development is affected by many factors, especially the

environmental conditions such as soil moisture and fertility status (*24-27*). Therefore,

more evidence is still needed to study the changes in *SS* and *SD* during C_4 evolution.

Stomatal development is controlled by a well-studied genetic program (*28*),

which involves a series of ligands and transmembrane receptors. The ligands and

receptors lead to downstream changes of mitogen-activated protein kinase (MAPK)

and transcription factors (*28*). TOO MANY MOUTHS (TMM) and ERECTA-family

(ERfs: ER, ERL1, ERL2) are the transmembrane receptor-like kinases that negatively

regulate stomatal development (*29, 30*), and the TMM interacts with ERfs (*29, 31*).

The ligands that bind to ERfs and TMM receptors are mainly EPIDERMAL

PATTERNING FACTOR/EPIDERMAL PATTERNING FACTOR-LIKE family

(EPF/EPFL-family) peptides (*32*). Overexpression of *EPF1* decreases *SD* (*33*). *EPF2*

has a similar genetic effect as *EPF1* (*34*). STOMAGEN prevents the phosphorylation

of downstream receptors, which results in the changed expression of basic Helix-loop-

Helix (bHLH) paralogs, SPCH, MUTE and FAMA, ultimately positively regulates

stomatal development (*35-38*).

C_4 plants evolved from C_3 ancestors through a number of intermediate stages,

which were shown in a number of genera (*7, 39-41*). Among these genera, we chose

Flaveria genus as the model for this study. The *Flaveria* genus contains close species

with multiple different photosynthetic types which cover the sequential stages of C_4

evolution, i.e. C_3, C_3-C_4 type I, C_3-C_4 type II, C_4-like and C_4 (*4, 41-45*). This genus is

the youngest genus in which C_4 photosynthesis evolved (*46*) and have many

intermediate species (*47*). A close relationship between *Flaveria* species can minimize

the impact of non-photosynthetic effects on stomatal traits (*11, 47*). The species in

Flaveria genus grows on similar growth habits and also have similar morphologies

(*13, 48-51*). These make *Flaveria* genus an ideal system to study the evolutionary

process from C_3 to C_4 (*39*). The physiology, structure, biochemistry and molecular

changes of the *Flaveria* genus along C_4 evolution have been systematically evaluated

(*41-45, 47, 51, 52*). In fact, the current C_4 evolutionary model illustrating the path

from C_3 to C_4 photosynthesis is largely based on studies on the *Flaveria* species (*39,*

53). The current model of C_4 evolution include many steps *(41, 45, 47, 48, 51, 52, 54)*.

First, bundle sheath cells and vein density showed C_4 characteristics (*39, 43, 55*).

Then, glycine decarboxylase (GDC) transfers from mesophyll cells to bundle sheath

cells (*56*), resulting in a modest CO_2 concentrating mechanism (*55*), which bridges the

formation of C_4 metabolism (*39, 41*). Although stomata have long been thought as a

part of C_4 syndrome (*17, 18*), the evolutionary changes of stomata traits in current C_4

evolutionary model have been much less studied.

This study aims to answer the following questions: 1) In the context of the

current C_4 evolution model, will stomatal traits and g_{smax} change with the evolution of

photosynthetic type? How and when do stomatal traits change along the path to C_4? 2)

What's the molecular mechanism underlying the change in the stomatal traits during

C_4 evolution? 3) What are the roles and drivers of stomatal evolvement in the

evolutionary process towards C_4 photosynthesis?

Result

Maximum stomatal conductance and stomatal pattern during the evolution of C_4

photosynthesis in *Flaveria*

As a model of C_4 evolution, there are distinct evolutionary phases in *Flaveria*

genus (Fig 1a). The maximum value of operational g_s (marked with circles on figure)

for C_3 *Flaveria* species under saturated light intensity was higher than that for C_4

species, indicating that the C_3 *Flaveria* species tends to have a higher g_s

(Supplementary Fig 1a). The pore length of stomata was approximately equal to

0.41×stomatal length, and the guard cell width was approximately equal to

0.33×stomatal length (Supplementary fig. 19). Through measuring stomatal density

and stomatal length, we calculated the g_{smax} for *Flaveria* species. The results show

that, whether growing under outdoor natural light or constant light in the greenhouse,

the g_{smax} decreased gradually from C_3 through intermediate species to C_4 species (Fig

1b,c,d,e). A significant correlation was found between the g_{smax} and g_s under saturated

light for the *Flaveria* species with different photosynthetic types (Fig 1f). Under

saturated light, the g_s of *son* (Type I) was extremely high (marked with circle on

figure), matching with its highest g_{smax} within the *Flaveria* genus (Fig1d,

Supplementary Fig 1a). Similar to that under the saturated light intensity, the WUE

(A/g_s) of *rob* (C_3) measured under unsaturated light intensity was higher than that of

bid (C_4)*,* and the g_s of *rob* was 71% higher than that of *bid* (Supplementary Fig. 1d).

In comparison to C_4 and C_4-like photosynthesis, the stomatal densities (*SD*) of C_3

and Type I photosynthesis on the abaxial leaf surface of *Flaveria* were nearly 150%

higher (Fig. 2a,b,c,d). With the progression from C_3 to C_4 species, there was a gradual

decrease in *SD* (Fig. 2a, c). The difference in *SD* and *SS* between C_3 and Type I

species wasn't as large as that between C_3 and C_4 species; and the *SD* and *SS* of Type

II species fell in between Type I and C_4-like species (Fig. 2b,d). The *SD* of C_4 and C_4-

like species were similar, although the *SD* of C_4-like species was slightly less than that

of C_4 (Fig. 2b,d). For the adaxial leaf surface, the *SD* of C_3 species was again higher

than that of the C_4 species, although the difference between *SD* in the intermediate

species was less significant (Supplementary fig. 3a, b, c, d). There was a strong

positive correlation of g_{smax} and *SD* either on the abaxial leaves (R^2=0.8955, p<0.0001)

(Fig. 2e) or on adaxial leaves (R^2=0.9165, p<0.0001) (Supplementary fig. 3e) and a

significant negative correlation of g_{smax} and *l* on abaxial leaves (R^2=0.2596, *p*<0.005)

(Fig. 2f), indicating the *SD* determined the change of g_{smax}, not stomatal size (SS). *SD*

and *l* showed a significant negative correlation on abaxial leaves (R^2=0.5093,

p<0.0001) (Fig. 2g) or on adaxial leaves (R^2=0.2055, *p*<0.05) (Supplementary fig. 3g).

Flaveria species, i.e. *rob*, *ram* and *bid,* were chosen to represent C_3, intermediate

and C_4 species, respectively, to analyse the ratio between stomatal number and the

number of epidermal cells, i.e. the stomatal index (*SI*). The *SI* of *rob* was nearly 50%

higher than *bid* and *ram* in plants grown in either greenhouse or outdoors, suggesting

that the proportion of stomata occupying the epidermis, contributed to the highest *SD*

in *rob* (Fig. 2h). In addition to the analysis of *Flaveria* plants grown from cuttings, we

also planted *Flaveria* germinated from seeds. Results show that the *SD* of C_3 species

was higher than C_4 species, consistent with the results from cuttings (Supplementary

Fig. 4).

Comparison of the sequence and transcription abundance of genes related to stomatal pattern between C_3 and C_4 *Flaveria* species

The signaling pathway for stomatal development has been studied extensively (*57*). We identified amino acid sequence encoded by homologs of stomatal development related genes in *Flaveria*. Amino acid sequence alignment analysis showed that epidermal patterning factor/epidermal patterning factor-Like gene family (*EPF*/*EPFL*-family) homologs was conserved at the C-terminal part of amino acid sequence (Fig. 4A, Supplementary fig. 5). Members of *EPF*/*EPFL*-family genes encode secretory proteins containing signal peptides at the N-terminus and functional domains at the C-terminus (*33, 34, 58*). Analysis shows that the hydrophobicity of EPFs/EPFL homologs in *Flaveria* was similar to EPF/EPFL in Arabidopsis (Supplementary fig 6), implying the homologs are also the secretory proteins and should have the same physiological function as EPF/EPFL. We hypothesized that either the amino acid sequence or the expression of stomatal development related genes changed during C_4 evolution. Analysis results reveals no differences within the functional region of amino acid sequence for STOMAGEN, EPF1 and EPF2 homologs between C_3 and C_4 species (Fig. 5a, Supplementary fig. 5a, b), indicating the change of amino acid sequence is unlikely to be a cause of the changed *SD* in C_4 evolution.

As up-regulation of transcription has been a major approach to increase the expression of many genes participating in the building of C_4 syndrome during C_3 to C_4 evolution in *Flaveria* (*41, 47*), we compared the transcript abundance of stomatal development related genes between C_3 and C_4 *Flaveria* species. The expression patterns of known C_4 related genes in our data analysis were same as the previous works, which indicates that our data collection and analysis procedure were reliable (Supplementary Fig. 11) (*41, 47*). We found that the homologs of most stomatal development related genes showed higher transcript abundance in developing tissues as compared to developed tissues (Fig. 3b), which is consistent with their function of regulating stomatal differentiation in developing leaf (*33, 36, 38, 59*). *STOMAGEN*,

EPF1, *SDD1*, MPK3 and *TMM* showed obviously lower expression levels in C_4

species than in C_3 species (Fig. 3c). Among these genes, *STOMAGEN* was the one

that changed the most (Fig. 3c) and showed the most significant difference (Fig. 3d).

EPF1, *SDD1, MPK3* and *TMM* are negative regulators for stomatal development

(Supplementary table 3) (*32-34, 60*). Actually *SD* in C_4 species was lower than that in

C_3 (Fig 2a,b,c,d), indicating these genes should not be the factors responsible for the

decrease in *SD* of C_4 species. *STOMAGEN* is a positive regulator of *SD*

(Supplementary table 3) (*32, 36, 37*), whose lower expression in C_4 species was in

line with the decreased *SD* in C_4 species (Fig.2a,b,c,d,Supplementary Fig. 3c, d). RT-

qPCR quantification of the expression levels of *STOMAGEN*, *EPF1*, *EPF2* homologs

between C_3 and C_4 species grown under our conditions also confirmed the results

from RNA-SEQ (Fig. 3e, Supplementary fig. 14). In this study, we renamed

STOMAGEN in *Flaveria* as *FSTOMAGEN*.

There are considerable differences in amino acid sequence of signal peptide of

STOMAGEN homologs, especially between species with large evolutionary distance

(Fig. 4a Fig. 5a). The functional domain of *STOMAGEN* is however relatively

conserved, not only between different *Flaveria* species (Fig. 5a), but also between

species that have a large evolutionary distance (Fig. 4a). A phylogenic tree

constructed for *STOMAGEN* from different plants suggest *STOMAGEN* homologs of

species in the genus *Flaveria* are extremely conserved (Supplementary fig. 18).

Experiments with exogenously applied STOMAGEN peptides shows that the

STOMAGEN induce increased SD extracellularly (*36, 37, 58, 59*). Previous studies

showed that overexpressing rice *STOMAGEN* homologs increases SD in Arabidopsis

and RNAi silencing of *STOMAGEN* homologs reduces SD in rice (*61-63*), implying

STOMAGEN homologs also regulate stomatal development in other species besides

Arabidopsis. When plants are grown under conditions that promote or inhibit stomatal

development, the expression of *STOMAGEN* is known to increase or decrease,

respectively (*24, 59, 64*). A similar modulation of *STOMAGEN* expression levels was

observed for *rob* when grown at different light conditions (Supplementary Fig 9).

To confirm that *FSTOMAGEN* can induce stomatal generation and assess the

ability of *FSTOMAGEN* to regulate stomatal development, we overexpressed the

FSTOMAGEN in *Arabidopsis*. Specifically, we recombined the signal peptide of

STOMAGEN from *Arabidopsis* to the functional region of *FSTOMAGEN* and named

the hybrid gene as AFSTO (Fig, 4e). Overexpressing either the AFSTO or the intact

FSTOMAGEN (FSTO) from *F.rob* into *Arabidopsis* resulted in similar increased *SD*

and could exceed 800 stomata per unit leaf area (800 mm^{-2}) (Fig. 4f,g,i,j,h,

Supplementary fig 15). Identification of transcription showed successful transgenes

and the expression level of the AFSTO or FSTO increased with the increase in SD

(Fig. 4f, g, i, j). Furthermore, overexpression of *FSTOMAGEN* also induced clustered

stomata (Fig. 4k, Supplementary fig 20), as also reported earlier for the over-

expression of *STOMAGEN* (*36*). To further confirm that *FSTOMAGEN* can increase

SD in *Flaveria*, we synthesized a peptide representing the FSTOMAGEN functional

region, consisting of 45 amino acids. In order to confirm our synthesized peptides that

have been folded were biologically active, the FSTOMAGEN peptides were applied

in vitro on the leaves of Arabidopsis and resulted in increased *SD* and clustered

stomata (Fig 4b, Supplementary fig 17b). Application of the peptides on leaves of *bid*

also increased *SD* and *SI* and caused stomatal cluster (Fig 4c, d, Supplementary fig

17a). These results demonstrate that, in addition to its *in vivo* function, *in vitro*

application of FSTOMAGEN also directly positively regulate *SD* in *Flaveria*. This is

consistent with the notion that *STOMAGEN* is expressed in mesophyll cells and

secreted out of cells to influence the stomatal development in the epidermal layer in

Arabidopsis (*36, 37*).

The transcript abundance of *FSTOMAGEN* during C$_4$ evolution in *Flaveria*

We examined the copy number of *FSTOMAGEN* in *Flaveria* using blastn with

the coding sequence of *FSTOMAGEN* as query sequence against the assembled

contigs of each *Flaveria* species from 1KP. Results show that there was only one hit

sequence with e-value less than 10^{-5} in each species, indicating that there is a single

copy of *FSTOMAGEN* in *Flaveria* (Supplementary fig. 12). Therefore we can ignore

the impact of other *FSTOMAGEN* paralogs on stomatal development, and there will
be no quantitative errors in the gene expression levels caused by other paralogs that
were not considered. Examination of the RNA-seq data of *Flaveria* shows that the
expression levels of *FSTOMAGEN* in C_3 species were markedly higher than those in
C_4 species along different developmental stages of leaves (Fig. 5c). During leaf
growth and development, the expression of *FSTOMAGEN* in both *bid* and *rob*
increased first, reaching the highest level at leaf development stage 3 (leaf length 1.5
cm) or 4 (leaf length 2.0 cm), and then declined (Fig. 5c). The similar patterns of
FSTOMAGEN expression in these two *Flaveria* species indicates that it would be
valid to compare the *FSTOMAGEN* expression levels in leaves at the same
development stage between *Flaveria* species. The expression level of *FSTOMAGEN*
in *bid* (C_4) was lower than that in *rob* (C_3), and that of *ram* (intermediate species) lied
between them (Fig. 5b). This encouraged us to test more *Flaveria* species with
different photosynthetic types. Similar results were obtained with either RNA-seq or
RT-qPCR (Fig. 5d, Supplementary Fig. 13), where we found the gradual decrease of
FSTOMAGEN expression from C_3 to intermediate species to C_4 species.

The expression of *FSTOMAGEN* homologs for species with different
photosynthetic type in multiple C_4 lineages.

To gain insight into whether *STOMAGEN* homolog might be a generic factor
leading to the difference in *SD* between C_3 and C_4 species in higher plants, we tested
its expression level in different C_4 lineages other than the *Flaveria genus*. The *SD* of
abaxial leaf in C_3 rice was 240% higher than that in C_4 maize, consistent with a 570%
higher g_s in rice compared to maize (Fig. 6a, c). The stomatal length (*l*) in maize was
133% longer (Fig. 6b). We analysed the transcript abundance of *STOMAGEN*
homologs, since there are two and three paralogs of *STOMAGEN* in rice and maize
respectively (*61*), only one in rice and two in maize are expressed. The *STOMAGEN*
homologs in rice had much higher transcript abundance than those in maize (Fig. 6f),
while the transcript abundance of other stomatal development related genes did not
differ to the same extent between rice and maize (Fig. 6e). This result encouraged us

to test more C_4 lineages (Fig. 6g). C_4 photosynthesis related genes were indeed higher

in C_4 species, suggesting the data and analysis pipeline can be used to compare the

differences in expression of genes between C_3 and C_4 photosynthetic types (Fig 6h). It

was showed that the expression levels of *STOMAGEN* homologs for C_4 were lower

than those for C_3, while intermediate had expression between them (Fig 6i).

Method

Plant materials, sample condition and data sources

All *Flaveria* species used in this study were kindly provided by Prof. Rowan F.

Sage (University of Toronto). Plants used for stomatal phenotyping were re-grown

from cuttings and placed in an environment humidified with humidifiers to generate

roots. After plants generated roots, they were transferred outdoors (Institute of Plant

Physiology and Ecology, Chinese Academy of Sciences, Shanghai, China), and

transplanted into 4-L pots filled with topsoil in April. In July, the number of stomata

on abaxial and adaxial surfaces of leaves were counted. Another batch of *Flaveria*,

which were also grown from cuttings, were transferred to the phytotron (Center of

Excellence for Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai,

China). Growth temperature of the phytotron was maintained at 25-28□, photoperiod

was maintained at 16h light / 8h dark, relative humidity (RH) was at 60-65% and

photosynthetic photon flux density (PPFD) was maintained at about 180 μmol m^{-2} s^{-1}.

Two months later, the numbers of stomata on the abaxial and adaxial surfaces of

leaves were counted. RNA was extracted from 1.5~3.0 cm long young leaves on the

axial shoot which were still expanding. At three hours before sampling, we provided

sufficient water to ensure that plants were not water limited, since drought and water

status can affect expression levels of *STOMAGEN* (*24, 26*). Gene expression was also

examined under normal growth conditions by RT-qPCR for the *rob*, *ram* and *bid*,

which are representative of C_3, intermediate and C_4 species in the *Flaveria* genus.

These three species were grown in the same phytotron with the same environmental

conditions, except that we did not water plants before sampling. To test whether

plants grown from cuttings or from seeds show difference in the *SD*, we further grew

Flaveria species from seeds in phytotron and examined stomatal properties. Same

environmental conditions were maintained for phytotron as above. Two months after

the germination of seeds, we counted stomatal number.

All plants were well watered to avoid drought stress during their growth. We

supplied the same amount of commercial nutrient solution (Peters Professional,

SCOTTS, N:P:K = 20:20:20+TE, at recommended dose for application) to the

different species to avoid nutrient deficiency.

Rice (*Oryza Sativa Nipponbare*) and maize (*Zea maize* B73) were grown in the

greenhouse (Center of Excellence for Molecular Plant Sciences, Chinese Academy of

Sciences, Shanghai, China) under a PPFD of 550 umol/m^2/sec, a photoperiod of 12h

light / 12h dark, and a daytime temperature of around 31℃ and a night-time

temperature of around 22℃. The relative humidity was maintained at around 50-55%.

The plants were regularly watered to avoid drought stress.

In addition to the expression data collected by in this study, we also used public

data for comparison of transcript abundance. Considering that stomatal development

related genes are mainly expressed in young tissues, in each of these used public data

source, samples for young leaves were included. For the *Flaveria*, data from four

sources for leaves of plants grown under normal condition were analysed. The first

data source is the One Thousand Plants (1KP) Consortium, and we subsequently

analyzed the data according to the methods described in previous work (*42, 65*). The

second and third data sources were Mallman et al (2014) (*41*) and Udo Gowik et al

(*47*). The fourth data source was Kumari Billakurthi et al (2018) (*66*). The last three

data sources are all from Heinrich Heine University. For the second and third data

source, the second and fourth visible leaves were sampled; the second leaf was not

completely mature, and stomatal developmental related genes were still actively

expressed. In addition to *Flaveria*, transcripts of C_3 and C_4 species from other C_4

lineage sequenced from 1KP were assembled with Trinity (*67*) with default

parameters. Transcript abundances were analyzed by mapping the reads to assembled

contigs for corresponding samples using the RSEM package with mapping option of

bowtie 1 (*68*), and outliers that less than Q1 (lower quartile) - 1.5IQR (interquartile

range) and greater than Q3 (upper quartile) + 1.5IQR (interquartile range) were

removed. Expression levels for stomatal development related genes in developing

leaves for rice and maize were obtained from Wang et al. (*69*).

Conservation and hydrophobicity analysis of proteins

Amino acid sequence of the homologs of *STOMAGEN*, *EPF1* and *EPF2* were

aligned by clustalx, and graphs for the multiple alignments were drawn with ESPrit

3.0 (*70*) (http://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi). Hydrophobicity of proteins

was determined by ProtScale (*71*) (https://web.expasy.org/protscale/).

Construction of the phylogenetic tree of *STOMAGEN*

In order to construct the gene evolutionary tree of *STOMAGEN*, we searched the

orthologs of *STOMAGEN* in 28 representative species among Viridisplantae as

demonstrated in (*72*). The protein sequences of representative species were

downloaded from Phytozome (https://phytozome.jgi.doe.gov/). Orthologs of

STOMAGEN were predicted applying OrthoFinder (V 0.7.1) (*73*) with default

parameters. Proteins sequences of STOMAGEN orthologs were then aligned with

MUSCLE (*74*). We then inferred the gene phylogeny using both maximum likelihood

(ML) and Bayesian-inference (BI) methods. The ML tree was constructed with

RAxML (V2.2.3) (*75*), where bootstrap scores were obtained by running 100,000

independent trees. BI tree was built with MrBayes package (v 3.2.1) (*76*), where

posterior probabilities were calculated by running 2000,000 generations with the first

2000 generations being used as burn-in. In both cases, PROTGAMMAILG model

(General Time Reversible amino acid substitution model with assumption that

variations in sites follow gamma distribution) was used to infer the gene phylogeny.

The protein sequence of AT4G14723 (*CLL2*) from *A. thaliana* was used as outgroup

as it showed the highest identity with STOMAGEN in *A. thaliana*, with a protein

identity of 26.97% and E-value of 0.037 based on Blastp (*77*). Protein sequences for

the *Flaveria* species were predicted using OrfPredictor (*78*) based on de novo assembled transcripts as described in (*79*).

Measurement of stomatal conductance (g_s)

The LI-6400 Portable Photosynthesis System (Li-Cor, Inc., Lincoln, NE, USA) was used to measure g_s on the youngest fully expanded leaves from approximately three-months-old *Flaveria* plants. The cuvette temperature was maintained at 25□, the PPFD was at ~500 µmol/m²s and the CO_2 concentration was at around 400 ppm. Chamber humidity was matched to ambient air humidity. Leaf-to-air vapor pressure difference (VPD) was maintained around 1.1 kPa. Before the g_s measurements, *Flaveria* plants were maintained under constant light with a PPFD of ~500 µmol/m^{2}s until g_s reached a steady state. External environments during the g_s measurements for rice and maize were controlled to be their growth conditions. In maize and rice, the16-day-old third leaves which were the youngest mature leaves, were used for gas exchange measurements (*69*).

Phenotyping stomatal traits and the calculation of g$_{smax}$

The youngest fully expanded leaves (usually the third or fourth leaves from the apex in *Flaveria*) at the main stem were used to phenotype stomata related traits. In *Flaveria* and maize, stomata related traits were observed using both nail polish method and epidermis directly peeled from leaves (*6, 80*). For the first method, nail polish was painted on both leaf surfaces to produce the impressions of epidermis. After the nail polish was dried, the nail polish was peeled off the leaf and placed on a microscope slide for examination. For the second method, leaf epidermis was peeled off manually and transferred onto microscope slide. For the count of SD in Arabidopsis, the rosette leaves was soaked in 9:1 (V/V) ethanol:acetic acid overnight, washed with water and cleared in chloral hydrate solution (glycerol:chloral hydrate:water, 1:8:1) overnight, and then stomata of Arabidopsis were stained with Safranin-O, finally leaves were placed on a microscope slide for imaging. Light microscope (smarte; CNOPTEC), camera and image processed software (OPTPRO;

388 CNOPTEC) were used to observe and image epidermis at 100x magnification in

389 *Flaveria*. Center of a leaf was chosen to count stomatal density and measure stomatal

390 length. Five stomata randomly chosen along a diagonal of the varnish peels or

391 epidermis were measured for stomatal length.

392   G$_{smax}$ was calculated with the follow equation (*16*):

393
$$g_{smax} = \frac{d}{v} \cdot D \cdot a_{max} / (e + \frac{\pi\sqrt{a_{max}/\pi}}{2})$$

394   The constants *d* is the diffusivity of H$_2$O in air; the constant *v* is the molar

395 volume of air; D is the stomatal density; a_{max} is the maximum stomatal pore area

396 which is calculated as $a_{max} = \pi p^2 / 4$ in *Flaveria* species, and the *p* is the stomatal

397 pore length; *e* is the stomatal pore depth, which is known as equal to guard cell width.

RNA isolation, semiquantitative RT-PCR and quantitative real-time PCR

The young leaf samples taken from the main stem in *Flaveria* or the developing

rosette leaves in Arabidopsis were immediately frozen in liquid nitrogen and used

directly for RNA extraction. Total RNA was extract from the young leaves using the

PureLink RNA Mini kit (Thermo Fisher Scientifc). The first-strand cDNA was

synthesized by the TransScript One-step gDNA Removal and cDNA Synthesis

SuperMix (TRANSGEN). Semiquantitative RT-PCR was performed with Phanta®

Max Super-Fidelity DNA Polymerase (vazyme). The actin8 was chosen as the loading

control. Real-time PCR was performed on the CFX connect^{TM} (Bio-Rad) with the

UNICON^{TM} qPCR SYBR ®Green Master Mix (YEASEN) according to the

manufacturer's instructions. The calculation of expression according to the $2^{-\Delta\Delta CT}$.

The expression stability of internal reference genes for the quantitative real-time

PCR was assessed by the software: geNORM and NormFinder (*81, 82*). Specifically,

we first calculated the coefficient of variations (CV) of all genes in different *Flaveria*

species. After that, all genes were sorted according to the calculated CV values. 150

genes with the lowest CV were selected as potential candidate reference genes.

Among these, those with mean expression level >300 were chosen as candidate

reference genes. We also included those commonly used house-keeping genes, which

include *UBC9* (AT4G27960), *UBQ11* (AT4G05050), *ACT7* (AT5G09810), *GAPDH*

(AT1G13440), *EF1a* (AT5G60390), *CACMS* (AT5G46630) (Supplementary fig. 6a, b,

d), as the internal reference genes in this study. The specificity of the primers

(Supplementary table 2) for all these candidate reference genes were identified by

agarose electrophoresis (Supplementary fig. 6c). With Genorm (Supplementary fig. 6e)

and Normfinder (Supplementary fig. 6f), we concluded that *ACT7*, *EF1a* and *HEME2*

showed stable expression levels across *rob*, *ram* and *bid*. *EF1a* and *ACT7* are classic

reference genes, and they showed minor variations under different conditions. The

geometrical means of the expression level of *ACT7* and *EF1a* were finally used as

normalization factors. All primers used for RT-qPCR are listed in Supplementary table

2. For each gene, three technical replicates and at least three biological replicates were

performed.

Vector construction and transformation

A list of primer sequences were used for plasmid construction (Supplementary

table 2). The *STOMAGEN* and *FSTOMAGEN* DNA sequences, named ATSTO-cDNA

and FloSTO-cDNA respectively, were PCR-amplified from cDNA in Arabidopsis and

rob. To obtain the AFSTO fragment, the signal peptide region of *STOMAGEN* was

cloned from ATSTO-cDNA by PCR-amplification, and the functional region of

FSTOMAGEN was PCR-amplified from FloSTO-cDNA. The signal peptide and

functional region were combined by fusion PCR. To obtain an intact *FSTOMAGEN*

(FSTO) gene, the whole gene in *rob*, including the signal peptide, propeptide and the

STOMAGEN functional region, was amplified from the cDNA of *rob*. To over-express

AFSTO and FSTO, the AFSTO and FSTO were cloned separately to a pcambia-3300

binary vector which has a CaMV 35S promoter. This construct was introduced into *A.*

tumefaciens strain GV3101, transformed into Arabidopsis using a floral dipping

procedure. Transgenic lines were selected on soil that had been treated with a diluted

solution of phosphinothricin (diluted 3000 times) over night. About a month after the

seeds were sowed on the soil, the stomata were observed on the biggest mature rosette

leaf. The phenotype of Arabidopsis T1 and T2 generations was observed at the 400x

magnification using the light microscope. For each epidermal peel, five fields were

counted for density.

Refolding of synthetic *FSTOMAGEN*

Chemically synthesized peptide (Synthesized peptide sequence: IGSVKPTCTY

NECRGCRSRCRAEQVPVEGNDPINSAYHYRCICHR, Sangon Biotech, Shanghai,

China) was dissolved in 20 mM Tris-HCl, pH 8.8, and 50 mM NaCl and then dialysed

(Sangon Biotech,0.5 KD-1 KD) for 1 day at 4□ against 0.5 mM glutathione (reduced

and oxidized forms, Macklin) and 200 mML-arginine (meilunbio) at pH 8.0, which

were further dialysed three times against 20 mM Tris-HCl, pH 8.8, and 50 mM NaCl

for 1.5 days to remove glutathione.

Stomata induction assay

When the first true leaf or cotyledons appeared in *bid* and Arabidopsis that had

germinated in 1/2 MS sterilized solid medium, F*STOMAGEN* peptide was applied on

the plants. After the *bid* and Arabidopsis were further incubated in 1/2 MS liquid

medium at 22℃ for 5 days, stomatal numbers on the abaxial surfaces for the first true

leaf and cotyledons were counted under a differential interference contrast microscope

(DIC)**.**

Plotting and statistical analysis

The correlation, linear regressions and t-test (a=0.05) were performed by

GraphPad Prism 8. Differences between means of multi-groups were assessed with

one-way analysis of variance (ANOVA). Duncan's multiple range test was performed

using the agricolae package of the R (https://www.r-project.org/) to detect differences

between two groups. The A and g_s at the saturated light intensity were digitized from

figure 1 and figure 3 in (*4, 83*) using the OriginPro 2018.

Accession numbers

The *Flaveria* read data have been submitted to the National Center for Biotechnology

Information Short Read Archive under accession numbers: SRP036880, SRP036881,

SRP036883, SRP036884, SRP036885, SRP037526, SRP037527, SRP037528,

SRP037529, SRP006166, SRP169568. The *Flaveria* RNA-Seq datasets from 1KP

Consortium are available at National Center for Biotechnology Information (NCBI)

Gene Expression Omnibus (GEO) under accession number: GSE54339. The rice and

maize RNA-seq datasets under accession number: GSE54274. Accession numbers of

genes in this study was shown in Supplementary table 3.

Discussion

Detailed characterization of stomatal traits in the *Flaveria* genus along C_4

evolution

Many recent studies have shown that stomatal change may play an important role

in the evolution of C_4 photosynthesis (*2, 3, 17*). The short-term regulation of stomata

function mainly involves the dynamic opening and closing (*84*). But on an

evolutionary scale, the adjustment of stomatal function mainly involves changes in

g_{smax} i.e. changes in stomatal density and stomatal size (*14-17*). The g_{smax} in *Flaveria* is

significantly correlated with the operating g_s (Supplementary fig 1b). In order to study

the stomatal evolution along C_4 evolution, we used the *Flaveria* genus, which

includes species with different photosynthetic pathways. Species in *Flaveria* also has

a close evolutionary distance, which can reduce the effects of factors other than

photosynthetic types on stomatal traits. In addition, *Flaveria* represents a C_4

evolutionary model that minimizes differences in habitat (6, 52, 53). The short

phylogenetic distance and similar growth habitats enabled us to better detect the

difference in stomatal traits between species with different photosynthetic pathways

(Fig 1, Fig 2). With data from species in the *Flaveria* genes, we consolidated the

earlier notion the g_{smax} is associated with photosynthetic pathway and there is a

decrease in g_{smax} and *SD* in C_4 plants compared to C_3 plants (*17*). Our data further

showed that the decrease in g_{smax} and *SD* also occurred in dicotyledon and this

decrease in *SD* is due to less stomata generation (Fig 2h). Photosynthesis and stomata

are closely related (*85, 86*), our work provide an excellent example for the co-
evolution between photosynthetic pathway and stomata.

We showed that the changes of stomata traits along the path to C_4 is gradual, not
abrupt (Fig 1, Fig 2). The stomata traits initially changed near the Type II stage, and
dramatically changed at the C_4-like stage (Fig 1, Fig 2). Previous studies have shown
that in C_4 grasses the decrease in *SD* might be due to the restriction of stomatal
development caused by the increased vein density (*17, 19*). In this study, we found no
significant change in stomatal traits in early stage in *Flaveria* (Fig 1, Fig2), while the
vein density at this stage, *e.g.* in *F. sonorensis*, was already increased (*43*). Therefore,
the decreased *SD* in C_4 evolution was not necessarily due to the constrained vein
development.

Along the road to C_4, stomata become less in number (lower SD, SI) and larger in
size (longer L) (Fig 2g). The opposite trends in *SD* and *SS* changes have been
repeatedly shown in different conditions (*14, 16*). The opposite trend may help
maintain relative constant allocation of epidermal space to stomata (*14, 16*).
Transgenic experiments of genes related to stomatal development have shown that *SS*
and *SD* are related, showing that changes in *SD* can directly cause opposite changes in
SS (*80, 87*). Therefore, the opposite trends of changes in *SD* and *SS* could be caused
by the stomatal development related genes, which is consistent with the changes in the
expression of *FSTOMAGEN* (Fig 5 b,c,d). Previous studies have shown that the *SS* is
related to the size of genome (*88*). Since the genome sizes increased gradually along
C_4 evolution in *Flaveria* (54), the increased genome size might be one reason for the
increased *SS* during C_4 evolution in this genus.

The role of decreased g_{smax} during C_4 photosynthesis evolution in *Flaveria*

Though water limitation might have been regarded as a primary driver for the C_4
evolution (*5-7*), data from this study suggests that CO_2 might have been the initial
driver for C_4 evolution. In the *F. sonorensis* and *F. angustifolia*, two Type I
intermediate *Flaveria* species, g_{smax} were still similar to those in C_3 *Flaveria* species,
i.e. *F. cronquistii, F. pringlei* and *F.robusta* (*4*) (Fig. 1b,c,d,e) (*89*); while in contrast,

the CO_2 concentration points of these Type I species were already dramatically

decreased (*90*) compared to their C_3 ancestors, as a result of an operating CO_2

concentrating mechanism supported by the photorespiration CO_2 pump (C_2

photosynthesis) (*91*). This is consistent with the notion that at the early stage of C_4

evolution, where the movement of glycine decarboxylase (GDC) from mesophyll cell

to bundle sheath cell occurred, the primary evolutionary driver should be to elevate

CO_2 levels around Rubisco, rather than saving H_2O (*92*), i.e. the carbon uptake, not

water loss, was the major selection pressure at the early stage of C_4 evolution in this

genus (*4, 83*). Only after this initial step, reducing water loss might become crucial.

This is because the enhancement of photosynthetic capacity by C_2 or C_4

photosynthesis can increase CO_2 uptake and evapotranspiration of canopies(*4*), hence

creating a higher water demand and raising the danger of embolism, especially in the

arid, semi-arid, hot and saline areas (*3*). Indeed, we found initial decline in g_{smax}

occurred near Type II (Fig 1b, c, d, e).

The gradually decline of g_{smax} is consistent with the gradual establishment of

photorespiratory C_2 pump and the shifting of photosynthetic facilities from mesophyll

tissue to bundle sheath tissue. The decrease of g_{smax} represents the negative regulation

on intercellular CO_2 concentration. The higher intercellular CO_2 concentration could

diminish the role of photorespiration CO_2 pump and photosynthetic capacity in bundle

sheath. Indeed, under higher CO_2, intermediate species will have no photosynthetic

advantage over C_3 plants (*89, 93*). Therefore, it is possible that the gradual decrease of

g_{smax} can help consolidate and strengthen C_2 photosynthesis during C_4 evolution. The

photorespiration CO_2 pump further magnified the ammonia imbalance, hence

promoting the evolution of C_4 metabolism, as in the case in Type II (*41*). Although the

C_4 photosynthesis is incomplete and the activities of C_4 enzymes are low in Type II

(*90, 94*), the increased C_4 cycle might alleviate the decrease in photosynthetic rate

caused by the decreased g_{smax}. In C_4-like and C_4 species, a complete and highly active

C_4 cycle (*90*), in particular caused by the increased expression levels and enzymes

activities of CA and PEPC, can enable maintenance of high photosynthetic rate even

though the reduced intercellular CO_2 concentration (*2, 4, 17*), which allows g_{smax}

further decreas. This is consistent with the earlier finding that when the limitation on

photosynthetic rate by CO_2 supply becomes less, g_{smax} dominated by SD will reduce

(*16*).

C_4 evolution select *STOMAGEN* homologs as a molecular trigger to alter g_{smax}

Stomatal development is controlled by a molecular signaling network composed

of many interacting components (*28*). In theory, any gene controlling stomatal

development could be potentially selected to alter stomatal patterning and affect g_{smax}

during the emergence of C_4 photosynthesis (*15, 80*). In fact, the alterations of stomata

patterning under different conditions are indeed caused by different molecular

pathways (*24, 95, 96*). Here we show that during C_4 evolution, the decreased

expression of *FSTOMAGEN* (Fig 3c,e, Fig 5b,c,d), a *STOMAGEN*-like gene, is

selected to decrease SD and g_{smax} in the *Flaveria* genus and likely in other lineage as

well (Fig 6). We used multiple evidences to support this notion. In our experiments,

FSTOMAGEN strongly and positively regulates stomatal development when either

synthetic FSTOMAGEN peptide was applied *in vitro* or *FSTOMAGEN* was over-

expressed *in vivo* (Fig 4). Besides *FSTOMAGEN*, we do notice that other genes

related to stomatal development were also differentially expressed between C_3 and C_4.

However, these genes showed either less changes or the pattern of changes did not

result in the observed changes in SD, as in the case of *EPF1* and *TMM* homologs (Fig

3c,d). For example, *EPF1* homologs negatively regulates stomatal development (*33,*

97-99), but its expression is lower in C_4, which contradicts with the lower SD in C_4.

The decrease in *TMM* and *EPF1* expression levels in C_4 may be due to the co-

regulation and feedback of these stomatal regulators, as previous work shows that

overexpression of *STOMAGEN* can increase the expression of other stomata

regulatory genes (*59*).

The decreased *FSTOMATGEN* expression might be a major factor for the

increased iWUE. The iWUE of C_4 relative to that of C_3 increased significantly, partly

contributed by a lower g_s (Supplementary fig 1c,d) (*4*). Since the magnitude of the

decrease in g_s is similar to the magnitude of the decrease in g_{smax}, the increased iWUE

of C$_4$ species can be at least partially attributed to the decreased g_{smax} (Fig 1;

Supplementary fig 1c,d). Because *FSTOMAGEN* induced changes in *SD* and hence

g_{smax}, the decreased *FSTOMAGEN* expression might have contributed to the increased

iWUE in C$_4$ plants. This is significant for improving iWUE in C$_4$ engineering. Many

studies suggest that genes related to stomatal development not only affect stomatal

development (*100, 101*), but also affect photosynthetic apparatus and the development

of photosynthetic tissues (*100*). Therefore, identifying genes that can alter stomata

without negatively influencing other parts in C$_4$ engineering is required. Results from

this study suggests that *FSTOMAGEN* is one such gene, since *FSTOMAGEN* was

already selected during the evolution of C$_4$ photosynthesis.

Reference

1. F. I. W. Alistair M. Hetherington, The role of stomata in sensing and driving environmental
change.pdf. *Nature* **424**, 901-908 (2003).

2. S. H. H. Taylor, S. P. Rees, M Ripley, B. S. Woodward, F. I. Osborne, C. P., Ecophysiological traits
in C$_3$ and C$_4$ grasses: a phylogenetically controlled screening experiment. *New Phytol* **185**,
780-791 (2010).

3. C. P. Osborne, L. Sack, Evolution of C$_4$ plants: a new hypothesis for an interaction of CO$_2$ and
water relations mediated by plant hydraulics. *Philos Trans R Soc Lond B Biol Sci* **367**, 583-600
(2012).

4. P. J. Vogan, R. F. Sage, Water-use efficiency and nitrogen-use efficiency of C$_3$-C$_4$ intermediate
species of Flaveria Juss. (Asteraceae). *Plant, cell & environment* **34**, 1415-1430 (2011).

5. H. Zhou, B. R. Helliker, M. Huber, A. Dicks, E. Akcay, C$_4$ photosynthesis and climate through
the lens of optimality. *Proc Natl Acad Sci U S A* **115**, 12057-12062 (2018).

6. G. Reeves *et al.*, Natural Variation within a Species for Traits Underpinning C$_4$ Photosynthesis.
Plant physiology **177**, 504-512 (2018).

7. B. P. Williams, I. G. Johnston, S. Covshoff, J. M. Hibberd, Phenotypic landscape inference
reveals multiple evolutionary paths to C$_4$ photosynthesis. *elife* **2**, e00961 (2013).

8. S. H. Taylor, B. S. Ripley, F. I. Woodward, C. P. Osborne, Drought limitation of photosynthesis
differs between C(3)and C(4)grass species in a comparative experiment. *Plant, cell &*
environment **34**, 65-75 (2011).

9. F. Kocacinar, Photosynthetic, hydraulic and biomass properties in closely related C3and
C4species. *Physiologia Plantarum* **153**, 454-466 (2015).

10. B. Uzilday, I. Turkan, R. Ozgur, A. H. Sekmen, Strategies of ROS regulation and antioxidant
defense during transition from C(3) to C(4) photosynthesis in the genus Flaveria under PEG-
induced osmotic stress. *J Plant Physiol* **171**, 65-75 (2014).

11. S. A. Aubry, O. Reyna-Llorens, I. Smith-Unna, R. D. Hibberd, J. M. Genty, B., A Specific
Transcriptome Signature for Guard Cells from the C_4 Plant Gynandropsis gynandra. *Plant*
physiology **170**, 1345-1357 (2016).

12. R. F. Sage, The evolution of C_4 photosynthesis. *New Phytologist* **161**, 341-270 (2004).

13. A. M. Powell, Systematics of Flaveria (Flaveriinae--Asteraceae). *Annals of the Missouri*
Botanical Garden **65**, 590-636 (178).

14. P. J. Franks, P. L. Drake, D. J. Beerling, Plasticity in maximum stomatal conductance
constrained by negative correlation between stomatal size and density: an analysis using
Eucalyptus globulus. *Plant, cell & environment* **32**, 1737-1748 (2009).

15. T. Doheny-Adams, L. Hunt, P. J. Franks, D. J. Beerling, J. E. Gray, Genetic manipulation of
stomatal density influences stomatal size, plant growth and tolerance to restricted water
supply across a growth carbon dioxide gradient. *Philos Trans R Soc Lond B Biol Sci* **367**, 547-
555 (2012).

16. P. J. F. a. D. J. Beerling, Maximum leaf conductance driven by CO_2 effects on stomatal size and
density over geologic time. *PNAS* **106**, 10343–10347 (2009).

17. S. H. Taylor *et al.*, Photosynthetic pathway and ecological adaptation explain stomatal trait
diversity amongst grasses. *New Phytol* **193**, 387-396 (2012).

18. W. M. Laetsch, The C4' syndrome: a structural analysis. *Ann. Rev. Plant Physiol.* **25**, 27-52
(1974).

19. D. A. Way, What lies between : the evolution of stomatal traits on the road to C4
photosynthesis. *New phytologist* **193**, 291-293 (2012).

20. T. W. Ocheltree, J. B. Nippert, P. V. Prasad, Changes in stomatal conductance along grass
blades reflect changes in leaf structure. *Plant, cell & environment* **35**, 1040-1049 (2012).

21. V. S. R. D. a. M. Santakumari, Stomatal characteristics of some dicotyledonous plants in
relation to the C_4 and C_3 pathways of photosynthesis. *Plant and cell physiology* **18**, 935-938
(1977).

22. E. J. Edwards, C. J. Still, Climate, phylogeny and the ecological distribution of C4 grasses. *Ecol*
Lett **11**, 266-276 (2008).

23. E. J. Edwards, C. J. Still, M. J. Donoghue, The relevance of phylogeny to studies of global
change. *Trends Ecol Evol* **22**, 243-249 (2007).

24. E. T. Hamanishi, B. R. Thomas, M. M. Campbell, Drought induces alterations in the stomatal
development program in Populus. *J Exp Bot* **63**, 4959-4971 (2012).

25. Q. Cai, C. Ji, Z. Yan, X. Jiang, J. Fang, Anatomical responses of leaf and stem of Arabidopsis
thaliana to nitrogen and phosphorus addition. *J Plant Res* **130**, 1035-1045 (2017).

26. Z. Xu, G. Zhou, Responses of leaf stomatal density to water status and its relationship with
photosynthesis in a grass. *J Exp Bot* **59**, 3317-3325 (2008).

27. N. Sekiya, K. Yano, Stomatal density of cowpea correlates with carbon isotope discrimination
in different phosphorus, water and CO2 environments. *New Phytol* **179**, 799-807 (2008).

28. N. Zoulias, E. L. Harrison, S. A. Casson, J. E. Gray, Molecular control of stomatal development.
Biochem J **475**, 441-454 (2018).

29. G. Lin *et al.*, A receptor-like protein acts as a specificity switch for the regulation of stomatal
development. *Genes Dev* **31**, 927-938 (2017).

30. E. D. Shpak, J. M. McAbee, L. J. Pillitteri, K. U. Torii, Stomatal patterning and differentiation by
synergistic interactions of receptor kinases. *Science* **309**, 290-293 (2005).

31. J. S. Lee *et al.*, Direct interaction of ligand-receptor pairs specifying stomatal patterning.
Genes Dev **26**, 126-136 (2012).
32. T. Shimada, S. S. Sugano, I. Hara-Nishimura, Positive and negative peptide signals control
stomatal density. *Cell Mol Life Sci* **68**, 2081-2088 (2011).
33. K. Hara, R. Kajita, K. U. Torii, D. C. Bergmann, T. Kakimoto, The secretory peptide gene EPF1
enforces the stomatal one-cell-spacing rule. *Genes Dev* **21**, 1720-1725 (2007).
34. L. Hunt, J. E. Gray, The signaling peptide EPF2 controls asymmetric cell divisions during
stomatal development. *Curr Biol* **19**, 864-869 (2009).
35. L. J. Pillitteri, K. U. Torii, Breaking the silence: three bHLH proteins direct cell-fate decisions
during stomatal development. *Bioessays* **29**, 861-870 (2007).
36. S. S. Sugano *et al.*, Stomagen positively regulates stomatal density in Arabidopsis. *Nature* **463**,
241-244 (2010).
37. T. Kondo *et al.*, Stomatal density is controlled by a mesophyll-derived signaling molecule.
Plant Cell Physiol **51**, 1-8 (2010).
38. L. Hunt, K. J. Bailey, J. E. Gray, The signalling peptide EPFL9 is a positive regulator of stomatal
development. *New Phytol* **186**, 609-614 (2010).
39. U. Schluter, A. P. Weber, The Road to C4 Photosynthesis: Evolution of a Complex Trait via
Intermediary States. *Plant Cell Physiol* **57**, 881-889 (2016).
40. D. Heckmann *et al.*, Predicting C4 photosynthesis evolution: modular, individually adaptive
steps on a Mount Fuji fitness landscape. *Cell* **153**, 1579-1588 (2013).
41. J. Mallmann *et al.*, The role of photorespiration during the evolution of C_4 photosynthesis in
the genus Flaveria. *Elife* **3**, e02478 (2014).
42. M. J. Lyu *et al.*, RNA-Seq based phylogeny recapitulates previous phylogeny of the genus
Flaveria (Asteraceae) with some modifications. *BMC Evol Biol* **15**, 116 (2015).
43. A. D. M. A. N. G. DENGLER, Key innovations in the evolution of Kranz anatomy and C4 vein
pattern in Flaveria (Asteraceae). *American Journal of Botany* **94**, 382–399 (2007).
44. S. Schulze *et al.*, Evolution of C4 photosynthesis in the genus flaveria: establishment of a
photorespiratory CO2 pump. *Plant Cell* **25**, 2522-2535 (2013).
45. M. Stata *et al.*, Mesophyll Chloroplast Investment in C3, C4 and C2 Species of the Genus
Flaveria. *Plant Cell Physiol* **57**, 904-918 (2016).
46. P. A. Christin, C. P. Osborne, R. F. Sage, M. Arakaki, E. J. Edwards, C(4) eudicots are not
younger than C(4) monocots. *J Exp Bot* **62**, 3171-3181 (2011).
47. U. Gowik, A. Bräutigam, K. L. Weber, A. P. M. Weber, P. Westhoff, Evolution of C_4
Photosynthesis in the Genus Flaveria: How Many and Which Genes Does It Take to Make C_4?
The Plant Cell **23**, 2087-2105 (2011).
48. A. D. MCKOWN, PHYLOGENY OF FLAVERIA (ASTERACEAE) AND INFERENCE OF C4
PHOTOSYNTHESIS EVOLUTION. *American Journal of Botany* **92**, 1911–1928 (2005).
49. E. A. Sudderth, R. M. Muhaidat, A. D. McKown, F. Kocacinar, R. F. Sage, Leaf anatomy, gas
exchange and photosynthetic enzyme activity in Flaveria kochiana. *Funct Plant Biol* **34**, 118-
129 (2007).
50. E. A. Sudderth, F. J. Espinosa-García, N. M. Holbrook, Geographic distributions and
physiological characteristics of co-existing Flaveria species in south-central Mexico. *Flora -
Morphology, Distribution, Functional Ecology of Plants* **204**, 89-98 (2009).

51.   F. Kocacinar, A. D. McKown, T. L. Sage, R. F. Sage, Photosynthetic pathway influences xylem
structure and function in Flaveria (Asteraceae). *Plant, cell & environment* **31**, 1363-1376
(2008).

52.   N. Nakamura, M. Iwano, M. Havaux, A. Yokota, Y. N. Munekage, Promotion of cyclic electron
transport around photosystem I during the evolution of NADP-malic enzyme-type C4
photosynthesis in the genus Flaveria. *New Phytol* **199**, 832-842 (2013).

53.   R. F. Sage, A portrait of the C4 photosynthetic family on the 50th anniversary of its discovery:
species number, evolutionary lineages, and Hall of Fame. *J Exp Bot* **68**, 4039-4056 (2017).

54.   Y. Y. Taniguchi *et al.*, Dynamic changes of genome sizes and gradual gain of cell-specific
distribution of C4 enzymes during C4 evolution in genus Flaveria. *Plant Genome*, e20095
(2021).

55.   R. F. Sage, R. Khoshravesh, T. L. Sage, From proto-Kranz to C4 Kranz: building the bridge to C4
photosynthesis. *J Exp Bot* **65**, 3341-3356 (2014).

56.   S. R. Christopher M. Hylton, Alison M. Smith, D. Alan Jones and Harold W. Woolhouse, Glycine
decarboxylase is confined to the bundle-sheath cells of leaves of C3-C4 intermediate species.
planta **175**, 452-459 (1988).

57.   L. J. Pillitteri, K. U. Torii, Mechanisms of stomatal development. *Annu Rev Plant Biol* **63**, 591-
614 (2012).

58.   S. Ohki, M. Takeuchi, M. Mori, The NMR structure of stomagen reveals the basis of stomatal
density regulation by plant peptide hormones. *Nat Commun* **2**, 512 (2011).

59.   M. Hronkova *et al.*, Light-induced STOMAGEN-mediated stomatal development in
Arabidopsis leaves. *J Exp Bot* **66**, 4621-4630 (2015).

60.   J. S. Lee *et al.*, Competitive binding of antagonistic peptides fine-tunes stomatal patterning.
Nature **522**, 439-443 (2015).

61.   J. Lu *et al.*, Homologous genes of epidermal patterning factor regulate stomatal development
in rice. *J Plant Physiol* **234-235**, 18-27 (2019).

62.   L. Xuxin, The Effects for Stomata Developmental on Arabidopsis by Overexpression of Rice
EPFL Family Genes. *Molecular Plant Breeding* **15**, 2042- 2047 (2017).

63.   X. Yin *et al.*, CRISPR-Cas9 and CRISPR-Cpf1 mediated targeting of a stomatal developmental
gene EPFL9 in rice. *Plant Cell Rep* **36**, 745-757 (2017).

64.   J. Y. Zhang, S. B. He, L. Li, H. Q. Yang, Auxin inhibits stomatal development through
MONOPTEROS repression of a mobile peptide gene STOMAGEN in mesophyll. *Proc Natl Acad*
Sci U S A **111**, E3015-3023 (2014).

65.   M. A. Lyu *et al.*, The coordination of major events in C4 photosynthesis evolution in the genus
Flaveria. *Sci Rep* **11**, 15618 (2021).

66.   K. Billakurthi *et al.*, Transcriptome dynamics in developing leaves from C_3 and C_4 Flaveria
species reveal determinants of Kranz anatomy. *bioRxiv*, (2018).

67.   M. G. Grabherr *et al.*, Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat Biotechnol* **29**, 644-652 (2011).

68.   B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

69.   L. Wang *et al.*, Comparative analyses of C_4 and C_3 photosynthesis in developing leaves of
maize and rice. *Nat Biotechnol* **32**, 1158-1165 (2014).

70.   X. Robert, P. Gouet, Deciphering key features in protein structures with the new ENDscript
763   server. *Nucleic Acids Res* **42**, W320-324 (2014).
764   71.   D. R. F. Kyte J, A simple method for displaying the hydropathic character of a protein. *Journal*
765   *of Molecular Biology* **157**, 105-132 (1982).
766   72.   S. Perveen *et al.*, Overexpression of maize transcription factor mEmBP-1 increases
767   photosynthesis, biomass, and yield in rice. *J Exp Bot* **71**, 4944-4957 (2020).
768   73.   D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
769   dramatically improves orthogroup inference accuracy. *Genome Biol* **16**, 157 (2015).
770   74.   R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high throughput.
771   *Nucleic Acids Res* **32**, 1792-1797 (2004).
772   75.   A. Stamatakis, RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
773   thousands of taxa and mixed models. *Bioinformatics* **22**, 2688-2690 (2006).
774   76.   F. Ronquist *et al.*, MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice
775   across a large model space. *Syst Biol* **61**, 539-542 (2012).
776   77.   S. A. Shiryev, J. S. Papadopoulos, A. A. Schaffer, R. Agarwala, Improved BLAST searches using
777   longer words for protein seeding. *Bioinformatics* **23**, 2949-2951 (2007).
778   78.   X. J. Min, G. Butler, R. Storms, A. Tsang, OrfPredictor: predicting protein-coding regions in
779   EST-derived sequences. *Nucleic Acids Res* **33**, W677-680 (2005).
780   79.   A. L. Ming-Ju *et al.*, The Coordination and Jumps along C$_4$ Photosynthesis Evolution in the
781   Genus Flaveria. *bioRxiv*, 460287 (2018).
782   80.   P. J. Franks, W. Doheny-Adams T, Z. J. Britton-Harper, J. E. Gray, Increasing water-use
783   efficiency directly through genetic manipulation of stomatal density. *New Phytol* **207**, 188-
784   195 (2015).
785   81.   J. L. J. a. T. F. Ø. Claus Lindbjerg Andersen, Normalization of real-time quantitative reverse
786   transcription-PCR data a model-based variance estimation approach to identify genes suited
787   for normalization, applied to bladder and colon cancer data sets. *Cancer Res* **64**, 5245-5250
788   (2004).
789   82.   D. P. K. Vandesompele J, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F., Accurate
790   normalization of real-time quantitative RT-PCR data by geometric averaging of multiple
791   internal control genes. *Genome Biology* **3**, research0034.0031–0034.0011 (2002).
792   83.   D. A. Way, G. G. Katul, S. Manzoni, G. Vico, Increasing water use efficiency along the C$_3$ to C$_4$
793   evolutionary pathway: a stomatal optimization perspective. *J Exp Bot* **65**, 3683-3693 (2014).
794   84.   A. V. a. D. C. Bergmann, Mechanisms of stomatal development: an evolutionary view.
795   *EvoDevo* **3**, (2012).
796   85.   P. J. Franks, D. J. Beerling, CO(2)-forced evolution of plant gas exchange capacity and water-
797   use efficiency over the Phanerozoic. *Geobiology* **7**, 227-236 (2009).
798   86.   I. R. C. G. D. F. S. C. Wong, Stomatal conductance correlates with photosynthetic capacity.
799   *Nature* **282**, 424-426 (1979).
800   87.   Y. Tanaka, S. S. Sugano, T. Shimada, I. Hara-Nishimura, Enhancement of leaf photosynthetic
801   capacity through increased stomatal density in Arabidopsis. *New Phytol* **198**, 757-764 (2013).
802   88.   J. M. Beaulieu, I. J. Leitch, S. Patel, A. Pendharkar, C. A. Knight, Genome size is a strong
803   predictor of cell size and stomatal density in angiosperms. *New Phytol* **179**, 975-986 (2008).

89.   P. J. Vogan, M. W. Frohlich, R. F. Sage, The functional significance of C_3-C_4 intermediate traits
in Heliotropium L. (Boraginaceae): gas exchange perspectives. *Plant, cell & environment* **30**,
1337-1345 (2007).
90.   J. W. Maurice S. B. Ku, Ziyu Dai, Rick A. Scott, Chun Chu, and Gerald E. Edwards,
Photosynthetic and Photorespiratory Characteristics of Flaveria species. *plant physiol* **96**, 518-
528 (1991).
91.   O. Keerberg, T. Parnik, H. Ivanova, B. Bassuner, H. Bauwe, C_2 photosynthesis generates about
3-fold elevated leaf CO_2 levels in the C_3-C_4 intermediate species Flaveria pubescens. *J Exp Bot*
65, 3649-3656 (2014).
92.   R. F. Sage, T. L. Sage, F. Kocacinar, Photorespiration and the evolution of C_4 photosynthesis.
Annu Rev Plant Biol **63**, 19-47 (2012).
93.   P. J. Vogan, R. F. Sage, Effects of low atmospheric CO2 and elevated temperature during
growth on the gas exchange responses of C3, C3-C4 intermediate, and C4 species from three
evolutionary lineages of C4 photosynthesis. *Oecologia* **169**, 341-352 (2012).
94.   R. K. M. MAURICE S. B. Ku, ROBERT. LITTLEJOHN, JR., HITOSHI NAKAMOTO, DONALD B.
FISHER, AND GERALD E. EDWARDS, Photosynthetic Characteristics of C3-C4 Intermediate
Flaveria Species. *Plant Physiol.* **71**, 944-948 (1983).
95.   C. B. Engineer *et al.*, Carbonic anhydrases, EPF2 and a novel protease mediate CO_2 control of
stomatal development. *Nature* **513**, 246-250 (2014).
96.   Y. Tanaka, T. Nose, Y. Jikumaru, Y. Kamiya, ABA inhibits entry into stomatal-lineage
development in Arabidopsis leaves. *Plant J* **74**, 448-457 (2013).
97.   C. Wang *et al.*, PdEPF1 regulates water-use efficiency and drought tolerance by modulating
stomatal density in poplar. *Plant Biotechnol J* **14**, 849-860 (2016).
98.   R. S. Caine *et al.*, Rice with reduced stomatal density conserves water and has improved
drought tolerance under future climate conditions. *New Phytol*, (2018).
99.   J. Hughes *et al.*, Reducing Stomatal Density in Barley Improves Drought Tolerance without
Impacting on Yield. *Plant physiology* **174**, 776-787 (2017).
100. G. J. Dow, J. A. Berry, D. C. Bergmann, Disruption of stomatal lineage signaling or
transcriptional regulators has differential effects on mesophyll development, but maintains
coordination of gas exchange. *New Phytol* **216**, 69-75 (2017).
101. H. Cui *et al.*, SPINDLY, ERECTA, and its ligand STOMAGEN have a role in redox-mediated cortex
proliferation in the Arabidopsis root. *Mol Plant* **7**, 1727-1739 (2014).
102. M. Stata, T. L. Sage, R. F. Sage, Mind the gap: the evolutionary engagement of the C4
metabolic cycle in support of net carbon assimilation. *Curr Opin Plant Biol* **49**, 27-34 (2019).

Acknowledgements

We thank Prof. Colin Osborne and Julie Gray for constructive comments on our
revised manuscript. The work was funded by Strategic Priority Research Program of
the Chinese Academy of Sciences (grant number: XDB27020105) and the general
program of National Science Foundation of China (31870214). We thank Prof. Rowan

F. Sage (University of Toronto) for providing the *Flaveria* cuttings. We thank Dr. Matt

stata for helping cutting of *Flaveria*.

Author contributions

YYZ and XGZ designed the study. YYZ observed and measured the stomatal

characteristic. YYZ carried out the molecular experiment and the treatment of

synthetic peptides in vivo. FFM conducted the transgene experiment. GYC guided the

treatment of peptides in vivo. Amy MJL analyzed the one kp RNA-seq data. YYZ and

XGZ interpreted results and wrote the article.

Competing interests

The authors declare no competing financial interests.

Figure Legend

Fig 1: Maximum stomatal conductance (g_{smax}) of *Flaveria* species

a, Phylogeny of the Flaveria genus according to (*42, 48*). Different colors represent

different photosynthetic types, orange indicates C_3, purple indicates Type I, gray

indicates Type II, green indicates C_4-like, blue indicates C_4. **b,c**, The trend of g_{smax} on

the abaxial surface in *Flaveria* species with different photosynthetic types from C_3 to

C_4 grown outdoors (**b**) and in greenhouse (**c**). **d,e**, The difference in g_{smax} of the

abaxial surface between species with different photosynthetic types in the *Flaveria*

genus grown outdoors (**d**) and in greenhouse (**e**). **f**, Correlation of g_{smax} and g_s for

Flaveria species with different photosynthetic types. The data of *gs* was from (*4*).

Error bar indicates s.d. The asterisks represent statistically significant differences,

One-way ANOVA: *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, ****, $P < 0.0001$;

Different letters represent statistically significant differences, Duncan's multiple range

test ($\alpha = 0.05$). Pearson correlation coefficient and the significance were noted in each

panel.

Fig 2: Stomatal density (*SD*) and its patterning.

a,c, The trend of *SD* of abaxial surface from C_3 to C_4 in *Flaveria* species grown

outdoors (**a**) and in greenhouse (**c**). **b,d**, The difference in *SD* and guard cell length (l)

of abaxial surface between species with different photosynthetic types in *Flaveria*

genus grown outdoors (**b**) and in greenhouse (**d**). **e,** Correlation of g_{smax} and *SD* for

different *Flaveria* species. Different color indicates different photosynthetic type. **f**,

Correlation of g_{smax} and *l* for different *Flaveria* species. Different color indicates

different photosynthetic type. **g**, Correlation of *l* and *SD* for different *Flaveria* species.

Different color indicates different photosynthetic type. **h,** The photographs of stomatal

patterning on abaxial surface in *rob*, *ram* and *bid*. The photographs from the left to

right are for *rob*, *ram* and *bid* grown outdoors (top) and in greenhouse (bottom). **i**,

Comparison of stomatal index (*SI*) between *rob*, *ram*, and *bid*. Different colours

represent plants grown outdoors (blue) or in greenhouse (green). *SD*, (n=4,

biologically independent replicates). *l*, (n=20, biologically independent replicates). *SI*,

(n=3, biologically independent replicates). Error bar indicates s.d. The asterisks

represent statistically significant differences, One-way ANOVA: *, $P < 0.05$, **, $P <$

0.01, ***, $P < 0.001$, ****, $P < 0.0001$; Different letters represent statistically

significant differences, Duncan's multiple range test ($\alpha = 0.05$). Pearson correlation

coefficient and the significance are noted in each panel.

Fig 3: Transcript abundance of stomatal developmental related genes between C_3 and

C_4 *Flaveria* species.

a, A model of the signaling pathway for the stomatal development in Arabidopsis (*28*).

A protodermal cell (PDC, white) becomes a meristemoid mother cell (MMC, orange),

which divides asymmetrically into two daughter cells, with one being a meristemoid

(M, red) and the other being a stomatal-lineage ground cell (SLGC). The M cell

develops into a guard mother cell (GMC, yellow); the SLGC develops into a

pavement cell (PC). The GMC conducts a symmetrical division to produce two equal-

sized guard cells (GC, green). EPF1, EPF2 and STOMAGEN competitively bind to

ER and TMM, which can deliver a signal to the YDA MAPK cascade. SPCH, MUTE

and FAMA ultimately are inactivated through phosphorylation by the YDA MAPK

cascade. **b**, Comparison of the expression of stomatal developmental related genes

between juvenile and mature leaves. **c**, The foldchanges of stomatal development

related genes for the C_3 and C_4 *Flaveria* species. Fold change means the ratio between

the gene expression in a C_3 *Flaveria* specie and gene expression in a C_4 *Flaveria*

specie under the same leaf development stage and growth condition as the C_3 *Flaveria*

species. The expression data were from (*66*), (*41*), (*47*) and One Thousand Plants

(1KP). **d,** The volcano plot for the expression of the stomatal development related

genes between C_3 and C_4 *Flaveria* species. The expression data were from (*41*) and

(*66*). **e**, Relative expression level determined by RT-qPCR and expression level

determined by RNA-seq of *FSTO* (*FSTOMAGEN*) in *rob* and *bid*. (n=4, biologically

independent replicates). Error bar indicates s.d. The asterisks represent statistically

significant differences ($P<0.05$, t-test, two-tailed distribution, unpaired).

Fig 4: The Function of *FSTOMAGEN* in controlling the *SD*

a, Protein conservation analysis of the homologs of FSTOMAGEN in different

species. The amino acid regions marked by green line and red line represent the signal

peptide and the functional domain of FSTOMAGEN, respectively. Red letters

represent conserved amino acids in most species, white letters with red backgrounds

represent amino acid conserved across all species. **b**, The stomatal density for

Arabidopsis leaves with FSTOMAGEN applied at different concentrations (n=8,

biologically independent replicates). **c**, The stomatal density for bid leaves with

FSTOMAGEN applied at different concentrations (n=8, biologically independent

replicates). **d**, The stomatal index (SI) for bid leaves with FSTOMAGEN applied at

different concentrations (n=6, biologically independent replicates). Error bar indicates

s.d. The asterisks represent statistically significant differences ($P<0.05$, t-test, two-

tailed distribution, unpaired). **e**, The structure of recombinant genes used to transform

Arabidopsis. AFSTO represents the recombination of the signal peptide of

STOMAGEN from Arabidopsis and the functional domain of FSTOMAGEN from

F.robusta. FSTO represents the intact STOMAGEN from *F.robusta*. **f**, **i**, The *SD* (**e**)

and RNA-level (**h**) of the different transgenic Arabidopsis (FSTO) lines. **g**, **j**, The *SD*

(**f**) and RNA-level (**i**) of the different transgenic Arabidopsis (AR) lines. **h**,

Comparison of the transgenic FSTO lines and AR lines. **k**, Photographs of stomata in

a leaf from Arabidopsis with FSTO and AFSTO over-expressed. Scar bar represents

20 um.   Error bar indicates s.d. The n.s. represent statistically no significant

differences (P>0.05, t-test, two-tailed distribution, unpaired).

Fig 5: Expression level of *FSTOMAGEN* at different stages of C_4 evolution.

a, Protein conservation analysis of the homologs of FSTOMAGEN between different

species in the *Flaveria* genus. The amino acid regions marked by green line and red

line represent the signal peptide and the functional domain of STOMAGEN,

respectively. Red letters represent conserved amino acids in most species; white

letters with red background represent amino acids conserved across all species. **b**,

Expression levels of *FSTOMAGEN* determined by RNA-seq at the different

developmental stages of leaves in *rob* and *bid*. RNA-seq data was from (*66*). P

denotes leaf primordia. Approximate leaf length for the different leaf stages was 0.3

cm (stage 1), 0.5 cm (stage 2), 1.5 cm (stage 3), 2 cm (stage 4), 3 cm (stage 5), 4 cm

(stage 6), 6 cm (stage7) , 8 cm (stage 8) and 10 cm (stage 9). **c**, Comparison of

relative expression levels of *FSTOMAGEN* between *rob*, *ram* and *bid*. (n=3,

biologically independent replicates). The asterisks represent statistically significant

differences (*P*<0.05, t-test, two-tailed distribution, unpaired). **d,** Comparison of the

expression levels of *FSTOMAGEN* between different species with different

photosynthetic types in *Flaveria* genus. For RT-qPCR, C_3 species incude *pri* (n=5)

and *rob* (n=13), intermediates species includes *ang* (n=3), *ram* (n=5) and *vag* (n=3),

C_4 species include *tri* (n=4), *aus* (n=3), *bid* (n=4) and *koc* (n=3). For RNA-seq, C_3

species include *pri* (n=4) and *rob* (n=4), intermediates species includes *ano* (n=4), *pub*

(n=4), *chl* (n=4), *bro* (n=4), C_4 speices include *tri* (n=4) and *bid* (n=4). RNA-seq data

was from (*41*). The number in parenthesis indicates biologically independent

replicates. The relative expression indicates the measurements with RT-qPCR, and the

others were measured with RNA-seq. The geometric means of *EF1a* and *ACT7* were

used as the reference in the calculation of expression levels for other genes. Error bars

indicate s.d. The asterisks represent statistically significant differences, One-way

ANOVA: *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, ****, $P < 0.0001$; Different letters

represent statistically significant differences, Duncan' s multiple range test (α = 0.05)).

Fig 6, Comparison of stomatal pattern and expression levels of stomatal development

related genes in the C_4 lineages other than *Flaveria*

a, Stomatal characteristics for rice and maize. *SD*, (n=8, biologically independent

replicates); *l*, (n=25, biologically independent replicates); *SI*, (n=3, biologically

independent replicates); g_s, (n=8, biologically independent replicates). Error bar

indicates s.d. The asterisks represent statistically significant differences (*P*<0.05, t-test,

two-tailed distribution, unpaired). **b**, Photographs of stomatal pattern in rice and

maize. Scale bars represent 50 μm. **c**, Comparison of transcript abundance of stomatal

development related genes between rice and maize. The developmental stages of

leaves were divided into 11 and 15 parts in rice and maize, respectively. R represents

the leaves of rice, M represents the leaves of maize. **d**, Comparison of the expression

levels of *STOMAGEN* homologs between rice and maize. There are one and two

paralogs of *STOMAGEN* that is expressing in rice and maize, respectively. *OsSTO*

represents the homologs of *STOMAGEN* in rice. *ZmSTO* represents the homologs of

STOMAGEN in maize. **e**, The families and numbers of species with different

photosynthetic type that we used to analyse. **f**, Comparison of the expression of C_4

related genes between species with different photosynthetic types. **g**, Comparison of

the expression of *STOMAGEN* homologs between species with different

photosynthetic types. The expression was determined by RNA-seq in rice and maize.

The RNA-seq data was from (*69*). The asterisks represent statistically significant

differences (P<0.05, t-test, two-tailed distribution, unpaired).

Fig 7：Schematic of the tactic change in the C_4 syndrome in the current C_4 model.

The position of the C_4 syndromes in the current C_4 model was shown. The arrows

represent the gradual enhancement or weaken of each feature over the entire

evolutionary gradient. a represents photosystem. b represents biochemistry and

metabolism. c represents structure and hydraulic feature. Darker color means

enhancement, lighter color means weaken. The yellow parts are related to the stomatal

traits shown by our data. The rest is the previous research on the C_4 related features in

Flaveria genus (4, 41, 43-45, 51, 54, 55, 102).

Legends for Supplemental Figures

Supplementary fig 1. a, Stomatal conductance (g_s) responses to net photosynthesis

rate (*A*) under saturated light. The circles on the graph indicate larger g_s values in

different photosynthetic types (Data from (*4, 83*)). **b**, The correlation of g_{smax} (abaxial

+ adaxial) and g_s. **c**, G_s, *A and iWUE* under saturated light between C_3 (rob, pri,cro)

and C_4 (bid, tri and koc) *Flaveria* species. (Data from (*4, 83*)). **d**, G_s, *A and iWUE*

under growth light between C_3 (rob) and C_4 (bid) *Flaveria* species (n=6, biologically

independent replicates). The asterisks represent statistically significant differences

(P<0.05, t-test, two-tailed distribution, unpaired).

Supplementary fig 2. g_{smax} on the adaxial surface of leaves in *Flaveria*. **a,b**, The

difference in g_{smax} in species with different photosynthetic types grown outdoors (**a**) or

in greenhouse (**b**). **c,d**, The changes of g_{smax} on the adaxial leaf surface from C_3 to C_4

species in the *Flaveria* genus grown outdoors (**c**) and in greenhouse (**d**). (n=3,

biologically independent replicates). **e**, Correlation of g_{smax} and g_s on the adaxial

surface of leaves. The asterisks represent statistically significant differences, one-way

ANOVA: *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, ****, $P < 0.0001$; Different letters

represent statistically significant differences, Duncan's multiple range test ($\alpha = 0.05$)).

1016

Supplementary fig 3. SD on the adaxial surface of leaves in *Flaveria*. **a,b**, The

difference in SD in species with different photosynthetic types grown outdoors (**a**) or

in greenhouse (**b**). **c,d**, The changes of SD on the adaxial leaf surface from C_3 to C_4

species in the *Flaveria* genus grown outdoors (**c**) and in greenhouse (**d**). (n=3,

biologically independent replicates). **e**, Correlation of g_{smax} and SD on the adaxial

surface of leaves. **f**, Correlation of g_{smax} and l on the adaxial surface of leaves. **g**,

Correlation of l and SD on the adaxial surface of leaves. The asterisks represent

statistically significant differences, one-way ANOVA: *, $P < 0.05$, **, $P < 0.01$, ***,

$P < 0.001$, ****, $P < 0.0001$; Different letters represent statistically significant

differences, Duncan's multiple range test ($\alpha = 0.05$)).

Supplementary fig 4. SD in *rob* (C_3), *ram* (intermediate) and *bid* (C_4) grown from

seeds. The SD on abaxial (ab) (n=5, biologically independent replicates) and adaxial

(ad) (n=4, biologically independent replicates) surfaces were observed in *rob*, *ram* and

bid grown in greenhouse. Different colours represent different photosynthetic types.

Different letters represent statistically significant differences (one-way ANOVA,

Duncan's multiple range test ($\alpha = 0.05$)).

Supplementary fig 5. Conservative analysis of amino acid sequences for EPF1 and EPF2. **a**. Comparison of amino acid sequences of homologs of EPF1 between different *Flaveria* species and Arabidopsis. **b**. Comparison of amino acid sequences of homologs of EPF2 between *Flaveria* species and Arabidopsis. White letters with red background represent amino acids conserved across all species. Red letters represent amino acids with similar biochemical property.

Supplementary fig 6. Hydrophobicity analysis of proteins. **a**, the protein of stomagen (left) and the protein of FSTOMAGEN (right). **b**, the protein of EPF1 (left) and the protein of FEPF1 (right). **c**, the protein of EPF2 (left) and the protein of FEPF2 (right).

Supplementary fig 7. Assessment of the stability of expression levels of reference genes. **a,b**, Comparison of the stability of expression levels of candidate reference genes in the *Flaveria* species by RNA-seq. **c**, Gel electrophoresis showing the specificity of the primers for candidate reference genes. From left to right, *ACT7*, *EF1a*, *J3*, *PNdO*, *CPN20*, *UBC9*, *GAPDH*, *F2N1.5*, *PSRP4*, *CACMS*, *HEME2*, and *UBQ11*. **d**, The coefficient of variations (CV) for the candidate reference genes. **e**, The expression stability for candidate reference genes calculated by Genorm. **f**, The expression stability for candidate reference genes calculated by Normfinder.

Supplementary fig 8. The expression levels of stomatal development related genes in *Flaveria* genus. Expression levels of stomatal development related genes in species with different photosynthetic types in the *Flaveria* genus by RNA-seq. Data from (*41*).

Supplementary fig 9. Relative expression levels of *FSTOMAGEN* and *SD* for *rob* (C$_3$) under different light conditions. (A) Relative expression levels for *FSTOMAGEN* under two different light conditions. (B) *SD* under two different light conditions. Natural light: the maximal photosynthetic photon flux density in the greenhouse was about 1300~1400 mmol m^{-2} s^{-1}. Light in the phytotron: 20~100 mmol m^{-2} s^{-1}. Reference gene used to calculate the expression level was *ACT7*. The asterisks represent statistically significant differences (*P*<0.05, t-test, two-tailed distribution, unpaired).

Supplementary fig 10. Relative expression levels of stomata development related genes determined with different reference genes and different primers. **a, b, c,**

Quantification with different primers and the same *EF1a* reference gene. Relative expressions shown in **a and b** were determined using the same primer pairs (Frob_EF1a, Fram_EF1a, Fbid_EF1a) while that in **c** used other primer pairs (EF1a) (See details of the primers used in Table S2). **d,** Relative expression levels determined with the gene *J3* used as the reference gene.

Supplementary fig 11. Transcript abundance of genes in the C_4 cycle and C_4 related transporters in *Flaveria* genus. Data was from the One Thousand Plants (1KP)(*42, 65*). J represents juvenile leaves and M represents mature leaves. Different colours represent species with different photosynthetic types in the *Flaveria* genus. Red colour represents C_3 species, grey colour represents C_3-C_4 species, green colour represents C_4-like species, and blue colour represents C_4 species.

Supplementary fig 12. Alignment of the functional domain of *STOMAGEN* with transcripts from different *Flaveria* species by Blast. From the top to bottom, *F. robusta*, *F. pringlei*, *F. angustifolia*, *F. ramosissima*, *F. vaginata*, *F. australasica*, *F. bidentis*.

Supplementary fig 13. The expression level of *FSTOMAGEN* in different *Flaveria* species based on RT-qPCR (**a**) or RNA-seq (**b**). For RT-qPCR, *pri* (n=5), *rob* (n=13), *ang* (n=3), *ram* (n=5), *vag* (n=3), *tri* (n=4), *aus* (n=3), *bid* (n=4) and *koc* (n=3). For RNA-seq, all species n=4. The relative expression indicates the measurements with RT-qPCR. RPKM indicates RNA-seq, data from (*41*). **c,** The relationship between *SD* and *FSTOMAGEN* expression in *Flaveria* species with different photosynthetic types.

Supplementary fig 14. Comparison of the expression level of *FEPF1* and *FEPF2* between *rob* and *bid Flaveria* species by RT-qPCR or RNA-seq. RNA-seq data from (*41*). *FEPF1* and *FEPF2* indicates the homologs of *EPF1* and *EPF2*.

Supplementary fig 15. The analysis for the *SD* of T2 generation of transgenic Arabidopsis (AFSTO and FSTO). **a**, Comparison of *SD* between WT and transgenic Arabidopsis AR, FSTO respectively. **b**, RNA level identification of transgenic Arabidospsis. Two independent lines was analyzed for each trangene event.

Supplementary fig 16. The increase of *SD* in cotyledon of *F. bidentis* (*bid*) with in vitro application of FSTOMAGEN. Photographs of the leaf surface in the untreated (control) and treated bid plants. scale bar, 20 mm.

Supplementary fig 17.a, Photographs of leaf surface taken with differential interference contrast microscope showing the increased stomata density in F. bidentis (bid) leaves with FSTOMAGEN applied. Scale bar represents 20 μm. **b**, Photographs of leaf surface taken with differential interference contrast microscope showing the increased stomata density in Arabidopsis leaves with FSTOMAGEN applied. Scale bar represents 20 μm.

Supplementary fig 18. Phylogenetic tree for FSTOMAGEN. blue represents dicotyledon, red represents monocotyledon and pink represents the *Flaveria* species. The gene tree of STOMAGEN based on bayesian-inference method.

Supplementary fig 19. a, Schematic of the progresssion of C_4 evolution in *Flaveria* genus. **b**, A stomatal graph taken with a light microscope showing stomatal parameters for *Flaveria* species which are needed to be measured to calculate the g_{smax}. The graph of stomata is *rob. l* (stomatal length), PL (pore length) and GCW (guard cell width). **c**, Correlation of the guard cell width and stomatal length (left), correlation of the stomatal length and pore length (right).

Supplementary fig 20. Photographs showing clustered stomata on a leaf for the transgenic Arabidopsis. Scar bar represents 10 um.

Supplementary table 1. Statistics for the mapping of the RNA-seq data of the juvenile and mature leaves in different species of the *Flaveria* genus.

Supplementary table 2. Primers used in this study

Supplementary table 3. The effect and accession number of genes related to stomatal development in Arabidopsis and rice and maize.

1120   **Supplementary table 4**. The N50 of the mapping of the RNA-SEQ data taken from

1121   1KP database for the juvenile and mature leaves

1122   **Supplementary table 5**. The mapping rates of the mapping of the RNA-SEQ data

1123   taken from 1KP database for the juvenile and mature leaves.

1124

C₃	Type I	Type II	C₄-like	C₄
Preconditioning; Proto-Kranz condition	Photorespiratory CO_2 pump (C_2 photosynthesis)	Consolidating C_2; Inducing C_4 cycles by rebalancing N metabolism	Establishment of C_4 cycle	Optimization; Efficient C_4 photosynthesis

a
The size of genome increase
PS I cyclic electron transport increase
PSII decrease

b
C_4 Metabolism enhancement
C_2 cycle enhancement
C_2 cycle weaken
Mesophyll GDC Decreases

c
BS Chloroplast Volume increase and relocate Centripetally
BS Mitochondria Increase
BS Mitos. Decrease
Mesophyll Tissue volume decreases
Vein density increases
iWUE increases
The slope of stomata responding to c_i enhances
FSTOMAGEN expression decreases
Stomatal density, g_{smax} decreases
Stomatal size increases