Analysis of Hippocampal Subfields in Sickle Cell Disease Using Ultrahigh Field MRI

Tales Santini[1], Minseok Koo[1], Nadim Farhat[1], Vinicius P. Campos[2], Salem Alkhateeb[1], Marcelo A. C. Vieira[2], Meryl A. Butters[3], Caterina Rosano[4], Howard J. Aizenstein[3,1], Joseph Mettenburg[5], Enrico M. Novelli[6,7*], Tamer S. Ibrahim[1,3,5*]

[1] Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA, United States

[2] Department of Electrical and Computer Engineering, University of São Paulo, São Carlos, SP, Brazil

[3] Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, United States

[4] Department of Epidemiology, University of Pittsburgh, Pittsburgh, PA, United States

[5] Department of Radiology, University of Pittsburgh, Pittsburgh, PA, United States

[6] Department of Medicine, University of Pittsburgh, Pittsburgh, PA, United States

[7] Department of Pharmacology and Chemical Biology, University of Pittsburgh, Pittsburgh, PA, United States

[*] Co-corresponding authors

E-mail: tibrahim@pitt.edu (TSI), noveex@upmc.edu (EN)

Word count: 2664

Abstract

Sickle cell disease (SCD) is an inherited hemoglobinopathy that causes organ dysfunction,

including cerebral vasculopathy and neurological complications. Hippocampal segmentation with

newer and advanced 7 Tesla (7T) MRI protocols has revealed atrophy in specific subregions in

other neurodegenerative and neuroinflammatory diseases, however, there is limited evidence of

hippocampal involvement in SCD. Thus, we explored whether SCD may be also associated with

abnormalities in hippocampal subregions. We conducted 7T MRI imaging in individuals with SCD,

including the HbSS, HbSC and HbS/beta thalassemia genotypes (n=37), and healthy race and

age-matched controls (n=40), using a customized head coil. Both T1 and T2 weighted images

were used for automatic segmentation of the hippocampus subfields. Individuals with SCD had

significantly smaller volume of the Dentate Gyrus and Cornu Ammonis (CA) 2 and 3 as compared

to the control group. Other hippocampal subregions also showed a trend towards smaller volumes

in the SCD group. These findings support previous reports of reduced volume in the temporal

lobe in SCD patients. Further studies are necessary to investigate the mechanisms that lead to

structural changes in the hippocampus subfields and their relationship with cognitive performance

in SCD patients.

Introduction

Sickle Cell Disease (SCD) consists of a group of heterogeneous syndromes that share the inheritance of a mutated sickle hemoglobin (hemoglobin S, or HbS) [1]. SCD is one of the most common genetic disorders, with more than 5 million newborns carrying the mutated gene and more than 300,000 individuals being born with the disease yearly [2]. The most predominant and severe type of SCD is caused by the homozygous inheritance of HbS (HbSS disease, or sickle cell anemia) [3]. SCD may also originate from co-inheritance of HbS with other hemoglobin mutations, such as hemoglobin C (HbSC) or β-thalassemia (HbSβ-thalassemia) [4]. β-thalassemia leads to absent (β^0) or reduced (β^+) synthesis of the beta-globin chains, and the coinheritance of HbS with β-thalassemia can be further categorized into two forms: HbSβ$^+$ thalassemia and HbSβ^0 thalassemia [5]. The genotypes HbSC and HbSβ$^+$ tend to result in milder phenotypes, while HbSβ^0 thalassemia may be as severe as HbSS disease [6].

Neurological complications of SCD include increased risk of stroke, silent cerebral infarction, and cognitive impairment, and are more common in individuals with HbSS disease [7, 8]; pediatric stroke, in particular, happens almost exclusively in individuals with HbSS. HbSS individuals also tend to display lower cognitive performance when compared with individuals with HbSC [9], possibly because anemia is less pronounced in HbSC. However, little research has been done on the impact of HbSC on the brain in adults.

Reduced cortical and subcortical volume has been previously reported in patients with SCD when compared with healthy controls. Kirk et al [10] showed cortical thinning in multiple brain regions in patients with SCD ranging from 5 to 21 years-old. They hypothesized that cortical thinning could be related with impaired cerebral perfusion. Mackin et al. [8] reported reduced cortical thickness in the temporal lobe in adults with SCD, while Kawadler et al. [11] found

significantly reduced volume in the hippocampus of SCD pediatric patients with silent cerebral

infarction. Overall, evidence of the impact of SCD on brain regions remains limited, particularly in

adults.

The hippocampal involvement in higher cognitive processing has been well investigated,

especially as pertains the hippocampal primary role in short-term and episodic long-term memory

and learning [12]. The hippocampus is a bilaminar brain structure situated in the medial temporal

lobe and it is comprised of the Cornu Ammonis (CA) – which is further divided into three

conventional histological subregions, CA1-CA3 – and the Dentate Gyrus (DG) [13]. Additionally,

the hippocampal formation includes the Subiculum (Sub) and the Entorhinal Cortex (ErC) [14,

15]. Hippocampal subfields are functionally interconnected, however functionalities are known to

vary among the different subfields. For example, DG is suggested to be a preprocessor of

incoming information, preparing it for subsequent processing in CA3, and potentially mediating

learning, memory, and spatial encoding [16]. ErC, on the other hand, serves as the major input-

output structure and mediates hippocampal connectivity with cortical regions [17]. Cells in the DG

receive excitatory input from ErC and send excitatory output to the CA3 region via the mossy

fibers [16]. Also, CA1 and CA3 are believed to contribute to episodic memory processing [18, 19].

CA2 is the smallest subfield but studies found that lesions in this area can lead to abnormal social

behavior by impairing social recognition memory [20].

As pathology within the hippocampus is increasingly linked to a number of neurological

and neuropsychiatric diseases such as Alzheimer's disease, epilepsy, and depression [21-25],

interest in the subfields of the hippocampus has grown. Subregions of the hippocampus are

differentially affected, depending on the disease [26]. Recent advances in MRI techniques, such

as 7 Tesla (T) MRI, have improved the spatial resolution and signal-to-noise ratio of MRI images

[27]. These improvements enable more accurate volumetric analysis of the hippocampus

subfields in *in vivo* studies, and improve automatic segmentation methods [28-32]. Manual

segmentation of hippocampal subregions is time-consuming and labor intensive, limiting data

collection and processing in large studies [23, 31]. To compensate for this shortcoming,

automated segmentation methods have been developed [29, 31, 33].

This study aims to investigate how SCD impacts the volumes of hippocampus subfields.

We used 7T MRI and an innovative radiofrequency (RF) coil to acquire high-resolution images of

the hippocampal formation from a cohort of SCD participants and controls (N=77). 7T tailored

preprocessing and automatic segmentation software [29] were used to compute the volumes of

the hippocampus subfields. We hypothesized that the participants with SCD would exhibit lower

hippocampal volume when compared with the control group and that the disease could affect

specific hippocampal subregions.

Methods

Participants

Patients with SCD were recruited from the University of Pittsburgh Medical Center (UPMC) Adult

Sickle Cell Program outpatient clinic under the University of Pittsburgh IRB protocols

PRO12040139 and PRO08110422. Data were collected from 53 patients and 47 controls as part

of a longitudinal study of neuroradiological biomarkers in SCD (ClinicalTrials.gov Identifier:

NCT02946905). All patients with HbSS, HbSC and HbSβ thalassemia older than 18 years old and

able to provide informed consent were informed about the study by staff members during their

routine clinic visit, and offered entry into the study if they were in steady-state SCD (i.e. two weeks

from an acute illness, including vaso-occlusive pain episodes). Eligibility criteria also included: 1)

English-speaking; and 2) currently receiving routine follow-up care at the UPMC Adult Sickle Cell

Program. Exclusion criteria included: 1) pregnancy, as determined by a positive urine human

chorionic gonadotropin test at the time of the of the MRI; 2) acute medical problems including, but

not limited to, acute vaso-occlusive episodes. Healthy controls, included in ClinicalTrials.gov

identifier NCT02946905, were recruited from the community via brochures and from the University

of Pittsburgh Clinical and Translational Science Institute Pitt+Me Registry if older than 18 and

able to provide informed consent. Healthy controls were age, race, and gender matched to the

participants with SCD. Both patients with SCD and controls were excluded if pregnant or lactating,

had any medical condition that could result in neurocognitive or brain dysfunction (other than

those resulting from SCD), including diabetes mellitus, coronary artery disease, peripheral

vascular disease, and other causes of cerebral vasculitis such as Systemic Lupus Erythematosus

(SLE), or contraindications to MRI scanning such as electronic implants, magnetically-activated

implants, tattoos above the shoulders, or brain implants.

MRI acquisition

MRI images were acquired using a 7T human MRI scanner (Siemens Magnetom,

Germany) with a customized radiofrequency (RF) head coil with 16 transmit channels and 32

receive channels [34]. This RF coil produces homogenous images and whole brain coverage at

7T [27, 35-38]. T1-weighted (T1w) Magnetization Prepared RApid Gradient Echo (MPRAGE) was

acquired with 0.75 mm isotropic resolution and the following parameters: TE/TI/TR =

2.17/1200/3000 ms, bandwidth = 391 Hz/Px, acceleration factor (Grappa) = 2, time of acquisition

= 5:02 min. T2w images were acquired with high resolution in the plane perpendicular to the main

axis of the hippocampus, slightly slanted from coronal acquisition. The sequence utilized was 2D

turbo spin echo (TSE) with the following parameters: resolution = 0.375 x 0.375 x 1.5 mm^3, TE/TR

= 61/10060 ms, bandwidth = 264, acceleration factor (Grappa) = 2, time of acquisition = 3:32 min.

Gradient echo (GRE) sequence – utilized as a secondary contrast for the calculations of the

intracranial volume masks – was acquired with the following parameters: TE/TR = 8.16/24 ms,

resolution 0.375 x 0.375 x 0.75 mm, acceleration factor 2, time of acquisition 8:20 min.

Image preprocessing and hippocampus segmentation

Both T1w and T2w images were denoised using an optimal forward and inverse variance-

stabilizing transformation (VST), specifically designed for the Rice distribution [39]. The forward

VST converts the Rician heteroscedastic noise into a homoscedastic one (approximately

Gaussian distributed), and thus the block-matching 4D (BM4D) denoising algorithm [40] was

applied, followed by the inverse VST, to obtain the denoised images [41]. The images were then

bias corrected with the SPM12 package using 30 mm of cutoff for the full width at half maximum

(against 60mm of the default) [42]. **Figure 1** shows the preprocessing methods applied in a

representative T2w image from this study. The bias correction and denoising methods improved

homogeneity and mitigated noise in the images.

The automatic segmentation of hippocampal subfields (ASHS) package [29] and a 7T

atlas from young adults' data [43] were utilized to segment hippocampal subfields. The images

from the atlas were denoised using the same technique described above for the T1w and T2w

images. The results of the segmentation were manually corrected when the fixes were obvious

and following the strategies described by Berron et al. [43]. The corrections were performed using

the software ITK-SNAP [44]. In 23 of the 100 subjects, either the automatic segmentation failed

or the manual fixes were not obvious (mostly due to excessive motion artifact); these subjects

were excluded from the analysis, yielding a final sample size of 77. **Figure 2** shows the

hippocampus segmentation in a representative SCD patient. The preprocessing techniques

applied before segmentation with the ASHS package produced a smoother and more accurate

segmentation of the hippocampus subfields. A summary of the methods is shown in **Figure 3**.

Intracranial volume is often used as a covariate in brain volumetric analysis; intracranial

masks were calculated with the SPM12 package using the MPRAGE and GRE acquisitions as

channels for the segmentations after a co-registration. The segmentations for the white matter,

gray matter, and cerebrospinal fluid were combined, and the missing pixels in the segmentation

were filled using the MATLAB function *imfill*. The imperfections in the segmentations were

manually corrected using the ITK-snap software [44].

Statistical Analysis

Sample characteristics were compared between SCD groups and controls using one-way

ANOVA. The volumes of the hippocampal subfields were compared for patients and controls

using analysis of covariance (ANCOVA), with gender, age, and intracranial volume as covariates.

Those measures that differed by SCD status (patients and controls) were considered for inclusion

in multivariable models testing the associations of SCD status and neuroimaging measures. The

p-value threshold for between-group differences was calculated based on the Bonferroni

correction. The statistical analysis was performed using SPSS (version 26, IBM, Chicago, IL).

Results

Demographics and hemoglobin values of the participants

There was no statistically significant difference in the age or gender between the groups

(**Table 1**). As expected, the hemoglobin levels were significantly different between groups, with

the severe SCD group (HbSS and HbSβ^0) having the lowest Hb level (9.10 ± 1.09 g/dL), the milder

SCD group (HbSC and HbSβ^+) having an intermediate level (11.25 ± 2.48 g/dL), and the controls

having the highest Hb level (13.94 ± 1.55 g/dL) at the time of the assessment (**Table 1**).

Neuroimaging characteristics of patients with SCD vs. controls

Results of the comparison of the hippocampal subfields volumes between SCD patients

and controls are tabulated in **Table 2**. **Figure 4** shows the bar plots with the individual volumetric

data points for all the subjects included in this study. Significant differences between the

hippocampal volumes of the SCD and control groups were observed bilaterally in the region that

encompasses the DG, CA2, and CA3: - 11.55% (F = 20.79, p = 0.020×10^{-3}) in the left

hippocampus, and - 11.36% (F = 14.09, p = 0.350×10^{-3}) in the right hippocampus. There was also

a trend towards a reduction of the left CA1 (- 10.52%, F = 5.69 , p-value 0.020), right CA1

(- 8.02%, F = 4.02, p-value 0.049), and left ErC (- 9.03%, F = 5.68, p-value 0.020), which was not

statistically significant after Bonferroni correction. The other hippocampal subregions, including

the right ErC, bilateral Tail, and bilateral Sub, did not significantly differ between SCD and control

groups. The whole left (- 7.29%, F = 7.29, p-value 0.009) and right hippocampus (- 6.57%,

F = 4.93, p-value 0.030) showed a trend that did not remain statistically significant after Bonferroni

correction. Between-group differences in the region containing DG, CA2, and CA3 were

marginally attenuated after adjustment for intracranial volume (from 11.55% to 10.10% for the left

hemisphere, and from 11.36% to 9.63% for the right hemisphere), and minimally attenuated with

the inclusion of age and gender in the model, as shown in **Table 3**. No significant differences

were found between the severe and milder SCD genotypes.

Discussion

In this study, 7T MRI and innovative radiofrequency coil developments allowed the

acquisition of high-resolution and high-quality images of the hippocampus. Preprocessing

techniques such as denoising and bias correction further improved the image quality. Manual

quality assurance and correction refined the automatic hippocampal segmentation method. The

use of the GRE sequence with the T1-weighted MPRAGE also improved the estimation of the

intracranial volume masks, although manual corrections were still necessary.

Compared with controls, our analysis shows that the region DG+CA2+CA3 is the most

affected hippocampal subregion in the SCD group, an effect observed with a mean difference of

-11.55% on the left hemisphere and -11.36% on the right hemisphere. After adjusting for age,

gender, and intracranial volume, the differences were marginally reduced, indicating that SCD is

the major contributor to the overall difference. Moreover, the hippocampal subregions CA1

bilaterally, left ErC, and hippocampus bilaterally also showed a trend towards smaller volumes in

the SCD group, but the differences were not statistically significant after Bonferroni correction.

These results are consistent with findings in pediatric populations with SCD. Kawadler et

al. [11] reported a volume difference of -6.74% in the right hippocampus and -10.26% in the left

hippocampus in pediatric HbSS subjects with silent cerebral infarction when compared with

healthy controls. If confirmed in future studies, our results indicate that the impact of SCD on

hippocampal morphology reported in pediatric patients persists in adulthood. Future studies

should assess the impact of these patterns of significant subregional hippocampal atrophy on

cognitive decline.

To the best of our knowledge, our study is the first to analyze hippocampal subregions

with high-resolution images obtained with 7T MRI, and to include both severe and milder

genotypes of SCD. Understanding how hippocampal subregion volumes differ in SCD may

identify biomarkers of disease severity and cognition. For instance, our analysis found that the

total hippocampal volume was not significantly different between patients and controls, whereas

the DG+CA2+CA3 subregion grouping showed a consistent volumetric difference between the

two groups. This hippocampal region is believed to mediate learning, memory, and spatial

encoding [16], and abnormalities in this area could be associated with deficits in these cognitive

domains in SCD.

The hippocampus is particularly vulnerable to hypoxia and inflammation [45], which are

common pathogenic mechanisms in SCD. Mouse models of SCD reveal that hippocampal

pathology is associated with cognitive deficits [46]. Interestingly, we did not observe an

association between the volume of hippocampal subregions and SCD genotype, despite evidence

that individuals with milder genotypes tend to have better cognitive functioning [47, 48]. It is

possible that compensatory mechanisms such as increased hippocampal activation and

connectivity [49, 50] may confound the effect of hippocampal atrophy on cognition.

One limitation of this study is that hippocampal subregions may be inaccurately labeled

as the adjacent subregions. In particular, CA2 and CA3 subregions are small and their borders

do not present clear landmarks in MRI; therefore, they are more prone to mislabeling errors. On

the other hand, the segmentation of DG is the most accurate [29], since it follows well defined

landmarks in the T2-weighted image. Nevertheless, only histological data could precisely identify

all specific subregions. To reduce the impact of mislabeling, we combined CA2 and CA3 with DG,

a common strategy [51]. To further enhance the precision of our measurements, we excluded 23

subjects whose scans presented excessive motion artifacts, mostly due to artifacts in the T2w

TSE acquisition. Future work in RF coil design, MRI sequence development, and image

processing may lead to a higher inclusion rate.

In summary, we found that between-group differences in the hippocampus followed a

distinct spatial distribution, being more pronounced in the DG+CA2+CA3 bilaterally as compared

to other subregions. These associations were attenuated after adjustment for intracranial volume

and demographics, but remained significant. Further studies will be necessary to clarify the

mechanisms that lead to volume reduction in the hippocampal subfields and elucidate their

significance as an imaging biomarker for cognitive deficits in individuals with SCD.

Acknowledgments

253　　　This work was supported by the National Institutes of Health under award numbers:

254　R01HL127107, R01MH111265, R01AG063525, and T32MH119168.　The first author was

255　partially supported by CAPES Foundation, Ministry of Education of Brazil, under the award

256　number 13385/13-5. This research was also supported in part by the University of Pittsburgh

257　Center for Research Computing (CRC) through the resources provided.

References

1. ter Maaten JC, Arogundade FA. Sickle cell disease. Comprehensive Clinical Nephrology: Elsevier; 2010. p. 596-608.

2. Piel FB, Patil AP, Howes RE, Nyangiri OA, Gething PW, Dewi M, et al. Global epidemiology of sickle haemoglobin in neonates: a contemporary geostatistical model-based map and population estimates. The Lancet. 2013;381(9861):142-51.

3. Davies SC, Gilmore A. The role of hydroxyurea in the management of sickle cell disease. Blood reviews. 2003;17(2):99-109.

4. Davies SC, Oni L. Fortnightly review: Management of patients with sickle cell disease. BMJ. 1997;315(7109):656-60.

5. Galanello R, Origa R. Beta-thalassemia. Orphanet journal of rare diseases. 2010;5(1):11.

6. Serjeant GR. Sickle-cell disease. The Lancet. 1997;350(9079):725-30.

7. Dowling MM, Quinn CT, Rogers ZR, Buchanan GR. Acute silent cerebral infarction in children with sickle cell anemia. Pediatr Blood Cancer. 2010;54(3):461-4. Epub 2009/10/09. doi: 10.1002/pbc.22242. PubMed PMID: 19813251; PubMed Central PMCID: PMCPMC2807470.

8. Mackin RS, Insel P, Truran D, Vichinsky EP, Neumayr LD, Armstrong FD, et al. Neuroimaging abnormalities in adults with sickle cell anemia: associations with cognition. Neurology. 2014;82(10):835-41. Epub 2014/02/14. doi: 10.1212/WNL.0000000000000188. PubMed PMID: 24523480; PubMed Central PMCID: PMCPMC3959758.

9. Jorgensen DR, Metti A, Butters MA, Mettenburg JM, Rosano C, Novelli EM. Disease severity and slower psychomotor speed in adults with sickle cell disease. Blood advances. 2017;1(21):1790-5.

10. Kirk GR, Haynes MR, Palasis S, Brown C, Burns TG, McCormick M, et al. Regionally specific cortical thinning in children with sickle cell disease. Cerebral Cortex. 2009;19(7):1549-56.

11. Kawadler JM, Clayden JD, Kirkham FJ, Cox TC, Saunders DE, Clark CA. Subcortical and cerebellar volumetric deficits in paediatric sickle cell anaemia. British journal of haematology. 2013;163(3):373-6.

12. Amaral DG, Witter MP. The three-dimensional organization of the hippocampal formation: A review of anatomical data. Neuroscience. 1989;31(3):571-91. doi: 10.1016/0306-4522(89)90424-7.

13. de Flores R, Berron D, Ding SL, Ittyerah R, Pluta JB, Xie L, et al. Characterization of hippocampal subfields using ex vivo MRI and histology data: Lessons for in vivo segmentation. Hippocampus. 2020;30(6):545-64. Epub 2019/11/02. doi: 10.1002/hipo.23172. PubMed PMID: 31675165; PubMed Central PMCID: PMCPMC7192771.

14. O'Mara SM, Commins S, Anderson M, Gigg J. The subiculum: a review of form, physiology and function. Progress in neurobiology. 2001;64(2):129-55.

15. Witter MP, Doan TP, Jacobsen B, Nilssen ES, Ohara S. Architecture of the entorhinal cortex a review of entorhinal anatomy in rodents with some comparative notes. Frontiers in Systems Neuroscience. 2017;11:46.

16. Jonas P, Lisman J. Structure, function, and plasticity of hippocampal dentate gyrus microcircuits. Front Neural Circuits. 2014;8:107. Epub 2014/10/14. doi: 10.3389/fncir.2014.00107. PubMed PMID: 25309334; PubMed Central PMCID: PMCPMC4159971.

17. Witter MP, Doan TP, Jacobsen B, Nilssen ES, Ohara S. Architecture of the Entorhinal Cortex A Review of Entorhinal Anatomy in Rodents with Some Comparative Notes. Front Syst Neurosci. 2017;11:46. Epub 2017/07/14. doi: 10.3389/fnsys.2017.00046. PubMed PMID: 28701931; PubMed Central PMCID: PMCPMC5488372.

18. Hunsaker MR, Lee B, Kesner RP. Evaluating the temporal context of episodic memory: the role of CA3 and CA1. Behav Brain Res. 2008;188(2):310-5. Epub 2008/01/08. doi: 10.1016/j.bbr.2007.11.015. PubMed PMID: 18178264; PubMed Central PMCID: PMCPMC2675273.

19. Zheng F, Cui D, Zhang L, Zhang S, Zhao Y, Liu X, et al. The Volume of Hippocampal
Subfields in Relation to Decline of Memory Recall Across the Adult Lifespan. Front Aging
Neurosci. 2018;10:320. Epub 2018/10/27. doi: 10.3389/fnagi.2018.00320. PubMed PMID:
30364081; PubMed Central PMCID: PMCPMC6191512.
20. Stevenson EL, Caldwell HK. Lesions to the CA 2 region of the hippocampus impair social
memory in mice. European Journal of Neuroscience. 2014;40(9):3294-301.
21. Mueller S, Yushkevich P, Wang L, Van Leemput K, Mezher A, Iglesias JE, et al.
Collaboration for a systematic comparison of different techniques to measure subfield volumes:
announcement and first results. Alzheimer's & Dementia: The Journal of the Alzheimer's
Association. 2013;9(4):P51.
22. Mueller S, Stables L, Du A, Schuff N, Truran D, Cashdollar N, et al. Measurement of
hippocampal subfields and age-related changes with high resolution MRI at 4 T. Neurobiology of
aging. 2007;28(5):719-26.
23. Wisse LE, Biessels GJ, Heringa SM, Kuijf HJ, Luijten PR, Geerlings MI, et al. Hippocampal
subfield volumes at 7T in early Alzheimer's disease and normal aging. Neurobiology of aging.
2014;35(9):2039-45.
24. Small SA, Schobel SA, Buxton RB, Witter MP, Barnes CA. A pathophysiological
framework of hippocampal dysfunction in ageing and disease. Nature Reviews Neuroscience.
2011;12(10):585-601.
25. de Flores R, La Joie R, Landeau B, Perrotin A, Mézenge F, de La Sayette V, et al. Effects
of age and Alzheimer's disease on hippocampal subfields: comparison between manual and
FreeSurfer volumetry. Human brain mapping. 2015;36(2):463-74.
26. West MJ, Kawas CH, Stewart WF, Rudow GL, Troncoso JC. Hippocampal neurons in pre-
clinical Alzheimer's disease. Neurobiology of aging. 2004;25(9):1205-12. Epub 2004/08/18. doi:
10.1016/j.neurobiolaging.2003.12.005. PubMed PMID: 15312966.
27. Krishnamurthy N, Santini T, Wood S, Kim J, Zhao T, Aizenstein HJ, et al. Computational
and experimental evaluation of the Tic-Tac-Toe RF coil for 7 Tesla MRI. PloS one.
2019;14(1):e0209663. Epub 2019/01/11. doi: 10.1371/journal.pone.0209663. PubMed PMID:
30629618.
28. Knierim JJ. The hippocampus. Current Biology. 2015;25(23):R1116-R21.
29. Yushkevich PA, Pluta JB, Wang H, Xie L, Ding SL, Gertje EC, et al. Automated volumetry
and regional thickness analysis of hippocampal subfields and medial temporal cortical structures
in mild cognitive impairment. Human brain mapping. 2015;36(1):258-87.
30. Boccardi M, Ganzola R, Bocchetta M, Pievani M, Redolfi A, Bartzokis G, et al. Survey of
protocols for the manual segmentation of the hippocampus: preparatory steps towards a joint
EADC-ADNI harmonized protocol. Journal of Alzheimer's disease. 2011;26(s3):61-75.
31. Iglesias JE, Augustinack JC, Nguyen K, Player CM, Player A, Wright M, et al. A
computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI:
application to adaptive segmentation of in vivo MRI. Neuroimage. 2015;115:117-37.
32. Giuliano A, Donatelli G, Cosottini M, Tosetti M, Retico A, Fantacci ME. Hippocampal
subfields at ultra high field MRI: A n overview of segmentation and measurement methods.
Hippocampus. 2017;27(5):481-94.
33. Yang Z, Zhuang X, Mishra V, Sreenivasan K, Cordes D. CAST: A multi-scale convolutional
neural network based automated hippocampal subfield segmentation toolbox. NeuroImage.
2020:116947.
34. Ibrahim T, Zhao Y, Krishnamurthy N, Raval S, Zhao T, Wood S, et al. 20-To-8 Channel
Tx Array with 32-Channel Adjustable Receive-Only Insert for 7T Head Imaging. International
Society of Magnetic Resonance in Medicine; Salt Lake City, Utah2013. p. 4408.
35. Ibrahim T ST, Raval S, Krishnamurthy N, Wood S, Kim J, Zhao Y, Wu X, Yacoub E,
Aizenstein H, Zhao T. Towards Homogenous 7T Neuro Imaging: Findings and Comparisons

between 7T TTT and NOVA RF Coil Systems. In Proc of the 25th International Society of
Magnetic Resonance in Medicine Annual Meeting; Honolulu, Hawaii, USA2017.
36.   Smagula SF, Karim HT, Rangarajan A, Santos FP, Wood SC, Santini T, et al. Association
of Hippocampal Substructure Resting-State Functional Connectivity with Memory Performance in
Older Adults. The American journal of geriatric psychiatry : official journal of the American
Association for Geriatric Psychiatry. 2018. doi: 10.1016/j.jagp.2018.03.003. PubMed PMID:
29628321.
37.   Santini T, Zhao Y, Wood S, Krishnamurthy N, Kim J, Farhat N, et al. In-vivo and numerical
analysis of the eigenmodes produced by a multi-level Tic-Tac-Toe head transmit array for 7 Tesla
MRI. PloS one. 2018;13(11):e0206127. doi: 10.1371/journal.pone.0206127. PubMed PMID:
30481187; PubMed Central PMCID: PMCPMC6258503 products in development to declare. This
does not alter our adherence to PLOS ONE policies on sharing data and materials.
38.   Santini T, Wood S, Krishnamurthy N, Martins T, Aizenstein HJ, Ibrahim TS. Improved 7
Tesla Transmit Field Homogeneity with Reduced Electromagnetic Power Deposition Using
Coupled   Tic   Tac   Toe   Antennas.   bioRxiv.   2020:2020.11.06.371328.   doi:
10.1101/2020.11.06.371328.
39.   Foi A. Noise estimation and removal in MR imaging: The variance-stabilization approach.
2011 IEEE International symposium on biomedical imaging: from nano to macro: IEEE; 2011. p.
1809-14.
40.   Maggioni M, Katkovnik V, Egiazarian K, Foi A. Nonlocal transform-domain filter for
volumetric data denoising and reconstruction. IEEE transactions on image processing.
2012;22(1):119-33.
41.   Santini T, Brito F, Wood S, Martins T, Mettenburgh J, Aizenstein H, et al. Noise mitigation
from high-resolution 7T MRI images. In Proc of the 26th International Society of Magnetic
Resonance in Medicine Annual Meeting; Paris, France2018.
42.   Ibrahim TS, Lee R, Abduljalil AM, Baertlein BA, Robitaille P-ML. Dielectric resonances
and B 1 field inhomogeneity in UHFMRI: computational analysis and experimental findings.
Magnetic resonance imaging. 2001;19(2):219-26.
43.   Berron D, Vieweg P, Hochkeppler A, Pluta J, Ding S-L, Maass A, et al. A protocol for
manual segmentation of medial temporal lobe subregions in 7 Tesla MRI. NeuroImage: Clinical.
2017;15:466-82.
44.   Yushkevich PA, Piven J, Hazlett HC, Smith RG, Ho S, Gee JC, et al. User-guided 3D
active contour segmentation of anatomical structures: significantly improved efficiency and
reliability. Neuroimage. 2006;31(3):1116-28.
45.   Iampietro M, Giovannetti T, Tarazi R. Hypoxia and inflammation in children with sickle cell
disease: implications for hippocampal functioning and episodic memory. Neuropsychol Rev.
2014;24(2):252-65. Epub 2014/04/20. doi: 10.1007/s11065-014-9259-4. PubMed PMID:
24744195.
46.   Wang L, Almeida LEF, de Souza Batista CM, Khaibullina A, Xu N, Albani S, et al. Cognitive
and behavior deficits in sickle cell mice are associated with profound neuropathologic changes in
hippocampus and cerebellum. Neurobiol Dis. 2016;85:60-72. Epub 2015/10/16. doi:
10.1016/j.nbd.2015.10.004. PubMed PMID: 26462816; PubMed Central PMCID:
PMCPMC4688201.
47.   Jorgensen DR, Metti A, Butters MA, Mettenburg JM, Rosano C, Novelli EM. Disease
severity and slower psychomotor speed in adults with sickle cell disease. Blood Adv.
2017;1(21):1790-5. Epub 2018/01/04. doi: 10.1182/bloodadvances.2017008219. PubMed PMID:
29296825; PubMed Central PMCID: PMCPMC5728099 interests.
48.   Schatz J, Finke RL, Kellett JM, Kramer JH. Cognitive functioning in children with sickle
cell disease: a meta-analysis. J Pediatr Psychol. 2002;27(8):739-48. Epub 2002/10/31. doi:
10.1093/jpepsy/27.8.739. PubMed PMID: 12403864.

49. Case M, Zhang H, Mundahl J, Datta Y, Nelson S, Gupta K, et al. Characterization of
functional brain activity and connectivity using EEG and fMRI in patients with sickle cell disease.
NeuroImage Clinical. 2017;14:1-17. Epub 2017/01/25. doi: 10.1016/j.nicl.2016.12.024. PubMed
PMID: 28116239; PubMed Central PMCID: PMCPMC5226854.
50. Hamalainen A, Pihlajamaki M, Tanila H, Hanninen T, Niskanen E, Tervo S, et al. Increased
fMRI responses during encoding in mild cognitive impairment. Neurobiology of aging.
2007;28(12):1889-903. Epub 2006/09/26. doi: 10.1016/j.neurobiolaging.2006.08.008. PubMed
PMID: 16997428.
51. Yushkevich PA, Amaral RS, Augustinack JC, Bender AR, Bernstein JD, Boccardi M, et al.
Quantitative comparison of 21 protocols for labeling hippocampal subfields and parahippocampal
subregions in in vivo MRI: towards a harmonized segmentation protocol. Neuroimage.
2015;111:526-41. Epub 2015/01/18. doi: 10.1016/j.neuroimage.2015.01.004. PubMed PMID:
25596463; PubMed Central PMCID: PMCPMC4387011.

Table 1. Sample Characteristics

	SCD HbSS, HbSβ⁰ (severe)	*SCD HbSC, HbSβ⁺ (milder)*	*Controls*	*F-statistic (p-value)*
N	*16*	21	*40*	-
Age, mean (SD)	34.31(10.23)	35.29(12.24)	36.15(13.24)	0.131 (0.88)
Gender, F/M	*7/9*	*9/12*	*16/24*	0.41 (0.96)
Hb¹, mean (SD)	**9.10 (1.09)**	**11.25 (2.48)**	**13.94 (1.55)**	**43.5 (6.9E-13) ***
¹ in g/dL *** p-value < 0.05**				

Table 2. Neuroimaging Characteristics. Mean volume in mm³, SD and percent difference

	SCD (severe and milder) -- Mean (SD)	Controls -- Mean (SD)	Percentage difference SCD vs controls[5]	F-statistic (p-value)
Intracranial volume[1]	1306040 (114740)	1336886 (130234)	-2.31%	2.813 (0.097786)[a]
Total hippocampal volume[1]	4635.69 (541.56)	4980.17 (557.54)	-6.92%	6.400 (0.013601)[b]
Asymmetry Ratio[2]	0.03655 (.02280)	0.03385 (.02242)	7.97%	0.475 (0.492749) [b]
Left Hemisphere				
Cornu Ammonis 1[1]	619.02 (115.08)	691.82 (122.65)	-10.52%	5.69 (0.019689) [b]
Dentate Gyrus[1,3]	**632.49** (70.68)	**715.05** (92.82)	**-11.55**%	**20.79 (0.000020) [b]***
Subiculum[1]	979.63 (116.29)	999.65 (133.50)	-2.00%	0.02 (0.899994) [b]
Entorhinal cortex[1]	743.54 (133.47)	817.36 (131.31)	-9.03%	5.68 (0.019773) [b]
Tail[3]	583.22 (123.34)	561.99 (96.57)	3.78%	1.34 (0.251466) [b]
Hippocampus[1]	2231.14 (250.92)	2406.52 (281.49)	-7.29%	7.29 (0.008650) [b]
Right Hemisphere				
Cornu Ammonis 1[1]	685.96 (116.09)	745.73 (108.93)	-8.02%	4.02 (0.048854) [b]
Dentate Gyrus[3]	**669.81 (94.72)**	**755.69 (103.89)**	**-11.36**%	**14.09 (0.000350) [b]***
Subiculum[1]	1048.77 (145.29)	1072.22 (131.94)	-2.19%	0.01 (0.918181) [b]
Entorhinal cortex[1]	799.98 (186.52)	841.61 (124.96)	-4.95%	0.81 (0.371777) [b]
Tail[1,4]	499.13 (84.39)	494.83 (69.95)	0.87%	0.46 (0.501763) [b]
Hippocampus[1]	2404.55 (300.00)	2573.64 (287.13)	-6.57%	4.93 (0.029504) [b]

[1] in mm³

[2] Defined as (right hippocampus - left hippocampus)/(right hippocampus + left hippocampus)*100

[3] Includes Dentate Gyrus, Cornu Ammonis 2 and 3

[4] Defined in the reference [23]

[5] Given 95% confidence interval

[a] Corrected for age and gender

[b] Corrected for age, gender, and intracranial volume

* Significantly different after Bonferroni correction (p-value threshold: 0.05 / 10 subfields / 2 groups = 0.0025 for significance)

Table 3. Multivariable Regression Models of Sickle Cell Disease Status predicting Dentate Gyrus volume

	Beta Coefficient (95% Confidence Intervals) in mm^3, Percentage change, p-value		
	Unadjusted model	*Model 1: adjusted for Intracranial volumes*	*Model 2: further adjusted for Age and Gender*
Dentate Gyrus, Right[1]	85.88 (40.62-131.13), 11.36%, 0.000312	72.76 (33.68-111.84), 9.63%, 0.000399	69.07 (32.39-105.75), 9.14%, **0.000350**
Dentate Gyrus, Left[1]	82.56 (44.87-120.24), 11.55%, 0.000040	72.25 (39.03-105.46), 10.10%, 0.000045	71.97 (40.51-103.43), 10.06%, **0.000020**
[1] Includes Dentate Gyrus, Cornu Ammonis 2 and 3			

427
428 **Figure 1: Preprocessing of the T2-weighted images in a representative SCD patient. In a),**
429 **raw image from the 7T scanner. In b), bias corrected image. In c), bias corrected and**
430 **denoised image**
431
432

Figure 2: Example of hippocampal subfields segmentation in a patient with SCD. a) coronal slice of the T2-weighted image; b,c) zoomed image showing details of the hippocampus structure, subject right and left, respectively; d,e) hippocampus subfields segmentations overlaying the T2-weighted image, subject right and left, respectively; f) 3D reconstruction of the hippocampal subfield segmentations. Abbreviations - cornu ammonis 1-3, CA1-3; dentate gyrus, DG; hippocampal tail, Tail; subiculum, Sub; entorhinal cortex, ErC.

441
442 **Figure 3: Flowchart of the method implemented for the hippocampal subfields**
443 **segmentation and analysis. T1w, T1-weighted contrast; T2w, T2-weighted contrast; GRE:**
444 **gradient echo sequence.**
445

Figure 4: Hippocampal subfield volume comparison between patients and controls. There was a significant volume difference between the patient and the control groups in the region formed by the Dentate Gyrus (DG), CA2, and CA3 bilaterally. Abbreviations - cornu ammonis 1-3, CA1-3; dentate gyrus, DG; entorhinal cortex, ErC; hippocampal Tail, Tail; subiculum, Sub. The error bars represent the standard deviation of the data. Controls (n=40), Milder SCD (n=21), Severe SCD (n=16).