A scale-free, fully connected global transition network underlies known microbiome diversity

Gongchao Jing [1,4][$], Yufeng Zhang [2][$], Lu Liu [1,4], Zengbin Wang [1,4], Zheng Sun [1,4], Rob Knight [3],

Xiaoquan Su [2][*] and Jian Xu [1,4*]

[1]Single-Cell Center, CAS Key Laboratory of Biofuels and Shandong Key Laboratory of Energy

Genetics, Qingdao Institute of BioEnergy and Bioprocess Technology, Chinese Academy of

Sciences, Qingdao, Shandong, China

[2]College of Computer Science and Technology, Qingdao University, Qingdao, Shandong, China

[3]University of California, San Diego, CA, USA

[4]University of Chinese Academy of Sciences, Beijing, China

[$]Contributed equally to this work

[*]Corresponding to suxq@qdu.edu.cn or xujian@qibebt.ac.cn

Abstract

Microbiomes are inherently linked by their structural similarity, yet the global features of such

similarity are not clear. Here we propose as solution a search-based microbiome transition network.

By traversing a composition-similarity based network of 177,022 microbiomes, we show that

although the compositions are distinct by habitat, each microbiome is on-average only seven

neighbors from any other microbiome on Earth, indicating the inherent homology of microbiome at

the global scale. This network is scale-free, suggesting a high degree of stability and robustness in

microbiome transition. By tracking the minimum spanning tree in this network, a global roadmap of

microbiome dispersal was derived that tracks the potential paths of formulating and propagating

microbiome diversity. Such search-based global microbiome networks, reconstructed within hours

on just one computing node, provide a readily expanded reference for tracing the origin and

evolution of existing or new microbiomes.

Background

Microbiome composition, a fundamental feature of all microbiota in nature, is shaped by a plethora of environmental factors such as habitats, geographic locations, temperature, oxygen level and even day length [1, 2]. However, it remains unclear whether and how compositional changes at the "community to community" level among microbiomes are linked to the origin and evolution of global microbiome diversity [3-5]. For example, did microbiomes from different environments emerge and develop separately, or did the global microbiome start homologically and then spread to other habitats with compositional dispersal and dynamics? Over recent years, a large number of microbiome samples (e.g., Human Microbiome Project [6], Earth Microbiome Project [1], Tara Ocean [7], etc), mainly in the form of 16S rRNA amplicons, have been produced and accumulated [4]; however, the ability to cluster and model microbiomes at the global scale has however been hindered by the enormous volume and sheer complexity of such data (e.g. a distance matrix of 100,000 microbiomes contains $\sim5*10^9$ elements).

Results

Microbiome transition model and search-based network

Here we describe the compositional dynamics and variation among microbial communities by a *microbiome transition* model. In this model, a microbial community is essentially a combination of microorganisms, and the structure of a community can be modified to another form by adding and/or removing species by compositional dispersal or fusion [8, 9] (**Fig. S1**). Theoretically, higher similarity between two communities indicates higher probability for such microbiome transition since fewer compositional exchanges are needed; however, it is not clear what level of similarity

may indicate such microbial transition with reasonable confidence. Based on a pairwise full

permutation of similarity calculation among all microbiome samples from the Microbiome Search

Engine database (MSE [10]; 177,022 samples in total; refer to "**Microbiome sample collection**" for

details) using Meta-Storms algorithm [11, 12] (**Table 1**), we consider that *direct transition* possibly

exists between sample pairs with significant similarities that cause permutation *p*-value < 0.01

(*equation 1;* **Fig. 1**). As the result, we define the Meta-Storms similarity of 0.868 as the threshold

for direct transition between microbiomes. By further analyzing the pairwise similarity in each

habitat, we found that the threshold similarity of 0.868 is significantly high in the between-habitat

similarity distribution (*p*-value=0.0022; **Fig 1B**); moreover it is higher than the upper boundary of

most within-habitats similarities (17 of 20; **Fig 1C**). Thus the similarity threshold of 0.868 is

sufficiently stringent for defining microbiome transition among the ecosystems.

Then for each of the input 177,022 microbiomes, we applied MSE to search against all other

samples and find the top matches with similarity higher than 0.868. Based on the search results we

constructed a transition network, in which each node is a microbiome, and each edge represents a

direct transition (*equation 2*; **Fig. S2**). Collectively the network consists of 177,022 nodes (samples)

and 11,175,742 edges (each called a *direct transition*). Notably, a pair of samples with low

similarity can be connected via multiple edges (i.e., via a series of direct transitions across

intermediate transfer samples), and such a sample pair is termed an *indirect transition* (*equation 3*).

The transition network predicts microbiome habitat at a global scale

At the global scale, it is still not clear whether (and to what degree) similarity in microbiome

structure implies similarity in ecosystem features [13, 14]. To quantitatively tackle this question, we

compared the direct transition frequency between within-habitat (transitions of sample pairs from

the same habitat) and between-habitat (transitions of sample pairs between two different habitats)

cases in the transition network. For each habitat, the direct transition frequency is calculated by the

average number of direct transitions per samples in this habitat. Notably, the direct transition exists

more frequently between samples in the same habitat (**Fig. 2A**; two-tailed pair *t*-test *p*-value < 0.01).

Thus the source environment of microbiomes dominates the microbial composition. We next used

the transition network to predict the habitat (mock samples were not included) of each sample by its

top neighbors (**Methods**). Via LOOCV (Leave-One-Out Cross Validation), 89.28% samples were

correctly assigned by their original habitats (**Fig. 2B**; **Table 2**). Therefore, at a global scale,

microbiome structure is strongly correlated with their environmental features.

"Mismatches", i.e., microbiomes that were assigned to an incorrect habitat as predicted by the

transition network, represent 10.72% (18,894 of 176,211) of all samples. Such mismatches are

interesting as they can be caused by, and thus potentially indicate, frequent contact and interchanges

of microbiota among the habitats. Those among human body sites are the most frequently observed

mismatches (1.86% of all samples, same as bellow) caused by the daily contact and exchange of

microbial composition [15]. Matches between non-mammal animal (sponge) and marine water are

the second most frequent mismatches (1.76%). Mismatches across "Human Skin", "Animal (pet)"

and "Building (indoor environment)" represent 1.68% of samples, likely due to their sharing of the

indoor environment (where the microbiome was largely sourced from humans [5, 16]). Moreover,

0.81% of the mismatches are between human-gut and mammal-animal-gut, which can be explained

by the close phylogenetic relationship between human and other mammals and the coevolution of

mammals and their gut microbiota [17]. Furthermore, mismatches are also observed (0.14%) where

lake water samples are predicted as river water (note that this is the source stream of lake), or vise

versa. Therefore, although the microbiome structure at the global scale is mainly shaped by their

habitat, microbiome structure can be altered by, and thus reflect, the contact and exchange of

microbiota from different environments.

Microbiomes are connected globally by the transition network

The beta diversity of global microbiome may have evolved via two scenarios: (*i*) "polyphyly",

where microbiomes from different environments were generated and developed separately (**Fig.**

S3A), or (*ii*) "monophyly", in which microbiomes started homologically and then were dispersed to

other habitats (e.g., via compositional transition, exchange or fusion; **Fig. S3B**). To distinguish

between the two scenarios, we used the transitive closure algorithm to examine the connectivity of

this transition network (**Methods**). A closure is a set of nodes (microbiomes), in which each

microbiome can traverse to any other one by direct or indirect transitions (with finite steps). Hence

being in a closure implies likelihood of transformation among samples via compositional exchange.

Traversing all nodes in the network via the transitive closure algorithm revealed that 98.31%

samples (174,032 of 177,022) can be clustered into a single closure (also named as the *main*

closure). Under a condition that microbiota composition is distinct by habitat for 89.28% of all

samples (**Fig. 2B**), such high connectivity suggests that the likelihood of polyphyly should be very

low (probability < 1.5e-05; estimated by *equation 5*), and supports the monophyletic origin of

global microbiomes and the formation of new microbiomes via such transitions (**Fig. S3B**). Notably,

1.69% (2,990 of 177,022) samples are still not included in the *main closure*, and they were mostly

due to statistical inaccuracy (1.47% exhibit a similarity level that is only slightly below the

threshold for being recruited into the *main closure*; *p*-value between 0.01 and 0.05) or curation

errors (e.g. 0.16% are labeled as microbiome but actually pure-cultures or 18S/ITS amplicon

samples). Therefore, the monophyly hypothesis best explains the origin and evolution of

present-day microbiome structures.

To size the global microbiome network, we computed the pairwise shortest transition steps of

all sample pairs in the *main closure* using the *Dijkstra* algorithm [18] (**Methods**). Interestingly, like

the "small world" principle for social network [19], the microbiome transition network follows the

"7-degree of separation" pattern (**Fig. S4A**). Specifically, any two microbiota in the *main closure*,

even if they were sampled from different habitats and exhibit low similarity, can traverse from one

to the other with only seven direct transitions on average [20], and 32 such steps at the maximum

(i.e., the network diameter; **Fig. S4B**). Such a pattern underscores the high connectivity and thus

surprisingly close interaction among microbiomes from diverse habitats at the planetary scale.

The global microbiome network is scale-free and the connectivity is robust

Notably, in this global transition network, for each node, its edge degree *k* (number of direct

transition neighbors) follows a Poisson distribution (**Fig. 3A**; Pearson *r*=-0.836 between *log(P(k))*

and *log(k)*, i.e., $P(k) \approx k^{-\gamma}$), suggesting that the network is scale-free [21, 22]. One key feature of

a scale-free network is the stability of topology, i.e., robustness to node removal from the network.

To test the robustness, we removed different numbers of randomly selected samples and their

associated edges, then assessed both size and leftover sample rate of the *main closure* in the residual

network (percentage of residual nodes ranged from 5% to 95%; each such removal procedure was

repeated for 10 times; **Methods**). When the number of nodes in the network reaches 80,000 (45% of

total nodes), the connectivity rate curve of the *main closure* already exhibits a flat trend with 97.19%

samples (98.31% prior to sample removal; **Fig. 3B**), and moreover, the mean transition steps and

maximum transition steps (diameter) converge to 8 and 33, respectively (which were 7 and 32

before sample removal; **Fig. 3C**). Thus these parameters are quite stable, and not dependent on the

increase of total sample number in the network. These findings suggest the robustness of

microbiome diversity and similarity pattern among ecosystems at the global scale.

The microbiome transition roadmap simulates the development of global microbial diversity

among multiple ecosystems

As microbiome compositions are dominantly determined by their habitats, the full connectivity

of global microbiomes in the network suggests the ability to reconstruct how the microbial diversity

spread among different habitats at a macroscopic scale. This "microbial dispersal" roadmap can be

simulated by a sub-network that *i)* covers and links all samples, and *ii)* consists of deterministic

finite transition steps without cyclic or redundant routes. Thus we derived such a roadmap (**Fig. 4A**)

by parsing the Minimum Spanning Tree (MST) of the *main closure* using the *Kurskal* algorithm [23]

(**Methods**). As the global optimum with the highest overall transition probability (similarity), the

MST maximally captures the transition pattern of world-wide microbial diversity among all the 19

habitats (with the "mock" samples excluded). For example, marine microbiomes most probably

exchange with two other environments: one is sand which is geographically close to the shore,

while the other is non-mammal animals such as fishes. These observations also suggest that sand

and freshwater microbiomes are the "gateways" to soil, plants and human-associated habitats such

as gut, oral, skin, and the human living environments.

This roadmap is verified by the isomorphic pattern of PCoA derived from a subset of 140

samples randomly selected from six habitats (**Fig. 4B**). Moreover, we employed a marine-gut route,

which represents one of the longest transitions in this subset, to illustrate the high-resolution

transition procedures (based on phylum-level compositional variations; urogenital and gut

microbiomes were combined as they are very close in the global scale; **Fig. 4C**). Zooming in of this

marine-gut route revealed a series of structure shifts that transform a freshwater microbiome to gut

samples. Starting from an actual freshwater sample [24], in each step, organisms enriched in

freshwater [25] (e.g. *Mycoplasma* and *Escherichia*) were removed/reduced, and organisms

abundant in gut [26] (e.g. *Bacteroides* and *Parabacteroides*) were added/increased. Although a

single step might have caused just slight modification on the microbiome structure, after several

iterations this sample can be smoothly transited to gut microbiomes [27] in the network, via a series

of transfer samples (**Fig. 4D**).

Microbiome transition over time and across geography

To test the feasibility of modeling microbial dynamics by the global microbiome transition

network, we used a longitudinal cohort to describe the transition of human microbiomes across time.

In this dataset, 1,963 samples were collected from three body sites (gut, oral cavity and skin) of two

individuals (I, male; II, female) over 396 time points [28]. Our search-based network analysis

revealed that the microbiome composition of each body site exhibits significant variations across

time (**Fig. 5A**, **B** and **C**; **Fig. S5**), while skin and oral microbiomes were clustered into the same

closure by direct transition (**Fig. 5D, E and F**). These suggest that microbiome transition is ongoing

within each site and between the skin and oral sites across different time points. In addition, for

both hosts, gut samples are "isolated" from the skin-oral closure, consistent with the global

microbiome transition map (**Fig. 4A**) where gut microbiomes are in a distinct route from skin and

oral ones. Therefore, although oral cavity and gut are both of the digestive tract and moreover

microbial translocation from oral to gut can occur [29], the oral microbiota have more likely been

derived from or more prominently shaped by the skin microbiota (and vice versa) than from the gut

microbiota. This seemingly counter-intuitive finding can actually lend support from the sharing of a

more aerobic and less acidic environment by skin and oral cavity (pH and oxygen level are known

to have large effect size in microbiome structure [1].

On the other hand, to validate the connectivity of microbiomes from varied geographical

locations, we constructed a search-based network by a single dataset [30] that contains 3,850

samples collected from six habitats (human gut, human oral, non-mammal animal, plant, soil and

freshwater) and locations at North America (Urbana IL, Columbia MO, Aurora CO, Ithaca NY and

Lansing MI, etc). With this dataset, the authors concluded that there was no overlap of abundant

bacterial taxa between the microbial communities from human gut and plant roots [30]. Consistent

with this conclusion, the network-based analysis based solely on this dataset, that samples were

distributed into three isolated closures of direct transitions (**Fig. 6A**). However, once extra 1,635

samples from MSE database that connected the different closures of this local network were added,

a single closure that covers 97.74% samples and integrates the original three closures emerged, with

the newly included samples serving as "transfer nodes" (i.e., samples that link two clusters in

network; *equation 3* in **Methods**) that provide additional indirect transitions(**Fig. 6B**; **Methods**). Notably, among such "transfer" microbiomes, most (96.89%) were from the same habitats as the original dataset, and others were mainly from sand and marine, which are found as the transfer nodes among non-mammal animal, plant and soil microbiomes in our global microbiome transition roadmap (**Fig. 4**). This example demonstrates that although microbiomes from diverse environments and isolated geographical locations can have very distinct structures, they can still be linked within a single closure in the microbiome transition network, i.e., evolve from each other, as long as global beta-diversity is adequately surveyed and covered. These results, which directly challenge the conclusion that multiple host microbiota compositions were independently evolved [30], underscores the importance of deriving or validating "local" datasets under the context of our global microbiome network, particularly when discussing similarity (i.e., beta-diversity), interaction or other kinds of relationship among microbiomes.

Conclusions and Discussions

Microbiota have been co-evolving with and shaping our planet, but their origin and evolution at the global scale remain elusive, due to the lack of fossils and the methodological challenges associated with integrating and mining such large-scale high-complexity data [14]. Although "species-to-species" interactions have been mapped by co-occurrence analysis on microbiomes across various habitats [31, 32], efforts to globally traverse and interrogate the vast microbiome data space at the "community to community" level have just started [10, 33, 34]. Here we propose a microbiome transition model and a network-based analysis framework to describe and simulate the variation and dispersal of the global microbial beta-diversity across multiple habitats. Benefited

from the extremely-high search speed of Microbiome Search Engine [10], we introduced a global

microbiome network with 177,022 microbiome samples that contains 11.3 billion sequences. By

traversing such network, we showed that the microbiome structures are world-widely connected by

significant similarity that follows the "small world" principle. This endeavor reveals the inherent

homology of the global microbiome diversity and supports the monophyletic origin of all

microbiomes on Earth. Then we drew the first global microbiome transition roadmap to illustrate

the potential yet most likely paths that can explain the evolution process of global microbiomes.

Due to the ongoing exponential growth of microbiome sequencing data, current beta-diversity

analysis approaches, which mainly rely on the $O(n^2)$-complexity pairwise relations (*n* is the number

of samples) such as Principle-Coordinate-Analysis (PCoA) and clustering, have become

increasingly stressed or even impractical, particularly when computational resources are limited.

Here we tackled this challenge via a search-based network, which is built on the "neighbors", i.e.,

those with the highest similarity for each sample; this strategy reduces the computational

complexity to $O(c*n)$ (c is constant, i.e., the number of neighbors), and thus enables deciphering the

pairwise similarity for > 100,000 microbiomes within 3 hours on a single computing node. As a

result, the global microbiome transition roadmap, which will be regularly updated as community

resource, can serve as a reference for interpreting or validating those existing or future observations

on inter-microbiome similarity, association or interaction, since local microbiome datasets can be

readily aligned to this global roadmap based on their shared nodes. Moreover, such a network-based

analysis framework, which can be extended to shotgun metagenome datasets, provides a new

perspective for tracking back or predicting microbiome evolution with fine resolution yet at the

global scale.

Methods and Materials

Microbiome samples collection

We used all the microbiome samples from Microbiome Search Engine database

(http://mse.ac.cn). Samples were collected from 572 studies/projects that including 20 habitats

(**Table S1**). OTUs were picked and annotated against Greengenes [35] full length 16S rRNA gene

sequences (version 13-8) on 97% similarity level by Parallel-META 3 [36] (version 3.4.4).

Variation of 16S rRNA gene copy number was normalized based on the IMG/M database [37]. We

set a minimum sequence number of 500 and minimum 16S rRNA mapping rate of 80% for each

sample to ensure high quality of the reference datasets. Finally, N=177,022 samples with

11,302,841,991 mapped sequences assed the quality control and curation (**Table S1**).

Calculation of pairwise microbiome similarity matrix for definition of direction transition

The pairwise similarity matrix of all N=177,022 samples was entirely permuted (totally (N *

N-1)/2 = 15,668,305,731 times) to examine the distribution of microbiome phylogeny similarity

using Meta-Storms algorithm [11, 12] in Parallel-META 3 software package. By setting a cutoff

p-value < 0.01 in the permutation of the similarity (rank of top 1%), we got the Meta-Storms

similarity 0.868 as the statistical threshold of the significant high value to define the direct transition

(this threshold is also referred to as $T_{d.t}$). Thus the transition model can be described in the

following form

$$P\big(direct\ transition(a,b)\big) = \begin{cases} sim(a,b), & if\ sim(a,b) \geq T_{d.t} \\ 0, & if\ sim(a,b) < T_{d.t} \end{cases} \qquad (1)$$

in which s_i and s_j are two arbitrary microbiomes, and $sim(s_i,\ s_j)$ represents their Meta-Storms

similarity.

Search-based microbiome network

The search-based microbiome network is built using Microbiome Search Engine (MSE) [10].

For each sample we searched it against all other samples for the top 100 matches, and connected it

with the matched samples that have similarity higher than the threshold of direct transition

($T_{d.t}$=0.868). By iterating such search with all samples, we constructed a global network G (*equation*

1).

$$G = (V = \{a, b, \dots\}, E = \{edge(a,b), \dots\})$$
$$(2)$$
$$edge(a,b) \rightarrow sim(a,b) \geq T_{d.t}$$

262   In this network, one node (e.g. *a* or *b* in *equation 2*) is a single microbiome, and edges (e.g.

263   *edge(a, b)* in *equation 2*) that link the nodes are direct transitions (**Fig. S2**). Finally in the network

264   there were 177,022 nodes (samples) and totally 11,175,742 edges (direct transitions). In this

265   network, a pair of samples with low similarity can be connected by a path of multiple edges, i.e., an

266   *indirect transition*:

$$indirect\ transition(a,b) \rightarrow sim(a,b) < T_{d.t}$$
$$(3)$$
$$\exists\ edge(a,x_1), edge(x_2,x_3), \dots edge(x_{i-1},x_i), edge(x_i,b)$$

Then here x_1, x_2, …, x_i are defined as the "transfer samples" for that underlie the indirect

transition from a to b.

Prediction of habitat using microbiome network

In the network G we predicted the source habitat of each microbiome by its top $n=10$ neighbor

samples and similarities. For an arbitrary microbiome sample a in the network, similarities to its top

10 neighbors are $S = \{s_1, s_2, ..., s_n\}$, while the n neighbor were from m $(1 \leq m \leq n)$ different

habitats as $H = \{h_1, h_2, ..., h_m\}$, then the probability for the predicted habitat of microbiome a as h_k

is calculated by

275
$$P(a \in h_k) = \frac{\sum_{j \in h_k}\left((n-j+1) * s_j\right)}{\sum_{i=1}^{n}\left((n-i+1) * s_i\right)} \qquad (4)$$

Here $j \in h_k$ means the habitat of neighbor j is h_k $(1 \leq k \leq m)$. Then the predicted habitat that has the

highest probability P will be assigned to the sample a as the prediction results.

Probability of global transition among all habitats in the network

By *equation 4* we calculated that at the global scale the microbiota composition is distinct by

habitat, that the probability of transition among the same habitat is 89.28% (**Fig. 2B**). To calculate

the overall probability of connecting all habitats in the transition network, we can start from

connecting arbitrary two habitats of which the probability $p_{transition}(n=2)$ = 1-89.28%. When one

more habitat added into the network, the probability of transitions among the three habitats can be

calculated as $p_{transition}(n=3) = p_{transition}(n=2)$ * (1-89.28%^{2}), where the square of 89.28% represents

the probability that no direct transition between the added habitat and the former two habitats. Then

we can expand such procedure to estimate the probability for connecting *n* habitats in the transition

network by

$$p_{transition}(n) = \begin{cases} 1 - 89.28\% & , \quad n = 2 \\ p_{transition}(n-1) \times (1 - 89.28\%^{(n-1)}), & \quad n > 2 \end{cases} \quad (5)$$

Transitive closure algorithm of microbiome network

In the microbiome transition network, a closure is a subset of nodes (microbiome) are fully

connected, that each microbiome is possible to link to any other sample by direct or indirect

transitions (with finite transfer nodes). Closures can be initialized by an arbitrary node in the

network, and then expanded by adding more external nodes that are directly connected with this

closure (**Fig. S6**). If two or more closures are connected by any edge, these closures can also be

merged as one closure. By the traversal among all nodes in the network *G* we get a *main closure C*

with 98.31% samples.

Size of the microbiome network

In the *main closure C*, there are always multiple routes between two indirectly connected nodes

(microbiomes). We count the edge between two directly linked nodes as 1, so the length of indirect

routes is the number of transfer nodes + 1 on this route (**Fig. S4**). We used the *Dijkstra* algorithm

[18] by Python package *igraph* (0.7.1 running inside Python 3.6.1) to find the pairwise shortest

transition steps (with smallest number of transfer nodes) between all indirectly linked node pairs in

the *main closure C*. Thus the number of maximal steps among the shortest route is the diameter of

the closure. The diameter means in this closure, any two microbiomes could be linked to each other

by a route with steps that smaller than the diameter.

Minimum spanning tree for the roadmap of microbiome network

306   In a transitive closure, a spanning tree is a sub-network that connects all nodes (microbiomes)

307   with no cycle. For two directly linked samples *a* and *b*, we define their distance as

308
$$dist(a,b) = 1 - sim(a,b) \quad (6)$$

the Minimum Spanning Tree (MST) could be consider as the global transition path of all samples

with the highest overall transition probability since it links all samples with the shortest total

distance. In the *main closure C* we used the Kruskal algorithm [23] to calculate the 2-level MST to

reflect the transition among different habitats from the global scale.

The first level MST was on "sample resolution", based on which we then make the second

MST on "habitat resolution". Initially we calculated the first level MST of the *main closure C*, and

then generated the habitat-based network *G'* (*equation 2*), each node represents one habitat, and

distance between two habitats h_i and h_j is the average distance of all edges that linked the two

habitats in the MST. Then we computed the second-level MST (*G'*), which illustrates the global

microbiome transition roadmap across multiple habitats.

$$G' = \left(V' = \{h_i, h_j, \dots\}, E' = \{average\left(edge(a,b)\right)\}\right)$$
$$(7)$$
$$a \in h_i, b \in h_j, edge(a,b) \in MST(C)$$

The significance of the roadmap (MST(*G'*)) was assessed by the permutation test of the

topologically equivalent sub-network in the *main closure C* of the original network. Specifically, in

a permutation, for each edge that connects two habitats (eg. *habitat$_i$* and *habitat$_j$*) in the roadmap,

we also randomly selected an edge that connects two samples (eg. sample *a* and *b*) respectively

from these two habitats (eg. $a \in habitat_i$ and $b \in habitat_j$). Since we iterated the permutation for

10,000 times, if the total distance of the roadmap is smaller than 99% of permutated network (also

means the total probability is in top 1%, *p*-value < 0.01), we can consider the roadmap MST(G') is

significant in the *main closure C*.

Search-based sample selection from reference database to link separated closures

To select transfer samples from a reference database to link two separated closures, we search

all samples of each closure against the referenced repository for top matches with higher similarity

than the direct transition cutoff ($T_{d.t}$=0.868), and the overlapped matches between the two closures

are the transfer microbiomes that link the two closures. If there is no overlap in the matches, we

then extend each of the closure by adding their matches, and repeat the search process until we find

any transfer sample. On the other hand, once closures cannot be furtherly extended by database

search but still no available transfer sample is found, this means that no sample in the reference

database is able to work as the transfer node to link the two separated closures by direct transition.

Acknowledgements

J.X. acknowledges support of grants 31327001, 31425002 and 81430011 from National Natural

Science Foundation of China (NSFC), KFZD-SW-219-4 and ZDBS-SSW-DQC002-03 from Chinese

Academy of Sciences (CAS). X.S. acknowledges support of grants 31771463 and 32070086 from

NSFC. R.K. acknowledges support from the U.S. National Institutes of Health, the National Science

Foundation, and the Alfred P. Sloan Foundation. We thank the Operations Research

Community-QIBEBT for the inspiring discussions.

Author contributions

J.X. and X.S. conceived the idea. X.S., and G.J., developed the software and algorithm. G.J., and

Y.Z. performed analysis. L.L., Z.X., Z.W. and Z.S. contributed to data collection and curation. X.S

and J.X. wrote the manuscript.

Competing interests

The authors declare that they have no competing interests.

Cod and data availability

All   the   source   code   and   source   data   used   in   this   work   are   available   at

https://github.com/qibebt-bioinfo/microbiomenetwork.

Figures

Fig. 1. Distribution of pairwise similarity in *N*=177,022 microbiome samples. (**A**) *P*-value<0.01 of significant similarity values in the permutation determine the threshold of 0.868 (under the shadow) for putative direct transition. (**B**) The threshold has *p*-value=0.0022 among between-habitat similarity distribution. (**C**) The threshold is higher than the upper boundary of within-habitat similarities for most habitats. Three panels use the same Y-axis. *P*-values are calculated by permutation test.

Fig. 2. Global microbiome network predicts the microbiome habitat. (**A**) Frequency of within-habitat direct transition is significantly higher than that of between-habitat. *P*-value is calculated by two-sided *t*-test. (**B**) Habitat of 89.28% samples is correctly predicted by the microbiome network. Inner ring represents the proportion of real habitats and the outer ring is the proportion of predicted habitats.

Fig. 3. Robustness of the global microbiome network. (**A**) Node degree (number of linked neighbors) of the network follows the Poisson distribution, suggesting the network is scale-free. The effect of random node removal on the *main closure* in (**B**) sample rate, (**C**) mean shortest transition step and maximum transition step (diameter).

Fig. 4. Roadmap of the global microbiome transition among habitats. (**A**) Bold arrow lines are the roadmap that represents the maximum overall similarity. The number of samples in each habitat is scaled by the node size, and the within-habitat transition frequency is represented by the node color depth (compared to the rim). (**B**) The PCoA parsed from a subset of 140 microbiomes demonstrates the roadmap by the equivalent topology. (**C**) The phylum-level compositional shift of a microbiome transition route for marine to gut environment. (**D**) The genus-level compositional shift of a transition case from a freshwater microbiome to gut samples.

Fig. 5. Transition of human microbiome across time and body sites. The within-habitat microbiome transition of gut (**A**), oral (**B**) and skin (**C**) of Individual I across 396 time points. (**D**) The oral-skin microbiome transition of individual I across time. Only selected samples were

shown. The transition patterns of Individual II are shown in **Fig. S5**. **E** and **F**, PCoA of the two individuals' time-series microbiomes: skin and

oral microbiomes are linked in a closure by direct transition (highlighted by grey dotted line) and gut samples forming another closure.

Fig. 6. Microbiome transition across habitats and geographical locations. (**A**) The 3,850 samples from six habitats are included in three isolated transition closures, of which the sample proportions are 43.22%, 7.90% and 45.53%, respectively. (**B**) After adding extra 1,635 bridge samples from the MSE database, three closures are merged into a single closure by direct transition, which contains 97.74% samples.

Tables

Table 1. The distribution of samples among the habitats.

Sample type	Habitat	Number of samples
Human associated	Gut	51,706
	Skin	19,455
	Oral	10,896
	Other human body-site	3,018
	Urogenital	1,204
	Nose	489
Animal associated	Mammal animal	29,918
	Non-mammal animal	11,172
Environmental	Building	11,248
	Soil	10,507
	Marine water	6,090
	Lake	4,234
	Plant	3,456
	Freshwater	3,112
	River	2,248
	Milk	1,636
	Sand	968
	Food	780
Other	Other	4,074
	Mock	811
Total		**177,022**

Table 2. Prediction of habitat based on the microbiome network.

Habitat	Number of samples	Number of correctly predicted samples	Accuracy	Within-habitat transition frequency	Between-habitat transition frequency
Gut	51,706	50,431	97.53%	66.89	5.40
Skin	19,455	17,464	89.77%	52.11	17.82
Oral	10,896	10,070	92.42%	55.10	10.98
Other human body-site	3,018	1,777	58.88%	24.64	37.92
Urogenital	1,204	1,046	86.88%	44.16	17.29
Nose	489	91	18.61%	5.00	55.29
Mammal animal	29,918	28,010	93.62%	50.67	8.87
Non-mammal animal	11,172	8,077	72.30%	19.81	19.95
Building	11,248	8,942	79.50%	40.24	24.54
Soil	10,507	9,978	94.97%	54.49	7.28
Marine water	6,090	3,960	65.02%	24.29	17.77

Lake	4,234	3,983	94.07%	49.64	7.35
Plant	3,456	3,127	90.48%	46.91	15.16
Freshwater	3,112	1,671	53.70%	22.51	28.66
River	2,248	2,011	89.46%	36.84	15.55
Milk	1,636	1,565	95.66%	55.21	8.23
Sand	968	864	89.26%	44.72	23.54
Food	780	677	86.79%	39.01	26.57
Other	4,074	3,573	87.70%	38.88	13.64

Reference

1.   Thompson, L.R., et al., *A communal catalogue reveals Earth's multiscale microbial diversity.* Nature, 2017. **551**(7681): p. 457-463.
2.   Caporaso, J.G., et al., *Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.* Proceedings of the National Academy of Sciences of the United States of America, 2011. **108**: p. 4516-4522.
3.   Mihaljevic, J.R., *Linking metacommunity theory and symbiont evolutionary ecology.* Trends Ecol Evol, 2012. **27**(6): p. 323-9.
4.   Brooks, A.W., et al., *Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host Evolutionary History.* PLoS Biol, 2016. **14**(11): p. e2000225.
5.   Groussin, M., et al., *Unraveling the processes shaping mammalian gut microbiomes over evolutionary time.* Nat Commun, 2017. **8**: p. 14319.
6.   Integrative, H.M.P.R.N.C., *The Integrative Human Microbiome Project.* Nature, 2019. **569**(7758): p. 641-648.
7.   Sunagawa, S., et al., *Ocean plankton. Structure and function of the global ocean microbiome.* Science, 2015. **348**(6237): p. 1261359.
8.   Charubin, K., et al., *Interspecies Microbial Fusion and Large-Scale Exchange of Cytoplasmic Proteins and RNA in a Syntrophic Clostridium Coculture.* mBio, 2020. **11**(5).
9.   Sarkar, A., et al., *Microbial transmission in animal social networks and the social microbiome.* Nature Ecology & Evolution, 2020. **4**(8): p. 1020-1035.
10.   Su, X., et al., *Identifying and Predicting Novelty in Microbiome Studies.* MBio, 2018. **9**(6).
11.   Su, X., et al., *GPU-Meta-Storms: computing the structure similarities among massive amount of microbial community samples using GPU.* Bioinformatics, 2014. **30**(7): p. 1031-3.
12.   Su, X., J. Xu, and K. Ning, *Meta-Storms: Efficient Search for Similar Microbial Communities Based on a Novel Indexing Scheme and Similarity Score for Metagenomic Data.* Bioinformatics, 2012.
13.   Shinichi, S., et al., *Ocean plankton. Structure and function of the global ocean microbiome.* Science, 2015. **348**(6237): p. 1261359.
14.   Kyrpides, N.C., E.A. Eloe-Fadrosh, and N.N. Ivanova, *Microbiome Data Science: Understanding Our Microbial Planet.* Trends in Microbiology, 2016. **24**(6): p. 425-427.
15.   Huttenhower, C., et al., *Structure, function and diversity of the healthy human microbiome.* Nature, 2012. **486**(7402): p. 207-214.
16.   Lax, S., et al., *Longitudinal analysis of microbial interaction between humans and the indoor environment.* Science, 2014. **345**(6200): p. 1048-1052.
17.   Muegge, B.D., et al., *Diet Drives Convergence in Gut Microbiome Functions Across Mammalian Phylogeny and Within Humans.* Science, 2011. **332**(6032): p. 970-974.
18.   Dijkstra, E.W., *A note on two problems in connexion with graphs.* Numerische mathematik, 1959. **1**(1): p. 269-271.
19.   Milgram, S., *The small world problem.* Psychology today, 1967. **2**(1): p. 60-67.
20.   Leskovec, J. and E. Horvitz, *Planetary-scale views on a large instant-messaging network*, in *Proceedings of the 17th international conference on World Wide Web*. 2008, Association for Computing Machinery: Beijing, China. p. 915–924.
21.   Barabasi, A.L., *Scale-free networks: a decade and beyond.* Science, 2009. **325**(5939): p. 412-3.

22. Jeong, H., et al., *The large-scale organization of metabolic networks.* Nature, 2000. **407**(6804): p. 651-4.

23. Kruskal, J.B., *On the shortest spanning subtree of a graph and the traveling salesman problem.* Proceedings of the American Mathematical society, 1956. **7**(1): p. 48-50.

24. Dunn, R.R., et al., *Home Life: Factors Structuring the Bacterial Diversity Found within and between Homes.* Plos One, 2013. **8**(5).

25. Lee, C.S., et al., *The Microbiota of Recreational Freshwaters and the Implications for Environmental and Public Health.* Front Microbiol, 2016. **7**: p. 1826.

26. Human Microbiome Project, C., *Structure, function and diversity of the healthy human microbiome.* Nature, 2012. **486**(7402): p. 207-14.

27. Song, S.J., et al., *Preservation Methods Differ in Fecal Microbiome Stability, Affecting Suitability for Field Studies.* mSystems, 2016. **1**(3).

28. Caporaso, J.G., et al., *Moving pictures of the human microbiome.* Genome Biol, 2011. **12**(5): p. R50.

29. Schmidt, T.S., et al., *Extensive transmission of microbes along the gastrointestinal tract.* Elife, 2019. **8**.

30. Hacquard, S., et al., *Microbiota and Host Nutrition across Plant and Animal Kingdoms.* Cell Host Microbe, 2015. **17**(5): p. 603-16.

31. Ma, B., et al., *Earth microbial co-occurrence network reveals interconnection pattern across microbiomes.* Microbiome, 2020. **8**(1): p. 82.

32. Faust, K., et al., *Microbial co-occurrence relationships in the human microbiome.* PLoS Comput Biol, 2012. **8**(7): p. e1002606.

33. Su, X., et al., *Multiple-Disease Detection and Classification across Cohorts via Microbiome Search.* mSystems, 2020. **5**(2): p. e00150-20.

34. Zhu, M., K. Kang, and K. Ning, *Meta-Prism: Ultra-fast and highly accurate microbial community structure search utilizing dual indexing and parallel computation.* Brief Bioinform, 2020.

35. McDonald, D., et al., *An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea.* ISME J, 2012. **6**(3): p. 610-8.

36. Jing, G., et al., *Parallel-META 3: Comprehensive taxonomical and functional analysis platform for efficient comparison of microbial communities.* Scientific Reports, 2017. **7**: p. 40371.

37. Markowitz, V.M., et al., *IMG/M: a data management and analysis system for metagenomes.* Nucleic Acids Research, 2008. **36**(suppl 1): p. D534-D538.