1 **FcγR responses to soluble immune complexes are governed by solubility and size.**

2 Haizhang Chen[1,2], Andrea Maul-Pavicic[3,4], Martin Holzer[5], Ulrich Salzer[3], Nina Chevalier[3], Reinhard E.

3 Voll[3,4], Hartmut Hengel[1,2,#] & Philipp Kolb[1,2,#]

[1]Institute of Virology, University Medical Center, Albert-Ludwigs-University Freiburg, Hermann-Herder-Str. 11, 79104 Freiburg, Germany
[2]Faculty of Medicine, Albert-Ludwigs-University Freiburg, 79104 Freiburg, Germany
[3]Department of Rheumatology and Clinical Immunology, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Hugstetterstr. 55, 79106 Freiburg, Germany
[4]Center for Chronic Immunodeficiency (CCI), Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, Breisacherstr. 115, 79106 Freiburg, Germany
[5]Institute for Pharmaceutical Sciences, Albert-Ludwigs-University Freiburg, Hermann-Herder-Str. 9, 79104 Freiburg, Germany
#corresponding authors

Summary

This study describes a novel cell-based reporter assay enabling the detection and

quantification of soluble multimeric IgG immune complexes (sICs). Receptor triggering of sICs

is restricted to specific FcγRs and depends on sIC size. The assay identifies sICs in sera from

SLE patients and autoimmune-prone mice as a novel biomarker of autoimmune disease.

Abstract

Fcγ-receptor (FcγR) activation by antibody formed soluble immune complexes (sICs) is

thought to be a major mechanism of inflammation in certain autoimmune diseases such as

systemic lupus erythematosus (SLE). A robust and scalable test system allowing for the

detection and quantification of sICs bioactivity is missing. We developed a comprehensive

reporter cell panel capable of measuring the sIC-mediated activation of individual human and

mouse FcγRs. We show that compared to human FcγRs IIB and III, human FcγRs I and IIA

lack sensitivity to sICs. Further, the assay proved to be sensitive to sIC stoichiometry and size

enabling us to demonstrate for the first time a complete translation of the Heidelberger-Kendall

precipitation curve to FcγR responsiveness. The approach was applied to quantify sICs-

mediated FcγR activation using sera from SLE patients and mouse models of lupus and

arthritis. Thus, in clinical practice, it might be employed as a test matrix toolbox for FcγR

activation evaluating sICs as a biomarker for disease activity in immune-complex mediated

diseases.

Introduction

Immunoglobulin G (IgG) is the dominant immunoglobulin isotype in chronic infections and in antibody-mediated autoimmune diseases. The multi-faced effects of the IgG molecule rely both on the F(ab) regions, which recognize a specific antigen to form immune complexes (ICs), and the constant Fc region (Fcγ), which is detected by Fcγ receptors (FcγRs) found on most cells of the immune system [1]. When IgG binds to its antigen ICs are formed, which, depending on the respective antigen, are either cell-bound or soluble (sICs). The composition of sICs is dependent on the number of epitopes recognized by IgG on a single antigen molecule and the ability of the antigen to form multimers. Fcγ-FcγR binding is necessary but not sufficient to activate FcγRs since physical receptor cross-linking generally underlyies receptor activation [2-5]. Firmly fixed cell bound ICs are readily able to cross-link FcγRs [4, 6]. This condition induces various signaling pathways [7-9] which in turn regulate immune cell effector functions [10, 11]. It is also suggested that sICs can dynamically tune FcγR triggering, implying that changes in sIC size directly impact FcγR responses [6]. However, the molecular requirements are largely unknown. Also, a functional reproduction of the paradigmatic Heidelberger-Kendall precipitation curve, describing that the molecular size of sICs determined by the antibody:antigen ratio dynamically tunes FcγR activation on and off [12, 13], is so far missing.

In contrast to FcγR binding of monomeric ligands without consequences, IC-mediated FcγR cross-linking initiates the full signal cascade followed by immune cell activation or inhibition, resp. [5, 8, 14, 15]. Human FcγRs are membrane resident receptors recognizing Fcγ. Among all type I FcγRs, FcγRIIB (CD32B) is the only inhibitory one signaling via immunoreceptor tyrosine-based inhibitory motifs (ITIMs) while the activating receptors are associated with immunoreceptor tyrosine-based activation motifs (ITAMs). Another exception is FcγRIIIB (CD16B), which is glycosylphosphatidylinositol (GPI)-anchored [16-19]. FcγRI (CD64) is the only receptor with high affinity binding to monomeric IgG not associated with antigen and is primarily tasked with phagocytosis linked to antigen processing and pathogen clearance [20, 21]. All the other FcγRs only efficiently bind to complexed, meaning antigen-bound IgG [1, 16, 17]. While FcγRI, FcγRIIB and FcγRIIIA are able to recognize sICs, this has not been reported for FcγRIIA

(CD32A), rather this receptor has recently been shown to depend on the neonatal Fc receptor

(FcRn) to do so [22, 23].

Activation of FcγRs leads to a variety of cellular effector functions such as antibody-dependent

cellular cytotoxicity (ADCC) by natural killer (NK) cells via FcγRIIIA, antibody-dependent

cellular phagocytosis (ADCP) by macrophages via FcγRI, cytokine and chemokine secretion

by NK cells and macrophages via FcγRIIIA. Further effector responses are reactive oxygen

species (ROS) production of neutrophils and neutrophil extracellular traps formation (NETosis)

via FcγRIIIB, dendritic cell (DC) maturation and antigen presentation via FcγRIIA and B cell

selection and differentiation via FcγRIIB [10, 24-30]. Consequently, FcγRs connect and regulate

both innate and adaptive branches of the immune system. Various factors have been indicated

to influence the IC-dependent FcγR activation profiles, including Fcγ-FcγR binding affinity and

avidity [31], IgG subclass, glycosylation patterns and genetic polymorphism [4, 24, 25, 32],

stoichiometry of antigen-antibody-ratio [6, 30, 33] and FcγR clustering patterns [34]. Specifically,

Asn297-linked glycosylation patterns of the IgG Fc domain initiate either pro- or anti-

inflammatory effector pathways by tuning the binding affinity to activating versus inhibitory

FcγRs, respectively [35]. However, despite being explored in proof-of-concept studies, the

functional consequences of these ligand features on a specific receptor are still not fully

understood. Therefore, an assay platform allowing for the systematic functional assessment

of IC-mediated FcγR activation is strongly required. sICs and immobilized ICs represent

different stimuli for the immune system [22, 29]. Soluble circulating ICs are commonly associated

with certain chronic viral or bacterial infections [36, 37] and autoimmune diseases, such as

systemic lupus erythematosus (SLE) or rheumatoid arthritis (RA) [38-40]. When accumulating and

deposited in tissues sICs can cause local damage due to inflammatory responses, classified

as type III hypersensitivity [41]. Typically, sICs related disorders are characterized by systemic

cytokine secretion which can be followed by immune cell exhaustion and senescence [42, 43]. In

order to study sIC-dependent activation of FcγRs in detail, we employed a cell-based assay

which has been previously utilized to study immobilized ICs [44] and adapted it into a sIC

sensitive reporter system capable of distinguishing the activation of individual FcγRs and their

responses to varying complex size. This approach allowed for the first time a complete

reproduction of the Heidelberger-Kendall precipitation curve measuring actual FcγR triggering.

The assay enables a quantification of clinically relevant sICs in sera from SLE patients and

autoimmune-prone mice with immune-complex-mediated arthritis and lupus using reporter

cells expressing chimeric mouse FcγRs.

Results

Experimental assay setup

The assay used in this study was adapted from a previously described cell-based FcγR

activation assay designed to measure receptor activation in response to opsonized virus

infected cells [44] or therapeutic Fc-fusion proteins [45]. We complemented the assay setup to

enable measurement of sICs when incubated with reporter cells stably expressing the

ectodomains of the human FcγR fused to the signaling module of the mouse CD3-ζ chain

(FcγRI: Acc# LT744984; FcγRIIA(R): Acc# M28697; FcγRIIB/C: Acc# LT737639;

FcγRIIIA/B(V): Acc# LT737365). Ectodomains of FcγRIIIA and FcγRIIIB as well as

ectodomains of FcγRIIB and FcγRIIC are identical. Second generation reporter cells were

generated to improve stable expression of chimeric FcγRs compared to the transfectants used

in the original assay [44]. To this end, BW5147 cells were transduced as described previously

via lentiviral transduction [44, 46]. Human FcγR expression on transduced cells after puromycin

selection is shown in Fig. 1A. FcγR triggering is translated into activation of the reporter cells

and measured by quantification of endogenous IL-2 (mIL-2) secretion into the cell culture

supernatant using an anti-IL-2 sandwich ELISA as described previously [44]. In order to employ

the original assay, designed to detect ICs on adherent virus-infected cells, for the detection of

soluble ICs we first determined the suspension of IgG achieved by pre-incubating a 96 well

ELISA microtiter plate with PBS/FCS blocking buffer. To this end, we compared graded

concentrations of FCS in the blocking reagent and measured the threshold at which IgG

(rituximab, Rtx) was no longer adsorbed to the plate and stayed abundantly in solution. Fig.

1B demonstrates that FCS supplementation to 1% (v/v) or higher is sufficient to keep IgG

antibodies in solution and prevents binding to the plastic surface. Using this adapted protocol,

the assay now allowed for the sensitive detection and characterization of FcγR interaction with

immobilized ICs versus sICs as shown schematically in Fig. 1C. First, we set out to test if

monomeric IgG upon immobilization becomes an operational surrogate for IgG-opsonized cells

or immobilized ICs with regard to FcγR activation as suggested before [47].

Fig. 1. Establishment of a cell-based reporter assay measuring FcγR activation in
response to sICs. A) BW5147 reporter cells stably expressing human FcγR-ζ chain chimeras

or BW5147 parental cells were stained with FcγR specific conjugated mAbs as indicated and
measured for surface expression of FcγRs via flow cytometry. Fcγ binding was determined
using a PE-TexasRed-conjugated human IgG-Fc fragment. B) FCS coating of an ELISA
microtiter plate allows for suspension of subsequently added IgG. Plate bound IgG was
quantified via ELISA. PBS supplemented with >1% FCS (v/v) avoids adhesion of IgG
(rituximab, Rtx) to the ELISA plate. C) Schematic of an immobilized IC or soluble IC setup.
BW5147 reporter cells expressing chimeric human FcγR receptors secrete IL-2 in response to
FcγR activation by clustered IgG. Soluble ICs are generated using mAbs and multivalent
antigens (blue). Solubility is achieved by pre-blocking an ELISA plate using PBS supplemented
with 10% FCS (FCS coat, orange).

As depicted in Fig. 2, there was no qualitative difference in FcγR activation between

immobilized Rtx, immobilized ICs (Rtx + rec. CD20) or Rtx-opsonized 293T-CD20 cells,

showing that FcγR cross-linking by clustered IgG alone is sufficient for receptor cross-linking

and activation. In contrast, sICs formed by monomeric CD20 peptide (aa 141-188) and Rtx

failed to activate FcγRs even at very high ligand concentrations. Based on this finding we

hypothesized that, in order to generate sICs able to activate FcγRs, antigens have to be

multivalent to achieve cross-linking of FcγRs. Of note, to reliably and accurately differentiate

between soluble and immobilized triggers using this assay, reagents for the generation of ICs

need to be of very high purity and consistent stability. Indeed, only combinations of

pharmaceutical therapy grade, i.e. ultra-pure mAbs and pure antigens (purified via size

exclusion chromatography) showed reproducible, dose-dependent and specific activation in

the reporter assay (data not shown).

Fig. 2. FcγRIIB/C and FcγRIIIA/B do not respond to small hetero-dimeric sICs but are
sensitive to immobilized IgG/ICs. Dose-dependent activation of FcγR-bearing reporter cells
by immobilized IC can be mimicked by immobilized IgG. Response curves of human FcγRIIB/C
and FcγRIIIA/B are similar between opsonized cells (293T cells stably expressing CD20 + Rtx),
immobilized IC (rec. soluble CD20 + Rtx) and immobilized IgG (Rtx). sIC formed by monovalent

antigen (rec. soluble CD20) do not activate human FcγRs. X-Axis shows sample concentration
determined by antibody molarity. Y-Axis shows FcγR activation determined by reporter cell IL-
2 production.

Quantification of human FcγR responsiveness to multimeric sICs

To date, there are only few commercially available human IgG-antigen pairs that meet both

the above mentioned high grade purity requirements while also providing a multimeric antigen.

In order to meet these stipulations of ultra-pure synthetic soluble immune complexes we

focused on three pairs of multivalent antigens and their respective mAbs that were available

in required amounts enabling large-scale titration experiments; trimeric rhTNFα:IgG1 infliximab

(TNFα:Ifx), dimeric rhVEGFA: IgG1 bevacizumab (VEGFA/Bvz) and dimeric rhIL-5: IgG1

mepolizumab (IL-5/Mpz). As lymphocytes express TNFα-receptors I and II while not

expressing receptors for IL-5 or VEGFA, we tested whether the mouse lymphocyte derived

BW5147 thymoma reporter cell line is sensitive to high concentrations of rhTNFα. Toxicity

testing revealed that even high concentrations up to 76.75 nM rhTNFα did not affect viability

of reporter cells (Fig. S1). Next, we measured the dose-dependent activation of human FcγRs

comparing immobilized IgG to soluble ICs using the FcγR reporter cell panel (Fig. 3). Soluble

antigen or mAb alone served as negative controls showing no background activation even at

high concentrations. Immobilized rituximab served as a positive control for inter-experimental

reference. We observed that all FcγRs are strongly activated in a dose-dependent manner

when incubated with immobilized IgG. Incubating the FcγR reporter cells with sICs at identical

molarities showed FcγRIIB/C and FcγRIIIA/B to be efficiently activated by sICs, while in

contrast, FcγRIIA and FcγRI did not respond to sICs. We furthermore observed FcγRIIIA/B to

be efficiently activated by sICs with responses even surpassing those achieved with

immobilized IgG for TNFα/Ifx and IL-5/Mpz ICs. FcγRIIB/C showed a generally weaker

reactivity towards sICs compared to immobilized ICs, especially at high concentrations

whereas an inversion of this order was seen for TNFα/Ifx and VEGFA/Bvz ICs at lower

concentrations. IL-5/Mpz, FcγRIIB/C and FcγRIIIA/B showed similar responsiveness towards

immobilized or sICs with a generally stronger activation on immobilized ICs. These findings

185   demonstrate that sICs of different composition vary in the resulting FcγR activation pattern,

186   most likely due to the antigens being either dimeric, trimeric or different in size.

Fig. 3. FcγRIIB/C and FcγRIIIA/B are activated by sICs formed from multivalent antigens.
Three different multivalent ultra-pure antigens (Ag) mixed with respective therapy-grade mAbs
were used to generate sICs as indicated for each set of graphs (top to bottom). IC pairs:
infliximab (Ifx) and rhTNFα; mepolizumab (Mpz) and rhIL-5; bevacizumab (Bvz) and rhVEGFA.
X-Axis: concentrations of stimulant expressed as molarity of either mAb or Ag monomer and
IC (expressed as mAb molarity) at a mAb:Ag ratio of 1:2. Soluble antigen or soluble antibody
alone served as negative controls and were not sufficient to activate human FcγRs. FcγR
responses were normalized to immobilized rituximab (Rtx) at 1µg/well (set to 1) and a medium
control (set to 0). All FcγRs show dose-dependent activation towards immobilized IgG. FcγRIIA
shows low activation at high sIC concentrations compared to immobilized IgG activation. FcγRI
shows no activation towards sICs. FcγRIIIA/B and FcγRIIB/C are dose-dependently activated
by sICs with responses comparable in strength to immobilized IgG stimulant. Experiments

200 performed in technical replicates. Error bars = SD. Error bars smaller than symbols are not
201 shown.
202

As we observed differences in responses to sICs vs. immobilized ICs for individual FcγR-ζ

reporter cells, this indicates that FcγR ectodomains alone are already able to differentiate

between these triggers. To validate this observation in principle, we determined FcγRIIIA

activation using primary human NK cells isolated from PBMCs of healthy donors. NK cells were

chosen as they mostly express only one type of FcγR similar to our reporter system. Measuring

a panel of activation markers and cytokine responses by flow cytometry, we observed a

differential activation pattern depending on ICs being soluble or immobilized at equal molarity

(Fig. 4A). We chose IL-5/Mpz sICs as NK cells do not express the IL-5 receptor. While MIP1-

β responses were comparable between the two triggers, degranulation (CD107a) and TNFα

responses showed a trend towards lower activation by sICs compared to immobilized IgG

(Mpz). Strikingly, IFNγ responses were significantly weaker when NK cells were incubated with

sICs compared to immobilized IgG. In order to confirm this hierarchy of responses and to

enhance the overall low activation by Fcγ compared to the PMA control, we changed the IC

setup by generating reverse-orientation sICs consisting of human FcγR-specific mouse mAbs

and goat-anti-mouse IgG F(ab)$_2$ fragments. NK cell activation by reverse sICs was compared

to NK cell activation by immobilized FcγR specific mAbs (Fig. 4B). Here, we not only confirm

our previous observations regarding MIP-1β and IFNγ, but we also confirm significantly lower

TNFα and CD107a responses towards soluble complexes compared to immobilized mAbs.

Importantly, these experiments validate that sICs readily activate primary NK cells and induce

immunological effector functions. As in roughly 10% of the population NK cells express

FcγRIIC [48-51], we also tested if this receptor plays a role in our measurements. Using the same

three donors and an FcγRII specific mAb as described above, we did not observe an FcγRII-

mediated response. Accordingly, we conclude that FcγRIIC expression did not play a role in

our experiments (Fig. 4C). Taken together, we show that multivalent but not dimeric soluble

immune complexes govern primary NK cell response and FcγRIIIA/B activation (Fig. 2A).

Fig. 4. The FcγRIII-dependent activation pattern of primary NK cells depends on IC solubility. Negatively selected primary NK cells purified from PBMCs of three healthy donors were tested for NK cell activation markers. Error bars = SD. Two-way ANOVA (Turkey). A) NK cells were incubated with immobilized IgG (mepolizumab, Mpz), soluble IC (Mpz:IL-5 = 1:1), soluble Mpz or soluble IL-5 (all at 200 nM, 10^6 cells). Incubation with PMA and Ionomycin (Iono) served as a positive control. Incubation with medium alone served as a negative control. B) NK cells were incubated for 4 h with immobilized FcγRIII-specific mAb, soluble mouse-anti-human IgG F(ab)₂ complexed FcγRIII-specific mAb (reverse sICs), immobilized IgG of non-FcγRIII-specificity (isotype control) or soluble F(ab)₂ complexed isotype control (all at 1µg, 10^6 cells). Incubation with PMA and Ionomycin served as a positive control. Incubation with medium alone served as a negative control. C) As in B using an FcγRII-specific mAb. NK cells from the tested donors in this study do not react to FcγRII activation.

Measurement of FcγR activation in response to changing sIC size.

We observed that the dimeric sIC CD20:Rtx molecule complex completely failed to trigger FcγRs, while potentially larger sICs based on multimeric antigens showed an efficient dose-dependent FcγR activation. In order to determine whether BW5147 FcγR-ζ reporter cells are able to respond to changes in sIC size, we tested cross-titrated amounts of antibody (mAb, infliximab, Ifx) and antigen (Ag, rhTNFα). To this end, the reporter cells were incubated with sIC of different mAb:Ag ratios by fixing one parameter and titrating the other. According to the Heidelberger-Kendall precipitation curve [12] describing sIC size as being dependent on the mAb:Ag ratio, this should result in varying sIC sizes as an excess of either antigen or antibody results in the formation of smaller complexes compared to the large molecular complexes formed at around equal molarity. Changes in sIC size due to a varying mAb:Ag ratio were quantified using asymmetrical flow-field flow fractionation (AF4) (Fig. 5A and Table S1). Fig. S2 shows a complete run of such an analysis. AF4 analysis identifies the highest sIC mean molecular being approximately 2130 kDa at a 1:3 ratio (mAb Ifx : Ag TNF-α) with sICs getting smaller with increasing excess of either antigen or antibody, recapitulating a Heidelberger-Kendall-like curve. Incubation of the FcγR reporter cells with ICs of varying size indeed shows that the assay is sensitive to exactly monitor changes in sIC size (Fig. 5B). Accordingly, both FcγR types showed the strongest responses at mAb:Ag ratios of approximately 1:3. We then set out to test the accuracy of our reporter cell assay as a surrogate for primary human immune cells expressing FcγRs. To this end, we tested primary NK cells from three individual donors and measured NK cell MIP1-β upregulation in response to synthetic sICs of varying size and composition again using a similar assay setup optimized for NK cell activation. We chose MIP-1β upregulation as a cell surface marker to measure NK cell activation as it showed the highest responsiveness in previous experiments (Fig. 4). We could observe that primary immune cells expressing FcγRIIIA respond to IC size, confirming our assay to be an accurate surrogate for primary immune cell responses to soluble ICs (Fig. 5C). Convincingly, NK cell responses to sICs generated from trimeric antigen (rhTNFα) peaked at a different mAb:Ag ratio compared to NK cell responses to sICs generated from dimeric antigens (rhIL-5 and rhVEGFA). Of note,

TNFα and VEGFA activate resting NK cells thus leading to higher MIP1-β positivity when NK

cells are incubated in the presence of excess antigen. As NK cells do not express IL-5 receptor,

this effect is not observed in the presence of excess IL-5. Regarding TNFα, NK cells still show

a stronger activation by sICs generated under optimal mAb:Ag ratios compared to conditions

where excess antigen is used. The data reveal a direct correlation between sIC size and

effector responses. Conversely, when changing antibody concentrations using fixed amounts

of antigen, a consistent reduction of NK cell activation is observed in the presence of excess

IgG for all three mAb/Ag pairs.

Fig. 5. FcγRIIB/C and FcγR-IIIA/B respond to sIC size reproducing a Heidelberger-Kendall like precipitation curve. A) infliximab (mAb) and rhTNFα (Ag) were mixed at different ratios (17 µg total protein, calculated from monomer molarity) and analysed via AF4. sIC size is maximal at a 1:3 ratio of mAb:Ag and reduced when either mAb or Ag are given in excess. $<M>_w$ = mass-weighted mean of the molar mass distribution. Three independent experiments. Error bars = SD. Data taken from Table S1. One complete run analysis is shown in Fig. S2. B) FcγR BW5147 reporter cell activation is sensitive to sIC size. sICs of different size were generated by cross-titration according to the AF4 determination. Reporter cells were incubated with fixed amounts of either mAb (infliximab, left) or Ag (rhTNFα, right) and titrated amounts of antigen or antibody, respectively. X-Axis shows titration of either antigen or antibody, respectively (TNFα calculated as monomer). IL-2 production of reporter cells shows a peak for FcγRIIB/C and FcγRIIIA/B activation at an antibody:antigen ratio between 1:2 and 1:4. FcγRs I and IIA show no activation towards sICs in line with previous observations, see Fig.3. Two independent experiments. Error bars = SD. C) Primary NK cells purified by negative selection magnetic bead separation from three different donors were incubated with cross-titrated sICs as in A. NK cells were measured for MIP-1β expression (% positivity). Incubation with PMA and Ionomycin served as a positive control. Incubation with medium alone served as a negative control. Measured in technical replicates. Error bars = SD.

Quantification of sIC bioactivity in sera of SLE patients.

In order to apply the assay to a clinically relevant condition associated with the occurrence of

sICs we measured circulating sICs present in the serum of SLE patients with variable disease

activity. Sera from 4 healthy donors and 25 SLE patients were investigated for FcγRIIIA/B and

FcγRIIB/C activation. Reporter cells readily produced IL-2 in response to patient sera in a dose-

dependent manner (shown exemplarily for some SLE patients, see Fig. 6A), which was not the

case when sera from healthy controls were tested. Consistent with the observation that FcγRI

and FcγRIIA do not respond to synthetic sICs, reporter cells expressing these receptors did

also not respond to the tested serum samples (Fig. 6A, lower panel). While this reaction pattern

already indicated that sICs are the reactive component measured in SLE patients' sera, we

further demonstrate that FcγRIIIA/B and FcγRIIB/C activation depends on the presence of

serum ICs by analyzing patient serum before and after polyethylene glycol (PEG) precipitation

and depletion of sICs (Fig. 6B). Next, we calculated the area under the curve (AUC) values for

all 25 SLE patient titrations and normalized them to the AUC values measured for healthy

individuals. The resulting index values were then correlated with established biomarkers of

SLE disease activity, such as anti-dsDNA titers (α-dsDNA) and concentrations of the

complement cleavage product C3d (Fig. 6C). We observed a significant correlation between

our FcγRIIIA/B activation index values and both of the determined disease activity markers,

anti-dsDNA titers and C3d concentration (p=0.0465 and p=0.0052, respectively). FcγRIIB/C

on the other hand showed no significant correlation with either biomarker. We assume these

interrelations may be due to the influence of IgG sialylation found to be reduced in active SLE

[52]. Generally, de-sialylation of IgG leads to stronger binding by the activating receptors FcγRI,

FcγRIIA and FcγRIII while it reduces the binding affinity of the inhibitory FcγRIIB [53]. In sum,

our assay allows the indirect quantification of clinically relevant sICs in sera of SLE patients.

Fig. 6. The reporter assay enables quantification of serum-derived sICs from SLE patients. Serum derived sIC from systemic lupus erythematosus (SLE) patients activate human FcγR reporter cells. 25 patients and 4 healthy control individuals were separated into three groups for measurement. A) Experiments shown for an exemplary group of 8 SLE patients and two healthy individuals. Dose-dependent reactivity of FcγRs IIIA/B and IIB/C was observed only for SLE patient sera and not for sera from healthy individuals. FcγRs I and IIA show no reactivity towards clinical IC in line with previous observations. B) Activation of FcγRs IIB/C and IIIA/B by patient serum is mediated by serum derived sICs. Patient serum samples were depleted of sICs by PEG precipitation and the supernatant (SN) was compared to untreated serum regarding FcγR activation (left). Performed in technical replicates. IgG concentration in the precipitate (PC), supernatant (SN) or unfractionated serum respectively is shown in the bar graph (right). IC precipitation did leave non-complexed IgG in the supernatant. C) FcγRIIIA/B activation, but not FcγRIIB/C activation, significantly correlates with known SLE disease markers. FcγR activation data from A was correlated to established SLE disease markers (α-dsDNA levels indicated as IU/ml or C3d concentrations indicated as mg/L). FcγR activation from a dose-response curve as in A was calculated as area under curve (AUC) for each SLE patient (n=25) or healthy individual (n=4) and expressed as fold change compared to the healthy control mean. SLE patients with α-dsDNA levels below 50 IU/ml and C3d values below 6 mg/L were excluded. One-tailed Spearman's.

Assay application to clinically relevant *in vivo* lupus and arthritis mouse models.

BW5147 reporter cells stably expressing chimeric mouse as well as rhesus macaque FcγRs have already been generated using the here described methodology and were successfully used to measure FcγR activation by opsonized adherent cells in previous studies [46, 54] (mFcγR reporter cells). As the human FcγR reporter cells described here are sensitive to certain sICs, we next aimed to translate the assay to clinically relevant animal models. To this end, we incubated previously described FcγR reporter cells expressing chimeric mouse FcγRs [46] with sera from lupus (NZB/WF1) or arthritis (K/BxN) mice with active autoimmune disease. The reporter assay was performed as described above. We chose to measure the stimulation of the activating receptors, mFcγRIII and mFcγRIV, that correspond to human FcγRIII and show a similar cellular distribution and immune function [16]. Incubation with synthetic sICs generated from rhTNFα and mouse-anti-hTNFα IgG1 showed both of the mFcγR reporter cells to be equally responsive to sICs (Fig. 7A). Parental BW5147 cells expressing no FcγRs served as a control. The sera of three mice per group were analysed and compared to sera from wildtype C57BL/6 mice, which served as a healthy control. C57BL/6 mice were chosen, as healthy K/BxN or NZB/WF1 mice cannot be reliably defined. This is due to the unpredictable disease

progression in these mouse models starting from a young age. We consistently detected

mFcγR activation by sera from K/BxN or NZB/WF1 compared to C57BL6 mice (Fig. 7B). While

the mFcγRIII responses were generally high and similar between K/BxN and NZB/WF1 mice,

mFcγRIV responsiveness attended to be lower and individually more variable. Altogether, the

assay enables the reliable detection of sICs in sera of mice with immune-complex mediated

diseases making it a promising novel tool to monitor sICs as a biomarker of disease activity.

Fig. 7. The reporter assay can be applied to mouse models of autoimmune disease. A) Reporter cells expressing mFcγRIII, mFcγRIV or parental BW5147 cells were incubated with titrated amounts of synthetic sICs generated from rhTNFα and mouse-anti-hTNFα at a 1:1 ratio by mass. One experiment in technical replicates. Error bars = SD. B) Titrations of 3 mouse sera per group (C57BL/6, K/BxN or NZB/WF1) were incubated with mFcγR reporter cells and FcγR activation was assessed as described above. Sera from BL/6 mice served as negative control. Performed in technical replicates. Error bars = SD.

Discussion

In this study we established, validated and applied a new assay system that is able for the first time to selectively detect soluble multimeric immune complexes as discrete ligands of FcγR. Major findings are i) that sICs cross-link and trigger human FcγRs IIB/C and IIIA/B but are neglected by activating FcγRI and IIA; ii) that sICs potency is strictly stoichiometry- and size-dependent and thus reproducing the classical Heidelberger-Kendall curve iii) that the overall functional design of FcγRs is adapted to physical cross-linking by soluble vs. membrane-bound immune complexes. The new test system has various obvious applications in medicine and pharmacy.

A novel assay for the quantification of disease-associated as well as synthetic sICs.

The new approach presents an important and implementable advancement in immunological methodology as it enables the sensitive detection of receptor-activating ICs by a relatively simple, scalable *in vitro* bioassay with high-throughput potential. Based on our pilot study demonstrating that the sIC-mediated FcγRIII activation correlated with SLE disease markers this is of great value for larger prospective clinical studies in patients with autoimmune diseases such as systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA), where circulating ICs have long been shown to crucially contribute to tissue damage and disease manifestations [39, 40, 55-57]. Disease-associated, endogenous sICs can also occur during infection, e.g. generated after antibody-mediated destruction of pathogenic viruses or microbes or probably more likely due to the oligomeric nature of numerous viral and bacterial structural proteins generated e.g. by SARS-CoV2 [58-60] HIV and hepatitis B virus (HBV) infection, during which circulating sICs are generated [36, 61].

As the assay also enables the sensitive and quantitative measurement of FcγR ligand bioactivity it allows the detection of sICs not only in clinical specimens but also pharmaceutical preparations of IgG and Fc-fusion proteins for therapeutic use [45]. The presence of ICs in therapeutic preparations or sIC formation following patient treatment is unwanted due to the

risk of side effects such as lupus-like syndrome which has been linked to mAb treatment in

patients receiving infliximab [62] or bevacizumab [63], both mAbs used in this study. As this assay

is highly sensitive to any aggregation of IgG, it also represents a tool to control the purity,

quality and stability of mAb preparations produced for therapeutic use in patients. In addition,

the assessment of sIC-mediated FcγR activation allows for optimization of mAbs and Fc-fusion

proteins regarding their molecular design and manufacturing. Specifically, Fc-molecules

targeting cytokines and soluble factors, which result in sIC formation, could be designed for

reduced or enhanced FcγR activation such as glyco-engineered mAbs or LALA-mutant mAbs

[64, 65]. Notably, the scalability of this cell-based test system does allow for large-scale screening

of samples.

sICs form FcγR ligands with distinct functional profiles.

While immobilized ICs decorating infected cells, viral particles or microbial surfaces restrict

their triggering effect to a single immune cell residing in very close vicinity, sICs can rapidly

disseminate reaching a high number of immune effector cells thus developing a systemic effect

with far reaching and long-lasting or even threatening consequences for the host as a whole.

This could explain the very tight control of sIC governed effector programs and the

unresponsiveness of activating receptors like FcγRI and IIA. We also observed a difference in

the response patterns for FcγRIIIA/B and FcγRII/B/C depending on the solubility of clustered

IgG (immobilized versus soluble ICs) which we validated for FcγRIIIA using primary human NK

cells (Fig. 4 and 5C). Importantly,only multimeric but not dimeric sICs can trigger FcγR

activation. This highlights the fundamental structural influence of the particular antigen as well

as the IgG molecule on sIC dimension and subsequent FcγR-mediated signal processing. The

ability of the here described assay to define and quantify the activation of individual FcγRs by

sIC ligands is not achievable using primary immune cells due to redundant immune responses

upon FcγR activation and complex, overlapping expression pattern of FcγRs. Finally, and in

contrast to primary human cells, murine BW5147 reporter cells are largely inert to human

cytokines, which provides a key advantage to measure FcγR activation selectively in human samples.

Notably, the two basically sIC-responsive human FcγR types, i.e. FcγRIIIA/B and FcγRIIB, are either highly activating versus strongly inhibitory and thus show a mutually exclusive expression pattern (ref. review, z.B. Ravetch und Falk). FcγRIIIA mediates ADCC elicited by NK cells and the induction of a pro-inflammatory cytokine profile by CD16$^+$ monocytes, while FcγRIIIB is a GPI-anchored receptor on neutrophils. FcγRIIB is an inhibitory receptor expressed by B cells and dendritic cells (DCs) regulating B cell activation, antibody production by plasma cells and the activation state of DCs, while the activating FcγRIIC is found on NK cells mediating ADCC. However, as FcγRIIC is only expressed by roughly 10% of the human population, its role is still poorly understood [48-51]. Given the here shown fundamental difference in FcγR reactivity towards multimeric sICs versus immobilized IgG it is tempting to speculate that FcγRIIIA/B- and FcγRIIB/C-positive immune cells might have evolved to differentially perceive these different FcγR ligands (sICs versus membrane bound insoluble ICs) and translate them into distinct reaction patterns. This could be achieved by differences in receptor density, signal transduction or regulation of receptor expression. Consulting the literature indeed supports our hypothesis with neutrophils, B cells and NK cells being efficiently activated by sICs via essentially identical receptor ectodomains [18, 28, 66, 67], while the immunological outcome of their triggering very much differs.

Revisiting the Heidelberger-Kendall curve: dynamic sIC size measurement and monitoring of bioactivity in sIC-associated diseases.

We provide for the first time a simultaneous functional and biophysical assessment of the paradigmatic Heidelberger-Kendall precipitation curve [12, 13]. While previous work already revealed that large and small sICs differentially impacts IL-6 production in PBMCs [6], the dynamics of FcγR activation resulting from constant changes in sIC size have not been explored on a systematic basis. This was undertaken in this study by directly analyzing synthetic ICsformed by highly pure recombinant components via AF4 (Fig. 5A, Fig. S2, Table

S1). Our data document that sIC size is indeed governed by antibody:antigen ratios covering

a wide range of sizes up to several megadaltons. In the presence of increasing amounts of

antibody or antigen deviating from an optimal antibody:ratio, sIC size steadily decreases.

Further, by the measurement of FcγR activation we now translate sIC size directly to asimple

but precise read-out. In doing so, we show that sIC size essentially tunes FcγR activation on

and off. Thus, our new test system can not only contribute to the functional detection and

quantification of clinically relevant sICs but also provides a starting point on how to avoid

pathological consequences by influencing the sIC size, for example by administering

therapeutic antibodies or recombinant antigens in optimized concentrations, thus becoming

relevant in clinical pharmacokinetics.

Limitations of the reporter system and conclusions

While it is known that FcRIIA requires CD16B or FcRn to efficiently respond to ICs [22, 23, 68],

unexpectedly, we found that FcγRI is not activated by sICs in our assay. We assume that FcγRI

interaction with sICs might require a native cellular environment given that a major function of

this FcγR is the uptake and processing of antigen via ICs even in the absence of a signaling

motif [21]. However, we find that FcγRI ectodomains alone are not responsive to sICs implying

a thoroughly different cross-linking threshold for FcγRI compared with FcγRIIB and IIIA/B. This

feature is possibly linked to its molecular architecture being the only high-affinity FcγR with

three extracellular Ig domains compared to the two domains found in other FcγRs with lower

affinity to monomeric IgG. This observation also reflects a general consideration regarding the

BW5147 reporter system. While providing a robust and uniform read-out using a scalable cell-

based approach, the assay is not able to reflect native immune cell functions governed by cell-

intrinsic signalling cascades. The major advancements of this reporter system include i) a

higher accuracy regarding FcγR activation compared to merely affinity measurements, ii) an

sIC size dependent quantification of FcγR responsiveness and iii) the identification of FcγR

activating sICs in autoimmunity and infection. Finally, this scalable, sensitive and robust

system to detect FcγR activating sICs in clinical samples might enable their identification in

diseases that have not been linked to sIC-mediated pathology, yet.

Materials and Methods

Cell culture

All cells were cultured in a 5% CO_2 atmosphere at 37°C. BW5147 mouse thymoma cells (BW,

obtained from ATCC: TIB-47) were maintained at 3×10^5 to 9×10^5 cells/ml in Roswell Park

Memorial Institute medium (RPMI GlutaMAX, Gibco) supplemented with 10% (vol/vol) fetal calf

serum (FCS, Biochrom), sodium pyruvate (1x, Gibco) and β-mercaptoethanol (0.1 mM, Gibco).

293T-CD20 (kindly provided by Irvin Chen, UCLA [69]) were maintained in Dulbecco's modified

Eagle's medium (DMEM, Gibco) supplemented with 10% (vol/vol) FCS.

FcγR receptor activation assay

FcγR activation was measured adapting a previously described cell-based assay [44, 70]. The

assay was modified to measure FcγR activation in solution. Briefly, 2×10^5 mouse BW-FcγR

(BW5147) reporter cells were incubated with synthetic sICs or diluted serum in a total volume

of 100 µl for 16 h at 37°C and 5% CO_2. Incubation was performed in a 96-well ELISA plate

(Nunc maxisorp) pre-treated with PBS/10% FCS (v/v) for 1 h at 4°C. Immobilized IgG was

incubated in PBS on the plates prior to PBS/10% FCS treatment. Reporter cell mIL-2 secretion

was quantified via ELISA as described [44].

Recombinant antigens and monoclonal antibodies to form sICs

Recombinant human (rh) cytokines TNF, IL-5, and VEGFA were obtained from Stem Cell

technologies. Recombinant CD20 was obtained as a peptide (aa141-188) containing the

binding region of rituximab (Creative Biolabs). FcγR-specific mAbs were obtained from Stem

Cell technologies (CD16: clone 3G8; CD32: IV.3). Reverse sICs were generated from these

receptor-specific antibodies using goat-anti-mouse IgG F(ab)$_2$ fragments (Invitrogen) in a 1:1

ratio. Pharmaceutically produced humanized monoclonal IgG1 antibodies infliximab (Ifx),

bevacizumab (Bvz), mepolizumab (Mpz) and rituximab (Rtx) were obtained from the University

Hospital Pharmacy Freiburg. Mouse anti-hTNFα (IgG2b, R&D Systems, 983003) was used to

generate sICs reactive with mouse FcγRs. sICs were generated by incubation of antigens and

antibodies in reporter cell medium or PBS for 2 h at 37°C.

Lentiviral transduction

Lentiviral transduction of BW5147 cells was performed as described previously [46, 54]. In brief,

chimeric FcγR-CD3ζ constructs [44] were cloned into a pUC2CL6IPwo plasmid backbone. For

every construct, one 10-cm dish of packaging cell line at roughly 70% density was transfected

with the target construct and two supplementing vectors providing the VSV gag/pol and VSV-

G-env proteins (6 μg of DNA each) using polyethylenimine (22.5 μg/ml, Sigma) and Polybrene

(4 μg/ml; Merck Millipore) in a total volume of 7 ml (2 ml of a 15-min-preincubated transfection

mix in serum-free DMEM added to 5 ml of fresh full DMEM). After a medium change, virus

supernatant harvested from the packaging cell line 2 days after transfection was then

incubated with target BW cells overnight (3.5 ml of supernatant on 10^6 target cells), followed

by expansion and pool selection using complete medium supplemented with 2 μg/ml of

puromycin (Sigma) over a one week culture period.

BW5147 toxicity test

Cell counting was performed using a Countess II (Life Technologies) according to supplier

instructions. Cell toxicity was measured as a ratio between live and dead cells judged by trypan

blue staining over a 16 h time frame in a 96well format (100 μl volume per well). BW5147 cells

were mixed 1:1 with Trypan blue (Invitrogen) and analysed using a Countess II. rhTNFα was

diluted in complete medium.

human IgG suspension ELISA

1 µg of IgG1 (rituximab in PBS, 50 µl/well) per well was incubated on a 96well microtiter plate

(NUNC Maxisorp) pre-treated (2h at RT) with PBS supplemented with varying percentages

(v/v) of FCS (PAN Biotech). IgG1 bound to the plates was detected using an HRP-conjugated

mouse-anti-human IgG mAb (Jackson ImmunoResearch).

BW5147 cell flow cytometry

BW5147 cells were harvested by centrifugation at 900 g and RT from the suspension culture.

1×10^6 cells were stained with PE- or FITC-conjugated anti-human FcγR mAbs (BD) or a PE-

TexasRed-conjugated human IgG-Fc fragment (Rockland) for 1h at 4°C in PBS/3%FCS. After

3 washing steps in PBS/3%FCS, the cells were transferred to Flow cytometry tubes (BD) and

analysed using BD LSR Fortessa and FlowJo (V10) software.

NK cell activation flow cytometry

PBMC were purified from donor blood using Lymphocyte separation Media (Anprotec). Primary

NK cells were separated from donor PBMCs via magnetic bead negative selection (Stem Cell

technologies) and NK cell purity was confirmed via staining of CD3 (Biolegend, clone HIT3a),

CD16 (Biolegend, clone 3G8) and CD56 (Miltenyi Biotec, clone AF12-7H3). 96well ELISA

plates (Nunc Maxisorp) were pre-treated with PBS/10% FCS (v/v) for 1 h at 4°C. NK cells were

stimulated in pre-treated plates and incubated at 37°C and 5% CO_2 for 4 h. Golgi Plug and

Golgi Stop solutions (BD) were added as suggested by supplier. CD107a (APC, BD, H4A3)

specific conjugated mAb was added at the beginning of the incubation period. Following the

stimulation period, MIP-1β (PE, BD Pharmigen), IFNγ (BV-510, Biolegends, 4SB3) and TNFα

(PE/Cy7, Biolegends, MAB11) production was measured via intracellular staining Cytokines

(BD, CytoFix/CytoPerm, Kit as suggested by the supplier). 50 ng/ml PMA (InvivoGen) + 0.5

µM Ionomycin (InvivoGen) were used as a positive stimulation control for NK cell activation.

After 3 washing steps in PBS/3%FCS, the cells were transferred to Flow cytometry tubes (BD)

and analysed using a BD FACS Fortessa and FlowJo (V10) software. FcγRII or FcγRIII block

was performed by addition of receptor specific mAbs (Stem cell technologies, IV.3 and 3G8)

at a 1:100 dilution at the beginning of the incubation period. Cells were transferred to Flow

cytometry tubes (BD) and analyzed using BD LSR Fortessa and FlowJo (V10) software.

Asymmetric flow field flow fractionation (AF4)

The AF4 system consisted of a flow controller (Eclipse AF4, Wyatt), a MALS detector (DAWN

Heleos II, Wyatt), a UV detector (1260 Infinity G1314F, Agilent) and the separation channel

(SC channel, PES membrane, cut-off 10 kDa, 490 µm spacer, wide type, Wyatt). Elution buffer:

1.15 g/L Na_2HPO_4 (Merck), 0.20 g/L NaH_2PO_4 x H_2O (Merck), 8.00 g/L NaCl (Sigma) and 0,20

g/L NaN_3 (Sigma), adjusted to pH 7.4, filtered through 0.1 µm. AF4 sequence (Vx = cross flow

in mL/min): (a) elution (2 min, Vx: 1.0); (b) focus (1 min, Vx: 1.0), focus + inject (1 min, Vx: 1.0,

inject flow: 0.2 mL/min), repeated three times; (c) elution (30 min, linear Vx gradient: 1.0 to

0.0); (d) elution (15 min, Vx: 0.0); (e) elution + inject (5 min, Vx: 0.0). A total protein mass of

17±0.3 µg (Ifx, rhTNFα or ICs, respectively) was injected. The eluted sample concentration

was calculated from the UV signal at 280 nm using extinction coefficients of 1.240 mL/(mg cm)

or 1.450 mL/(mg cm) in the case of TNFα or Ifx, respectively. For the ICs, extinction coefficients

were not available and difficult to calculate as the exact stoichiometry is not known. An

extinction coefficient of 1.450 mL/(mg cm) was used for calculating the molar masses of all

ICs. Especially in the case of ICs rich in TNFα, the true coefficients should be lower, and the

molar masses of these complexes are overestimated by not more than 14 %. The determined

molar masses for TNFα-rich complexes are therefore biased but the observed variations in

molar mass for the different ICs remain valid. The mass-weighted mean of the distribution of

molar masses for each sample was calculated using the ASTRA 7 software package (Wyatt).

SLE patient cohort

Sera from patients with SLE were obtained from the Immunologic, Rheumatologic Biobank (IR-

B) of the Department of Rheumatology and Clinical Immunology. Biobanking and the project

were approved by the local ethical committee of the University of Freiburg (votes 507/16 and

624/14). All patients who provided blood to the biobank had provided written informed consent.

Ethical Statement: The study was designed in accordance with the guidelines of the

Declaration of Helsinki (revised 2013). Patients with SLE (n = 25) and healthy controls (n = 4)

were examined. All patients met the revised ACR classification criteria for SLE. Disease activity

was assessed using the SLEDAI-2K score. C3d levels were analyzed in EDTA plasma using

rocket double decker immune-electrophoresis with antisera against C3d (Polyclonal Rabbit

Anti-Human C3d Complement, Agilent) and C3c (Polyclonal Rabbit Anti-Human C3c

Complement Agilent) as previously described [71]. Anti-human dsDNA antibodies titers were

determined in serum using an anti-dsDNA IgG ELISA kit (diagnostik-a GmbH).

Patient serum IC precipitation

For polyethylene glycol (PEG) precipitation human sera were mixed with PEG 6000 (Sigma-

Aldrich) in PBS at a final concentration of 10% PEG 6000. After overnight incubation at 4°C,

ICs were precipitated by centrifugation at 2000 x g for 30 min at 4 °C, pellets were washed

once with PEG 6000 and then centrifugated at 2000 x g for 20 min at 4 °C. Supernatants were

harvested and precipitates re-suspended in pre-warmed PBS for 1 h at 37 °C. IgG

concentrations of serum, precipitates and supernatants obtained after precipitation were

quantified by Nanodrop (Thermo Scientific™) measurement.

Mice and Models

Animal experiments were approved by the local governmental commission for animal

protection of Freiburg (Regierungspräsidium Freiburg, approval no. G16/59 and G19/21).

Lupus-prone (NZBxNZW)F1 mice (NZB/WF1) were generated by crossing NZB⁄BINJ mice with

NZW/LacJ mice, purchased from The Jackson Laboratory. KRNtg mice were obtained from F.

Nimmerjahn (Universität Erlangen-Nürnberg) with the permission of D. Mathis and C. Benoist

(Harvard Medical School, Boston, MA), C57BL/6 mice (BL/6) and NOD/ShiLtJArc (NOD/Lt)

mice were obtained from the Charles River Laboratories. K/BxN (KRNtgxNOD)F1 mice

(K/BxN) were obtained by crossing KRNtg mice and NOD/Lt mice. All mice were housed in a

12-h light/dark cycle, with food and water ad libitum. Mice were euthanized and blood collected

for serum preparation from 16 weeks old BL/6 animals, from 16 weeks old arthritic K/BxN

animals and from 26 – 38 weeks old NZB/WF1 mice with established glomerulonephritis.

Statistical analyses

Statistical analyses were performed using Graphpad Prism software (v6) and appropriate

tests.

Funding

This work was supported by an intramural junior investigator fund of the Faculty of Medicine

to PK (EQUIP - Funding for Medical Scientists, Faculty of Medicine, University of Freiburg), by

the German Research foundation (DFG) (FOR2830 HE 2526/9-1) to HH, by the DFG research

grant TRR130 to REV and the Ministry of Science, Research, and Arts Baden-Württemberg

(Margarete von Wrangell Programm) to NC.

Acknowledgement

We thank T. Schleyer (IR-B Biobank) for providing patient samples. We are indebted to Falk.

Nimmerjahn (Universität Erlangen-Nürnberg) for providing KRNtg mice.

Competing interests
We declare no financial and non-financial competing interests.
644

References

1.   Lu, L.L., Suscovich, T.J., Fortune, S.M. & Alter, G. Beyond binding: antibody effector functions in infectious diseases. *Nat Rev Immunol* **18**, 46-61 (2018).

2.   Ravetch, J.V. & Bolland, S. IgG Fc receptors. *Annu Rev Immunol* **19**, 275-290 (2001).

3.   van der Poel, C.E., Spaapen, R.M., van de Winkel, J.G. & Leusen, J.H. Functional characteristics of the high affinity IgG receptor, FcgammaRI. *J Immunol* **186**, 2699-2704 (2011).

4.   Bruhns, P. et al. Specificity and affinity of human Fcgamma receptors and their polymorphic variants for human IgG subclasses. *Blood* **113**, 3716-3725 (2009).

5.   Urlaub, D. et al. Activation of natural killer cells by rituximab in granulomatosis with polyangiitis. *Arthritis Res Ther* **21**, 277 (2019).

6.   Lux, A., Yu, X., Scanlan, C.N. & Nimmerjahn, F. Impact of immune complex size and glycosylation on IgG binding to human FcgammaRs. *J Immunol* **190**, 4315-4323 (2013).

7.   Kiefer, F. et al. The Syk protein tyrosine kinase is essential for Fc gamma receptor signaling in macrophages and neutrophils. *Mol Cell Biol* **18**, 4209-4220 (1998).

8.   Luo, Y., Pollard, J.W. & Casadevall, A. Fcgamma receptor cross-linking stimulates cell proliferation of macrophages via the ERK pathway. *J Biol Chem* **285**, 4232-4242 (2010).

9.   Greenberg, S., Chang, P. & Silverstein, S.C. Tyrosine phosphorylation of the gamma subunit of Fc gamma receptors, p72syk, and paxillin during Fc receptor-mediated phagocytosis in macrophages. *J Biol Chem* **269**, 3897-3902 (1994).

10.   Bournazos, S., Wang, T.T., Dahan, R., Maamary, J. & Ravetch, J.V. Signaling by Antibodies: Recent Progress. *Annu Rev Immunol* **35**, 285-311 (2017).

11.   Nimmerjahn, F. & Ravetch, J.V. Antibody-mediated modulation of immune responses. *Immunol Rev* **236**, 265-275 (2010).

12.   Heidelberger, M. & Kendall, F.E. A Quantitative Study of the Precipitin Reaction between Type Iii Pneumococcus Polysaccharide and Purified Homologous Antibody. *J Exp Med* **50**, 809-823 (1929).

13.   Heidelberger, M. & Kendall, F.E. The Precipitin Reaction between Type Iii Pneumococcus Polysaccharide and Homologous Antibody : Iii. A Quantitative Study and a Theory of the Reaction Mechanism. *J Exp Med* **61**, 563-591 (1935).

14.   Duchemin, A.M., Ernst, L.K. & Anderson, C.L. Clustering of the High-Affinity Fc Receptor for Immunoglobulin-G (Fc-Gamma-Ri) Results in Phosphorylation of Its Associated Gamma-Chain. *J Biol Chem* **269**, 12111-12117 (1994).

15.   Getahun, A. & Cambier, J.C. Of ITIMs, ITAMs, and ITAMis: revisiting immunoglobulin Fc receptor signaling. *Immunol Rev* **268**, 66-73 (2015).

16.   Bruhns, P. & Jonsson, F. Mouse and human FcR effector functions. *Immunol Rev* **268**, 25-51 (2015).

17.   Bruhns, P. Properties of mouse and human IgG receptors and their contribution to disease models. *Blood* **119**, 5640-5649 (2012).

18.   Nimmerjahn, F. & Ravetch, J.V. Fcgamma receptors as regulators of immune responses. *Nat Rev Immunol* **8**, 34-47 (2008).

19.   Nimmerjahn, F. & Ravetch, J.V. Fcgamma receptors: old friends and new family members. *Immunity* **24**, 19-28 (2006).

20.   Guilliams, M., Bruhns, P., Saeys, Y., Hammad, H. & Lambrecht, B.N. The function of Fcgamma receptors in dendritic cells and macrophages. *Nat Rev Immunol* **14**, 94-108 (2014).

21.   Indik, Z.K. et al. The high affinity Fc gamma receptor (CD64) induces phagocytosis in the absence of its cytoplasmic domain: the gamma subunit of Fc gamma RIIIA imparts phagocytic function to Fc gamma RI. *Exp Hematol* **22**, 599-606 (1994).

22.   Fossati, G., Bucknall, R.C. & Edwards, S.W. Insoluble and soluble immune complexes activate neutrophils by distinct activation mechanisms: changes in functional responses induced by priming with cytokines. *Ann Rheum Dis* **61**, 13-19 (2002).

23. Hubbard, J.J. et al. FcRn is a CD32a coreceptor that determines susceptibility to IgG
immune complex-driven autoimmunity. *J Exp Med* **217** (2020).
24. Pincetic, A. et al. Type I and type II Fc receptors regulate innate and adaptive immunity.
Nat Immunol **15**, 707-716 (2014).
25. Vidarsson, G., Dekkers, G. & Rispens, T. IgG subclasses and allotypes: from structure
to effector functions. *Front Immunol* **5**, 520 (2014).
26. Tay, M.Z., Wiehe, K. & Pollara, J. Antibody-Dependent Cellular Phagocytosis in
Antiviral Immune Responses. *Front Immunol* **10**, 332 (2019).
27. Laborde, E.A. et al. Immune complexes inhibit differentiation, maturation, and function
of human monocyte-derived dendritic cells. *J Immunol* **179**, 673-681 (2007).
28. Kang, S. et al. IgG-Immune Complexes Promote B Cell Memory by Inducing BAFF. *J
Immunol* **196**, 196-206 (2016).
29. Granger, V., Peyneau, M., Chollet-Martin, S. & de Chaisemartin, L. Neutrophil
Extracellular Traps in Autoimmunity and Allergy: Immune Complexes at Work. *Front
Immunol* **10**, 2824 (2019).
30. Berger, S., Ballo, H. & Stutte, H.J. Immune complex-induced interleukin-6, interleukin-
10 and prostaglandin secretion by human monocytes: A network of pro- and anti-
inflammatory cytokines dependent on the antigen:antibody ratio. *Eur J Immunol* **26**,
1297-1301 (1996).
31. Koenderman, L. Inside-Out Control of Fc-Receptors. *Front Immunol* **10**, 544 (2019).
32. Plomp, R. et al. Subclass-specific IgG glycosylation is associated with markers of
inflammation and metabolic health. *Sci Rep* **7**, 12325 (2017).
33. Pierson, T.C. et al. The stoichiometry of antibody-mediated neutralization and
enhancement of West Nile virus infection. *Cell Host Microbe* **1**, 135-145 (2007).
34. Patel, K.R., Roberts, J.T. & Barb, A.W. Multiple Variables at the Leukocyte Cell Surface
Impact Fc gamma Receptor-Dependent Mechanisms. *Front Immunol* **10**, 223 (2019).
35. Bohm, S., Kao, D. & Nimmerjahn, F. Sweet and sour: the role of glycosylation for the
anti-inflammatory activity of immunoglobulin G. *Current topics in microbiology and
immunology* **382**, 393-417 (2014).
36. Wang, T.T. & Ravetch, J.V. Immune complexes: not just an innocent bystander in
chronic viral infection. *Immunity* **42**, 213-215 (2015).
37. Yamada, D.H. et al. Suppression of Fcgamma-receptor-mediated antibody effector
function during persistent viral infection. *Immunity* **42**, 379-390 (2015).
38. Antes, U., Heinz, H.P., Schultz, D., Brackertz, D. & Loos, M. C1q-bearing immune
complexes detected by a monoclonal antibody to human C1q in rheumatoid arthritis
sera and synovial fluids. *Rheumatol Int* **10**, 245-250 (1991).
39. Zubler, R.H. et al. Circulating and intra-articular immune complexes in patients with
rheumatoid arthritis. Correlation of 125I-Clq binding activity with clinical and biological
features of the disease. *J Clin Invest* **57**, 1308-1319 (1976).
40. Koffler, D., Agnello, V., Thoburn, R. & Kunkel, H.G. Systemic lupus erythematosus:
prototype of immune complex nephritis in man. *J Exp Med* **134**, 169-179 (1971).
41. Rajan, T.V. The Gell-Coombs classification of hypersensitivity reactions: a re-
interpretation. *Trends in immunology* **24**, 376-379 (2003).
42. Tahir, S. et al. A CD153+CD4+ T follicular cell population with cell-senescence features
plays a crucial role in lupus pathogenesis via osteopontin production. *J Immunol* **194**,
5725-5735 (2015).
43. Bano, A. et al. CD28 (null) CD4 T-cell expansions in autoimmune disease suggest a
link with cytomegalovirus infection. *F1000Res* **8** (2019).
44. Corrales-Aguilar, E. et al. A novel assay for detecting virus-specific antibodies
triggering activation of Fcgamma receptors. *Journal of immunological methods* **387**,
21-35 (2013).
45. Lagasse, H.A.D., Hengel, H., Golding, B. & Sauna, Z.E. Fc-Fusion Drugs Have
FcgammaR/C1q Binding and Signaling Properties That May Affect Their
Immunogenicity. *AAPS J* **21**, 62 (2019).

46. Van den Hoecke, S. et al. Hierarchical and Redundant Roles of Activating FcgammaRs
in Protection against Influenza Disease by M2e-Specific IgG1 and IgG2a Antibodies. *J*
Virol **91** (2017).
47. Tanaka, M. et al. Activation of Fc gamma RI on monocytes triggers differentiation into
immature dendritic cells that induce autoreactive T cell responses. *J Immunol* **183**,
2349-2355 (2009).
48. Lisi, S., Sisto, M., Lofrumento, D.D., D'Amore, S. & D'Amore, M. Advances in the
understanding of the Fc gamma receptors-mediated autoantibodies uptake. *Clin Exp*
Med **11**, 1-10 (2011).
49. Anania, J.C., Chenoweth, A.M., Wines, B.D. & Hogarth, P.M. The Human FcgammaRII
(CD32) Family of Leukocyte FcR in Health and Disease. *Front Immunol* **10**, 464 (2019).
50. Metes, D. et al. Expression of functional CD32 molecules on human NK cells is
determined by an allelic polymorphism of the FcgammaRIIC gene. *Blood* **91**, 2369-
2380 (1998).
51. Breunis, W.B. et al. Copy number variation of the activating FCGR2C gene predisposes
to idiopathic thrombocytopenic purpura. *Blood* **111**, 1029-1038 (2008).
52. Vuckovic, F. et al. Association of systemic lupus erythematosus with decreased
immunosuppressive potential of the IgG glycome. *Arthritis Rheumatol* **67**, 2978-2989
(2015).
53. Kaneko, Y., Nimmerjahn, F. & Ravetch, J.V. Anti-inflammatory activity of
immunoglobulin G resulting from Fc sialylation. *Science* **313**, 670-673 (2006).
54. Kolb, P. et al. Identification and Functional Characterization of a Novel Fc Gamma-
Binding Glycoprotein in Rhesus Cytomegalovirus. *J Virol* **93** (2019).
55. Nydegger, U.E. & Davis, J.S.t. Soluble immune complexes in human disease. *CRC Crit*
Rev Clin Lab Sci **12**, 123-170 (1980).
56. Levinsky, R.J., Cameron, J.S. & Soothill, J.F. Serum immune complexes and disease
activity in lupus nephritis. *Lancet* **1**, 564-567 (1977).
57. Levinsky, R.J. Role of circulating soluble immune complexes in disease. *Arch Dis Child*
53, 96-99 (1978).
58. Briant, L., Coudronniere, N., Robert-Hebmann, V., Benkirane, M. & Devaux, C. Binding
of HIV-1 virions or gp120-anti-gp120 immune complexes to HIV-1-infected quiescent
peripheral blood mononuclear cells reveals latent infection. *J Immunol* **156**, 3994-4004
(1996).
59. Oh, S.K. et al. Identification of HIV-1 envelope glycoprotein in the serum of AIDS and
ARC patients. *J Acquir Immune Defic Syndr (1988)* **5**, 251-256 (1992).
60. Vuitton, D.A., Vuitton, L., Seilles, E. & Galanaud, P. A plea for the pathogenic role of
immune complexes in severe Covid-19. *Clin Immunol* **217**, 108493 (2020).
61. Madalinski, K., Burczynska, B., Heermann, K.H., Uy, A. & Gerlich, W.H. Analysis of
viral proteins in circulating immune complexes from chronic carriers of hepatitis B virus.
Clin Exp Immunol **84**, 493-500 (1991).
62. Wetter, D.A. & Davis, M.D. Lupus-like syndrome attributable to anti-tumor necrosis
factor alpha therapy in 14 patients during an 8-year period at Mayo Clinic. *Mayo Clin*
Proc **84**, 979-984 (2009).
63. MacDonald, D.A. et al. Aflibercept exhibits VEGF binding stoichiometry distinct from
bevacizumab and does not support formation of immune-like complexes. *Angiogenesis*
19, 389-406 (2016).
64. Saunders, K.O. Conceptual Approaches to Modulating Antibody Effector Functions and
Circulation Half-Life. *Front Immunol* **10**, 1296 (2019).
65. Li, T. et al. Modulating IgG effector function by Fc glycan engineering. *Proceedings of*
the National Academy of Sciences of the United States of America **114**, 3485-3490
(2017).
66. Goodier, M.R. et al. Sustained Immune Complex-Mediated Reduction in CD16
Expression after Vaccination Regulates NK Cell Function. *Front Immunol* **7**, 384
(2016).
67. Mayadas, T.N., Tsokos, G.C. & Tsuboi, N. Mechanisms of immune complex-mediated
neutrophil recruitment and tissue injury. *Circulation* **120**, 2012-2024 (2009).

68.   Nagarajan, S. et al. Cell-specific, activation-dependent regulation of neutrophil CD32A
811   ligand-binding function. *Blood* **95**, 1069-1077 (2000).
812   69.   Morizono, K. et al. Redirecting lentiviral vectors pseudotyped with Sindbis virus-derived
813   envelope proteins to DC-SIGN by modification of N-linked glycans of envelope
814   proteins. *J Virol* **84**, 6923-6934 (2010).
815   70.   Corrales-Aguilar, E. et al. Human cytomegalovirus Fcgamma binding proteins gp34
816   and gp68 antagonize Fcgamma receptors I, II and III. *PLoS pathogens* **10**, e1004131
817   (2014).
818   71.   Rother, E., Lang, B., Coldewey, R., Hartung, K. & Peter, H.H. Complement split product
819   C3d as an indicator of disease activity in systemic lupus erythematosus. *Clin
820   Rheumatol* **12**, 31-35 (1993).

Supplemental Data

Fig. S1. rhTNFα is not toxic to mouse lymphocyte BW5147 cells even at high concentrations. Cell count and percentage of live cells were unaltered over a 16 h time frame of reporter cell culture in the presence of indicated rhTNFα concentrations and comparable to regular growth in complete medium. Experiments were conducted in 3 replicates. Error bars = SD.

833
Fig. S2. AF4 elution profiles of Ifx/TNFα-immune complexes.
The elution profiles from one of three independent runs are shown. Protein
concentration in the eluate is shown in black (UV signal at λ = 280 nm, normalized to
the highest UV signal found in this experiment), molar masses determined by MALS
for a given retention time in red. Horizontal red lines indicate the range of molar masses
used to calculate the mass-weighted mean of molar masses <Mw>. A) Overlay of the
elution profiles obtained for TNFα and Ifx, respectively; B1 to B5) Elution profiles for
sICs formed after incubation of TNFα and Ifx at different molar ratios.
842

Sample	Range of assigned molar masses [kDa]			Mass-weighted mean of assigned molar masses [kDa]			
	Run 1	Run 2	Run 3	Run 1	Run 2	Run 3	Mean ± SD
Infliximab, IFX	158 – 182	153 – 164	159 – 193	162	156	163	160 ± 4
TNF -alpha	52 – 55	51 – 61	52 – 62	52	52	52	52 ± 0
Immune complexes							
IFX/TNF 3:1	$182 - 1.16 \cdot 10^3$	$164 - 1.11 \cdot 10^3$	$193 - 1.10 \cdot 10^3$	409	398	518	442 ± 66
IFX/TNF 1:1	$182 - 2.06 \cdot 10^3$	$164 - 1.31 \cdot 10^3$	$193 - 1.42 \cdot 10^3$	801	589	681	690 ± 106
IFX/TNF 1:3	$182 - 5.05 \cdot 10^3$	$164 - 6.89 \cdot 10^3$	$193 - 10.8 \cdot 10^3$	$1.77 \cdot 10^3$	$2.00 \cdot 10^3$	$2.61 \cdot 10^3$	$2.13 \cdot 10^3 \pm 435$
IFX/TNF 1:9	$182 - 5.36 \cdot 10^3$	$164 - 3.38 \cdot 10^3$	$193 - 3.51 \cdot 10^3$	$1.66 \cdot 10^3$	$1.18 \cdot 10^3$	$1.17 \cdot 10^3$	$1.34 \cdot 10^3 \pm 279$
IFX/TNF 1:27	$182 - 1.68 \cdot 10^3$	$164 - 768$	$193 - 1.01 \cdot 10^3$	689	464	521	558 ± 117

Table S1. Analysis of the molar mass distribution of ICs from AF4 data.

For a given elution time, the AF4 profiles provide the concentration (UV) at which a given molar
mass (MALS) of a protein is present in the sample. The molar mass distribution of Ifx, TNFα
and their immune complexes (sICs) was obtained by plotting the cumulative frequency as a
function of molar mass. For a selected range of molar masses, a mass-weighted mean value
($<M_w>$) was calculated. All detected molar masses were selected in the case of Ifx and TNFα
whereas only molar masses larger than the maximal molar mass found for Ifx were assigned
to sICs. The table shows the range of assigned molar masses and the calculated $<M_w>$ for
each AF4 run (n = 3).