Detection and functional resolution of soluble multimeric immune complexes by a comprehensive FcγR reporter cell panel

Haizhang Chen[1,2], Andrea Maul-Pavicic[3,4], Martin Holzer[5], Magdalena Huber[1,2], Ulrich Salzer[3], Nina Chevalier[3], Reinhard E. Voll[3,4], Hartmut Hengel[1,2,*] & Philipp Kolb[1,2,*]

[1]Institute of Virology, University Medical Center, Albert-Ludwigs-University Freiburg, Hermann-Herder-Str. 11, 79104 Freiburg, Germany

[2]Faculty of Medicine, Albert-Ludwigs-University Freiburg, 79104 Freiburg, Germany

[3]Department of Rheumatology and Clinical Immunology, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Hugstetterstr. 55, 79106 Freiburg, Germany

[4]Center for Chronic Immunodeficiency (CCI), Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, Breisacherstr. 115, 79106 Freiburg, Germany

[5]Institute for Pharmaceutical Sciences, Albert-Ludwigs-University Freiburg, Hermann-Herder-Str. 9, 79104 Freiburg, Germany

*corresponding authors (Hartmut.hengel@uniklinik-freiburg.de)

One Sentence Summary: In this study we established a comprehensive FcγR reporter cell assay enabling the detection and quantification of soluble immune complexes generated in experimental and clinical settings.

Abstract: Fc-gamma receptor (FcγR) activation by soluble IgG immune complexes (sICs) represents a major mechanism of inflammation in certain autoimmune diseases such as systemic lupus erythematosus (SLE). A robust and scalable test system allowing for the detection and quantification of sIC bioactivity is missing. Previously described FcγR interaction assays are limited to certain FcγRs, lack scalability and flexibility, are not indicative of receptor activation or lack sensitivity towards sIC size. We developed a comprehensive reporter cell panel detecting individual activation of FcγRs from humans and the mouse. The reporter cell lines were integrated into an assay format that provides flexible read-outs enabling the quantification of sIC reactivity via ELISA or a fast detection using flow cytometry. This identified FcγRIIA(H) and FcγRIIIA as the most sIC-sensitive FcγRs in our test system. Applying the assay we demonstrate that sICs versus immobilized ICs are fundamentally different FcγR-ligands with regard to FcγR preference and signal strength. Reaching a detection limit in the very low nanomolar range, the assay proved also to be sensitive to sIC stoichiometry and size enabling for the first time a complete reproduction of the Heidelberger-Kendall precipitation curve in terms of immune receptor activation. Analyzing sera from SLE patients and mouse models of lupus and arthritis proved that sIC-dependent FcγR activation has predictive capabilities regarding severity of SLE disease. The new methodology provides a sensitive, scalable and comprehensive tool to evaluate the size, amount and bioactivity of sICs in all settings.

Introduction

Immunoglobulin G (IgG) is the dominant immunoglobulin isotype in chronic infections and in various antibody-mediated autoimmune diseases. The multi-faceted effects of the IgG molecule rely both on the F(ab) regions, which recognize a specific antigenic determinant to form immune complexes (ICs), and the constant Fc region (Fcγ), which is detected by effector molecules like the Fcγ receptors (FcγRs) found on most cells of the immune system or C1q of

the complement system. When IgG binds to its antigen ICs are formed, which, depending on the respective antigen, are either matrix/cell-bound or soluble (sICs). The composition of sICs is dependent on the number of epitopes recognized by IgG on a single antigen molecule and the ability of the antigen to form multimers. Fcγ-FcγR binding is necessary but not sufficient to activate FcγRs since physical receptor cross-linking is required for receptor triggering (Duchemin et al, 1994; Luo et al, 2010; Patel et al, 2019). IgG opsonized infected cells or microorganisms are readily able to cross-link FcγRs (Bruhns et al, 2009; Lux et al, 2013). This initiates various signalling pathways (Greenberg et al, 1994; Kiefer et al, 1998; Luo et al, 2010) which in turn regulate immune cell effector functions (Bournazos et al, 2017; Nimmerjahn & Ravetch, 2010). It is also suggested that sICs can dynamically tune FcγR triggering, implying that changes in sIC size directly impact strength and duration of FcγR responses (Lux et al, 2013). However, the molecular requirements are largely unknown and a translation to bioactivity of the paradigmatic Heidelberger-Kendall precipitation curve, describing that sIC size depends on the antigen:antibody ratio, is so far missing (Heidelberger & Kendall, 1929; Heidelberger & Kendall, 1935).

Among all type I FcγRs, FcγRIIB (CD32B) is the only inhibitory receptor, signalling via immunoreceptor tyrosine-based inhibitory motifs (ITIMs), while the activating receptors are associated with immunoreceptor tyrosine-based activation motifs (ITAMs). Another exception is FcγRIIIB (CD16B), which is glycosylphosphatidylinositol (GPI)-anchored and lacks a signalling motif (Bruhns, 2012; Bruhns & Jonsson, 2015; Nimmerjahn & Ravetch, 2006; Nimmerjahn & Ravetch, 2008). Still, FcγRIIIB is widely accepted to be a neutrophil activating receptor, e.g. by cooperating with other FcγRs such as FcγRIIA (Vossebeld et al, 1997). FcγRI (CD64) is the only high affinity FcγR binding also to monomeric IgG, while all other FcγRs only efficiently bind to complexed, i.e. antigen-bound IgG (Bruhns, 2012; Bruhns & Jonsson, 2015; Lu et al, 2018). Activation of FcγRs leads to a variety of cellular effector functions elicited by several immune cells such as natural killer (NK) cells via FcγRIIC/FcγRIIIA,

monocyte-derived   cells   via   FcγRI/FcγRIIB/FcγRIIIA,   granulocytes   via

FcγRI/FcγRIIA/FcγRIIIB, platelets via FcγRIIA and B cells via FcγRIIB. Consequently, FcγRs

connect and regulate both the innate and adaptive branches of the immune system. Various

factors have been shown to influence IC-dependent FcγR activation profiles, including FcγR-

Fcγ binding affinity and avidity (Koenderman, 2019), IgG subclass, glycosylation patterns and

genetic polymorphism (Bruhns et al, 2009; Pincetic et al, 2014; Plomp et al, 2017; Vidarsson

et al, 2014), stoichiometry of antigen-antibody-ratio (Berger et al, 1996; Lux et al, 2013; Pierson

et al, 2007) and FcγR clustering patterns (Patel et al, 2019). Specifically, Asn297-linked

glycosylation patterns of the IgG Fc domain initiate either pro- or anti-inflammatory effector

pathways by tuning the binding affinity to activating versus inhibitory FcγRs, respectively

(Bohm et al, 2014). However, despite being explored in proof-of-concept studies, the functional

consequences of these ligand features on a given FcγR are still not fully understood. Therefore,

there is an obvious need for an assay platform allowing for the systematic assessment of IC-

mediated FcγR activation.

sICs and immobilized ICs represent unequal and discrete stimuli for the immune system

(Fossati et al, 2002; Granger et al, 2019). Soluble circulating ICs are commonly associated with

certain chronic viral or bacterial infections (Wang & Ravetch, 2015; Yamada et al, 2015) and

autoimmune diseases, such as systemic lupus erythematosus (SLE) or rheumatoid arthritis (RA)

(Antes et al, 1991; Koffler et al, 1971; Zubler et al, 1976). When deposited and accumulating

in tissues, sICs can cause local damage due to inflammatory responses, classified as type III

hypersensitivity (Rajan, 2003). Compared with immobilized local ICs, which recruit immune

cells causing tissue damage (Mayadas et al, 2009; Mulligan et al, 1991; Ward et al, 2016), sIC

related disorders are characterized by systemic inflammation which is reflected by immune cell

exhaustion and senescence (Bano et al, 2019; Chauhan, 2017; Tahir et al, 2015). In order to

resolve sIC-dependent activation of FcγRs in greater detail, we developed a scalable reporter

system suited for two high throughput readouts, capable of quantifying and distinguishing the

activation of single FcγRs. As the assay is also sensitive to stoichiometry and sIC size, we were able to translate the Heidelberger-Kendall precipitation curve to FcγR bioactivity. Compared to currently available ELISA assays detecting sICs by their affinity to C1q-CIC (circulating immune complexes) or C3d, the assay system presented below is strictly specific for IgG immune complexes and integrates sICs of all sizes into single Fcγ receptor bioactivity. Applying reporter cell lines enables very high sensitivity in the low nanomolar range, as signals are biologically amplified compared to biochemical binding based read-outs. Finally, we applied the assay to a clinical setting, measuring sICs in sera from SLE patients. A reporter cell panel expressing murine FcγRs revealed the detection of sICs in the serum of autoimmune-prone diseased mice in preclinical models of lupus and arthritis. Prospectively, this methodology could be instrumental as an experimental and clinical toolbox to unveil sIC-mediated FcγR activation in various autoimmune or infectious diseases.

Results

Experimental assay setup

The assay used in this study was adapted from a previously described cell-based FcγR activation test system designed to measure receptor activation in response to opsonized virus infected cells (Corrales-Aguilar et al, 2013; Kolb et al, 2021) and therapeutic Fc-fusion proteins (Lagasse et al, 2019). We refined the assay to enable selective detection of sICs and expanded the reporter cell line-up (FcγRI: Acc# LT744984; FcγRIIA (131R): Acc# M28697; FcγRIIA(131H): Acc# XP_011507593; FcγRIIB/C: Acc# LT737639; FcγRIIIA(176V): Acc# LT737365; FcγRIIIB(176V): Acc# O75015). Ectodomains of FcγRIIB and FcγRIIC are identical. Second generation reporter cells were generated to improve stable expression of chimeric FcγRs compared to the transfectants used in the original assay (Corrales-Aguilar et al, 2013). To this end, mouse BW5147 cells were transduced as described previously via lentiviral transduction (Corrales-Aguilar et al, 2013; Halenius et al, 2011; Van den Hoecke et al, 2017). Human FcγR

expression on transduced cells after puromycin selection and two consecutive cell sorting steps

was assessed by flow cytometry (Fig. 1A). FcγR activation is measured by surface CD69

expression after 4h of incubation using high thoughput flow cytometry or by quantification of

IL-2 secretion after 16 h of incubation using ELISA. Suspension of IgG or sICs in the liquid

phase is enforced by pre-incubation of a 96 well ELISA microtiter plate with PBS/FCS blocking

buffer (Fig. 1B). To this end, we compared graded concentrations of FCS in the blocking

reagent and measured the threshold at which IgG was no longer adsorbed to the plate and stayed

abundantly in solution. FCS supplementation to 1% (v/v) or higher is sufficient to keep IgG

antibodies in solution. We then set out to test if immobilized IgG can be used as an operational

surrogate for IgG-opsonized cells or immobilized ICs with regard to FcγR activation as

suggested previously (Tanaka et al, 2009). We found no qualitative difference in FcγR

activation between immobilized Rtx, immobilized ICs (Rtx + rec. CD20) or Rtx-opsonized

293T-CD20 cells (Fig. 1C). In contrast, sICs formed by monomeric CD20 antigen (aa 141-188)

and Rtx failed to activate FcγRs even at very high ligand concentrations. We concluded that

FcγR-crosslinking by sICs is only achieved by multivalent antigens but not dimeric ICs. Of

particular note, to reliably and accurately differentiate between strictly soluble and immobilized

including aggregated triggers using this assay, reagents for the generation of synthetic ICs

needed to be of therapy-grade purity. The assay setup is depicted in Fig. 1D.

Detection of human FcγR activation by multimeric sICs

Next, we generated synthetic sICs from recombinant ultrapure molecules to evaluate the assay.

We aimed to avoid the use non-human molecules, misfolded IgG aggregates or IgG-IgG

complexes to generate a most native and defined ligand. To date, there are still few

commercially available human IgG-antigen pairs that meet both the above mentioned high

grade purity requirements while also consisting of at least two antigen monomers. In order to

meet these stipulations we focused on three pairs of multivalent antigens and their respective

mAbs that were available in required amounts enabling large-scale titration experiments;

trimeric rhTNFα:IgG1 infliximab (TNFα:Ifx), dimeric rhVEGFA: IgG1 bevacizumab

(VEGFA/Bvz) and dimeric rhIL-5: IgG1 mepolizumab (IL-5/Mpz). As lymphocytes express

TNFα-receptors I and II while not expressing receptors for IL-5 or VEGFA, we tested whether

the mouse lymphocyte derived BW5147 thymoma reporter cell line is sensitive to high

concentrations of rhTNFα. Toxicity testing revealed that even high concentrations of up to

76.75 nM rhTNFα did not affect viability of reporter cells (Fig. S1). Next, we measured the

dose-dependent activation of human FcγRs comparing immobilized IgG to sICs (TNFα:Ifx)

using the full FcγR reporter cell panel (Fig. 2). Soluble antigen or mAb alone served as negative

controls showing no background activation even at high concentrations. Immobilized rituximab

(Rtx, human IgG1) and immobilized FcγR-specific mouse mAbs served as positive controls for

inter-experimental reference. All FcγRs were activated by immobilized IgG. Only FcγRI failed

to respond when reporter cells were incubated with sICs. Both the IL-2 ELISA as well as the

CD69 expression read-out gave comparable results. Further, using an IL-2 standard, we were

able to quantify FcγR activation (right y-axis). This revealed that the reporter cell lines differ

regarding reactivity which did not correlate with receptor expression. Although the low signals

for FcγRI might be linked to receptor expression, responsiveness was markedly lower compared

to other reporter cell lines (Fig. 1A). Attempts to increase and equalize receptor expression by

repeated cell sorting steps failed, indicating that individual receptors only tolerate limited

molecule densities on the reporter cell surfaces.

Evaluation of human FcγR activation by multimeric sICs

The assay proved to be sensitive to sICs in the nanomolar range. Regarding immobilized IgG,

the detection limit was between 1 to 3 nM. sICs were detected with the following limits

regarding the IL-2 readout: FcγRI – no detection; FcγRIIA(R) – 3 nM; FcγRIIA(H) – 0.2 nM;

FcγRIIB/C – 3 nM; FcγRIIIA – 0.2 nM; FcγRIIIB - 25 nM. We observed that sICs and

immobilized ICs induce largely different signal strength in individual reporter cell lines. FcγRI

and FcγRIIIB were more efficiently activated by immobilized IgG compared to sICs.

Conversely, FcγRIIA(H) and FcγRIIIA were more efficiently activated by sICs. FcγRIIB/C

showed discrepant results when comparing the IL-2 read-out (16 h) with the CD69 read-out (4

h). Here, it seems that a longer activation leads to a stronger signal on immobilized IgG, while

shorter activation slightly favours sIC reactivity. FcγRII(R) looked similar to FcγRIIB/C with

a slightly higher response to sICs at low stimulant concentrations. Nevertheless, the response

to immobilized IgG was higher for both read-outs at higher concentrations. FcγRIIA(H) and

FcγRIIIA proved to be the most sensitive towards sICs stimulation. Notably, the reported

superior interactivity of sICs with FcγRII (H) over FcγRII (R) (Shashidharamurthy et al, 2009)

was not only confirmed using our assay but we also show that this difference is limited to sIC

reactivity and is not seen in immobilized IC reactivity (Fig. 2). Additionally, we measured the

response of select reporter cell lines towards sICs of different composition (VEGFA/Bvz and

IL-5/Mpz). As these sICs incorporate dimeric antigens, we tested if reporter responses were

still comparable (Fig. S2). We observed that responses to sICs were generally lower for

FcγRIIA(R) but comparable for FcγRI, FcγRIIB/C and FcγRIIIA. Of note, FcγRI showed slight

reactivity towards VEGFA/Bvz sICs. Based on the universal transmembrane and cytosolic part

of the FcγR chimeras in our assay we concluded that FcγR ectodomains are intrinsically able

to differentiate between different conformations of sICs and immobilized monomeric IgG

ligands. To validate the data generated by the reporter assay, we determined FcγRIIIA

activation using primary human NK cells isolated from PBMCs of three healthy donors. We

chose NK cells as they mostly express only one type of FcγR similar to the reporter system and

used IL-5/Mpz sICs as NK cells do not respond to IL-5. Measuring a panel of activation markers

and cytokine responses by flow cytometry, we observed a differential activation pattern

depending on ICs being soluble or immobilized at equal molarity (Fig. 3A). While MIP1-β

responses were comparable between the two triggers, degranulation (CD107a) and TNFα

responses showed a trend towards lower activation by sICs compared to immobilized IgG

(Mpz). Strikingly, IFNγ responses were significantly weaker when NK cells were incubated

with sICs compared to immobilized IgG. Next, in order to confirm this to be due to specific

activation of FcγRIIIA, we changed the sIC setup by generating reverse-orientation sICs

consisting of human FcγR-specific mouse mAbs and goat-anti-mouse IgG F(ab)$_2$ fragments

(Fig. S3A). NK cell activation by reverse sICs was compared to NK cell activation by

immobilized FcγR specific mAbs. This confirmed our previous observations. As in roughly

10% of the population NK cells express FcγRIIC (Anania et al, 2019; Breunis et al, 2008; Lisi

et al, 2011; Metes et al, 1998), we also tested reverse sIC activation using an FcγRII specific

mAb. As we did not observe an FcγRII-mediated response, we conclude that FcγRIIC

expression did not play a role in our experiments (Fig. S3B). Importantly, these experiments

validate that all our experimentally synthesized sICs readily activate primary NK cells and

induce immunological effector functions.

Next, primary neutrophils isolated from whole blood samples of four individual donors were

analysed, measuring upregulation of CD11B, CD66B and shedding of L-selectin as markers of

immune complex mediated adhesion and activation (Ilton et al, 1999; Khawaja et al, 2019; Lard

et al, 1999; Zarbock & Ley, 2009) (Fig. 3B). Again, immobilized IgG and sICs activated

primary neutrophils with different efficiency. While both, sICs and immobilized IgG, strongly

induced the shedding of L-selectin, the upregulation of CD11B and CD66B showed a tendency

towards lower activation by sICs compared to immobilized IgG. As neutrophils express both

FcγRIIA and FcγRIIIB, we also individually activated these receptors on neutrophils using

immobilized FcγR-specific mAbs. This revealed that neutrophil activation by FcγRs is mostly

driven by FcγRIIIB. In light of our previous tests (Fig. 2), this explains the reduced activation

by sICs. Taken together, we conclude that primary cells differentiate between opsonized targets

and sICs via inherent features of individual FcγRs as well as by the co-expression of FcγRs

with different sensitivity towards sICs.

Measurement of FcγR activation in response to the molecular size of sICs

We observed that the dimeric CD20:Rtx molecule complex completely failed to trigger FcγR activation (Fig. 1C) while potentially larger sICs, based on multimeric antigens, showed an efficient dose-dependent FcγR activation (Fig. 2, S2). In order to determine whether FcγR signalling responds to changes in sIC size, we cross-titrated amounts of antibody (mAb, infliximab, Ifx) and antigen (Ag, rhTNFα). Specifically, the reporter cells were incubated with sICs of varying mAb:Ag ratio by fixing one parameter and titrating the other. According to the Heidelberger-Kendall precipitation curve (Heidelberger & Kendall, 1929), sIC size depends on the mAb:Ag ratio. sICs of varying sizes result from an excess of either antigen or antibody, leading to the formation of smaller complexes compared to the large molecular complexes formed at around equal molarity. Presumed changes in sIC size were quantified using asymmetrical flow-field flow fractionation (AF4) (Fig. 4A and Table S1). Fig. S4 shows a complete run of an exemplary analysis. Af4 analysis revealed a sIC mean molecular weight of approximately 2130 kDa at a 1:3 ratio (Ifx/TNF-α) with sICs getting smaller with increasing excess of either antigen or antibody, recapitulating a Heidelberger-Kendall-like curve. Incubation of the FcγR reporter cells with sICs of varying size indeed shows that the assay is highly sensitive to changes in sIC size (Fig. 4B). Accordingly, FcγRs showed the strongest responses at mAb:Ag ratios of approximately 1:3. Next, we validated the accuracy of our reporter cell data by subjecting primary human NK cells to the same variation of mAb:Ag stoichiometry. NK cells from three individual donors were measured for MIP1-β upregulation in response to synthetic sICs of varying size and composition (Fig. 4C). Indeed, primary NK cells equally responded to sIC size at the same nanomolar range of stimulating ligand, confirming that the reporter system accurately measures immune cell responses to sICs. Convincingly, NK cell responses to sICs generated from trimeric antigen (TNFα) peaked at a different mAb:Ag ratio compared to NK cell responses to sICs generated from dimeric antigens

(IL-5 and VEGFA). TNFα and VEGFA contribute to the activation of resting NK cells, thus

leading to higher MIP1-β positivity when NK cells are incubated in the presence of excess

antigen. As NK cells do not express IL-5 receptor, this effect is not observed in the presence of

excess IL-5. The data reveal a direct correlation between sIC dimension and effector responses.

Conversely, when changing antibody concentrations using fixed amounts of antigen, a

consistent reduction of NK cell activation is observed in the presence of excess IgG for all three

mAb:Ag pairs.

Quantification of sIC bioactivity in sera of SLE patients

In order to apply the assay to a clinically relevant setting associated with the occurrence of sICs,

we measured circulating sICs present in the serum of SLE patients with varying disease activity.

Sera from 4 healthy donors and 25 SLE patients were investigated for FcγRIIIA and FcγRIIB/C

activation to compare an activating and an inhibitory receptor. Reporter cells readily secreted

mIL-2 in response to patient sera in a dose-dependent manner (Fig. 5A), which was not the case

when sera from healthy controls were tested. We confirm that FcγRIIIA and FcγRIIB/C

activation depends on the presence of serum sICs by comparing the bioactivity of patient serum

before and after polyethylene glycol (PEG) precipitation which is known to deplete sICs (Lux

et al, 2013) (Fig. 5B). Next, we calculated the area under the curve (AUC) values for all 25 SLE

patient titrations and normalized them to the AUC values measured for healthy individuals. The

resulting index values were then correlated with established biomarkers of SLE disease activity,

being anti-dsDNA titers (α-dsDNA) and concentrations of the complement cleavage product

C3d (Fig. 5C). We observed a significant correlation between our FcγRIIIA activation index

values and both disease activity markers (p=0.0465 and p=0.0052, respectively). FcγRIIB/C

activation showed no significant correlation with either biomarker. We assume these

interrelations may be due to the influence of IgG sialylation found to be reduced in active SLE

(Vuckovic et al, 2015). Generally, de-sialylation of IgG leads to stronger binding by the

activating receptors FcγRI, FcγRIIA and FcγRIII while it reduces the binding affinity of the

inhibitory FcγRIIB (Kaneko et al, 2006). In practice, our assay allows the detection and

quantification of clinically relevant sICs in sera from SLE patients as shown here or in synovial

fluid of rheumatoid arthritis patients (Zhao et al, 2021).

Assay application to in vivo mouse models of lupus and arthritis

BW5147 reporter cells stably expressing chimeric mouse as well as rhesus macaque FcγRs have

already been generated using the here described methodology (Kolb et al, 2019; Van den

Hoecke et al, 2017). Next we aimed to translate the assay to clinically relevant mouse models.

FcγR reporter cells expressing chimeric mouse FcγRs were incubated with sera from lupus

(NZB/WF1)(Dubois et al, 1966) or arthritis (K/BxN)(Kouskoff et al, 1996) mice with

symptomatic disease. We chose to determine the stimulation of the activating receptors,

mFcγRIII and mFcγRIV. Incubation with synthetic sICs generated from rhTNFα and mouse-

anti-hTNFα IgG1 showed both of the reporters to be equally responsive to sICs (Fig. 6A).

Parental BW5147 cells expressing no FcγRs served as a control. The sera of three mice per

group were analysed and compared to sera from wildtype C57BL/6 mice, which served as a

healthy control. C57BL/6 mice were chosen, as K/BxN or NZB/WF1 mice show temporal

variability in disease onset and presymptomatic phase. We consistently detected mFcγR

activation by sera from K/BxN or NZB/WF1 but not healthy C57BL/6 mice (Fig. 6B). While

the mFcγRIII responses were generally high and similar between K/BxN and NZB/WF1 mice,

mFcγRIV responsiveness tended to be lower and individually more variable. Altogether, the

assay enables the reliable detection of sICs in sera of mice with immune-complex mediated

diseases making it a promising novel research tool to study the role of sIC formation and FcγR

activation in preclinical mouse models.

Discussion

In this study we established, validated and applied a new assay system that is able to selectively detect soluble multimeric immune complexes as discrete ligands of FcγRs. Our system is sensitive to the size, concentration and composition of sICs. The assay is scalable and supports measurement with human and mouse FcγRs. It provides two readouts suitable for high-throughput analysis: fast CD69 surface expression and quantifiable IL-2 secretion.

A novel assay for the quantification of individual FcγR activation by experimental and clinical sICs.

Our methodology provides a comprehensive system, supporting the assessment of essentially all FcγRs, which presents an advantage over previously developed sIC detection and FcγR activation assays (Aoyama et al, 2019; Cheng et al, 2014; Hsieh et al, 2017; Stopforth et al, 2018; Szittner et al, 2016; Tada et al, 2014). In contrast to currently available commercial assays detecting sICs by C1q-CIC or C3d ELISA in the micromolar range, our assay measures overall sIC bioactivity in the nanomolar range and has a sole specificity for IgG sICs. The new approach presents with hands on technical advances as it allows for the measurement of small or large amounts of samples by a relatively simple *in vitro* assay with high-throughput potential. Favourably, BW5147 reporter cells are largely inert to human cytokines, which provides a key advantage to measure their responsiveness after contact with human samples. Our pilot study demonstrates that sIC-mediated FcγRIIIA activation correlates with conventional SLE disease markers. This is of great value as a recent analysis shows that circulating sICs and IL-6 can predict SLE activity with the higher accuracy compared to conventional clinical SLE biomarkers (Thanadetsuntorn et al, 2018). However, circulating immune complexes in this study were determined using a commercial C1q-binding ELISA, lacking information on immune cell bioactivity of the measured sICs. Our assay should therefore be explored as an addition to the clinician's toolbox which may allow better disease management. Due to the scalability and high-throughput readouts, the assay can also be of use for larger prospective

clinical studies in patients with autoimmune diseases such as SLE or rheumatoid arthritis, where

circulating sICs have long been shown to crucially contribute to tissue damage and disease

manifestations (Koffler et al, 1971; Levinsky, 1978; Levinsky et al, 1977; Nydegger & Davis,

1980; Zubler et al, 1976). Disease-associated, endogenous sICs can also be formed from

multimeric viral and bacterial structural proteins generated during infection (Briant et al, 1996;

Oh et al, 1992; Vuitton et al, 2020), where circulating sICs strongly impact pathogenesis

(Madalinski et al, 1991; Wang & Ravetch, 2015).

Dynamic sIC size measurement and monitoring of bioactivity in sIC-associated diseases

The new sIC approach allowed for a simultaneous functional and biophysical assessment of the

paradigmatic Heidelberger-Kendall precipitation curve (Heidelberger & Kendall, 1929;

Heidelberger & Kendall, 1935). While previous work already revealed that large and small sICs

differentially impact IL-6 production in PBMCs (Lux et al, 2013), the dynamics of FcγR

activation resulting from constant changes in sIC size have not been explored systematically

and lacked resolution of defined FcγR types. We analyzed synthetic sICs formed by highly pure

recombinant components via AF4. Our data document that sIC size is indeed governed by

antibody:antigen ratios covering a wide range of sizes up to several megadaltons. In the

presence of increasing amounts of antibody or antigen deviating from an optimal antibody:ratio,

sIC size steadily decreases. Further, by the measurement of FcγR activation we now translate

physical sIC size directly to a simple but precise biological read-out. In doing so, we show that

sIC size essentially tunes FcγR activation on and off. Thus, our new test system can not only

contribute to the functional detection and quantification of clinically relevant sICs but also

provides a starting point on how to avoid pathological consequences by influencing the sIC

size, for example by administering and monitoring of therapeutic antibodies or recombinant

antigens in controlled amounts, thus becoming relevant in clinical pharmacokinetics.

Limitations of the reporter system and conclusions

There is a wide range of factors, regulating and influencing the sIC-FcγR interaction. These include Fcγ-FcγR binding affinity and avidity (Koenderman, 2019), IgG subclass, IgG glycan profiles and genetic polymorphism (Bruhns et al, 2009; Pincetic et al, 2014; Plomp et al, 2017; Vidarsson et al, 2014), stoichiometry of antigen-antibody-ratio (Berger et al, 1996; Lux et al, 2013; Pierson et al, 2007), FcγR clustering patterns (Patel et al, 2019), downstream signaling (Bournazos et al, 2017; Getahun & Cambier, 2015) and the interaction of FcγR with other receptors (Douek et al, 2009; Ortiz-Stern & Rosales, 2003; Urbaczek et al, 2014; van Egmond et al, 2015; Vanderbruggen et al, 1994). Our assay is sensitive to amount, size and glycosylation of sICs and can readily be adapted to include more FcγR genotypes and polymorphisms by generation of additional reporter cell lines.

The major advancements of this reporter system include i) a high accuracy and resolution regarding FcγR type-specific activation compared to traditional indirect assessment via affinity measurements, ii) a scalable and quantifiable assay providing flexible high-throughput readouts in the nanomolar range, iii) an sIC size sensitive reporter system and iv) a comprehensive panel including all human FcγRs. In practice, the platform is suitable to be implemented into small- or large-scale screening setups in research as well as routine laboratories. Prospectively, the reporter cell approach allows for future adaptation as the cells can be equipped with alternative reporter modules to optimize the methodology for specific applications.

Materials and Methods

Cell culture: All cells were cultured in a 5% CO_2 atmosphere at 37°C. BW5147 mouse thymoma cells (BW, kindly provided by Ofer Mandelboim, Hadassah Hospital, Jerusalem, Israel) were maintained at 3×10^5 to 9×10^5 cells/ml in Roswell Park Memorial Institute medium (RPMI GlutaMAX, Gibco) supplemented with 10% (vol/vol) fetal calf serum (FCS, Biochrom),

sodium pyruvate (1x, Gibco) and β-mercaptoethanol (0.1 mM, Gibco). 293T-CD20 (kindly

provided by Irvin Chen, UCLA (Morizono et al, 2010)) were maintained in Dulbecco's

modified Eagle's medium (DMEM, Gibco) supplemented with 10% (vol/vol) FCS.

BW5147 cell flow cytometry: BW5147 cells were harvested by centrifugation at 900 g and RT

from the suspension culture. $1x10^6$ cells were stained with PE- conjugated anti-human FcγR

mAbs (BD) or a PE-TexasRed-conjugated human IgG-Fc fragment (Rockland) for 1h at 4°C in

PBS/3%FCS. After 3 washing steps in PBS/3%FCS, the cells were transferred to Flow

cytometry tubes (BD) and analysed using BD LSR Fortessa and FlowJo (V10) software. Cells

sorting was performed at the Lighthouse core facility of the University Hospital Freiburg using

receptor staining (BD Pharmingen, PE-conjugated).

Lentiviral transduction: Lentiviral transduction of BW5147 cells was performed as described

previously (Halenius et al, 2011; Kolb et al, 2019; Van den Hoecke et al, 2017). In brief,

chimeric FcγR-CD3ζ constructs (Corrales-Aguilar et al, 2013) were cloned into a

pUC2CL6IPwo plasmid backbone. For every construct, one 10-cm dish of packaging cell line

at roughly 70% density was transfected with the target construct and two supplementing vectors

providing the VSV gag/pol and VSV-G-env proteins (6 µg of DNA each) using

polyethylenimine (22.5 µg/ml, Sigma) and Polybrene (4 µg/ml; Merck Millipore) in a total

volume of 7 ml (2 ml of a 15-min-preincubated transfection mix in serum-free DMEM added

to 5 ml of fresh full DMEM). After a medium change, virus supernatant harvested from the

packaging cell line 2 days after transfection was then incubated with target BW cells overnight

(3.5 ml of supernatant on 10^6 target cells), followed by expansion and pool selection using

complete medium supplemented with 2 µg/ml of puromycin (Sigma) over a one week culture

period.

human IgG suspension ELISA: 1 µg of IgG1 (rituximab in PBS, 50 µl/well) per well was incubated on a 96well microtiter plate (NUNC Maxisorp) pre-treated (2h at RT) with PBS supplemented with varying percentages (v/v) of FCS (PAN Biotech). IgG1 bound to the plates was detected using an HRP-conjugated mouse-anti-human IgG mAb (Jackson ImmunoResearch).

Recombinant antigens and monoclonal antibodies to form sICs: Recombinant human (rh) cytokines TNF, IL-5, and VEGFA were obtained from Stem Cell technologies. Recombinant CD20 was obtained as a peptide (aa141-188, Acc# P11836) containing the binding region of rituximab (Creative Biolabs). FcγR-specific mAbs were obtained from Stem Cell technologies (CD16: clone 3G8; CD32: IV.3). Reverse sICs were generated from these receptor-specific antibodies using goat-anti-mouse IgG F(ab)$_2$ fragments (Invitrogen) in a 1:1 ratio. Pharmaceutically produced humanized monoclonal IgG1 antibodies infliximab (Ifx), bevacizumab (Bvz), mepolizumab (Mpz) and rituximab (Rtx) were obtained from the University Hospital Pharmacy Freiburg. Mouse anti-hTNFα (IgG2b, R&D Systems, 983003) was used to generate sICs reactive with mouse FcγRs. sICs were generated by incubation of antigens and antibodies in reporter cell medium or PBS for 2 h at 37°C.

FcγR receptor activation assay: FcγR activation was measured adapting a previously described cell-based assay (Corrales-Aguilar et al, 2014; Corrales-Aguilar et al, 2013). The assay was modified to measure FcγR activation in solution. Briefly, 2×10^5 mouse BW-FcγR (BW5147) reporter cells were incubated with synthetic sICs or diluted serum in a total volume of 100 µl for 16 h at 37°C and 5% CO_2. Incubation was performed in a 96-well ELISA plate (Nunc maxisorp) pre-treated with PBS/10% FCS (v/v) for 1 h at 4°C. Immobilized IgG was incubated in PBS on the plates prior to PBS/10% FCS treatment. After 4h incubation, surface mouse CD69 expression was measured using a high throughout sampler (HTS)-FACS. Reporter cell

mouse IL-2 secretion was quantified after 16 h of incubation via anti-IL-2 ELISA as described

earlier (Corrales-Aguilar et al, 2013).

High throughout sampler flow cytometry (HTS-FACS): After 4h of stimulation, $1x10^5$ BW5147

reporter cells were stained with APC-conjugated anti-mCD69 (Biolegend; CD69: H1.2F3;

1:100) for 30min at 4°C in PBS/3%FCS. Cells were transferred to a U Form 96well Microplate

(Greiner 650101) and analysed by flow cytometry (BD Fortessa). High Throughput mode was

designed within BD FACSDiva software using HTS mode with the following parameters:

sample flow rate 2µl/s, sample volume 10µl, mixing volume 50µl, mixing speed 200µl/s,

number of mixes 2 cycles and wash volume 200µl.

BW5147 toxicity test: Cell counting was performed using a Countess II (Life Technologies)

according to supplier instructions. Cell toxicity was measured as a ratio between live and dead

cells judged by trypan blue staining over a 16 h time frame in a 96well format (100 µl volume

per well). BW5147 cells were mixed 1:1 with trypan blue (Invitrogen) and analysed using a

Countess II. rhTNFα was diluted in complete medium.

NK cell activation flow cytometry: PBMC were purified from donor blood using Lymphocyte

separation Media (Anprotec). Blood draw and PBMC purification from donors was approved

by vote 474/18 (ethical review committee, University of Freiburg). Primary NK cells were

separated from donor PBMCs via magnetic bead negative selection (Stem Cell technologies)

and NK cell purity was confirmed via staining of CD3 (Biolegend, clone HIT3a), CD16

(Biolegend, clone 3G8) and CD56 (Miltenyi Biotec, clone AF12-7H3). 96well ELISA plates

(Nunc Maxisorp) were pre-treated with PBS/10% FCS (v/v) for 1 h at 4°C. NK cells were

stimulated in pre-treated plates and incubated at 37°C and 5% CO_2 for 4 h. Golgi Plug and

Golgi Stop solutions (BD) were added as suggested by supplier. CD107a (APC, BD, H4A3)

specific conjugated mAb was added at the beginning of the incubation period. Following the

stimulation period, MIP-1β (PE, BD Pharmingen), IFNγ (BV-510, Biolegends, 4SB3) and

TNFα (PE/Cy7, Biolegends, MAB11) production was measured via intracellular staining

Cytokines (BD, CytoFix/CytoPerm, Kit as suggested by the supplier). 50 ng/ml PMA

(InvivoGen) + 0.5 µM Ionomycin (InvivoGen) were used as a positive stimulation control for

NK cell activation. After 3 washing steps in PBS/3%FCS, the cells were transferred to Flow

cytometry tubes (BD) and analysed using a BD FACS Fortessa and FlowJo (V10) software.

Neutrophil adhesion and activation flow cytometry: Human primary neutrophil granulocytes

were isolated from whole blood of healthy donors via magnetic bead negative selection

(Stemcell #19666). 96well ELISA plates (Nunc Maxisorp) were pre-treated with PBS/10% FCS

(v/v) for 1 h at 4°C. Per reaction, $2x10^5$ cells/ml neutrophils were stimulated with ICs in Roswell

Park Memorial Institute medium (RPMI GlutaMAX, Gibco) supplemented with 10% (vol/vol)

fetal calf serum (FCS, Biochrom) and incubated at 37°C and 5% CO_2 for 30 min. Adhesion and

activation markers of neutrophils were measured by surface staining of CD11B (APC,

Biolegend, ICRF44), CD66B (FITC, Stemcell, G10F5) and L-selectin (PE, Biolegend, DREG-

56)(Ilton et al, 1999; Khawaja et al, 2019; Lard et al, 1999; Veen et al, 1998). Cells were then

analysed by flow cytometry. FcγRII or FcγRIII cross-linking controls were performed by

immobilization of receptor specific mAbs (Stem cell technologies, IV.3 and 3G8) before the

ELISA plate was blocked.

Asymmetric flow field flow fractionation (AF4): The AF4 system consisted of a flow controller

(Eclipse AF4, Wyatt), a MALS detector (DAWN Heleos II, Wyatt), a UV detector (1260

Infinity G1314F, Agilent) and the separation channel (SC channel, PES membrane, cut-off 10

kDa, 490 µm spacer, wide type, Wyatt). Elution buffer: 1.15 g/L Na_2HPO_4 (Merck), 0.20 g/L

NaH_2PO_4 x H_2O (Merck), 8.00 g/L NaCl (Sigma) and 0,20 g/L NaN_3 (Sigma), adjusted to pH

7.4, filtered through 0.1 μm. AF4 sequence (Vx = cross flow in mL/min): (a) elution (2 min,

Vx: 1.0); (b) focus (1 min, Vx: 1.0), focus + inject (1 min, Vx: 1.0, inject flow: 0.2 mL/min),

repeated three times; (c) elution (30 min, linear Vx gradient: 1.0 to 0.0); (d) elution (15 min,

Vx: 0.0); (e) elution + inject (5 min, Vx: 0.0). A total protein mass of 17±0.3 μg (Ifx, rhTNFα

or ICs, respectively) was injected. The eluted sample concentration was calculated from the UV

signal at 280 nm using extinction coefficients of 1.240 mL/(mg cm) or 1.450 mL/(mg cm) in

the case of TNFα or Ifx, respectively. For the ICs, extinction coefficients were not available

and difficult to calculate as the exact stoichiometry is not known. An extinction coefficient of

1.450 mL/(mg cm) was used for calculating the molar masses of all ICs. Especially in the case

of ICs rich in TNFα, the true coefficients should be lower, and the molar masses of these

complexes are overestimated by not more than 14 %. The determined molar masses for TNFα-

rich complexes are therefore biased but the observed variations in molar mass for the different

ICs remain valid. The mass-weighted mean of the distribution of molar masses for each sample

was calculated using the ASTRA 7 software package (Wyatt).

SLE patient cohort: Sera from patients with SLE were obtained from the Immunologic,

Rheumatologic Biobank (IR-B) of the Department of Rheumatology and Clinical Immunology.

Biobanking and the project were approved by the local ethical committee of the University of

Freiburg (votes 507/16 and 624/14). All patients who provided blood to the biobank had

provided written informed consent. Ethical Statement: The study was designed in accordance

with the guidelines of the Declaration of Helsinki (revised 2013). Patients with SLE ($n = 25$)

and healthy controls ($n = 4$) were examined. All patients met the revised ACR classification

criteria for SLE. Disease activity was assessed using the SLEDAI-2K score. C3d levels were

analyzed in EDTA plasma using rocket double decker immune-electrophoresis with antisera

against C3d (Polyclonal Rabbit Anti-Human C3d Complement, Agilent) and C3c (Polyclonal

Rabbit Anti-Human C3c Complement Agilent) as previously described (Rother et al, 1993).

Anti-human dsDNA antibodies titers were determined in serum using an anti-dsDNA IgG

ELISA kit (diagnostik-a GmbH).

Patient serum IC precipitation: For polyethylene glycol (PEG) precipitation human sera were

mixed with PEG 6000 (Sigma-Aldrich) in PBS at a final concentration of 10% PEG 6000. After

overnight incubation at 4°C, ICs were precipitated by centrifugation at 2000 x g for 30 min at

4 °C, pellets were washed once with PEG 6000 and then centrifuged at 2000 x g for 20 min at

4 °C. Supernatants were harvested and precipitates re-suspended in pre-warmed PBS for 1 h at

37 °C. IgG concentrations of serum, precipitates and supernatants obtained after precipitation

were quantified by Nanodrop (Thermo Scientific™) measurement.

Mice and Models: Animal experiments were approved by the local governmental commission

for animal protection of Freiburg (Regierungspräsidium Freiburg, approval no. G16/59 and

G19/21). Lupus-prone (NZBxNZW)F1 mice (NZB/WF1) were generated by crossing

NZB/BlNJ mice with NZW/LacJ mice, purchased from The Jackson Laboratory. KRNtg mice

were obtained from F. Nimmerjahn (Universität Erlangen-Nürnberg) with the permission of D.

Mathis and C. Benoist (Harvard Medical School, Boston, MA), C57BL/6 mice (BL/6) and

NOD/ShiLtJArc (NOD/Lt) mice were obtained from the Charles River Laboratories. K/BxN

(KRNtgxNOD)F1 mice (K/BxN) were obtained by crossing KRNtg mice and NOD/Lt mice.

All mice were housed in a 12-h light/dark cycle, with food and water ad libitum. Mice were

euthanized and blood collected for serum preparation from 16 weeks old BL/6 animals, from

16 weeks old arthritic K/BxN animals and from 26 – 38 weeks old NZB/WF1 mice with

established glomerulonephritis.

Statistical analyses: Statistical analyses were performed using Graphpad Prism software (v6)

and appropriate tests.

Supplementary Materials

Fig. S1. rhTNFα is not toxic to mouse lymphocyte BW5147 cells even at high concentrations.

Fig. S2. FcγRs are activated by VEGFA and IL-5 sICs

Fig. S3. Distinct activation patterns of NK cells incubated with inverse sICs

Fig. S4. AF4 elution profiles of Ifx/TNFα-immune complexes.

Table S1. Analysis of the molar mass distribution of ICs from AF4 data.

Figure legends

Fig. 1. Establishment of a cell-based reporter assay measuring FcγR activation in response to sICs. A) BW5147 reporter cells stably expressing human FcγR-ζ chain chimeras or BW5147 parental cells (grey/dashed) were stained with FcγR specific conjugated mAbs as indicated and measured for surface expression of FcγRs via flow cytometry. B) FCS coating of an ELISA microtiter plate allows for suspension of subsequently added IgG. Plate bound IgG was quantified via ELISA. C) Immobilized IC, immobilized IgG and IgG opsonized cells represent qualitatively similar ligands for FcγRs. Response curves of human FcγRs activated by opsonized cells (293T cells stably expressing CD20 + Rituximab [Rtx]), immobilized IC (rec. soluble CD20 + Rtx) and immobilized IgG (Rtx). sICs formed using monovalent antigen (rec. soluble CD20 + Rtx) do not activate human FcγRs. X-Axis shows sample concentration determined by antibody molarity. Y-Axis shows FcγR activation determined by reporter cell mouse IL-2 production (OD 450nm). Two independent experiments performed in technical duplicates. Error bars = SD. D) Schematic of used assay setups. BW5147 reporter cells expressing chimeric human FcγR receptors express endogenous CD69 or secrete mouse IL-2 in response to FcγR activation by clustered IgG. sICs are generated using mAbs and multivalent antigens. sIC suspension requires pre-blocking of an ELISA plate using PBS supplemented with 10% FCS (FCS coat, grey-dashed).

Fig. 2. FcγRs are activated by sICs formed from multivalent antigens. Ultra-pure antigen (Ag, TNF-α) mixed with therapy-grade mAb (infliximab, Ifx) was used to generate sICs. X-Axis: concentrations of stimulant expressed as molarity of either mAb or Ag monomer and IC (expressed as mAb molarity) at a mAb:Ag ratio of 1:2. Soluble antigen or soluble antibody alone served as negative controls and were not sufficient to activate human FcγRs. Immobilized IgG (Rtx) or immobilized FcγR-specific mAbs served as positive controls. Two independent

experiments performed in technical duplicates. Error bars = SD. Error bars smaller than

symbols are not shown. Left panel: IL-2 quantification 16 h after reporter cell activation.

Background (blank) was subtracted (dashed line). IL-2 was measured via anti-IL-2 ELISA

(A_{450nm}) and IL-2 concentrations were calculated from an IL-2 standard. Right panel: Reporter

cell CD69 expression 4 h post trigger was measured using flow cytometry. MFI were

normalized to untreated cells (ctrl.) and are presented as fold-change increase.

Fig. 3. The FcγR-dependent activation pattern of primary NK cells or primary

neutrophils depends on IC solubility. A) Negatively selected primary NK cells purified from

PBMCs of three healthy donors were tested for activation markers using flow cytometry. NK

cells were incubated with immobilized IgG (mepolizumab, Mpz), soluble IC (Mpz:IL-5 = 1:1),

soluble Mpz or soluble IL-5 (all at 200 nM, 10^6 cells). Incubation with PMA and Ionomycin

(Iono) served as a positive control. Incubation with medium alone served as a negative control.

Means of technical duplicates. Error bars = SD. One-way ANOVA (Tukey); *p<0.05,

**p<0.01, ***p<0.001, ****p<0.0001. B) Negatively selected primary neutrophils purified

from whole blood of four healthy donors were tested for adhesion and activation markers using

flow cytometry. Neutrophils were incubated with immobilized IgG (Mpz), soluble IC (Mpz:IL-

5 = 1:1), soluble Mpz or soluble IL-5 (all at 200 nM, $2*10^5$ cells). Incubation with PMA or

immobilized rituximab served as positive controls. Incubation with medium served as a

negative control. Immobilized FcγRII and FcγRIII specific mAbs served as functional controls.

Mean florescence intensity (MFI) values at t=30 minutes of incubation are presented as increase

over t=0 min. Means of technical duplicates. Error bars = SD. Two-way ANOVA compared to

medium (Dunnett); *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Fig. 4. FcγRIIB/C and FcγR-IIIA respond to sIC size reproducing a Heidelberger-Kendall

like precipitation curve. A) infliximab (mAb) and rhTNFα (Ag) were mixed at different ratios

(17 μg total protein, calculated from monomer molarity) and analysed via AF4. sIC size is

maximal at a 1:3 ratio of mAb:Ag and reduced when either mAb or Ag are given in excess.

$<M>_w$ = mass-weighted mean of the molar mass distribution. Three independent experiments.

Error bars = SD. Data taken from Table S1. One complete run analysis is shown in Fig. S2. B)

sICs of different size were generated by cross-titration according to the AF4 determination.

Reporter cells were incubated with fixed amounts of either mAb (infliximab, left) or Ag

(rhTNFα, right) and titrated amounts of antigen or antibody, respectively. X-Axis shows

titration of either antigen or antibody, respectively (TNFα calculated as monomer). Two

independent experiments performed in technical duplicates. Error bars = SD. C) Purified

primary NK cells from three different donors were incubated with cross-titrated sICs as in A.

NK cells were measured for MIP-1β expression (% positivity). Incubation with PMA and

Ionomycin served as a positive control. Incubation with medium alone served as a negative

control. Measured in technical duplicates. Error bars = SD.

Fig. 5. The reporter assay enables quantification of serum-derived sICs from SLE

patients. Serum derived sIC from systemic lupus erythematosus (SLE) patients activate human

FcγR reporter cells. 25 patients and 4 healthy control individuals were separated into three

groups for measurement. A) Experiments shown for an exemplary group of 8 SLE patients and

two healthy individuals. Dose-dependent reactivity of FcγRs IIIA and IIB/C was observed only

for SLE patient sera and not for sera from healthy individuals. One exemplary experiment

performed in technical duplicates. Error bars = SD. B) Activation of FcγRs IIB/C and IIIA by

patient serum is mediated by serum derived sICs. Patient serum samples were depleted of sICs

by PEG precipitation and the supernatant (SN) was compared to untreated serum regarding

FcγR activation (left). One experiment performed in technical duplicates. Error bars =SD. IgG

concentration in the precipitate (PC), supernatant (SN) or unfractionated serum respectively is

shown in the bar graph (right). C) FcγR activation data from A was compared to conventional

SLE disease markers (α-dsDNA levels indicated as IU/ml or C3d concentrations indicated as

mg/L). FcγR activation from a dose-response curve as in A was calculated as area under curve

(AUC) for each SLE patient (n=25) or healthy individual (n=4) and expressed as fold change

compared to the healthy control mean. SLE patients with α-dsDNA levels below 50 IU/ml and

C3d values below 6 mg/L were excluded. One-tailed Spearman's.

Fig. 6. The reporter assay can be applied to mouse models of autoimmune disease. A)

Reporter cells expressing mFcγRIII, mFcγRIV or parental BW5147 cells were incubated with

titrated amounts of synthetic sICs generated from rhTNFα and mouse-anti-hTNFα at a 1:1 ratio

by mass. One experiment performed in technical duplicates. Error bars = SD. B) Titrations of

3 mouse sera per group (C57BL/6, K/BxN or NZB/WF1) were incubated with mFcγR reporter

cells and FcγR activation was assessed as described above. Sera from BL/6 mice served as

negative control. Two independent experiments in technical duplicates. Error bars = SD.

References

Anania JC, Chenoweth AM, Wines BD, Hogarth PM (2019) The Human FcgammaRII (CD32) Family of Leukocyte FcR in Health and Disease. Front Immunol 10**:** 464

Antes U, Heinz HP, Schultz D, Brackertz D, Loos M (1991) C1q-bearing immune complexes detected by a monoclonal antibody to human C1q in rheumatoid arthritis sera and synovial fluids. Rheumatol Int 10**:** 245-250

Aoyama M, Tada M, Ishii-Watabe A (2019) A Cell-Based Reporter Assay Measuring the Activation of Fc Gamma Receptors Induced by Therapeutic Monoclonal Antibodies. Methods Mol Biol 1904**:** 423-429

Bano A, Pera A, Almoukayed A, Clarke THS, Kirmani S, Davies KA, Kern F (2019) CD28 (null) CD4 T-cell expansions in autoimmune disease suggest a link with cytomegalovirus infection. F1000Res 8

Berger S, Ballo H, Stutte HJ (1996) Immune complex-induced interleukin-6, interleukin-10 and prostaglandin secretion by human monocytes: A network of pro- and anti-inflammatory cytokines dependent on the antigen:antibody ratio. Eur J Immunol 26**:** 1297-1301

Bohm S, Kao D, Nimmerjahn F (2014) Sweet and sour: the role of glycosylation for the anti-inflammatory activity of immunoglobulin G. Current topics in microbiology and immunology 382**:** 393-417

Bournazos S, Wang TT, Dahan R, Maamary J, Ravetch JV (2017) Signaling by Antibodies: Recent Progress. Annu Rev Immunol 35**:** 285-311

Breunis WB, van Mirre E, Bruin M, Geissler J, de Boer M, Peters M, Roos D, de Haas M, Koene HR, Kuijpers TW (2008) Copy number variation of the activating FCGR2C gene predisposes to idiopathic thrombocytopenic purpura. Blood 111**:** 1029-1038

Briant L, Coudronniere N, Robert-Hebmann V, Benkirane M, Devaux C (1996) Binding of HIV-1 virions or gp120-anti-gp120 immune complexes to HIV-1-infected quiescent peripheral blood mononuclear cells reveals latent infection. J Immunol 156**:** 3994-4004

Bruhns P (2012) Properties of mouse and human IgG receptors and their contribution to disease models. Blood 119**:** 5640-5649

Bruhns P, Iannascoli B, England P, Mancardi DA, Fernandez N, Jorieux S, Daeron M (2009) Specificity and affinity of human Fcgamma receptors and their polymorphic variants for human IgG subclasses. Blood 113**:** 3716-3725

Bruhns P, Jonsson F (2015) Mouse and human FcR effector functions. Immunol Rev 268**:** 25-51

Chauhan AK (2017) Editorial: Immune Complexes in Disease Pathology. Front Immunol 8**:** 173

Cheng ZJ, Garvin D, Paguio A, Moravec R, Engel L, Fan F, Surowy T (2014) Development of a robust reporter-based ADCC assay with frozen, thaw-and-use cells to measure Fc effector function of therapeutic antibodies. J Immunol Methods 414**:** 69-81

Corrales-Aguilar E, Trilling M, Hunold K, Fiedler M, Le VT, Reinhard H, Ehrhardt K, Merce-Maldonado E, Aliyev E, Zimmermann A et al (2014) Human cytomegalovirus Fcgamma binding

proteins gp34 and gp68 antagonize Fcgamma receptors I, II and III. PLoS pathogens 10: e1004131

Corrales-Aguilar E, Trilling M, Reinhard H, Merce-Maldonado E, Widera M, Schaal H, Zimmermann A, Mandelboim O, Hengel H (2013) A novel assay for detecting virus-specific antibodies triggering activation of Fcgamma receptors. Journal of immunological methods 387: 21-35

Douek DC, Rittirsch D, Flierl MA, Day DE, Nadeau BA, Zetoune FS, Sarma JV, Werner CM, Wanner GA, Simmen H-P et al (2009) Cross-Talk between TLR4 and FcγReceptorIII (CD16) Pathways. Plos Pathog 5: e1000464

Dubois EL, Horowitz RE, Demopoulos HB, Teplitz R (1966) NZB/NZW mice as a model of systemic lupus erythematosus. JAMA 195: 285-289

Duchemin AM, Ernst LK, Anderson CL (1994) Clustering of the High-Affinity Fc Receptor for Immunoglobulin-G (Fc-Gamma-Ri) Results in Phosphorylation of Its Associated Gamma-Chain. J Biol Chem 269: 12111-12117

Fossati G, Bucknall RC, Edwards SW (2002) Insoluble and soluble immune complexes activate neutrophils by distinct activation mechanisms: changes in functional responses induced by priming with cytokines. Ann Rheum Dis 61: 13-19

Getahun A, Cambier JC (2015) Of ITIMs, ITAMs, and ITAMis: revisiting immunoglobulin Fc receptor signaling. Immunol Rev 268: 66-73

Granger V, Peyneau M, Chollet-Martin S, de Chaisemartin L (2019) Neutrophil Extracellular Traps in Autoimmunity and Allergy: Immune Complexes at Work. Front Immunol 10: 2824

Greenberg S, Chang P, Silverstein SC (1994) Tyrosine phosphorylation of the gamma subunit of Fc gamma receptors, p72syk, and paxillin during Fc receptor-mediated phagocytosis in macrophages. J Biol Chem 269: 3897-3902

Halenius A, Hauka S, Dolken L, Stindt J, Reinhard H, Wiek C, Hanenberg H, Koszinowski UH, Momburg F, Hengel H (2011) Human cytomegalovirus disrupts the major histocompatibility complex class I peptide-loading complex and inhibits tapasin gene transcription. J Virol 85: 3473-3485

Heidelberger M, Kendall FE (1929) A Quantitative Study of the Precipitin Reaction between Type Iii Pneumococcus Polysaccharide and Purified Homologous Antibody. J Exp Med 50: 809-823

Heidelberger M, Kendall FE (1935) The Precipitin Reaction between Type Iii Pneumococcus Polysaccharide and Homologous Antibody : Iii. A Quantitative Study and a Theory of the Reaction Mechanism. J Exp Med 61: 563-591

Hsieh YT, Aggarwal P, Cirelli D, Gu L, Surowy T, Mozier NM (2017) Characterization of Fc gamma RIIIA effector cells used in in vitro ADCC bioassay: Comparison of primary NK cells with engineered NK-92 and Jurkat T cells. Journal of Immunological Methods 441: 56-66

Ilton MK, Langton PE, Taylor ML, Misso NL, Newman M, Thompson PJ, Hung J (1999) Differential expression of neutrophil adhesion molecules during coronary artery surgery with cardiopulmonary bypass. J Thorac Cardiovasc Surg 118: 930-937

Kaneko Y, Nimmerjahn F, Ravetch JV (2006) Anti-inflammatory activity of immunoglobulin G resulting from Fc sialylation. Science 313: 670-673

Khawaja AA, Pericleous C, Ripoll VM, Porter JC, Giles IP (2019) Autoimmune rheumatic disease IgG has differential effects upon neutrophil integrin activation that is modulated by the endothelium. Sci Rep 9**:** 1283

Kiefer F, Brumell J, Al-Alawi N, Latour S, Cheng A, Veillette A, Grinstein S, Pawson T (1998) The Syk protein tyrosine kinase is essential for Fc gamma receptor signaling in macrophages and neutrophils. Mol Cell Biol 18**:** 4209-4220

Koenderman L (2019) Inside-Out Control of Fc-Receptors. Front Immunol 10**:** 544

Koffler D, Agnello V, Thoburn R, Kunkel HG (1971) Systemic lupus erythematosus: prototype of immune complex nephritis in man. J Exp Med 134**:** 169-179

Kolb P, Hoffmann K, Sievert A, Reinhard H, Merce-Maldonado E, Le-Trilling VTK, Halenius A, Gutle D, Hengel H (2021) Human Cytomegalovirus antagonizes activation of Fcgamma receptors by distinct and synergizing modes of IgG manipulation. Elife 10

Kolb P, Sijmons S, McArdle MR, Taher H, Womack J, Hughes C, Ventura A, Jarvis MA, Stahl-Hennig C, Hansen S et al (2019) Identification and Functional Characterization of a Novel Fc Gamma-Binding Glycoprotein in Rhesus Cytomegalovirus. J Virol 93

Kouskoff V, Korganow AS, Duchatelle V, Degott C, Benoist C, Mathis D (1996) Organ-specific disease provoked by systemic autoimmunity. Cell 87**:** 811-822

Lagasse HAD, Hengel H, Golding B, Sauna ZE (2019) Fc-Fusion Drugs Have FcgammaR/C1q Binding and Signaling Properties That May Affect Their Immunogenicity. AAPS J 21**:** 62

Lard LR, Mul FPJ, de Haas M, Roos D, Duits AJ (1999) Neutrophil activation in sickle cell disease. J Leukocyte Biol 66**:** 411-415

Levinsky RJ (1978) Role of circulating soluble immune complexes in disease. Arch Dis Child 53**:** 96-99

Levinsky RJ, Cameron JS, Soothill JF (1977) Serum immune complexes and disease activity in lupus nephritis. Lancet 1**:** 564-567

Lisi S, Sisto M, Lofrumento DD, D'Amore S, D'Amore M (2011) Advances in the understanding of the Fc gamma receptors-mediated autoantibodies uptake. Clin Exp Med 11**:** 1-10

Lu LL, Suscovich TJ, Fortune SM, Alter G (2018) Beyond binding: antibody effector functions in infectious diseases. Nat Rev Immunol 18**:** 46-61

Luo Y, Pollard JW, Casadevall A (2010) Fcgamma receptor cross-linking stimulates cell proliferation of macrophages via the ERK pathway. J Biol Chem 285**:** 4232-4242

Lux A, Yu X, Scanlan CN, Nimmerjahn F (2013) Impact of immune complex size and glycosylation on IgG binding to human FcgammaRs. J Immunol 190**:** 4315-4323

Madalinski K, Burczynska B, Heermann KH, Uy A, Gerlich WH (1991) Analysis of viral proteins in circulating immune complexes from chronic carriers of hepatitis B virus. Clin Exp Immunol 84**:** 493-500

Mayadas TN, Tsokos GC, Tsuboi N (2009) Mechanisms of immune complex-mediated neutrophil recruitment and tissue injury. Circulation 120**:** 2012-2024

Metes D, Ernst LK, Chambers WH, Sulica A, Herberman RB, Morel PA (1998) Expression of
functional CD32 molecules on human NK cells is determined by an allelic polymorphism of the
FcgammaRIIC gene. Blood 91**:** 2369-2380

Morizono K, Ku A, Xie Y, Harui A, Kung SK, Roth MD, Lee B, Chen IS (2010) Redirecting
lentiviral vectors pseudotyped with Sindbis virus-derived envelope proteins to DC-SIGN by
modification of N-linked glycans of envelope proteins. J Virol 84**:** 6923-6934

Mulligan MS, Hevel JM, Marletta MA, Ward PA (1991) Tissue-Injury Caused by Deposition of
Immune-Complexes Is L-Arginine Dependent. P Natl Acad Sci USA 88**:** 6338-6342

Nimmerjahn F, Ravetch JV (2006) Fcgamma receptors: old friends and new family members.
Immunity 24**:** 19-28

Nimmerjahn F, Ravetch JV (2008) Fcgamma receptors as regulators of immune responses.
Nat Rev Immunol 8**:** 34-47

Nimmerjahn F, Ravetch JV (2010) Antibody-mediated modulation of immune responses.
Immunol Rev 236**:** 265-275

Nydegger UE, Davis JSt (1980) Soluble immune complexes in human disease. CRC Crit Rev
Clin Lab Sci 12**:** 123-170

Oh SK, Cruikshank WW, Raina J, Blanchard GC, Adler WH, Walker J, Kornfeld H (1992)
Identification of HIV-1 envelope glycoprotein in the serum of AIDS and ARC patients. J Acquir
Immune Defic Syndr (1988) 5**:** 251-256

Ortiz-Stern A, Rosales C (2003) Cross-talk between Fc receptors and integrins. Immunol Lett
90**:** 137-143

Patel KR, Roberts JT, Barb AW (2019) Multiple Variables at the Leukocyte Cell Surface Impact
Fc gamma Receptor-Dependent Mechanisms. Front Immunol 10**:** 223

Pierson TC, Xu Q, Nelson S, Oliphant T, Nybakken GE, Fremont DH, Diamond MS (2007) The
stoichiometry of antibody-mediated neutralization and enhancement of West Nile virus
infection. Cell Host Microbe 1**:** 135-145

Pincetic A, Bournazos S, DiLillo DJ, Maamary J, Wang TT, Dahan R, Fiebiger BM, Ravetch
JV (2014) Type I and type II Fc receptors regulate innate and adaptive immunity. Nat Immunol
15**:** 707-716

Plomp R, Ruhaak LR, Uh HW, Reiding KR, Selman M, Houwing-Duistermaat JJ, Slagboom
PE, Beekman M, Wuhrer M (2017) Subclass-specific IgG glycosylation is associated with
markers of inflammation and metabolic health. Sci Rep 7**:** 12325

Rajan TV (2003) The Gell-Coombs classification of hypersensitivity reactions: a re-
interpretation. Trends in immunology 24**:** 376-379

Rother E, Lang B, Coldewey R, Hartung K, Peter HH (1993) Complement split product C3d as
an indicator of disease activity in systemic lupus erythematosus. Clin Rheumatol 12**:** 31-35

Shashidharamurthy R, Zhang F, Amano A, Kamat A, Panchanathan R, Ezekwudo D, Zhu C,
Selvaraj P (2009) Dynamics of the interaction of human IgG subtype immune complexes with
cells expressing R and H allelic forms of a low-affinity Fc gamma receptor CD32A. J Immunol
183**:** 8216-8224
876

877 Stopforth RJ, Oldham RJ, Tutt AL, Duriez P, Chan HTC, Binkowski BF, Zimprich C, Li D,
878 Hargreaves PG, Cong M et al (2018) Detection of Experimental and Clinical Immune
879 Complexes by Measuring SHIP-1 Recruitment to the Inhibitory FcgammaRIIB. J Immunol 200**:**
880 1937-1950
881

882 Szittner Z, Bentlage AEH, Rovero P, Migliorini P, Lorand V, Prechl J, Vidarsson G (2016)
883 Label-free detection of immune complexes with myeloid cells. Clin Exp Immunol 185**:** 72-80
884

885 Tada M, Ishii-Watabe A, Suzuki T, Kawasaki N (2014) Development of a cell-based assay
886 measuring the activation of FcgammaRIIa for the characterization of therapeutic monoclonal
887 antibodies. Plos One 9**:** e95787
888

889 Tahir S, Fukushima Y, Sakamoto K, Sato K, Fujita H, Inoue J, Uede T, Hamazaki Y, Hattori M,
890 Minato N (2015) A CD153+CD4+ T follicular cell population with cell-senescence features
891 plays a crucial role in lupus pathogenesis via osteopontin production. J Immunol 194**:** 5725-
892 5735
893

894 Tanaka M, Krutzik SR, Sieling PA, Lee DJ, Rea TH, Modlin RL (2009) Activation of Fc gamma
895 RI on monocytes triggers differentiation into immature dendritic cells that induce autoreactive
896 T cell responses. J Immunol 183**:** 2349-2355
897

898 Thanadetsuntorn C, Ngamjanyaporn P, Setthaudom C, Hodge K, Saengpiya N, Pisitkun P
899 (2018) The model of circulating immune complexes and interleukin-6 improves the prediction
900 of disease activity in systemic lupus erythematosus. Sci Rep 8**:** 2620
901

902 Urbaczek AC, Toller-Kawahisa JE, Fonseca LM, Costa PI, Faria CM, Azzolini AE, Lucisano-
903 Valim YM, Marzocchi-Machado CM (2014) Influence of FcgammaRIIIb polymorphism on its
904 ability to cooperate with FcgammaRIIa and CR3 in mediating the oxidative burst of human
905 neutrophils. Hum Immunol 75**:** 785-790
906

907 Van den Hoecke S, Ehrhardt K, Kolpe A, El Bakkouri K, Deng L, Grootaert H, Schoonooghe
908 S, Smet A, Bentahir M, Roose K et al (2017) Hierarchical and Redundant Roles of Activating
909 FcgammaRs in Protection against Influenza Disease by M2e-Specific IgG1 and IgG2a
910 Antibodies. J Virol 91
911

912 van Egmond M, Vidarsson G, Bakema JE (2015) Cross-talk between pathogen recognizing
913 Toll-like receptors and immunoglobulin Fc receptors in immunity. Immunol Rev 268**:** 311-327
914

915 Vanderbruggen T, Kok PTM, Raaijmakers JAM, Lammers JWJ, Koenderman L (1994)
916 Cooperation between Fc-Gamma Receptor-Ii and Complement Receptor-Type-3 during
917 Activation of Platelet-Activating-Factor Release by Cytokine-Primed Human Eosinophils.
918 Journal of Immunology 153**:** 2729-2735
919

920 Veen JCCMI, Grootendorst DC, Bel EH, Smits HH, van der Keur M, Sterk PJ, Hiemstra PS
921 (1998) CD11b and L-selectin expression on eosinophils and neutrophils in blood and induced
922 sputum of patients with asthma compared with normal subjects. Clinical and Experimental
923 Allergy 28**:** 606-615
924

925 Vidarsson G, Dekkers G, Rispens T (2014) IgG subclasses and allotypes: from structure to
926 effector functions. Front Immunol 5**:** 520
927

928 Vossebeld PJ, Homburg CH, Roos D, Verhoeven AJ (1997) The anti-Fc gamma RIII mAb 3G8
929 induces neutrophil activation via a cooperative actin of Fc gamma RIIIb and Fc gamma RIIa.
930 Int J Biochem Cell Biol 29**:** 465-473
931

932 Vuckovic F, Kristic J, Gudelj I, Teruel M, Keser T, Pezer M, Pucic-Bakovic M, Stambuk J,
933 Trbojevic-Akmacic I, Barrios C et al (2015) Association of systemic lupus erythematosus with
934 decreased immunosuppressive potential of the IgG glycome. Arthritis Rheumatol 67**:** 2978-
935 2989
936
937 Vuitton DA, Vuitton L, Seilles E, Galanaud P (2020) A plea for the pathogenic role of immune
938 complexes in severe Covid-19. Clin Immunol 217**:** 108493
939
940 Wang TT, Ravetch JV (2015) Immune complexes: not just an innocent bystander in chronic
941 viral infection. Immunity 42**:** 213-215
942
943 Ward PA, Fattahi F, Bosmann M (2016) New Insights into Molecular Mechanisms of Immune
944 Complex-Induced Injury in Lung. Front Immunol 7**:** 86
945
946 Yamada DH, Elsaesser H, Lux A, Timmerman JM, Morrison SL, de la Torre JC, Nimmerjahn
947 F, Brooks DG (2015) Suppression of Fcgamma-receptor-mediated antibody effector function
948 during persistent viral infection. Immunity 42**:** 379-390
949
950 Zarbock A, Ley K (2009) Neutrophil adhesion and activation under flow. Microcirculation 16**:**
951 31-42
952
953 Zhao S, Grieshaber-Bouyer R, Rao DA, Kolb P, Chen H, Andreeva I, Tretter T, Lorenz HM,
954 Watzl C, Wabnitz G et al (2021) JAK inhibition prevents the induction of pro-inflammatory HLA-
955 DR(+) CD90(+) RA synovial fibroblasts by IFN. Arthritis Rheumatol
956
957 Zubler RH, Nydegger U, Perrin LH, Fehr K, McCormick J, Lambert PH, Miescher PA (1976)
958 Circulating and intra-articular immune complexes in patients with rheumatoid arthritis.
959 Correlation of 125I-Clq binding activity with clinical and biological features of the disease. J
960 Clin Invest 57**:** 1308-1319
961
962
963

Acknowledgements: We thank T. Schleyer (IR-B Biobank) for providing patient samples. We

are indebted to Falk Nimmerjahn (Universität Erlangen-Nürnberg) for providing KRNtg mice.

Funding: This work was supported by an intramural junior investigator fund of the Faculty of

Medicine to PK (EQUIP - Funding for Medical Scientists, Faculty of Medicine, University of

Freiburg), by the German Research foundation (DFG) (FOR2830 HE 2526/9-1) to HH, by the

DFG research grant TRR130 to REV and the Ministry of Science, Research, and Arts Baden-

Württemberg (Margarete von Wrangell Programm) to NC. **Competing interests:** The authors

declare no financial and non-financial competing interests. **Data and materials availability:**

All data associated with this study are present in the paper or Supplementary Materials.

Author contribution

Conceived and designed the experiments: P.K., H.C., A.M-P., M.H.

Performed the experiments: P.K., H.C., M.H., A.M-P, U.S.

Analyzed the data: P.K., H.C., A.M-P, U.S.

Contributed/reagent/sample material: RE.V.

Writing and original draft preparation: P.K., H.C.

Review and editing: H.H., RE.V., P.K.

Figure 2

Figure 3

Figure 4

Figure 5

999   **Figure 6**

1000

1001

1002

1003

1004 **Supplementary material**

1005

1006 **Figure S1**

1007

1008

1009

1010 **Fig. S1. rhTNFα is not toxic to mouse lymphocyte BW5147 cells even at high**

1011 **concentrations**. Cell count and percentage of live cells were unaltered over a 16 h time frame

1012 of reporter cell culture in the presence of indicated rhTNFα concentrations and comparable to

1013 regular growth in complete medium. Experiments were conducted in 3 replicates. Error bars =

1014 SD.

1015

1016

Fig. S2. FcγRs are activated by sICs formed from multivalent antigens. Two different

multivalent ultra-pure antigens (Ag) mixed with respective therapy-grade mAbs were used to

generate sICs as indicated for each set of graphs (top to bottom). IC pairs: mepolizumab (Mpz)

and rhIL-5; bevacizumab (Bvz) and rhVEGFA. X-Axis: concentrations of stimulant expressed

as molarity of either mAb or Ag monomer and IC (expressed as mAb molarity) at a mAb:Ag

ratio of 1:2. Soluble antigen or soluble antibody alone served as negative controls and were not

sufficient to activate human FcγRs. FcγR responses were normalized to immobilized rituximab

(Rtx) at 1 µg/well (set to 1) and a medium control (set to 0). Two independent experiments

performed in technical duplicates. Error bars = SD. Error bars smaller than symbols are not

shown.

1027

1028

Fig. S3. Distinct activation patterns of NK cells incubated with inverse sICs. Negatively selected primary NK cells purified from PBMCs of three healthy donors were tested for NK cell activation markers. Error bars = SD. One-way ANOVA (Tukey); *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. A) NK cells were incubated for 4 h with immobilized FcγRIII-specific mAb, soluble mouse-anti-human IgG F(ab)$_2$ complexed FcγRIII-specific mAb (reverse sICs), immobilized IgG of non-FcγRIII-specificity (isotype control) or soluble F(ab)$_2$ complexed isotype control (all at 1µg, 10^6 cells). Incubation with PMA and Ionomycin served as a positive control. Incubation with medium alone served as a negative control. B) As in A using an FcγRII-specific mAb. NK cells from the tested donors in this study do not react to FcγRII activation.

1039 **Figure S4**

1040

1041

1042

 Fig. S4. AF4 elution profiles of Ifx/TNFα-immune complexes.

The elution profiles from one of three independent runs are shown. Protein concentration in the eluate is shown in black (UV signal at λ = 280 nm, normalized to the highest UV signal found in this experiment), molar masses determined by MALS for a given retention time in red. Horizontal red lines indicate the range of molar masses used to calculate the mass-weighted mean of molar masses <M_w>. A) Overlay of the elution profiles obtained for TNFα and Ifx, respectively; B1 to B5) Elution profiles for sICs formed after incubation of TNFα and Ifx at different molar ratios.

Table S1

Sample	Range of assigned molar masses [kDa]			Mass-weighted mean of assigned molar masses [kDa]			
	Run 1	Run 2	Run 3	Run 1	Run 2	Run 3	Mean ± SD
Infliximab, IFX	158 – 182	153 – 164	159 – 193	162	156	163	160 ± 4
TNF -alpha	52 – 55	51 – 61	52 – 62	52	52	52	52 ± 0
Immune complexes							
IFX/TNF 3:1	182 – 1.16·10³	164 – 1.11·10³	193 – 1.10·10³	409	398	518	442 ± 66
IFX/TNF 1:1	182 – 2.06·10³	164 – 1.31·10³	193 – 1.42·10³	801	589	681	690 ± 106
IFX/TNF 1:3	182 – 5.05·10³	164 – 6.89·10³	193 – 10.8·10³	1.77·10³	2.00·10³	2.61·10³	2.13·10³ ± 435
IFX/TNF 1:9	182 – 5.36·10³	164 – 3.38·10³	193 – 3.51·10³	1.66·10³	1.18·10³	1.17·10³	1.34·10³ ± 279
IFX/TNF 1:27	182 – 1.68·10³	164 – 768	193 – 1.01·10³	689	464	521	558 ± 117

Table S1. Analysis of the molar mass distribution of ICs from AF4 data.

For a given elution time, the AF4 profiles provide the concentration (UV) at which a given molar mass (MALS) of a protein is present in the sample. The molar mass distribution of Ifx, TNFα and their immune complexes (sICs) was obtained by plotting the cumulative frequency as a function of molar mass. For a selected range of molar masses, a mass-weighted mean value (<M_w>) was calculated. All detected molar masses were selected in the case of Ifx and TNFα whereas only molar masses larger than the maximal molar mass found for Ifx were assigned to sICs. The table shows the range of assigned molar masses and the calculated <M_w> for each AF4 run (n = 3).