Chromosome-length genome assembly and structural variations of the primal Basenji dog (*Canis lupus familiaris*) genome

Richard J. Edwards[1†] , Matt A. Field[2,3†], James M. Ferguson[4†], Olga Dudchenko[5,6,7†], Jens

Keilwagen[8], Benjamin D. Rosen[9], Gary S. Johnson[10], Edward S. Rice[11], LaDeanna Hillier[12],

Jillian M. Hammond[4], Samuel G. Towarnicki[1], Arina Omer[5,6], Ruqayya Khan[5,6], Ksenia

Skvortsova[13,14], Ozren Bogdanovic[1,13], Robert A. Zammit[15], Erez Lieberman Aiden[5,6,7,16,17*],

Wesley C. Warren[10*], J. William O. Ballard[1,18*]

[†] Richard Edwards, Matt Field, James Ferguson and Olga Dudchenko contributed equally to

this work.

* Correspondence: jwoballard@gmail.com, warrenwc@missouri.edu erez@erez.com

1   School of Biotechnology and Biomolecular Sciences, University of New South Wales,

        Sydney NSW 2052, Australia. Richard.edwards@unsw.edu.au,

        o.bogdanovic@garvan.org.au, s.towanicki@unsw.edu.au, jwoballard@gmail.com

2   Centre for Tropical Bioinformatics and Molecular Biology, Australian Institute of

        Tropical Health and Medicine, James Cook University, Cairns, QLD 4878, Australia.

        matt.field@jcu.edu.au

3   John Curtin School of Medical Research, Australian National University, Canberra,

        ACT 2600, Australia. matt.field@jcu.edu.au

4   Kinghorn Center for Clinical Genomics, Garvan Institute of Medical Research, Victoria

        Street, Darlinghurst, NSW 2010, Australia. j.ferguson@garvan.org.au,

        j.hammond@garvan.org.au

25   5   The Center for Genome Architecture, Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA. Olga.Dudchenko@bcm.edu, Arina.Omer@bcm.edu, Ruqayya.Khan@bcm.edu, Zane.Colaric@bcm.edu, erez@erez.com

    6   Department of Computer Science, Rice University, Houston, TX, USA.

30   Olga.Dudchenko@bcm.edu, Arina.Omer@bcm.edu, Ruqayya.Khan@bcm.edu, Zane.Colaric@bcm.edu, erez@erez.com

    7   Center for Theoretical and Biological Physics, Rice University, Houston, TX, USA. Olga.Dudchenko@bcm.edu, erez@erez.com

    8   Julius Kühn-Institut, Erwin-Baur-Str. 27 06484 Quedlinburg, Germany.

35   jens.keilwagen@julius-kuehn.de

    9   Animal Genomics and Improvement Laboratory, Agricultural Research Service USDA, Beltsville, MD 20705. ben.rosen@usda.gov

    10   College of Agriculture, Food & Natural Resources, University of Missouri, MO 65211 USA johnsongs@missouri.edu, warrenwc@missouri.edu

40   11   Bond Life Sciences Center, University of Missouri, Columbia, MO 65211. Edward.rice@missouri.edu

    12   Genome Sciences, University of Washington, WA 98195 USA. lhillier@uw.edu

    13   Genomics and Epigenetics Division, Garvan Institute of Medical Research, Victoria Street, Darlinghurst, NSW 2010, Australia. k.skvortsova@garvan.org.au,

45   o.bogdanovic@garvan.org.au

    14   St Vincent's Clinical School, Faculty of Medicine, University of New South Wales, Sydney, NSW, 2010, Australia, k.skvortsova@garvan.org.au

    15   Vineyard Veterinary Hospital, 703 Windsor Rd, Vineyard, NSW, 2765. Robert@vineyardvet.com.au

50   16   Faculty of Science, UWA School of Agriculture and Environment, University of

        Western Australia, Perth WA 6009, Australia. erez@erez.com

        17   Shanghai Institute for Advanced Immunochemical Studies, ShanghaiTech University,

        Shanghai, China. erez@erez.com

        18   School of Biosciences, University of Melbourne, Royal Parade, Parkville, Victoria

55   3052, Australia. jwoballard@gmail.com

Abstract

Background: Basenjis are considered an ancient dog breed of central African origins that still live and hunt with tribesmen in the African Congo. Nicknamed the barkless dog, Basenjis possess unique phylogeny, geographical origins and traits, making their genome structure of great interest. The increasing number of available canid reference genomes allows us to examine the impact the choice of reference genome makes with regard to reference genome quality and breed relatedness.

Results: Here, we report two high quality *de novo* Basenji genome assemblies: a female, China (CanFam_Bas), and a male, Wags. We conduct pairwise comparisons and report structural variations between assembled genomes of three dog breeds: Basenji (CanFam_Bas), Boxer (CanFam3.1) and German Shepherd Dog (GSD) (CanFam_GSD). CanFam_Bas is superior to CanFam3.1 in terms of genome contiguity and comparable overall to the high quality CanFam_GSD assembly. By aligning short read data from 58 representative dog breeds to three reference genomes, we demonstrate how the choice of reference genome significantly impacts both read mapping and variant detection.

Conclusions: The growing number of high-quality canid reference genomes means the choice of reference genome is an increasingly critical decision in subsequent canid variant analyses. The basal position of the Basenji makes it suitable for variant analysis for targeted applications of specific dog breeds. However, we believe more comprehensive analyses across the entire family of canids is more suited to a pangenome approach. Collectively this work highlights the importance the choice of reference genome makes in all variation studies.

Keywords: canine genome, domestication, comparative genomics, artificial selection

Background

Dogs were the first animals to be domesticated by humans some 30,000 years ago [1] and exhibit exceptional levels of breed variation as a result of extensive artificial trait selection [2]. It is not clear whether dogs were domesticated once or several times, though the weight of accumulating evidence suggests multiple events [3-9]. By establishing genome resources for more ancient breeds of dog, we can explore genetic adaptations perhaps unique to the modern dog breeds. Basenjis are an ancient breed that sits at the base of the currently accepted dog phylogeny [10]. Basenji-like dogs are depicted in drawings and models dating back to the Twelfth Dynasty of Egypt [11] and they share many unique traits with pariah dog types. Like dingoes and New Guinea Singing dogs (NGSD), Basenjis come into oestrus annually—as compared to most other dog breeds, which have two or more breeding seasons every year. Basenjis, dingoes and NGSDs are prone to howls, yodels, and other vocalizations over the characteristic bark of modern dog breeds. One explanation for the unusual vocalisation of the Basenji is that the larynx is flattened [12]. The shape of the dingo and NGSD larynx is not reported.

Basenjis were originally indigenous to central Africa, wherever there was tropical forest. Primarily, what is now the DRC Congo, Southern Sudan, Central African Republic and the small countries on the central Atlantic coast. Today their territory has shrunk to the more remote parts of central Africa. The Basenji probably made its debut in the western world in around 1843. In a painting of three dogs belonging to Queen Victoria and Prince Albert entitled "Esquimaux, Niger and Neptune", Niger is clearly a Basenji. In total, 71 Basenjis have been exported from Africa and, to date, ~ 56 have been incorporated into the registered Basenji breeding population.

The first dog genome to be sequenced was of Tasha the Boxer [13], which was a tremendous advance and continues to be the resource guiding canine genomics research

today. The Boxer is a highly derived brachycephalic breed that has been subjected to generations of artificial selection. Further, due to its discontiguous sequence representation it has been difficult to accurately detect structural variations (SVs) in other domestic dog

110   breeds. Now, a new generation of breed-specific chromosome-level genome reference assemblies are becoming available (5 breeds in October 2020 according to the NCBI assembly archive). For example, we previously published a chromosome-level German Shepherd dog (GSD) genome assembly (CanFam_GSD) that is comprised of only 410 scaffolds and 716 contigs [14].

115   Here, we first report the sequence and *de novo* assembly of two Basenji genomes, female and male. We then compare these assemblies with the Boxer (CanFam3.1) [15] and GSD (CanFam_GSD) [14]. We conduct pairwise comparisons and report single-nucleotide variants (SNVs) and SVs between Basenji, Boxer and GSD. We distinguish an SNV as a variation in a single nucleotide without any limitations on its frequency. SV comprises a

120   major portion of genetic diversity and its biological impact is highly variable. Chen et al. [16] used high-resolution array comparative genome hybridization to create the first map of DNA copy number variation (CNV) in dogs. Many canine CNVs were shown to effect protein coding genes, including disease and candidate disease genes, and are thus likely to be functional. In this study, we find all types of genetic variation are impacted by the choice of

125   reference genome. The basal position of the Basenji makes it useful as a general reference for variant analysis, but the generation of clade-specific genomes is likely to be important for canine nutrition and disease studies. We recommend a pan-genome approach for comprehensive analyses of canid variation.

Results

Basenji female assembly, CanFam_Bas

The female Basenji, China (Fig. 1A), was initially assembled from 84.5 Gb Oxford Nanopore

Technologies (ONT) PromethION reads (approx. 35x depth based on a 2.41 Gb genome size)

using Flye (v2.6) [17, 18] and subjected to long read polishing with Racon v1.3.3 [19] and

Medaka 1.03 [20] (Supplementary Fig 1A, Additional File 1). Additional short read Pilon

[21] error-correction was performed with 115.1 Gb (approx. 47.7x) BGIseq data. Hi-C

proximity ligation was used with the DNA zoo pipeline [22-24] to scaffold 1,657 contigs into

1,456 scaffolds, increasing the N50 from 26.3 Mb to 63.1 Mb and decreasing the L50 from

33 to 14 (Figs 2 and 3, Supplementary Table 1, Additional File 2). Scaffolds were gap-filled

by PBJelly (pbsuite v.15.8.24) [25] using the ONT data, reducing the number of gaps from

348 to 148 and the number of scaffolds to 1,407. Following a final round of Pilon [21]

BGIseq-polishing, scaffolds were mapped onto the CanFam3.1 [13] using PAFScaff v0.40

[14, 26]. Diploidocus v0.9.0 vector filtering [27] removed one 5.7kb contig and masked a

3.3kb region of Chromosome X as lambda phage (J02459.1) contamination. Seven rounds of

iterative Diploidocus tidying of the remaining sequences removed 277 (832 kb) as low

coverage/quality and 481 (1.58 Mb) as probable haplotigs, retaining 483 core scaffolds and

165 probable repeat-heavy sequences [14] as China v1.0 (Fig 3, Supplementary Fig 2,

Additional File 1, Supplementary Table 2, Additional File 2).

150   **Figure 1. The Basenji dogs included in the study. A**. China. Is registered as Australian
Kennel Club Supreme Champion Zanzipow Bowies China Girl. Her registration is
#2100018945. She was born in 2016 and she is free of all known genetic diseases. Her sire
and dam are Australian bred and her most recent ancestor from Africa was 18 generations
ago. **B**. Wags. Is registered as American Kennel Club Champion Kibushi the Oracle, born in
155   in 2008. His registration number is HP345321/01. His sire is an American bred dog while his
dam was imported from the Haut-Ule district of the DRC Congo, 3°24'04.0"N 27°19'04.6"E,
in 2006 .

160   **Figure 2. Contact matrices generated by aligning the CanFam_Bas (China) Hi-C data
set to the genome assembly A**. before the Hi-C upgrade (draft assembly). **B.** After Hi-C
scaffolding (End-to-end assembly).

Figure 3. Key contiguity, quality and completeness metrics for different assembly stages and comparison dog genomes. Square, pre-scaffolded China; Diamond, scaffolded China; Triangle, complete assembly; Circle, main chromosome scaffolds only; Blue, China; Purple, Wags; Red, CanFam_GSD; Green, CanFam3.1. **A.** Contig (open) and scaffold (filled) numbers. **B.** Contig (open) and scaffold (filled) N50. **C.** Contig (open) and scaffold (filled) L50. **D.** Genome completeness estimated by BUSCO v3 (filled) and Merqury (open). **E.** The percentage of missing BUSCO genes (filled) and BUSCOMP genes (those found to be Complete in any assembly) (open). **F.** Schematic of China assembly workflow. CanFam_Bas is China v1.2.

175 *Genome assembly correction*. Two pairs of fused chromosomes in China v1.0 were

incorrectly joined by PBJelly. Pre-gap-filled HiC scaffolds were mapped onto the assembly

using Minimap2 v2.17 [28] and parsed with GABLAM v2.30.5 [29] to identify the gap

corresponding to the fusion regions. These were manually separated into four individual

chromosomes, gap lengths standardised, and scaffolds re-mapped onto CanFam3.1 using

180 PAFScaff v0.4.0. D-GENIES [30] analysis against CanFam_GSD chromosomes confirmed

that PBJelly had incorrectly fused two pairs of chromosomes: chromosomes 8 with 13, and

chromosome 18 with 30. These were manually separated and the assembly re-mapped onto

CanFam3.1 as China v1.1. PAFScaff assigned 112 scaffolds to chromosomes, including 39

nuclear chromosome-length scaffolds.

185   It was observed that the mitochondrial chromosome was missing and China v1.1

Chromosome 29 contained a 33.2 kb region consisting of almost two complete copies of the

mitochondrial genome that were not found in other dog genome assemblies. The 26 ONT

reads that mapped onto both flanking regions were reassembled with Flye v.2.7.1 [17, 18]

into a 77.2 kb chromosome fragment, which was polished with Racon v1.3.3 [19] and

190 Medaka 1.03 [20]. This was mapped back on to the Chromosome 29 scaffold with GABLAM

v2.30.5 [29] (blast+ v2.9.0 megablast [31, 32]) and the mitochondrial artefact replaced with

the corrected nuclear mitochondrial DNA (NUMT) sequence. Finally, scaffolds under 1 kb

were removed to produce the China v1.2 nuclear genome that we name CanFam_Bas.

195 *Mitochondrial genome assembly.* In total, 4,740 ONT reads (52.1 Mb) mapping on to

mtDNA were extracted. To avoid NUMT contaminants, a subset of 80 reads (1.32 Mb) with

99%+ mtDNA assignment and 99%+ mtDNA coverage, ranging in size from 16,197 kb to

17,567 kb, were assembled with Flye 2.7b-b1526 [17, 18] (genome size 16.7 kb) into a

33,349 bp circular contig consisting of two mtDNA copies. This contig was polished with

200   Racon [19] and Medaka [20], before being re-circularised to a single-copy matching the

CanFam3.1 mtDNA start position. After final Pilon [21] correction of SNPs and indels, the

16,761 bp mitochondrial genome was added to the CanFam_Bas assembly.

CanFam_Bas (China) reference genome. The resulting chromosome-length CanFam_Bas

205   reference genome is 2,345,002,994 bp on 632 scaffolds with 149 gaps (76,431 bp gaps)

(Table 1). The 39 nuclear plus mitochondrial chromosome scaffolds account for 99.3% of the

assembly and show a high level of synteny with CanFam3.1 and CanFam_GSD (Fig 4).

CanFam_Bas represents the most contiguous dog chromosomes to date, with a contig N50 of

37.8 Mb and contig L50 of 23, which is slight improvement over CanFam_GSD and

210   considerably more contiguous than the standard dog reference genome, CanFam3.1 (Fig 3,

Table 1). The completeness and accuracy of the genome as measured by BUSCO v3 [33]

(laurasiatherian, *n*=6253) is also superior to CanFam3.1 and approaches that of

CanFam_GSD (92.9% Complete, 3.75% Fragmented, 3.34% Missing).

215   **Methylomic identification of putative regulatory elements**

Additionally, we profiled whole genome methylation of Basenji's blood DNA using MethylC-

seq [34]. Numbers of unmethylated and highly methylated CpG sites in Basenji's genome were

similar to that of GSD (Supplementary Fig. 3A, Additional File 1). Importantly, high resolution

DNA methylation data can be utilised to identify the putative regulatory elements in a given

220   tissue type. That is, CpG-rich unmethylated regions (UMRs) mostly correspond to gene

promoters, while CpG-poor low-methylated regions (LMRs) correspond to distal regulatory

elements, both characterised by DNAse I hypersensitivity [35]. Using MethylSeekR algorithm

[36], we performed the segmentation of Basenji DNA methylome and discovered 20,660

UMRs and 54,807 LMRs (Supplementary Fig. 3B,C, Additional File 1), in line with previously

reported numbers in mammalian genomes [14, 36, 37]. Genome-wide and locus-specific CpG

methylation called by MethylC-seq correlated strongly with that called directly from the ONT

data (Supplementary Fig. 3D-F, Additional File 1), confirming the robustness of determined

DNA methylation profile of the blood DNA.

Table 1. Genome assembly and annotation statistics for Basenji assemblies vs CanFam3.1 and CanFam_GSD.

	CanFam_Bas (China)	Wags	CanFam3.1	CanFam_GSD
Total sequence length	2,345,002,994	2,410,429,933	2,410,976,875	2,407,308,279
Total ungapped length	2,344,926,563	2,410,291,233	2,392,715,236	2,401,163,822
Number of scaffolds	632	2,243	3,310	430
Scaffold N50	64,291,023	61,087,166	45,876,610	64,346,267
Scaffold L50	14	16	20	15
Number of contigs	780	3,630	27,106	736
Contig N50	37,759,230	3,131,423	267,478	20,914,347
Contig L50	23	217	2,436	37
No. chromosomes	40	40	40	40
Percentage genome in main chromosomes	99.3%	94.8%	98.3%	96.5%
BUSCO complete (genome)	92.9% (1.14% Duplicated)	91.5% (1.31% Duplicated)	92.2% (1.17% Duplicated)	93.0% (1.38% Duplicated)
BUSCO fragmented (genome)	3.74%	4.53%	4.03%	3.73%
BUSCO missing (genome)	3.34%	3.98%	3.73%	3.37%
BUSCO complete (proteome)	98.5% (1.9% Duplicated)	97.8% (2.4% Duplicated)	95.1% (1.0% Duplicated)	98.9% (2.4% Duplicated)
BUSCO fragmented (proteome)	1.2%	1.5%	1.9%	1.0%
BUSCO missing (proteome)	0.3%	0.7%	3.0%	0.1%

Figure 4. D-GENIES synteny plots of main chromosome scaffolds for three dog genome assemblies against CanFam_Bas. In each case, CanFam_Bas (China v1.2) is on the x-axis and the comparison assembly on the y-axis. Gridlines demarcate scaffolds. Thick black lines indicate regions of genomic alignment. **A**. All-by-all main chromosome scaffold alignments with **(i)** CanFam_GSD, **(ii)** CanFam_3.1, and **(iii)** Wags. **B**. Main chromosome 9 scaffold alignment with **(i)** CanFam_GSD, **(ii)** CanFam_3.1, and **(iii)** Wags.

Male Basenji assembly, Wags

For the male Basenji, Wags, (Fig 1B) we generated Pacific Bioscience Single Molecule Real

245   Time (SMRT) sequences to approximately 45x genome coverage and assembled the genome

to ungapped size of 2.41 Gb (Supplementary Fig 1B, Additional File 1). Assembly contiguity

metrics of 3,630 total contigs show N50 contig and scaffold lengths of 3.1 and 61 Mb length,

respectively (Table 1). Wags alignment to China revealed a high level of synteny. However,

the Wags assembly of the X chromosome is smaller in size (59 Mb vs 125 Mb) and shows

250   multiple rearrangements as a result of lower sequence coverage on the sex chromosomes

(~21x). We were unable to accurately place 124.4 Mb of Wags sequence on 2,204 scaffolds

(2,210 contigs), including 651 contigs with a total length of 45.6 Mb mapped on to the

CanFam3.1 X chromosome by PAFScaff. Therefore, all comparative analyses reported herein

were done with CanFam_Bas. In addition, the Wags assembly includes 3.6 Mb of the Basenji

255   dog Y for future comparative studies of this unique chromosome.

Table 2. Predicted copy numbers for CanFam_Bas GeMoMa genes based on A. assembly mapping, and B. long read mapping.

A. Dog	Missing	Partial (<50%)	Single (1n)	Duplicate (2n)	3n+
CanFam_Bas	0	0	27129	0	0
	(0)	(0)	(25788)	(0)	(0)
CanFam_Wags	302	120	26103	486	118
	(108)	(58)	(25035)	(472)	(115)
CanFam3.1	211	167	26404	251	96
	(187)	(161)	(25125)	(223)	(92)
CanFam_GSD	95	48	26586	299	101
	(80)	(42)	(25304)	(266)	(96)

B. Data	0n	0.5n	1n	1.5n	2n	2.5n+
CanFam_Bas	2	1257	24116	1508	80	166
(ONT)	(2)	(1201)	(22954)	(1403)	(68)	(160)
CanFam_Wags	7	4717	21140	1028	79	158
(PacBio)	(2)	(3476)	(21049)	(1024)	(79)	(158)
CanFam_GSD	21	1893	22412	2503	109	191
(ONT)	(18)	(1795)	(21350)	(2349)	(98)	(178)

Figures in brackets exclude predicted genes on X chromosome.

Genome annotation

The CanFam_Bas and Wags assemblies were annotated using the homology-based gene prediction program GeMoMa v1.6.2beta [38] and nine reference species [14]. In total, CanFam_Bas and Wags had similar numbers of predicted protein-coding genes at 27,129 (68,251 transcripts) and 27,783 (65,511) transcripts, respectively (Supplementary Table 3, Additional File 2). Analysing the longest protein isoform per gene with BUSCO v3 [33] (laurasiatherian, n=6253, proteins mode), CanFam_Bas was measured to be 98.5 % complete (1.9% duplicated) and Wags was measured as 97.8% complete (2.4% duplicated). Both proteomes compare favourably with CanFam3.1 in terms of completeness (Table 1). To correct for differences introduced by the annotation method, CanFam3.1 was annotated with the same GeMoMa pipeline. Approximately 90% of the Quest For Orthologues (QFO) reference dog proteome [39] is covered by each GeMoMa proteome, confirming comparable levels of completeness (Supplementary Table 3, Additional File 2).

When the CanFam_Bas GeMoMa proteome was compared to Wags, CanFam3.1 and CanFam_GSD, over 91% genes had reciprocal best hits for at least one protein isoform (Supplementary Table 3, Additional File 2). To investigate this further, the Wags, CanFam3.1 and CanFam_GSD genomes were mapped onto CanFam_Bas and the coverage for each gene calculated with Diploidocus v0.10.0. Of the 27,129 predicted genes, 26,354 (97.1%) are found at least 50% covered in all four dogs, whilst only 30 (0.11%) are completely unique to CanFam_Bas. In total, Wags is missing 302 predicted genes, CanFam_GSD is completely missing 95 predicted genes, and CanFam3.1 is missing 211 predicted genes (Table 2). A considerably greater proportion of the missing genes in Wags (64.2% versus 11.4% in CanFam3.1 and 15.8% in CanFam_GSD) were on the X chromosome. To test for artefacts due to assembly errors we mapped the long read data for Wags and CanFam_GSD onto CanFam_Bas. Only 7 of the 302 missing Wags genes (2.3%) had no long read coverage,

285   whilst 21/95 (22.1%) of genes missing in CanFam_GSD were confirmed by an absence of

mapped long reads.

Amylase copy number

Two copies of the *Amy2B* gene were identified in a tandem repeat on Chromosome 6 of the

290   CanFam_Bas assembly. The single-copy read depth for CanFam_Bas, calculated as the

modal read depth across single copy complete genes identified by BUSCO v3 [33], was

estimated to be 34x. This was verified by BUSCO complete genes, which gave mean

predicted copy numbers of 1.008±0.005 (95% C.I.) (Supplementary Fig 4, Additional File 1).

The two complete *Amy2B* coding sequence copies had a mean depth of 97.5x, equating to

295   2.87N, or a total copy number estimate of 5.78N (2 x 97.5 / 34). The full CanFam_GSD

Amy2B repeat region was also found in two complete copies with a mean depth of 98.1x,

estimating 5.77 copies (2 x 98.1 / 34). Similar results were obtained restricting analysis to

reads at least 5 kb (6.01 gene copies; 5.98 region copies) or 10 kb (6.18 gene copies; 6.04

region copies) to minimise repeat-based fluctuations in read depth. In contrast, droplet digital

300   PCR (ddPCR) estimated that the Basenji China had 4.5 copies per individual chromosome.

This slight difference suggests that the raw sequence data slightly overestimated copies or the

ddPCR primers did not capture all the genetic variation.

        Wags assembly has a single copy of the *Amy2B* region, which includes 90% of the

Amy2B coding sequence (data not shown). Single-copy depth analysis of Wags estimates

305   4.97 (90% at 253.8x) and 5.51 (100% at 253.5x) copies of the *AMY2B* coding sequence and

tandem repeat unit, respectively. To estimate copy number in other Basenji dogs, short read

data was downloaded from SRA for 11 Basenjis (Supplementary Table 4, Additional File 2)

and mapped onto CanFam_BAS. AMY2B copy number was estimated as the ratio of mean

read depth per AMY2B gene versus the mean depth for the whole genome. Estimated copy

310   numbers ranged from 3.93 to 6.79, with a mean of 5.25 (Supplementary Table 4, Additional

File 2). The same analysis was performed on China BGI data, yielding an estimate of 4.38

copies, consistent with the ddPCR results.

Nuclear mitochondrial DNA fragments

315   During the assembly of the female Basenji genome (China v1.0), the mitochondrial genome

was erroneously assembled into a NUMT fragment on chromosome 29. A blastn search (e <

10-4 [40]) identified 291 putative NUMT fragments, ranging in size from 34 bp to 6,580 bp,

forming 212 NUMT blocks from 34 bp to 16,577 bp (Supplementary Table 5,6, Additional

File 2). Fragments total 190.5 kb (approx. 11.4 mtDNA copies) and span the entire mtDNA

320   with reasonably even coverage and no obvious bias (Supplementary Fig. 5A, Additional File

1), except for low coverage in a region of the D-loop as has been previously reported in

primates [41]. All 291 NUMT fragments are well-supported with at least 3 reads that span the

entire NUMT plus at least 5 kb each side (Supplementary Table 5, Additional File 2). Only 1

NUMT was not found to be full-length in CanFam_GSD (Supplementary Fig. 5B, Additional

325   File 1). An additional 26 NUMTs are partially covered in CanFam3.1 and 9 are entirely

absent. Whilst this could represent a breed difference, 19 of the 35 additional incomplete

NUMTs in CanFam3.1 (65.5%) are also incomplete in Wags, whilst Wags has a further ten

incomplete NUMTs that are present in CanFam3.1 (Supplementary Table 5, Additional File

2). This is consistent with these regions being generally harder to assemble and/or align.

330   Further analyses of these regions may provide insight into domestic dog genealogies.

Whole genome assembly comparisons

To discover unique large-scale structural differences in assembled genomes of the three

breeds – Basenji, German Shepard and Boxer – we performed pairwise alignments of

335   CanFam_Bas, CanFam3.1 and CanFam_GSD. Overall, genome synteny was maintained and

there were limited large scale genomic rearrangements observed (Fig 4A). There was,

however, a large inversion in CanFam3.1 that was not present in CanFam_Bas or

CanFam_GSD (Fig 4B). To investigate this further, we aligned CanFam_Bas against Wags

(Fig 4A iii). As expected, there was no inversion on Chromosome 9 (Fig 4B iii), suggesting

340   the inversion or perhaps an assembly error, occurs in CanFam3.1.

Long read structural variant detection

To generate a conservative set of high-quality structural variants, the consensus of ONT and

SMRT long read sequences data from both Basenji (China female ONT and Wags male

345   SMRT) and GSD (Nala female ONT and SMRT [14]) was mapped onto Basenji

(CanFam_BAS), Boxer (CanFam3.1) and GSD (CanFam_GSD) reference genomes

(Supplementary Table 7, Additional File 2). One difference in the data sets is that both ONT

and SMRT reads from GSD were sourced from the same individual dog whereas for Basenji

the ONT and SMRT reads are from different individual Basenji dogs. These high-quality SVs

350   were additionally annotated for their overlap to both genes and exons. Of these high-quality

ONT/SMRT consensus SVs, the Basenji long reads overlap 814 CanFam3.1 exons and 568

CanFam_GSD exons while the GSD long reads overlap 825 CanFam3.1 exons and 495

CanFam_Bas exons. In total, these SVs represent 92.19 Mb, 97.21 Mb, and 78.99 Mb of

deleted sequence and 4.11 Mb, 4.09 Mb, and 7.69 Mb of inserted sequence for the

355   CanFam_Bas, CanFam3.1, and CanFam_GSD assemblies respectively.

A. Consensus SVs >100 bp

B. Consensus deletions >100 bp vs CanFam3.1

C. Consensus insertions >100 bp vs CanFam3.1

Figure 5. Consensus structural variant calls for combined ONT and PacBio data. High-quality set of consensus structural variant (SV) calls generated from the intersection of the ONT and PacBio SV calls for each breed versus reference comparison, limited to SVs >100 bp long. **A.** Total numbers of SVs called from Basenji CanFam_Bas reads (red) versus CanFam3.1 and CanFam_GSD, and German Shepherd long reads (green) versus CanFam3.1 and CanFam_Bas. **B.** Numbers and overlap of consensus deletion calls for Basenji reads (blue) and GSD reads (green) versus CanFam3.1. **B.** Numbers and overlap of consensus insertion calls for Basenji reads (blue) and GSD reads (green) versus CanFam3.1.

To reduce the impact of small indels arising from local mis-assembly errors, the high-quality consensus SVs were further reduced to those over 100 bp in length (Fig 5). Overall, we observe similar number of total SV calls from the Basenji long reads relative to CanFam_GSD and the GSD long reads relative to CanFam_BAS. Both breeds had a substantially larger number of consensus SV calls against the CanFam3.1 reference, with Basenji long reads generating more SVs calls than GSD long reads. We next overlapped the CanFam SV calls relative to CanFam3.1 and found 18,063 long read deletion calls overlapped between Basenji and GSD. For Basenji this represented 70.00% of the total

25,260 Basenji deletions while for GSD this represented 73.25% of the total 24,138 GSD

deletions (Fig. 5B). Insertions were fewer in number and degree of overlap, however we still

found 5,146 overlapping insertions between Basenji and GSD long reads representing

33.33% of the total 15,434 Basenji insertions and 36.46% of the total 14,111 GSD insertions

(Fig. 5C). The high degree of overlap in SVs from GSD and Basenji relative to CanFam3.1

represent Boxer-specific SVs or potential issues with the current canid reference assembly.

Short read mapping, SNV / small indel detection

Mapping multiple individuals to a reference genome for variant calling is standard practice in

population genomics and is known to be prone to biases in the reference genome. To

investigate whether differences identified in variant analyses were due to the Basenji being a

basal breed or due to assembly quality difference, short read data from 58 different dog

breeds belonging to sixteen different well-supported clades from Parker et al. [10]

(Supplementary Table 8, Additional File 2) were mapped on to three reference genomes

Basenji (CanFam_BAS), Boxer (CanFam3.1) and GSD (CanFam_GSD). Large-scale

structural differences between breeds would be expected to significantly affect read mapping

efficiencies for closely-related versus distantly-related breeds, whilst missing assembly data

would be expected to result in a systematic reduction in mapping across all breeds. In our

analysis, we observe such systematic and breed specific changes in both the number of

mapped reads and variants detected.

For the systematic changes, overall trends are exhibited in the total percentage of

reads mapped across the three references, with the highest percent of reads mapping to

CanFam_GSD, followed closely by CanFam_BAS and the lowest percent of reads mapping

to CanFam3.1 (Fig 6A). ANOVA shows these differences are significant ($F_{2, 171}=819.53$,

$P<0.0001$). To investigate this result further and test for interactions we focused upon breeds

within each of the monophyletic clades close to or associated with the three reference

400   genomes. Specifically, we included the short-read sequences from the four breeds in the

Asian Spitz clade as this close to the basal Basenji lineage. We also included the six breeds

European Mastif clade containing the Boxer and three breeds within the New World clade

containing the GSD. Overall, the CanFam_Bas and CanFam_GSD performed equally well

while the relative mapping was lowest for CanFam3.1 (Fig 6B). Once again ANOVA shows

405   this result is significant ($F_{8, 30}$=32.01, P<0.0001). Next, we considered the capacity of each

reference genome to detect SNV's and indels. In this case CanFam_BAS detected higher

number of changes than did either CanFam3.1 or CanFam_GSD (Fig 6C and D; ANOVA $F_{2, 171}$=30.71, and $F_{2, 171}$= 12.08, respectively with P<0.0001 for each). In combination these data

attest to the quality of the CanFam_Bas assembly. There is a noticeable depletion of variant

410   calls for the reference breed (Supplementary Fig 6, Additional File 1), but there were no

significant interactions between clades and reference genome (data not shown).

Figure 6. Comparative short read mapping and single nucleotide variant calling for 58
415   **dog breeds versus three reference genomes: CanFam_Bas (Bas), CanFam3.1 (3.1) and**
CanFam_GSD (GSD). A. Relative percentage of reads mapped to each reference calculated
by subtracting the mean percentage of reads for the three reference genomes from the number
of reads for each reference. **B.** Relative percentage of reads mapped to each reference for
three closely related clades determined from Parker et al. [10]. The Basenji is closely related
420   to the Asian Spitz clade. The Boxer is a member of the European Mastiff clade and the GSD
is a member of the New world clade. **C.** The number of SNVs. **D.** The number of small
indels.

Discussion

In this manuscript we present a reference-quality assembly of a female Basenji (China),
which we designate CanFam_Bas. We also present a second Basenji genome, Wags. The
Wags build is of high-quality but not as good as CanFam_Bas, partly because Wags was a
male. In total, 64.2% of the missing genes are on the X chromosome, compared to under 16%
in the other two individuals. Analysis of long read data mapped onto the CanFam_Bas
genome annotation further supports assembly issues as a contributor, with only two
autosomal genes lacking any coverage in Wags SMRT data. Equivalently, two other
autosomal genes also lacked any coverage in China ONT data, suggesting that even high-
quality assemblies can have missing genes in an assembly that may not be biologically lost.

CanFam_Bas is considerably more contiguous and complete than CanFam3.1
(Boxer breed) and slightly more contiguous (Contig N50 37.8 kb vs 20.9 kb) but has slightly
lower completeness (BUSCO completeness 92.9% vs 93.0%) than CanFam_GSD [14] (Table
1, Supplementary Table 1, Additional File 2). The ungapped nuclear chromosomes lengths of
CanFam_Bas, CanFam3.1, and CanFam_GSD and 2.33 Gb, 2.32 Gb, 2.36 Gb and,
respectively. Likely, both assembly quality and breed differences contribute to the 211
missing genes in CanFam3.1. CanFam_GSD had fewer predicted genes missing (95) but a
higher percentage (22.0%) also lacking long read coverage. This uncertainty notwithstanding,
our analysis highlights the need to consider multiple reference genomes in a pan-genomic
approach when a comprehensive analysis is required. These results also highlight the need for
considering missing genes carefully on a case-by-case basis as, even with high quality
genome assemblies such as those compared in this study, an apparent absence may reflect
missing data rather than missing biology.

We noted that the mitochondrial chromosome was missing from the China 1.0 Basenji
assembly. An exhaustive search of the genome detected a 33.2kb region consisting of almost

two complete copies of the mitochondrial genome on Chromosome 29 that was not present in the other dog assemblies analysed. This assembly error occurred at a real 4,032 bp NUMT. A

450   similar false incorporation of complete mtDNA has been previously reported in the little brown bat [42]. In total, 212 NUMT blocks (215.1 kb) in 291 fragments (190.5 kb) were detected across the genome, including a previously detected almost full-length (16,577 bp) NUMT on Chromosome 11 [43]. NUMTs are insertions that are universal in nuclear DNA, but they are not ubiquitous in nuclear genome assemblies [44]. Very often NUMTs are edited

455   out during the genome annotation process [45], but it is becoming increasingly clear that that such organelle insertions must be properly accounted for in nuclear genome assemblies as they have the potential to provide unique insight into population histories and animal well-being [46]. In domestic cats, the nuclear cat genome includes 7.9 kb fragment that is duplicated in tandem 38–76 times [47]. In humans NUMTs are polymorphic at 141 loci.

460   NUMT polymorphisms have been used to estimate the age of insertions in human and other primate lineages [46] while five insertions have been implicated in human disease [44, 48]. Only one Basenji NUMT was incomplete in the GSD genome. To ease future NUMT comparisons between breeds, we have wrapped up the NUMT discovery and merging methods into an open source tool, NUMTFinder [49].

465   Sequence analyses estimated six copies of *Amy2B* in China and five in Wags. Previous studies have shown that Basenjis may have 4-18 copies [6], placing these estimates at the lower end of the range. Short read analysis of 11 other Basenji dogs gave similar copy number estimates of four to seven copies, and 4.38 copies in China. This is consistent with the ddPCR estimate of 4.5 copies, although the uneven coverage of short reads is expected to

470   make this less accurate than the long-read approach. Additional work is needed to establish the source of the differences between ddPCR and read depth estimates. The high variation in *Amy2B* copy number suggests at least three possible evolutionary histories of the gene in

Basenjis. First, *Amy2B* copies may have differentially accumulated in specific lineages since the divergence from other dog breeds. Second, the ancestral founding population of the modern Basenji may have been polymorphic for *Amy2B*. Third, it remains possible that the *Amy2B* region from other breeds have differentially introgressed into the modern breed dog. Basenji-like dogs are depicted in In Egyptian drawings and models dating back to 1550-1900 BC [11]. Possibly, the Basenji is derived from the Abuwtiyuw, which was a lightly built hunting dog with erect ears and a curly tail that died before 2280BCE [50]. Likely, sequencing the region around *Amy2B* in Basenjis with a higher copy numbers will aid resolving the alternate hypotheses. This large genetic diversity in Basenji *Amy2B* copy has important dietary and veterinary implications. Veterinarians frequently recommend a rice-based diet following complex intestinal surgeries and following diarrhoea because it is thought to be bland and easily digestible. However, dogs with low *Amy2B* copy numbers may not be able to digest rice readily as their serum amylase levels are lower [51].

CanFam_Bas and CanFam_GSD represent two high-quality reference genomes from different breeds. The availability of corresponding Basenji and GSD long read data, provided an excellent opportunity to further investigate the contributions of breed and assembly differences, as real SVs will be represented in the raw read data even if assembly errors have introduced false positives or negatives. Possessing ONT and SMRT data for both GSD and Basenji allows us to overlap the disparate SV call sets to generate a conservative list of SV calls relative to CanFam3.1. This analysis identified over 70,000 SVs in CanFam_Bas relative to CanFam3.1 and over 64,000 SVs in GSD relative to CanFam3.1 (Supplementary Table 7, Additional File 2). There is a high degree of overlap in SVs from GSD and Basenji relative to CanFam3.1 (68% of total basenji SV calls and 77% of GSD SV calls), which highlights potential issues with the current canid reference assembly. Further, each consensus set contains several hundred SVs overlapping annotated exons, highlighting

the importance of the selection of appropriate reference genome for analysis of specific genomic regions.

500   Next, we examined the overlap of the consensus calls for SVs over 100 bp of GSD and Basenji relative to CanFam3.1. We find a high degree of agreement for deletions, with 70.1% of GSD calls and 74.75% of Basenji calls overlapping (Fig 5B), compared to insertions where 33.3% of basenji calls and 36.4% of calls overlap (Fig 5C). Basenji and GSD long reads have 25.9% and 19.1% more SVs called when CanFam3.1 is used as the

505   reference, respectively, and the high degree of overlap is consistent with Boxer-specific SVs and/or assembly issues. GSD-specific and Basenji-specific SV calls are also evident.

        To get additional insight into breed differences and the influence of reference genome selection, we mapped short read data from 58 breeds onto CanFam_Bas, CanFam3.1, and CanFam_GSD. If assembly differences are dominated by real differences between breeds

510   then we might expect different breeds to map with different efficiencies onto the three genomes. Differences to due quality, on the other hand, should be reflected across all breeds. With some minor exceptions, short read data from the different breeds consistently mapped better onto CanFam_GSD than CanFam_Bas, which was in turn better than CanFam3.1 (Fig 6A). This is consistent with assembly quality being a dominating factor.

515   After adjusting for read mapping difference, different reference genomes also produce different SNV and small indel densities for the 58 mapped breeds (Supplementary Fig 6, Additional File 1). Using a Basenji reference genome consistently identifies more variants than either GSD or Boxer (Fig 6, Supplementary Fig 6, Additional File 1). This probably reflects the basal position of Basenji in the breed phylogeny [10]. The basal position

520   of the Basenji makes it useful as a reference for variant analysis as there are clear biases affecting related breeds seen for both the GSD and Boxer reference genomes. On the other

hand, CanFam_GSD has slightly higher read mapping levels across breeds, which may provide better total coverage.

Sequencing and assembly efforts are increasingly moving from species reference genomes to breed-specific assemblies, such as those recently published for Great Dane [52], Labrador Retriever [53], and a second GSD [54]. Together, our data suggest that a single high-quality reference should be sufficient for most general analyses, but the generation of breed-specific genomes is likely to be important for canine nutrition and disease studies. The most severe and common ailment in Basenjis is Fancioni Syndrome, in which the renal tubes fail to reabsorb electrolytes and nutrients [55]. Approximately, 10-30% of Basenjis in North America are affected. In 2011, it was shown that Basenji Fanconi Syndrome is caused by a 370bp deletion on canine chromosome 3 [56]. To date no other breeds have been recoded with this same deletion, although other mutations can cause the disease. Likely complex diseases in dogs belonging to different clades may have different underlying causes. For example, it remains unclear whether the same suite of mutations causes hip dysplasia in the GSD (New World clade) and the Labrador (Retriever/ Spaniel clade).

Conclusions

Here, we present two high quality *de novo* Basenji genome assemblies: CanFam_Bas (China, female) and Wags (male). CanFam_Bas offers improved genome contiguity relative to CanFam3.1 and can serve as a representative basal breed in future canid studies. We generate core genomic information for the Basenji that has the potential to inform future studies of population history and aid disease management. We generate high-quality variants (SNVs, small indels, and SVs) relative to CanFam_Bas, CanFam3.1, and CanFam_GSD. We demonstrate the impact that the reference genome makes on both read mapping and variant

detection, illustrating the importance of either selecting the appropriate reference genome or employing a pan-genome approach in future canid studies.

Methods

550 **Sequencing and genome assembly of female Basenji, China**

Sampling. China or Zanzipow Bowies China Girl is an Australian Supreme Champion Kennel Club show dog. Her Australian National Kennel Council registration number is 2100018945. She is bred primarily from Australian lines, with her most recent common ancestor coming from Africa 18 generations previously. She was born on 14 Jan 2016 and is

555 free from all known currently reported diseases.

Sequencing. High molecular weight DNA was extracted from 100 µl of blood using the DNeasy Blood and Tissue kit (Qiagen). For long read (Oxford Nanopore) sequencing, 1 µg of DNA was prepared via the Genomic DNA by Ligation kit (SQK-LSK109) as per the

560 manufacturer's protocol. Prepared DNA (180 ng) was loaded onto a PromethION (FLO-PRO002) flowcell and sequenced with standard parameters. After 48 hours, a nuclease flush was performed and an additional 115 ng was loaded onto the run. Base-calling was performed after sequencing with the GPU-enabled guppy basecaller (v3.0.3) using the PromethION high accuracy flip-flop model with config 'dna_r9.4.1_450bps_hac.cfg'.

565 For short read sequencing whole blood was shipped to BGI, Hong Kong. High molecular weight DNA was extracted, a paired-end library prepared, and the sample run on the BGISEQ-500 sequencing system to generate high quality PE100 data. A total of 767,111,208 clean reads (115.1 Gb) were produced with a lower base call accuracy (Q20) of 95.92%.

Assembly. An overview of the China assembly workflow is given in Supplementary Fig 1A, Additional File 1. The ONT reads were assembled with the Flye (v2.6-release) assembler [17, 18]. The resulting contigs were polished with ONT reads using four rounds of Racon (v1.4.3) [19] followed by Medaka (v0.10.0) [20] to minimise error propagation. BGI-seq reads were

575   aligned to the polished assembly with BWA-mem (v 0.7.17) [57] and Pilon (v1.23) (diploid) [21] was used for further error correction. A second assembly was performed using Canu assembler (v1.8.0) [58] and correcting the sequencing errors using 2 rounds of Arrow polishing [59]. The Flye assembly was considered more contiguous and therefore selected as the core assembly.

Scaffolding. The ONT assembly was scaffolded to chromosome-length by the DNA Zoo following standard DNA zoo methodology (www.dnazoo.org/methods). Briefly, an *in situ* Hi-C library was prepared [24] from a blood sample from the same individual and sequenced to ~30x coverage (assuming 2.4 Gb genome size). The Hi-C data was processed using Juicer

585   [60], and used as input into the 3D-DNA pipeline [61] to produce a candidate chromosome-length genome assembly. We performed additional finishing on the scaffolds using Juicebox Assembly Tools [22]. The matrices are visualized in Juicebox.js, a cloud-based visualization system for Hi-C data [23] and are available for browsing at multiple resolutions on https://www.dnazoo.org/post/basenji-the-african-hunting-dog. After scaffolding, all ONT

590   reads were aligned to the assembly with Minimap2 (v2.16) (-ax map-ont) [28] and used by PBJelly (pbsuite v.15.8.24) [25] to fill gaps. BGI reads were re-mapped with BWA-mem (v 0.7.17) and another round of polishing was performed with Pilon (v1.23) (diploid, SNP and indel correction) [21] .

595   *Final clean-up.* The Pilon-polished genome underwent a final scaffold clean-up to generate a

high-quality core assembly following Field and colleagues' processing of Canfam_GSD [14]. Scaffolds were reoriented and assigned to CanFam3.1 chromosomes [15] using PAFScaff (v0.3.0) [14] (Minimap2 v2.16 mapping) [28]. Diploidocus (v0.9.0) [27] was used to screen contamination, remove low-coverage artefacts and haplotig sequences, and

600   annotate remaining scaffolds with potential issues as described in [14]. ONT reads were mapped onto the assembly using Minimap2 (v2.17) (-ax map-ont --secondary=no) [28] and read depth summaries calculated with BBMap (v38.51) pileup.sh [62]. Any scaffolds with median coverage less than three (e.g., less than 50% of the scaffold covered by at least three reads) were filtered out as low-coverage scaffolds. Single-copy read depth was estimated

605   using the modal read depth of 35X across the 5,5578 single copy complete genes identified by BUSCO (v3.0.2b) [33]. This was used to set low-, mid- and high-depth thresholds at 8x, 25x and 68x for PurgeHaplotigs v20190612 [63] (implementing Perl v5.28.0, BEDTools v2.27.1 [64, 65], R v3.5.3, and SAMTools v1.9 [66]). PurgeHaplotig coverage was adjusted to exclude gap regions and any scaffolds with 80%+ bases in the low/haploid coverage bins

610   and 95%+ of their length mapped by PurgeHaplotigs onto another scaffold were filtered as haplotigs or assembly artefacts. Any other scaffolds with 80%+ low coverage bases were filtered as Low Coverage. Remaining scaffolds were further classified based on read depth profiles. Scaffolds with <20% diploid coverage and 50%+ high coverage were marked as probable collapsed repeats. Scaffolds with "Diploid" depth as the dominant PurgeHaplotigs

615   coverage bin and >50% match to another scaffold were marked as a possible repeat sequences [14].

Genome assembly correction. Main chromosome scaffold integrity was checked using D-GENIES [30] comparisons of different assembly stages with CanFam_GSD chromosomes

620   [14]. Two pairs of fused chromosomes were identified following incorrect joins made by

PBJelly (pbsuite v.15.8.24) [25]. Pre-gap-filled HiC scaffolds were mapped onto the assembly using Minimap2 (v2.17) [28] and parsed with GABLAM (v2.30.5) [29] to identify the gap corresponding to the fusion regions. These were manually separated into four individual chromosomes, gap lengths standardised, and scaffolds re-mapped onto CanFam3.1

625   using PAFScaff (v0.4.0) [26]. Finally, scaffolds under 1 kb were removed following correction of Chromosome 29 (below).

Correction of mitochondrial insertion into Chromosome 29. NUMT analysis identified a 33.2 kb region consisting of almost two complete copies of the mitochondrial genome, not

630   present in other dog genome assemblies. GABLAM (v2.30.5) [29] was used to confirm that this region was also absent from the Canu assembly of China v1.1. ONT reads that mapped onto both flanking regions of the 33.2 kb putative NUMT were extracted and reassembled with Flye (v2.7.1) [17, 18]. The new NUMT was approx. 2.8 kb long. To avoid repeated problems with mitochondrial reads mis-polishing the sequence, a subset of 4.82M ONT reads

635   (72.7 Gb, ~30X) was extracted and mapped onto the assembled region with Minimap (v2.17) [28]. Reads mapping to at least 5 kb of the assembled region including some immediate flanking sequence were extracted (66 reads, 1.50 Mb) and polished with one round of Racon (v1.4.5) [19] (-m 8 -x -6 -g -8 -w 500) and Medaka (v0.7.1) [20] (model r941_prom_high). The polished NUMT region was mapped on to the Chromosome 29 scaffold with GABLAM

640   (v2.30.5) [29] (blast+ v2.9.0 [67] megablast) and stretches of 100% sequence identity identified each side of the NUMT. The mtDNA sequence between these regions of identity was replaced with the re-assembled NUMT sequence.

Mitochondrial genome assembly. To assemble the mitochondrion, ONT reads were mapped

645   onto a construct of three tandem copies of the CanFam3.1 mtDNA with minimap2 (v2.17)

[28] (-ax map-ont --secondary=no). Reads with hits were extracted using SAMTools (v1.9) [66] fasta and mapped onto a double-copy CanFam3.1 mtDNA with GABLAM (v2.30.5) [29] (blast+ v2.9.0 [67] megablast). "Pure complete" mtDNA ONT reads were identified as those with 99%+ of their length mapping to mtDNA and 99%+ coverage of the mtDNA.

650 These reads were assembled with Flye (v2.7b-b1526) [17, 18] (genome size 16.7 kb) and polished with Racon (v1.4.5) [19] (-m 8 -x -6 -g -8 -w 500) followed by Medaka (v0.7.1) [20] (model r941_prom_high). The polished mtDNA assembly was mapped onto CanFam3.1 mtDNA with GABLAM (v2.30.5) [29] (blast+ v2.9.0 [67] megablast) and circularised by extracting a region from the centre of the assembly corresponding to a single full-length copy

655 with the same start and end positions. Final correction of SNPs and indels was performed by adding the mtDNA to the nuclear assembly, mapping BGI reads with BWA (v0.7.17) and polishing the mtDNA with Pilon (v1.23) [21]. The polished mtDNA was then added back to the nuclear genome for the final *China* assembly.

660 *Genome assembly quality assessment.* At each stage of the assembly, summary statistics were calculated with SLiMSuite SeqList (v1.45.0) [68, 69], quality was assessed with Merqury (v20200318) (Meryl v20200313, bedtools v2.27.1 [64, 65], SAMTools v1.9 [66], java v8u45, igv v2.8.0) and completeness assessed with BUSCO (v3.0.2b) [33] (BLAST+ v2.2.31 [67], HMMer v3.2.1 [70], Augustus v3.3.2, EMBOSS v6.6.0, laurasiatherian lineage

665 (n=6253)). To account for fluctuations in BUSCO ratings, presence of complete BUSCO genes across assembly stages was also assessed with BUSCOMP (v0.9.4) [71, 72]. Final assembly scaffold statistics and quality assessment was performed with Diploidocus (v0.10.2) (KAT v2.4.2, perl v5.28.0, BEDtools v2.27.1 [64, 65], SAMTools v1.9 [66], purge_haplotigs v20190612, java v8u231-jre, bbmap v38.51, minimap2 v2.17 [28], BLAST+

670   v2.9.0 [67]). To get a sense of final quality, comparisons were made with the two other dog

genomes published at the time of analysis: CanFam3.1 [15] and CanFam_GSD [14].

DNA methylation calling. China's blood DNA methylome libraries were sequenced on the

Illumina HiSeq X platform (150 bp, PE), generating 336 million read pairs and yielding 14x

675   sequencing coverage. Sequenced reads were trimmed using Trimmomatic [73] and mapped

to the China v1.0 genome reference using WALT [74] with the following parameters: -m 10 -

t 24 -N 10000000 -L 2000. The mappability of the MethylC-seq library was 86%. Duplicate

reads were removed using Picard Tools (v2.3.0). Genotype and methylation bias correction

were performed using MethylDackel with additional parameters: minOppositeDepth 5 --

680   maxVariantFrac 0.5 --OT 10,140,10,140 --OB 10,140,10,140. The numbers of methylated

and unmethylated calls at each CpG site were determined using MethylDackel

(https://github.com/dpryan79/MethylDackel). Bisulphite conversion efficiency was 99.71%,

estimated using unmethylated lambda phage spike-in control. Nanopore reads were aligned to

the generated reference genome using Minimap2 v2.17 [28], and CpG methylation sites were

685   called with f5c v0.6 [75], which is an accelerated version of nanopolish [76]. Methylation

frequency was then collated for each CpG site.

UMR and LMR calling. Segmentation of basenji's blood DNA methylome into CpG-rich

unmethylated regions (UMRs) and CpG-poor low-methylated regions (LMRs) was

690   performed using MethylSeekR [36] (segmentUMRsLMRs(m=meth, meth.cutoff=0.5,

nCpG.cutoff=5, PMDs = NA, num.cores=num.cores, myGenomeSeq=build,

seqLengths=seqlengths(build), nCpG.smoothing = 3, minCover = 5).

Sequencing and genome assembly of male Basenji, Wags

695   Wags DNA was derived from blood of a single male. He is registered as American Kennel

Club Champion Kibushi the Oracle, born on December 3, 2008. His registration number is

HP345321/01. Sire is AM Ch C-Quests Soul Driver, HM827502/02, and his dam is

Avongara Luka, HP345312/01, a native female dog imported from the Haut-Ule district of

the DRC Congo, 3°24'04.0"N 27°19'04.6"E, in 2006. SMRT sequences for Wags (Fig. 1B)

700   were generated on the Pacific Biosciences Sequel instrument (V2 chemistry) to

approximately 45x genome coverage based on a genome size estimate of 2.5 Gb. An

overview of the China assembly workflow is given in Supplementary Fig 1B, Additional File

1. All SMRT sequences were assembled with the HGAP4 algorithm, a Falcon based

assembly pipeline available through the SMRT Link interphase (SMRT Link v5.0.1.9585)

705   [77]. The assembly was then error corrected with the original SMRT sequences using the

Arrow error-correction module [77]. Additional polishing of the assembly for residual indels

was done by aligning 32x coverage of Illumina data and the Pilon algorithm [21].

Chromosomal level scaffolds were generated with the same DNA source using the

Proximo™ Hi-C genome scaffolding software (Phase Genomics Inc) and finalized by

710   alignment to the CanFam3.1 reference.

Locus copy number estimation

Copy numbers for specific assembly regions were calculated using Diploidocus (v0.10.0)

(runmode=regcnv) [27]. For each animal, long reads were mapped onto the assembly with

715   Minimap2 (v2.17) (no secondary mapping) [28] and the modal read depth across single-copy

Complete genes identified by BUSCO v3 [33] (laurasiatherian_db) calculated using

SAMTools (v1.9) [66] mpileup. This set the expected single-copy read depth, X_{SC}. Copy

number for a region, N_{reg} was estimated by dividing the mean read depth across that region, X_{reg}, by X_{SC}. The variance and standard deviation of the estimate was calculated using X_{reg} for

720   all single copy BUSCO genes. For Wags (a male), genes on the X chromosome were excluded from this analysis.

Amylase copy number. The copy number of the beta amylase gene *Amy2B* was calculated using Diploidocus (v0.10.0) (runmode=regcnv) [27] using a modification of the single locus

725   copy number estimation (above) to account for multiple copies of the gene in the assembly. First, the AMY2B protein sequence from CanFam3.1 (UniprotKB: J9PAL7) was used as a query and searched against the genome with Exonerate (v2.2.0) [78] to identify assembled copies of the *Amy2B* gene. A full-length (14.8 kb) *Amy2B* gene repeat was also extracted from the CanFam_GSD assembly and mapped onto the assembly with Minimap2 (v2.17) (-x

730   asm5) [28]. Estimated N_{reg} values were converted into a number of copies by multiplying by the proportion of the query found covered by that region. The total genome *Amy2B* copy number was then calculated as the summed copy number over each hit. To further investigate the robustness of the method and improve the *Amy2B* copy number estimate in CanFam_Bas, analysis was repeated with ONT reads at least 5 kb in length and at least 10 kb in length.

735   These reads should be less susceptible to poor mapping at repeat sequences, but at a cost of reduced coverage. A simpler approach was also applied to short read data for China and eleven Basenji dogs with data available on SRA (Supplementary Table 4, Additional File 2). Reads were mapped onto the China genome with BWA mem [57] and sequencing depth calculated across the annotated AMY2B genes from GeMoMa (Chromosome 6: 46,943,564-

740   46,950,679 and 46,928,705-46,935,822) using samtools v1.11 [66]. The mean sequencing depth per base per AMY2B copy was then compared to the mean sequencing depth across the

genome, calculated with BBMap (v38.51) pileup.sh [62]. AMY2B copy number was estimated using the AMY2B:genome read depth ratio.

We also used droplet digital PCR (ddPCR) to directly quantify the *Amy2B* copy number of China DNA [79]. ddPCR was performed using a QX100 ddPCR system (Bio-rad). Each reaction was performed in a 20µl reaction volume containing 10 µl of 2x ddPCR Supermix (Bio-rad), 1µl of each 20x primer/probe, 1µl of DraI restriction enzyme (New England BioLabs #R0129S), 5µl of DNA template (4ng/µl) and 2µl ddH$_2$O. Primer sequence for Amy2B: forward 5′-CCAAACCTGGACGGACATCT-3′ and reverse 5′-TATCGTTCGCATTCAAGAGCAA-3′ with FAM probe: 6FAM–TTTGAGTGGCGCTGGG-MGBNFQ. Primer sequence for C7orf28b-3: 5′-GGGAAACTCCACAAGCAATCA-3′ and reverse 5′-GAGCCCATGGAGGAAATCATC-3′ with HEX probe HEX-CACCTGCTAAACAGC-MGBNFQ.

Nuclear mitochondrial DNA (NUMT) fragment analysis

To make sure that any contiguous NUMTs were identified as a single region, a double-copy CanFam3.1 mtDNA sequence was constructed and then searched against the CanFam_Bas nuclear genome and compressed to unique hits using GABLAM (v2.30.5) [29] (blast+ v2.9.0 [32] blastn, localunique) with a blastn e-value cut-off of 1e-4 [40]. For comparison with published dog NUMTs [43], NUMT fragments with 8 kb were merged into NUMT blocks using NUMTFinder v0.1.0 [49]. Predicted copy number was calculated for each NUMT fragment in China v1.1 using the method described above. Diploidocus (v0.10.0) was also used to calculate number of reads spanning each entire NUMT fragment plus flanking regions of 0 bp, 100 bp, 1 kb and 5 kb. In addition, assembly coverage for each NUMT fragment was calculated for Wags, CanFam3.1 and CanFam_GSD. Each genome was split

into 1 Mb tiled fragments and mapped onto CanFam_Bas with Minimap2 (v2.17) [28]. Each

BAM file was used for Diploidocus (v0.10.0) regcnv [27] analysis with a single-copy read

depth of 1x. Mitochondrial genome coverage was analysed by extracting all 291 NUMT

fragment regions with SeqSuite (v1.23.3) [69] and mapping them onto the CanFam3.1

770 mtDNA chromosome using GABLAM (v2.30.5) [29] (blast+ v2.10 [32] tblastn).

Genome annotation

Following Field et al. [14] each genome was annotated using the homology-based gene

prediction program GeMoMa [38] (v1.6.2beta) and nine reference organisms. For each

775 reference organism, coding exons of full-length transcript were extracted and translated to

peptides using the GeMoMa module Extractor. These peptides were searched in each Basenji

genome using mmseqs2 [80] (v5877873cbcd50a6d954607fc2df1210f8c2c3a4b). Based on

the results of mmseqs2 and Extractor, transcripts were predicted for each Basenji genome

from each reference organism independently. These nine gene annotation sets were then

780 combined into a final gene annotation using the GeMoMa module GAF. To make a fair

comparison of the influence of genome quality and completeness on annotation, CanFam3.1

was annotated with the same pipeline. Annotation of CanFam_GSD using the same pipeline

was obtained from Field et al. [14].

785 ***Annotation summary and quality assessment.*** Annotation summary statistics and the longest

protein isoform per gene were generated with SAAGA (v0.4.0) [81]. Annotation

completeness was estimated using BUSCO v3 [33] (laurasiatherian, $n=6253$, proteins mode),

run on a reduced annotation consisting of the longest protein per gene. To check for truncated

or fragmented protein predictions, predicted proteins were mapped onto the Quest For

790 Orthologues reference dog proteome [39] with mmseqs2 v [80]. The best protein hit for each

gene was used to calculate a protein length ratio (length of predicted protein / length of reference protein). Percentage coverage of query and hit proteins was calculated with mmseqs2 v [80]. A reciprocal search identified the closest predicted protein for each reference protein. Any reciprocal best hits were marked as predicted orthologues.

Annotation copy number and coverage analysis. Predicted copy number was calculated for every protein-coding gene in CanFaBas using the method described above. In addition, assembly coverage for each CanFam_Bas gene was calculated for Wags, CanFam3.1 and CanFam_GSD. Each genome was split into 1 Mb tiled fragments and mapped onto

800   CanFam_Bas with Minimap2 (v2.17) (-ax asm5 -L) [28]. Each BAM file was used for Diploidocus (v0.10.0) regcnv analysis with a single-copy read depth of 1x. In addition, SMRT reads from Wags and ONT reads from the GSD were mapped onto CanFam_Bas with Minimap2 (v2.17) (--secondary=no -L -ax map-pb or -ax map-ont) [28] and the standard predicted copy number calculation applied. Genes with zero coverage were marked 0n. Other

805   genes were binned according to coverage: for mapped assemblies, coverage was rounded to the nearest integer; for long read mapping, coverage was rounded to the nearest 0.5n. Genes with greater than zero but less than 50% coverage were assigned to 0.5n. Any genes exceeding a rounded coverage of 2n were grouped as "3+".

810   ***Ribosomal RNA prediction.*** For each genome, genes for rRNA were predicted with Barrnap (v0.9) [82] (eukaryotic mode, implementing Perl v5.28.0, HMMer v3.2.1 [70] and BEDTools v2.27.1 [64, 65]).

Whole genome assembly comparisons

815   Whole genome synteny analysis were performed for the main chromosome scaffolds using

the D-GENIES [30] web portal.

Long read structural variant detection

Structural variant calls were generated using a combination of minimap2 (v2.17-r943-dirty)

820   [28], SAMTools (v1.9) [66], and sniffles (v1.0.11) [83]. In total, four sets of long reads from

three samples were analyzed consisting of China the Basenji (Oxford Nanopore), Wags the

Basenji (SMRT) and Nala the German Shepherd (Oxford Nanopore and SMRT [14]). Reads

were mapped against China v1.0, CanFam3.1 and CanFam_GSD. Analysis was restricted to

the main nuclear chromosome scaffolds. Variants for the Basenji and for Nala were annotated

825   with gene model predictions generated using GeMoMa [38] (v1.6.2beta) while CanFam3.1

variants were annotated with Ensembl gene annotations v100 for CanFam3.1 [84]

Short read mapping, SNV / small indel detection

Representative Illumina data was identified from [85]. All Sequence Read Archive (SRA)

830   IDs associated with the DBVDC bioproject on NCBI (SRP144493) were downloaded along

with their metadata and reduced to 126 samples representing the biggest sequencing run (no.

bases) per annotated breed. These 126 samples were used for initial read mapping and variant

calling (Supplementary Table 9, Additional File 2). Following removal of

unknown/mixed/village dog samples, canids other than domestic dogs, and duplicates the

835   remaining breeds were mapped onto those used by Parker et al. [10]. In total, 58 breeds in the

significantly monophyletic clades (greater than 70% bootstrap) designated by Parker et al.

[10] were considered for analyses (Supplementary Table 8, Additional File 2).

SNVs and small indels were called from the Illumina reads of the 58 representative

breeds against three reference genomes (Basenji China v1.0, CanFam3.1, and

840   CanFam_GSD). All Illumina reads were downloaded from the short read archive using the

SRA toolkit (v2.10.4) [86]. All samples were analysed using a modified version of an

existing variant detection pipeline [87]. Briefly, the pipeline employs BWA (v0.7.17-r1188)

for read alignment [88], SAMTools (v1.9) [66] and picard (v2.4.1)

(http://broadinstitute.github.io/picard) for binary alignment map (BAM) file preprocessing,

845   and Genome Analysis ToolKit (v3.6) (GATK) for calling SNVs and small indels [89]. The

workflow follows GATK best-practices using default parameters throughout. Additionally,

SRA reads from each reference genome were aligned using BWA and SAMTools and a

consensus variant list generated using an approach described previously [90]. The consensus

variant lists for each reference genome were utilized in GATK's BaseRecalibrator step as the

850   '-knownSites' argument to serve as the required variant truthset. Variant alignment statistics

were generated using SAMTools flagstat. Joint variant calls were generated for the larger

dataset of 126 samples relative to the three reference genomes and the total number of SNVs

and small indels for individual samples tabulated. Variants for the Basenji and CanFam_GSD

were annotated with gene model predictions generated using GeMoMa (see above) while

855   CanFam variants were annotated with ENSEMBL gene annotations v100 for CanFam3.1.

Read mapping statistics for each sample were calculated using SAMTools to remove

secondary mapping and then summarized with BBTools (v38.51) pileup.sh [62]. Numbers

were then converted into relative values for each reference by averaging the score for each

breed over the three reference genomes and then calculating the difference from the mean.

860   Breeds were considered individually, but mean values for each clade were also calculated.

Three clades are of particular importance as they are closely related, or include, the three

reference genome assemblies. The Asian Spitz clade is considered closely related to the

Basenji. This clade contained the Alaskan Malamute, Shar-Pei, Shiba Inu and Tibetan

Mastiff. The European Mastiff Clade contained the Boxer, Bull Terrier, Cane Corseo, Great

865   Dane, Mastiff and Rhodesian Ridgeback. The New World clade contained the Berger Picard, Chinook and German Shepherd. One-way ANOVA's were employed to detect significant differences between groups. As the same short read samples were examined relative to the three reference genomes statistical significance was set to be $P<0.01$.

Abbreviations

BAM, binary alignment map; bp, base pair; ddPCR, digital droplet PCR; CNV, copy number variation; GATK, Genome Analysis Toolkit; Gb, gigabase; GSD, German Shepherd Dog; kb, kilobase; LMR, low-methylated region; Mb, megabase; NGSD, New Guinea Singing Dog; NUMT, nuclear mitochondrial DNA; ONT, Oxford Nanopore Technologies; QFO, Quest For Orthologues; SMRT, Single Molecule Real Time; SNV, single nucleotide variant; SRA, Sequence Read Archive; SV, structural variant; UMR, unmethylated region.

Declarations

Ethics approval and consent to participate

880   For China, all experimentation was performed under the approval of the University of New South Wales Ethics Committee (ACEC ID: 18/18B) and with the owner's written consent.

Experiments at the University of Missouri were done with approval from the University of Missouri Animal Care and Use Committee (ACUC 8833) and with owner's written consent.

885 **Consent for publication**

Not applicable.

Availability of data and materials

Chromosome-length genome assemblies for both Basenji are available at NCBI

890 (CanFam_Bas (*China*), GCA_013276365.1 (v1.0) and GCA_013276365.2 (v1.2); *Wags*, GCA_004886185.2). The CanFam_Bas mitochondrial genome is available at NCBI Genbank: MW051511. Raw read data is available in the Sequence Read Archive. To review GEO accession GSE159396: Go to

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE159396

895 and enter token ylclqswyvhitlcr into the box. Supplementary data accompanies this paper at http://www.slimsuite.unsw.edu.au/research/basenji/ and Open Science Foundation (https://osf.io/r3jfm/). Additional datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

900 **Competing interests**

JF has received travel and accommodation expenses to speak at Oxford Nanopore Technologies conferences. Otherwise, the authors declare that the submitted work was carried out in the absence of any professional or financial relationships that could potentially be construed as a conflict of interest.

Funding

This work was supported by the University of New South Wales/ School of Biotechnology and Biomolecular Sciences Genomics Initiative and the Basenji Health Endowment Inc, Poynette, WI and. The DNA Zoo initiative funded the Hi-C data collection and analyses. RJE

910   is funded by ARC LP160100610 and ARC LP180100721. MF is funded by NHMRC APP5121190. ELA was supported by an NSF Physics Frontiers Center Award (PHY1427654), the Welch Foundation (Q-1866), a USDA Agriculture and Food Research Initiative Grant (2017-05741), and an NIH Encyclopedia of DNA Elements Mapping Center Award (UM1HG009375).

Authors' contributions

JWOB coordinated the project. RJE, MAF, and JWOB designed the study. JWOB and WW funded the project. For China, RAZ provided the sample. JF and BDR performed the initial assembly and polishing. AO performed the Hi-C experiment, and OD, RK and ELA

920   conducted the Hi-C analyses. RJE performed the final assembly polishing, curation, correction, analyses of genome completeness, mitochondrial genome assembly, NUMT analysis, and read-depth gene copy number analyses. SGT completed the digital droplet PCR. KS and OB conducted the DNA methylation analyses. MAF performed the structural comparisons, dog breed short read mapping and variant calling. MAF, RJE and JWOB

925   performed the dog breed variant analysis. GSJ provided Wags samples. ER, LH, and WW performed the Wags assembly. JK performed the genome annotation. RJE, JF, MAF, BDR, OD, OB and JWOB wrote the manuscript. All authors edited and approved the final manuscript.

Acknowledgements

We would like to thank Jennifer Power for introducing us to China and Barbara Reisinger, Jon Curby and Jared Reisinger for access to Wags. For China, Brooke Morgan-Burke at the Vineyard Veterinary Hospital provided constant encouragement. ONT data were collected at the Garvan Institute and the nuclease flush by Jillian Hammond, the BGI -seq data was provided by BGI and the Hi-C data collected at Baylor College of Medicine. For Wags, DNA sequencing was completed at the The Hudson Alpha Institute and assembly at the The McDonnell Genome Institute, Washington University School of Medicine.

References

1. Botigue LR, Song S, Scheu A, Gopalan S, Pendleton AL, Oetjens M, Taravella AM, Seregely T, Zeeb-Lanz A, Arbogast RM *et al*: **Ancient European dog genomes reveal continuity since the Early Neolithic**. *Nat Commun* 2017, **8**:16082.
2. Drake AG, Klingenberg CP: **Large-scale diversification of skull shape in domestic dogs: disparity and modularity**. *Am Nat* 2010, **175**(3):289-301.
3. Thalmann O, Shapiro B, Cui P, Schuenemann VJ, Sawyer SK, Greenfield DL, Germonpre MB, Sablin MV, Lopez-Giraldez F, Domingo-Roura X *et al*: **Complete mitochondrial genomes of ancient canids suggest a European origin of domestic dogs**. *Science* 2013, **342**(6160):871-874.
4. Serres-Armero A, Povolotskaya IS, Quilez J, Ramirez O, Santpere G, Kuderna LFK, Hernandez-Rodriguez J, Fernandez-Callejo M, Gomez-Sanchez D, Freedman AH *et al*: **Similar genomic proportions of copy number variation within gray wolves and modern dog breeds inferred from whole genome sequencing**. *BMC Genomics* 2017, **18**(1):977.
5. Ostrander EA, Wayne RK, Freedman AH, Davis BW: **Demographic history, selection and functional diversity of the canine genome**. *Nat Rev Genet* 2017, **18**(12):705-720.
6. Freedman AH, Gronau I, Schweizer RM, Ortega-Del Vecchyo D, Han E, Silva PM, Galaverni M, Fan Z, Marx P, Lorente-Galdos B *et al*: **Genome sequencing highlights the dynamic early history of dogs**. *PLoS Genet* 2014, **10**(1):e1004016.
7. Zhang S-j, Wang G-D, Ma P, Zhang L-l, Yin T-T, Liu Y-h, Otecko NO, Wang M, Ma. Y-p, Wang L *et al*: **Genomic regions under selection in the feralization of the dingoes** *Nat Comm* 2020, **11**:671.
8. Wang GD, Zhai W, Yang HC, Wang L, Zhong L, Liu YH, Fan RX, Yin TT, Zhu CL, Poyarkov AD *et al*: **Out of southern East Asia: the natural history of domestic dogs across the world**. *Cell Res* 2016, **26**(1):21-33.

9. Gopalakrishnan S, Sinding MS, Ramos-Madrigal J, Niemann J, Samaniego Castruita JA, Vieira FG, Caroe C, Montero MM, Kuderna L, Serres A *et al*: **Interspecific gene flow shaped the evolution of the genus *Canis***. *Curr Biol* 2018, **28**(21):3441-3449 e3445.

10. Parker HG, Dreger DL, Rimbault M, Davis BW, Mullen AB, Carpintero-Ramirez G, Ostrander EA: **Genomic analyses reveal the influence of geographic origin, migration, and hybridization on modern dog breed development**. *Cell Rep* 2017, **19**(4):697-708.

11. Dollman G: **The Basenji dog**. *J R Africa Soc* 1937, **36**:148-149.

12. Ashdown RR, Lea T: **The larynx of the Basenji dog**. *J Small Anim Pract* 1979, **20**(11):675-679.

13. Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, Clamp M, Chang JL, Kulbokas EJ, 3rd, Zody MC *et al*: **Genome sequence, comparative analysis and haplotype structure of the domestic dog**. *Nature* 2005, **438**(7069):803-819.

14. Field MA, Rosen BD, Dudchenko O, Chan EKF, Minoche AE, Barton K, Lyons RJ, Tuipulotu DE, Edwards RJ, Hayes VM *et al*: **Canfam_GSD: *De novo* chromosome-length genome assembly of the German Shepherd Dog (*Canis lupus familiaris*) using a combination of long reads, optical mapping and Hi-C**. *GiGaScience* 2020, **accepted 14/02/20**.

15. Hoeppner MP, Lundquist A, Pirun M, Meadows JR, Zamani N, Johnson J, Sundstrom G, Cook A, FitzGerald MG, Swofford R *et al*: **An improved canine genome and a comprehensive catalogue of coding genes and non-coding transcripts**. *PLoS One* 2014, **9**(3):e91172.

16. Chen WK, Swartz JD, Rush LJ, Alvarez CE: **Mapping DNA structural variation in dogs**. *Genome Res* 2009, **19**(3):500-509.

17. Kolmogorov M, Yuan J, Lin Y, Pevzner PA: **Assembly of long, error-prone reads using repeat graphs**. *Nat Biotechnol* 2019, **37**(5):540-546.

18. Lin Y, Yuan J, Kolmogorov M, Shen MW, Chaisson M, Pevzner PA: **Assembly of long error-prone reads using de Bruijn graphs**. *Proc Natl Acad Sci U S A* 2016, **113**(52):E8396-E8405.

19. Vaser R, Sovic I, Nagarajan N, Sikic M: **Fast and accurate *de novo* genome assembly from long uncorrected reads**. *Genome Res* 2017, **27**(5):737-746.

20. ONT: **Medaka.** [https://github.com/nanoporetech/medaka] 2018. v1.03.

21. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK *et al*: **Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement**. *PLoS One* 2014, **9**(11):e112963.

22. Dudchenko O, M.S. S, Batra SS, Durand NC, Musial NT, Mostofa R, Pham M, St Hilaire BG, Yao W, Stamenova E *et al*: **The Juicebox Assembly Tools module facilitates *de novo* assembly of mammalian genomes with chromosome-length scaffolds for under $1000**. *bioRxiv* 2018:1-26.

23. Robinson JT, Turner D, Durand NC, Thorvaldsdottir H, Mesirov JP, Aiden EL: **Juicebox.js provides a cloud-based visualization system for Hi-C data**. *Cell Syst* 2018, **6**(2):256-258 e251.

24. Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL, Machol I, Omer AD, Lander ES *et al*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping**. *Cell* 2014, **159**(7):1665-1680.

1015   25.   English AC, Richards S, Han Y, Wang M, Vee V, Qu J, Qin X, Muzny DM, Reid JG, Worley KC *et al*: **Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology**. *PLoS One* 2012, **7**(11):e47768.

26.   Edwards RJ: **PAFScaff: Pairwise mApping Format reference-based scaffold anchoring and super-scaffolding.** [https://github.com/slimsuite/pafscaff] 2019.

1020   v0.4.0.

27.   Edwards RJ: **Diploidocus: Diploid genome assembly analysis tools.** [https://github.com/slimsuite/diploidocus] 2020. v0.9.0 & v0.10.0.

28.   Li H: **Minimap2: pairwise alignment for nucleotide sequences**. *Bioinformatics* 2018, **34**(18):3094-3100.

1025   29.   Davey NE, Shields DC, Edwards RJ: **SLiMDisc: short, linear motif discovery, correcting for common evolutionary descent**. *Nucleic Acids Res* 2006, **34**(12):3546-3554.

30.   Cabanettes F, Klopp C: **D-GENIES: dot plot large genomes in an interactive, efficient and simple way**. *PeerJ* 2018, **6**:e4958.

1030   31.   Morgulis A, Coulouris G, Raytselis Y, Madden TL, Agarwala R, Schaffer AA: **Database indexing for production MegaBLAST searches**. *Bioinformatics* 2008, **24**(16):1757-1764.

32.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL: **BLAST+: architecture and applications**. *BMC Bioinformatics* 2009, **10**:421.

1035   33.   Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM: **BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs**. *Bioinformatics* 2015, **31**(19):3210-3212.

34.   Urich MA, Nery JR, Lister R, Schmitz RJ, Ecker JR: **MethylC-seq library preparation for base-resolution whole-genome bisulfite sequencing**. *Nat Protoc*

1040   2015, **10**(3):475-483.

35.   Stadler MB, Murr R, Burger L, Ivanek R, Lienert F, Scholer A, van Nimwegen E, Wirbelauer C, Oakeley EJ, Gaidatzis D *et al*: **DNA-binding factors shape the mouse methylome at distal regulatory regions**. *Nature* 2011, **480**(7378):490-495.

36.   Burger L, Gaidatzis D, Schubeler D, Stadler MB: **Identification of active regulatory**

1045   **regions from DNA methylation data**. *Nucleic Acids Res* 2013, **41**(16):e155.

37.   Mo A, Mukamel EA, Davis FP, Luo C, Henry GL, Picard S, Urich MA, Nery JR, Sejnowski TJ, Lister R *et al*: **Epigenomic signatures of neuronal diversity in the mammalian brain**. *Neuron* 2015, **86**(6):1369-1384.

38.   Keilwagen J, Hartung F, Grau J: **GeMoMa: Homology-based gene prediction**

1050   **utilizing intron position conservation and RNA-seq data**. *Methods Mol Biol* 2019, **1962**:161-177.

39.   Altenhoff AM, Garrayo-Ventas J, Cosentino S, Emms D, Glover NM, Hernandez-Plaza A, Nevers Y, Sundesha V, Szklarczyk D, Fernandez JM *et al*: **The Quest for Orthologs benchmark service and consensus calls in 2020**. *Nucleic Acids Res*

1055   2020, **48**(W1):W538-W545.

40.   Richly E, Leister D: **NUMTs in sequenced eukaryotic genomes**. *Mol Biol Evol* 2004, **21**(6):1081-1084.

41.   Tsuji J, Frith MC, Tomii K, Horton P: **Mammalian NUMT insertion is non-random**. *Nucleic Acids Res* 2012, **40**(18):9073-9088.

1060   42.   Shi H, Xing Y, Mao X: **The little brown bat nuclear genome contains an entire mitochondrial genome: Real or artifact?** *Gene* 2017, **629**:64-67.

43.   Verscheure S, Backeljau T, Desmyter S: **In silico discovery of a nearly complete mitochondrial genome Numt in the dog (Canis lupus familiaris) nuclear genome**. *Genetica* 2015, **143**(4):453-458.

1065   44.   Hazkani-Covo E, Martin WF: **Quantifying the Number of Independent Organelle DNA Insertions in Genome Evolution and Human Health**. *Genome Biol Evol* 2017, **9**(5):1190-1203.

45.   Pesole G, Allen JF, Lane N, Martin W, Rand DM, Schatz G, Saccone C: **The neglected genome**. *EMBO Rep* 2012, **13**(6):473-474.

1070   46.   Dayama G, Emery SB, Kidd JM, Mills RE: **The genomic landscape of polymorphic human nuclear mitochondrial insertions**. *Nucleic Acids Res* 2014, **42**(20):12640-12649.

47.   Lopez JV, Yuhki N, Masuda R, Modi W, O'Brien SJ: **Numt, a recent transfer and tandem amplification of mitochondrial DNA to the nuclear genome of the**
1075   **domestic cat**. *J Mol Evol* 1994, **39**(2):174-190.

48.   Chen JM, Chuzhanova N, Stenson PD, Ferec C, Cooper DN: **Meta-analysis of gross insertions causing human genetic disease: novel mutational mechanisms and the role of replication slippage**. *Hum Mutat* 2005, **25**(2):207-221.

49.   Edwards RJ: **NUMTFinder: Nuclear mitochondrial fragment (NUMT) search**
1080   **tool.** [https://github.com/slimsuite/numtfinder] 2021. v0.1.0.

50.   Martin EC: **Dr. Johnson's Apple Orchard: The Story of America's First Pet Cemetery**: Hartsdale Canine Cemetery; 1997.

51.   Arendt M, Fall T, Lindblad-Toh K, Axelsson E: **Amylase activity is associated with AMY2B copy numbers in dog: implications for dog domestication, diet and**
1085   **diabetes**. *Anim Genet* 2014, **45**(5):716-722.

52.   Halo JV, Pendleton AL, Shen F, Doucet AJ, Derrien T, Hitte C, Kirby LE, Myers B, Sliwerska E, Emery S *et al*: **Long-read assembly of a Great Dane genome highlights the contribution of GC-rich sequence and mobile elements to canine genomes**. *bioRxiv* 2020:2020.2007.2031.231761.

1090   53.   Player RA, Forsyth ER, Verratti KJ, Mohr DW, Scott AF, Bradburne CE: **A novel canis lupus familiaris reference genome improves variant resolution for use in breed-specific GWAS**. *Life Sci Alliance* 2021, **4**(4).

54.   Wang C, Wallerman O, Arendt ML, Sundstrom E, Karlsson A, Nordin J, Makelainen S, Pielberg GR, Hanson J, Ohlsson A *et al*: **A novel canine reference genome**
1095   **resolves genomic architecture and uncovers transcript complexity**. *Commun Biol* 2021, **4**(1):185.

55.   Mainka SA: **Fanconi syndrome in a Basenji**. *Can Vet J* 1985, **26**(10):303-305.

56.   Johnson GS, Farias F: **Progress in a 15 year search for the genetic causes for Fanconi syndrome in Basenjs**. In*: 2011; Direct Fanconi Syndrome DNA test FAQ*:
1100   Basenji Club of Ameica Website; 2011.

57.   Li H: **Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. . 1303.** . *ArXiv* 2013, **1303**:1-3.

58.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM: **Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and**
1105   **repeat separation**. *Genome Res* 2017, **27**(5):722-736.

59.   Pacific Biosciences: **GenomicConsensus: Genome polishing and variant calling.** [https://github.com/PacificBiosciences/GenomicConsensus] 2019. 1 November 2019.

60.   Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL: **Juicebox provides a visualization system for Hi-C contact maps with unlimited**
1110   **zoom**. *Cell Syst* 2016, **3**(1):99-101.

61.   Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol I, Lander ES, Aiden AP *et al*: ***De novo* assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds**. *Science* 2017, **356**(6333):92-95.

1115 62. Bushnell B: **BBMap short read aligner, and other bioinformatic tools.** [https://sourceforge.net/projects/bbmap/] 2019 v38.51.

 63. Roach MJ, Schmidt SA, Borneman AR: **Purge Haplotigs: allelic contig reassignment for third-gen diploid genome assemblies**. *BMC Bioinformatics* 2018, **19**(1):460.

1120 64. Quinlan AR: **BEDTools: The Swiss-Army Tool for Genome Feature Analysis**. *Curr Protoc Bioinformatics* 2014, **47**:11 12 11-34.

 65. Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing genomic features**. *Bioinformatics* 2010, **26**(6):841-842.

 66. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
1125  Durbin R, Genome Project Data Processing S: **The Sequence Alignment/Map format and SAMtools**. *Bioinformatics* 2009, **25**(16):2078-2079.

 67. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool**. *J Mol Biol* 1990, **215**(3):403-410.

 68. Edwards RJ, Paulsen K, Aguilar Gomez CM, Perez-Bercoff A: **Computational**
1130  **Prediction of Disordered Protein Motifs Using SLiMSuite**. *Methods Mol Biol* 2020, **2141**:37-72.

 69. Edwards RJ: **SLiMSuite v1.9.1.** [https://github.com/slimsuite/SLiMSuite] 2020.

 70. Wheeler TJ, Eddy SR: **nhmmer: DNA homology search with profile HMMs**. *Bioinformatics* 2013, **29**(19):2487-2489.

1135 71. Edwards RJ: **BUSCOMP: BUSCO compilation and comparison – Assessing completeness in multiple genome assemblies.** [https://github.com/slimsuite/buscomp] 2019. v0.9.4.

 72. Edwards RJ: **BUSCOMP: BUSCO compilation and comparison – Assessing completeness in multiple genome assemblies [version 1; not peer reviewed]**.
1140  *F1000Res* 2019, **8**:995.

 73. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina sequence data**. *Bioinformatics* 2014, **30**(15):2114-2120.

 74. Chen H, Smith AD, Chen T: **WALT: fast and accurate read mapping for bisulfite sequencing**. *Bioinformatics* 2016, **32**(22):3507-3509.

1145 75. Gamaarachchi H, Lam CW, Jayatilaka G, Samarakoon H, Simpson JT, Smith MA, Parameswaran S: **GPU accelerated adaptive banded event alignment for rapid comparative nanopore signal analysis**. *BMC Bioinformatics* 2020, **21**(1):343.

 76. Loman NJ, Quick J, Simpson JT: **A complete bacterial genome assembled de novo using only nanopore sequencing data**. *Nat Methods* 2015, **12**(8):733-735.

1150 77. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE *et al*: **Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data**. *Nat Methods* 2013, **10**(6):563-569.

 78. Slater GS, Birney E: **Automated generation of heuristics for biological sequence**
1155  **comparison**. *BMC Bioinformatics* 2005, **6**:31.

 79. Ollivier M, Tresset A, Bastian F, Lagoutte L, Axelsson E, Arendt ML, Balasescu A, Marshour M, Sablin MV, Salanova L *et al*: **Amy2B copy number variation reveals starch diet adaptations in ancient European dogs**. *Roy Soc Open Sci* 2016, **3**(11).

 80. Steinegger M, Soding J: **MMseqs2 enables sensitive protein sequence searching**
1160  **for the analysis of massive data sets**. *Nat Biotechnol* 2017, **35**(11):1026-1028.

 81. Edwards RJ: **SAAGA: Summarise, Annotate & Assess Genome Annotations.** [https://github.com/slimsuite/saaga] 2020. v0.4.0.

 82. Seeman T: **Barrnap: BAsic Rapid Ribosomal RNA Predictor.** [https://github.com/tseemann/barrnap] 2018. v0.9.

1165   83.   Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, von Haeseler A, Schatz MC: **Accurate detection of complex structural variations using single-molecule sequencing**. *Nat Methods* 2018, **15**(6):461-468.

     84.   Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Azov AG, Bennett R *et al*: **Ensembl 2020**. *Nucleic Acids Res* 2020,
1170   **48**(D1):D682-D688.

     85.   Jagannathan V, Drogemuller C, Leeb T, Dog Biomedical Variant Database C: **A comprehensive biomedical variant catalogue based on whole genome sequences of 582 dogs and eight wolves**. *Anim Genet* 2019, **50**(6):695-704.

     86.   Leinonen R, Sugawara H, Shumway M, International Nucleotide Sequence Database
1175   C: **The sequence read archive**. *Nucleic Acids Res* 2011, **39**(Database issue):D19-21.

     87.   Field MA, Cho V, Andrews TD, Goodnow CC: **Reliably Detecting Clinically Important Variants Requires Both Combined Variant Calls and Optimized Filtering Strategies**. *PLoS One* 2015, **10**(11):e0143199.

     88.   Li H, Durbin R: **Fast and accurate short read alignment with Burrows-Wheeler**
1180   **transform**. *Bioinformatics* 2009, **25**(14):1754-1760.

     89.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M *et al*: **The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data**. *Genome Res* 2010, **20**(9):1297-1303.

1185   90.   Waardenberg AJ, Field MA: **consensusDE: an R package for assessing consensus of multiple RNA-seq algorithms with RUV correction**. *PeerJ* 2019, **7**:e8206.

Supplementary Figures and Tables

1190   **Additional file 1 (.pdf): Supplementary Figures**

Supplementary Figure 1. The workflow used to construct the de novo genomes.

Supplementary Figure 2. KAT kmer analysis of China assembly.

Supplementary Figure 3. Identification of putative regulatory elements in basenji's genome

using whole-genome bisulphite sequencing.

1195   Supplementary Figure 4. Predicted copy number of BUSCO Complete genes for three dogs

based on long-read read depth.

Supplementary Figure 5. CanFam_Bas nuclear mitochondrial DNA (NUMT) coverage.

Supplementary Figure 6. Comparative short read mapping and single nucleotide variant

calling for 58 dog breeds versus three reference genomes: CanFam_Bas, CanFam_GSD

1200   (GSD) and CanFam3.1 (BOX).

Additional file 2 (.xlsx): Supplementary Tables

Supplementary Table 1. Assembly statistics for China assembly stages and other dog genomes used in the study.

Supplementary Table 2. Diploidocus statistics and ratings for China scaffolds during tidying.

Supplementary Table 3. GeMoMa summary statistics and annotation comparisons.

Supplementary Table 4. AMY2B short read mapping and estimated copy number in Basenji dogs.

Supplementary Table 5. CanFam_Bas nuclear mitochondrial sequence (NUMT) fragments.

Supplementary Table 6. CanFam_Bas nuclear mitochondrial sequences (NUMTs) merging fragments within 8 kb.

Supplementary Table 7. Structural variant calls for Basenji and German Shepherd long read data versus three reference genomes.

Supplementary Table 8. Dog breeds used for read mapping and SNV analysis.

Supplementary Table 9. SRA runs for representative dog breeds from DBVDC.