The chemotherapeutic drug methotrexate selects for antibiotic resistance

Jónína S. Guðmundsdóttir[1*], Elizabeth G. A. Fredheim[1], Catharina I. M. Koumans[2], Joachim Hegstad[1,3], Po-Cheng Tang[4], Dan I. Andersson[4], Ørjan Samuelsen[1,5] and Pål J. Johnsen[1]

[1]Department of Pharmacy, Faculty of Health Sciences, UiT The Arctic University of Norway, Tromsø, Norway.

[2]Institute of Biology, Leiden University, Leiden, The Netherlands.

[3]Research and Development Division, Department of Microbiology and Infection Control, University Hospital of North Norway, Tromsø, Norway.

[4]Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden.

[5]Norwegian National Advisory Unit on Detection of Antimicrobial Resistance, Department of Microbiology and Infection Control, University Hospital of North Norway, Tromsø, Norway.

*Correspondence to: jonina.gudmundsdottir@uit.no

Abstract

Background: Understanding drivers of antibiotic resistance evolution is fundamental for

designing optimal treatment strategies and interventions to reduce the spread of antibiotic

resistance. Various cytotoxic drugs used in cancer chemotherapy have antibacterial properties,

but how bacterial populations are affected by these selective pressures is unknown. Here we

test the hypothesis that the widely used cytotoxic drug methotrexate affects the evolution and

selection of antibiotic resistance.

Methods: First, we determined methotrexate susceptibility (IC_{90}) in a collection of clinical

strains with and without pre-existing trimethoprim resistance determinants and the co-selective

abilities of methotrexate by growth curve measurements. We constructed fluorescently labelled

isogenic pairs of *E. coli* MG1655 differing only in trimethoprim resistance determinants

(chromosomal *folA* mutations and plasmid-mediated *dfrA* genes) and determined the minimum

selective concentrations of methotrexate using flow-cytometry. We further used an

experimental evolution approach to investigate the effects of methotrexate on *de novo*

trimethoprim resistance evolution in a clinical *E. coli* strain.

Findings: We show that methotrexate can select for acquired trimethoprim resistance

determinants located on the chromosome or a plasmid in clinical strains of *E. coli*. Additionally,

methotrexate co-selects for genetically linked resistance determinants when present together

with trimethoprim resistance on a multi-drug resistance clinical plasmid. These selective effects

occur at concentrations 40- to >320-fold below the methotrexate minimal inhibitory

concentration for *E. coli* and is independent of the well characterized efflux mediated intrinsic

resistance mechanisms.

Interpretation: Our results strongly suggest a selective role of methotrexate for virtually any

antibiotic resistance determinant when present together with trimethoprim resistance on a

multi-drug resistance plasmid. The presented results may have significant implications for

patient groups strongly depending on effective antibiotic treatment.

Funding: UiT The Arctic University of Norway, Northern Norway Regional Health Authority

and the Swedish Research Council.

Research in context

Evidence before this study: We searched PubMed and Web of Science for the following

search terms: Methotrexate + trimethoprim + resistan*; Cytostatic* + resistan* + antibiotic*;

Cytostatic* + cross + resistan* + antibiotic; Chemotherapy + driving + resistan* + antibiotic*;

drivers/driving + antimicro* + resistan* in May 2020. The evidence indicate that cytostatic

drugs, including methotrexate, a widely used drug for treatment of cancer and inflammatory

diseases, share molecular targets with commonly used antibiotics (e.g. trimethoprim). Further,

it is proposed that cytostatics may drive antibiotic resistance evolution due to microbiome

alterations, overlapping intrinsic resistome, and SOS induced mutagenesis. Important questions

remain regarding direct evidence for cross-resistance development between cytostatics and

antibiotics, and how these affect the selection, and spread of acquired antibiotic resistance

determinants.

Added value: To our knowledge, we show for the first time that methotrexate directly selects

for acquired trimethoprim resistance determinants on the chromosome as well as on a clinical

multi-drug resistance plasmid. These selective and co-selective effects occur at low

methotrexate concentration ranges expected to represent intestinal concentrations during

clinical use.

Implications: The presented data show that methotrexate has the potential to select for virtually

any given antibiotic resistance gene when genetically linked to trimethoprim resistance**.** This

study highlights the need for increased awareness of the presence of acquired antibiotic

resistance determinants in the gut of patients with impaired immunity undergoing methotrexate

treatment to preserve the effects of downstream antibiotic treatments.

Introduction

Antibiotic resistance is a major risk factor for patients with impaired immunity, such as cancer

patients, and often a patient's survival depends on antibiotic treatment to reduce the risk for

hospital-acquired infections during chemotherapy[1]. Several cytotoxic drugs used in cancer

chemotherapy are known to both elevate bacterial mutation rates and have direct antimicrobial

properties[2]. It has been proposed that cancer chemotherapy may drive *de novo* antibiotic

resistance evolution through SOS induced mutagenesis[3], and some reports have provided

support for this hypothesis[4,5]. Recently, the effects of non-antibacterial drugs on bacteria

typically found in the human gut were thoroughly explored and cytotoxic drugs were reported

to cause the most severe alterations of the microbiota[6]. Taken together, these studies suggest

that cytotoxic drugs affect survival of human gut commensals, they may increase the

evolvability of bacterial populations, and lead to reduced susceptibility towards drugs used to

treat cancer in bacteria. How bacterial populations respond to selective and co-selective

pressures exerted by individual cytotoxic drugs and the implications for antibiotic resistance

selection and spread is unknown. Thus, there is an urgent need to understand these potential

collateral effects of cancer chemotherapy to ensure effective antibiotic treatment for a large

group of immunocompromised patients. Moreover, cytotoxic drugs may constitute a previously

unrecognized target for intervention to limit the selection and spread of antibiotic resistance.

Methotrexate (MTX) is widely used in treatments including but not limited to; cancer of the

breast, skin, head, neck, and lung as well as many inflammatory diseases, such as rheumatoid

arthritis[7]. We specifically targeted resistance towards trimethoprim (TMP), since both drugs act

through inhibiting the dihydrofolate reductase enzyme in bacteria and eukaryotic cells, central

in DNA synthesis[8]. TMP in combination with sulfamethoxazole is among the most frequently

used antibiotics in the treatment of urinary tract infections and is recommended as first line

treatment internationally[9]. Our target organism in this study is *Escherichia coli*, the most

common agent of nosocomial infections world-wide[10]. *E. coli* is known to display intrinsic

resistance towards methotrexate through AcrAB-TolC mediated efflux[11], however TMP is not

a substrate for this efflux system.

Previous studies have focused on the abilities of MTX and other non-antibiotics to inhibit

bacterial growth[6]. These approaches have provided valuable insights on the effects of non-

antibiotics as modulators of the intestinal flora but lacked the necessary resolution to detect

more subtle selective effects on acquired antibiotic resistance determinants.

Here, we hypothesize that despite the demonstrated *E. coli* intrinsic MTX resistance[11], MTX

can affect antibiotic resistance evolution in *E. coli,* due to the shared molecular target with

TMP. We show that MTX selects for acquired bacterial TMP resistance (TMP^R) and co-selects

for other antibiotic resistance determinants when co-residing on a mobile genetic element.

Exposure to a wide concentration range of MTX selects for mutations identical to those

emerging during TMP selection in clinical isolates of *E. coli*. Moreover, we show that the

minimum selective concentrations (MSCs) of MTX and positive selection for chromosomal

and plasmid-mediated TMP^R determinants occurs at concentrations 40- and >320-fold below

the MTX minimum inhibitory concentrations (MICs), respectively.

Methods

Bacterial strains and growth conditions: Bacterial strains used in this study and their

assigned reference numbers are listed in Table S1. Most strain constructs were derived from

the *E. coli* K-12 strain MG1655 (DA4201/DA5438), or for cloning purposes from *E. coli* DH5α

(MP18-09). Two isogenic pairs of clinical *E. coli* strains, where one strain is trimethoprim

susceptible (TMP^S) and the other is TMP^R, were used for fitness measurements. In the first pair,

TMP^R in *E. coli* K56-2 (MP06-01) is caused by one intragenic point mutation T>A (W30R) in

the *folA* gene, and one in its promotor region (P$_{folA}$, C>T 58 base pairs (bp) upstream of the

gene)[12,13] (MP06-05). In the second pair, *E. coli* K56-75 (MP06-41) harbouring the pG06-VIM-

1 multi-drug resistance plasmid represents the TMP^R counterpart (MP05-31). This plasmid

encodes, amongst seven other antibiotic resistance genes, two different *dfrA* genes (*dfrA1* and

dfrA12) as putative TMP^R determinants as well as a carbapenem resistance determinant (*bla*$_{VIM-}$

$_1$)[14]. For accurate MSC measurements we constructed pairs of fluorescently tagged (*yfp* and

bfp) *E. coli* MG1655 strains containing the same resistance markers as described above. For

details concerning growth conditions, susceptibility testing and strain constructions see

protocols in supplementary information (SI).

Fitness measurements: Growth rates were determined using a BioscreenC MBR reader (Oy

Growth Curves Ab, Ltd). The growth rate calculations were done using the Bioscreen Analysis

Tool BAT 2.1[15] (see SI for detailed protocol). Competition experiments were performed in six

biological replicates, including dye swaps, using the fluorescently tagged strain pairs described

above, containing *folA* or pG06-VIM-1 mediated TMP^R. Both competition experiments, MSC

calculations and competitions performed at different starting ratios were done as previously

described in[16] (see SI for detailed protocols).

Experimental evolution: To examine the effect of MTX presence on TMP^R evolution, strain

K56-2 (MP06-01) was serially passaged in liquid cultures with MTX supplemented at a

concentration slightly above the estimated MSC. Initially, ten independent overnight cultures

were started from independent colonies on separate agar plates. From the original overnight

cultures, ~10^3 cells were used to start ten independent lineages in 1 mL batch cultures

containing 400 μg/mL MTX (lineages 1-10). Every 12 h for 25 days, the lineages were serially

passaged by 1,000-fold dilution, allowing for ~500 generations of growth. Every ~50

generations the populations were frozen down at –80°C for downstream analysis. For detailed

protocols concerning experimental evolution, selective plating on high concentrations of MTX

and whole genome sequencing see SI.

Results

Methotrexate selects for pre-existing trimethoprim resistance determinants

We determined the MICs of MTX in clinical and laboratory strains of *E. coli* (Table S1, SI).

Initial experiments revealed variable, but high MTX MICs (Table S2) consistent with previous

reports demonstrating that *E. coli* displays intrinsic resistance towards MTX due to AcrAB-

TolC mediated efflux[6,11]. We also observed that the MTX MIC was dependent on the presence

of TMP[R] determinants. All isolates with a functional TMP resistance determinant and increased

TMP MIC showed consistently higher MTX MICs (>32 mg/mL) than TMP susceptible isolates

(4-32 mg/mL), indicating possible co-selective abilities of the two drugs.

Antibiotic resistance selection and co-selection have traditionally been assumed to occur

between the MICs of susceptible and resistant isolates within a bacterial population (known as

the selective window)[16]. However, several reports unequivocally show that antibiotic resistance

selection and co-selection can occur at concentrations several hundredfold below the MIC of a

susceptible isolate (known as sub-MIC)[16,17]. To test how sub-MICs of MTX affect bacterial

fitness, we measured exponential growth rates for two pairs of clinical isogenic TMP[R] and

TMP[S] *E. coli* across a wide MTX concentration span. One pair with TMP[R] located on the

chromosome (two mutations associated with *folA*)[12] and one pair with TMP[R] (*dfrA*) located on

the MDR plasmid pG06-VIM-1[14]. TMP[S] strains displayed sharply declining growth rates

between 1 and 2 mg/mL of MTX, whereas the TMP[R] strains remained unaffected (Figure S1).

These results suggest a selective benefit during MTX exposure for TMP[R] strains at

concentrations below the observed MTX MIC of the TMP[S] clinical isolates.

The MSC describes pharmacodynamically the lowest concentration where selection for

resistance occurs[16]. To determine the MSC for MTX, we constructed a fluorescently tagged

pair of *E. coli* MG1655 strains to enable accurate separation between the two in mixed

populations. In these backgrounds, we introduced TMP[R], either through mutations (*folA*) using

genome engineering or the pG06-VIM-1 plasmid. The isogenic TMP[R] and TMP[S] strain pairs

were competed head-to-head by serial passage for 30 generations and the ratio of TMP[R]:TMP[S]

was determined over time using flow cytometry. From this data the MSC was estimated (Tables

S3-S5)[16]. *E. coli* with chromosomal *folA* mutations had an MSC of 200 µg/mL (1/40 of the

MIC of MTX) whereas those harboring the plasmid were close to the detection limit of the

assay[16] and we conservatively estimated the MSC to be <25 µg/mL (less than 1/320 of the MIC

of MTX) (Figure 1). Taken together, our data strongly suggest that selection for TMP[R] occurs

at MTX concentrations far below the estimated MTX MIC.

Figure 1: Selection coefficients as functions of MTX concentrations from competition

experiments between TMP[R] and TMP[S] isogenic strains. The MSC is defined as the

concentration where the selection coefficient equals zero. The MSC of *E. coli* MG1655

harboring **(A)** two chromosomal *folA* mutations (MP18-04 and MP18-07) is set to 200 µg/mL,

and **(B)** the MDR pG06-VIM-1 plasmid encoding *dfrA12* (MP18-05 and MP18-08) is

conservatively set at 25 µg/mL. Dashed lines represent the set MSC, bullets the average

selection coefficients and error bars the standard deviations.

Sub-MICs of methotrexate promotes invasion of trimethoprim resistance determinants

even when rare in *E. coli* populations

Exploring MTX-selective dynamics further, we asked if TMP^R determinants could invade the

population at lower initial densities to exclude potential bias from the 1:1 ratio in the

competition experiments. We started competition experiments from frequencies as low as 10^{-4}

of the TMP^R strains, at concentrations slightly above the estimated MSC of MTX (400 µg/mL

for *folA* mediated resistance and 75 µg/mL for pG06-VIM-1 mediated resistance) and followed

the change in ratios over 30 generations of growth (Figure 2). Both chromosomal and plasmid

mediated TMP^R determinants were able to invade, even when initially rare in their respective

populations, strongly suggesting that the MTX selective effects are independent on initial

frequencies of resistant and susceptible strains during competition experiments.

Figure 2: Competition experiments during sub-MIC MXT exposure at different initial

frequencies of TMP^R strains. The change in TMP^R:TMP^S ratios over 30 generations of growth

at MTX concentrations slightly above the MSCs where *E. coli* MG1655 harbors **(A)** two *folA*

mutations (MP18-07), and **(B)** the pG06-VIM-1 plasmid (MP18-08) were competed against a

differently tagged, isogenic susceptible strain (DA56507) at 1:1, $1:10^{-2}$, $1:10^{-3}$ and $1:10^{-4}$

starting ratios. Error bars represent the standard deviation of the average ratio.

Methotrexate co-selects for resistance determinants on a multi-drug resistance plasmid

The *dfr*-genes represent a common TMP^R mechanism in *E. coli* and these genes are frequently

located on mobile genetic elements such as integrons and plasmids. Given that MTX selects for

dfr-mediated TMP^R, co-selection of other genetically linked resistance genes is likely. To show

this, we used the MDR pG06-VIM-1 plasmid harboring *dfrA1* and *dfrA12* along with multiple

resistance determinants including four aminoglycoside resistance genes and the bla_{VIM-1}

carbapenemase gene conferring resistance to broad-spectrum β-lactams including

carbapenems[14]. To assess the stability of pG06-VIM-1 in our strains competing in the presence

of MTX, *E. coli* K56-75 harboring the plasmid (MP05-31) was serially passaged in batch

cultures with 400 μg/mL MTX supplemented for 50 generations. The lineages were then plated

on non-selective agar and 100 colonies from each lineage tested for reduced susceptibility

towards ampicillin, TMP, streptomycin and spectinomycin. The results revealed complete

phenotypic stability across all three lineages, confirming MTX mediated co-selection of

plasmid mediated MDR.

To verify that the TMP^R determinants on the MDR pG06-VIM-1 plasmid is the primary

mediators of MTX resistance and selection (Table S2), both *dfrA1* and *dfrA12* were isolated

from the plasmid and cloned onto an expression vector and the effects of the individual genes

measured. Of the two genes, only *dfrA12* was shown to give the same resistance pattern for

TMP as well as MTX as the pG06-VIM-1 plasmid (Figure 3), and the lack of detectable

phenotype for *dfrA1* (MP18-11) is likely due to a start codon frameshift mutation[14].

Figure 3: MTX and TMP MIC of *E. coli* DH5α expressing *dfrA1* or *dfrA12*. The MTX **(A)** and TMP **(B)** MIC for the wild type (WT) *E. coli* DH5α (MP18-09) compared to the strain harboring the empty pBAD30 (MP18-10) expression vector as well as strains with pBAD30 with different *dfrA* genes expressed under the inducible expression control by the pBAD promotor (MP18-11 and MP18-12). The detection limit of the assay is 64 µg/mL for TMP and 32 mg/mL for MTX. For both drugs, the MIC of *E. coli* DH5α expressing *dfrA*12 (MP18-12) exceeded the detection limit whereas the strain expressing *dfrA1* (MP18-11) has the same MICs as the WT strain. Showing that MTX and TMP resistance conferred by the pG06-VIM-1 plasmid is caused by the *dfrA12* gene.

Methotrexate selects for *de novo* trimethoprim resistance

We further examined whether exposure to MTX could lead to *de novo* TMP[R] evolution. We selected spontaneous mutants from overnight cultures with and without exposure to MTX, plated on selective agar at high MTX concentrations and tested for TMP cross-resistance (Figure S2, Table S6). *E. coli* K56-2 isolated at 16 and 32 mg/mL MTX (MP18-17 to MP18-

28) displayed increased MICs of TMP close to or above the clinical breakpoint[18], clearly

demonstrating selection for TMP^R by MTX. DNA sequencing of the resistant isolates revealed

two different mutations in the *folA* promoter, previously reported to result in TMP^{R19}, as well

as a single mutation in the *marR* gene (Table S6-S11).

Finally, we asked if exposure to sub-MICs of MTX close to the estimated MSCs would select

for *de novo* TMP^R mutations in a susceptible *E. coli* population. Starting from 1,000 cells to

minimize the probability of pre-existing mutants, we grew ten independent lineages of the *E.*

coli K56-2 strain at 400 μg/mL MTX for 500 generations. The frequency of TMP^R was

determined every 50 generations. TMP^R ascended in frequency in 2/10 lineages at different

rates and time-points during the first 250 generations before they were outcompeted by a

different set of mutants with reduced susceptibility to MTX and no cross-resistance to TMP

(Figure 4, Table S12). These experiments show that MTX exposure can select for *de novo*

TMP^R, both at high and sub-MIC concentrations.

Figure 4: Evolution of TMP^R during MTX exposure for 500 generations. (A) Sub-MIC

evolution experimental set-up. Ten biological replicates of K56-2 (MP06-01) were evolved for

~500 generations with 400 μg/mL MTX and three biological replicates without drug. All

lineages were screened for TMP^R every 50 generations. After 500 generations all end-point

populations were plated on 32 mg/mL MTX. All populations were able to grow at 32 mg/mL

MTX, but not a single clone isolated conferred TMP^R, strongly suggesting that reduced

susceptibility to MTX with no cross-resistance to TMP evolved in the endpoint populations.

(B) Fractions of TMP^R *folA* mutants isolated every 50 generations from the lineages where

these were detected. The detection limit of the assay was ~2x10^{-9}. Solid lines represent the two

lineages where TMP[R] emerged and ascended in frequency whereas dotted lines indicate

spontaneous mutants.

Pharmacokinetic approximations

To assess pharmacokinetic relevance, we attempted to estimate the MTX concentration range

likely to be found in the intestine of patients undergoing MTX treatment. Limited information

is available on gut MTX concentrations following intravenous administration during cancer

treatment, as pointed out by others[6]. Pharmacokinetic data reveal that up to 90% of administered

MTX is renally excreted[20] and we assume that the remaining ~10% of the dose constitutes the

upper limit of the concentration range present in the human intestine. The lower limit is set to

2% of the dose based on the mean ^{3}H labelled MTX concentrations measured in stool samples

from nine patients receiving MTX intravenously[21]. From this, we set a 24 hour transition time

in a total volume of 0.6 L[6] and calculated the dose (d) required to achieve MSC in the human

intestine from:

289
$$\frac{d}{0.6L} \; x \; (0.1 \; or \; 0{,}02) = MSC$$

Estimated doses needed to reach intestinal MSCs assuming 2% and 10% fecal MTX

concentrations were from 0.15g to 0.75g for plasmid-mediated TMP[R] and from 1.2g to 6g for

chromosomal *folA* mutations. Thus, assuming close to 2m^2 body surface in grown-up patients[22]

estimates of MSC for plasmid-mediated TMP[R] translates to dosing regimens from 75-375

mg/m^2 and from 0.6-3 g/m^2 for the chromosomal *folA* mutations. These approximations indicate

that our MSC estimates are relevant for patients receiving high dose MTX treatment (1-12

g/m^2)[23]. A recent study used a different approach estimating gut MTX concentrations following

oral administration during treatment of rheumatoid arthritis[24]. Their data suggested MTX

concentrations as high as 100 µg/mL are found in the gut, suggesting that our estimated MSC

for plasmid-mediated antibiotic resistance determinants (25 µg/mL) is well within this

concentration range.

Discussion

Here we show that exposure to the cytotoxic drug MTX affects selection and evolution of TMP[R]

determinants at clinically relevant concentrations. Notably, MTX can mediate selection of any

antibiotic resistance determinant when TMP[R] is co-localized on a mobile genetic element across

a wide concentration gradient. Arguably, this potentially important side-effect of MTX

treatment has been previously unrecognized, as studies on the effects of non-antibiotic drugs,

including MTX, have either focused on bacterial growth inhibition or used drug concentrations

around the MIC[6,25-27], with a few exceptions[17,28].

Using the approaches outlined here, including high resolution mixed culture competition

experiments, allow determination of the true MTX selective window ranging from the MSC to

the MIC[16]. This is particularly relevant for non-antibiotics for which bacteria display reduced

susceptibility. In *E. coli,* MTX is a substrate for the AcrAB-TolC efflux pump[11] and selective

effects as those demonstrated here would not have been detected in classical susceptibility

and/or growth assays in bacterial monocultures. This was recently supported in an *E. coli*

chemical genetic screen where clear growth inhibitory effects of MTX, as well as for a range

of other non-antibiotics, were only demonstrated in a *tolC* knock-out mutant (i.e. in a mutant

lacking the intrinsic mechanism of resistance)[6].

Given that many cytotoxic drugs are structurally similar to antibiotics (e.g.

doxorubicin/tetracyclines), or target similar key processes as the major antibiotic groups (e.g.

DNA/protein synthesis) it is possible that cancer chemotherapy may lead to increased levels of

antibiotic resistance in a vulnerable patient group that very often rely on efficient antibiotic

treatment for survival. To acquire a deeper understanding of the evolutionary potential of novel,

non-antibiotic drivers of antibiotic resistance the approaches presented here are essential. These

approaches need to be combined with an improved understanding of the intestinal

pharmacokinetics of MTX and other cytotoxic drugs, possible effects of co-administered drugs

such as leucovorin mediated MTX rescue[29], and their interactions with the human microbiome.

Such knowledge could allow identification of antibiotic + non-antibiotic drug combinations

that should be avoided to preempt resistance evolution.

Recent studies have showed that non-antibiotics can increase mutation rates[4,5] and promote

horizontal gene transfer[28,30]. Taken together with the data presented here it is clear that non-

antibiotic drugs have the potential to affect the evolution, selection, and spread of antibiotic

resistance determinants. It is also clear that without a better understanding of these potential

drivers of antibiotic resistance, interventions including those targeting human consumption of

antibiotics may be negatively affected.

Acknowledgments: We thank: Fredrik Sund, MD at The University Hospital North Norway

for valuable input on medical aspects of MTX. Ass. Prof. Daniel Rozen, Leiden University for

valuable discussions. Prof. Natasa Skalko-Basnet and Ass. Prof. Ann Mari Holsæter, UiT The

Arctic University of Norway, for training in handling of cytostatic drugs. Funding: PJJ was

supported by a joint grant from UiT The Arctic University of Norway and the Northern Norway

Regional Health Authority (A23270). DIA was supported by the Swedish Research Council

(grant 2017-01527).

Author contributions: The study was designed by JSG, EGAF, ØS and PJJ. Strain

constructions were done by PT and JSG. Experiments were conducted by JSG, EGAF and

CIMK. WGS analysis was done using a bioinformatic pipeline designed by JH. WGS data was

verified by JH and JSG. Experimental data was verified by JSG and PJJ. Formal data analysis

and data visualization was done by JSG. The first draft of the manuscript was written by JSG

and revised by PJJ. All authors revised subsequent version and provided key edits to the

manuscript. Funding acquisition and resources were provided by PJJ and DIA.

Competing Interests: The authors declare no competing interests. **Data availability:** WGS

data are available at NCBI (BioProject PRJNA677979).

References:

1.   Teillant A, Gandra S, Barter D, Morgan DJ, Laxminarayan R. Potential burden of
antibiotic resistance on surgery and cancer chemotherapy antibiotic prophylaxis in the USA:
a literature review and modelling study. *The Lancet Infectious Diseases* 2015; **15**(12): 1429-
37.
2.   Nakamura S-i, Oda Y, Shimada T, Oki I, Sugimoto K. SOS-inducing activity of chemical
carcinogens and mutagens in *Salmonella typhimurium* TA1535/pSK1002: examination with
151 chemicals. *Mutation Research Letters* 1987; **192**(4): 239-46.
3.   Papanicolas LE, Gordon DL, Wesselingh SL, Rogers GB. Not Just Antibiotics: Is Cancer
Chemotherapy Driving Antimicrobial Resistance? *Trends in Microbiology* 2018; **26**(5): 393-
400.
4.   Meunier A, Nerich V, Fagnoni-Legat C, *et al*. Enhanced emergence of antibiotic-
resistant pathogenic bacteria after *in vitro* induction with cancer chemotherapy drugs.
Journal of Antimicrobial Chemotherapy 2019; **74**(6): 1572-7.
5.   Hobson CA, Bonacorsi S, Hocquet D, *et al.* Impact of anticancer chemotherapy on the
extension of beta-lactamase spectrum: an example with KPC-type carbapenemase activity
towards ceftazidime-avibactam. *Scientific Reports* 2020; **10**(1): 589.
6.   Maier L, Pruteanu M, Kuhn M, *et al*. Extensive impact of non-antibiotic drugs on
human gut bacteria. *Nature* 2018; **555**: 623-8.
7.   Jolivet J, Cowan KH, Curt GA, Clendeninn NJ, Chabner BA. The Pharmacology and
Clinical Use of Methotrexate. *New England Journal of Medicine* 1983; **309**(18): 1094-104.
8.   Mauldin RV, Carroll MJ, Lee AL. Dynamic dysfunction in dihydrofolate reductase
results from antifolate drug binding: modulation of dynamics within a structural state.
Structure 2009; **17**(3): 386-94.
9.   Bergstrom P, Holm S, Grankvist K, Henriksson R. Interaction between antibiotics and
antineoplastic drugs on antibacterial activity *in vitro*: Estramustine phosphate sensitizes
pneumococci to amikacin. *International Journal of Oncology* 1994; **4**(2): 435-9.
10.   Struelens MJ, Denis O, Rodriguez-Villalobos H. Microbiology of nosocomial infections:
progress and challenges. *Microbes and Infection* 2004; **6**(11): 1043-8.
11.   Kopytek SJ, Dyer JC, Knapp GS, Hu JC. Resistance to methotrexate due to AcrAB-
dependent export from *Escherichia coli*. *Antimicrobial Agents and Chemotherapy* 2000;
44(11): 3210-2.

12.   Podnecky NL, Fredheim EGA, Kloos J, et al. Conserved collateral antibiotic
susceptibility networks in diverse clinical strains of *Escherichia coli*. *Nature Communications*
2018; **9**(1): 3673.
13.   Kahlmeter G. The ECO.SENS Project: a prospective, multinational, multicentre
epidemiological survey of the prevalence and antimicrobial susceptibility of urinary tract
pathogens--interim report. *The Journal of antimicrobial chemotherapy* 2000; **46 Suppl 1**: 15-
22; discussion 63-5.
14.   Di Luca MC, Samuelsen Ø, Starikova I, *et al.* Low biological cost of carbapenemase-
encoding plasmids following transfer from *Klebsiella pneumoniae* to *Escherichia coli*. *Journal*
of Antimicrobial Chemotherapy 2016; **72**(1): 85-9.
15.   Thulin M. "BAT: an online tool for analysing growth curves". 2018. (Available from:
http://www.mansthulin.se/bat/2019).
16.   Gullberg E, Cao S, Berg OG, *et al*. Selection of Resistant Bacteria at Very Low
Antibiotic Concentrations. *PLOS Pathogens* 2011; **7**(7): e1002158.
17.   Gullberg E, Albrecht LM, Karlsson C, Sandegren L, Andersson DI. Selection of a
Multidrug Resistance Plasmid by Sublethal Levels of Antibiotics and Heavy Metals. *mBio*
2014; **5**(5): e01918-14.
18.   The European Committee on Antimicrobial Susceptibility Testing "Breakpoint tables
for interpretation of MICs and zone diameters, Version 10.0". EUCAST, 2020.
19.   Eliopoulos GM, Huovinen P. Resistance to Trimethoprim-Sulfamethoxazole. *Clinical*
Infectious Diseases 2001; **32**(11): 1608-14.
20.   Seideman P, Beck O, Eksborg S, Wennberg M. The pharmacokinetics of methotrexate
and its 7-hydroxy metabolite in patients with rheumatoid arthritis. *British Journal of Clinical*
Pharmacology 1993; **35**(4): 409-12.
21.   Henderson ES, Adamson RH, Oliverio VT. The metabolic fate of tritiated
methotrexate. II. Absorption and excretion in man. *Cancer Research* 1965; **25**(7): 1018-24.
22.   Sacco JJ, Botten J, Macbeth F, Bagust A, Clark P. The Average Body Surface Area of
Adult Cancer Patients in the UK: A Multicentre Retrospective Study. *PLOS ONE* 2010; **5**(1):
e8933.
23.   Howard SC, McCormick J, Pui C-H, Buddington RK, Harvey RD. Preventing and
Managing Toxicities of High-Dose Methotrexate. *Oncologist* 2016; **21**(12): 1471-82.
24.   Nayak RR, Alexander M, Deshpande I, et al. Methotrexate impacts conserved
pathways in diverse human gut bacteria leading to decreased host immune activation. *Cell*
Host Microbe 2021.
25.   Kruszewska H, Zareba T, Tyski S. Antimicrobial activity of selected non-antibiotics -
Activity of methotrexate against *staphylococcus aureus* strains. *Acta Poloniae Pharmaceutica*
2000.
26.   Moody MR, Morris MJ, Young VM, Moyé LA, Schimpff SC, Wiernik PH. Effect of Two
Cancer Chemotherapeutic Agents on the Antibacterial Activity of Three Antimicrobial
Agents. *Antimicrobial Agents and Chemotherapy* 1978; **14**(5): 737-42.
27.   Nyhlén A, Ljungberg B, Nilsson-Ehle I, Odenholt I. Bactericidal Effect of Combinations
of Antibiotic and Antineoplastic Agents against *Staphylococcus aureus* and *Escherichia coli*.
Chemotherapy 2002; **48**(2): 71-7.
28.   Wang Y, Lu J, Mao L, *et al.* Antiepileptic drug carbamazepine promotes horizontal
transfer of plasmid-borne multi-antibiotic resistance genes within and across bacterial
genera. *The ISME Journal* 2019; **13**(2): 509-22.

29.   Van der Beek JN, Oosterom N, Pieters R, de Jonge R, van den Heuvel-Eibrink MM, Heil
SG. The effect of leucovorin rescue therapy on methotrexate-induced oral mucositis in the
treatment of paediatric ALL: A systematic review. *Critical Reviews in Oncology/Hematology*
2019; **142**: 1-8.
30.   Wang Y, Lu J, Engelstädter J, *et al*. Non-antibiotic pharmaceuticals enhance the
transmission of exogenous antibiotic resistance genes through bacterial transformation. *The*
ISME Journal 2020; **14**(8): 2179-96.
441