1
7   **Shapes and Genescapes: Mapping Multivariate Phenotype-Biological Process**
8   **Associations for Craniofacial Shape**
9
10
11 November 12, 2020
12

J. David Aponte[1], David C. Katz[1], Wei Liu[1], Fernando Andrade[2], Charles C. Roseman[3],

Stephen A. Murray[4], James Cheverud[2], Ralph S. Marcucio[5], Benedikt Hallgrímsson[1*]

1) Department of Cell Biology & Anatomy, Alberta Children's Hospital Research Institute and

McCaig Bone and Joint Institute, Cumming School of Medicine, University of Calgary,

AB, Canada

2) Department of Biology, Loyola University Chicago, Chicago, IL, USA

3) Department of Animal Biology, University of Illinois Urbana Champaign, Urbana, IL, USA

4) The Jackson Laboratory, Bar Harbor, ME, USA

5) Department of Orthopaedic Surgery, School of Medicine, University of California San

Francisco, San Francisco, CA, USA

* Correspondence to bhallgri@ucalgary.ca

Abstract

The importance of coordinated gene effects on morphological phenotypes is clear from the

intertwining of gene actions in signaling pathways, gene regulatory networks, and developmental

processes underlying the development of shape and size. However, genome wide association

studies identify and localize the contribution of individual genes to phenotypic variation, seldom

targeting the coordinated influence of multiple genes on phenotypes. Here, we explicitly model

the joint effect of biologically coherent collections of genes on craniofacial shape in a sample of

$n = 1,145$ mice from the Diversity Outbred (DO) experimental line. We use biological process

gene ontology (GO) annotations to select skeletal and facial development gene sets and solve for

the axis of shape variation that maximally covaries with gene set marker variation. Our process-

centered view of cranial shape variation allows us to consider fundamental issues that are

difficult to address with existing methods, such as the extent to which variation in different

processes relevant to the same phenotype correspond to similar axes of shape variation, and

whether the direction of the effect of experimental mutations to a pathway gene is similar to the

primary axis of shape variation associated with that pathway. Finally, we introduce an online

application which provides users the means to customize their own process-centered craniofacial

shape analyses in the DO. The process-centered approach is generally applicable to any

continuously varying phenotype and thus has wide-reaching implications for complex-trait

genetics.

64 **Keywords**: multivariate genotype-phenotype map, complex traits, mouse, craniofacial, diversity

65 outbred

Introduction

Variation in human craniofacial shape is moderately to highly heritable (~30-70% (Cole et al., 2017; Tsagkrasoulis et al., 2017)), and resemblances among close relatives as well as twins underscore the strong relationship between shared genetics and shared phenotype (Johannsdottir et al., 2005; Nakata, 1985). Despite many studies in humans and in mice (Claes et al., 2018; Cole et al., 2016; Shaffer et al., 2016), we know very little about the genetic basis for variation in craniofacial shape. This is likely due to genetic complexity (Katz et al., 2019; Richtsmeier and Flaherty, 2013; Visscher, 2008; Wood et al., 2014; Wray et al., 2013). Like many aspects of morphological variation, craniofacial shape is extraordinarily polygenic. Genes with major mechanistic roles in facial development such as *Fgf8* often contribute little to standing phenotypic variation (Green et al., 2017) while genetic influences without obvious connections to craniofacial development emerge as significant contributors (Kenney-Hunt et al., 2008; Klingenberg and Leamy, 2001; Maga et al., 2015; Pallares et al., 2015, 2014). The effects of genetic variants on phenotype often depend on genetic background (Mackay and Moore, 2014; Percival et al., 2017) and many mutations have variably penetrant effects even when background is controlled (Hallgrimsson et al, 2009; Rendel, 1967). These issues likely arise because genetic influences act through multiple layers of interacting developmental processes to influence phenotypic traits, resulting in complex patterns of epistasis and variance heterogeneity (Hallgrimsson et al., 2018, 2014; Kawauchi et al., 2009; Wagner and Zhang, 2011). Solutions that go beyond studies of single gene effects are required to overcome these significant challenges in complex-trait genetics. Here, we implement an enhanced form of the more general candidate gene approach to evaluate the conjoint effects of multiple genes on a complex trait – craniofacial shape.

There are two basic approaches to mapping genetic effects on to phenotypic variation. A candidate gene approach measures genotypic values with known physiological and biochemical relationships to the phenotypes of interest (Cheverud and Routman, 1993). In contrast, a random marker or genome-wide approach seeks to associate any potential genetic variant with variation in the trait of interest. There are advantages and disadvantages to these two approaches. The candidate gene approach is blind to the unknown – phenotypic variation is often associated with genes not expected to be important. On the other hand, a candidate gene approach allows direct

measurement of genotypic values and produces results that are interpretable in terms of trait

physiology or development. A genome-wide or random marker approach can produce

unexpected insight by revealing novel gene-phenotype associations. However, this comes at a

great cost in power (Visscher et al., 2017). For highly polygenic traits, this approach often

produces a "tip of the iceberg" effect in which studies reveal a small and often incoherent subset

of the genes that actually determine variation in the trait of interest (Broman and Sen, 2009, p.

123-124).

Several strategies have been developed that partially overcome these tradeoffs. One

solution is the use of polygenic risk scores. Polygenic risk scores assess the overall genetic

influence on a trait without regard to the genome-wide significance of individual SNP effects

(Dudbridge, 2013; Wray et al., 2007). Approaches such as meta-analyses of genome-wide

association studies (GWAS) or studies based on extreme phenotypes (Morozova et al., 2015)

have expanded gene lists for a variety of complex traits. However, lengthy lists of genes or

overall genomic risk for specific phenotypes do not necessarily constitute tractable genetic

explanations for phenotypic variation. When 1000s of genes are required to explain heritable

variation in stature, for instance, it is not clear what such lists tell you beyond the obvious fact

that stature is heritable and polygenic (Yang et al., 2010; Wood et al., 2014). This tension

between hypothesis-driven and hypothesis-free approaches and their attendant tradeoffs between

statistical power and interpretability is, arguably, a major issue within complex trait genetics. To

resolve this conceptual conflict, approaches are needed that integrate quantitative genetics with

biological insights regarding the cellular and developmental processes through which genes

influence phenotypic variation.

Building on Mitterocker et al.'s (2016) multivariate genotype-phenotype (MGP) method, we

extend the candidate-gene framework to evaluate the combined contributions of genes that share

ontological information such as membership in pathways that influence craniofacial development

in mice. Our process-centered, multivariate approach brings traditional GWAS together with a

simplified model of the hierarchical genotype-phenotype (GP) map. GP maps describe the

relationship between genetic and phenotypic measurements (Lewontin, 1974). Understanding the

genetic determinants of craniofacial variation, as with most complex traits, represents a many-to-

many GP map problem (Fig 1). Both phenotypic and genotypic measurements have complex

within-set covariance structures. On the genetic side, the covariance structure is represented by

pathway/biochemical interactions, as well as chromosomal structure like linkage, chromatin, and

3D chromosomal organization. For shape-related phenotypes, the covariance matrix is structured

by the chosen set of landmarks and their resulting coordinates. The functional relationship from

genotype to phenotype is then described by a between-set covariance (Klingenberg and Leamy,

2001; Mitteroecker et al., 2016). To dissect these relationships, we use a regularized partial least

squares (PLS) (Lorenzo et al., 2019) approach to estimate a low-dimensional mapping from the

alleles in our sample to variation in adult mouse craniofacial shape. We focus on how allelic

variation in processes relevant to craniofacial development maps to craniofacial shape variation.

To do so, we estimate the joint effects on multivariate craniofacial variation of markers for

groups of genes that share functionally relevant gene ontology terms. We ask the following four

questions:

1) How much shape variation is communally accounted for by genes contributing to a

145   process, e.g., chondrocyte differentiation?

146   2) How similar are the effects of different processes on shape? For instance, do cell

147   proliferation genes affect face shape in a similar way to genes in the bone morphogenic

148   protein pathway?

149   3) How similar are mutant model effects and process effects? For example, do chondrocyte

150   mutant effects align with the effects of natural variants in chondrocyte differentiation

151   genes?

152   4) What is the overarching structure of process effects? Do process effects align with major

153   axes of variation or other interesting directions in morphospace?

Results

Process Multivariate Genotype-Phenotype (MGP) Mapping

We demonstrate regularized-PLS MGP mapping with two examples. The first estimates the primary axis of skull shape covariation with genes involved in chondrocyte differentiation (Fig 2). Differentiation of chondrocytes is one of several key developmental milestones involved in endochondral ossification. Endochondral bones form the majority of the cranial base through a cartilage model of bone formation (Percival and Richtsmeier, 2013). There are 38 genes annotated to chondrocyte differentiation in the Ensembl database (Yates et al., 2020). Genetic effects are shown as zero-centered bars that span the range of estimated allele effects across the 8 DO founders; individual founder allele effects—8 per marker—are color-coded within those bars (Fig 2A). Among chondrocyte differentiation genes, *Nov*, *Mapk14*, and *Bmpr1b* (*Alk6*) most heavily drive covariation with craniofacial shape. The phenotypic effects at each landmark— magnified 4x— primarily relate to antero-posterior positioning of the zygomatic arches and dorso-ventral jugal position (Fig 2B, 2C). The chondrocyte differentiation GP map explains 2.15% of the total variance in craniofacial shape. Compared to 1000 randomly generated marker sets of the same size (38), chondrocyte differentiation explains substantially more variation in phenotype than random markers (Supp fig 1A).

Figure 2B and 2C also compare the direction of the chondrocyte differentiation MGP axis to the axis of shape variation of a relevant mutant phenotype. We chose homozygous *Bmpr1b* mutants for this comparison because Bmpr1b heterodimerization with other bone morphogenic protein pathway receptors is essential for chondrocyte differentiation and proliferation (Liu et al., 2005; Yoon et al., 2005). The overall phenotypic directions of *Bmpr1b* mutant variation and chondrocyte differentiation variation are moderately correlated at r = 0.312, but the direction at landmarks with large effects in mutant and MGP are clearly coincident. Over the landmarks we measured, the chondrocyte differentiation effect is less global than the *Bmpr1b* effect, likely due to the difference in severity of the mutant phenotype.

The second example quantifies shape associations with the 41 genes annotated to
fibroblast growth factor (FGF) receptor signaling. The FGF pathway contributes to several
aspects of craniofacial development, from the epithelial to mesenchymal transition of neural
crest cell precursors in early facial formation, to the outgrowth of the later facial prominences
and specification of intramembranous and endochondral bone (Jin et al., 2012; Ornitz and Marie,
2015; Richman et al., 1997). Alleles from the WSB founder background for *Fgf2* and *Flrt2*
contribute the most variance to the phenotypic effect (Fig 3C), primarily the relative proportions
of vault and face (Fig 3A, 3B). The phenotype associated the *Fgfr* alleles is predominately
related to a larger neurocranium volume relative to the outgrowth of the face. FGFR signaling
loci explain 1.4% of the total variance in craniofacial shape, which greatly exceeds 1000
randomly selected marker sets of the same size (Supp fig 1B).
199
Finally, we compared the estimated FGF signaling MGP effect with the direction of an
Fgf10 homozygous mutant. FGF10 is a key ligand in early development, directing proliferation
as well as differentiation for many craniofacial components, including the palate, teeth, and
bones (Hilliard et al., 2005; Prochazkova et al., 2018; Watson and Francavilla, 2018). The vector
correlation between the FGF mutant and the estimated FGF signaling effect is 0.623. Both
phenotypes strongly affect the cranial base, and the vector correlation between base landmarks is
0.94.
207
In all, we have shown how sets of allelic variation in the genome can map to distinct
phenotypic directions. Each MGP model shows the relative contributions of different genotypes
for craniofacial variation. For both examples, we focus only on the first PLS axis, so other alleles
for other genes may contribute to novel phenotypic directions in lower PLS axes. In both cases,
we were able to find mouse mutant phenotypes with moderate to strongly correlated craniofacial
phenotypes. In the next sections we will examine how MGP phenotypes relate to each other, as
well as the phenotypic directions of many mutant mouse models.
215
216   **Pairwise comparison of craniofacial development processes**
217

We chose 15 processes integral to craniofacial development and compared the pairwise

similarity of effect on craniofacial shape using a heatmap based on clustering of the correlation

matrix (R Core Team, 2020). Processes with similar effects on craniofacial shape will be highly

correlated, while processes that affect distinct aspects of craniofacial variation will be

uncorrelated to each other. The clustering algorithm resulted in two main blocks of strongly

correlated effects (Fig 4A). The largest block of highly correlated phenotypic effects includes

neural crest cell migration, epithelial to mesenchymal transition, forebrain development, as well

as some of the most general developmental processes like cell proliferation, bone development,

apoptosis, A/P pattern specification, and FGFR signaling. In addition, there is a general BMP

block, with Bmp signaling, dorsoventral pattern formation, endochondral ossification, and

positive regulation of skeletal muscle tissue growth. Interestingly, phenotypic variation

associated with cranial suture morphogenesis, neural tube patterning, and intramembranous

ossification is largely uncorrelated with the other craniofacial developmental processes included

here.

To assess the stability of the clustering result, we estimated the vector correlation

between the cluster distances—also known as the cophenetic distance— and the original

correlation matrix. A high vector correlation suggests reliable clustering, whereas a low

correlation suggests a random clustering result. The correlation between the cophenetic distance

matrix and the correlation matrix is 0.648 (t = 8.64, df = 103, p = 7.6^{-14}), suggesting a moderate,

though significant structure in the similarity of effects amongst this set of MGP processes.

Comparison of processes to principal component directions

Almost a third of the 15 pairwise process comparisons showed a vector correlation > 0.5,

suggesting that many processes may feed into a limited set of directions in morphospace. To

assess the extent to which different processes affect the same aspects of facial shape we

randomly chose 1,000 process annotations, fit individual regularized PLS models to each set of

markers for a given annotation term, and then compared the direction of phenotypic effects for

each model to principal components 1-4 of the DO shape data (Fig 4B). Doing so with principal

components allows us to highlight similarities in directions of process effects. Process effects

range from completely uncorrelated to PC1 to highly correlated (0.0 - 0.8). However, the central

tendency of randomly selected process effects is one of moderate-to-high correlation with PC1.

Moderate correlations (0.55 - 0.6) with PC1 are more common than uncorrelated effects.

Supplemental table 1 contains the 10 most highly correlated processes with PC1 as well as the

corresponding correlations to PC 2-4. The most highly correlated process to PC1 is "zinc ion

binding", which is 0.86 correlated to the PC1 direction. The remaining 9 processes most highly

correlated to PC1 includes "sensory perception of sound", "Calcium ion transport", "Protein

homooligomerization", "Dendrite morphogenesis", "Neuropeptide signaling pathway", "Focal

adhesion", "Chromosome segregation", "Sarcomere organization", and "Integral component of

endoplasmic reticulum membrane".

Process correlations with PCs 2-4 are generally less strong. The maximum correlated

process with PCs 2-4 was "V(D)J recombination", "n-terminal protein myristoylation", and

"branching morphogenesis of an epithelial tube" with vector correlations of 0.72, 0.89, and 0.56,

respectively. Processes with high vector correlations for a given PC tend to be uncorrelated with

other PCs (Supp table 1), although some processes load moderately high across several PCs. For

example, "negative regulation of I-kappaB kinase/NF-kappaB signaling" shows vector

correlations between 0.25 - 0.54 for the first four PCs (Supp table 2).

Process effects in the mutant morphospace

To assess the extent to which craniofacial shape variation associated with developmental

processes aligns with variation from mutants of major effect, we projected 7 process effects onto

the first two principal components (PCs) of a dataset containing the DO sample, and samples

from 30 mutant genotypes (Fig 5A). Each black label represents the mean shape score of the

listed mutant genotype. The shaded ellipse with an orange border displays the 95% confidence

ellipse of PCs 1 and 2 of DO cranial shape variation. The DO mean shape is contrasted by the

mutant variation along PC1. The first PC describes vault size relative to the length of the face.

The phenotype shown along the x-axis of figure 5A depicts the maximum positive PC1 shape,

while the heatmap drawn on the crania represents the local deformations towards the minimum

PC1 shape. The positive direction of PC2 describes coordinated variation that includes a

relatively wider vault, narrower zygomatic, and shorter premaxilla (Fig 5A, y-axis margin).

Process effects—highlighted with orange vectors originating at the DO mean shape —

are necessarily of smaller magnitude than the total variation in the DO sample. Therefore, to

better compare the direction of process effects the vector magnitudes were magnified 4x. Several

process effects align in distinct directions of mutant effects, such as bmp signaling pathway and

endochondral ossification in the direction of *Shh*, *Nipbl*, and *Ift88* mutants. Neurotransmitter

transport and Wnt signaling pathway is similar in direction to *Mceph* and *B9d1* mutant effects.

Execution phase of apoptosis and intracellular transport both show similar effects to a cluster of

Bmp mutants.

Finally, we show the similarity of 30 process MGP effects to 30 mouse mutant models in

figure 5B. The heatmap shows the correlation in direction with yellow/green denoting higher

correlation and teal/blue denoting lower correlation. The bottom right of the heatmap

(highlighted by a white border) shows a block of mutants for which there are strong process

correlations. These are among the most extreme phenotypes along PC1 (Fig 5A) and include

mutants for *Nosip, Bmp2, Grm1, Bmp2; Bmp7* transheterozygote, *Bmp7, Ghrhr, Fgf10, and*

Papps2. The processes most strongly correlated to these mutants are histone methylation,

dendrite morphogenesis, chromosome segmentation, vasodilation, and fibroblast growth factor

binding.

There are a set of mutant phenotypes that have generally low correlations to the set of

processes chosen. These mutants include *Fgf3, Shh, Nipbl, Disp, Pten, Hhat, and Alk2; Alk3*

transheterozygote. Interestingly, this group of mutants vary more along PC2 than PC1 (Fig 5A).

Notably, regulation of intracellular protein transport and regulation of cell death are strongly

uncorrelated with the majority of mutant directions.

Real-time process GP mapping

Finally, we provide an online tool to visualize process effects and make comparisons to mutant

effects in real time. This application is found at **genopheno.ucalgary.ca/MGP** and can be used

for analyses similar to those described in this paper. When the user selects gene ontology terms,

the program searches for genotype markers adjacent to each gene listed and uses the selected

markers to fit a regularized PLS model. The result is an estimate of the many-to-many

relationship between the selected markers and cranial shape variation. The visual outputs include

barplots depicting the relative allele effect sizes for each gene in the process and a 3D plot of the

corresponding axis of shape variation. Users can compare the effects of different processes and

also compare process effects to mutant effects from a provided database of 30 mutant genotypes.

To illustrate how to use this application, we have provided the graphical user interface

used to select the parameters (Fig 6). As an example, in the "Process text" entry field, supply a

starting term; we chose "brain." The GO database is then filtered, returning a user-selectable

subset of biological process ontology annotation terms in the "Process filter" field. We chose

"forebrain morphogenesis," which has 11 associated genes. We chose to magnify the process

phenotype vectors 4x and compare the effect to a heterozygous *Ift88* mutant. *Ift88* is a core

component of the primary cilia, which are responsible for promoting developmental signals

involved in many facets of facial development (Tian et al., 2017). Further, the plots that are

generated are interactive. For example, marker loadings can be highlighted and subset by genes

of interest (Plotly, 2015). There is further information about using this online tool in the "About

this app" tab.

Discussion and Conclusion

A key goal in genomics is to create tractable genetic explanations for phenotypic variation. In

this study, we used a regularized PLS approach to model the joint effects of genomic markers on

multivariate craniofacial shape. This innovative approach allows us to address the joint

contributions of multiple genes that share ontological characteristic such as pathway membership

on craniofacial shape as a multivariate trait. Specifically, we chose markers adjacent to genes

annotated under a developmental process of interest. We showed two process MGP analyses in

depth, each with distinct phenotypic effects. The chondrocyte differentiation MGP effect mainly

showed effects on the shape of the zygomatic and jugal bones with *Ccn3/Nov* as the most highly

loaded corresponding marker effect. The FGF signaling MGP effect was less localized in

comparison to the chondrocyte differentiation effect. The FGF signaling phenotype was

primarily a smaller cranial vault volume with a longer facial outgrowth, broadly similar to the

primary axis of shape change during mouse growth and development (Gonzalez et al., 2013).

The most highly loaded markers for the FGF signaling effect were related to *Fgf2* and *Flrt2*. We

then compared process MGP phenotypic effects to each other, to mutant phenotypes, and the first

four principal components of the diversity outbred sample. Each of these comparisons

highlighted the integrated structure of phenotypic variation in mouse craniofacial shape. We

found that while there are processes with distinct and localized effects, genetic effects generally

converge on a limited set of directions in phenotype space. Further, these process effects often

correspond with the directions of major mutations known to affect these same processes.

Many recent studies have addressed the genetics of craniofacial shape in humans and

mice (reviewed in: Roosenboom et al., 2016; Weinberg et al., 2018). While these studies are

yielding a growing list of genes, suggesting that facial shape is highly polygenic, they have left

the vast majority of heritable variation unexplained. Existing studies have either used univariate

measures of facial shape such as linear measurements or univariate summaries of multivariate

shape (eg. Procrustes distances or PC scores). In addition, most genomic studies of craniofacial

shape quantify the effects of each genomic marker independently, with notable exceptions

focusing on epistatic effects (e.g. Varón-González et al., 2019). Our approach shares common

features with some predecessor GP mapping strategies in which candidate genes/SNPs are

selected a priori because of common involvement in a pathway (or other mechanistic cluster)

(Claes et al., 2014; Liu et al., 2012; Wang et al., 2010, 2007). In particular, Wang and colleagues

selected SNPs based on proximity to genes of interest and effect size to jointly model the

pathway-level effects on Parkinson disease data. Their approach is similar to gene-set

enrichment analysis, weighing over-representation of statistical effects related to case-control

group membership. In contrast, our approach focuses on estimating a multivariate continuous set

of craniofacial responses. Importantly, our approach jointly identifies genotype-phenotype axes

that maximally covary. This differs significantly from approaches that determine phenotypes for

analysis *a priori* or based on a pre-determined method of data reduction such as PCA. Our

approach also differs from methods that associate single locus effects with a multivariate

phenotype (Claes et al., 2018).

A key finding of our application of the MGP method to craniofacial shape is that

multivariate phenotypic variation aligns nonrandomly to genetic markers associated with

pathways or developmental processes. For most process MGP maps, multiple markers for gene

sets with known developmental relationships covary in their relationship with craniofacial shapes

(Supp fig 2). These covarying effects represent joint genetic effects of multiple contributors to

phenotypic variance. While these patterns of multivariate genotype-phenotype covariation may

include genetic variants that do not actually affect the phenotype, many others will be

contributors that we lack statistical power to detect under a typical univariate approach (Pitchers

et al., 2019; Varón-González et al., 2019). Here, the overall pattern of genotype-phenotype

covariance is the level of genetic explanation for phenotypic variation. When such patterns

involve genes that are ontologically linked in meaningful ways, they provide a level of insight

into the developmental-genetics of phenotypic variation that is beyond reach for most genome-

wide association studies for complex traits.

In aggregate, our results show substantial covariance in the directions of phenotypic

effects among different developmental processes (Figure 4A). The largest of these captures

processes general to development such as cell proliferation or pattern specification. The second

captures processes more specific to craniofacial development such as cranial suture/ossification

and neural tube patterning. While processes are structured in their effects, our data suggest that

many processes likely "add up" to produce variation. Thus, of 1000 randomly selected processes,

25.7% had a PC1 correlation higher than 0.6, supporting not only a highly polygenic model of

facial variation, but one in which hundreds or even thousands of developmental processes that

contribute to craniofacial variation. Importantly, this result shows how many processes and

pathways converge to produce central axes of variation in craniofacial shape.

The explicit modeling of multivariate relationships between phenotypes and genotypes

also allows a focus on pleiotropy. Developmental studies in mice demonstrate widespread

craniofacial morphological effects from localized developmental perturbations (Martínez-

Abadías et al., 2012; Stelzer et al., 2007; Young et al., 2010). Perturbations to specific processes

in development generally produce effects on multiple aspects of phenotype due to knock-on

effects at later stages or to interactions at the level of tissues or anatomical structures

(Hallgrimsson et al, 2007). A change in cartilage growth in basicranial synchondroses produces a

global change in craniofacial form, for example (Parsons et al, 2015). Remarkably, enhancers

with highly specific temporospatial effects on gene expression also produce global rather than

localized changes in craniofacial shape (Attanasio et al., 2013). Given that pleiotropy is likely

ubiquitous (Hill and Zhang, 2012; Wagner et al, 2008), explicitly multivariate approaches to

understanding genotype-phenotype maps are clearly needed.

This convergence of genetic effects on axes of covariation is also reflected in our finding

that mutations to major developmental genes produce effects that tend to align with the

directions of effect associated with the corresponding broader pathways or ontological groups.

Our analysis focused on two specific processes— FGF signaling and chondrocyte differentiation.

There was a strong correlation between the *Fgf10* mutant and the FGF signaling pathway effect,

while the *Alk6* mutant effect was moderately correlated to the chondrocyte differentiation

direction. For both process MGP maps, other mouse models in the same pathway showed

significant but weaker correlations in direction of effect. These results suggest that perturbations

that are developmentally similar tend to move the phenotype in the same direction in multivariate

space (Figure 5B). Even so, both mutational and higher-level pathway/process effects tend to

converge on a few directions of variation suggesting that multiple pathways and processes lead

to common developmental outcomes. This conclusion is further supported by our finding that the

genetic axes of covariance for individual processes/pathways can align with multiple directions

of mutational effect. For example, the process MGP phenotypes highlighted in the white

rectangle in figure 5B are all highly correlated with a set of BMP and growth hormone-related

mutants.

In some cases, mutants and MGP map directions do not correspond. There several

reasons why this can occur. The first is that the DO population may simply lack alleles as

deleterious as found in mutant lines. A small effect allele in the DO may not align with the

direction of a mutant almost completely lacking expression of the target gene. Further, there are

many examples where a mutation may have different and sometimes even opposite effects

depending on genetic background (Mackay, 2014; Percival et al., 2017). Mutations of major

effect may also differ in direction from variants in related genes that have smaller phenotypic

effects due to underlying nonlinearities in development (Green et al., 2017). Investigating how

variants in genes that are functionally related vary in phenotypic effect is an important avenue of

inquiry that is revealed by analyses such as those we have performed here. Additionally,

relationships between process and mutant effects may stimulate hypotheses about previously

unknown or unvalidated interactions between loci or pathways.

A second potential reason that MGP effects may not correspond to major mutation effects

is the use of only one PLS axis for each process analysis. With only one axis, we select the

phenotypic direction with greatest covariance with genetic marker variation. If there are multiple

large marker effects that do not covary, the weaker marker effect will be masked in the analysis.

For instance, there may be a PLS axis for "FGF signaling" that corresponds more strongly with

the *Fgf3* mutant phenotype (Supp fig 3). This phenomenon may be particularly prominent for

pathways with substantially different mutant effects, like FGF (Fig 5A).

Finally, our analysis shares the limitation of all approaches based on gene annotation

data. Incomplete annotation may contribute to lead to faulty or incomplete groupings of genes

when defining pathway/process hypotheses. Gene annotation is a huge undertaking, and there is

substantial variation in the completeness of different process annotations. Many process

annotations are manually assigned using inference from the literature, while most are a

combination of automated efforts based on transcript similarity and human curation (Mudge and

Harrow, 2015). Related to this, we assign gene annotation data to genetic markers based on the

closest protein-coding region. While this is a reasonable proxy, there will be regulatory sites that

affect genes other than the one immediately adjacent and this is a potential source of uncertainty

in our analysis.

The MGP method represents a deliberate decision to trade higher level insight from

genotype-phenotype association data at the expense of statistical certainty about the significance

of individual gene effects. The current implementation of the method also does not allow for

quantification of individual epistatic effects. Epistasis occurs when the genotypic trait value for a

locus is altered by the genotype of a different locus. Such effects generate nonlinear genotype

phenotype maps, but when considered genome-wide, contribute mainly to additive variance

(Cheverud and Routman, 1995; Hill, 2017). The MGP method is additive in that it models only

the linear effects of genes. However, since it captures the covariances among genotypic effects,

much of this "additive" variation is likely epistatic in origin.

Complex traits present a massive challenge in genomics because so many are turning out

to be enormously polygenic. To generate tractable explanations of the genetic basis for such

traits, methods are needed that extract higher-level representation of genotype-phenotype

relationships than those that emerge from single-locus focused approaches. Here, we present a

hypothesis-driven framework for deriving such higher-level genetic explanations for phenotypic

variation. Our approach leverages the biological tendency for developmental processes to

produce covariation among aspects of a multivariate phenotypic trait (Hallgrimsson et al., 2009;

Wagner et al., 2007). The underlying assumption in this approach is that there are latent variables

within high-dimensional genotype-phenotype data that correspond to developmental architecture.

We believe that analyses aimed at defining and characterizing such latent variables represent a

level of genetic explanation for phenotypic variation that is complementary to genetic analyses

designed to establish the significance of single locus effects. Pursuing such questions will help

bridge the gap between emerging mechanistic accounts of morphogenesis and our growing

understanding of the genetics of morphological variation.

Methods

DO mice

We use a sample ($n = 1,145$) of Diversity Outbred mice (DO; Jackson Laboratory, Bar Harbor,

ME) to map GP relationships for craniofacial shape (Churchill et al., 2012, 2004). The DO is a

multiparental outcross population derived from the eight founding lines of the Collaborative

Cross (CC). Each animal's genome is a unique mosaic of the genetic diversity found in the CC—

more than 45 million segregating SNPs (Collaborative Cross Consortium, 2012). Random

outcrossing over many DO generations maintains this diversity and, with recombination,

increases mapping resolution.

Our DO sample was sourced from three separate laboratories and seven DO generations.

386 are from the Jackson Laboratory (JAX), 287 from the University of North Carolina (UNC),

and 472 come from the Scripps Research Institute. Supplemental figure 4 shows the distribution

of the sample by lab source and generation of breeding. Imaging of mice at the University of

Calgary was performed under IACUC protocol AC13-0268.

Genotyping

Genotyping was performed by Neogen (Lincoln, NE). Ear clippings were used to extract DNA

for all samples. Mice from generations 9, 10, and 15 were genotyped using the MegaMUGA

genotyping array (77,808 markers); mice from generations 19, 21, 23, and 27 were genotyped

using the larger GigaMUGA array (143,259 markers) (Morgan et al., 2016). To pool the

genotype data from these two SNP arrays with differing numbers of markers, we imputed

markers between the two genotyping arrays using the "calc_genoprob" function in the qtl2

package (Broman et al., 2018). The function uses a hidden Markov model to estimate genotype

probabilities and missing genotype data (Gatti et al., 2014). After imputation, the merged genetic

dataset consists of 123,309 SNPs which vary among CC founders. Each animal's genetic record

is a 123,309*8 matrix of estimated diplotype contributions of each CC founder to each marker.

Scanning and landmarking

We used micro-computed tomography to acquire 3D scans of the full heads of the mice.

Scanning was done at the University of Calgary at .035 mm voxel resolution (Scanco vivaCT40).

One of us (WL) then acquired 54 3D landmarks (Fig 7) manually on each volume using Analyze

3D. A discussion of the error associated with manual landmarking can be found in Percival et al

(Percival et al., 2019).

Landmark registration

We symmetrized landmarks along the midline of the skull using Klingenberg et al.'s method for

object symmetry which configures landmark pairs into a common orientation with reflection and

subsequently removes variation associated with translation, scale, and rotation, using

Generalized Procrustes Analysis (Adams and Otárola-Castillo, 2013; Klingenberg et al., 2002;

Mardia, 2000; Schlager, 2017). To focus on shared, within-generation patterns in our

multigenerational DO sample, we regressed symmetric shape on DO generation, and used the

residual shapes with the grand mean added as the observations for analysis.

Genetic relatedness

Adjustment of phenotypes for the influence of genetic relatedness is a common approach in

genomic studies to prevent spurious associations. However, it is not necessary in all cases, such

as situations with low genetic relatedness and little variation in relatedness. We evaluated

whether accounting for genetic relatedness was important for our sample. To do so, we estimated

a kinship matrix based on DO genotype correlations (Cheng et al., 2013; Broman et al., 2019).

The kinship values in our sample have a mean of 0 and a standard deviation of .047 (Supp fig 5).

As a result of these findings, we performed all subsequent analyses on the within-generation

symmetric shape data, without an adjustment for relatedness.

Regularized PLS analysis

Multivariate genotype-phenotype methods for explicitly modeling multivariate phenotypes and

for overcoming the limitations of simple linear regression are increasingly common in mapping

studies. Claes et al. (Claes et al., 2018) used canonical correlation analysis to quantify individual

SNP effects for a multivariate measurement of facial shape. Each test returns a vector of the

linear combination of phenotypic effects that maximally correlates to the alleles at a given locus.

Mitteroecker et al. (Mitteroecker et al., 2016) developed a multivariate strategy around a singular

value decomposition (SVD) of GP covariation. Partial least squares (PLS) describes a family of

approaches that use SVD to decompose cross covariance matrices (Lee et al., 2011; Mitteroecker

and Gunz, 2009; Singh et al., 2016). PLS is increasingly used with large genetic datasets in order

to model how genomic effects extend to multiple traits (BJØRNSTAD et al., 2004; Mehmood et

al., 2011; Tyler et al., 2017). However, its implementation for MGP mapping is, thus far, much

more limited.

SVD decomposes the covariance matrix into three matrices:

$$Y = UDV'$$

Where $\mathbf{Y}$ is the mean-centered covariance matrix, $\mathbf{U}$ denotes the left singular vectors, a set of

vectors of unit length describing the relative weighting of each variable on each axis, and $\mathbf{D}$

denotes the variance along each axis. $\mathbf{V}$ denotes the set of right singular vectors. For a full

(square, symmetric) covariance matrix, $\mathbf{U}$ and $\mathbf{V}$ are identical, and the decomposition is

equivalent to PCA. For a non-symmetric matrix of covariances, i.e., one describing covariance

between two distinct blocks of traits, each successive column of $\mathbf{U}$ and $\mathbf{V}$ provide a pair of

singular vectors describing the best least squares approximation of covariance between the two

blocks, in order of greatest covariance explained to least.

PLS is most often used to find low-rank linear combinations that maximize covariance

between two sets of features. Here, we use a data-driven regularized PLS model implemented in

the mddsPLS package to find paired axes that maximize covariance between allelic and shape

variation (Lorenzo et al., 2019). The model uses a lasso penalty to minimize the coefficients

(loadings) towards zero to prevent overfitting (James et al., 2013). Overfitting can occur in when

many genotypic markers are included in the model, particularly when markers are colinear. The

genotype block is composed of the full set of DO founder probabilities for each selected marker.

Thus, an analysis of 20 markers would estimate 160 genotype coefficients. The phenotype block

consists of the full set of 54 3-dimensional landmarks (162 phenotype coefficients). In all

biological process analyses undertaken herein, we used a regularization parameter of 0.06 and

report only the first paired axes of the PLS model, i.e., the genotype and phenotype axes which

explain the most covariance.

Process-specific analysis

For process-specific MGP analyses, we used the mouse genome informatics database (Bult et al.,

2018) to identify genes annotated to a given process. Each annotation term has an associated GO

ID. For example, "chondrocyte differentiation" has GO ID GO:000206 (Fig 2, box 1). We cross-

reference the GO ID with the Ensemble genome database (GRCm38.p6) to find the name,

chromosome, and base pair start/end position for each gene (Fig 2, box 2) annotated to the

process. For genes with multiple splice variants, we select the full transcript start/end positions.

For each gene, we compare marker base pair positions and select the closest upstream and

downstream markers to the center of each gene. The 8-state genotype probability is then

calculated as the average founder allele probabilities between the two selected markers. (Fig 2,

box 3). After marker selection, we fit the regularized PLS model using the founder allele

probabilities (8 variables/marker) and full landmark data set (Fig 2, box 4).

We generate graphical displays of process results using the R packages ggplot2

(Wickham, 2016) and Morpho (Schlager, 2017). An example script to reproduce the analyses is

provided at **github.com/j0vid**.

Statistical results and comparisons

We estimate the magnitude and direction of MGP process effects using R^2 and vector

correlations, respectively. R^2 is calculated as the ratio of trace of the predicted model covariance

to the trace of the phenotypic covariance matrix. We contextualize the MGP process R^2 by

comparing it to the R^2 value of 1000 randomly drawn marker sets of the same size. For instance,

a process annotated with 40 genes would be compared to 1000 40-gene MGP analyses with

random markers selected in each iteration. Random marker selection for permutation is

constrained to follow similar patterns of linkage disequilibrium to the observed marker set of

interest. The null expectation in this scenario is that gene annotation does not provide better

information about coordinated marker effects than a randomly selected set of markers.

Vector correlations between process MGP effects are calculated by taking the Pearson

product-moment correlation of the two sets of process PLS1 phenotypic loadings. Vector

correlations between process effects and mutant effects are calculated by taking the correlation

between the process PLS1 phenotypic loadings and mutant MANOVA coefficients. The

MANOVA compares the mutant group phenotype with the DO sample specified as the reference

group.

Visualization tools

We introduce an interactive web application that allows the user to select processes of interest

with a graphical user interface and see the resulting craniofacial effect at

genopheno.ucalgary.ca/MGP. The web apps were written using the shiny package in R (Chang

et al., 2018). The application dynamically filters the MGI GO database based on the initial user

input. Queries will only list GO terms with exact matches. For example, "chond" will return GO

terms that incorporate either "chondrocyte" and "mitochondria".

Multiple queries can be selected. An analysis of "chondrocyte differentiation" and

"chondrocyte hypertrophy" will select the joint gene set of both processes. Processes with

different names can be jointly queried with the pipe operator "|", which is interpreted as an OR

(union) operator. For example, to generate the list of GO terms associated with either apoptosis

or WNT, we used the "apoptosis|WNT" query and selected the processes "Wnt signaling

pathway" and "execution phase of apoptosis" to perform the analysis on the joint set of

associated genes (Supp fig 6).

Several other parameters can be specified by the user including the type of plot to be

generated for the genetic loadings, the amount of magnification applied to the phenotype effect

vectors, the regularization parameter, and the option to overlay a mutant phenotype for

comparison. The comparative database currently includes craniofacial shape contrast data (wild-

type vs. mutant) for 30 mutant genotypes. If a mutant comparison is selected, the full set of DO

specimens are registered with the mutants added (with size removed). We then provide the

vector correlation between the process effect and the mutant effect (see Fig 6). The database also

includes PC1 of the DO sample for comparison.

The app enables users to save results. A save request will generate and download an

HTML report of the analysis which includes several versions of the genetic effect plot and an

interactive 3D model of the estimated phenotypic effect. If a mutant comparison is selected, it

will also appear in the report.

The application tracks recent searches by the user for their reference. A heatmap of

process vector correlations of the PLS phenotype loadings is also available under the "recent

searches" tab. The user can select between a heatmap of the processes in their search history or a

random assortment of process correlations from past, anonymous user searches.

Acknowledgements:

Grants: NIH - 2R01DE019638 to RM, BH and JC, NSERC 238992-17, CIHR Foundation grant to BH

and CFI grant #36262 to BH. JDA is supported by an Eyes High fellowship and an Alberta Children's

Hospital Research Institute scholarship.

References

Attanasio C, Nord AS, Zhu Y, Blow MJ, Li Z, Liberton DK, Morrison H, Plajzer-Frick I, Holt A, Hosseini R, Phouanenavong S, Akiyama JA, Shoukry M, Afzal V, Rubin EM, FitzPatrick DR, Ren B, Hallgrímsson B, Pennacchio LA, Visel A. 2013. Fine Tuning of Craniofacial Morphology by Distant-Acting Enhancers. *Science* **342**:1241006. doi:10.1126/science.1241006

Brazil DP, Church RH, Surae S, Godson C, Martin F. 2015. BMP signalling: agony and antagony in the family. *Trends Cell Biol* **25**:249–264. doi:10.1016/j.tcb.2014.12.004

Broman K, Sen Ś. (2009). A Guide to QTL Mapping with R/qtl. Springer. ISBN 978-0-387-92125-9. https://doi.org/10.1007/978-0-387-92125-9

Broman K, Gatti D, Simecek P, Furlotte N, Prins P, Sen Ś, Yandell B, Churchill G. (2018). R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multi-parent Populations Genetics 211(2), genetics.301595.2018. https://dx.doi.org/10.1534/genetics.118.301595

Chai Y. 2015. Craniofacial Development. *Current Topics in Dev Bio.* Academic Press. Volume 115. ISBN 9780124081413.

Chebib J, Guillaume F. 2017. What affects the predictability of evolutionary constraints using a G-matrix? The relative effects of modular pleiotropy and mutational correlation. *Evol Int J Org Evol* **71**:2298–2312. doi:10.1111/evo.13320

Cheverud JM, Routman EJ. 1996. EPISTASIS AS A SOURCE OF INCREASED ADDITIVE GENETIC VARIANCE AT POPULATION BOTTLENECKS. *Evolution* **50**:1042–1051. doi:10.1111/j.1558-5646.1996.tb02345.x

Churchill GA, Gatti DM, Munger SC, Svenson KL. 2012. The diversity outbred mouse population. *Mamm Genome* **23**:713–718. doi:10.1007/s00335-012-9414-2

Claes P, Liberton DK, Daniels K, Rosana KM, Quillen EE, Pearson LN, McEvoy B, Bauchet M, Zaidi AA, Yao W, Tang H, Barsh GS, Absher DM, Puts DA, Rocha J, Beleza S, Pereira RW, Baynam G, Suetens P, Vandermeulen D, Wagner JK, Boster JS, Shriver MD. 2014. Modeling 3D Facial Shape from DNA. *Plos Genet* **10**:e1004224. doi:10.1371/journal.pgen.1004224

Claes, P., Roosenboom, J., White, J. D., Swigut, T., Sero, D., Li, J., Lee, M. K., Zaidi, A., Mattern, B. C., Liebowitz, C., Pearson, L., González, T., Leslie, E. J., Carlson, J. C., Orlova, E., Suetens, P., Vandermeulen, D., Feingold, E., Marazita, M. L., Shaffer, J. R., Wysocka, J., Shriver, M. D., & Weinberg, S. M. (2018). Genome-wide mapping of global-to-local genetic effects on human facial shape. *Nature Genetics,* **50**(3), 414-423. doi:10.1038/s41588-018-0057-4

Cole, J. B., Manyama, M., Kimwaga, E., Mathayo, J., Larson, J. R., Liberton, D. K., Lukowiak,
    K., Ferrara, T. M., Riccardi, S. L., Li, M., Mio, W., Prochazkova, M., Williams, T., Li, H.,
    Jones, K. L., Klein, O. D., Santorico, S. A., Hallgrimsson, B., & Spritz, R. A. (2016).
    Genomewide Association Study of African Children Identifies Association of SCHIP1 and
    PDE8A with Facial Size and Shape. *PLoS Genetics*, **12**(8), e1006174.
    doi:10.1371/journal.pgen.1006174

Cole JB, Manyama M, Larson JR, Liberton DK, Ferrara TM, Riccardi SL, Li M, Mio W, Klein
    OD, Santorico SA, Hallgrímsson B, Spritz RA. 2017. Human Facial Shape and Size
    Heritability and Genetic Correlations. *Genetics* **205**:967–978.
    doi:10.1534/genetics.116.193185

Collaborative Cross Consortium. 2012. The Genome Architecture of the Collaborative Cross
    Mouse Genetic Reference Population. *Genetics* **190**:389–401.
    doi:10.1534/genetics.111.132639

Dudbridge F. 2013. Power and Predictive Accuracy of Polygenic Risk Scores. *Plos Genet*
    9:e1003348. doi:10.1371/journal.pgen.1003348

FANTOM Consortium, Forrest ARR, Kawaji H, Rehli M, Baillie JK, Hoon MJL de, Haberle V,
    Lassmann T, Kulakovskiy IV, Lizio M, Itoh M, Andersson R, Mungall CJ, Meehan TF,
    Schmeier S, Bertin N, Jørgensen M, Dimont E, Arner E, Schmidl C, Schaefer U, Medvedeva
    YA, Plessy C, Vitezic M, Severin J, Semple CA, Ishizu Y, Young RS, Francescatto M, Alam
    I, Albanese D, Altschuler GM, Arakawa T, Archer JAC, Arner P, Babina M, Rennie S,
    Balwierz PJ, Beckhouse AG, Pradhan-Bhatt S, Blake JA, Blumenthal A, Bodega B, Bonetti
    A, Briggs J, Brombacher F, Burroughs AM, Califano A, Cannistraci CV, Carbajo D, Chen Y,
    Chierici M, Ciani Y, Clevers HC, Dalla E, Davis CA, Detmar M, Diehl AD, Dohi T, Drabløs
    F, Edge ASB, Edinger M, Ekwall K, Endoh M, Enomoto H, Fagiolini M, Fairbairn L, Fang
    H, Farach-Carson MC, Faulkner GJ, Favorov AV, Fisher ME, Frith MC, Fujita R, Fukuda S,
    Furlanello C, Furuno M, Furusawa J, Geijtenbeek TB, Gibson AP, Gingeras T, Goldowitz D,
    Gough J, Guhl S, Guler R, Gustincich S, Ha TJ, Hamaguchi M, Hara M, Harbers M,
    Harshbarger J, Hasegawa A, Hasegawa Y, Hashimoto T, Herlyn M, Hitchens KJ, Sui SJH,
    Hofmann OM, Hoof I, Hori F, Huminiecki L, Iida K, Ikawa T, Jankovic BR, Jia H, Joshi A,
    Jurman G, Kaczkowski B, Kai C, Kaida K, Kaiho A, Kajiyama K, Kanamori-Katayama M,
    Kasianov AS, Kasukawa T, Katayama S, Kato S, Kawaguchi S, Kawamoto H, Kawamura YI,
    Kawashima T, Kempfle JS, Kenna TJ, Kere J, Khachigian LM, Kitamura T, Klinken SP,
    Knox AJ, Kojima M, Kojima S, Kondo N, Koseki H, Koyasu S, Krampitz S, Kubosaki A,
    Kwon AT, Laros JFJ, Lee W, Lennartsson A, Li K, Lilje B, Lipovich L, Mackay-sim A,
    Manabe R, Mar JC, Marchand B, Mathelier A, Mejhert N, Meynert A, Mizuno Y, Morais DA
    de L, Morikawa H, Morimoto M, Moro K, Motakis E, Motohashi H, Mummery CL, Murata
    M, Nagao-Sato S, Nakachi Y, Nakahara F, Nakamura T, Nakamura Y, Nakazato K,
    Nimwegen E van, Ninomiya N, Nishiyori H, Noma S, Nozaki T, Ogishima S, Ohkura N,
    Ohmiya H, Ohno H, Ohshima M, Okada-Hatakeyama M, Okazaki Y, Orlando V,
    Ovchinnikov DA, Pain A, Passier R, Patrikakis M, Persson H, Piazza S, Prendergast JGD,
    Rackham OJL, Ramilowski JA, Rashid M, Ravasi T, Rizzu P, Roncador M, Roy S, Rye MB,
    Saijyo E, Sajantila A, Saka A, Sakaguchi S, Sakai M, Sato H, Satoh H, Savvi S, Saxena A,

Schneider C, Schultes EA, Schulze-Tanzil GG, Schwegmann A, Sengstag T, Sheng G,
Shimoji H, Shimoni Y, Shin JW, Simon C, Sugiyama D, Sugiyama T, Suzuki M, Suzuki N,
Swoboda RK, Hoen PAC 't, Tagami M, Takahashi N, Takai J, Tanaka H, Tatsukawa H,
Tatum Z, Thompson M, Toyoda H, Toyoda T, Valen E, Wetering M van de, Berg LM van
den, Verardo R, Vijayan D, Vorontsov IE, Wasserman WW, Watanabe S, Wells CA,
Winteringham LN, Wolvetang E, Wood EJ, Yamaguchi Y, Yamamoto M, Yoneda M,
Yonekura Y, Yoshida S, Zabierowski SE, Zhang PG, Zhao X, Zucchelli S, Summers KM,
Suzuki H, Daub CO, Kawai J, Heutink P, Hide W, Freeman TC, Lenhard B, Bajic VB, Taylor
MS, Makeev VJ, Sandelin A, Hume DA, Carninci P, Hayashizaki Y. 2014. A promoter-level
mammalian expression atlas. *Nature* **507**:462. doi:10.1038/nature13182

Finger JH, Smith CM, Hayamizu TF, McCright IJ, Xu J, Law M, Shaw DR, Baldarelli RM, Beal
JS, Blodgett O, Campbell JW, Corbani LE, Lewis JR, Forthofer KL, Frost PJ, Giannatto SC,
Hutchins LN, Miers DB, Motenko H, Stone KR, Eppig JT, Kadin JA, Richardson JE,
Ringwald M. 2017. The mouse Gene Expression Database (GXD): 2017 update. *Nucleic*
Acids Res **45**:D730–D736. doi:10.1093/nar/gkw1073

Francis-West P, Crespo-Enriquez I. 2016. Vertebrate Embryo: Craniofacial Development. *eLS*
1–15. doi:10.1002/9780470015902.a0026602

Gasch AP, Payseur BA, Pool JE. 2016. The Power of Natural Variation for Model Organism
Biology. *Trends Genet* **32**:147–154. doi:10.1016/j.tig.2015.12.003

Gatti DM, Svenson KL, Shabalin A, Wu L-Y, Valdar W, Simecek P, Goodwin N, Cheng R,
Pomp D, Palmer A, Chesler EJ, Broman KW, Churchill GA. 2014. Quantitative Trait Locus
Mapping Methods for Diversity Outbred Mice. *G3 Genes Genomes Genetics* **4**:1623–1633.
doi:10.1534/g3.114.013748

Gonzalez, PN, Kristensen, E, Morck, DW, Boyd, S, Hallgrímsson, B. 2013. Effects of growth
hormone on the ontogenetic allometry of craniofacial bones. *Evolution & Development* **15**:
133-145. doi:10.1111/ede.12025

Green RM, Fish JL, Young NM, Smith FJ, Roberts B, Dolan K, Choi I, Leach CL, Gordon P,
Cheverud JM, Roseman CC, Williams TJ, Marcucio RS, Hallgrímsson B. 2017.
Developmental nonlinearity drives phenotypic robustness. *Nat Commun* **8**:1970.
doi:10.1038/s41467-017-02037-7

Hallgrimsson, B, Lieberman, DE, Liu, W, Ford-Hutchinson, AF, Jirik, FR. 2007. Epigenetic
interactions and the structure of phenotypic variation in the cranium. *Evol Dev* **9**(1), 76-91.

Hallgrimsson B, Green RM, Katz DC, Fish JL, Bernier FP, Roseman CC, Young NM, Cheverud
JM, Marcucio RS. 2018. The developmental-genetics of canalization. *Semin Cell Dev Biol*.
doi:10.1016/j.semcdb.2018.05.019

Hallgrímsson B, Jamniczky H, Young NM, Rolian C, Parsons TE, Boughner JC, Marcucio RS.
2009. Deciphering the Palimpsest: Studying the Relationship Between Morphological

Integration and Phenotypic Covariation. *Evol Biol* **36**:355–376. doi:10.1007/s11692-009-
9076-5

Hallgrímsson B, Lieberman DE. 2008. Mouse models and the evolutionary developmental
biology of the skull. *Integr Comp Biol* **48**:373–384. doi:10.1093/icb/icn076

Hallgrimsson B, Mio W, Marcucio RS, Spritz R. 2014. Let's Face It—Complex Traits Are Just
Not That Simple. *Plos Genet* **10**:e1004724. doi:10.1371/journal.pgen.1004724

Hilliard SA, Yu L, Gu S, Zhang Z, Chen YP. 2005. Regional regulation of palatal growth and
patterning along the anterior–posterior axis in mice. *J Anat* **207**:655–667. doi:10.1111/j.1469-
7580.2005.00474.x

Houle D, Fierst J. 2012. Properties of spontaneous mutational variance and covariance for wing
size and shape in Drosophila melanogaster. *Evol Int J Org Evol* **67**:1116–30.
doi:10.1111/j.1558-5646.2012.01838.x

Houle D, Govindaraju DR, Omholt S. 2010. Phenomics: the next challenge. *Nat Rev Genet*
11:855. doi:10.1038/nrg2897

Jin Y-R, Han XH, Taketo MM, Yoon JK. 2012. Wnt9b-dependent FGF signaling is crucial for
outgrowth of the nasal and maxillary processes during upper jaw and lip development.
Development **139**:1821–1830. doi:10.1242/dev.075796

Johannsdottir B, Thorarinsson F, Thordarson A, Magnusson TE. 2005. Heritability of
craniofacial characteristics between parents and offspring estimated from lateral
cephalograms. *Am J Orthod Dentofac* **127**:200–207. doi:10.1016/j.ajodo.2004.07.033

Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. 2017. KEGG: new perspectives on
genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**:D353–D361.
doi:10.1093/nar/gkw1092

Katz DC, Aponte JD, Liu W, Green RM, Mayeux JM, Pollard KM, Pomp D, Munger SC,
Murray SA, Roseman CC, Percival CJ, Cheverud J, Marcucio RS, Hallgrímsson B. 2019.
Facial shape and allometry quantitative trait locus intervals in the Diversity Outbred mouse
are enriched for known skeletal and facial development genes. *Biorxiv* 787291.
doi:10.1101/787291

Kawauchi S, Calof AL, Santos R, Lopez-Burks ME, Young CM, Hoang MP, Chua A, Lao T,
Lechner MS, Daniel JA, Nussenzweig A, Kitzes L, Yokomori K, Hallgrimsson B, Lander
AD. 2009. Multiple Organ System Defects and Transcriptional Dysregulation in the Nipbl+/−
Mouse, a Model of Cornelia de Lange Syndrome. *Plos Genet* **5**:e1000650.
doi:10.1371/journal.pgen.1000650

Kenney-Hunt JP, Wang B, Norgard EA, Fawcett G, Falk D, Pletscher LS, Jarvis JP, Roseman C,
Wolf J, Cheverud JM. 2008. Pleiotropic Patterns of Quantitative Trait Loci for 70 Murine
Skeletal Traits. *Genetics* **178**:2275–2288. doi:10.1534/genetics.107.084434

Klingenberg CP, Leamy LJ. 2001. QUANTITATIVE GENETICS OF GEOMETRIC SHAPE IN
THE MOUSE MANDIBLE. *Evolution* **55**:2342–2352. doi:10.1554/0014-
3820(2001)055[2342:qgogsi]2.0.co;2

Lewontin R. 1974. The Genetic Basis of Evolutionary Change. New York, Columbia University
Press, 1974.

Liu F, Lijn F van der, Schurmann C, Zhu G, Chakravarty MM, Hysi PG, Wollstein A, Lao O,
Bruijne M de, Ikram MA, Lugt A van der, Rivadeneira F, Uiterlinden AG, Hofman A,
Niessen WJ, Homuth G, Zubicaray G de, McMahon KL, Thompson PM, Daboul A, Puls R,
Hegenscheid K, Bevan L, Pausova Z, Medland SE, Montgomery GW, Wright MJ, Wicking
C, Boehringer S, Spector TD, Paus T, Martin NG, Biffar R, Kayser M. 2012. A Genome-
Wide Association Study Identifies Five Loci Influencing Facial Morphology in Europeans.
Plos Genet **8**:e1002932. doi:10.1371/journal.pgen.1002932

Liu W, Sun X, Braut A, Mishina Y, Behringer RR, Mina M, Martin JF. 2005. Distinct functions
for Bmp signaling in lip and palate fusion in mice. *Development* **132**:1453–1461.
doi:10.1242/dev.01676

Lorenzo H, Saracco J, Thiébaut R. 2019. Supervised Learning for Multi-Block Incomplete Data.
arXiv preprint *arXiv*:1901.04380. http://arxiv.org/abs/1901.04380

Mackay, T. Epistasis and quantitative traits: using model organisms to study gene–gene
interactions.*Nat Rev Genet* **15,** 22–33 (2014). https://doi.org/10.1038/nrg3627

Mackay TF, Moore JH. 2014. Why epistasis is important for tackling complex human disease
genetics. *Genome Med* **6**:42. doi:10.1186/gm561

Maga AM, Navarro N, Cunningham ML, Cox TC. 2015. Quantitative trait loci affecting the 3D
skull shape and size in mouse and prioritization of candidate genes in-silico. *Front Physiol*
6:92. doi:10.3389/fphys.2015.00092

Martínez-Abadías N, Mitteroecker P, Parsons TE, Esparza M, Sjøvold T, Rolian C, Richtsmeier
JT, Hallgrímsson B. 2012. The Developmental Basis of Quantitative Craniofacial Variation in
Humans and Mice. *Evol Biol* **39**:554–567. doi:10.1007/s11692-012-9210-7

Mattick JS, Taft RJ, Faulkner GJ. 2010. A global view of genomic information – moving beyond
the gene and the master regulator. *Trends Genet* **26**:21–28. doi:10.1016/j.tig.2009.11.002

Mitteroecker P, Cheverud JM, Pavlicev M. 2016. Multivariate Analysis of Genotype–Phenotype
Association. *Genetics* **202**:1345–1363. doi:10.1534/genetics.115.181339

Morozova TV, Huang W, Pray VA, Whitham T, Anholt RRH, Mackay TFC. 2015.
Polymorphisms in early neurodevelopmental genes affect natural variation in alcohol
sensitivity in adult drosophila. *Bmc Genomics* **16**:865. doi:10.1186/s12864-015-2064-5

Mudge JM, Harrow J. 2015. Creating reference gene annotation for the mouse C57BL6/J
genome assembly. *Mamm Genome* **26**:366–378. doi:10.1007/s00335-015-9583-x

Nakata M. 1985. Twin Studies in Craniofacial Genetics: A Review. *Amg Acta Geneticae*
Medicae Et Gemellologiae Twin Res **34**:1–14. doi:10.1017/s0001566000004876

Ornitz DM, Marie PJ. 2015. Fibroblast growth factor signaling in skeletal development and
disease. *Gene Dev* **29**:1463–1486. doi:10.1101/gad.266551.115

Pallares LF, Carbonetto P, Gopalakrishnan S, Parker CC, Ackert-Bicknell CL, Palmer AA, Tautz
D. 2015. Mapping of Craniofacial Traits in Outbred Mice Identifies Major Developmental
Genes Involved in Shape Determination. *Plos Genet* **11**:e1005607.
doi:10.1371/journal.pgen.1005607

Pallares LF, Harr B, Turner LM, Tautz D. 2014. Use of a natural hybrid zone for genomewide
association mapping of craniofacial traits in the house mouse. *Mol Ecol* **23**:5756–5770.
doi:10.1111/mec.12968

Parsons, T. E., Downey, C. M., Jirik, F. R., Hallgrimsson, B., & Jamniczky, H. A. (2015). Mind
the Gap: Genetic Manipulation of Basicranial Growth within Synchondroses Modulates
Calvarial and Facial Shape in Mice through Epigenetic Interactions. *PloS One, 10*(2),
e0118355. doi:10.1371/journal.pone.0118355

Percival CJ, Marangoni P, Tapaltsyan V, Klein O, Hallgrímsson B. 2017. The Interaction of
Genetic Background and Mutational Effects in Regulation of Mouse Craniofacial Shape. *G3*
Genes Genomes Genetics **7**:1439–1450. doi:10.1534/g3.117.040659

Percival CJ, Richtsmeier JT. 2013. Angiogenesis and intramembranous osteogenesis. *Dev*
Dynam **242**:909–922. doi:10.1002/dvdy.23992

Pitchers W, Nye J, Márquez EJ, Kowalski A, Dworkin I, Houle D. 2019. A Multivariate
Genome-Wide Association Study of Wing Shape in Drosophila melanogaster. *Genetics*
211:genetics.301342.2018. doi:10.1534/genetics.118.301342

Plotly Technologies Inc. Collaborative data science. Montréal, QC, 2015. https://plot.ly.

Prochazkova M, Prochazka J, Marangoni P, Klein OD. 2018. Bones, Glands, Ears and More:
The Multiple Roles of FGF10 in Craniofacial Development. *Frontiers Genetics* **9**:542.
doi:10.3389/fgene.2018.00542

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Reed LK, Baer CF, Edison AS. 2017. Considerations when choosing a genetic model organism
for metabolomics studies. *Curr Opin Chem Biol* **36**:7–14. doi:10.1016/j.cbpa.2016.12.005

Rendel, J. M. (1967). *Canalization and gene control*. London: Logos Press.

Richman JM, Herbert M, Matovinovic E, Walin J. 1997. Effect of Fibroblast Growth Factors on
Outgrowth of Facial Mesenchyme. *Dev Biol* **189**:135–147. doi:10.1006/dbio.1997.8656

Richtsmeier JT, Flaherty K. 2013. Hand in glove: brain and skull in development and
dysmorphogenesis. *Acta Neuropathol* **125**:469–489. doi:10.1007/s00401-013-1104-y

Rie D de, Abugessaisa I, Alam T, Arner E, Arner P, Ashoor H, Åström G, Babina M, Bertin N,
Burroughs AM, Carlisle AJ, Daub CO, Detmar M, Deviatiiarov R, Fort A, Gebhard C,
Goldowitz D, Guhl S, Ha TJ, Harshbarger J, Hasegawa A, Hashimoto K, Herlyn M, Heutink
P, Hitchens KJ, Hon CC, Huang E, Ishizu Y, Kai C, Kasukawa T, Klinken P, Lassmann T,
Lecellier C-H, Lee W, Lizio M, Makeev V, Mathelier A, Medvedeva YA, Mejhert N,
Mungall CJ, Noma S, Ohshima M, Okada-Hatakeyama M, Persson H, Rizzu P, Roudnicky F,
Sætrom P, Sato H, Severin J, Shin JW, Swoboda RK, Tarui H, Toyoda H, Vitting-Seerup K,
Winteringham L, Yamaguchi Y, Yasuzawa K, Yoneda M, Yumoto N, Zabierowski S, Zhang
PG, Wells CA, Summers KM, Kawaji H, Sandelin A, Rehli M, Consortium TF, Hayashizaki
Y, Carninci P, Forrest ARR, Hoon MJL de. 2017. An integrated expression atlas of miRNAs
and their promoters in human and mouse. *Nat Biotechnol* **35**:nbt.3947. doi:10.1038/nbt.3947

Roosenboom J, Hens G, Mattern BC, Shriver MD, Claes P. 2016. Exploring the Underlying
Genetics of Craniofacial Morphology through Various Sources of Knowledge. *Biomed Res*
Int **2016**:1–9. doi:10.1155/2016/3054578

Shaffer, J. R., Orlova, E., Lee, M. K., Leslie, E. J., Raffensperger, Z. D., Heike, C. L.,
Cunningham, M. L., Hecht, J. T., Kau, C. H., Nidey, N. L., Moreno, L. M., Wehby, G. L.,
Murray, J. C., Laurie, C. A., Laurie, C. C., Cole, J., Ferrara, T., Santorico, S., Klein, O., Mio,
W., Feingold, E., Hallgrimsson, B., Spritz, R. A., Marazita, M. L., & Weinberg, S. M. (2016).
Genome-Wide Association Study Reveals Multiple Loci Influencing Normal Human Facial
Morphology. *PLoS Genetics*, **12**(8), e1006149. doi:10.1371/journal.pgen.1006149

Schoenfelder S, Javierre B-M, Furlan-Magaril M, Wingett SW, Fraser P. 2018. Promoter
Capture Hi-C: High-resolution, Genome-wide Profiling of Promoter Interactions. *J Vis Exp*
Jove 57320. doi:10.3791/57320

Sprinzak E, Margalit H. 2001. Correlated sequence-signatures as markers of protein-protein
interaction1 1Edited by G. von Heijne. *J Mol Biol* **311**:681–692. doi:10.1006/jmbi.2001.4920

Stelzer C, Brimmer A, Hermanns P, Zabel B, Dietz UH. 2007. Expression profile of Papss2 (3′-
phosphoadenosine 5′-phosphosulfate synthase 2) during cartilage formation and skeletal
development in the mouse embryo. *Dev Dynam* **236**:1313–1318. doi:10.1002/dvdy.21137

Tian H, Feng J, Li J, Ho T-V, Yuan Y, Liu Y, Brindopke F, Figueiredo JC, Magee W, Sanchez-
Lara PA, Chai Y. 2017. Intraflagellar transport 88 (IFT88) is crucial for craniofacial
development in mice and is a candidate gene for human cleft lip and palate. *Hum Mol Genet*
ddx002. doi:10.1093/hmg/ddx002

Tsagkrasoulis D, Hysi P, Spector T, Montana G. 2017. Heritability maps of human face
morphology through large-scale automated three-dimensional phenotyping. *Sci Rep-uk*
7:45885. doi:10.1038/srep45885

Uller T, Moczek AP, Watson RA, Brakefield PM, Laland KN. 2018. Developmental Bias and
Evolution: A Regulatory Network Perspective. *Genetics* **209**:949–966.
doi:10.1534/genetics.118.300995

Varón-González C, Pallares LF, Debat V, Navarro N. 2019. Mouse Skull Mean Shape and Shape
Robustness Rely on Different Genetic Architectures and Different Loci. *Frontiers Genetics*
10:64. doi:10.3389/fgene.2019.00064

Vidal M, Cusick ME, Barabási A-L. 2011. Interactome Networks and Human Disease. *Cell*
144:986–998. doi:10.1016/j.cell.2011.02.016

Visscher PM. 2008. Sizing up human height variation. *Nat Genet* **40**:ng0508-489.
doi:10.1038/ng0508-489

Visscher PM, Wray NR, Zhang Q, Sklar P, McCarthy MI, Brown MA, Yang J. (2017). 10 Years
of GWAS Discovery: Biology, Function, and Translation. *Am J Hum Genetics.* **101**:1:5-22.
https://doi.org/10.1016/j.ajhg.2017.06.005

Wagner, G., Pavlicev, M., Cheverud, J. (2007). The road to modularity. *Nat Rev Genet* **8**(12),
nrg2267. https://dx.doi.org/10.1038/nrg2267

Wagner GP, Zhang J. 2011. The pleiotropic structure of the genotype–phenotype map: the
evolvability of complex organisms. *Nat Rev Genet* **12**:204. doi:10.1038/nrg2949

Wang K, Li M, Bucan M. 2007. Pathway-Based Approaches for Analysis of Genomewide
Association Studies. *Am J Hum Genetics* **81**:1278–1283. doi:10.1086/522374

Wang K, Li M, Hakonarson H. 2010. Analysing biological pathways in genome-wide association
studies. *Nat Rev Genet* **11**:843. doi:10.1038/nrg2884

Watson J, Francavilla C. 2018. Regulation of FGF10 Signaling in Development and Disease.
Frontiers Genetics **9**:500. doi:10.3389/fgene.2018.00500

Weinberg SM, Cornell R, Leslie EJ. 2018. Craniofacial genetics: Where have we been and where
are we going? *Plos Genet* **14**:e1007438. doi:10.1371/journal.pgen.1007438

Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S, Chu AY, Estrada K, Luan J,
Kutalik Z, Amin N, Buchkovich ML, Croteau-Chonka DC, Day FR, Duan Y, Fall T,
Fehrmann R, Ferreira T, Jackson AU, Karjalainen J, Lo KS, Locke AE, Mägi R, Mihailov E,
Porcu E, Randall JC, Scherag A, Vinkhuyzen AAE, Westra H-J, Winkler TW, Workalemahu
T, Zhao JH, Absher D, Albrecht E, Anderson D, Baron J, Beekman M, Demirkan A, Ehret
GB, Feenstra B, Feitosa MF, Fischer K, Fraser RM, Goel A, Gong J, Justice AE, Kanoni S,
Kleber ME, Kristiansson K, Lim U, Lotay V, Lui JC, Mangino M, Leach IM, Medina-Gomez
C, Nalls MA, Nyholt DR, Palmer CD, Pasko D, Pechlivanis S, Prokopenko I, Ried JS, Ripke
S, Shungin D, Stancáková A, Strawbridge RJ, Sung YJ, Tanaka T, Teumer A, Trompet S,
Laan SW van der, Setten J van, Vliet-Ostaptchouk JVV, Wang Z, Yengo L, Zhang W, Afzal
U, Ärnlöv J, Arscott GM, Bandinelli S, Barrett A, Bellis C, Bennett AJ, Berne C, Blüher M,
Bolton JL, Böttcher Y, Boyd HA, Bruinenberg M, Buckley BM, Buyske S, Caspersen IH,
Chines PS, Clarke R, Claudi-Boehm S, Cooper M, Daw EW, Jong PAD, Deelen J, Delgado
G, Denny JC, Dhonukshe-Rutten R, Dimitriou M, Doney ASF, Dörr M, Eklund N, Eury E,
Folkersen L, Garcia ME, Geller F, Giedraitis V, Go AS, Grallert H, Grammer TB, Gräßler J,
Grönberg H, Groot LCPGM de, Groves CJ, Haessler J, Hall P, Haller T, Hallmans G,
Hannemann A, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hemani
G, Henders AK, Hillege HL, Hlatky MA, Hoffmann W, Hoffmann P, Holmen O, Houwing-
Duistermaat JJ, Illig T, Isaacs A, James AL, Jeff J, Johansen B, Johansson Å, Jolley J,
Juliusdottir T, Junttila J, Kho AN, Kinnunen L, Klopp N, Kocher T, Kratzer W, Lichtner P,
Lind L, Lindström J, Lobbens S, Lorentzon M, Lu Y, Lyssenko V, Magnusson PKE, Mahajan
A, Maillard M, McArdle WL, McKenzie CA, McLachlan S, McLaren PJ, Menni C, Merger
S, Milani L, Moayyeri A, Monda KL, Morken MA, Müller G, Müller-Nurasyid M, Musk
AW, Narisu N, Nauck M, Nolte IM, Nöthen MM, Oozageer L, Pilz S, Rayner NW, Renstrom
F, Robertson NR, Rose LM, Roussel R, Sanna S, Scharnagl H, Scholtens S, Schumacher FR,
Schunkert H, Scott RA, Sehmi J, Seufferlein T, Shi J, Silventoinen K, Smit JH, Smith AV,
Smolonska J, Stanton AV, Stirrups K, Stott DJ, Stringham HM, Sundström J, Swertz MA,
Syvänen A-C, Tayo BO, Thorleifsson G, Tyrer JP, Dijk S van, Schoor NM van, Velde N van
der, Heemst D van, Oort FVA van, Vermeulen SH, Verweij N, Vonk JM, Waite LL,
Waldenberger M, Wennauer R, Wilkens LR, Willenborg C, Wilsgaard T, Wojczynski MK,
Wong A, Wright AF, Zhang Q, Arveiler D, Bakker SJL, Beilby J, Bergman RN, Bergmann S,
Biffar R, Blangero J, Boomsma DI, Bornstein SR, Bovet P, Brambilla P, Brown MJ,
Campbell H, Caulfield MJ, Chakravarti A, Collins R, Collins FS, Crawford DC, Cupples LA,
Danesh J, Faire U de, Ruijter HM den, Erbel R, Erdmann J, Eriksson JG, Farrall M,
Ferrannini E, Ferrières J, Ford I, Forouhi NG, Forrester T, Gansevoort RT, Gejman PV,
Gieger C, Golay A, Gottesman O, Gudnason V, Gyllensten U, Haas DW, Hall AS, Harris TB,
Hattersley AT, Heath AC, Hengstenberg C, Hicks AA, Hindorff LA, Hingorani AD, Hofman
A, Hovingh GK, Humphries SE, Hunt SC, Hypponen E, Jacobs KB, Jarvelin M-R, Jousilahti
P, Jula AM, Kaprio J, Kastelein JJP, Kayser M, Kee F, Keinanen-Kiukaanniemi SM,
Kiemeney LA, Kooner JS, Kooperberg C, Koskinen S, Kovacs P, Kraja AT, Kumari M,
Kuusisto J, Lakka TA, Langenberg C, Marchand LL, Lehtimäki T, Lupoli S, Madden PAF,
Männistö S, Manunta P, Marette A, Matise TC, McKnight B, Meitinger T, Moll FL,
Montgomery GW, Morris AD, Morris AP, Murray JC, Nelis M, Ohlsson C, Oldehinkel AJ,
Ong KK, Ouwehand WH, Pasterkamp G, Peters A, Pramstaller PP, Price JF, Qi L, Raitakari
OT, Rankinen T, Rao DC, Rice TK, Ritchie M, Rudan I, Salomaa V, Samani NJ, Saramies J,
Sarzynski MA, Schwarz PEH, Sebert S, Sever P, Shuldiner AR, Sinisalo J, Steinthorsdottir V,

Stolk RP, Tardif J-C, Tönjes A, Tremblay A, Tremoli E, Virtamo J, Vohl M-C, Consortium TEMR and G (eMERGE), Consortium TMig, Consortium TP, Study TLC, Amouyel P, Asselbergs FW, Assimes TL, Bochud M, Boehm BO, Boerwinkle E, Bottinger EP, Bouchard C, Cauchi S, Chambers JC, Chanock SJ, Cooper RS, Bakker PIW de, Dedoussis G, Ferrucci L, Franks PW, Froguel P, Groop LC, Haiman CA, Hamsten A, Hayes MG, Hui J, Hunter DJ, Hveem K, Jukema JW, Kaplan RC, Kivimaki M, Kuh D, Laakso M, Liu Y, Martin NG, März W, Melbye M, Moebus S, Munroe PB, Njølstad I, Oostra BA, Palmer CNA, Pedersen NL, Perola M, Pérusse L, Peters U, Powell JE, Power C, Quertermous T, Rauramaa R, Reinmaa E, Ridker PM, Rivadeneira F, Rotter JI, Saaristo TE, Saleheen D, Schlessinger D, Slagboom PE, Snieder H, Spector TD, Strauch K, Stumvoll M, Tuomilehto J, Uusitupa M, Harst P van der, Völzke H, Walker M, Wareham NJ, Watkins H, Wichmann H-E, Wilson JF, Zanen P, Deloukas P, Heid IM, Lindgren CM, Mohlke KL, Speliotes EK, Thorsteinsdottir U, Barroso I, Fox CS, North KE, Strachan DP, Beckmann JS, Berndt SI, Boehnke M, Borecki IB, McCarthy MI, Metspalu A, Stefansson K, Uitterlinden AG, Duijn CM van, Franke L, Willer CJ, Price AL, Lettre G, Loos RJF, Weedon MN, Ingelsson E, O'Connell JR, Abecasis GR, Chasman DI, Goddard ME, Visscher PM, Hirschhorn JN, Frayling TM. 2014. Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat Genet* **46**:ng.3097. doi:10.1038/ng.3097

Wray NR, Goddard ME, Visscher PM. 2007. Prediction of individual genetic risk to disease from genome-wide association studies. *Genome Res* **17**:1520–1528. doi:10.1101/gr.6665407

Wray NR, Yang J, Hayes BJ, Price AL, Goddard ME, Visscher PM. 2013. Pitfalls of predicting complex traits from SNPs. *Nat Rev Genet* **14**:nrg3457. doi:10.1038/nrg3457

Yang, J., Benyamin, B., McEvoy, B. P., Gordon, S., Henders, A. K., Nyholt, D. R., Madden, P. A., Heath, A. C., Martin, N. G., Montgomery, G. W., Goddard, M. E., & Visscher, P. M. (2010). Common SNPs explain a large proportion of the heritability for human height. *Nat Genet,* **42**(7), 565-569. doi:ng.608

Andrew D Yates, Premanand Achuthan, Wasiu Akanni, James Allen, Jamie Allen, Jorge Alvarez-Jarreta, M Ridwan Amode, Irina M Armean, Andrey G Azov, Ruth Bennett, Jyothish Bhai, Konstantinos Billis, Sanjay Boddu, José Carlos Marugán, Carla Cummins, Claire Davidson, Kamalkumar Dodiya, Reham Fatima, Astrid Gall, Carlos Garcia Giron, Laurent Gil, Tiago Grego, Leanne Haggerty, Erin Haskell, Thibaut Hourlier, Osagie G Izuogu, Sophie H Janacek, Thomas Juettemann, Mike Kay, Ilias Lavidas, Tuan Le, Diana Lemos, Jose Gonzalez Martinez, Thomas Maurel, Mark McDowall, Aoife McMahon, Shamika Mohanan, Benjamin Moore, Michael Nuhn, Denye N Oheh, Anne Parker, Andrew Parton, Mateus Patricio, Manoj Pandian Sakthivel, Ahamed Imran Abdul Salam, Bianca M Schmitt, Helen Schuilenburg, Dan Sheppard, Mira Sycheva, Marek Szuba, Kieron Taylor, Anja Thormann, Glen Threadgold, Alessandro Vullo, Brandon Walts, Andrea Winterbottom, Amonida Zadissa, Marc Chakiachvili, Bethany Flint, Adam Frankish, Sarah E Hunt, Garth IIsley, Myrto Kostadima, Nick Langridge, Jane E Loveland, Fergal J Martin, Joannella Morales, Jonathan M Mudge, Matthieu Muffato, Emily Perry, Magali Ruffier, Stephen J Trevanion, Fiona Cunningham, Kevin L Howe, Daniel R Zerbino, Paul Flicek, Ensembl (2020). *Nucleic Acids Research*, **48**, D1, D682–D688, https://doi.org/10.1093/nar/gkz966

Yoon BS, Ovchinnikov DA, Yoshii I, Mishina Y, Behringer RR, Lyons KM. 2005. Bmpr1a and
Bmpr1b have overlapping functions and are essential for chondrogenesis in vivo. *P Natl Acad*
Sci Usa **102**:5062–5067. doi:10.1073/pnas.0500031102

Young NM, Chong HJ, Hu D, Hallgrímsson B, Marcucio RS. 2010. Quantitative analyses link
modulation of sonic hedgehog signaling to continuous variation in facial growth and shape.
Development **137**:3405–3409. doi:10.1242/dev.052340

Yue F, Cheng Y, Breschi A, Vierstra J, Wu W, Ryba T, Sandstrom R, Ma Z, Davis C, Pope BD,
Shen Y, Pervouchine DD, Djebali S, Thurman RE, Kaul R, Rynes E, Kirilusha A, Marinov
GK, Williams BA, Trout D, Amrhein H, Fisher-Aylor K, Antoshechkin I, DeSalvo G, See L-
H, Fastuca M, Drenkow J, Zaleski C, Dobin A, Prieto P, Lagarde J, Bussotti G, Tanzer A,
Denas O, Li K, Bender MA, Zhang M, Byron R, Groudine MT, McCleary D, Pham L, Ye Z,
Kuan S, Edsall L, Wu Y-C, Rasmussen MD, Bansal MS, Kellis M, Keller CA, Morrissey CS,
Mishra T, Jain D, Dogan N, Harris RS, Cayting P, Kawli T, Boyle AP, Euskirchen G,
Kundaje A, Lin S, Lin Y, Jansen C, Malladi VS, Cline MS, Erickson DT, Kirkup VM,
Learned K, Sloan CA, Rosenbloom KR, Sousa BL de, Beal K, Pignatelli M, Flicek P, Lian J,
Kahveci T, Lee D, Kent WJ, Santos MR, Herrero J, Notredame C, Johnson A, Vong S, Lee
K, Bates D, Neri F, Diegel M, Canfield T, Sabo PJ, Wilken MS, Reh TA, Giste E, Shafer A,
Kutyavin T, Haugen E, Dunn D, Reynolds AP, Neph S, Humbert R, Hansen RS, Bruijn MD,
Selleri L, Rudensky A, Josefowicz S, Samstein R, Eichler EE, Orkin SH, Levasseur D,
Papayannopoulou T, Chang K-H, Skoultchi A, Gosh S, Disteche C, Treuting P, Wang Y,
Weiss MJ, Blobel GA, Cao X, Zhong S, Wang T, Good PJ, Lowdon RF, Adams LB, Zhou X-
Q, Pazin MJ, Feingold EA, Wold B, Taylor J, Mortazavi A, Weissman SM,
Stamatoyannopoulos JA, Snyder MP, Guigo R, Gingeras TR, Gilbert DM, Hardison RC, Beer
MA, Ren B, Consortium TME. 2014. A comparative encyclopedia of DNA elements in the
mouse genome. *Nature* **515**:355. doi:10.1038/nature13992

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074 Figure 1. Schematic of the many-to-many relationship between genetic and phenotypic variation.
1075 From left to right: Allelic variation (colored dots) at individual genes is organized into
1076 developmental processes. Processes differ in start/end and duration during development. Genes
1077 are reused for different processes at different times. Processes are substantially pleiotropic in
1078 their effects contributing to global variation as well as local variation.
1079

Figure 2. Process MGP for chondrocyte differentiation. A) PLS1 genetic loadings are shown for each gene in the model. Individual founder allele effect sizes are colored within each bar. B-C) Estimated chondrocyte differentiation MGP phenotype is shown with black vectors multiplied 4x. A *Bmpr1b* (*Alk6*) homozygous mutant is shown with red vectors for comparison. The vector correlation between chondrocyte differentiation MGP and *Bmpr1b* is shown below the phenotypic effects.

Chondrocyte differentiation map -> Bmpr1b -/-
r = .312

1124 Figure 3. Process MGP for Fgf signaling. A) PLS1 genetic loadings are shown for each gene in
1125 the model. Individual founder allele effect sizes are colored within each bar. B-C) Estimated Fgf
1126 signaling MGP phenotype is shown with black vectors multiplied 4x. An *Fgf10* homozygous
1127 mutant is shown with red vectors for comparison. The vector correlation between Fgf signaling
1128 MGP and the *Fgf10* mutant is shown below the phenotypic effects.

Figure 4. Pairwise MGP vector correlations and random process correlations to principal components. A) Pairwise correlations of phenotypic effects for 15 process MGP analyses. Scale on the right denotes color correspondences to vector correlation, where yellows are high correlations, greens are moderate, and blues are low. B) The densities of vector correlations between 1000 randomly chosen process MGP effects and PCs 1-4.

Figure 5. Comparisons of MGP and mouse mutant directions. A) Seven MGP phenotypes
projected onto a PCA of the DO and a sample of 30 mutant mouse genotypes. Mutant means are
labeled in black. The directions of MGP effects are shown with orange vectors from the DO
mean to the associated process MGP. The range of DO variation on PCs 1 and 2 is shown with
the shaded ellipse with an orange border. B) A heatmap of vector correlations between 30 mutant
effects and 30 process MGP effects.
1167

1179 Figure 6. Example screenshot of web version of process analysis. Analyses include a barplot of
1180 the relative effect sizes of each selected marker and the associated phenotype shown with black
1181 vectors at each landmark. If a mutant comparison is selected, the vector correlation is provided
1182 and the mutant phenotype is shown with red vectors. Selecting "send me the results" generates
1183 an HTML report with an interactive 3D model.

Figure 7. 54 3D landmark configuration. A) Sagittal view of representative scan with landmarks shown as red spheres. B) Dorsal view of landmark configuration. C) Ventral view of landmark configuration.

1210 Figure 8. Process MGP schematic. Once a process is selected, we cross-reference the known
1211 gene locations with the locations of the genotyped markers in the DO sample. The founder
1212 probabilities of the nearest upstream and downstream markers are averaged for each gene. The
1213 compiled founder probabilities and landmark coordinates are then used in a regularized PLS
1214 model to estimate latent axes of covariation.

1227 Supplemental figure 1. Permutation of marker sets of fixed size. A) The permuted R^2 distribution
1228 of 1000 38-marker MGP analyses is shown in blue. The estimated R^2 of the chondrocyte
1229 differentiation MGP is shown as a black vertical line. B) The permuted R^2 distribution of 1000
1230 41-marker MGP analyses is shown in blue. The estimated R^2 of the Fgf signaling MGP is shown
1231 as a black vertical line.
1232

A)

B)

1249   Supplemental figure 2. Genetic effect loadings for multiple process MGPs. Selected processes
1250   are listed above their respective plots. Marker loadings are not scaled to a common range.

A)

B)

C)

1251
1252
1253
1254

1255 Supplemental figure 3. Additional process-to-mutant comparisons. A) The estimated
1256 chondrocyte differentiation MGP phenotype magnified 4x (black) with *Bmp2* homozygous
1257 mutant (red). B) The estimated Fgf signaling MGP phenotype magnified 4x (black) with *Fgf3*
1258 homozygous mutant (red).
1259
1260

1280   Supplemental figure 4. Demographic plots for the DO sample. A) The distribution of the sample
1281   by generation and data source (lab). B) Distribution of sex by source (lab).

A)

B)

1296   Supplemental figure 5. Histogram of kinship matrix values in the DO sample.
1297

Supplemental figure 6. Combining queries in the MGP shiny app with the pipe operator. In order to filter the GO database with multiple terms, the pipe operator can be used as shown. Here, the user has selected processes associated with either the apoptosis or Wnt pathway process. The barplot shows the relative effect sizes for markers associated to both "Wnt signaling pathway" and "execution phase of apoptosis" GO terms.

1354 Supplemental table 1. Top 10 MGP vector correlations with PC1. Corresponding PC2- 4 vector
1355 correlations are also provided.

PROCESS	PC1 CORRELATION	PC2 CORRELATION	PC3 CORRELATION	PC4 CORRELATION
ZINC ION BINDING	0.859241	0.01277426	0.07172876	0.20641736
SENSORY PERCEPTION OF SOUND	0.84834632	0.10996389	0.0407517	0.20432551
CALCIUM ION TRANSPORT	0.84816486	0.16123711	0.05231719	0.11159004
PROTEIN HOMOOLIGOMERIZATION	0.84782484	0.20793974	0.0156349	0.16244344
DENDRITE MORPHOGENESIS	0.84752457	0.06832718	0.10026087	0.07711775
NEUROPEPTIDE SIGNALING PATHWAY	0.84620616	0.21526031	0.02883174	0.13173442
FOCAL ADHESION	0.84611744	0.13458614	0.04380431	0.21903587
CHROMOSOME SEGREGATION	0.84610062	0.10383492	0.03360725	0.16218602
SARCOMERE ORGANIZATION	0.83893574	0.15368816	0.02581707	0.02880551
INTEGRAL COMPONENT OF ENDOPLASMIC RETICULUM MEMBRANE	0.83775207	0.1673053	0.03517157	0.02691882

1356
1357
1358 Supplemental table 2. Processes with moderate vector correlations to PC 1-4.

PROCESS	PC1 CORRELATION	PC2 CORRELATION	PC3 CORRELATION	PC4 CORRELATION
NEGATIVE REGULATION OF I-KAPPAB KINASE/NF-KAPPAB SIGNALING	0.54158492	0.25581449	0.49244383	0.25894853
LOCOMOTORY EXPLORATION BEHAVIOR	0.37993095	0.2110723	0.68937988	0.24189284
AU-RICH ELEMENT BINDING	0.57006676	0.32997589	0.40318313	0.20843601
TRANSCRIPTION REGULATORY REGION SEQUENCE-SPECIFIC DNA BINDING	0.70056311	0.09294266	0.44398467	0.26285711
DYNEIN COMPLEX	0.68525872	0.24644459	0.31380962	0.19363051
PRESYNAPTIC CYTOSOL	0.39216186	0.38329479	0.13557636	0.50766778
NEGATIVE REGULATION OF FAT CELL DIFFERENTIATION	0.39080052	0.24460715	0.73049567	0.02976212
CELL BODY	0.80176806	0.25146354	0.06871634	0.2628449
ADULT BEHAVIOR	0.74617475	0.3923876	0.03450902	0.21024944
PROTEASOMAL UBIQUITIN-INDEPENDENT PROTEIN CATABOLIC PROCESS	0.61654501	0.39974619	0.09993121	0.26252902