Molecular characterisation of, and antimicrobial resistance in, *Clostridioides difficile* from Thailand, 2017-2018

Korakrit Imwattana,[1,2] Papanin Putsathit,[3] Daniel R Knight,[1,4] Pattarachai Kiratisin,[2] and Thomas V Riley[1,3,4,5,]*

[1] School of Biomedical Sciences, The University of Western Australia, Western Australia, [2] Faculty of Medicine Siriraj Hospital, Mahidol University, Thailand, [3] School of Medical and Health Sciences, Edith Cowan University, Western Australia, [4] School of Veterinary and Life Sciences, Murdoch University, Western Australia, [5] Department of Microbiology, PathWest Laboratory Medicine, Queen Elizabeth II Medical Centre, Western Australia

* Corresponding author - Professor Thomas V Riley (thomas.riley@uwa.edu.au), School of Biomedical Sciences, The University of Western Australia, Nedlands, Western Australia, Australia

Running title: *Clostridioides difficile* in Thailand

Synopsis

Background: Antimicrobial resistance (AMR) plays an important role in the pathogenesis and spread of *Clostridioides difficile* infection (CDI). Many antimicrobials, such as fluoroquinolones, have been associated with outbreaks of CDI globally.

Objectives: This study aimed to characterise AMR among clinical *C. difficile* strains in Thailand, a country where the use of antimicrobials remains inadequately regulated.

Methods: Stool samples were screened for *tcdB* and positives were cultured*. C. difficile* isolates were characterised by toxin profiling and PCR ribotyping. Antimicrobial susceptibility testing was performed using an agar incorporation method, and whole-genome sequencing and AMR genotyping performed on a subset of strains.

Results: A total of 321 *C. difficile* strains were isolated from 326 stool samples. The most common toxigenic ribotype (RT) was RT 017 (18%), followed by RTs 014 (12%) and 020 (7%). There was a high resistance prevalence (≥ 10%) to clindamycin, erythromycin, moxifloxacin and rifaximin, and resistance prevalence was greatest among RT 017 strains. AMR genotyping revealed a significant correlation between resistance genotype and phenotype for moxifloxacin and rifampicin. The presence of *erm*-class genes was associated with high-level clindamycin and erythromycin resistance. Point substitutions on the penicillin-binding proteins (PBP1 and PBP3) were not sufficient to confer meropenem resistance, however, a Y721S substitution in PBP3 was associated with a slight increase in meropenem MIC. No resistance to metronidazole, vancomycin or fidaxomicin was observed.

Conclusion: There was a large proportion of *C. difficile* RT 017 in Thailand and significant AMR among these strains*.* The concordance between AMR phenotype and genotype was significant.

Introduction

Clostridioides (*Clostridium*) *difficile* is a major cause of antimicrobial-associated diarrhoea globally (1). *C. difficile* infection (CDI) is mediated by toxins and, so far, there have been three different major toxins identified: toxin A (TcdA), toxin B (TcdB) and binary toxin (*C. difficile* transferase, CDT). The genes encoding TcdA and TcdB are located on the 19.6 kb pathogenicity locus (PaLoc) (2) and the genes for CDT are located in a different part of the chromosome, the CDT locus (3). In non-toxigenic *C. difficile* (NTCD), the PaLoc is replaced by a fixed 115 bp locus (2). The toxin genes (*tcdA*, *tcdB*, *cdtA* and *cdtB*) can be detected by PCR (4, 5), however, some *C. difficile* strains, such as *C. difficile* ribotype (RT) 017, have a deletion at the repeating region of the *tcdA* gene, resulting in a truncated and non-functional toxin A (6). Several methods have been developed to detect this deletion (7, 8).

C. difficile can be separated into different RTs by amplifying the intergenic spacer region between the 16S and 23S rRNA genes (9). This classification method has been used worldwide due to its simplicity and high discriminating power (10). Important *C. difficile* RTs include *C. difficile* RT 027, an A+B+CDT+ strain associated with outbreaks of severe CDI in North America and Europe in the early 2000s (11), *C. difficile* RT 078, another A+B+CDT+ strain associated with zoonotic disease (12), and *C. difficile* RT 017, a *tcdA*-negative (A-B+CDT-) strain associated with global outbreaks since 1995 (6).

Although resistance to antimicrobials used for the treatment of CDI is relatively rare (13), resistance to other commonly used antimicrobials plays an important role in the pathogenesis of CDI and the spread of *C. difficile*. Resistance to clindamycin, new generation fluoroquinolones, rifampicin and tetracycline has been associated with outbreaks of CDI (14), and these antimicrobials are also associated with an increased risk of developing CDI in general (15). Strict regulation of antimicrobials can be a successful measure to control CDI. In the US, such regulation resulted in a significant decrease in CDI cases and CDI-related deaths (16). Fluoroquinolone regulation in Australia is also associated with a relatively low prevalence of fluoroquinolone-resistant organisms (17), including *C. difficile* (18).

In previous studies, the epidemiology of CDI in Thailand has been characterised by a high prevalence of A-B+CDT- and NTCD, and an absence of A+B+CDT+ strains (19-21). *C. difficile* strains isolated in Thailand, especially *C. difficile* RT 017, were resistant to many antimicrobial groups, reflecting the use and misuse of these antimicrobials in the country (22). This study provides an update on the characterisation and antimicrobial susceptibility of *C. difficile* isolated from a tertiary hospital in Bangkok, Thailand.

Materials and Methods

Isolation and characterisation of *C. difficile*

The study included 326 diarrhoeal stools samples collected from patients being treated at Siriraj Hospital, a large teaching hospital in Bangkok, Thailand, during 2017 – 2018. All stool samples were screened for the presence of the *tcdB* gene using the BD Max Cdiff assay (Becton Dickinson, US) as a part of routine investigations at Siriraj Hospital, and stools that were *tcdB* positive were then sent to a reference laboratory in Perth, Western Australia, for further investigation.

At the reference laboratory, stool samples were processed as previously described, including enrichment culture in cooked meat broth supplemented with gentamicin, cefoxitin and cycloserine (23). Characterisation of *C. difficile* was performed by PCR ribotyping and toxin gene profiling. PCR ribotyping was performed using the method described by Stubbs *et al* (9). The banding patterns were compared to a local database consisting of more than 80 internationally recognised RTs, including 15 reference RTs from the European Centre for Disease Prevention and Control. Patterns that did not match strains in the database were given an internal nomenclature. Detection of *tcdA* and *tcdB,* and the binary toxin genes (*cdtA* and *cdtB*), was performed using the methods described by Kato

et al (7) and Stubbs *et al* (5), respectively. All NTCD isolates in this study were confirmed as such by
the absence of the PaLoc using the method described by Braun *et al* (*lok* PCR) (2).

All stool samples were screened also for colonisation with multiple *C. difficile* strains. Briefly,
DNA extraction was performed on all enrichment broths. DNA was then screened with either *tcdB* (7)
or *lok* PCR (2)*,* based on the toxin profile of the first *C. difficile* strain isolated from the specimen (2,
7). For example, a specimen previously positive for toxigenic *C. difficile* (TCD) was screened with
lok PCR for NTCD and vice versa. All screening-positive broths were re-cultured and the second
C. difficile strain characterised as described above.

<u>Antimicrobial susceptibility testing</u>

Antimicrobial susceptibility testing (AST) was performed using the agar incorporation method as
described by the Clinical and Laboratory Standards Institute (CLSI) against nine antimicrobials listed in
Table 1 (24). *C. difficile* ATCC 700057, *Bacteroides fragilis* ATCC 25285, *Eubacterium lentum* ATCC
43055 and *B. thetaiotamicron* ATCC 29741 were included as control strains. Susceptibility results were
interpreted using the minimal inhibitory concentration (MIC) breakpoints listed in **Table 1** (24-28).
C. difficile strains that were resistant to at least three different antimicrobial classes were classified as
multidrug-resistant (MDR). Resistance to clindamycin and erythromycin was considered as resistance
to a single class (macrolide-lincosamide-streptogramin B; MLS_B).

Table 1 – List of antimicrobials, test ranges and susceptibility breakpoints used in this study.

Antimicrobial	Test range (mg/l)	Breakpoint (mg/l)			Reference
		S	I	R	
Clindamycin	0.008 – 256	≤ 2	4	≥ 8	(24)
Erythromycin	0.06 – 256	-	-	> 8	(25)
Moxifloxacin	0.06 – 64	≤ 2	4	≥ 8	(24)
Amoxicillin/clavulanate	0.015 – 16	≤ 4	8	≥ 16	(24)
Meropenem	0.12 – 16	≤ 4	8	≥ 16	(24)
Rifaximin	0.008 – 64	-	-	> 32	(26)
Metronidazole	0.008 – 4	≤ 2	-	> 2	(27)
Vancomycin	0.03 – 4	≤ 2	-	> 2	(27)
Fidaxomicin	0.002 – 2	-	-	> 1	(28)

<u>Whole-genome sequencing, high-resolution typing and antimicrobial resistance characterisation</u>

A subset of *C. difficile* strains (n = 37) was selected for whole-genome sequencing (WGS) to explore
possible antimicrobial resistance (AMR) genotypes. Genomic DNA was extracted, sequenced on an
Illumina HiSeq platform which generated 150 bp pair-end reads with a median coverage of 73X and
characterised by multi-locus sequence typing (MLST) as previously described (29). Clade assignment
of a newly defined sequence type (ST) was confirmed by comparing the average nucleotide identity
(ANI) with *C. difficile* strains 630 (clade 1, GenBank accession AM180355) and R20291 (clade 2,
GenBank accession FN545816) using FastANI (30). Known accessory AMR genes were identified by
interrogating the read files with SRST2 version 0.2.0 against ARGannot database version 3 (31, 32).
Draft annotated genomes were interrogated on Artemis version 17.0.1 to look for additional accessory
genes (33). The genomes were interrogated also for the presence of known point substitutions
associated with resistance to carbapenems (substitution in penicillin-binding proteins PBP1 and PBP3),
fluoroquinolones (substitution in GyrA and GyrB subunit of the gyrase enzyme) and rifaximin
(substitution in RpoB enzyme) (34, 35). The genotypes were then compared with phenotypic
susceptibility data. All sequence data generated in this study have been submitted to the European
Nucleotide Archive under BioProject PRJEB40974, sample accessions ERS5247348 – ERS5247384.
Details of the genomes are available in the **Supplementary Table S1**.

 Statistical analysis

All statistical analyses were performed using online tools by Social Science Statistics available at
https://www.socscistatistics.com/. A p-value ≤ 0.05 was considered statistically significant.

Human research ethics approval

This study was approved by the Human Research Ethics Committee of The University of Western
Australia (reference file RA/4/20/4704) and the Siriraj Institutional Review Board (protocol number
061/2558 [EC1]).

Results

Characterisation of Thai *C. difficile*

A total of 296 *C. difficile* strains were initially isolated from the stool samples. Forty-four of the original
326 PCR positive stool samples were negative for TCD by culture; 30 contained no *C. difficile* while
from 14 only NTCD was cultured. The enrichment broths for these samples were re-screened for TCD
and were all negative by *tcdB* PCR. Another 25 strains were identified from the co-colonisation
screening process, yielding a total of 321 *C. difficile* strains. Of these, 221 (68.85%) were positive for
tcdA and *tcdB* (A+B+CDT-), 58 (18.07%) were positive for *tcdB* only and had a deletion in *tcdA* (A-
B+CDT-), three (0.93%) were positive for *tcdA*, *tcdB*, as well as *cdtA* and *cdtB* (A+B+CDT+) and 39 strains
(12.15%) were negative for all toxin genes (A-B-CDT-, NTCD).

The 321 *C. difficile* strains belonged to 63 different RTs, 19 of which were internationally
recognised. The remaining RTs were given internal nomenclature beginning with either "QX" or "KI".
The prevalence of the most common RTs is summarised in **Table 2**. The most common TCD strain was
C. difficile RT 017 (A-B+CDT-), followed by RTs 014 and 020 (both A+B+CDT-). The most common NTCD
was *C. difficile* RT 010.

Characterisation of a novel binary toxin-positive *C. difficile* isolate

One of the *C. difficile* strains isolated in this study was positive for all three toxin genes (A+B+CDT+)
and had a unique ribotyping pattern compared to the local reference library. According to the MLST
scheme, this isolate was characterised as the novel ST 692 within evolutionary clade 1. However,
pairwise ANI analysis showed that this strain was more closely related to *C. difficile* R20291 (clade 2,
ANI = 99.17%) than *C. difficile* 630 (clade 1, ANI = 98.89%).

Antimicrobial susceptibility of Thai *C. difficile*

Overall antimicrobial susceptibility results are shown in **Table 3** and the MIC distribution of selected
six antimicrobial classes is displayed in **Figure 1**. Based on the MIC value, clindamycin-resistant
C. difficile strains could be divided into two groups: those with MIC ≥ 32 mg/l (n = 97) and those with
MIC < 32 mg/l (n = 166). There was a strong correlation between high-level clindamycin resistance and
erythromycin resistance: 95 strains (97.94%) that had clindamycin MIC ≥ 32 mg/l were also resistant
to erythromycin while only 16 strains (9.64%) in the other group were resistant to erythromycin
(Cohen's kappa = 0.857). There was also a clear separation in MIC value between strains with and
without rifaximin resistance. The separation was less clear for moxifloxacin and was not observed in
amoxicillin/clavulanate and meropenem.

156 **Table 2** – Ribotypes of 321 *C. difficile* strains from Thailand, by toxin profile.

Toxin Profile	Ribotype	Number	% of toxigenic strains	% of all
A+B+CDT-	**Total**	**221**	**78%**	**69%**
	014	40	14%	12%
	020	24	9%	7%
	046	18	6%	6%
	QX517	17	6%	5%
	297	13	5%	4%
	QX026	11	4%	3%
	043	10	4%	3%
	Others	88	31%	27%
A-B+CDT-	**Total**	**58**	**21%**	**18%**
	017	58	21%	18%
A+B+CDT+	**Total**	**3**	**1%**	**1%**
	078	1	<1%	<1%
	QX273	1	<1%	<1%
	KI008	1	<1%	<1%
A-B-CDT-	**Total**	**39**	-	**12%**
	010	7	-	2%
	QX002	5	-	2%
	009	4	-	1%
	QX011	4	-	1%
	Others	19	-	6%

157 Note: RTs designated with "QX" and "KI" were RTs that did not match the internationally recognised
158 RTs in the database and were given internal nomenclature.

160 **Table 3** – Antimicrobial susceptibility of 321 *C. difficile* strains from Thailand.

Antimicrobial	MIC range (mg/l)	MIC$_{50}$ (mg/l)	MIC$_{90}$ (mg/l)	Susceptibility (%)		
				S	I	R
Clindamycin	0.5 – >256	8	>256	26 (8%)	32 (10%)	263 (82%)
Erythromycin	0.5 – >256	2	>256	-	-	109 (34%)
Moxifloxacin	0.5 – >64	2	32	242 (75%)	3 (1%)	76 (24%)
Amoxicillin/clavulanate	0.25 – 8	0.5	1	320 (>99%)	1 (<1%)	0
Meropenem	1 – 16	4	8	280 (87%)	39 (12%)	2 (1%)
Rifaximin	≤0.008 – >64	0.03	2	-	-	31 (10%)
Metronidazole	0.06 – 1	0.25	0.25	321 (100%)	-	0
Vancomycin	0.5 – 2	1	2	321 (100%)	-	0
Fidaxomicin	0.03 – 0.25	0.125	0.25	-	-	0

When classified by toxin gene profiles, resistance to clindamycin, erythromycin, moxifloxacin and rifaximin were more prevalent among A-B+CDT- *C. difficile*, all of which belonged to RT 017, than A+B+CDT- and NTCD (**Figure 1**). A total of 29 (9.03%) *C. difficile* isolates were MDR, 26 (8.10%) of which were *C. difficile* RT 017 (A-B+CDT-). The remaining isolates were NTCD (n=2) and A+B+CDT- *C. difficile* (n=1). All MDR isolates were resistant to the MLS_B group (both clindamycin and erythromycin), moxifloxacin and rifaximin. One MDR isolate was also resistant to meropenem (A-B+CDI-, RT 017, MIC = 16 mg/l).

Figure 1 – MIC distribution for six antimicrobials against 321 strains of *C. difficile* in Thailand. *C. difficile* strains were classified according to their toxin gene profiles: A+B+CDT-, red; A-B+CDT-, yellow; A-B-CDT-, green. The number of A+B+CDT+ *C. difficile* strains was low (n = 3) and these were excluded. Breakpoints for intermediate resistance (I) and resistance (R) are shown by broken and solid lines, respectively.

 Genotypic resistance determinants in Thai *C. difficile*

A summary of MIC values and genotypic resistance determinants of 37 sequenced *C. difficile* strains is
available in **Supplementary Table S2**. Of these strains, 31 had high-level resistance to clindamycin: 23
strains carried an *erm*(B) gene, five carried an *erm*(G) gene and three carried a gene encoding an rRNA
adenine N(6)-methyltransferase family protein [GenBank accession WP_002571018.1], the closest
match to which in the CARD database was the Erm(42) protein [39% identity, E-value = 1.19E-53] (36).
Of the 23 *erm*(B)-positive strains, 19 carried the *erm*(B) gene on transposon Tn*6194* (82.61%), while
the other four (17.39%) carried the gene on Tn*6189*. No *erm*-class genes were identified among strains
with low-level clindamycin resistance. The concordance between the presence of *erm*-class gene and
high-level clindamycin resistance was 91.89%. The concordance increased to 100% when the
uncharacterised *erm*-class gene was included. A gene encoding a macrolide efflux protein [accession
WP_009271522.1] was identified in two strains with high-level erythromycin resistance (MIC > 256
mg/l). No significant genotypic resistance determinants were identified in strains with low-level
clindamycin resistance.

A T82I substitution in GyrA was found in 23 strains and a D426V substitution in GyrB was found
in one strain with a moxifloxacin MIC of 16 – 32 mg/l. No known point substitutions were found in the
remaining 14 strains, 13 of which were moxifloxacin-susceptible; one had low-level moxifloxacin
resistance (MIC 8 mg/l) [97.37% concordance]. H502N and R505K substitutions in RpoB were found in
all 23 rifaximin-resistant strains and none of the susceptible strains [100% concordance].

Twelve strains had an A555T substitution in penicillin-binding protein 1 (PBP1) and another
seven had a Y721S substitution in PBP3. A multiple linear regression analysis was performed to assess
the association between the presence of these substitutions and the MIC values for meropenem. The
Y721S substitution on the PBP3 was associated with an increase in meropenem MIC (adjusted R^2 =
0.516, t = 5.521, p < 0.0001), while the A555T substitution on the PBP1 was not associated with the
change in meropenem MIC (t = -1.127, p = 0.268).

Discussion

This study provides an update on the molecular epidemiology and antimicrobial susceptibility of
C. difficile strains circulating in Thailand. It also explores the genomic basis of important AMR in these
strains. As the focus was on patients suspected of having CDI and not on the prevalence of *C. difficile*
in the general population, the stool samples included in this study were first screened for the *tcdB*
gene by PCR before culture. Thus the overall prevalence of TCD was higher than previous studies in
Thailand (19-21), however, the common strains found were similar. The majority of A+B+CDT- strains
belonged to *C. difficile* RT 014/020 group, all A-B+CDT- strains belonged to *C. difficile* RT 017 and most
NTCD belonged to *C. difficile* RTs 009, 010 and 039. Three binary toxin-positive isolates were found in
this study, one of which was *C. difficile* RT 078. The epidemic *C. difficile* RT 027 remained absent in
Thailand despite its successful spread in other regions (37).

Why *C. difficile* RT 027 has failed to spread and to establish in Thailand, and Southeast Asia in
general, remains unknown. One possible reason is that the successful spread of this RT was mainly
due to its resistance to fluoroquinolones which provided an advantage over other less resistant RTs
(38). Although there is high consumption of fluoroquinolones, such as levofloxacin, in the country (39),
Thailand already harbours *C. difficile* RT 017, another epidemic RT with a high prevalence of resistance
to fluoroquinolones, as well as other antimicrobials (14). Thus, it may have been difficult for *C. difficile*
RT 027 to compete with this local RT compared to other regions.

Though *C. difficile* RT 027 was not identified in this study, a possible relative of this
hypervirulent strain was identified (ST 692). The MLST result for this strain was unusual, as it was
classified into clade 1 despite containing binary toxin genes in a complete CDT locus. The presence of

binary toxin genes is a feature common in *C. difficile* clades 2 and 5 but rare in clade 1 (40). Thus, an
ANI analysis was performed, which indicated that this newly characterised strain was more related to
clade 2 than clade 1 *C. difficile*, as expected from the toxin gene profile. Clades 1 and 2 *C. difficile* are
closely related and share a large proportion of housekeeping gene alleles used in the MLST scheme.
As a result, it may be difficult to properly discriminate these two clades by this method. The use of ANI
analysis, which involves the whole genome rather than a specific set of housekeeping genes, can help
in the correct classification of some borderline strains as shown in a previous study (40). According to
the ANI analysis, it is likely that this newly described *C. difficile* ST 692 belongs to clade 2 and is related
to *C. difficile* RT 027.

A discordance between culture results and the result of a conventional real-time *tcdB* PCR was
observed in 44 stool samples. The false-positive rate of the real-time PCR method (13.50%) was
comparable to the previous report comparing *tcdB* PCR with a similar enrichment culture method but
without the colonisation screening step (23). This suggests that the additional screening step does not
significantly increase the yield of the culture method, although it may help identify stool samples with
multiple *C. difficile* strains. This false-positive rate also highlights the importance of patient clinical
data or additional tests to improve the accuracy of CDI diagnosis. In the latest guidelines for the
treatment and diagnosis of CDI, *tcdB* PCR is no longer recommended as a stand-alone test unless the
patient has symptoms suggestive of CDI (41).

AMR in *C. difficile* affects both the pathogenesis and treatment of CDI. To cause the disease,
C. difficile must tolerate the presence of antimicrobials in the intestinal lumen while the microbiota
perishes (14). Many successful *C. difficile* lineages have indeed been characterised with increased
resistance to at least one major drug group (14). In this study, *C. difficile* RT 017, which was the most
common RT, had greater resistance to MLS$_B$ (both clindamycin and erythromycin), moxifloxacin and
rifaximin than other RTs. It was also the most common MDR *C. difficile* strain in this study. *C. difficile*
RT 017 has been reported also to be the most prevalent RT with significant resistance to many
antimicrobials in other parts of Thailand (21). This particular RT has been associated with resistance
to at least six antimicrobial groups (14), which may have accounted for its successful global spread (6).
As regulation of antimicrobial use has reduced the impact of *C. difficile* in many countries (16, 18), a
similar approach should effective for the control CDI in Thailand.

All *erm*(B)-positive *C. difficile* strains carried the gene on two well-characterised *erm*(B)-
positive transposons: Tn*6189* and Tn*6194*, the latter being found also in *C. difficile* M68, a *C. difficile*
RT 017 strain widely used as a reference in genomic studies (14). Tn*6194*, the most prevalent
transposon in this study, is capable of inter-species transfer, most notably between *C. difficile* and
Enterococcus faecalis (42). This emphasises another aspect of AMR in *C. difficile*; its possible role as a
reservoir of AMR genes for other pathogenic bacteria residing in the colon.

Previously, it has been reported that concordance between the presence of the *erm*(B) gene
and the MLS$_B$ resistance phenotype was low (43). This is likely due to the presence of multiple
resistance mechanisms. In a previous study, however, the *erm*(B) gene was found in all *C. difficile*
isolates with high-level resistance to both clindamycin and erythromycin, hinting that the gene may
be associated with high-level MLS$_B$ resistance (44). We also observed separation between *C. difficile*
strains with high-level and low-level clindamycin and erythromycin resistance (**Figure 1**). Upon
genomic analysis of a subset of strains, we found a correlation between the presence of an *erm*-class
gene (*erm*(B), *erm*(G) and another possible *erm*-class gene) with high-level resistance to clindamycin.
Many of these strains also had high-level erythromycin resistance, supporting the previous study (44).
A gene encoding a macrolide efflux protein was identified also in strains with high-level erythromycin
resistance and not in erythromycin-susceptible strains, suggesting an association between the
presence of this efflux protein and phenotypic erythromycin resistance, although the function of this
gene was not characterised. Resistance determinants were not identified among strains with low-level
clindamycin resistance, however, this underestimation is likely irrelevant, as the median clindamycin

MIC in this population (8 mg/l) remained significantly lower than the clindamycin level in stools (approximately 240 mg/g of stool) (45). Besides MLS$_B$, a separation between strains resistant and susceptible to rifaximin and fluoroquinolones was observed (**Figure 1**). The concordance between resistant phenotype and known genotype was also high, similar to a previous study (43).

Compared to the study at the same hospital in 2015, there was no difference in the overall resistance prevalence (25), however, there was a slight increase in meropenem MICs and the emergence of carbapenem resistance. In a previous study, high-level imipenem resistance was associated with point substitutions on PBP1 and PBP3 (35). It appears that these substitutions only confer high-level resistance to imipenem and not to meropenem, although the linear regression analysis suggests that the Y721S substitution on PBP3 may have contributed to a slight increase in meropenem MIC in this study.

C. difficile remained susceptible to metronidazole, vancomycin and fidaxomicin, similar to the other parts of the world (46). Thus, these antimicrobials should remain an effective treatment for CDI. There was a slight increase in vancomycin MIC reaching the clinical breakpoint, which is consistent with a previous study (25), however, this should have little impact on the treatment of CDI given that the faecal vancomycin concentration remains far greater than the MIC (>2,000 mg/l vs 2 mg/l) (47). The increase in vancomycin MIC in this study is in contrast to other hospitals in Thailand and this could reflect the usage of vancomycin at the study site (21). The use of vancomycin can lead to the emergence of vancomycin-resistant *Enterococcus* spp., which can have a devastating effect on patients (48, 49). Therefore, the use of vancomycin should be carefully monitored.

Conclusion

A-B+CDT- and NTCD were prevalent in Thailand. Few binary toxin-positive strains (A+B+CDT+) were identified; one belonged to a known epidemic lineage and another was a novel strain related to *C. difficile* RT 027. The most common RT in this study was *C. difficile* RT 017 (A-B+CDT-), a large proportion of which was resistant to MLS$_B$, moxifloxacin and rifaximin. Many strains were also MDR. Such resistance may have played a role in the success of *C. difficile* RT 017 in Thailand. There was a strong concordance between the presence of *erm*-class genes and high-level clindamycin resistance, as well as significant concordance between point substitutions on gyrase subunits and RpoB protein with fluoroquinolone and rifaximin resistance, respectively. Resistance to antimicrobials for treatment of CDI was not detected.

Acknowledgement

Funding

This work was supported by Mahidol University (Mahidol Scholarship to K.I.) and the National Health and Medical Research Council of Australia (Peter Doherty Biomedical Early Career Fellowship [APP1138257] to D.R.K).

Transparency declaration

T.V.R. has received grants from Cepheid; Merck; Otsuka; Roche; Sanofi and Summit for work outside that in this report. Other authors have no conflicts of interest to declare.

References

1.   Leffler DA, Lamont JT. *Clostridium difficile* Infection. N Engl J Med. 2015;373(3):287-8.

2.   Braun V, Hundsberger T, Leukel P, Sauerborn M, von Eichel-Streiber C. Definition of the single integration site of the pathogenicity locus in *Clostridium difficile*. Gene. 1996;181(1-2):29-38.

3.   Perelle S, Gibert M, Bourlioux P, Corthier G, Popoff MR. Production of a complete binary toxin (actin-specific ADP-ribosyltransferase) by *Clostridium difficile* CD196. Infect Immun. 1997;65(4):1402-7.

4.   Kato N, Ou CY, Kato H, Bartley SL, Brown VK, Dowell VR, Jr., et al. Identification of toxigenic *Clostridium difficile* by the polymerase chain reaction. J Clin Microbiol. 1991;29(1):33-7.

5.   Stubbs S, Rupnik M, Gibert M, Brazier J, Duerden B, Popoff M. Production of actin-specific ADP-ribosyltransferase (binary toxin) by strains of *Clostridium difficile*. FEMS Microbiol Lett. 2000;186(2):307-12.

6.   Imwattana K, Knight DR, Kullin B, Collins DA, Putsathit P, Kiratisin P, et al. *Clostridium difficile* ribotype 017 - characterization, evolution and epidemiology of the dominant strain in Asia. Emerg Microbes Infect. 2019;8(1):796-807.

7.   Kato H, Kato N, Watanabe K, Iwai N, Nakamura H, Yamamoto T, et al. Identification of toxin A-negative, toxin B-positive *Clostridium difficile* by PCR. J Clin Microbiol. 1998;36(8):2178-82.

8.   Lemee L, Dhalluin A, Testelin S, Mattrat M-A, Maillard K, Lemeland J-F, et al. Multiplex PCR targeting *tpi* (triose phosphate isomerase), *tcdA* (Toxin A), and *tcdB* (Toxin B) genes for toxigenic culture of *Clostridium difficile*. J Clin Microbiol. 2004;42(12):5710-4.

9.   Stubbs SL, Brazier JS, O'Neill GL, Duerden BI. PCR targeted to the 16S-23S rRNA gene intergenic spacer region of *Clostridium difficile* and construction of a library consisting of 116 different PCR ribotypes. J Clin Microbiol. 1999;37(2):461-3.

10.   Elliott B, Androga GO, Knight DR, Riley TV. *Clostridium difficile* infection: Evolution, phylogeny and molecular epidemiology. Infect Genet Evol. 2017;49:1-11.

11.   Warny M, Pepin J, Fang A, Killgore G, Thompson A, Brazier J, et al. Toxin production by an emerging strain of *Clostridium difficile* associated with outbreaks of severe disease in North America and Europe. Lancet. 2005;366(9491):1079-84.

12.   Bakker D, Corver J, Harmanus C, Goorhuis A, Keessen EC, Fawley WN, et al. Relatedness of human and animal *Clostridium difficile* PCR ribotype 078 isolates determined on the basis of multilocus variable-number tandem-repeat analysis and tetracycline resistance. J Clin Microbiol. 2010;48(10):3744-9.

13.   Banawas SS. *Clostridium difficile* infections: a global overview of drug sensitivity and resistance mechanisms. Biomed Res Int. 2018;2018:8414257.

14.   Imwattana K, Knight DR, Kullin B, Collins DA, Putsathit P, Kiratisin P, et al. Antimicrobial resistance in *Clostridium difficile* ribotype 017. Expert Rev Anti Infect Ther. 2020;18(1):17-25.

15.   Slimings C, Riley TV. Antibiotics and hospital-acquired *Clostridium difficile* infection: update of systematic review and meta-analysis. J Antimicrob Chemother. 2014;69(4):881-91.

16.   Centers for Disease Control and Prevention (CDC). Antibiotic resistance threats in the United States, 2019. Atlanta, GA: U.S. Department of Health and Human Services, CDC; 2019.

17.   Cheng AC, Turnidge J, Collignon P, Looke D, Barton M, Gottlieb T. Control of fluoroquinolone resistance through successful regulation, Australia. Emerg Infect Dis. 2012;18(9):1453-60.

18.   Lew T, Putsathit P, Sohn KM, Wu Y, Ouchi K, Ishii Y, et al. Antimicrobial susceptibilities of *Clostridium difficile* isolates from 12 Asia-Pacific countries in 2014 and 2015. Antimicrob Agents Chemother. 2020;64(7).

19.   Putsathit P, Maneerattanaporn M, Piewngam P, Kiratisin P, Riley TV. Prevalence and molecular epidemiology of *Clostridium difficile* infection in Thailand. New Microbes New Infect. 2017;15:27-32.

20.   Ngamskulrungroj P, Sanmee S, Putsathit P, Piewngam P, Elliott B, Riley TV, et al. Molecular epidemiology of *Clostridium difficile* infection in a large teaching hospital in Thailand. PLoS One. 2015;10(5):e0127026.

21.   Imwattana K, Wangroongsarb P, Riley TV. High prevalence and diversity of *tcdA*-negative and
tcdB-positive, and non-toxigenic, *Clostridium difficile* in Thailand. Anaerobe. 2019;57:4-10.
22.   Zellweger RM, Carrique-Mas J, Limmathurotsakul D, Day NPJ, Thwaites GE, Baker S, et al. A
current perspective on antimicrobial resistance in Southeast Asia. J Antimicrob Chemother.
2017;72(11):2963-72.
23.   Putsathit P, Morgan J, Bradford D, Engelhardt N, Riley TV. Evaluation of the BD Max Cdiff
assay for the detection of toxigenic *Clostridium difficile* in human stool specimens. Pathology.
2015;47(2):165-8.
24.   Clinical and Laboratory Standards Institute (CLSI). Methods for Antimicrobial Susceptibility
Testing of Anaerobic Bacteria, Approved Standard M11-A8. 8th ed. Wayne, PA: Clinical and
Laboratory Standards Institute; 2012.
25.   Putsathit P, Maneerattanaporn M, Piewngam P, Knight DR, Kiratisin P, Riley TV.
Antimicrobial susceptibility of *Clostridium difficile* isolated in Thailand. Antimicrob Resist Infect
Control. 2017;6:58.
26.   O'Connor JR, Galang MA, Sambol SP, Hecht DW, Vedantam G, Gerding DN, et al. Rifampin
and rifaximin resistance in clinical isolates of *Clostridium difficile*. Antimicrob Agents Chemother.
2008;52(8):2813-7.
27.   European Committee on Antimicrobial Susceptibility Testing. Clinical breakpoint tables,
version 8.1 London, United Kingdom [updated 16 May 2018. Available from:
http://www.eucast.org/clinical_breakpoints/.
28.   Goldstein EJC, Babakhani F, Citron DM. Antimicrobial activities of fidaxomicin. Clin Infect Dis.
2012;55 Suppl 2:S143-8.
29.   Knight DR, Squire MM, Collins DA, Riley TV. Genome analysis of *Clostridium difficile* PCR
ribotype 014 lineage in Australian pigs and humans reveals a diverse genetic repertoire and
signatures of long-range interspecies transmission. Front Microbiol. 2016;7:2138.
30.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI
analysis of 90K prokaryotic genomes reveals clear species boundaries. Nature Communications.
2018;9.
31.   Inouye M, Dashnow H, Raven LA, Schultz MB, Pope BJ, Tomita T, et al. SRST2: Rapid genomic
surveillance for public health and hospital microbiology labs. Genome Med. 2014;6(11):90.
32.   Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, et al. ARG-
ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes.
Antimicrob Agents Chemother. 2014;58(1):212-20.
33.   Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. Artemis: an integrated platform for
visualization and analysis of high-throughput sequence-based experimental data. Bioinformatics.
2012;28(4):464-9.
34.   Spigaglia P. Recent advances in the understanding of antibiotic resistance in *Clostridium
difficile* infection. Ther Adv Infect Dis. 2016;3(1):23-42.
35.   Isidro J, Santos A, Nunes A, Borges V, Silva C, Vieira L, et al. Imipenem resistance in
Clostridium difficile ribotype 017, Portugal. Emerg Infect Dis. 2018;24(4):741-5.
36.   Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A, et al. CARD 2020:
antibiotic resistome surveillance with the comprehensive antibiotic resistance database. Nucleic
Acids Res. 2020;48(D1):D517-D25.
37.   Valiente E, Cairns MD, Wren BW. The *Clostridium difficile* PCR ribotype 027 lineage: a
pathogen on the move. Clin Microbiol Infect. 2014;20(5):396-404.
38.   He M, Miyajima F, Roberts P, Ellison L, Pickard DJ, Martin MJ, et al. Emergence and global
spread of epidemic healthcare-associated *Clostridium difficile*. Nat Genet. 2013;45(1):109-13.
39.   Prakobsrikul N, Malathum K, Santanirand P, Chumnumwat S, Piebpien P, Montakantikul P.
Correlation between antimicrobial consumption and the prevalence of carbapenem-resistant
Escherichia coli and carbapenem-resistant *Klebsiella pneumoniae* at a university hospital in Thailand.
J Clin Pharm Ther. 2019;44(2):292-9.

40.   Knight DR, Imwattana K, Kullin B, Guerrero-Araya E, Paredes-Sabja D, Didelot X, et al. The
Clostridioides difficile species problem: global phylogenomic analysis uncovers three ancient,
toxigenic, genomospecies. BioRxiv. 2020.
41.   McDonald LC, Gerding DN, Johnson S, Bakken JS, Carroll KC, Coffin SE, et al. Clinical practice
guidelines for *Clostridium difficile* infection in adults and children: 2017 update by the Infectious
Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). Clin
Infect Dis. 2018;66(7):987-94.
42.   Wasels F, Monot M, Spigaglia P, Barbanti F, Ma L, Bouchier C, et al. Inter- and intraspecies
transfer of a *Clostridium difficile* conjugative transposon conferring resistance to MLS$_B$. Microb Drug
Resist. 2014;20(6):555-60.
43.   Knight DR, Kullin B, Androga GO, Barbut F, Eckert C, Johnson S, et al. Evolutionary and
genomic insights into *Clostridioides difficile* sequence type 11: a diverse zoonotic and antimicrobial-
resistant lineage of global one health importance. mBio. 2019;10(2).
44.   Solomon K, Fanning S, McDermott S, Murray S, Scott L, Martin A, et al. PCR ribotype
prevalence and molecular basis of macrolide-lincosamide-streptogramin B (MLS$_B$) and
fluoroquinolone resistance in Irish clinical *Clostridium difficile* isolates. J Antimicrob Chemother.
2011;66(9):1976-82.
45.   Kager L, Liljeqvist L, Malmborg AS, Nord CE. Effect of clindamycin prophylaxis on the colonic
microflora in patients undergoing colorectal surgery. Antimicrob Agents Chemother. 1981;20(6):736-
40.
46.   Freeman J, Vernon J, Pilling S, Morris K, Nicholson S, Shearman S, et al. The ClosER study:
results from a three-year pan-European longitudinal surveillance of antibiotic resistance among
prevalent *Clostridium difficile* ribotypes, 2011-2014. Clin Microbiol Infect. 2018;24(7):724-31.
47.   Gonzales M, Pepin J, Frost EH, Carrier JC, Sirard S, Fortier LC, et al. Faecal pharmacokinetics
of orally administered vancomycin in patients with suspected *Clostridium difficile* infection. BMC
Infect Dis. 2010;10:363.
48.   Chotiprasitsakul D, Santanirand P, Thitichai P, Rotjanapan P, Watcharananan S, Siriarayapon
P, et al. Epidemiology and control of the first reported vancomycin-resistant *Enterococcus* outbreak
at a tertiary-care hospital in Bangkok, Thailand. Southeast Asian J Trop Med Public Health.
2016;47(3):494-502.
49.   Hemapanpairoa J, Changpradub D, Thunyaharn S, Santimaleeworagun W. Vancomycin-
resistant enterococcal infection in a Thai university hospital: clinical characteristics, treatment
outcomes, and synergistic effect. Infect Drug Resist. 2019;12:2049-57.