Title

Phylodynamic analysis of an emergent *Mycobacterium bovis* outbreak in an area with no

previously known wildlife infections

Authors

Gianluigi Rossi[1], Joseph Crispell[2], Tanis Brough[3], Samantha J. Lycett[1], Piran C. L. White[4],

Adrian Allen[5], Richard J. Ellis[6], Stephen V. Gordon[2,7], Roland Harwood[8], Eleftheria

Palkopoulou[3], Eleanor L. Presho[5], Robin Skuce[5], Graham C. Smith[9], and Rowland R. Kao[1*]

[1] Roslin Institute and Royal (Dick) School of veterinary Studies, University of Edinburgh,

EH25 9RG, UK

[2] School of Veterinary Medicine, University College Dublin, Dublin D04, Ireland

[3] Department of Bacteriology, APHA, Weybridge, New Haw, Surrey, KT15 3NB

[4] Department of Environment and Geography, University of York, Wentworth Way, York

YO10 5NG, UK

[5] Bacteriology Branch, Veterinary Sciences Division, Agri-food and Biosciences Institute,

Belfast, BT4 3SD, UK

[6] Surveillance and Laboratory Services Department, APHA Weybridge, New Haw, Surrey,

KT15 3NB, UK

[7] UCD Conway Institute, University College Dublin, Dublin D04, Ireland

[8] Veterinary Service, Department of Agriculture, Environment and Rural Affairs, Northern

Ireland, UK

[9] National Wildlife Management Centre, APHA, Sand Hutton, York, YO41 1LZ, UK

*Corresponding author: Rowland.Kao@ed.ac.uk

Abstract

Understanding how an emergent pathogen successfully establishes itself and persists in a

previously unaffected population is a crucial problem in disease ecology. In multi-host

pathogen systems this problem is particularly difficult, as the importance of each host species

to transmission is often poorly characterised, and the epidemiology of the disease is complex.

Opportunities to observe and analyse such emergent scenarios are few.

Here, we exploit a unique dataset combining densely-collected data on the epidemiological

and evolutionary characteristics of an outbreak of *Mycobacterium bovis* (*M. bovis,* the

causative agent of bovine tuberculosis, bTB) in a population of cattle and badgers in an area

considered low-risk for bTB, that has no previous record of either persistent infection in

cattle, or of any infection in wildlife.

We analyse the outbreak dynamics using a combination of mathematical modelling, machine

learning and Bayesian evolutionary analyses. Comparison to *M. bovis* whole-genome

sequences from Northern Ireland confirmed this to be a single introduction of the pathogen

from the latter region, with evolutionary analysis supporting an introduction directly into the

local cattle population at least six years prior to its first discovery in badgers. Once

introduced, the evidence supports *M. bovis* epidemiological dynamics passing through two

phases, the first dominated by cattle-to-cattle transmission before becoming established in the

local badger population.

These findings emphasise the importance of disease surveillance for early containment of

outbreaks, in particular for pathogens not causing immediately evident symptoms in the

infected host, and highlight the utility of combining dynamic modelling and phylogenetic

analyses for understanding the often complex infection dynamics associated with emergent

outbreaks.

1. Introduction

Pathogens able to spread at the interfaces between livestock, wildlife, and humans are one of the most serious threats to human health, wildlife conservation, and livestock economic sustainability [1,2]. Generally, the spread of a pathogen is enhanced when it co-circulates in multiple sympatric host species, as interspecific and intraspecific transmissions can complement each other, resulting in pathogen persistence [3,4].

M. bovis, a member of the *Mycobacterium tuberculosis* complex (MTBC) [5], is responsible for bovine (or animal) tuberculosis (bTB) in domestic cattle and a range of wild mammals [6], including European badgers and deer in Great Britain and Ireland [7–9], deer and wild boar in the Iberian Peninsula [10,11], deer and elk in Michigan, US [12], possums in New Zealand [13,14], and water buffalo in South Africa [15].

In Great Britain and Ireland several studies have established an association between the presence of infected badger populations and the persistence of bTB in cattle [16–18]. More recently, researchers have been able to demonstrate that the same *M. bovis* strains are co-circulating in domestic cattle and sympatric badgers in endemic bTB areas, first using the pathogen's DNA genotyping techniques [19,20], and later using whole-genome sequencing [21,22]. Despite the efforts made by governments to control and eradicate bTB, the last decades have seen an increase in the number of cases and substantial expansion of bTB endemic areas, in particular in England and Wales [5,23]. The eradication efforts of this disease in England alone, costs the UK government around 100 million pounds per year [8,24,25].

Collecting reliable and up-to-date data about wildlife populations can present many challenges at broader scales [26] and while over the years many studies focused on characterising specific badger populations (see [27], and references therein), the lack of information in some areas might prevent the design of effective disease control practices

when bTB is introduced. In addition, broad surveys across England have shown that, since

the mid-1980s, the estimated number of badger social groups has been increasing by 2.6%

annually, contributing to the uncertainty around their level of contributions potential as bTB

reservoir in different regions [28]. However, reliable estimates of badger density, movement,

and potentially infectious contact patterns are poorly recorded in most regions of Great

Britain and Ireland, with only few populations such as at Woodchester Park (Gloucestershire,

England) subject to denser sampling, in this case since the 1980s [29].

Further complications arise from the difficulty of estimating the true prevalence of *M. bovis*

infection in badgers, as well as in domestic cattle. First is the elusive nature of *M. bovis*: the

bacillus is characterised by slow replication with the potential for latent periods of variable

length within the host [30,31]. In addition, the accuracy of currently available diagnostic tests

is suboptimal in both cattle [32] and badgers [33]. These factors contribute to obscuring the

relative roles of the two species in bTB maintenance and spread, and hampering the control

and surveillance strategies. In particular, if both species are able to maintain the disease,

control efforts focused on only one will be ineffective in achieving eradication [22].

One of the main goals of the current control and eradication strategy in GB is to prevent bTB

from becoming established in new non-endemic areas, in particular those officially

categorised as "low-risk" [34]. Within the Low Risk Area of England (LRA), the eastern part

of the Cumbria county in north-west England (hereafter referred to as 'East Cumbria'), has

recently experienced a bTB outbreak of unusual magnitude and duration for this area. The

outbreak began in 2014 and by mid-2019, through enhanced TB surveillance testing of all

cattle herds in the affected area of East Cumbria, it had resulted in the detection of 39

breakdowns (positive cattle herds) across 33 premises [35]. The outbreak was caused by a

strain of *M. bovis* (genotype) new to England, but previously observed in Northern Ireland,

which was shown by preliminary molecular analyses to be the likely area of origin (Skuce,

personal communication). Surveillance of 'found dead' wildlife (badgers and wild deer) for

M. bovis infection was initiated in September 2016 in the area by the Animal and Plant

Health Agency (APHA) [35]. By August 2018 three (out of 52 inspected) roadkill badgers

had been found to be infected, all of them with the same bacterium genotype previously

isolated from local cattle herds. As a result of this epidemiological link, a badger cull was

licensed in a defined area within the affected part of East Cumbria with the aim of eradicating

the disease in badgers and cattle. During the first season of culling operations in the autumn

of 2018, 11% of all the removed badgers were found to be infected, all animals with the same

genotype, suggesting that *M. bovis* infection has become established in the local badger

population.

The aim of this study was to shed light on the dynamic spread of *M. bovis* when introduced in

a two-host system in a non-endemic area, for which the outbreak in East Cumbria provided

us with a unique opportunity to closely study. Here, we describe the East Cumbria outbreak's

spatial and temporal characteristics, and identify the factors which led *M. bovis* infection

becoming established in a wildlife population, while estimating the number of intraspecies

and interspecies transmission events. Our approach includes the use of forensic molecular

epidemiology [36], since over 60 isolates of *M. bovis* from the outbreak with usable whole

genomic sequences were available at the time of writing; complete with precise metadata

including dates, locations and, for cattle, animal and farm identifications.

Results of this study are an important step toward a deeper understanding of bTB

introduction and establishment into non-endemic areas, thus assisting with the process of

disease risk assignment and future policy decision making by the animal health authorities.

2. Results

2.1.Outbreak description

In November 2014, typical bTB lesions were detected via routine slaughterhouse inspection of a seven month-old male calf from a dairy herd in East Cumbria. Bacteriological culture of the lesions yielded an unusual genotype of *M. bovis*, designated 17:z by APHA (Figure 1, A). Following this first report, 23 more cattle were confirmed to be infected with the same strain in East Cumbria, with the last of these detected in November 2018 (at the time of writing). Further cattle were declared bTB positive (using the tuberculin skin test and/or supplementary interferon-gamma blood tests) during this period in the outbreak area, although an *M. bovis* bacilli could not be isolated. The 24 cattle infected with *M. bovis* 17:z genotype included three animals detected outside the outbreak area but still in Cumbria, as well as three in the neighbouring counties of Lancashire (two) and Yorkshire (one), all deemed likely to be part of the same outbreak due to associations through contact tracing. The index animal in this outbreak was a homebred calf that had never left its birth farm until it was moved to slaughter. Therefore, this animal could not have been the "case zero" of this outbreak. We attempted to trace back the first infected individual introduced in the outbreak area by analysing all the Cattle Trace System dataset records that included animals born in Northern Ireland or in the Republic of Ireland from 2009 to 2014. Tracing back the direct movements from Northern Ireland to the outbreak area indicated a limited number of "first arrival" premises (on average 9.3 per year, range 5-15), but unfortunately it did not reveal an obvious first introduction. Conversely, searching for indirect links between Northern Irish farms by selecting other British farms with links to the Cumbrian outbreak which previously imported animals from Northern Ireland, provided too many potential "arrival" premises (on average 216.5 and range 165-247, farms in the outbreak area per year).

During 2016 and 2017, respectively, two and 35 roadkill badger carcasses were reported to

the local authorities within the designated outbreak area of East Cumbria. Three of the badger

carcasses, retrieved respectively in January, February, and April 2017, were positive for *M.*

bovis on culture (while two carcasses were unsuitable for inspection) (Figure S1.1), and all

three positive animals were infected with the 17:z genotype.

The identification of infected badgers led Defra, following a public consultation, to issue a

badger culling licence in a specified section of the outbreak area in the autumn of 2018.

Culling operations from September to November 2018 resulted in 602 culled badgers, of

which 369 were submitted for post mortem inspection and laboratory testing (Defra 2019). In

total, 42 were culture positive for *M. bovis* and of those 38 isolates yielded a whole genome

sequence of sufficient quality to enable phylogenetic analysis [37].

Data on found-dead surveillance of 2018 and 2019, but prior to the second culling season,

included an additional 42 retrieved carcasses (29 and 13, respectively), all negative to *M.*

bovis, except for 15 that were unsuitable for post-mortem inspection, and three still pending

at the time of data gathering.

2.2. Outbreak phylogeny and Northern Ireland isolates

The phylodynamic tree of the East Cumbria outbreak is reported in Figure 1 (B). Early

analyses identified a genotype usually found in Northern Ireland; further evidence showed

the existence of cattle movements from this area to England and Wales. Coincidentally, the

origin area included the recently completed Test, Vaccinate or Remove (TVR) trial area [38]

in Northern Ireland (Skuce, personal communication) where extensive *M. bovis* whole

genome sequencing had already been done – these isolates were included in the current

analyses.

The complete phylogenetic tree (Figure 2) confirmed the association between the *M. bovis*

circulating in the Northern Irish TVR area and in East Cumbria; thus it appeared that the East

Cumbrian outbreak likely originated from the dominant strain circulating in or around the

TVR area (Figure 2, orange branches) imported by movement of infected cattle, though the

first introduction was not identified.

2.3. Epidemiological signatures in genetic data

Following our previous approach [22], epidemiological signatures in the sequence distances

were identified using Boosted Regression Trees (BRT) [39], which combines decision trees

and boosting techniques [40].

As previously, the dependent variable was the genetic distance between *M. bovis* strains,

expressed as single nucleotide variants (SNVs); for explanatory variables we calculated 18

relational covariates for each pair of sampled animals. These covariates are listed in Table 1

and are divided into four categories: temporal (1 covariate), spatial (2 covariates), group (3

covariates), and contact networks (12 covariates).

The BRT model on the full dataset (1,711 observations) was able to explain 41% (pseudo R^2)

of the variability on the test dataset, while the Root Mean Squared Error (RMSE) was 0.94.

To test the robustness of this model we ran the same analysis on the same dataset but with an

increasing percentage of randomly reassigned values for the dependent variable (SNV

distance) observations. The results (Figure S3.1) showed that even for a limited percentage of

re-assigned data (10%) the model underperformed substantially, explaining only 28% of the

variation on average. The full model results (Figure 3) showed that the most important

covariate to explain the SNV distance between isolates was the time between isolate

samplings (25.2%), followed by the hosts populations' size (14.3%) and the geographical

distance (considering both isolate sampling locations and between isolates land parcels,

respectively 11.3 and 9.3%). Among the contact network covariates, the degree in multi-layer

(9.2%) and land parcels networks (9.1%) were the most important, while the covariates

related to the single-species networks were not important (all lower than 4%). Partial

dependency plots are reported in Figure S3.1.

We ran a further three BRT models: two of them considered the within-species interactions

only, cattle-to-cattle (153 observations), and badger-to-badger (820 observations).

Conversely, the third model considered cross-species interactions (738 observations). These

models were able to explain respectively 16%, 43% and 36% of the variation, while the

respective RMSE values were 0.82, 1.04, and 0.96.

The covariates influence rankings in the badger and inter-species models were similar to the

full model (sampling time distance the most influent, respectively, 19.6 and 26.7%), with the

exception that in both cases the degree in the land parcels network was the third most

influential covariate (11.4 and 12.0%). In the badger model the population size dropped to

sixth (9.7%), while the geographic distance between sampling locations was second (14.5%).

The cattle model showed the most differences with the full model, with group size and degree

in the land-parcel network being the most influential covariates (22.2 and 12.9%), while

sampling time distance was third (11.0%).

2.4. Pairwise transmission probability and most likely transmission tree

The pairwise transmission probability was calculated using the Kolmogorov Forward

Equations (KFEs)[41], and the pairwise transmission probability matrix was reported in

Figure 4. The KFEs methodology was used to calculate the probability of observing a pair of

bTB infected hosts given their sampling time and the genetic divergence between their

bacterial isolates, assuming that a direct transmission occurred between the pair.

The animal that infected the first detected cattle (C1) with *M. bovis* might have escaped

detection, since the transmission probabilities from other sampled animals are low (median

0.13×10^{-3}, range $0 - 1.98 \times 10^{-3}$). Similarly, two out of three roadkill badgers (B44 and B18)

had, respectively, the lowest and the third lowest average and maximum transmission

probability from other sampled animals (see Figure 5). This indicates that animals infected

early in the outbreak likely escaped detection (either the 4-year herd testing or carcass

inspection), in particular the "case zero" (i.e. the first cow imported infected with the 17:z

genotype of *M. bovis*).

In general, within-species transmission probabilities were higher than between-species ones

(Figure S4.1), and the phylogenetic root-to-tip temporal signal was strong as well ($R^2 = 0.39$,

p-value ~0; see Figure S4.2). This was consistent with the BRT model results, where the

temporal signal was identified as the most important factor to predict the SNV distance.

We computed 10,000 "random trees" built by selecting random transmissions (except the

ones for which the probability was zero because the cattle's lifespans did not overlap).

Results showed that random trees had a median [95th quantile] of 17[13–21] cattle-to-cattle,

26[21–31] badger-to-badger, 14[10–19] cattle-to-badger, and 6[2–10] badger-to-cattle

transmissions (Figure S4.3). We computed the most likely transmission tree by selecting the

transmissions with highest probability within the sampled animals while avoiding loops in

the tree (see [41]). The best tree (Figure 4) showed that most of the transmissions in this

system likely happened within-species, i.e. 20 cattle-to-cattle and 29 badger-to-badger,

respectively. Conversely, inter-species transmissions comprised 12 cattle-to-badgers

incidents, and 3 badgers-to-cattle. When comparing these results with the randomly

computed trees, the most-likely tree showed a lower number of cross-species transmissions

and a higher number of within-species transmissions, although the estimates fell in the 95th

quantile of the random trees ones.

2.5. BASTA analysis

The transmission rates estimated by Bayesian Structured coalescent Approximation

(BASTA) [42] on the 10 different sub-samples suggest that transmission from cattle-to-

badgers occurred much more frequently (at least an order of magnitude) than transmission

from badgers-to-cattle in Cumbria (Figure 7). In addition, there is little support for the

inclusion of badgers-to-cattle transmission in the structured population model. In contrast,

there is strong support for transmission of *M. bovis* from the sampled Northern Ireland area

into the Cumbria area via the cattle population. Figure S5.1 shows the rate estimates that

BASTA produced when no genomic data was provided, therefore only sampling dates were

used. These analyses were conducted to determine how much signal there was in the genomic

data to support the transmission rates being estimated. Given the contrasting rates shown in

Figure 7 and Figure S5.1, there was strong evidence that there was sufficient signal in the *M.*

bovis genomic data to estimate the transmission rates. Lastly, there is good agreement across

the 10 sub-samples, suggesting that the estimated rates were robust to any inter-sub-sample

variation.

Analyses in BASTA leveraging the temporal signal in the *M. bovis* genomic data were used

to estimate the timing of *M. bovis* transmission from the cattle in the TVR area to cattle in

Cumbria (Figure S4.4). Whilst credible intervals around these estimates were very broad, the

transmission event was estimated to have occurred in March 2011 (lower 2.5% bound

estimate: August 2001; upper 97.5% bound estimate: April 2014). This reflect the slow and

variable replication rate characteristic of *M. bovis*.

3. Discussion

While it is accepted that cattle movements are responsible for transmission on a wide spatial scale [23,43,44], uncertainty remains around the relative roles of cattle and badger in maintaining and spreading *M. bovis* infection at local scales, complicating the formulation and execution of control policies. Our analyses of the East Cumbria outbreak highlight this dynamic, with the introduction of *M. bovis* most likely being caused by a cattle movement from Northern Ireland, followed by a more complex spread among local cattle and badgers. Identifying the source of infection for the index case in cattle in this outbreak (marked as C1 in the figures) would be a crucial piece of information. Should a badger be the most likely infection source then this would imply an earlier establishment of the disease in wildlife (i.e. before or during 2014). Our results suggest another infected bovid as the most likely bTB source for the index case, but the transmission from known cases is poorly supported as the transmission probabilities from the other sampled animals to the index are generally low (Figure 5).

Overall, both the likely transmission tree selection (obtained with the KFEs) and the BASTA analysis indicate that most transmissions happened within-species, and that cattle-to-badger transmission has played a more important role than badger-to-cattle transmission, with similar outcomes obtained using two different analyses.

Our hypothesis, which is supported by the described results, is that after an initial seeding of *M. bovis* into the local East Cumbria cattle population, the infection subsequently became established in local badgers. Once that happened, rapid spread of the pathogen led to the observed 11% prevalence in the autumn of 2018, with peaks up to 20.9% in the core area [37]. Given the historically low prevalence of *M. bovis* in this area [45], the immunological naivete of this specific badgers population might have facilitated a quicker spread, although more research in this direction is needed. From a disease dynamic point of view, the

relatively higher number of predicted cattle-to-badger transmission events suggests that the

establishment in badgers may happen when infection pressure from the sympatric cattle

population reaches a threshold, rather than a single transmission event.

A concern might be that transmission tree outcomes have been affected by the sampling

timeline, since early in the outbreak there are less *M. bovis* sampled sequences and most

come from cattle, while later on the majority of sampled isolates come from badgers. While

the BASTA analysis is designed to specifically reduce the effects of unbalanced sampling

[42], our conclusions are also supported by a low genetic diversity in the recovered badger *M.*

bovis isolates, which points to a relatively recent outbreak in this population. Nonetheless, the

BASTA analysis was able to exploit this low genetic diversity, as it is shown by the different

results of this model when neglecting the genetic information.

Similar to a previous study in a separate population [22], the Boosted Regression Trees

(BRT) analysis indicated temporal bacterial isolation differences, and spatial distance

between hosts, as the most important factors to predict inter-*M. bovis* genetic distance

(SNVs). Another important variable was host population sizes, with bigger groups (farms or

setts) linked to higher SNV distances. This result was particularly strong for the BRT models

which included cattle isolates, and it is consistent with farms size being a risk factor for bTB

[46,47].

Including different contact networks metrics variables in the BRT model can help to

understand which transmission mechanism is more relevant in this system. Among these

variables, the number of adjacent land parcels, corresponding to the spatial network's degree,

was equal (full model) or more important (cattle, badger and inter-species models) than other

metrics. This points to fine-scale spatial effects which cannot be entirely explained by the

simple distance between samples locations. On the contrary, the single-species network

metrics (animal movements for cattle and sett adjacency for badgers) were not significant in

any models. This might be the result of complex interactions between the two species that

cannot be explained when considered individually, as previously suggested [48]. However,

these interactions become evident when both species contact networks are included in a

single framework such as multi-layer network [49].

The badger-to-badger model was able to predict the SNV distances in the test dataset slightly

better than the full model (43% to 42%), despite reducing the sample size (from 1,711

observations in the full model, to 820 in the badger only model). Furthermore, we observed

that the SNV distance variability explained by the cattle only model was comparable to that

of the full model when 20% of the data were randomly shuffled. This was surprising since

more data are collected on domestic animals and this should provide a better picture than the

equivalent wildlife data. One potential explanation is that the selected variables for this

model might better explain *M. bovis* dynamics in badgers than in cattle, at the local scale.

However, when we consider this and the good explanatory power of the land parcels network

metrics, we could speculate that the cattle dataset might hide some contact patterns. This

could occur due to the inability to isolate *M. bovis* in all herd breakdowns or in all cattle, or

due to unrecorded movements, with cattle grazing in several land parcels belonging to the

same farm but not contiguous to the farm [50,51]. In general, the landscape and the farmland

fragmentation and distribution in space, which is ignored when considering farms' main

building locations only, might play a crucial role in disease dissemination. This calls for

surveillance and control strategies to be adapted to specific contexts and informed by detailed

veterinary investigation and spatial information, such as land parcel distribution.

By comparing the East Cumbria outbreak with other bTB in other contexts leads to some

important insights. In East Cumbria, badgers have played a lesser role in the local persistence

of the disease, the converse to what observed in a bTB endemic area in Woodchester Park

[22]. This may be due to the "age" of the outbreak. Whilst in newly infected areas, and

therefore non-endemic, cattle-to-cattle and cattle-to-wildlife transmissions may dominate, in

endemic areas the dynamic may have shifted towards a more complex dynamic, where

wildlife can play a more important role. While in both cases breaking the transmission at the

wildlife/livestock interface is critical, in outbreaks within low risk areas (non-endemic) it is

crucial to prevent the establishment of bTB.

Questions remains as to how likely bTB is to get established in non-endemic areas, and how

long it would take to detect it. From this perspective, the fact that the first introduction in this

area (estimated to be 2011), and that the initial infected cattle were not detected until

inspection at slaughter, indicates limitations of the bTB surveillance strategy in low risk areas

and the importance of good biosecurity to reduce the risk of onward transmission to wildlife

from introductions of cattle with undetected *M. bovis* infection. In order to mitigate this risk,

in April 2016 Defra adopted mandatory post movement bTB testing of all cattle moved from

high-risk to lower-risk areas [52].

To conclude, our analyses of the recent East Cumbria outbreak highlight how the

transmission dynamics of *M. bovis* can change during the establishment in a non-endemic

area, and how these changes affect the relative roles of wildlife and domestic animals in

establishing and maintaining the infection. Our results suggest how genomic data and

phylogenetic analyses are becoming fundamental to disentangle bacterial pathogen outbreaks,

in particular when combined with epidemiological and network models [53]. Therefore,

infrastructures for genomic surveillance can definitely help inform bTB and other endemic

diseases control policies.

Finally, we highlight how local spatial dynamics might affect pathogen spread during the

early phases of an outbreak. This makes the case for different diseases control strategies in

endemic and non-endemic areas that take into account detailed characteristics about the

system landscape and rearing practices, and for improving biosecurity, in order to achieve

minimal transmission at the cattle/wildlife interface and therefore preventing the

establishment of newly introduced diseases.

Methods

4.1. Data description

4.1.1. Sequences and metadata

Test positive cattle, found-dead badgers and most culled badgers in the Cumbria area were subject to post-mortem and culture of suitable tissues at the Animal and Plant Health Agency (APHA), with positive cultures subjected to genotyping and whole genome sequencing at the Central Sequencing Unit in Weybridge. 65 *M. bovis* whole-genome sequences were available from East Cumbria (65 in total, 24 from cattle, 3 from roadkill badgers, and 38 from culled badgers). The sampling timeline of the available sequences is reported in Figure S1.1. The metadata included a unique identifier, the sampling date, location coordinates (for badger isolates), and the farm's county-parish-holding (CPH) code (for cattle sequences only). The isolates and raw sequence data were processed using the same pipeline as described by Crispell et al. [22].

The East Cumbria outbreak dataset included a further sequence from the same *M. bovis* genotype sampled in Scotland (Figure S1.2). The epidemiological investigation showed that the animal was imported from Northern Ireland for slaughter only, thus we did not consider it in the analyses.

4.1.2. Badger population

A total of 160 badger setts were identified in and around the outbreak area in 2017/18 by the APHA, 117 of them were in the 2018 culling permit area. Badgers were both shot and trapped, with trapped animals subjected to post-mortem analysis, and population data (number of badgers removed, TB positive, negative, and TB status not determined) were available for 99 setts.

4.1.3. Cattle population and outbreak area definition

To obtain all infected cattle life histories (movements, birth and death) we first matched the sequences' unique identifier in the SAM dataset, then we extracted the data from the Cattle Tracing System (CTS) using the animal unique identifier.

The outbreak area was defined as the area within the minimum circle around the sequences sampling locations and all 160 badger setts, and adding to that all the parcels assigned to the infected farms which are contiguous to the above described circle (Figure 1, A). This study included all the farms active between 01/01/2010 and 31/12/2018 (which reported any cattle movements, births or deaths) directly located in the area or which owned a parcel in the outbreak area. The total number of selected cattle farms was 336.

4.1.4. Northern Ireland TVR data

The Test, Vaccinate or Remove (TVR) trial in Northern Ireland ran from 2014 to 2018, and it was designed to determine whether a combination of vaccination and an animal side TB test could be an effective means of controlling *M. bovis* infection in badger populations. During this period *M. bovis* samples were taken from infected cattle and badgers in the area for culturing.

All positive cultures were sequenced at the Agri-Food and Biosciences Institute in Belfast (AFBI-NI). In addition, archived *M. bovis* isolates that were stored as part of routine surveillance operations in the TVR area prior to the start of the trial were selected for sequencing. These additional isolates were sourced from routine test and slaughter surveillance of cattle or road killed badgers.

From the TVR area in Northern Ireland, there were 544 *M. bovis* genomes sourced from infected cattle (479) and badgers (65), sampled from 1996 to 2017. The distribution of sampling times for the *M. bovis* genomes is shown in Figure S1.1.

4.2. Epidemiological information and machine learning analysis

An important part of this analysis involved investigating the population, temporal, spatial, and contact network signatures in the sampled *M. bovis* genomic data [22]. We computed the inter-sequence Single Nucleotide Variants (SNV) distance and then we fitted a Poisson regression model using the Boosted Regression Trees [39] model in R [54].

As mentioned above, we used population, temporal, spatial, and network covariates, which are listed in Table 1. Network covariates (shortest distance, same community, degree, and number of infected nodes in the shortest path) were calculated for three contact networks: single species, spatial, and multi-layer.

The *single species network* accounted for only the within-species potential contacts. The cattle population's nodes corresponded to farms, and edges correspond to movements. In order to build these networks for each pair of cattle sequences, we only accounted for movements spanning from the previous year of the first of the two sequences sampled, to the year of the second one (i.e. if the two sequences were sampled respectively in 2015 and 2017, we computed the network by using movements recorded between 2014 and 2017). The badger population's nodes corresponded to setts, and since most of the trapped badgers occurred near a sett, their sequences were assigned to the closest one. We used the Voronoi partition (or Dirichlet tessellation) to create the edges between setts, in order to avoid relying on an assumption for the distance at which two setts were connected. In this network configuration the inter-species sequences are considered not connected.

The *spatial network* was built by considering each land parcel as a single node. Two land parcels were considered connected by the proximity criterion: if they shared a border or if the borders were closer than 100 meters. In this case, the degree of a node in this network (i.e. land parcel) corresponds to the number of neighbouring parcels.

Finally, the *multi-layer network* accounted for all the previously described contacts

combined. In this case, nodes are defined as land parcels as well, but on top of the proximity

criterion, two land parcels could also be connected if they include two farms or setts which

were connected in the single species network.

The network communities were defined using the Louvain method which optimises the

modularity, as provided in the R package *igraph* [55].

The model was run on the complete dataset (All Samples, Table 1 and Figure 3), on the cattle

and badger isolates only, and on the inter-species isolates. In all cases the train and test

dataset were half of the observations, while in the cattle dataset we used 60% for training and

40% for testing, given the reduced observations compared to the others. We evaluated the

models using the pseudo-R^2 calculation comparing the observed and predicted SNV distances

on the test datasets, and the Root Mean Squared Error (RMSE). These were both calculated

using the package *caret* [56]. For BRT the relative influence of the covariates is determined

by the times each variable is selected to split the data in a decision tree, which in turn is

weighted by the improvement to the model fit that resulted from that variable being used at

each split [39]. All models were fitted with a 10-fold cross validation.

Preliminary runs of the models were used to tune the BRT parameters in order to improve the

predictions. These parameters were the learning rate, which controls the contribution of each

tree to the final model, and the tree complexity, which corresponds to the number of nodes in

the tree. For the full dataset model the learning rate was set to 0.005 and the tree complexity

to 5. For the other model we set the learning rate to 0.003, 0.009, and 0.0075 for the cattle,

badger and inter-species dataset, respectively and the tree complexity to 7, 1, and 10.

4.3. Pairwise KFE and transmission tree reconstruction

We calculated the transmission probability between pairs of animals where the *M. bovis*

sequences have been sampled. We calculated this probability using the Kolmogorov-Forward

Equations (KFE) methodology [41]. The KFEs consist of a set of ordinary differential

equations which track the probability of a system to be in a given state through time [57–59].

In the pairwise transmission case, the system state was given by the combination of the two

hosts disease progression state and, once an individual is infected, by the number of SNVs on

its *M. bovis* strain. The underlying assumption was that, at the time of infection, the two

pathogen strains found in the source and recipient hosts are identical. After the infection, the

two strains start to replicate, and thus substitution on the pathogen DNA happens at a rate μ

(*substitution rate*), generating SNVs. Because the two strains are diverging, we will call the

SNVs found in a strain sampled in the source host A [recipient host B] as *divergent* SNVs, or

divA[*divB*]. The sum of *divA* and *divB* results in the SNV distance between the two strains.

In order to use this methodology, we have to provide three main pieces of information: the

pathogens sequences, the two hosts sampling time, and an underlying model representing the

disease progression. For bTB, we chose to use a simple Susceptible-Exposed-Infectious

model [23,60], where susceptible can become exposed (or latent) after contact with an

infectious host with *infection rate β*, and exposed hosts move to the infectious state with

transition rate σ (or after a latency period of average *$1/\sigma$*). In this study, we provided the

birth date of the cattle as well, in order to limit the time span where each cattle could have

been infected first. For badgers we assumed a constant death rate, based on the observation

that less than 0.1% of individuals would survive past 8-years of age [27]. The

epidemiological parameters were chosen according the most recent literatures (see [41]), and

in order to account for their variability, for each pairwise transmission we tested a 1'000

combination of randomly selected parameters combinations and chose the one returning the

highest probability.

Following Rossi et al. [41], we assembled the most likely transmission tree by progressively

selecting the pairs with the highest transmission probability, and excluding those not possible

given the previously selected ones (e.g. if A→B and B→C are selected, B→A, C→B and

C→A were going to be a priori excluded).

4.4. BASTA analysis

Given that the phylogenetic evidence suggests a large amount of inter-species transmission is

occurring in East Cumbria, the next critical question is in what direction? We used the

Bayesian Structured coalescent Approximation (BASTA) package [42] with the Bayesian

evolutionary analysis platform BEAST2 (Bayesian Evolutionary analysis by Sampling Trees;

[61]) to estimate *M. bovis* inter-species transmission rates. The BASTA package was able to

estimate these rates whilst accounting for the known structure and sampling biases in the

study population. In the current study, the sampled *M. bovis* population was split into four

different sub-populations based on host species (badger or cow) and location (Cumbria or

TVR) (Figure S4.1). In addition, to estimate transmission rates in a structured population,

BASTA is robust to sampling biases, which are likely to be present in the current *M. bovis*

dataset. Importantly, it is assumed that transmission from TVR to Cumbria only occurred in

one direction, as the Cumbria clade is monophyletic within the larger TVR phylogeny

(Figure 2).

The evolutionary analyses using BASTA require that there is a temporal signal in the

M. bovis genomic data. With the presence of a temporal signal, the accumulation of

substitutions will be tied to the evolutionary processes of the sampled population, making it

possible to leverage genetic variation to estimate evolutionary dynamics such as transmission

rates. A root-to-tip versus sampling time regression was used to determine whether a

measurable temporal signal was present in the *M. bovis* genomic data (Figure S4.2). The

positive trend observed in this regression indicates the presence of a weak temporal signal,

therefore it was possible to proceed with the evolutionary analyses in BASTA.

The computational complexity of the analyses to be conducted using BASTA meant that the

large number of *M. bovis* genomes available within the outbreak clade (orange clade in

Figure 2) had to be sub-sampled.

In addition, it was only from 2014 to 2017 that the sampling of cattle and badgers in the TVR

area can be considered approximately equal (in terms of effort). Therefore, only genomes

sourced from infected cattle and badgers sampled from 2014 to 2018 were included in the

BASTA analyses. The window was extended to 2018 to include the genomes sourced from

Cumbria. The sub-sampling was then weighted to include samples from as many years as

possible and from equal numbers of cattle and badgers. The sub-sampling was conducted 10

times, each time selecting 20 badgers and 20 cattle-derived *M. bovis* genomes from Cumbria

and 40 badger and 40 cattle-derived genomes from the TVR area (an example of a sub-

sample is shown in Figure S4.3). A root-to-tip versus sampling time regression was

conducted for each sub-sample and found to be positive in all cases. Each sub-sample was

then analysed separately in BASTA.

References

1. Gortazar C, Diez-Delgado I, Barasona JA, Vicente J, De La Fuente J, Boadella M. The Wild Side of Disease Control at the Wildlife-Livestock-Human Interface: A Review. Front Vet Sci. 2015;1: 1–12. doi:10.3389/fvets.2014.00027

2. Wiethoelter AK, Beltrán-Alcrudo D, Kock R, Mor SM. Global trends in infectious diseases at the wildlife–livestock interface. Proc Natl Acad Sci. 2015;112: 9662–9667. doi:10.1073/pnas.1422741112

3. Haydon DT, Cleaveland S, Taylor LH, Laurenson MK. Identifying reservoirs of infection: A conceptual and practical challenge. Emerg Infect Dis. 2002;8: 1468–1473. doi:10.3201/eid0812.010317

4. Craft ME, Hawthorne PL, Packer C, Dobson AP. Dynamics of a multihost pathogen in a carnivore community. J Anim Ecol. 2008;77: 1257–1264. doi:10.1111/j.1365-2656.2008.01410.x

5. Smith NH, Gordon S V., de la Rua-Domenech R, Clifton-Hadley RS, Hewinson RG. Bottlenecks and broomsticks: The molecular evolution of *Mycobacterium bovis*. Nat Rev Microbiol. 2006;4: 670–681. doi:10.1038/nrmicro1472

6. Palmer M V. *Mycobacterium bovis*: Characteristics of wildlife reservoir hosts. Transbound Emerg Dis. 2013;60: 1–13. doi:10.1111/tbed.12115

7. Skuce R, Breadon E, Allen A, Milne G, McCormick C, Hughes C, et al. Longitudinal dynamics of herd-level *Mycobacterium bovis* MLVA type surveillance in cattle in Northern Ireland 2003–2016. Infect Genet Evol. 2020;79: 104131. doi:10.1016/j.meegid.2019.104131

8. Godfray HCJ, Donnelly CA, Kao RR, Macdonald DW, McDonald R, Petrokofsky G, et al. A restatement of the natural science evidence base relevant to the control of bovine tuberculosis in Great Britain. Proc R Soc B Biol Sci. 2013;280: 1–18.

doi:10.1098/rspb.2013.1634

9.   Crispell J, Cassidy S, Kenny K, McGrath G, Warde S, Cameron H, et al.

Mycobacterium bovis genomics reveals transmission of infection between cattle and

deer in Ireland. Microb genomics. 2020;6. doi:10.1099/mgen.0.000388

10.   Anderson LG, Gortázar C, Vicente J, Hutchings MR, White PCL. Modelling the

effectiveness of vaccination in controlling bovine tuberculosis in wild boar. Wildl Res.

2013;40: 367–376. doi:10.1071/WR12139

11.   Madeira S, Manteigas A, Ribeiro R, Otte J, Fonseca AP, Caetano P, et al. Factors that

Influence *Mycobacterium bovis* Infection in Red Deer and Wild Boar in an

Epidemiological Risk Area for Tuberculosis of Game Species in Portugal. Transbound

Emerg Dis. 2017;64: 793–804. doi:10.1111/tbed.12439

12.   Salvador LCM, O'Brien DJ, Cosgrove MK, Stuber TP, Schooley A, Crispell J, et al.

Disease management at the wildlife-livestock interface: using whole-genome

sequencing to study the role of elk in *Mycobacterium bovis* transmission in Michigan,

USA. Mol Ecol. 2019; 1–14. doi:10.1111/mec.15061

13.   Crispell J, Zadoks RN, Harris SR, Paterson B, Collins DM, de-Lisle GW, et al. Using

whole genome sequencing to investigate transmission in a multi-host system: bovine

tuberculosis in New Zealand. BMC Genomics. 2017;18: 180. doi:10.1186/s12864-

017-3569-x

14.   Anderson DP, Ramsey DSL, Nugent G, Bosson M, Livingstone P, Martin PAJ, et al. A

novel approach to assess the probability of disease eradication from a wild-animal

reservoir host. Epidemiol Infect. 2013;141: 1509–1521.

doi:10.1017/S095026881200310X

15.   Fitzgerald SD, Kaneene JB. Wildlife Reservoirs of Bovine Tuberculosis Worldwide:

Hosts, Pathology, Surveillance, and Control. Vet Pathol. 2013;50: 488–499.

doi:10.1177/0300985812467472

16.   Martin SW, Eves J a, Dolan L a, Hammond RF, Griffin JM, Collins JD, et al. The

association between the bovine tuberculosis status of herds in the East Offaly Project

Area, and the distance to badger setts, 1988-1993. Prev Vet Med. 1997;31: 113–25.

doi:10.1016/S0167-5877(96)01111-7

17.   Donnelly CA, Woodroffe R, Cox DR, Bourne FJ, Cheeseman CL, Clifton-Hadley RS,

et al. Positive and negative effects of widespread badger culling on tuberculosis in

cattle. Nature. 2006;439: 843–846. doi:10.1038/nature04454

18.   Donnelly CA, Woodroffe R, Cox DR, Bourne J, Gettinby G, Le Fevre AM, et al.

Impact of localized badger culling on tuberculosis incidence in British cattle. Nature.

2003;426: 834–837. doi:10.1038/nature02192

19.   Olea-Popelka FJ, Flynn O, Costello E, McGrath G, Collins JD, O'Keeffe J, et al.

Spatial relationship between *Mycobacterium bovis* strains in cattle and badgers in four

areas in Ireland. Prev Vet Med. 2005;71: 57–70. doi:10.1016/j.prevetmed.2005.05.008

20.   Woodroffe R, Donnelly CA, Cox DR, Gilks P, Jenkins HE, Thomas Johnston W, et al.

Bovine tuberculosis in cattle and badgers in localized culling areas. J Wildl Dis.

2009;45: 128–143. doi:10.7589/0090-3558-45.1.128

21.   Biek R, O'Hare A, Wright D, Mallon T, McCormick C, Orton RJ, et al. Whole

Genome Sequencing Reveals Local Transmission Patterns of *Mycobacterium bovis* in

Sympatric Cattle and Badger Populations. PLoS Pathog. 2012;8.

doi:10.1371/journal.ppat.1003008

22.   Crispell J, Benton CH, Balaz D, De Maio N, Akhmetova A, Allen A, et al. Combining

genomics and epidemiology to analyse bi-directional transmission of *Mycobacterium*

bovis in a multi-host system. Elife. 2019; 1–36.

doi:https://doi.org/10.7554/eLife.45833.001

23.   Brooks-Pollock E, Roberts GO, Keeling MJ. A dynamic model of bovine tuberculosis

614   spread and control in Great Britain. Nature. 2014;511: 228–231.

615   doi:10.1038/nature13529

616   24.   Tildesley MJ, Brand S, Brooks Pollock E, Bradbury N V., Werkman M, Keeling MJ.

617   The role of movement restrictions in limiting the economic impact of livestock

618   infections. Nat Sustain. 2019;2: 834–840. doi:10.1038/s41893-019-0356-5

619   25.   Defra. Bovine TB Eradication Programme for England. 2011.

620   26.   Craft ME. Infectious disease transmission and contact networks in wildlife and

621   livestock. Philosophical Transactions of the Royal Society B: Biological Sciences.

622   2015. pp. 20140107–20140107. doi:10.1098/rstb.2014.0107

623   27.   Roper T. Badger. Collins; 2010. Available:

624   https://books.google.co.uk/books?id=FZhrPgAACAAJ

625   28.   Judge J, Wilson GJ, Macarthur R, Delahay RJ, McDonald RA. Density and abundance

626   of badger social groups in England and Wales in 2011-2013. Sci Rep. 2014;4: 1–8.

627   doi:10.1038/srep03809

628   29.   Delahay RJ, Walker N, Smith GS, Wilkinson D, Clifton-Hadley RS, Cheeseman CL,

629   et al. Long-term temporal trends and estimated transmission rates for *Mycobacterium*

630   *bovis* infection in an undisturbed high-density badger (*Meles meles*) population.

631   Epidemiol Infect. 2013;141: 1445–1456. doi:10.1017/S0950268813000721

632   30.   Pollock JM, Neill SD. *Mycobacterium bovis* infection and tuberculosis in cattle. Vet J.

633   2002;163: 115–127. doi:10.1053/tvjl.2001.0655

634   31.   Cassidy JP. The pathogenesis and pathology of bovine tuberculosis with insights from

635   studies of tuberculosis in humans and laboratory animal models. Veterinary

636   Microbiology. 2006. pp. 151–161. doi:10.1016/j.vetmic.2005.11.031

32.   Nuñez-Garcia J, Downs SH, Parry JE, Abernethy DA, Broughan JM, Cameron AR, et

al. Meta-analyses of the sensitivity and specificity of ante-mortem and post-mortem

diagnostic tests for bovine tuberculosis in the UK and Ireland. Prev Vet Med.

2018;153: 94–107. doi:10.1016/j.prevetmed.2017.02.017

33.   Drewe JA, Tomlinson AJ, Walker NJ, Delahay RJ. Diagnostic accuracy and optimal

use of three tests for tuberculosis in live badgers. PLoS One. 2010;5.

doi:10.1371/journal.pone.0011196

34.   Perrin LD, Harris KA, Reynolds M, Lawes JR, Frost S, Brouwer A, et al. Bovine TB

infection status in cattle in Great Britain in 2017. Vet Rec. 2019;184: 371–378.

doi:10.1136/vr.l1321

35.   Defra. An update on TB surveillance in wildlife. 2019 [cited 26 May 2020] p. 6.

Available:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachme

nt_data/file/830810/surveillance-wildlife-2018.pdf

36.   Kao RR, Haydon DT, Lycett SJ, Murcia PR. Supersize me: How whole-genome

sequencing and big data are transforming epidemiology. Trends Microbiol. 2014;22:

282–291. doi:10.1016/j.tim.2014.02.011

37.   Defra. TB surveillance in badgers during year 1 badger control operations in eastern

Cumbria, Low Risk Area (2018). 2019 [cited 31 Oct 2020]. Available:

https://www.gov.uk/government/publications/bovine-tb-surveillance-in-wildlife-in-

england/tb-surveillance-in-badgers-during-year-1-badger-control-operations-in-

eastern-cumbria-low-risk-area-2018

38.   DAERA-NI. The test and vaccinate or remove (TVR) wildlife intervention research

project - year 2 report. www.dardni.gov.uk/ test-and-vaccinate-or-remove.htm. 2015.

Available: https://www.daera-ni.gov.uk/articles/test-and-vaccinate-or-remove-tvr-

wildlife-intervention-research

39.   Elith J, Leathwick JR, Hastie T. A working guide to boosted regression trees. J Anim

Ecol. 2008;77: 802–813. doi:10.1111/j.1365-2656.2008.01390.x

40.   Brock PM, Fornace KM, Grigg MJ, Anstey NM, William T, Cox J, et al. Predictive

analysis across spatial scales links zoonotic malaria to deforestation. Proc R Soc B

Biol Sci. 2019;286. doi:10.1098/rspb.2018.2351

41.   Rossi G, Crispell J, Balaz D, Lycett SJ, Delahay RJ, Kao RR. Identifying likely

transmission pairs with pathogen sequence data using Kolmogorov Forward

Equations; an application to *M.bovis* in cattle and badgers. BiorXiv. 2020; 1–32.

doi:10.1101/2020.06.11.146894

42.   De Maio N, Wu CH, O'Reilly KM, Wilson D. New Routes to Phylogeography: A

Bayesian Structured Coalescent Approximation. PLoS Genet. 2015;11: 1–22.

doi:10.1371/journal.pgen.1005421

43.   Green DM, Kiss IZ, Mitchell AP, Kao RR. Estimates for local and movement-based

transmission of bovine tuberculosis in British cattle. Proc R Soc B Biol Sci. 2008;275:

1001–1005. doi:10.1098/rspb.2007.1601

44.   Gilbert M, Mitchell A, Bourn D, Mawdsley J, Clifton-Hadley R, Wint W. Cattle

movements and bovine tuberculosis in Great Britain. Nature. 2005;435: 491–496.

doi:10.1038/nature03548

45.   Atkins PJ, Robinson PA. Bovine tuberculosis and badgers in Britain: Relevance of the

past. Epidemiol Infect. 2013;141: 1437–1444. doi:10.1017/S095026881200297X

46.   Salvador LCM, Deason M, Enright J, Bessell PR, Kao RR. Risk-based strategies for

surveillance of tuberculosis infection in cattle for low-risk areas in England and

Scotland. Epidemiol Infect. 2018;146: 107–118. doi:10.1017/S0950268817001935

47.   Brooks-Pollock E, Keeling M. Herd size and bovine tuberculosis persistence in cattle

farms in Great Britain. Prev Vet Med. 2009;92: 360–365.

doi:10.1016/j.prevetmed.2009.08.022

48.   Silk MJ, Drewe JA, Delahay RJ, Weber N, Steward LC, Wilson-Aggarwal J, et al.

Quantifying direct and indirect contacts for the potential transmission of infection

between species using a multilayer contact network. Behaviour. 2018.

doi:10.1163/1568539X-00003493

49.   Kinsley AC, Rossi G, Silk MJ, VanderWaal K. Multilayer and Multiplex Networks:

An Introduction to Their Use in Veterinary Epidemiology. Frontiers in Veterinary

Science. 2020. p. 596. doi:10.3389/fvets.2020.00596

50.   Hamilton L, Evans N, Allcock J. "i don't go to Meetings": Understanding farmer

perspectives on bovine TB and biosecurity training. Vet Rec. 2019;184: 1–8.

doi:10.1136/vr.104995

51.   Campbell EL, Byrne AW, Menzies FD, McBride KR, McCormick CM, Scantlebury

M, et al. Interspecific visitation of cattle and badgers to fomites: A transmission risk

for bovine tuberculosis? Ecol Evol. 2019;9: 8479–8489. doi:10.1002/ece3.5282

52.   Defra. Bovine TB Information Note 03 / 16 Reducing TB risks from the sale of cattle

from 4-yearly testing herds in England. 2016. Available:

https://www.gov.uk/government/publications/bovine-tb-information-note-0116-post-

movement-testing

53.   Campbell F, Strang C, Ferguson N, Cori A, Jombart T. When are pathogen genome

sequences informative of transmission events? PLoS Pathog. 2018;14: 1–21.

doi:10.1371/journal.ppat.1006885

54.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna,

Austria; 2018. Available: https://www.r-project.org/

55.   Csárdi G, Nepusz T. The igraph software package for complex network research. J

Comput Appl. 2014;Complex Sy: 9. doi:10.3724/SP.J.1087.2009.02191

56.   Kuhn M, Weston S, Keefer C, Engelhardt A, Cooper T, Mayer Z, et al. Classification

714   and Regression Training. 2016. p. 198. Available: https://cran.r-

715   project.org/package=caret

716   57.   Stollenwerk N, Jansen VAA. Meningitis, pathogenicity near criticality: The

717   epidemiology of meningococcal disease as a model for accidental pathogens. J Theor

718   Biol. 2003;222: 347–359. doi:10.1016/S0022-5193(03)00041-9

719   58.   Sharkey KJ. Deterministic epidemiological models at the individual level. J Math Biol.

720   2008;57: 311–331. doi:10.1007/s00285-008-0161-7

721   59.   Keeling MJ, Ross J V. On methods for studying stochastic disease dynamics. J R Soc

722   Interface. 2008;5: 171–181. doi:10.1098/rsif.2007.1106

723   60.   Rossi G, De Leo GA, Pongolini S, Natalini S, Vincenzi S, Bolzoni L. Epidemiological

724   modelling for the assessment of bovine tuberculosis surveillance in the dairy farm

725   network in Emilia-Romagna (Italy). Epidemics. 2015;11: 62–70.

726   doi:10.1016/j.epidem.2015.02.007

727   61.   Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, et al. BEAST 2: A

728   Software Platform for Bayesian Evolutionary Analysis. PLoS Comput Biol. 2014;10:

729   1–6. doi:10.1371/journal.pcbi.1003537

732 **Tables**

Category	Covariates	Description	All samples model	Cattle model	Badger model	Inter-species model
Temporal	SampleTimeDist	Time distance (in days) between the isolates sample dates	YES	YES	YES	YES
Spatial	SampleGeoDist	Geographical Euclidean distance between the isolates sampling locations	YES	YES	YES	YES
Spatial	ParcelsDist	Minimum geographical Euclidean distance between the land parcels where isolates have been sampled	YES	YES	YES	YES
Group	SameSpecies	Isolates from same species animals (YES/NO)	YES	NO	NO	NO
Group	SameGroup	Isolates from same herd or sett animals (YES/NO)	YES	YES	YES	NO
Group	Sizes	Size coefficient (squared root of product) of the isolates herd/social groups	YES	YES	YES	YES
Network	ShortestPathSNet	Shortest path length between isolates nodes in single-species network	YES	YES	YES	NO
Network	InfNodesInPathSNet	Number of infected nodes in shortest path between isolates in single-species network	YES	YES	YES	NO
Network	DegreeSNet	Degree coefficient (squared root of product) of isolates nodes in single-species network	YES	YES	YES	NO
Network	SameCommSNet	Isolates' nodes in same community in single-species network (YES/NO)	YES	YES	YES	NO
Network	ShortestPathLPNet	Shortest path length between isolates nodes in spatial (land parcels) network	YES	YES	YES	YES
Network	InfParcInPathLPNet	Number of infected nodes in shortest path between isolates in spatial network	YES	YES	YES	YES
Network	DegreeLPNet	Degree coefficient (squared root of product) of isolates nodes in spatial network	YES	YES	YES	YES
Network	SameCommLPNet	Isolates' nodes in same community in spatial network (YES/NO)	YES	YES	YES	YES
Network	ShortestPathML	Shortest path length between isolates nodes in multi-layer network	YES	YES	YES	YES
Network	InfNodesInPathML	Number of infected nodes in shortest path between isolates in multi-layer network	YES	YES	YES	YES
Network	DegreeML	Degree coefficient (squared root of product) of isolates nodes in multi-layer network	YES	YES	YES	YES
Network	SameCommML	Isolates' nodes in same community in multi-layer network (YES/NO)	YES	YES	YES	YES

734   **Table 1. Epidemiological covariates tested in the BRT algorithm.** List of epidemiological

735   covariates tested in the boosted regression trees (BRT) algorithm against the isolates genetic

736   distance, calculated in number of SNVs. The four columns indicate in which model the

737   covariate has been used.

Figures

Figure 1. East Cumbria outbreak area and phylogenetic tree. A: East Cumbria outbreak

area (black) location in Cumbria (dark grey). B: *M. bovis* genomes phylogenetic tree

(distance calculated in Single Nucleotide Variants, SNVs). Red dots represent badgers, and

blue square cattle, and the label report the code assigned to each individual, sampling

location and sample date.

~ 10 SNPs

● Badger
▲ Cow

Figure 2. East Cumbria and TVR *M. bovis* phylogenies. A maximum likelihood

phylogeny of the *M. bovis* genomes sourced from Cumbria and the Test, Vaccinate, or

Release (TVR) area in Northern Ireland. The tree is rooted with the *M. bovis* reference

genome AF2122/97. The *M. bovis* genomes sourced from infected cattle and badgers in

Cumbria are highlighted with a black semi-circle at the top left. The branches of the clade

containing the *M. bovis* genomes sourced from Cumbria, and those from the TVR area that

are most similar is highlighted in orange.

Figure 3. Covariate influence in the BRT models. The relative influence of the 18

epidemiological covariates calculated by the Boosted Regression Trees (BRT) model. Bars

colours correspond to the four models run with different sub-samples of the dataset: dark-red

for the full model account all samples, light red for the badger-to-badger model, light-blue for

the cattle-to-cattle only model, and dark-blue for the interspecies (badger-to-cattle) model.

Figure 4. Transmission matrix. Pairwise transmission probabilities between infected

animals in the East Cumbria outbreak. Animals are reported in *x* (source animal) and *y*

(infected animal) axes in the order they have been sampled, and they are labelled from one to

65 and the species name (B for badgers and C for Cattle). Different colours correspond to

transmission directions (red: badger-to-badger, green: badger-to-cow, light-blue: cattle-to-

badger, and magenta: cattle-to-cattle).

Figure 5. Transmission probability to sequenced animals. Average (dots/triangles) and

range (line) of infection probabilities (x axis) from all sampled infection sources to each

individual animal (y axis). Red lines/dots correspond to badgers and blue lines/triangles to

cattle.

Figure 6. East Cumbria *M. bovis* outbreak transmission tree. Most likely transmission for

the sampled infectious individuals in the East Cumbria outbreak (transmissions from top to

bottom). Red circles correspond to badgers, and blue squares to cattle. Edge thickness is

proportional to the pair transmission probability and the edge label (dark green) indicates the

SNV distance between the two individuals.

Figure 7. Transmission rates and number of transmissions estimated with BASTA. a)

Inter-species and Northern Ireland to Cumbria *M. bovis* transmission rates (y axes, log scale)

estimated using BASTA, based on analyses that used the genomic data and compared with

the results for Woodchester Park [22]; b) estimates from BASTA of the number of

transmission events (y axes) between the sampled cattle and badgers.