Sensitizing *Staphylococcus aureus* to antibacterial host defense by decoding and blocking

the lipid flippase MprF

Christoph J. Slavetinsky[1,2,3,4], Janna N. Hauser[1,3,4], Cordula Gekeler[1,3,4], Jessica Slavetinsky[1,3,4], André

Geyer[1,3], Alexandra Kraus[5], Doris Heilingbrunner[5], Samuel Wagner[3,4,6], Michael Tesar[5], Bernhard

Krismer[1,3,4], Sebastian Kuhn[1,3], Christoph M. Ernst[1,3,7] & Andreas Peschel[1,3,4]

[1]Department of Infection Biology, Interfaculty Institute for Microbiology and Infection Medicine

Tübingen (IMIT), University of Tübingen, Tübingen, Germany.

[2]Pediatric Gastroenterology and Hepatology, University Children's Hospital Tübingen, Eberhard Karls

University Tübingen, Tübingen, Germany.

[3]German Centre for Infection Research (DZIF), Partner Site Tübingen, Tübingen, Germany.

[4]Cluster of Excellence "Controlling Microbes to Fight Infections," University of Tübingen, Tübingen,

Germany.

[5]MorphoSys AG, Martinsried, Germany.

[6]Section of Cellular and Molecular Microbiology, Interfaculty Institute for Microbiology and Infection

Medicine Tübingen (IMIT), University of Tübingen, Tübingen, Germany.

[7]Present address: Christoph M. Ernst, Department of Molecular Biology and Center for

Computational and Integrative Biology, Massachusetts General Hospital, Harvard Medical School,

Boston, Massachusetts, USA.

Correspondence   to   **Christoph   J.   Slavetinsky**,   University   Children's   Hospital;   Mail:

christoph.slavetinsky@med.uni-tuebingen.de or **Christoph M. Ernst**, Harvard University, Broad

Institute; Mail: cmernst@broadinstitute.org.

Conflict of interest statement

Antibodies disclosed in the manuscript are part of patent "Anti-staphylococcal antibodies" (US9873733B2 / EP2935324B1) from A.P., C.M.E., A.K., M.T., assigned to the Universitaetsklinikum Tuebingen and MorphoSys AG. The authors have declared that no further conflict of interest exists.

Abstract

The pandemic of antibiotic resistance represents a major human health threat demanding new antimicrobial strategies. MprF is the synthase and flippase of the phospholipid lysyl-phosphatidylglycerol that increases virulence and resistance of methicillin-resistant *Staphylococcus aureus* (MRSA) and other pathogens to cationic host defense peptides and antibiotics. With the aim to design MprF inhibitors that could sensitize MRSA to both, human antimicrobials and antibiotics and support the clearance of staphylococcal infections with minimal selection pressure, we developed MprF-targeting monoclonal antibodies, which bound and blocked the MprF flippase subunit. Antibody M-C7.1 targeted a specific loop in the flippase domain that proved to be exposed at both sides of the bacterial membrane, thereby enhancing the mechanistic understanding into bacterial lipid translocation. M-C7.1 rendered MRSA susceptible to host antimicrobial peptides and antibiotics such as daptomycin. Moreover, it impaired MRSA survival in human phagocytes, which recommends MprF inhibitors for new anti-MRSA approaches. MprF-directed monoclonal antibodies provide a proof of concept for development of precisely targeted anti-virulence approaches, which block bacterial antimicrobial resistance mechanisms.

Introduction

The continuous increase of antibiotic resistance rates undermines the significance and efficacy of available antibiotics against bacterial infections [1]. Several opportunistic antibiotic-resistant bacterial pathogens including methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant enterococci, and extended-spectrum beta-lactam or carbapenem-resistant proteobacteria impose a continuously growing pressure on modern healthcare systems [2]. MRSA is responsible for a large percentage of superficial and severe bacterial infections and the available last-resort antibiotics are much less effective than beta-lactams [3]. Unfortunately, no new class of antibiotics has entered the clinical phase since the introduction of the lipopeptide antibiotic daptomycin in 2003

[1]. Novel anti-infective strategies that would circumvent on one hand the difficulties in identifying new microbiota-preserving small-molecule antimicrobials and, on the other hand, the enormous selection pressures exerted by broad-spectrum antibiotics, are discussed as potential solutions against a looming post-antibiotic era [4]. Such strategies could be based for instance on therapeutic antibodies or bacteriophages, which usually have only a narrow activity spectrum. A possible direction could be the inhibition of bacterial targets that are of viable importance only during infection [5]. Blocking such targets by so-called anti-virulence or anti-fitness drugs would preserve microbiome integrity and create selection pressure for resistance-conferring mutations only on invading pathogens. Interfering with bacterial virulence factors should ameliorate the course of infection and enable more effective bacterial clearance by the immune system or by antibiotics.

While anti-virulence and anti-fitness therapies are discussed for several of the notoriously antibiotic-resistant bacterial pathogens, none has passed an early experimental stage, probably because most of the major pharmaceutical companies have strongly reduced their investments into new antibacterials [1, 6]. For instance, a recent approach showed promising effects reducing host colonization and invasion in meningococcal disease using phenothiazines, a family of compounds applied for treatment of psychotic disorders, to target Type IV pili, important bacterial adhesion and virulence factors [7]. Most of the documented anti-virulence approaches appear to rely on small-molecule inhibitors, which are difficult to identify and often have problematic pharmacological, toxicological, and resistance-inducing properties [6]. Monoclonal antibodies (mABs) directed against anti-virulence targets could be interesting alternatives provided the target can be reached by comparatively large antibody molecules. Therapeutic mABs are used in several malignant and inflammatory diseases [8] and have proven efficacy in toxin-mediated bacterial infections such as anthrax or clostridial toxin-mediated diseases [4, 9, 10]. Apart from toxin neutralization, however, mABs have hardly been applied in antimicrobial development programs.

The Multiple Peptide Resistance Factor (MprF), a large integral membrane protein, is crucial for the capacity of bacterial pathogens such as *S. aureus* to resist cationic antimicrobial peptides (CAMPs) of

the innate immune system and CAMP-like antibiotics such as daptomycin [11-13]. MprF is highly

conserved and can be found in various Gram-positive or Gram-negative pathogens [13]. MprF

proteins proved to be crucial for *in vivo* virulence of various pathogens in infection models and when

exposed to human phagocytes [13]. Some parts of the protein are located at the outer surface of the

cytoplasmic membrane and could in principle be reached by mABs [14, 15]. MprF forms oligomers

and it is a bifunctional enzyme, which can be separated into two distinct domains [15]. The C-

terminal domain synthesizes positively charged lysyl-phosphatidylglycerol (LysPG) from a negatively

charged phosphatidylglycerol (PG) acceptor and a Lys-tRNA donor substrate, while the N-terminal

domain translocates newly synthesized LysPG from the inner to the outer leaflet of the cytoplasmic

membrane [14, 16]. The exposure of LysPG at the outer surface of the membrane reduces the affinity

for CAMPs and other antimicrobials [14]. Notably, *mprF* is a major hot spot for point mutations that

lead to daptomycin resistance acquired during therapy of *S. aureus* infections [17].

In order to assess the suitability of MprF as a target for anti-virulence agents we developed mABs

targeting several epitopes of potential extracellular loops of its transmembrane part and analyzed

their capacity to bind specifically to *S. aureus* MprF. We identified a collection of mABs, which did not

only bind to but also inhibited the LysPG flippase domain of MprF. Our results suggest that a specific

loop between two of the transmembrane segments (TMS) of MprF is exposed at both sides of the

membrane suggesting an unusual, potentially flexible topology of this protein part, which may be

involved in LysPG translocation. Accordingly, targeting this loop with a specific mAB inhibited the

MprF flippase function, rendered *S. aureus* susceptible to killing by antimicrobial host peptides and

daptomycin, and reduced *S. aureus* survival when challenged by human CAMP-producing

polymorphonuclear leukocytes (PMNs).

Results

1.   Generation of mABs binding to putative extracellular loops of MprF

The hydrophobic part of *S. aureus* MprF appears to include 14 TMS connected by loops with

predicted lengths between two and 56 amino acids [15]. Several of the loops are located at the outer

surface of the cytoplasmic membrane, accessible to mABs (Figure 1A). Peptides representing four

loops with a minimum length of 13 amino acids were synthesized with N- and C-terminal cysteine

residues to allow cyclization (Table S1). The peptides corresponded to three loops predicted to be at

the outer membrane surface (loops one, nine, and thirteen) and loop seven, the location of which

has remained ambiguous due to conflicting computational and experimental findings [15]. The N-

terminal amino groups of cyclized peptides were linked to biotin to facilitate their recovery and

detection. The antigen peptides were incubated with MorphoSys´s Human Combinatorial Antibody

Library (HuCAL), a phage display library expressing human Fab fragments with highly diverse variable

regions at the phage surface [18]. Antigen-binding phages were enriched in three iterative rounds of

panning in solution and antigen-phage complexes were captured with streptavidin-coated beads. The

bound phages were extensively washed to remove unspecifically binding phages, eluted, and

propagated in *Escherichia coli* for a subsequent panning round. Washing steps were prolonged and

antigen concentrations reduced from round one to round three to increase stringency and discard

antibodies with low specificity and affinity. DNA of the eluted, antigen-specific phages was isolated

and subcloned in specific *E. coli* expression vectors to yield His-tagged Fab fragments. 368 individual

colonies per antigen were picked and Fab fragments were expressed and purified. A representative

selection of 24 unique Fabs against all four peptides were converted to human IgG by cloning in an

IgG1 expression vector system and expression in human HKB11 cells and IgGs were purified via

protein A chromatography, as recently described [18] (see graphical workflow in Fig. S1).

The IgGs were analyzed for binding to the corresponding antigen peptides and also to the three non-

cognate peptides to assess their selectivity, by ELISA with streptavidin-coated microtiter plates (Fig.

S2). Antibodies directed against the four targeted MprF loops were selected based on affinity and

selectivity and analyzed for binding to *S. aureus* SA113 cells expressing or not expressing MprF. Both,

the "wild-type" and *mprF* deletion mutant strains lacked the gene for protein A (*spa*), which would

otherwise unspecifically bind to IgG [19]. Bacteria were adsorbed to microtiter plates, blocked with

bovine serum albumin (BSA), and incubated with IgGs, which were then detected with goat anti-

human IgG after extensive washing. Antibodies M-C1, M-C7.1, M-C7.3, M-C9.1, M-C13.1, and M-

C13.2 bound significantly stronger to MprF-expressing *S. aureus* (SA113Δ*spa*) compared to MprF-

deficient *S. aureus* (SA113Δ*spa*Δ*mprF*), while the humanized isotype control mAB L-1 showed no

specific binding (Fig. 1B). These findings are in agreement with the location of the loops one, nine,

and thirteen at the outer surface of the cytoplasmic membrane, confirming the overall topology of

MprF (Fig. 1A). Of note, loop seven between potential TMS 7 and 8 whose location had previously

remained controversial was detected by antibodies M-C7.1 and M-C7.3 (Fig. 1B) indicating that loop

seven is accessible from the outside. Antibodies M-C1, M-C7.1, M-C9.1, and M-C13.1 showed the

strongest binding to MprF according to mean fluorescence intensities.

2. The MprF epitope bound by M-C7.1 is located at both, outer and inner surface of the

cytoplasmic membrane

The MprF loop seven between TMS 7 and 8 bound by M-C7.1 seemed to have an ambiguous position

within the cytoplasmic membrane because the flanking TMS have a comparatively low content of

hydrophobic amino acids. Accordingly, only some topology analysis algorithms predict its location at

the outer surface of the cytoplasmic membrane and a previous experimental topology investigation

using a set of translational fusions with enzymes that are active only at intracellular or extracellular

location has revealed a preferential location at the inner cytoplasmic membrane surface in *E. coli*

[15]. Since our M-C7.1 and M-C7.3 binding experiments indicated accessibility of loop seven from the

outside, we revisited its location in *S. aureus* with two experimental strategies.

To confirm that M-C7.1 has access to its cognate MprF antigen epitope at the outer surface of the

cytoplasmic membrane in intact bacterial cells, *S. aureus* SA113Δ*spa*Δ*mprF* expressing MprF (pRB-

MprF) was grown in the presence of M-C7.1. Cells were then disrupted, membranes were solubilized

by treatment with the mild non-ionic detergent n-dodecyl-ß-D-maltoside (DDM), and proteins were

separated in non-denaturing PAGE gels (blue native PAGE). If M-C7.1 bound before cell disruption

and remained tightly attached to MprF, it should shift the MprF bands in Western blots of the blue

native gels and be detectable after Western blotting of the blue native gels with a human IgG-specific

secondary antibody. In order to detect MprF independently of M-C7.1 an MprF variant translationally

fused to green-fluorescent protein (MprF-GFP, expressed from plasmid pRB-MprF-GFP), which is

detectable by a GFP-specific primary antibody [15], was also used and treated in the same way. To

detect potential non-shifted MprF proteins in the native MprF variant (pRB-MprF), M-C7.1 was used

as additional primary antibody. *S. aureus* SA113*ΔspaΔmprF* bearing the empty vector (pRB) was used

as a negative control. In addition to M-C7.1, all three *S. aureus* strains were also incubated with the

control antibody L-1. *S. aureus* cells expressing either unmodified MprF or MprF-GFP yielded a

protein band migrating at a molecular weight of around 900 kDa when preincubated with M-C7.1 but

not with L-1 or in cells with the empty vector control (pRB) (Fig. 2A; Fig. S4). In contrast, the empty-

vector control (pRB) strain showed an unspecific band at 300 kDa when preincubated with M-C7.1 or

at 150 kDa when preincubated with L-1 but no MprF-specific band (Fig. 2A; Fig. S4). The 900-kDa

band of MprF-GFP was detected by both M-C7.1 and anti-GFP confirming the identity of MprF (Fig.

2A; Fig. S4). We could recently show that MprF forms oligomers in the staphylococcal membrane,

which were migrating at ca. 300 and 600 kDa [15]. Bands migrating at similar heights (ca. 250 and

500 kDa) were detected specifically in the MprF-GFP lanes after preincubation with either M-C7.1 or

L-1 and only by anti-GFP primary antibodies but not by M-C7.1 as primary antibody (Fig. 2A; Fig. S4)

suggesting that they represent the oligomerized but not the mAB-complexed MprF proteins [15].

Therefore, the 900-kDa bands probably represent a complex formed by MprF and M-C7.1, which

confirms that M-C7.1 specifically binds MprF loop seven in live *S. aureus* cells. The fact, that M-C7.1

used as primary antibody was not able to detect 250 and 500-kDa bands of non-complexed MprF

proteins neither after preincubation with M-C7.1 nor L-1 suggests that either antibody detection is

prevented by the Western blot procedure or that M-C7.1 binding only occurs in the living

staphylococcal cell potentially due to MprF conformational changes during flippase activity. Such

conformational changes during Lys-PG translocation in live *S. aureus* would render MprF loop seven

accessible to M-C7.1. Of note, the MprF-mAB complex migrating at around 900 kDa indicates that

higher-order MprF multimers were shifted by complex formation with M-C7.1.

The position of MprF loop seven was further investigated by inserting a cysteine residue into the loop

and analyzing the capacity of the membrane-impermeable agent Na*-(3-maleimidylpropionyl)-

biocytin (MPB) to label cysteines covalently, using the Substituted Cysteine Accessibility Method

(SCAM) [20]. MPB treatment of intact *S. aureus* cells should only lead to labelling of extracellular

protein portions while blocking cysteines at the outside with 4-acetamido-4'-maleimidylstilbene-2,2'-

disulfonic acid (AMS), followed by cell homogenization and addition of MPB should allow to label

only protein parts at the inner surface of the membrane according to a previously established

method in *E. coli* [20]. An MprF variant lacking all native cysteine residues was generated to exclude

background signals. Cysteine residues in MprF were exchanged against serine or alanine residues to

minimize structural or functional changes. Native *S. aureus* MprF contains six cysteines none of which

is conserved in MprF proteins from other bacteria suggesting that they do not have critical functions.

The cysteine-deficient mutant protein was found to be indeed functional because it decreased the

susceptibility of the *S. aureus mprF* mutant (SA113*ΔmprF*) to daptomycin (Fig. S3), which depends on

intact MprF synthase and flippase activities [14]. Cysteines were then inserted into MprF loop seven

(T263) and, as a control, into the first intracellular loop (loop two, A99) and the last extracellular loop

(loop thirteen, T480), the localization of which had been consistently confirmed by previous

computational and experimental analyses [15] (Fig. 1A & 2C). Amino acids for exchange were chosen

according to a predicted weak effect for functional changes by respective substitution with cysteine

using https://predictprotein.org [21]. The prediction is based on a machine learning program

integrating both evolutionary information and structural features such as predicted secondary

structure and solvent accessibility to evaluate the effect of amino acid exchanges in a protein

sequence [21]. The SCAM approach further confirmed the topology of intracellular loop two and

extracellular loop thirteen (Fig. 2B), thereby demonstrating that the technique can lead to reliable

results in *S. aureus*. Notably, MprF loop seven was found in both locations, at the inner and outer

surface of the membrane. This finding corroborates the ambiguous position of loop seven and

suggests that this loop may have some degree of mobility in the membrane, which may reflect the

lipid translocation process. The finding also clarifies why M-C7.1 has the capacity to bind MprF loop

seven at the outer surface of the cytoplasmic membrane.

3.   MABs binding to putative extracellular loops of MprF render *S. aureus* susceptible to CAMPs

MprF confers resistance to cationic antimicrobials such as the bacteriocin nisin by reducing the

negative net charge of the membrane outer surface [11, 14]. If the mABs would not only bind to but

also inhibit the function of MprF, *S. aureus* should become more susceptible to nisin. The six mABs

with confirmed specific binding to MprF and the isotype control mAB L-1 were analyzed for their

capacity to increase the susceptibility of *S. aureus* SA113 to nisin. For those initial screening

experiments protein A (*spa*) mutants were used to diminish effects of unspecific IgG binding to

protein A. Bacteria were grown in the presence of one of the mABs and then incubated with nisin at

the IC_{50} followed by quantification of viable bacterial cells. Two of the MprF-specific mABs targeting

loops seven or thirteen increased the sensitivity of *S. aureus* SA113*Δspa* to nisin while the other

mABs and the isotype control mAB L-1 had no significant impact (Fig. 3A). MAB M-C7.1 caused the

strongest sensitization and was selected for further analysis using the highly prevalent and virulent

community-associated methicillin-resistant *S. aureus* (CA-MRSA) USA300 clone [22].

For all following experiments *S. aureus* wild type (WT) strains with intact protein A were used to

make sure that the observed sensitization to CAMPs was not affected by unspecific antibody binding

to protein A.

M-C7.1 was found to also increase the susceptibility of USA300 wild type to the human CAMP LL-37,

a host defense peptide produced by epithelial and phagocyte cells [23], and to daptomycin, a

lipopeptide antibiotic in clinical use sharing physicochemical and antibacterial properties with CAMPs [24], in addition to nisin (Fig. 3B-D). Of note, M-C7.1 also increased the antimicrobial activity of daptomycin against the daptomycin-resistant clinical MRSA isolate DAP-R 703 [25] (Fig. 3E), suggesting that M-C7.1 may potentially be able to overcome daptomycin resistance during therapy. M-C7.1 but not the isotype control mAB L-1 was able to inhibit growth of USA300 in the presence of subinhibitory concentrations of nisin (Fig. 3F). Thus, mABs specific for certain extracellular loops of MprF may not only bind to MprF but also inhibit its function.

4. M-C7.1 inhibits the flippase function of MprF

M-C7.1 binds MprF at the junction between the LysPG synthase and flippase domains (Fig. 1). The lipid patterns of SA113 wild type treated with M-C7.1 at concentrations that increased the susceptibility to nisin or with the isotype control mAB L-1 were compared but showed no differences indicating that the synthase function of MprF was not inhibited by M-C7.1 (Fig. 4A). The flippase activity of MprF promotes the exposure of positively charged LysPG at the outer surface of the cytoplasmic membrane thereby reducing the affinity for the small cationic protein cytochrome C [26], which binds preferentially to negatively charged PG, or for calcium-bound annexin V [27]. These model proteins have been shown to allow a sensitive assessment of changes in the surface charge of *S. aureus* in several previous studies [14, 17, 28]. Treatment of SA113 wild type with M-C7.1 at concentrations that increased the susceptibility to nisin led to a significant increase in the capacity to bind cytochrome C or annexin V compared to treatment with the isotype control mAB L-1 (Fig. 4B, C) thereby indicating that M-C7.1 inhibits the flippase function of MprF. It is tempting to speculate that the protein region of MprF loop seven and adjacent TMSs may accomplish a crucial function in the process of phospholipid translocation, as suggested by the dynamic localization of loop seven on the cytoplasmic and external faces of the membrane.

5. M-C7.1-treatment abrogates *S. aureus* survival in phagocytes

The capacity of PMNs to kill phagocytosed bacteria does not only rely on the oxidative burst but also on the activity of LL-37 and other CAMPs and antimicrobial proteins [29]. The increased susceptibility of M-C7.1-treated *S. aureus* to CAMPs should therefore alter its survival ability in PMNs. CA-MRSA wild-type strain USA300 was pretreated with mABs, opsonized with normal human serum, and exposed to human PMNs. Treatment with M-C7.1 or the isotype control mAB L-1 did not alter the rate of PMN phagocytosis, but M-C7.1-treated USA300 cells were significantly more rapidly killed by PMNs than those treated with the isotype control antibody (Fig. 5A, B). This finding reflects our previous reports on reduced survival of MprF-deficient *S. aureus* in PMNs [11, 30]. Thus, *S. aureus* treatment with M-C7.1 might reduce the capacity to persist in infections and may help to blunt the virulence of *S. aureus* in invasive infections.

Discussion

Monoclonal therapeutic antibodies have proven efficacy for neutralization of bacterial toxins such as *Clostridium botulinum* or *Clostridioides difficile* toxins, and mAB-based therapies targeting the *S. aureus* alpha toxins and leukotoxins are currently developed [4, 31]. Moreover, therapeutic mABs binding to *S. aureus* surface molecules to promote opsonic phagocytosis have been assessed in preclinical and clinical trials [32]. In contrast, mABs inhibiting crucial cellular mechanisms of bacterial pathogens have hardly been assessed so far [33]. We developed specific mABs, which can block the activity of *S. aureus* MprF, the first described bacterial phospholipid flippase. Our mABs did not mediate increased internalization of *S. aureus* cells, most probably because the bacterial cell wall is too thick to allow binding of the mAB FC part to phagocyte FC receptors. However, the inhibition of MprF by M-C7.1 sensitized *S. aureus* to CAMPs and daptomycin and promoted killing by human PMNs, which use CAMPs as an important component of their antimicrobial arsenal. Specific inhibition of MprF could therefore promote the capacity of human host defense to clear or prevent a

S. aureus infection and would, at the same time, increase the susceptibility of *S. aureus* to CAMP-like

antibiotics such as daptomycin. Thus, targeting bacterial defense mechanisms provides a promising

concept for anti-virulence therapy. M-C7.1 sensitized S. aureus also in the presence of protein A

indicating that the unspecific IgG-binding by protein A does not interfere with the M-C7.1 capacity to

block MprF.

Specific binding of MprF antibody M-C7.1 to the extracellular loop seven inhibited the flippase

function of MprF, which indicates that this loop plays a crucial role in the lipid translocation process.

Loop seven is located between predicted TMS 7 and 8, and its presence at the outer surface of the

cytoplasmic membrane had remained elusive since computational and experimental analyses had

yielded conflicting results [15]. Surprisingly, we found loop seven to be accessible from both, the

outside and inside of the cytoplasmic membrane. This suggests that the protein part formed by loop

seven and adjacent TMS 7 and 8 may change its position in the protein complex, moving between the

two membrane surfaces to accomplish LysPG translocation (Fig. 6), or it could be in a unique position

of the large MprF protein complex that allows access from both sides. This protein part might

represent the center of the MprF flippase establishing the previously reported MprF-dependent

distribution of charged LysPG in the inner and outer leaflet of the cytoplasmic membrane of *S. aureus*

[14] (Fig. 6). Interestingly, in living staphylococcal cells, M-C7.1 bound to putative higher-order MprF

multimers as shown via blue native Western blotting suggesting that such multimers might represent

the active lipid translocating state of MprF. Replacement of conserved amino acids in the M-C7.1-

targeted protein part has been shown to inhibit the flippase function of MprF [15], which

underscores the crucial function of this domain. Notably, one of the conserved and essential amino

acids in TMS 7, D254, is negatively charged. It is tempting to speculate that D254 may interact with

the positively charged head group of LysPG during the translocation process. Mutations on loop

seven and adjacent TMS have recently been found to cause specific resistance to the structurally

related lipopeptide antibiotics daptomycin and friulimicin B but not to other antimicrobials [17].

Indirect evidence has suggested that these mutations allow the flippase to translocate these

lipopeptide antibiotics instead of or together with LysPG [17], which is in agreement with a direct

role of loop seven in the translocation process. A mAB binding loop thirteen also sensitized *S. aureus*

to nisin in a similar but less pronounced way as M-C7.1 suggesting that this loop may also have a

critical role for MprF activity.

With the severe shortage of new antibiotic target and drug candidates, crucial cellular machineries

conferring fitness benefits during infection rather than accomplishing essential cellular functions

should be considered as targets for the development of new therapeutics. The phospholipid synthase

and flippase MprF may become a role model for further targeting of bacterial defense mechanisms

for such anti-fitness or anti-virulence drugs. Compared to other potential targets, MprF has the

advantage that essential parts of it are exposed at the external surface of the cytoplasmic

membrane. Moreover, MprF is found in many other bacterial pathogens for which similar

therapeutic mABs could be developed [14]. On the other hand, only a few amino acids in loop seven

are conserved in other bacteria, which would limit the impact of loop-seven-directed mABs on other

microbiome members thereby minimizing the resistance selection pressure and the risk of dysbiosis.

Materials and Methods

Bacterial strains, maintenance and mutagenesis of mprF

We used commonly used strains, the methicillin-susceptible laboratory strain *S. aureus* SA113 (ATCC

35556), methicillin-resistant clinical clone *S. aureus* USA300, and the SA113 *mprF* knock-out

derivative SA113*ΔmprF*, which has been described recently [26, 34] (Tab. S2).

For the construction of a protein A mutant (Δ*spa*) the *E. coli/S. aureus* shuttle vector pKOR1 [35] was

used, which allows allelic replacement with inducible counter-selection in staphylococci. Flanking

regions of *spa* were amplified from chromosomal DNA of *S. aureus* COL with primer pairs Spa-

del_attB1 (ggggacaagtttgtacaaaaaagcaggc**caatattccatggtccagaact; bold: spa sequence)** and Spa-del

for BglII (gtcg*agatct*ataaaaacaaacaatacacaacg, restriction site italic) as well as Spa-del_attB2

(ggggaccactttgtacaagaaagctggg**atcagcaagaaaacacacttcc; bold: spa flanking sequence)** and Spa-del

rev BglII (aaa*agatct*aacgaattatgtattgcaata, restriction site italic). Both PCR products were digested

with BglII and subsequently ligated. Without further purification, the ligation product was mixed with

equimolar amounts of pKOR1 and *in vitro* recombination was performed with BP clonase Mix

(Invitrogen) according to the manufacturer's instructions. The recombination mixture was

transferred to chemically competent *E. coli* DH5α and isolated plasmids from the resulting

transformants were analyzed by restriction digest. The correct plasmid was isolated from *E. coli*

DH5α and used to electroporate competent *S. aureus* RN4220 from which it was again isolated and

transferred into *S. aureus* SA113 by electroporation. Allelic replacement was essentially conducted as

previously described [35] and resulting deletion mutants were confirmed by PCR.

The *S. aureus* SA113 *spa mprF* double mutant (SA113*ΔspaΔmprF*) was constructed by transducing

the gene deletion cassette of the *S. aureus* SA113 *mprF* deletion mutant (SA113*ΔmprF*) to the

marker-less *spa*-deficient *S. aureus* SA113 mutant (SA113*Δspa*) using standard transduction

protocols. The resulting *S. aureus* strain SA113*ΔspaΔmprF* was identified by screening for

erythromycin resistance conferred by the gene deletion cassette and confirmed by PCR of the

deleted genome section. Bacteria were maintained on tryptic soy agar plates.

Cysteine substitutions in *mprF* were accomplished by site-directed mutagenesis in *Escherichia coli*

using *E. coli/S. aureus* shuttle vector pRB474 bearing *mprF,* using the QuickChange II Site-Directed

Mutagenesis Kit (Agilent), as described recently [17]. Mutated *mprF* derivatives in pRB474*mprF* were

transferred into SA113*ΔmprF* and *mprF* expression was mediated by the plasmid-encoded

constitutive *Bacillus subtilis* promoter *vegII*. 10 µg/ml chloramphenicol served for maintenance in all

plasmid-based studies. A99, T263, and T480 were chosen for exchange against a cysteine residue

because of prediction of a weak effect on protein structure in an analysis of functional changes given

a single point mutation according to https://predictprotein.org [21].

Plasmids and primers used in this study are given in Table S3 and S4, respectively. Unless otherwise

stated, bacteria were cultivated in Mueller-Hinton Broth (MHB, Sigma-Aldrich) with appropriate

antibiotics for all experiments.

Antigen selection and antibody production

MprF-derived peptides of interest (Fig. 1 and Tab. S1) were custom-synthesized by JPT Peptide

Technologies GmbH (Berlin). An N-terminal and a C-terminal cysteine were added to enable

cyclization. Biotin was coupled to the peptide via a 4,7,10-trioxa-1,13-tridecanediamine succinic acid

(Ttds)-linker [36]. Cyclization and coupling of linker and biotin were performed by JPT Peptide

Technologies.

Recombinant antibodies were generated from the HuCAL PLATINUM® library, as described recently,

by three iterative rounds of panning on the antigen peptides in solution and antigen-antibody-phage

complexes were captured with streptavidin-coated beads (Dynal M-280) [18, 37]. Fab-encoding

inserts of the selected HuCAL PLATINUM phagemids were cloned and expressed in *E. coli* TG1 F-cells,

purified chromatographically by IMAC (Bio-Rad) and subcloned in an IgG1 expression vector system

to obtain full-length IgG1s by expression in eukaryotic HKB11 cells, as described recently [18]. IgGs

were purified by protein A affinity chromatography (MabSelect SURE, GE Healthcare) [18].

The humanized antibody MOR03207 [38] directed against chicken lysozyme serves as isotype control

and was called L-1 in this manuscript.

Peptide and whole-cell ELISAs of MprF specific mABs

Overnight *S. aureus* SA113*Δspa* or SA113*ΔspaΔmprF* cultures grown in MHB were adjusted to OD_{600}

0.1 in fresh MHB and grown to OD_{600} of at least 1.0. Cells were harvested, adjusted to OD_{600} 0.5 with

0.9% NaCl, and used to coat 96 well NUNC Maxisorp Immuno Plates for 1 hour (50 µl/well). After

three washing steps with Tris-buffered saline (TBS; 50 mM Tris, 150 mM NaCl in 800 mL of H_2O, pH

7.4) containing 0.05% Tween 20 (TBS-T), the cells were blocked with PBS containing 1 x ROTI®Block

(Carl Roth) for 1 h. Primary antibodies were added after three washing steps with TBS-T at an

indicated final concentration of 1, 10 or 100 nM and incubated for 1 h. After three more washing

steps with TBS-T, anti-human IgG conjugated to alkaline phosphatase (Sigma-Aldrich A8542) diluted

1:10,000 in TBS-T was applied. Finally, after three washing steps with 2x TBS-T and 1x TBS, the cells

were incubated with 100 µl p-nitrophenyl phosphate solution as advised by the manufacturer

(Sigma-Aldrich N1891). Fluorescence was measured at an excitation of 440±25 nm and an emission

of 550±35 nm.

Detection of antibody binding to MprF by blue native Western blotting

To detect binding of the anti-MprF antibody to MprF we performed blue native polyacrylamide gel

electrophoresis (BN-PAGE) as previously described [15] with slight modifications. Briefly, 20 ml of *S.*

aureus cells grown in MHB with anti-MprF antibodies (100 µg/ml) to OD_{600} 1 were incubated with

750 µl of lysis buffer (100 mM EDTA [pH 8.0], 1 mM $MgCl_2$, 5 µg/ml lysostaphin, 10 µg/ml DNase,

proteinase inhibitor [cocktail set III from Calbiochem] diluted 1:100 in PBS) for 30 min at 37°C and

homogenized three times with 500 µl zirconia-silica beads (0.1 mm diameter from Carl-Roth) using a

FastPrep 24 homogenizer (MP Biomedicals) for 30 s at a speed of 6.5 m/s. After removing the beads,

cell lysate was centrifuged 20 min at 14,000 rpm and 4°C to sediment cell debris and supernatant

was transferred to microcentrifuge polypropylene tubes (Beckman Coulter) and cytoplasmic

membranes were precipitated by ultracentrifugation for 45 min at 55,000 rpm and 4°C (Beckman

Coulter rotor TLA 55). Membrane fractions resuspended in resuspension buffer (750 mM

aminocaproic acid, 50 mM Bis-Tris [pH 7.0]) were incubated with the MprF- or lysozyme-specific

humanized IgG mAB and/or a GFP-specific rabbit IgG mAB (Invitrogen) for 30 min shaking at 37°C.

Dodecyl maltoside was added to a final concentration of 1% for 1 h at 4°C to solubilize MprF-

antibody complexes. Insoluble material was removed by ultracentrifugation for 30 min at 40,000 rpm

and 4°C. 20 µl supernatant was mixed 1:10 with 10 × BN loading dye (5% [wt/vol] Serva Blue G, 250

mM aminocaproic acid, 50% glycerol) and run on a Novex NativePAGE 4 to 16% Bis-Tris gel

(Invitrogen). Separated proteins were transferred to a polyvinyl difluoride membrane and detected

via either, goat anti-human IgG DyLight 700 (Pierce) or goat anti-rabbit IgG DyLight 800 (Pierce) as

secondary antibodies in an Odyssey imaging system from LI-COR.

Substituted cysteine accessibility method (SCAM) to localize MprF loops in inner and/or outer membrane leaflet

Substituted cysteine accessibility method (SCAM) to localize MprF loops in inner and/or outer

membrane leaflet

SCAM was adapted for use in *S. aureus* from the protocol recently established for *E. coli* [20].

Variants of *S. aureus* SA113*ΔmprF* were expressing cysteine-deficient and/or -altered MprF

derivatives bearing a FLAG® tag at the C-terminus via the plasmid pRB474. Bacteria were grown at

37°C to an OD_{600} 0.7-0.8 in 100 ml TSB, split into two equal aliquots, and harvested. Pellets were

resuspended in a mixture of buffer A (100 mM HEPES, 250 mM sucrose, 25 mM $MgCl_2$, 0.1 mM KCl

[pH 7.5]) supplemented with 1 mM $MgSO_4$, 1 mM EDTA, 2.1 µg lysostaphin, 0.7 µg DNase and 1% of a

protease inhibitor cocktail (Roche) and incubated for 45 min at 37°C. For staining of cysteine residues

in the outer leaflet of the cytoplasmic membrane, cells of the first aliquot were treated with N*-(3-

maleimidylpropionyl) biocytin (MPB, Thermo Scientific) at a final concentration of 107 µM for 20 min

on ice. In the second aliquot, external cysteine residues were blocked with 107 µM 4-acetamino-4´-

maleimidylstilbene-2,2´-disulfonic acid (AMS, Thermo Scientific) for 20 min on ice, bacterial cells

were subsequently disrupted in a bead mill as described above and internal cysteines were labeled

with MPB for 5 min. MPB labeling was quenched in both aliquots by the addition of 21 mM β-

mercaptoethanol. Cell debris was removed by several centrifugation and washing steps, membrane

fractions were collected via ultracentrifugation at 38,000 x g, and labeled membrane samples were

stored at -80°C.

For protein enrichment and precipitation, labeled membrane samples were thawed on ice and

suspended in 20 mM β-mercaptoethanol containing buffer A. Proteins were solubilized by the

addition of an equal volume of solubilization buffer (50 mM Tris-HCl [pH 9], 1 mM EDTA, 2% SDS) and

vigorous vortexing for 30 min at 4°C, followed by an incubation step of 30 min at 37°C and vortexing

for 30 min at 4°C. After solubilization, one and a half volumes of immunoprecipitation buffer 1 (50

mM Tris-HCl [pH 9], 150 mM NaCl, 1 mM EDTA, 2% Thesit, 0.4 % SDS [pH 8.1]) was added and

samples were incubated for 2.5 h with magnetic FLAG®beads (Sigma-Aldrich) in a rotation wheel at

4°C. After several washing steps with immunoprecipitation buffer 1 and immunoprecipitation buffer

2 (50 mM Tris-HCl [pH 9], 1 M NaCl, 1 mM EDTA, 2% Thesit, 0.4 % SDS [pH 8.1]), FLAG®-tagged MprF

was eluted by the addition of 0.1 M glycine-HCl (pH 2.2) and neutralized by adding Tris-HCl (pH 9).

For detection of cysteine-labelled MprF, samples were analyzed by denaturing SDS-PAGE and

Western blotting. To this end, 12-μl samples were mixed with 4 x Laemmli sample buffer (Bio-Rad),

loaded onto a 10%-polyacrylamide gel, and separated via electrophoresis. Proteins were transferred

onto a polyvinylidene difluoride (PVDF) membrane (Immobilon-PSQ PVDF membrane, Merck) by

semi-dry turbo Western blot procedure (Trans-Blot Turbo Transfer System, Bio-Rad). FLAG®-tagged

MprF was detected both by a mouse anti-FLAG® primary antibody (Sigma-Aldrich) and goat anti-

mouse secondary antibody (LI-COR) at 700 nm, while MBP-labelled cysteine residues were detected

with streptavidin DyLight conjugate (Thermo Scientific) at 800 nm in an Odyssey imaging system from

LI-COR.

Determination of susceptibility to antimicrobial agents

Overnight cultures of *S. aureus* SA113*Δspa* or USA300 WT were diluted in fresh MHB and adjusted to

OD_{600} 0.25 (~1.5 x 10^7 cells). Antibodies were adjusted to a concentration of 1 mg/ml and 10 μl per

well of a 96 well plate were added to 90 μl of the adjusted cell suspension (final antibody

concentration: 100 μg/ml). Cells were grown in the presence of the isotype control antibody L-1 or

with the respective anti-MprF antibodies. After 3 hours of incubation at 37°C under shaking, optical

density was determined, and cells were adjusted to OD$_{600}$ 0.025 in 500 µl cold PBS. 80 µl of the

adjusted cell suspension were mixed with 20 µl of antimicrobial substances to final concentrations of

22.5 µg/ml nisin for SA113Δspa or of 5 µg/ml nisin for USA300, 1.5 µg/ml daptomycin for USA300, 11

µg/ml daptomycin for daptomycin resistant CA-MRSA strain 703, and 45 µg/ml LL-37 for USA300 or

20 µl PBS as control. After incubation for 2 hours under shaking at 37°C, the cell suspensions were

diluted 1:2,000 and 100 µl of each duplicate was plated in triplicates on TSB agar plates to obtain a

representative value of bacterial survival.

MICs of daptomycin were determined using MIC test strips (Liofilchem) according to the

manufacturer's advice.

To analyze the capacity of mAB M-C7.1 to inhibit *S. aureus* growth, overnight cultures of CA-MRSA

strain USA300 grown in MHB medium were diluted to OD$_{600}$ 0.01 with fresh MHB medium containing

1 µM of antibodies and 4 µg/ml nisin (Sigma-Aldrich) or PBS as a control in 96 well plates. Plates were

incubated for 24 hours at 37°C under shaking conditions using a Bioscreen C (Oy Growth Curves Ab

Ltd). Optical density at 600 nm was determined and compared to growth without nisin.

Isolation and quantification of polar lipids

S. aureus cultures (1 ml in MHB) were grown to the exponential phase (OD$_{600}$ 1) in the presence of

antibodies (final concentration: 100 µg/ml) as described above for the determination of susceptibility

to antimicrobials. Polar lipids were extracted using the Bligh-Dyer method [39], with

chloroform/methanol/sodium acetate buffer (20 mM) (1:1:1, by vol.), vacuum-dried, and

resuspended in chloroform/methanol (2:1, by vol.). Extracts were filled into a 100-µl Hamilton

syringe and spotted onto silica gel 60 F254 high-performance thin-layer chromatography (HPTLC)

plates (Merck) with a Linomat 5 sample application unit (Camag) and run in a developing chamber

ADC 2 (Camag) with a running solution composed of chloroform-methanol-water (65:25:4, by vol.).

Phosphate group-containing lipids were detected by molybdenum blue staining and phospholipids

were quantified in relation to total phospholipid content by determining lipid spot intensities

densitometrically with ImageJ (http://rsbweb.nih.gov/ij/docs/guide/index.html) as described

recently [28].

Determination of S. aureus cell surface charge

For analysis of the cytochrome C repulsion capacity of _S. aureus_, bacterial cells were grown in the

presence of antibodies (100 µg/ml) as described above for the analysis of antimicrobial susceptibility.

Bacteria were adjusted to OD_{600} 1 in 1 ml PBS, pelleted, resuspended in 100 µl 0.05 mg/ml

cytochrome C (Sigma-Aldrich) followed by incubation at $37^{o}C$ under shaking. The cells were then

pelleted by centrifugation and absorption of the supernatant containing unbound cytochrome C was

determined at OD_{410}

To determine annexin V binding to _S. aureus,_ bacterial cells were grown in the presence of antibodies

(100 µg/ml) as described above at 37°C with shaking for 3 hours, harvested, washed twice in PBS

buffer, and resuspended in PBS containing $CaCl_2$ to OD_{600} 0.5 in 1 ml. Bacteria were gently mixed with

5 µl of allophycocyanin (APC)-labeled annexin V (Thermo Scientific) and incubated at room

temperature for 15 min in the dark, as described recently [17]. At least 50,000 bacterial cells per

antibody were analyzed by flow cytometry to quantify surface-bound fluorophore (FL-4).

Phagocytosis and killing of S. aureus by human polymorphonuclear leukocytes (PMNs)

Human PMNs are the major human phagocytes to counteract bacterial infection and form the largest

subgroup (~95%) of PMNs [29]. PMNs were isolated from fresh human blood of healthy volunteers

by standard Ficoll/Histopaque gradient centrifugation as described recently [40]. Cells were

resuspended in HBSS containing 0.05% human serum albumin (HBSS-HSA; HBSS with 0.05%

Albiomin®, Biotest AG).

CA-MRSA strain USA300 was prepared for PMN experiments as described for the determination of

susceptibility to antimicrobials, grown in the presence of antibodies (100 µg/ml) at 37°C under

shaking for 3 hours, harvested, washed twice in HBSS, and adjusted to a density of 5×10^7 CFU/ml.

Pooled serum from healthy human volunteers (blood bank of the University Hospital Tübingen) was

added to a final concentration of 5% and bacteria were opsonized for 10 min at 37°C, as described

recently [11]. Prewarmed bacteria and PMNs were mixed to final concentrations of 5×10^6 CFU/ml

and 2.5×10^6 PMNs/ml in flat-bottom 96 well plates together with antibodies (final concentration

100 µg/ml). Samples of 50 µl were shaken at 37°C.

For analysis of the PMN killing capacity, incubation was stopped at different time points by the

addition of 100-fold volumes of ice-cold distilled water to disrupt the PMNs. Triplets of appropriate

sample volumes were spread on LB agar plates and colonies were counted after 24 h incubation at

37°C. Numbers of live bacteria did not change during the 60-min incubation period in the absence of

PMNs compared to the initial bacteria counts.

For phagocytosis studies, overnight cultures of bacteria were labeled with 0.1 mg/ml fluorescein-5-

isothiocyanate (FITC) at 37°C for 1 h, as described previously [11]. After washing with PBS, the

bacteria were resuspended in HBSS-HSA, adjusted to 5×10^6 CFU/ml, mixed with PMNs (2.5×10^6

PMNs/ml), and opsonized as described above (shaking at 37°C). Incubation was stopped by addition

of 100 µl ice-cold 1% paraformaldehyde. The percentage of PMNs bearing FITC-labeled bacteria was

determined by flow cytometric analysis of 20,000 cells.

Statistics

Statistical analyses were performed with the Prism 5.0 package (GraphPad Software) and Group

differences were analyzed for significance with the two-tailed Student's *t* test. A *P* value of ≤0.05 was

considered statistically significant.

Author contributions

C.J.S, A.K., S.W., M.T., B.K., C.M.E., and A.P. conceived and designed the experiments. C.J.S., J.N.H.,

C.G., J.S., A.G., D.H., B.K., and C.M.E. performed the experiments. C.J.S., J.N.H., J.S., A.G., A.K., S.W.,

S.K., C.M.E., and A.P. analyzed the data. S.W., M.T., and A.P. contributed material / analysis tools.

C.J.S., J.N.H., S.W., C.M.E., and A.P. wrote the paper.

Acknowledgements

This work was financed by Grants from the German Research Foundation TRR34, SFB766, TRR156,

TRR261, and GRK1708 to A.P.; and the German Center of Infection Research (DZIF) to A.P. C.J.S. was

supported by the intramural Experimental Medicine program of the Medical Faculty at the University

of Tübingen and received grants from the DZIF and the cluster of Excellence EXC2124 Controlling

Microbes to Fight Infection (CMFI). The authors acknowledge infrastructural support by the cluster of

Excellence EXC2124 CMFI.

Figures

Figure 1. MprF topology and binding of mABs to targeted MprF-expressing *S. aureus* cells. (A) MprF

membrane topology is given according to our recent study [15] showing synthase and flippase

domains in gray and black, respectively. Amino acid (aa) positions of TMS and the C-terminal

hydrophilic domain are indicated. TMSs from aa 245-265 and aa 278-298 are shown in two

alternative positions as computational and experimental results of transmembrane topology have

been contradictory [15]. Localizations of MprF's TMS-connecting loops are numbered starting from

the N-terminus, antibody-targeted loops are indicated by red circles and antibody symbols. (B)

Specific binding of mABs (100 nM) to *S. aureus* was analyzed by ELISA using SA113 strains deficient in

the IgG-binding protein A (Spa) comparing the SA113 *spa* mutant (*Δspa*) and *spa mprF* double

knockout mutant (*ΔspaΔmprF*). The red line indicates the mean fluorescence measured at A 440 nm

(affinity) of the isotype control mAB L-1 bound to *S. aureus* SA113*Δspa.* Means and SEM of at least

 three experiments are shown. Significant differences between SA113*Δspa* and SA113*ΔspaΔmprF*

were calculated by Student's paired t-test (ns, not significant; *, *P* < 0.05; ***, *P* < 0.0001).

Figure 2. Binding of M-C7.1 to MprF and membrane localization of the M-C7.1-targeted MprF loop

seven. (A) Detection of M-C7.1 binding to MprF. Plasmid-encoded native and GFP-tagged MprF

variants were expressed in *S. aureus* SA113*ΔspaΔmprF* and living cells were preincubated with M-

C7.1 or the isotype control mAB L-1 (in order to form MprF-mAB complexes). MprF

variants/complexes were detected by blue native Western blotting via primary (anti-GFP or M-C7.1)

and secondary antibodies (red for secondary antibody detection of M-C7.1; green for secondary

antibody detection of anti-GFP). SA113*ΔspaΔmprF* expressing the empty vector (pRB) served as

negative control. Molecular masses in kDa of marker proteins are given on the left of the gel. (B)

Cellular localization of the antigen epitope of M-C7.1 using the substituted cysteine accessibility

method (SCAM) for specific loops between the MprF TMSs. The substituted cysteine T263C is

localized in M-C7.1´s target peptide sequence in MprF. Substitution of A99C served as inside control,

substitution of T480C served as outside control (see topology model, part C). *S. aureus* SA113*ΔmprF*

expressing the empty vector (pRB) and an MprF variant lacking all native cysteines (WT (-Cys)) served

as additional negative controls. All MprF variants were plasmid-encoded, FLAG®-tagged at the C-

terminus to allow immunoprecipitation and detection, and were expressed in *S. aureus* SA113*ΔmprF*.

Substituted extracellular cysteine residues were labeled with Na*-(3-maleimidylpropionyl)-biocytin

(MPB) (outer leaflet signal, green in overlay), while labeling of substituted internal cysteine with MPB

was performed after the blocking of external cysteines with 4-acetamino-4'-maleimidylstilbene-2,2'-

disulfonic acid (AMS) (inner leaflet signal, green in overlay). MprF was detected via antibody staining

by an anti-FLAG® antibody (red in overlay). (C) MprF topology showing location and amino acid

exchanges of artificial cysteine residues for SCAM detection in green.

Figure 3. Killing and growth inhibition of *S. aureus* by antimicrobial peptides and antibiotics in the

presence of M-C7.1. (A) Survival of *S. aureus* SA113Δ*spa* in the presence of nisin and anti-MprF

antibodies compared to control mAB L-1. Surviving CFU of each antibody (and nisin) containing

sample were analyzed after 3 h incubation and the negative control (isotype mAB L-1) was set to

100% survival. (B) Survival of CA-MRSA WT strain USA300 in the presence of nisin and M-C7.1

compared to the isotype control mAB L-1. (C) Survival of USA300 WT in the presence of LL-37 and M-

C7.1 compared to the isotype control mAB L-1. (D) Survival of USA300 WT in the presence of

daptomycin and M-C7.1 compared to the isotype control mAB L-1. (E) Survival of daptomycin-

resistant CA-MRSA strain 703 [25] in the presence of daptomycin and M-C7.1 compared to the

isotype control mAB L-1. (F) Growth inhibition of USA300 WT in the presence of subinhibitory nisin

concentrations and M-C7.1 compared to the isotype control mAB L-1. Wells without nisin and/or

antibodies served as additional negative controls. The means plus SEM of results from at least three

independent experiments are shown in A-E. Technical duplicates were employed for each analysis. F

shows means plus SEM of technical triplicates from a representative experiment of three

independent experiments. Values for M-C7.1 or other anti-MprF antibodies that were significantly

different from those for the isotype control mAB L-1, calculated by Student's paired t-test are

indicated (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.0001$).

Figure 4. M-C7.1 inhibits the MprF LysPG flippase but not the LysPG synthase. (A) Detection of phospholipids from *S. aureus* SA113*ΔmprF* and WT treated or not treated with M-C7.1 or the isotype control mAB L-1. Polar lipids were separated by TLC and stained with the phosphate group-specific dye molybdenum blue to detect the well-documented PG and LysPG pattern of *S. aureus* WT and *mprF* deletion mutant [28]. Percentages of LysPG in relation to total phospholipid content are given below respective LysPG spots. (B) The repulsion of positively charged cytochrome C corresponds to MprF LysPG synthase plus flippase activity while the synthase activity alone does not affect repulsion. To assess MprF flippase efficiency, unbound cytochrome C in the supernatant was quantified photometrically after incubation with the *S. aureus* SA113 WT pretreated with M-C7.1 or the isotype control mAB L-1 (the latter set to 100%). SA113*ΔmprF* treated with M-C7.1 served as negative

control. The means plus SEM of results from three independent experiments are shown. (C) Annexin

V binding to *S. aureus* SA113 WT after incubation with M-C7.1 or the isotype control mAB L-1 was

quantified by measuring cell-bound annexin V by FACS and L-1 treated samples were set to 100%.

Data are expressed in relative fluorescence units. The means plus SEM of results from three

independent experiments are shown in B and C. Significant differences between M-C7.1 and L-1-

treated samples were calculated by Student's paired t-test (*, P < 0.05; ***, P < 0.0001).

Figure 5. M-C7.1 supports *S. aureus* clearance by isolated human PMNs. (A) Kinetics of killing of CA-

MRSA strain USA300 WT treated with M-C7.1 compared to isotype control mAB L-1 by freshly

isolated human PMNs. Viable bacteria (CFU) after incubation with PMNs are shown as percentage of

initial CFU counts. The means plus SEM of results from three independent experiments are shown.

(B) Kinetics of phagocytosis of USA300 WT treated with M-C7.1 compared to isotype control mAB L-1

by freshly isolated human PMNs. Percentages of PMNs bearing FITC-labeled USA300 are given.

Means of three counts from a representative experiment are shown.

Figure 6. Proposed model for MprF inhibition by M-C7.1. Flipping of positively charged LysPG probably by TMS 7 and 8 of MprF results in a more positively charged staphylococcal membrane able to repulse AMPs or daptomycin (DAP). Binding of M-C7.1 to MprF loop seven blocks the flippase, which results in a more negatively charged staphylococcal membrane and subsequently in an increased *S. aureus* membrane disruption by AMPs and daptomycin.

644 **Supplemental figures**

Workflow

Supplemental Figure S1. Workflow for the development of MprF specific mABs. Biotinylated MprF

peptide loops were incubated with the HuCal phage display library expressing single-chain human

Fab fragments [18], antigen-binding phages were enriched in three iterative panning rounds, bound

antigen-specific phages were isolated and respective Fab fragments were subcloned in *E. coli*

expression vectors to yield His-tagged Fab fragments, 24 unique Fabs against all peptides were

converted to human IgG by cloning in an IgG1 expression vector system for human HKB11 cells, and

IgGs were purified via protein A chromatography.

Supplemental Figure S2. Specific binding of selected mABs to cyclic MprF-derived target peptides

was analyzed by ELISA. Biotinylated cyclic peptides corresponding to the MprF loops 1, 7, 9, and 13

were incubated with eight anti-MprF IgGs in PBS compared to the control mAB L-1. (A-D) show

binding of mABs at increasing concentrations to cyclic MprF loops 1, 7, 9, and 13, respectively. Mean

fluorescence measured at A 440 nm and SEM of three independent experiments are shown.

Supplemental Figure S3. Effects of cysteine replacement and insertion on MprF function, assessed by

measuring daptomycin susceptibility. Minimal inhibitory concentrations (MICs) of daptomycin

667 against the indicated *S. aureus* strains are shown. The *mprF* deletion mutant with empty pRB474

668 plasmid served as a negative control. The means plus SEM of results from three independent

669 experiments are shown. Values that are significantly different from the values determined for *S.*

670 *aureus* SA113Δ*mprF* bearing pRB474 (pRB), calculated by Student´s paired t-test, are indicated (**,

671 *P* < 0.01; ***, *P* < 0.0001).

675 **Supplemental Figure S4.** Detection of M-C7.1 binding to MprF. Single channels are showed in black

676 and white from figure 3A. Further explanations are found in figure legend of figure 3A.

679 **Supplemental tables**

681 **Table S1**. MprF-directed antibodies and its target peptides.

Antibody	Antigen	Peptide sequence	Framework
M-C1	Cyclic peptide, loop TMS 1-2	ELSGINFKDTLVEFSKINR	VH3-23 kappa 3
M-C7.1	Cyclic peptide, loop TMS 7-8	LGFKTLGVPEEKV	VH1A kappa 1
M-C7.2	Cyclic peptide, loop TMS 7-8	LGFKTLGVPEEKV	VH3-23 kappa 1
M-C7.3	Cyclic peptide, loop TMS 7-8	LGFKTLGVPEEKV	VH1A kappa 1
M-C9.1	Cyclic peptide, loop TMS 9-10	DALYDGNHLT	VH1A kappa 1
M-C9.2	Cyclic peptide, loop TMS 9-10	DALYDGNHLT	VH3-23 kappa 1
M-C13.1	Cyclic peptide, loop TMS 13-14	DIYTIEMHTSVLR	VH1A kappa 1
M-C13.2	Cyclic peptide, loop TMS 13-14	DIYTIEMHTSVLR	VH1A kappa 1

683   **Table S2.** Bacterial strains used in this study.

Strain name	Characteristics
S. aureus SA113 WT (ATCC 35556)	Restriction-deficient *S. aureus* strain derived from NCTC 8325 [41]
S. aureus SA113Δ*mprF*	MprF deletion mutant of SA113, Erm[r] [11]
S. aureus SA113Δ*spa*	Spa deletion mutant of SA113. Constructed in this study. Markerless.
S. aureus SA113Δ*spa*Δ*mprF*	Spa and MprF double deletion mutant of SA113. Constructed in this study. Erm[r]
S. aureus DAP-R MRSA 703	Daptomycin resistant clinical MRSA isolate possessing a single point mutation in *mprF* (S295L) [25]
S. aureus USA300 LAC	CA-MRSA wild type strain [34]
E. coli TG1	Strain for phage display usage [18]

685   **Table S3.** Plasmids used in this study.

Plasmid	Characteristics	Short name in figures
pKOR1	*E. coli/S. aureus* shuttle vector to allow allelic replacement with inducible counter-selection in staphylococci [35]	-
pRB474	*E. coli/S. aureus* shuttle vector pRB474 [42]	pRB
pRB474mprF	*mprF* cloned in *E. coli/S. aureus* shuttle vector pRB474 [11]	pRB-MprF

pRB474mprF-GFP	N-terminally GFP tagged *mprF* cloned in *E. coli/S. aureus* shuttle vector pRB474 [15]	pRB-MprF-GFP
pRB474mprFdelCys flag	C-terminally FLAG tagged, cysteine depleted *mprF* cloned in *E. coli/S. aureus* shuttle vector pRB474; constructed in this study	WT (-Cys)
pRB474mprFdelCys T263C flag	C-terminally FLAG tagged, cysteine depleted *mprF* with artificial cysteine insertion cloned in *E. coli/S. aureus* shuttle vector pRB474; as indicated, each plasmid bears another amino acid substituted against cysteine (e.g. T263C)	pRB-T263C
pRB474mprFdelCys A99C flag		pRB-A99C
pRB474mprFdelCys T480C flag		pRB-T480C

688 **Table S4.** Primers used in this study.

Primer	5' → 3' sequence	Usage
A99C fw	GCATTGAATTGTATTGTAGGTTTCGGTGGCTTTATTGGTGCAGGCG	Forward primer for construction of pRB-A99C by site-directed mutagenesis
A99C rev	CCGAAACCTACAATACAATTCAATGCATTGATGATATAACTTACTC	Reverse primer for construction of pRB-A99C by site-directed mutagenesis
T263C fw	GTTGTATTACTAGGATTTAAATGTTTAGGTGTCCCTGAGGAAAAAG	Forward primer for construction of pRB-T263C by site-directed mutagenesis
T263C rev	CTTTTTCCTCAGGGACACCTAAACATTTAAATCCTAGTAATACAAC	Reverse primer for construction of pRB-T263C by site-directed mutagenesis
T480C fw	GGAACGTTATATGCATTAGATATTTATTGTATTGAAATGCATACATCTGTATTGCG	Forward primer for construction of pRB-T480C by site-directed mutagenesis
T480C rev	CGCAATACAGATGTATGCATTTCAATACAATAAATATCTAATGCATATAACGTTCC	Reverse primer for construction of pRB-T480C by site-directed mutagenesis
mprF C199+204S fw	TACTCTACTTTAGTGTCGTCTGTTGAATGGTTAGCAG	
mprF C199+204S rev	AACAGACGACACTAAAGTAGAGTACAATCCTACAAAACG	
mprF C217A fw	TTCGCTGGTGTAATTGTTGACGC	Primer for cysteine depletion of pRB474 encoded native *mprF* by site-directed mutagenesis
mprF C217A rev	ACCAGCGAAATATAATACAACTGC	
mprF C380A fw	GCTGCTTTATTACTTTTACTGAATGTAGTTGG	
mprF C380A rev	TAAAGCAGCACTAGTATGAATTGCC	

mprF C526S fw	GATAGCGAGGAGATTATTAATCAG	
mprF C526S rev	CTCGCTATCTTCAATTTTAGAAG	
mprF C717S fw	GTAATTGCATTTAGTAGTTTAATGCCAACATACTTTAATGATG	
mprF C717S rev	CTACTAAATGCAATTACTTCATTTTCTTCATTTCGCATTACACC	
Spa-del_attB1	GGGGACAAGTTTGTACAAAAAAGCAGGCCAATATTCCATGGTCCAGAACT	Construction of a markerless *spa* knockout mutant using the pKOR1 vector system [35]
Spa-del	GTCGAGATCTATAAAAACAAACAATACACAACG	
Spa-del_attB2	GGGGACCACTTTGTACAAGAAAGCTGGGATCAGCAAGAAAACACACTTCC	
Spa-del rev	AAAAGATCTAACGAATTATGTATTGCAATA	

References

1.   Ardal, C., et al., *Antibiotic development - economic, regulatory and societal challenges.* Nat Rev Microbiol, 2020. **18**(5): p. 267-274.

2.   Tacconelli, E., et al., *Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis.* Lancet Infect Dis, 2017.

3.   Lee, A.S., et al., *Methicillin-resistant Staphylococcus aureus.* Nat Rev Dis Primers, 2018. **4**: p. 18033.

4.   Dickey, S.W., G.Y.C. Cheung, and M. Otto, *Different drugs for bad bugs: antivirulence strategies in the age of antibiotic resistance.* Nat Rev Drug Discov, 2017. **16**(7): p. 457-471.

5.   Lakemeyer, M., et al., *Thinking Outside the Box-Novel Antibacterials To Tackle the Resistance Crisis.* Angew Chem Int Ed Engl, 2018. **57**(44): p. 14440-14475.

6.   Theuretzbacher, U., et al., *The global preclinical antibacterial pipeline.* Nat Rev Microbiol, 2019.

7.   Denis, K., et al., *Targeting Type IV pili as an antivirulence strategy against invasive meningococcal disease.* Nat Microbiol, 2019. **4**(6): p. 972-984.

8.   Qu, X., Y. Tang, and S. Hua, *Immunological Approaches Towards Cancer and Inflammation: A Cross Talk.* Front Immunol, 2018. **9**: p. 563.

9.   Chen, Z., M. Moayeri, and R. Purcell, *Monoclonal antibody therapies against anthrax.* Toxins (Basel), 2011. **3**(8): p. 1004-19.

10.   Posteraro, B., et al., *Actoxumab + bezlotoxumab combination: what promise for Clostridium difficile treatment?* Expert Opin Biol Ther, 2018. **18**(4): p. 469-476.

11.   Peschel, A., et al., *Staphylococcus aureus resistance to human defensins and evasion of neutrophil killing via the novel virulence factor MprF is based on modification of membrane lipids with l-lysine.* J Exp Med, 2001. **193**(9): p. 1067-76.

12.   Ernst, C.M. and A. Peschel, *Broad-spectrum antimicrobial peptide resistance by MprF-mediated aminoacylation and flipping of phospholipids.* Mol Microbiol, 2011. **80**(2): p. 290-9.

13.   Slavetinsky, C., S. Kuhn, and A. Peschel, *Bacterial aminoacyl phospholipids - Biosynthesis and role in basic cellular processes and pathogenicity.* Biochim Biophys Acta, 2016.

14.   Ernst, C.M., et al., *The bacterial defensin resistance protein MprF consists of separable domains for lipid lysinylation and antimicrobial peptide repulsion.* PLoS Pathog, 2009. **5**(11): p. e1000660.

15.   Ernst, C.M., et al., *The lipid-modifying multiple peptide resistance factor is an oligomer consisting of distinct interacting synthase and flippase subunits.* MBio, 2015. **6**(1).

16. Roy, H. and M. Ibba, *Broad range amino acid specificity of RNA-dependent lipid remodeling*
by multiple peptide resistance factors. J Biol Chem, 2009. **284**(43): p. 29677-83.
17. Ernst, C.M., et al., *Gain-of-Function Mutations in the Phospholipid Flippase MprF Confer*
Specific Daptomycin Resistance. MBio, 2018. **9**(6).
18. Prassler, J., et al., *HuCAL PLATINUM, a synthetic Fab library optimized for sequence diversity*
and superior performance in mammalian expression systems. J Mol Biol, 2011. **413**(1): p. 261-
78.
19. Kim, H.K., et al., *Recurrent infections and immune evasion strategies of Staphylococcus*
aureus. Curr Opin Microbiol, 2012. **15**(1): p. 92-9.
20. Bogdanov, M., et al., *Transmembrane protein topology mapping by the substituted cysteine*
accessibility method (SCAM(TM)): application to lipid-specific membrane protein topogenesis.
Methods, 2005. **36**(2): p. 148-71.
21. Yachdav, G., et al., *PredictProtein--an open resource for online prediction of protein structural*
and functional features. Nucleic Acids Res, 2014. **42**(Web Server issue): p. W337-43.
22. Otto, M., *Community-associated MRSA: what makes them special?* Int J Med Microbiol, 2013.
303(6-7): p. 324-30.
23. Pinheiro da Silva, F. and M.C. Machado, *The dual role of cathelicidins in systemic*
inflammation. Immunol Lett, 2017. **182**: p. 57-60.
24. Bayer, A.S., T. Schneider, and H.G. Sahl, *Mechanisms of daptomycin resistance in*
Staphylococcus aureus: role of the cell membrane and cell wall. Ann N Y Acad Sci, 2013. **1277**:
p. 139-58.
25. Jones, T., et al., *Failures in clinical treatment of Staphylococcus aureus Infection with*
daptomycin are associated with alterations in surface charge, membrane phospholipid
asymmetry, and drug binding. Antimicrob Agents Chemother, 2008. **52**(1): p. 269-78.
26. Peschel, A., et al., *Inactivation of the dlt operon in Staphylococcus aureus confers sensitivity*
to defensins, protegrins, and other antimicrobial peptides. J Biol Chem, 1999. **274**(13): p.
8405-10.
27. Yount, N.Y., et al., *Selective reciprocity in antimicrobial activity versus cytotoxicity of hBD-2*
and crotamine. Proc Natl Acad Sci U S A, 2009. **106**(35): p. 14972-7.
28. Slavetinsky, C.J., A. Peschel, and C.M. Ernst, *Alanyl-phosphatidylglycerol and lysyl-*
phosphatidylglycerol are translocated by the same MprF flippases and have similar capacities
to protect against the antibiotic daptomycin in Staphylococcus aureus. Antimicrob Agents
Chemother, 2012. **56**(7): p. 3492-7.
29. Spaan, A.N., et al., *Neutrophils versus Staphylococcus aureus: a biological tug of war.* Annu
Rev Microbiol, 2013. **67**: p. 629-50.
30. Kristian, S.A., et al., *MprF-mediated lysinylation of phospholipids in Staphylococcus aureus*
leads to protection against oxygen-independent neutrophil killing. Infect Immun, 2003. **71**(1):
p. 546-9.
31. Tabor, D.E., et al., *Staphylococcus aureus Alpha-Toxin Is Conserved among Diverse Hospital*
Respiratory Isolates Collected from a Global Surveillance Study and Is Neutralized by
Monoclonal Antibody MEDI4893. Antimicrob Agents Chemother, 2016. **60**(9): p. 5312-21.
32. Missiakas, D. and O. Schneewind, *Staphylococcus aureus vaccines: Deviating from the carol.* J
Exp Med, 2016. **213**(9): p. 1645-53.
33. Zheng, Y., et al., *Virulence Determinants Are Required for Brain Abscess Formation Through*
Staphylococcus aureus Infection and Are Potential Targets of Antivirulence Factor Therapy.
Front Microbiol, 2019. **10**: p. 682.
34. Wang, R., et al., *Identification of novel cytolytic peptides as key virulence determinants for*
community-associated MRSA. Nat Med, 2007. **13**(12): p. 1510-4.
35. Bae, T. and O. Schneewind, *Allelic replacement in Staphylococcus aureus with inducible*
counter-selection. Plasmid, 2006. **55**(1): p. 58-63.

36. Ieronymaki, M., et al., *Structure-Activity Relationship Studies, SPR Affinity Characterization,*
and Conformational Analysis of Peptides That Mimic the HNK-1 Carbohydrate Epitope.
ChemMedChem, 2017. **12**(10): p. 751-759.
37. Tiller, T., et al., *A fully synthetic human Fab antibody library based on fixed VH/VL framework*
pairings with favorable biophysical properties. MAbs, 2013. **5**(3): p. 445-70.
38. Neuber, T., et al., *Characterization and screening of IgG binding to the neonatal Fc receptor.*
MAbs, 2014. **6**(4): p. 928-42.
39. Bligh, E.G. and W.J. Dyer, *A rapid method of total lipid extraction and purification.* Can J
Biochem Physiol, 1959. **37**(8): p. 911-7.
40. Hanzelmann, D., et al., *Toll-like receptor 2 activation depends on lipopeptide shedding by*
bacterial surfactants. Nat Commun, 2016. **7**: p. 12304.
41. Iordanescu, S. and M. Surdeanu, *Two restriction and modification systems in Staphylococcus*
aureus NCTC8325. J Gen Microbiol, 1976. **96**(2): p. 277-81.
42. Bruckner, R., *A series of shuttle vectors for Bacillus subtilis and Escherichia coli.* Gene, 1992.
122(1): p. 187-92.