Functional correlates of immediate early gene expression in mouse visual cortex

David Mahringer[1,2], Pawel Zmarz[1,2,3], Hiroyuki Okuno[4], Haruhiko Bito[5] & Georg B. Keller[1,2,6]

[1] *Friedrich Miescher Institute for Biomedical Research, Basel, Switzerland.*
[2] *Faculty of Natural Sciences, University of Basel, Basel, Switzerland.*
[3] *Current address: Department of Organismic and Evolutionary Biology and Center for Brain Science,*
Harvard University, Cambridge, MA 02138, USA.
[4] *Department of Biochemistry and Molecular Biology, Kagoshima University Graduate School of Medical*
and Dental Sciences, Kagoshima, Kagoshima 890-8544, Japan.
[5] *Department of Neurochemistry, Graduate School of Medicine, The University of Tokyo, Hongo 7-3-1,*
Bunkyo-ku, Tokyo 113-0033, Japan.
[6] *Lead contact: georg.keller@fmi.ch*

During visual development, response properties of layer 2/3 neurons in visual cortex are shaped by

experience. Both visual and visuomotor experience are necessary to coordinate the integration of

bottom-up visual input and top-down motor-related input. Whether visual and visuomotor

experience engage different plasticity mechanisms, possibly associated with the two separate input

pathways, is still unclear. To begin addressing this, we measured the expression level of three

different immediate early genes (IEG) (*c-fos*, *egr1* or *Arc*) and neuronal activity in layer 2/3 neurons of

visual cortex before and after a mouse's first visual exposure in life, and subsequent visuomotor

learning. We found that expression levels of all three IEGs correlated positively with neuronal activity,

but that first visual and first visuomotor exposure resulted in differential changes in IEG expression

patterns. In addition, IEG expression levels differed depending on whether neurons exhibited

primarily visually driven or motor-related activity. Neurons with strong motor-related activity

preferentially expressed EGR1, while neurons that developed strong visually driven activity

preferentially expressed Arc. Our findings are consistent with the interpretation that bottom-up visual

input and top-down motor-related input are associated with different IEG expression patterns and

hence possibly also with different plasticity pathways.

*** Dear reader, please note this manuscript is formatted in a standard submission format. ***

INTRODUCTION

During first visuomotor exposure in life, experience with coupling between movement and visual

feedback is thought to coordination inputs onto layer 2/3 neurons in primary visual cortex (V1) such that

individual neurons receive balanced and opposing top-down motor-related and bottom-up visual input

(Attinger et al., 2017; Jordan and Keller, 2020; Leinweber et al., 2017). While visual input without

visuomotor coupling is sufficient to establish normal visual responses in layer 2/3 neurons, the

emergence of visuomotor mismatch responses is contingent on experience with visuomotor coupling

(Attinger et al., 2017) and requires on normal NMDA receptor dependent plasticity in the local V1 circuit

(Widmer and Keller, 2021). How top-down and bottom-up inputs are coordinated during visual and

visuomotor experience and whether plasticity mechanisms in the bottom-up visual driven pathway are

the same as those engaged in the top-down motor-related input is still unclear. Here we set out to test

for differences in plasticity mechanisms related to first exposure to visual input and first exposure to

normal visuomotor coupling using concurrent measurement of neuronal activity and the expression

levels of different immediate early gene (IEG) products as markers of plasticity in layer 2/3 of mouse

visual cortex.

IEG products play a critical role in synaptic and neuronal plasticity during learning (Chowdhury et al.,

2006; Fleischmann et al., 2003; Gandolfi et al., 2017; Jones et al., 2001; Messaoudi et al., 2007; Rial

Verde et al., 2006; Shepherd and Bear, 2011; Shepherd et al., 2006; Tzingounis and Nicoll, 2006;

Vazdarjanova et al., 2006; Veyrac et al., 2014; Waung et al., 2008) and are necessary for long-term

memory consolidation (Bozon et al., 2003; Fleischmann et al., 2003; Guzowski, 2002; Guzowski and

McGaugh, 1997; Guzowski et al., 2000; Jones et al., 2001; Ploski et al., 2008; Yasoshima et al., 2006).

Ever since the discovery that the expression of the transcription factor c-Fos can be induced by electrical

or chemical stimulation in neurons (Greenberg and Ziff, 1984), the expression of IEGs has been used as a

marker for neuronal activity (Bullitt, 1990; Guzowski et al., 1999; Jarvis et al., 2000; Knapska and

Kaczmarek, 2004; Minatohara et al., 2015; Morgan et al., 1987; Ramírez-Amaya et al., 2005;

Reijmers et al., 2007; Wang et al., 2021). Based on the discovery that certain forms of episodic

memory can be reactivated by artificially activating an ensemble of neurons characterized by high IEG

expression levels during memory acquisition (Denny et al., 2014; Garner et al., 2012; Liu et al., 2012;

Ramirez et al., 2013), it has been speculated that IEG expression is related not simply to neuronal

activity *per se*, but to the induction of activity-dependent plasticity (Holtmaat and Caroni, 2016; Josselyn

et al., 2015; Kaplan et al., 1996). Assuming IEG expression is indeed related to the induction of neuronal

plasticity, it is conceivable that different IEGs are preferentially involved in plasticity of different synapse

types or input pathways. c-Fos and EGR1 expression levels in visual cortex, for example, are differentially

regulated by visual experience and exhibit a differential dependence on neuromodulatory input

(Yamada et al., 1999). Consistent with a pathway-specific role of Arc and EGR1 in visual cortex, it has

been shown that Arc is necessary for different forms of plasticity of bottom-up visual input, including

ocular dominance plasticity (Gao et al., 2010; Jenks et al., 2017; McCurry et al., 2010; Wang et al., 2006),

while a knockout of *egr1* has been shown to leave ocular dominance plasticity unaffected (Mataga et al.,

2001). It is still unclear however, whether IEG expression is differentially regulated by changes in

bottom-up and top-down input and if there is preferential expression of different IEGs in neurons that

are predominantly excited by top-down motor-related input as compared to neurons that are

predominantly excited by bottom-up visual input.

RESULTS

To measure both IEG expression levels and neuronal activity chronically, we used a combination of

transgenic mice that express GFP under the control of an IEG promoter and viral delivery of a red variant

of a genetically encoded calcium indicator. We did this for three different IEGs (*c-fos*, *egr1*, and *Arc*), in

three groups of mice separately. EGFP-Arc and c-Fos-GFP mice are transgenic mice that express a fusion

protein of Arc or c-Fos, and GFP downstream of either an *Arc* or a *c-fos* promoter, respectively (Barth et

al., 2004; Okuno et al., 2012), while the EGR1-GFP mouse expresses GFP under an *egr1* promotor (Xie et

al., 2014). Although there are a number of caveats to using GFP levels in these mouse lines as a proxy for

IEG expression levels (see discussion), there is a strong overlap between post-mortem antibody staining

for the respective IEG and GFP expression in all three mouse lines (Barth et al., 2004; Okuno et al., 2012;

Xie et al., 2014; Yassin et al., 2010). Throughout the manuscript, we will use IEG expression to mean GFP

expression levels in these mice. To measure calcium activity, we used an AAV2/1-Ef1a-jRGECO1a viral

vector to express the genetically encoded red calcium indicator jRGECO1a (Dana et al., 2016). This

biased our recordings to excitatory neurons, as in the first few weeks after the injection, the Ef1a

promoter restricts expression mainly to excitatory neurons (Attinger et al., 2017).

To quantify the correlation between neuronal activity and IEG expression in layer 2/3 of visual cortex in

adult mice, we first used a paradigm of dark adaptation and subsequent brief visual exposure (**Figure

1A**). We did this in three groups of adult mice separately (4 EGFP-Arc mice, 4 c-Fos-GFP mice, and 4

EGR1-GFP mice, between 100 and 291 days old). We dark-adapted all three groups of mice for 24 hours

and subsequently head-fixed them, while still in complete darkness, under a two-photon microscope on

a spherical treadmill (**Figure 1A**). We then measured calcium activity and IEG levels every 15 minutes for

six hours (**Figures 1B-1D**; see Methods). Between the first and second measurement, mice were exposed

to visual input for 15 minutes. This paradigm, which is a combination of light exposure and exposure of

the mouse to head-fixation, resulted in transient and modest increases in Arc and EGR1 expression

levels, and a decrease in c-Fos expression levels (**Figure S1A**). Given that mean neuronal activity levels in

V1 are rapidly stabilized across light-dark transitions (Hengen et al., 2016), and only a prolonged dark

adaption (of 60 hours) results in a substantial increase of neuronal activity upon light exposure (Torrado

Pacheco et al., 2019), the absence of a strong IEG response is perhaps not surprising.

To quantify the relationship between neuronal activity and IEG expression levels more directly, we then

computed the correlation between average neuronal activity and IEG expression levels for each neuron

as a function of time between neuronal activity measurement and IEG expression measurement (**Figures**

1E-1G). Correlation peaked at a time lag of approximately 3.5 h ± 0.5 h (mean ± SEM) between neuronal

activity measurement and IEG measurement for Arc and c-Fos, and was positive but relatively stable in a

window from -2 hours to +3 hours for EGR1 consistent with previous results (Wang et al., 2021)(Arc:

1382 neurons, c-Fos: 1070 neurons, EGR1: 1319 neurons; **Figures 1E-1G**). At peak, the correlation

between neuronal activity and IEG expression was highest for c-Fos, intermediate for Arc, and lowest for

EGR1 (**Figures 1H-1J**; correlation coefficients for c-Fos: 0.39 ± 0.07, Arc: 0.26 ± 0.05, EGR1: 0.21 ± 0.03,

mean ± SEM; comparisons between c-Fos vs. Arc: $p < 3 \times 10^{-4}$, Arc vs. EGR1: p = 0.0188, c-Fos vs. EGR1: p

$< 10^{-8}$; 4 mice per group, t-test with bootstrapping, see Methods). The positive correlation and the time

lag of the correlation peak would be consistent with the idea that neuronal activity induces IEG

expression, but the fact that correlations with mean activity were relatively weak could mean that it is

specific patterns or types of activity that induce IEG expression. It is often assumed that IEG expression

is also a correlate of neuronal plasticity (Holtmaat and Caroni, 2016; Josselyn et al., 2015; Kaplan et al.,

1996). Given that certain forms of neuronal plasticity are associated with bursts of activity, we first

tested whether maximum activity was a better predictor of IEG expression levels than mean activity.

Indeed, we found that the correlation with maximum neuronal activity was higher than the correlation

with mean activity for all three IEGs, but only significantly so for Arc and EGR1 (**Figure S1B**). Thus, the

fact that we could detect a positive, albeit weak, relationship between neuronal calcium activity and IEG

expression, would suggest that the method of using transgenic mice that drive GFP under an IEG

promoter has sufficient resolution to detect physiologically relevant changes in IEG levels.

We thus proceeded to apply the same methods of measuring IEG dynamics and neuronal calcium

activity during a mouse's first exposure to visual input and subsequent first exposure to normal

visuomotor coupling. To investigate whether first visual and first visuomotor exposure results in a

differential expression of immediate early genes, we used a virtual reality system to experimentally

separate first visual exposure and first exposure to normal visuomotor coupling. To do this, we reared

mice in complete darkness and quantified both IEG expression levels and neuronal activity before and

after mice were exposed to visual input for the first time in life as well as during a subsequent phase of

visuomotor learning. Under normal conditions, first visual exposure is coincident with exposure to

normal visuomotor coupling. At eye opening, mice are capable of moving eyes, head, and body and thus

immediately experience self-generated visual feedback. To experimentally separate the moment of first

visual exposure from first exposure to normal visuomotor coupling, we recorded neuronal activity and

IEG expression as mice transitioned through three different experimental conditions. Prior to

experiments, three groups of mice were reared in complete darkness until postnatal day 40 (7 EGFP-Arc

mice, 5 c-Fos-GFP mice, and 4 EGR1-GFP mice). We then imaged neuronal activity and IEG expression

levels every 12 hours for a total of 6 days. During all two-photon imaging experiments, mice were head-

fixed on a spherical treadmill. During the first four recording sessions, mice were kept on the setup in

darkness to measure locomotion-related and non-visual activity and remained dark housed in between

recording sessions (condition 1). At the beginning of the 5^{th} recording session, mice were then exposed

to visual input for the first time in their life. In the subsequent four recording sessions, mice were

exposed to different virtual environments but remained housed in darkness in the time between the

recording sessions (condition 2). In addition to recording activity in darkness, recording sessions in

condition 2 also contained 8 min of closed-loop feedback during which visual flow on the walls of a

virtual corridor was coupled to the mouse's locomotion on the spherical treadmill. During closed-loop

feedback, we added brief halts of visual flow to probe for visuomotor mismatch responses (Keller et al.,

2012). This was followed by a phase of open-loop feedback during which the visual flow generated by

the mouse during the closed-loop feedback was replayed independently of the locomotion of the

mouse. Lastly, we presented a series of drifting gratings to the mouse to quantify visual responses (see

Methods). Following recording session 8, mice were introduced to a normal 12 h light / 12 h dark cycle.

At this time, mice first experienced normal visuomotor coupling in their home cage. We continued

recording for an additional four sessions (condition 3) with the same series of closed-loop, open-loop

and grating stimulation phases as in condition 2 (**Figure 2A**). Recording sessions lasted on average 12

min ± 0.5 min (mean ± SEM) in condition 1, and 83 min ± 1 min (mean ± SEM) in conditions 2 and 3

(**Figure S2**). Note, to probe for different visuomotor mismatch responses, condition 2 already contains a

short period of head-fixed, closed-loop feedback. Given the short duration and the fact that coupling

here is only in a small subspace (forward locomotion coupled to backward visual flow and normal

coupling for eye movements) of the total space of visuomotor coupling, we would expect the exposure

to normal visuomotor coupling in condition 3 to have a stronger influence on visuomotor integration.

The aim of this paradigm was to experimentally separate the first visual exposure at the beginning of

condition 2, from the first exposure to normal visuomotor coupling at the beginning of condition 3.

It has been shown that visual input can drive the expression of different IEGs in a subset of neurons in

visual cortex (Kaminska et al., 1996; Kawashima et al., 2013; Rosen et al., 1992; Tagawa et al., 2005;

Wang et al., 2006). Based on this it is sometimes assumed that neuronal activity in visual cortex is higher

with visual input than it is in darkness. Differences in neuronal activity between light and dark, however,

are small and predominantly transient (Fiser et al., 2004; Hengen et al., 2016; Torrado Pacheco et al.,

2019). To test whether in our paradigm first visual exposure or first exposure to normal visuomotor

coupling results in an increase of average neuronal activity in visual cortex, we quantified average

neuronal activity in each recording session (in condition 1 this only included recordings in darkness,

while in conditions 2 and 3 this included recordings in darkness, closed and open-loop feedback, as well

as drifting gratings). Consistent with a strong motor-related drive in visual cortex (Keller et al., 2012;

Saleem et al., 2013) and rapid homeostatic restoration of average activity following removal of visual

input (Keck et al., 2013), we found no evidence of an increase of average neuronal activity at the onset

of either condition 2 (first visual exposure) or condition 3 (first exposure to normal visuomotor coupling)

(**Figure 2B**). To the contrary, following the first visual exposure, there was a trend for decreasing activity

levels (p = 0.0293, R^2 = 0.371, linear trend analysis, see Methods). We next quantified average

expression of Arc, c-Fos and EGR1 over the same time course. Consistent with the absence of a change

in average activity levels, we found no significant changes in the expression levels of any of the three

IEGs following the first visual exposure at the beginning of condition 2 (**Figure 2C**). Note, we cannot

exclude that there is a transient increase in IEG expression between 1 h and 12 h following first visual

exposure, as we only recorded for 1 hour every 12 hours. We did however find that the first exposure to

normal visuomotor coupling at the beginning of condition 3, resulted in an increase in the expression of

Arc and a decrease in the expression of EGR1 in the absence of a measurable change in average

neuronal activity levels (**Figure 2C**). To test for changes in the pattern of IEG expression, we quantified

the similarity of IEG expression patterns by computing the correlation of IEG expression vectors

between imaging time points (see Methods). We found that the pattern of Arc expression changed both

with the first visual exposure (onset of condition 2) and the first exposure to normal visuomotor

coupling (onset of condition 3) (**Figure 2D**). The pattern of c-Fos expression exhibited no detectable

discontinuous changes (**Figure 2E**), while the pattern of EGR1 expression exhibited a marked transition

with the first exposure to normal visuomotor coupling (onset of condition 3) (**Figure 2F**). This suggests

that the expression patterns of IEGs are differentially and dynamically regulated by visuomotor

experience, also in absence of population mean expression level changes.

Neurons in layer 2/3 of primary visual cortex are driven differentially by visual and motor-related inputs

(Attinger et al., 2017; Keller et al., 2012; Leinweber et al., 2017). Given that the expression patterns of

the three IEGs are differentially altered by first visual exposure and first exposure to normal visuomotor

coupling, we speculated that the different IEGs could be preferentially expressed in different functional

types of excitatory neurons in layer 2/3. Neurons that are more strongly visually driven, likely by

bottom-up visual input, could have a different IEG expression profile than neurons that are more

strongly driven by top-down motor-related signals (Leinweber et al., 2017; Makino and Komiyama,

2015). To test this, we quantified the functional properties of the neurons with the highest IEG

expression levels immediately after the first exposure to normal visuomotor coupling where we

observed the largest mean IEG expression level changes (**Figure 2C**). We selected the 10 % of neurons

with the highest Arc, c-Fos and EGR1 expression, respectively, at the beginning of condition 3 (Arc: 197

neurons, c-Fos: 189 neurons, EGR1: 121 neurons) and tested whether these neurons were more strongly

driven by visual or motor-related input. As a measure of the strength of the motor-related input, we

used the magnitude of the neuronal response during running onsets in darkness. We found that neurons

with high EGR1 expression levels developed higher motor-related responses than the rest of the

population in both condition 2 and condition 3. Conversely, neurons with high Arc expression levels

developed motor-related responses that are lower than the rest of the population following exposure to

normal visuomotor coupling. Responses in neurons with high c-Fos expression levels were not different

from responses in the rest of the population (**Figure 3A**). To quantify the strength of visual input we

used the magnitude of the neuronal response to drifting grating stimuli. Consistent with the fact that

Arc expression can be selectively induced by visual stimuli in a stimulus-specific manner (Kawashima et

al., 2013), we found that neurons with high Arc expression levels developed responses to drifting grating

stimuli that were stronger than the rest of the population after exposure to normal visuomotor

coupling. The drifting grating responses of neurons with high EGR1 or c-Fos expression levels were not

different from the mean population response (**Figure 3B**). Thus, neurons with high levels of EGR1

expression after first exposure to normal visuomotor coupling were more strongly driven by motor-

related input, while those with high levels of Arc expression were more strongly driven by visual input.

One of the signals that has been speculated to be computed in mouse primary visual cortex that

combines visual and motor-related input is visuomotor mismatch (Attinger et al., 2017; Keller et al.,

2012; Zmarz and Keller, 2016). Neurons that respond to mismatch, or negative prediction errors, are

thought to receive excitatory motor-related input and inhibitory visual input (Attinger et al., 2017; Keller

and Mrsic-Flogel, 2018). We speculated that given the increased motor-related activity in neurons that

express high levels of EGR1, neuronal activity in these neurons should correlate positively with running,

while activity in neurons that express high levels of Arc should correlate positively with visual flow. To

quantify this, we computed the correlation of neuronal activity with either running or visual flow during

the open-loop phases in conditions 2 and 3 for the three groups of neurons with high IEG expression

levels. We found that the activity of neurons expressing high levels of EGR1 correlated most strongly

with running, while the activity of neurons with high levels of Arc expression correlated positively with

visual flow (**Figure 4A**). Consistent with this we found that visuomotor mismatch responses were larger

in neurons with high EGR1 expression than in the rest of the population, while they were lower in

neurons with high Arc expression than in the rest of the population (**Figure 4B**). This indicates that, at

the onset of normal visuomotor coupling, EGR1 is preferentially expressed in mismatch neurons or,

more generally, in neurons that are driven by excitatory top-down input, while Arc is preferentially

expressed in neurons that are driven by bottom-up visual input.

DISCUSSION

It is well established that both neuronal activity and plasticity are linked to the expression of immediate

early genes (Dudek, 2008; Mahringer et al., 2019; Minatohara et al., 2015; Wang et al., 2021; Yap and

Greenberg, 2018). Comparably little, however, is known about how specific functional characteristics of

neurons relate to the expression of immediate early genes. Here we investigated the relationship

between the expression of three IEGs (Arc, c-Fos, and EGR1) and functional responses in excitatory layer

2/3 neurons of mouse visual cortex. We found that during visuomotor learning following a mouse's first

visual exposure in life, Arc was preferentially expressed in neurons that are driven by excitatory bottom-

up visual input, while EGR1 was preferentially expressed in neurons that are driven by motor-related

input. In addition, we found that neurons expressing high levels of EGR1 exhibit visuomotor mismatch

responses higher than the rest of the population, while neurons expressing high levels of Arc exhibit

visuomotor mismatch responses weaker than the rest of the population.

Such a relationship between a neuron's IEG expression profile and its functional properties could be

explained by differences in the contribution of different IEGs to different types of input synapses. Arc, c-

Fos, and EGR1 all have unique cellular functions, and it is conceivable that they make different

contributions to different synapse types. Genes for a subset of $GABA_A$ receptor subunits, for example,

are transcriptional targets of EGR1 (Mo et al., 2015). If the postsynaptic subunit composition of the

GABA receptor is correlated with the presynaptic inhibitory cell type, EGR1 expression could

preferentially upregulate specific inhibitory input pathways. Similar input pathway-specific roles have

been described for other IEGs. Neuronal activity-regulated pentraxin (NARP) is secreted by pyramidal

neurons and exclusively accumulates at parvalbumin-positive inhibitory neurons where it regulates

excitatory synapses onto these cells (Chang et al., 2010; Gu et al., 2013). The activity-dependent

transcription factor NPAS4 has been found to restrict the number of synapses of mossy-fiber input

specifically onto CA3 pyramidal cells during learning (Weng et al., 2018).

Our data would be consistent with the interpretation that the IEG expression pattern of a given neuron

correlates with its pattern of synaptic inputs. Neurons that predominantly receive excitatory bottom-up

drive likely require a different distribution and type of input synapses compared to neurons that receive

mainly top-down excitatory drive. Layer 2/3 neurons that exhibit strong motor-related and mismatch

responses are thought to be driven by top-down excitatory inputs (Leinweber et al., 2017), which

predominantly target apical dendrites (Petreanu et al., 2009). Conversely, layer 2/3 neurons with strong

visual responses are thought to be driven by bottom-up visual inputs, which predominantly target basal

dendrites (Petreanu et al., 2009). We have speculated that mismatch neurons that receive motor-

related input also receive matched bottom-up inhibitory input from a specific subset of somatostatin

(SST)-positive interneurons (Attinger et al., 2017). Thus, EGR1 expression may be preferentially

increased in neurons that are driven by excitatory top-down input and SST mediated bottom-up

inhibition, while Arc expression may be preferentially increased in neurons that are driven by excitatory

bottom-up visual input. This may explain why a change to the visual input alone at first visual exposure

primarily resulted in a rearrangement of the Arc expression pattern (**Figure 2D**) but left the EGR1

expression pattern relatively unaffected (**Figure 2F**), while first exposure to normal visuomotor coupling

resulted in a rearrangement of the expression pattern of both Arc and EGR1.

When interpreting our results, it should be kept in mind that both the method we use to approximate

IEG expression levels and the method we use to approximate neuronal activity levels come with a series

of caveats. In the case of the transgenic mice used for the IEG expression measurements, two of these

express a fusion protein (Arc and c-Fos), where the IEG is likely overexpressed (Steward et al., 2017), and

it is possible that the decay kinetics of the fusion protein differ from those of the native protein. In the

case of the GFP driven by the *egr1* promoter, the GFP decay kinetics are likely different from the decay

kinetics of EGR1. However, these potential differences in decay kinetics and expression levels do not

completely mask the correlation between IEG expression levels and reporter proteins. In post-mortem histological stainings the expression of GFP in these mouse lines overlaps well with the expression levels of the IEGs (Barth et al., 2004; Okuno et al., 2012; Wang et al., 2021; Xie et al., 2014; Yassin et al., 2010). Thus, reporter protein levels reflect a filtered version of IEG expression levels, and the two are likely related by a monotonic function. Given that all our analyses rely only on relative expression levels among populations of simultaneously recorded neurons or relative changes of expression levels in time, the lack of a direct measurement of IEG expression levels should not change our conclusions. A second caveat concerns the genetically encoded calcium indicator used to measure neuronal activity. Our activity measures are biased towards bursts of neuronal activity, as single spikes are probably not always detectable using calcium indicators *in vivo*. However, even though the transfer function from neuronal activity to calcium signal is non-linear, it is also monotonic. Thus, we may be underestimating the correlation between neuronal activity and IEG expression, but neither caveat would bias the results towards finding specific correlations between different IEGs and functional cell types.

In summary, our results suggest that the expression of Arc and EGR1 in layer 2/3 neurons in mouse visual cortex may be a correlate of the type of functional input the neuron receives. Such a preference for expression in a functionally specific subset of neurons would be consistent with differential changes in the ratio of the expression of different IEGs under conditions that result in identical mean levels of neuronal activity (Bailey and Wade, 2003; Farina and Commins, 2016; Guzowski et al., 2006) that are difficult to explain if IEG expression were simply driven by mean activity. In future experiments, it will be important to establish a more detailed picture of how immediate early genes could orchestrate or stabilize the pattern of functionally distinct input streams a neuron receives.

Figure 1. Simultaneous imaging of neuronal activity and immediate early gene expression in visual

cortex.

(**A**) Left: Schematic of the virtual reality setup used for imaging experiments. Right: Schematic of the
experimental timeline. Mice were dark-adapted for 24 hours. Neuronal activity and IEG expression levels
were recorded in 25 imaging sessions starting immediately before and continuing until 6 hours after
visual stimulation in intervals of 15 minutes.

(**B**) Example two-photon images of neurons in primary visual cortex labelled with jRGECO1a (red, top
left), Arc (green, bottom left), and the overlay (right). Scale bar is 50 μm.

(**C**) Same as in (**B**), but for c-Fos.

(**D**) Same as in (**B**), but for EGR1.

(**E**) Correlation of average activity and IEG expression level as a function of the time difference between
the two measurements. Dotted line indicates average correlation, shading indicates standard error of
the mean (SEM) across mice (n = 4).

(**F**) Same as in (**E**), but for c-Fos mice (n = 4).

(**G**) Same as in (**E**), but for EGR1 mice (n = 4).

(**H**) Scatter plot of Arc expression 3.5 hours after visual stimulation and average neuronal activity during
visual stimulation (1382 neurons in 4 mice, 83 neurons outside of plot range). Shown in the panel is the
average correlation coefficient across mice (mean ± SEM, n= 4).

(**I**) Same as in (**H**), but for c-Fos (1070 neurons in 4 mice, 28 neurons outside of plot range).

(**J**) Same as (**H**), but for EGR1 (1319 neurons in 4 mice, 18 neurons outside of plot range).

Figure 2. IEG expression dynamics during visuomotor learning.

(**A**) Schematic of the experimental timeline. Mice were born and reared in complete darkness. jRGECO1a was injected 10 to 12 days prior to the start of imaging experiments. We then imaged calcium activity and IEG expression levels every 12 h over the course of 6 days both before and after first visual exposure and first exposure to normal visuomotor coupling. On the first two days (condition 1) activity in visual cortex was recorded in complete darkness while mice were head-fixed and free to run on a spherical treadmill. On the third day of recording mice were exposed to visual feedback (first visual exposure) in a virtual reality environment. Outside of the recording sessions mice were still housed in complete darkness (condition 2). Starting on day 5, mice were subjected to a 12 h / 12 h light/dark cycle (condition 3).

(**B**) Average calcium activity during all conditions (condition 1 vs. 2: p = 0.2183, condition 2 vs. 3: p = 0.527, condition 1 vs. 3: p = 0.0123, 5067 neurons, paired t-test). Shading is SEM over mice. Dashed line indicates linear fit to the data of conditions 2 and 3. The linear fit to the data from conditions 2 and 3 exhibited a significant negative slope (p = 0.0293, R^2 = 0.371, linear trend analysis, see Methods).

(**C**) Normalized mean IEG expression levels during all conditions. Expression level of Arc (blue, 1969 neurons in 7 mice) significantly increased after first exposure to visuomotor coupling, decreased for EGR1 (green, 1213 neurons in 4 mice) and remained unchanged for c-Fos (orange, 1885 neurons in 5 mice). Change in IEG expression level between conditions 1 and 2 for Arc: 0.1764 ± 0.1556, p = 0.2775; c-Fos: -0.0536 ± 0.1877, p = 0.7816; EGR1: -0.0371 ± 0.1246, p = 0.7745 (mean ± SEM, paired t-test). Change in IEG expression level between conditions 2 and 3 for Arc: 1.2628 ± 0.5012, p = 0.0256; c-Fos: 0.01612 ± 0.1372, p = 0.2702; EGR1: -0.4568 ± 0.1130, p = 0.0049 (mean ± SEM, t-test). Shading indicates SEM over mice.

352   (**D**) Average rank order correlation coefficients for Arc expression during visuomotor learning (7 mice).

353   The expression pattern changes both at the onset of conditions 2 and 3.

354   (**E**) Same as in (**D**), but for c-Fos (5 mice). The expression pattern exhibits no apparent transitions.

355   (**F**) Same as in (**D**), but for EGR1 (4 mice). The expression pattern changes at the onset of condition 3.

Figure 3. Differential relationship between IEG expression and motor-related and visual responses.

(**A**) Average running onset response during darkness for the top 10 % IEG expressing neurons (Arc: 197 neurons, c-Fos: 189 neurons, EGR1: 121 neurons). Neuronal responses were pooled from all experimental sessions for each condition, subtracted by the mean and normalized by the standard deviation of the response of all neurons (Z-score). Error bars are SEM over neurons. Statistics above the plot indicate comparisons against 0, statistics to the right are between-group comparisons. n.s.: p > 0.05, *: p < 0.05, **: p < 0.01, *** p < 0.001, t-test.

(**B**) Average grating onset response for the top 10 % IEG expressing neurons (Arc: n = 197, c-Fos: n = 189, EGR1: n = 121). Neuronal responses were pooled from all experimental sessions for conditions 2 and 3, subtracted by the mean and normalized by standard deviation of the response of all neurons (Z-score). Error bars are SEM over neurons. Statistics above the plot indicate comparisons against 0, statistics to the right are between-group comparisons. n.s.: p > 0.05, *: p < 0.05, **: p < 0.01, *** p < 0.001, t-test.

Figure 4. Functional cell type specific expression of IEGs in visual cortex.

(**A**) Correlation of neuronal activity with running and of neuronal activity with visual flow during open-loop phases of conditions 2 and 3 for the top 10 % IEG expressing neurons (Arc: 197 neurons, c-Fos: 189 neurons, EGR1: 121 neurons). Inset: Average correlation coefficient for the three groups of high IEG expressing neurons. High Arc expressing neurons had the highest correlation with visual flow (Arc vs. c-Fos: $p < 10^{-10}$, Arc vs. EGR1: $p < 10^{-9}$, c-Fos vs. EGR1: $p = 0.0791$, t-test), while high EGR1 expressing neurons had the highest correlation with running (Arc vs. c-Fos: $p < 10^{-10}$, Arc vs. EGR1: $p < 10^{-8}$, c-Fos vs. EGR1: $p < 10^{-10}$, t-test).

(**B**) Mismatch responses in condition 3 were significantly higher for the top 10 % EGR1 expressing neurons and significantly lower for the top 10 % Arc expressing neurons than the rest of the respective population (Arc: 197 neurons, c-Fos: 189 neurons, EGR1: 121 neurons). Arc: $p = 0.0461$, c-Fos: $p = 0.2273$, EGR1: $p = 0.0234$; Arc vs. c-Fos: $p = 0.0101$, c-Fos vs. EGR1: $p = 0.048$, Arc vs. EGR1: $p = 0.0057$, t-test.

388 **Supplementary Figures**

(**A**) Time course of normalized IEG expression levels following 24 h dark adaptation and 15 min visual

stimulation at time 0. Shading indicates SEM over neurons.

(**B**) Correlation coefficient of mean and maximum activity (average across or peak within a recording

session, respectively) with IEG expression 3.5 h after stimulation or recording onset (Arc: 11 mice, c-Fos:

9 mice, EGR1: 8 mice). Box whisker plot: red line indicates median, box marks 25th to 75th percentiles

and whiskers extended to the next most extreme datapoint within a range of 1.5 times the interquartile

distance (rank sum test, Arc: $p = 0.0086$, c-Fos: $p = 0.1359$, EGR1: $p = 0.0207$).

Figure S2. Duration of recording sessions. Related to Figure 2.

Histogram of the durations of the recording sessions. On average, one recording session lasted for

approximately 12 min during condition 1 (solid line) and, due to the addition of closed-loop, open-loop,

and grating stimulation phases, 83 min during conditions 2 and 3 (dashed line).

METHODS

Animals and surgery. All animal procedures were approved by and carried out in accordance with guidelines of the Veterinary Department of the Canton Basel-Stadt, Switzerland. We used imaging data from a total of 11 EGFP-Arc mice (Okuno et al., 2012), 9 c-Fos-GFP mice (Barth et al., 2004) and 8 EGR1-GFP mice (Xie et al., 2014), aged 40 days at the start of visuomotor learning (**Figures 2 - 4**) or aged 100-104 (Arc), 279-291 (c-Fos) or 120-124 (EGR1) days (**Figure 1**). Sample sizes were chosen according to the standards in the field and no statistical methods were used to predetermine sample sizes. Mice were group-housed in a dark cabinet and in a vivarium (light/dark cycle: 12 h / 12 h). Viral injections and window implantation were performed as previously described (Dombeck et al., 2010; Leinweber et al., 2014). Briefly, for sensorimotor learning experiments, mice (aged 29 d ± 1 d, mean ± SEM) were anesthetized in darkness using a mix of fentanyl (0.05 mg/kg), medetomidine (0.5 mg/kg) and midazolam (5 mg/kg), and additionally their eyes were covered with a thick, black cotton fabric during all surgical procedures. A 3 mm to 5 mm craniotomy was made above visual cortex (2.5 mm lateral of lambda (Paxinos and Franklin, 2013)) and AAV2/1-Ef1a-NES-jRGECO1a-WPRE ((Dana et al., 2016); titer: between 7.2×10^{10} GC/ml and 6.8×10^{12} GC/ml) was injected into the target region. The craniotomy was sealed with a fitting cover slip. A titanium head bar was attached to the skull and stabilized with dental cement.

Imaging and virtual reality. Imaging commenced 10 – 12 (visuomotor learning experiments, **Figures 2 - 4**) or 12 – 29 (**Figure 1**) days following virus injection and was carried out using a custom-built two-photon microscope. Illumination source was a Chameleon Vision laser (Coherent) tuned to a wavelength of either 950 nm, 990 nm or 1030 nm. Imaging was performed using an 8 kHz resonance scanner (Cambridge Technology) resulting in frame rates of 40 Hz at a resolution of 400 × 750 pixels. In addition, we used a piezo actuator (Physik Instrumente) to move the objective (Nikon 16x, 0.8 NA) in steps of 15 µm between frames to acquire images at four different depths, thus reducing the effective frame rate to 10 Hz. The behavioral imaging setup was as previously described (Leinweber et al., 2014). After brief isoflurane anesthesia mice were head-fixed in complete darkness and the setup was light-shielded before every imaging session. Mice were free to run on an air-supported polystyrene ball, the motion of which was restricted to the forward and backward directions by a pin. The ball's rotation was coupled to linear displacement in the virtual environment that was projected onto a toroidal screen surrounding the mouse. The screen covered a visual field of approximately 240 degrees horizontally and 100 degrees vertically. All displayed elements of the tunnel or sinusoidal gratings were calibrated to be isoluminant.

Experimental design. For experiments shown in **Figure 1**, mice were dark-adapted for 24 h and 17 min ±

10 min (mean ± SEM, 12 mice) before head fixation under the microscope in darkness. Activity and

immediate early gene expression were recorded every 15 minutes for 6 hours. Except for the time of

visual stimulation with sinusoidal gratings moving in 8 different directions (a total of 80 presentation in

random order), mice were kept in complete darkness under the microscope for the duration of the

entire experiment. For visuomotor learning experiments (**Figures 2 - 4**) mice were born and reared in

complete darkness until P44 and then transferred to a vivarium with a 12 h /12 h light/dark cycle.

Experimental sessions started on P40 and occurred twice per day, spaced 12 h apart. In condition 1, all

imaging was done in complete darkness and experiments consisted of recording approximately 8 min of

neuronal activity during which mice were free to run on the spherical treadmill. IEG expression level

measurements were taken before and after each activity recording. In conditions 2 and 3, neuronal

activity measurements consisted of 7 recordings of approximately 8 minutes each. Each recording

session started with a recording in darkness, followed by a closed-loop recording. In the closed-loop

recording, the movement of the mouse in a linear virtual corridor (sinusoidal vertical grating) was

coupled to the locomotion of the mouse on the spherical treadmill. During the closed-loop session we

included brief (1 s) halts of visual flow to induce mismatch events (Attinger et al., 2017). The subsequent

two recordings were of the open-loop type and consisted of a playback of the visual flow the mouse had

generated during the preceding closed-loop recording. Subsequently, mice were exposed to a second

recording in darkness, followed by a visual stimulation recording. During the visual stimulation

sinusoidal moving grating stimuli (2 second standing grating, 3 second drifting grating, 8 different

orientations, 10 presentations of each orientation, in a randomized order) were presented. Finally, mice

were exposed to a third recording in darkness. In early phases of the experiment mice were encouraged

to run by applying occasional mild air puffs to the neck.

Data analysis. Imaging data were full-frame registered using a custom-written software (Leinweber et

al., 2014). Neurons were selected manually based on their mean fluorescence or maximum projection in

the red channel (jRGECO1a). This biased our selection towards active neurons. Fluorescence traces were

calculated as the mean pixel value in each region of interest per frame, and were then median-

normalized to calculate ΔF/F. ΔF/F traces were filtered as previously described (Dombeck et al., 2007).

GFP intensities were calculated as the mean pixel value in each region of interest (ROI) for mean

fluorescence projections. To compensate for expression level differences between different IEG mouse

lines as well as for image quality differences between different mice we normalized the GFP level

measurements as follows: For each mouse, all ROI measurements were subtracted by the minimum

calculated over all ROIs and timepoints, and normalized by the median over all ROIs and timepoints.

$$ROI^{i,tp}_{normalized} = (ROI^{i,tp} - \min_{i,tp} ROI^{i,tp}) \, / \, (\underset{i,tp}{\text{median}} \, ROI^{i,tp} - \min_{i,tp} ROI^{i.tp})$$

This ensured that the minimum value of IEG expression was 0 and the median 1. No blinding of

experimental condition was performed in any of the analyses. Statistical tests were used as stated in the

figure legends.

Figure 1. Examples images (**Figures 1B-1D**) are average projections of the recorded channel. IEG

expression was normalized as described above (**Figures 1E-1G**). Correlation coefficients (**Figures 1E-1G**)

were calculated based on the neuronal population vectors of average activity and IEG expression per

measurement timepoint, for each mouse. For the statistical comparison of the correlation coefficients of

IEG expression levels with neural activity between the three different groups (4 mice per group), data

were bootstrapped 5 times with random replacement and then a t-test was performed on the

bootstrapped data.

Figure 2. To compare changes in neural activity and IEG expression levels between conditions we

averaged data from the last two recording sessions of the previous condition and the first two recording

sessions of the following condition (**Figures 2B and 2C**). Linear trend analysis (**Figure 2B**) was performed

using the MATLAB regress function. To quantify the significance of the linear trend we report the R^2

statistic and p-value of the F statistic. Linear fits were performed for each mouse individually using the

MATLAB polyfit and polyval functions. Rank order correlation coefficients (**Figures 2D-2F**) were

determined based on the population vectors of average IEG expression per measurement timepoint and

mouse, and then averaged.

Figure 3. For plots of event-triggered activity changes ΔF/F traces were baseline-subtracted by the

average ΔF/F in a window −500 ms to −100 ms preceding the event onset. Z-scores were obtained on a

population vector with average stimulus onset values calculated over a response window of 1.5 s. High

IEGs neurons were selected as the top 10% of IEG expressing neurons based on average expression level

on the first day of condition 3.

Figure 4. Correlation coefficients (**Figure 4A**) were calculated by correlating each neuron's activity trace

with either the running trace or the visual flow trace during open-loop phases. High IEG neurons were

selected with the same criteria used for **Figure 3**. Stimulus-triggered fluorescence changes (**Figure 4B**)

were mean-subtracted in a window −500 ms to −100 ms preceding the stimulus onset. Responses were

quantified in a window of 1.5 s.

Figure S1. Correlation coefficients of mean or maximum activity with IEG expression were calculated for

each mouse (**Figure S1B**). Mice from visual stimulation experiments (**Figure 1**) and sensorimotor learning

experiments (**Figures 2-4**) were pooled for this analysis. For mice from the sensorimotor learning

experiments the calculation was done using mean or maximum activity of the first recording segment

and the last IEG measurement within a session. Shown is the average correlation across all sessions.

Code and data availability. All imaging and image processing code can be found online at

https://sourceforge.net/projects/iris-scanning/ (IRIS, imaging software package) and

https://sourceforge.net/p/iris-scanning/calliope/HEAD/tree (Calliope, image processing software

package). All the raw data and analysis code used in this study can be downloaded from the following

website: http://data.fmi.ch/PublicationSupplementRepo/.

ACKNOWLEDGEMENTS

We thank the entire Keller lab for helpful discussion and comments on earlier versions of this

manuscript. We thank Daniela Gerosa-Erni for production of the AAV vectors. This work was supported

by the Swiss National Science Foundation (GBK), the Novartis Research Foundation (GBK), the Human

Frontier Science Program (GBK), and JSPS-Kakenhi and AMED (HO and HB).

AUTHOR CONTRIBUTIONS

D.M. and P.Z. performed the experiments, D.M. analyzed the data. H.O. and H.B. made the EGFP-Arc

mouse. All authors wrote the manuscript.

REFERENCES

Attinger, A., Wang, B., and Keller, G.B. (2017). Visuomotor Coupling Shapes the Functional Development
of Mouse Visual Cortex. Cell *169*, 1291-1302.e14.

Bailey, D.J., and Wade, J. (2003). Differential expression of the immediate early genes FOS and ZENK
following auditory stimulation in the juvenile male and female zebra finch. Brain Research. Molecular
Brain Research *116*, 147–154.

Barth, A.L., Gerkin, R.C., and Dean, K.L. (2004). Alteration of Neuronal Firing Properties after In Vivo
Experience in a FosGFP Transgenic Mouse. The Journal of Neuroscience *24*, 6466–6475.

Bozon, B., Kelly, A., Josselyn, S.A., Silva, A.J., Davis, S., and Laroche, S. (2003). MAPK, CREB and zif268 are
all required for the consolidation of recognition memory. Philosophical Transactions of the Royal Society
of London. Series B, Biological Sciences *358*, 805–814.

Bullitt, E. (1990). Expression of C-fos-like protein as a marker for neuronal activity following noxious
stimulation in the rat. The Journal of Comparative Neurology *296*, 517–530.

Chang, M.C., Park, J.M., Pelkey, K.A., Grabenstatter, H.L., Xu, D., Linden, D.J., Sutula, T.P., McBain, C.J.,
and Worley, P.F. (2010). Narp regulates homeostatic scaling of excitatory synapses on parvalbumin-
expressing interneurons. Nat Neurosci *13*, 1090–1097.

Chowdhury, S., Shepherd, J.D., Okuno, H., Lyford, G., Petralia, R.S., Plath, N., Kuhl, D., Huganir, R.L., and
Worley, P.F. (2006). Arc/Arg3.1 Interacts with the Endocytic Machinery to Regulate AMPA Receptor
Trafficking. Neuron *52*, 445–459.

Dana, H., Mohar, B., Sun, Y., Narayan, S., Gordus, A., Hasseman, J.P., Tsegaye, G., Holt, G.T., Hu, A.,
Walpita, D., et al. (2016). Sensitive red protein calcium indicators for imaging neural activity. ELife *5*.

Denny, C.A., Kheirbek, M.A., Alba, E.L., Tanaka, K.F., Brachman, R.A., Laughman, K.B., Tomm, N.K., Turi,
G.F., Losonczy, A., and Hen, R. (2014). Hippocampal Memory Traces Are Differentially Modulated by
Experience, Time, and Adult Neurogenesis. Neuron *83*, 189–201.

Dombeck, D.A., Khabbaz, A.N., Collman, F., Adelman, T.L., and Tank, D.W. (2007). Imaging Large-Scale
Neural Activity with Cellular Resolution in Awake, Mobile Mice. Neuron *56*, 43–57.

Dombeck, D.A., Harvey, C.D., Tian, L., Looger, L.L., and Tank, D.W. (2010). Functional imaging of
hippocampal place cells at cellular resolution during virtual navigation. Nature Neuroscience *13*, 1433–
1440.

Dudek, S. (2008). Transcriptional Regulation by Neuronal Activity (Boston, MA: Springer US).

Farina, F.R., and Commins, S. (2016). Differential expression of immediate early genes Zif268 and c-Fos
in the hippocampus and prefrontal cortex following spatial learning and glutamate receptor antagonism.
Behavioural Brain Research *307*, 194–198.

Fiser, J., Chiu, C., and Weliky, M. (2004). Small modulation of ongoing cortical dynamics by sensory input
during natural vision. Nature *431*, 573–578.

Fleischmann, A., Hvalby, O., Jensen, V., Strekalova, T., Zacher, C., Layer, L.E., Kvello, A., Reschke, M.,
Spanagel, R., Sprengel, R., et al. (2003). Impaired Long-Term Memory and NR2A-Type NMDA Receptor-
Dependent Synaptic Plasticity in Mice Lacking c-Fos in the CNS. The Journal of Neuroscience *23*, 9116–
9122.

Gandolfi, D., Cerri, S., Mapelli, J., Polimeni, M., Tritto, S., Fuzzati-Armentero, M.-T., Bigiani, A., Blandini,
F., Mapelli, L., and D'Angelo, E. (2017). Activation of the CREB/c-FosPathway during Long-Term Synaptic
Plasticity in the Cerebellum Granular Layer. Frontiers in Cellular Neuroscience *11*, 184.

Gao, M., Sossa, K., Song, L., Errington, L., Cummings, L., Hwang, H., Kuhl, D., Worley, P., and Lee, H.-K.
(2010). A Specific Requirement of Arc/Arg3.1 for Visual Experience-Induced Homeostatic Synaptic
Plasticity in Mouse Primary Visual Cortex. The Journal of Neuroscience *30*, 7168–7178.

Garner, A.R., Rowland, D.C., Hwang, S.Y., Baumgaertel, K., Roth, B.L., Kentros, C., and Mayford, M.
(2012). Generation of a Synthetic Memory Trace. Science (New York, N.Y.) *335*, 1513–1516.

Greenberg, M.E., and Ziff, E.B. (1984). Stimulation of 3T3 cells induces transcription of the c-fos proto-
oncogene. Nature *311*, 433–438.

Gu, Y., Huang, S., Chang, M.C., Worley, P., Kirkwood, A., and Quinlan, E.M. (2013). Obligatory Role for
the Immediate Early Gene NARP in Critical Period Plasticity. Neuron *79*, 335–346.

Guzowski, J.F. (2002). Insights into immediate-early gene function in hippocampal memory consolidation
using antisense oligonucleotide and fluorescent imaging approaches. Hippocampus *12*, 86–104.

Guzowski, J.F., and McGaugh, J.L. (1997). Antisense oligodeoxynucleotide-mediated disruption of
hippocampal cAMP response element binding protein levels impairs consolidation of memory for water
maze training. Proceedings of the National Academy of Sciences of the United States of America *94*,
2693–2698.

Guzowski, J.F., McNaughton, B.L., Barnes, C.A., and Worley, P.F. (1999). Environment-specific expression
of the immediate-early gene Arc in hippocampal neuronal ensembles. Nature Neuroscience *2*, 1120–
1124.

Guzowski, J.F., Lyford, G.L., Stevenson, G.D., Houston, F.P., McGaugh, J.L., Worley, P.F., and Barnes, C. a
(2000). Inhibition of activity-dependent arc protein expression in the rat hippocampus impairs the
maintenance of long-term potentiation and the consolidation of long-term memory. The Journal of
Neuroscience : The Official Journal of the Society for Neuroscience *20*, 3993–4001.

Guzowski, J.F., Miyashita, T., Chawla, M.K., Sanderson, J., Maes, L.I., Houston, F.P., Lipa, P., McNaughton,
B.L., Worley, P.F., and Barnes, C.A. (2006). Recent behavioral history modifies coupling between cell
activity and Arc gene transcription in hippocampal CA1 neurons. Proceedings of the National Academy
of Sciences of the United States of America *103*, 1077–1082.

Hengen, K.B., Torrado Pacheco, A., McGregor, J.N., Van Hooser, S.D., and Turrigiano, G.G. (2016).
Neuronal Firing Rate Homeostasis Is Inhibited by Sleep and Promoted by Wake. Cell *165*, 180–191.

Holtmaat, A., and Caroni, P. (2016). Functional and structural underpinnings of neuronal assembly
formation in learning. Nature Neuroscience *19*, 1553–1562.

Jarvis, E.D., Ribeiro, S., Da Silva, M.L., Ventura, D., Vielliard, J., and Mello, C. V. (2000). Behaviourally
driven gene expression reveals song nuclei in hummingbird brain. Nature *406*, 628–632.

Jenks, K.R., Kim, T., Pastuzyn, E.D., Okuno, H., Taibi, A. V, Bito, H., Bear, M.F., and Shepherd, J.D. (2017).
Arc restores juvenile plasticity in adult mouse visual cortex. Proceedings of the National Academy of
Sciences *114*, 9182–9187.

Jones, M.W., Errington, M.L., French, P.J., Fine, A., Bliss, T.V.P., Garel, S., Charnay, P., Bozon, B., Laroche,
S., and Davis, S. (2001). A requirement for the immediate early gene Zif268 in the expression of late LTP
and long-term memories. Nature Neuroscience *4*, 289–296.

Jordan, R., and Keller, G.B. (2020). Opposing Influence of Top-down and Bottom-up Input on Excitatory
Layer 2/3 Neurons in Mouse Primary Visual Cortex. Neuron *108*, 1194-1206.e5.

Josselyn, S.A., Köhler, S., and Frankland, P.W. (2015). Finding the engram. Nature Reviews Neuroscience
16, 521–534.

Kaminska, B., Kaczmarek, L., and Chaudhuri, A. (1996). Visual Stimulation Regulates the Expression of
Transcription Factors and Modulates the Composition of AP-1 in Visual Cortexa. Journal of
Neuroscience *16*, 3968–3978.

Kaplan, I. V, Guo, Y., and Mower, G.D. (1996). Immediate early gene expression in cat visual cortex
during and after the critical period: differences between EGR-1 and Fos proteins. Brain Research.
Molecular Brain Research *36*, 12–22.

Kawashima, T., Kitamura, K., Suzuki, K., Nonaka, M., Kamijo, S., Takemoto-Kimura, S., Kano, M., Okuno,
H., Ohki, K., and Bito, H. (2013). Functional labeling of neurons and their projections using the synthetic
activity–dependent promoter E-SARE. Nature Methods *10*, 889–895.

Keck, T., Keller, G.B., Jacobsen, R.I., Eysel, U.T., Bonhoeffer, T., and Hübener, M. (2013). Synaptic scaling
and homeostatic plasticity in the mouse visual cortex in vivo. Neuron *80*.

Keller, G.B., and Mrsic-Flogel, T.D. (2018). Predictive Processing: A Canonical Cortical Computation.
Neuron *100*, 424–435.

Keller, G.B., Bonhoeffer, T., and Hübener, M. (2012). Sensorimotor mismatch signals in primary visual
cortex of the behaving mouse. Neuron *74*, 809–815.

Knapska, E., and Kaczmarek, L. (2004). A gene for neuronal plasticity in the mammalian brain:
Zif268/Egr-1/NGFI-A/Krox-24/TIS8/ZENK? Progress in Neurobiology *74*, 183–211.

Leinweber, M., Zmarz, P., Buchmann, P., Argast, P., Hübener, M., Bonhoeffer, T., and Keller, G.B. (2014).
Two-photon calcium imaging in mice navigating a virtual reality environment. Journal of Visualized
Experiments : JoVE e50885.

Leinweber, M., Ward, D.R., Sobczak, J.M., Attinger, A., and Keller, G.B. (2017). A Sensorimotor Circuit in
Mouse Cortex for Visual Flow Predictions. Neuron *95*, 1420-1432.e5.

Liu, X., Ramirez, S., Pang, P.T., Puryear, C.B., Govindarajan, A., Deisseroth, K., and Tonegawa, S. (2012).
Optogenetic stimulation of a hippocampal engram activates fear memory recall. Nature *484*, 381–385.

Mahringer, D., Petersen, A., Fiser, A., Okuno, H., Bito, H., Perrier, J.-F., and Keller, G. (2019). Expression
of c-Fos and Arc in hippocampal region CA1 marks neurons that exhibit learning-related activity changes.
BioRxiv 644526.

Makino, H., and Komiyama, T. (2015). Learning enhances the relative impact of top-down processing in
the visual cortex. Nature Neuroscience *18*, 1116–1122.

Mataga, N., Fujishima, S., Condie, B.G., and Hensch, T.K. (2001). Experience-Dependent Plasticity of
Mouse Visual Cortex in the Absence of the Neuronal Activity-Dependent Markeregr1/zif268 . The
Journal of Neuroscience *21*, 9724–9732.

McCurry, C.L., Shepherd, J.D., Tropea, D., Wang, K.H., Bear, M.F., and Sur, M. (2010). Loss of Arc renders
the visual cortex impervious to the effects of sensory experience or deprivation. Nature Neuroscience
13, 450–457.

Messaoudi, E., Kanhema, T., Soulé, J., Tiron, A., Dagyte, G., da Silva, B., and Bramham, C.R. (2007).
Sustained Arc/Arg3.1 synthesis controls long-term potentiation consolidation through regulation of local
actin polymerization in the dentate gyrus in vivo. The Journal of Neuroscience : The Official Journal of
the Society for Neuroscience *27*, 10445–10455.

Minatohara, K., Akiyoshi, M., and Okuno, H. (2015). Role of Immediate-Early Genes in Synaptic Plasticity
and Neuronal Ensembles Underlying the Memory Trace. Frontiers in Molecular Neuroscience *8*, 78.

Mo, J., Kim, C.-H., Lee, D., Sun, W., Lee, H.W., and Kim, H. (2015). Early growth response 1 (Egr-1)
directly regulates GABA receptor α2, α4, and θ subunits in the hippocampus. Journal of Neurochemistry
133, 489–500.

Morgan, J.I., Cohen, D.R., Hempstead, J.L., and Curran, T. (1987). Mapping patterns of c-fos expression in
the central nervous system after seizure. Science *237*, 192–197.

Okuno, H., Akashi, K., Ishii, Y., Yagishita-Kyo, N., Suzuki, K., Nonaka, M., Kawashima, T., Fujii, H.,
Takemoto-Kimura, S., Abe, M., et al. (2012). Inverse synaptic tagging of inactive synapses via dynamic
interaction of Arc/Arg3.1 with CaMKIIβ. Cell *149*, 886–898.

Paxinos, G., and Franklin, K.B.J. (2013). Paxinos and Franklin's the mouse brain in stereotaxic coordinates
(Academic Press).

Petreanu, L., Mao, T., Sternson, S.M., and Svoboda, K. (2009). The subcellular organization of neocortical
excitatory connections. Nature *457*, 1142–1145.

Ploski, J.E., Pierre, V.J., Smucny, J., Park, K., Monsey, M.S., Overeem, K.A., and Schafe, G.E. (2008). The
activity-regulated cytoskeletal-associated protein (Arc/Arg3.1) is required for memory consolidation of
pavlovian fear conditioning in the lateral amygdala. Journal of Neuroscience *28*, 12383–12395.

Ramirez, S., Liu, X., Lin, P.-A., Suh, J., Pignatelli, M., Redondo, R.L., Ryan, T.J., and Tonegawa, S. (2013).
Creating a false memory in the hippocampus. Science (New York, N.Y.) *341*, 387–391.

Ramírez-Amaya, V., Vazdarjanova, A., Mikhael, D., Rosi, S., Worley, P.F., and Barnes, C.A. (2005). Spatial
Exploration-Induced Arc mRNA and Protein Expression: Evidence for Selective, Network-Specific
Reactivation. The Journal of Neuroscience *25*, 1761–1768.

Reijmers, L.G., Perkins, B.L., Matsuo, N., and Mayford, M. (2007). Localization of a stable neural
correlate of associative memory. Science (New York, N.Y.) *317*, 1230–1233.

Rial Verde, E.M., Lee-Osbourne, J., Worley, P.F., Malinow, R., and Cline, H.T. (2006). Increased
Expression of the Immediate-Early Gene Arc/Arg3.1 Reduces AMPA Receptor-Mediated Synaptic
Transmission. Neuron *52*, 461–474.

Rosen, K.M., McCormack, M.A., Villa-Komaroff, L., and Mower, G.D. (1992). Brief visual experience
induces immediate early gene expression in the cat visual cortex. Proceedings of the National Academy
of Sciences of the United States of America *89*, 5437–5441.

Saleem, A.B., Ayaz, A., Jeffery, K.J., Harris, K.D., and Carandini, M. (2013). Integration of visual motion
and locomotion in mouse visual cortex. Nature Neuroscience *16*, 1864–1869.

Shepherd, J.D., and Bear, M.F. (2011). New views of Arc, a master regulator of synaptic plasticity. Nature
Neuroscience *14*, 279–284.

Shepherd, J.D., Rumbaugh, G., Wu, J., Chowdhury, S., Plath, N., Kuhl, D., Huganir, R.L., and Worley, P.F.
(2006). Arc/Arg3.1 Mediates Homeostatic Synaptic Scaling of AMPA Receptors. Neuron *52*, 475–484.

Steward, O., Matsudaira Yee, K., Farris, S., Pirbhoy, P.S., Worley, P., Okamura, K., Okuno, H., and Bito, H.
(2017). Delayed Degradation and Impaired Dendritic Delivery of Intron-Lacking EGFP-Arc/Arg3.1 mRNA
in EGFP-Arc Transgenic Mice. Frontiers in Molecular Neuroscience *10*, 435.

Tagawa, Y., Kanold, P.O., Majdan, M., and Shatz, C.J. (2005). Multiple periods of functional ocular
dominance plasticity in mouse visual cortex. Nature Neuroscience *8*, 380–388.

Torrado Pacheco, A., Tilden, E.I., Grutzner, S.M., Lane, B.J., Wu, Y., Hengen, K.B., Gjorgjieva, J., and
Turrigiano, G.G. (2019). Rapid and active stabilization of visual cortical firing rates across light-dark
transitions. Proc Natl Acad Sci U S A *116*, 18068–18077.

Tzingounis, A. V., and Nicoll, R.A. (2006). Arc/Arg3.1: Linking Gene Expression to Synaptic Plasticity and
Memory. Neuron *52*, 403–407.

Vazdarjanova, A., Ramirez-Amaya, V., Insel, N., Plummer, T.K., Rosi, S., Chowdhury, S., Mikhael, D.,
Worley, P.F., Guzowski, J.F., and Barnes, C.A. (2006). Spatial exploration induces ARC, a plasticity-related
immediate-early gene, only in calcium/calmodulin-dependent protein kinase II-positive principal
excitatory and inhibitory neurons of the rat forebrain. The Journal of Comparative Neurology *498*, 317–
329.

Veyrac, A., Besnard, A., Caboche, J., Davis, S., and Laroche, S. (2014). The Transcription Factor
Zif268/Egr1, Brain Plasticity, and Memory. Progress in Molecular Biology and Translational Science *122*,
89–129.

Wang, G., Xie, H., Hu, Y., Chen, Q., Liu, C., Liu, K., Yan, Y., and Guan, J.-S. (2021). Egr1-EGFP transgenic
mouse allows in vivo recording of Egr1 expression and neural activity. J Neurosci Methods *363*, 109350.

Wang, K.H., Majewska, A., Schummers, J., Farley, B., Hu, C., Sur, M., and Tonegawa, S. (2006). In vivo
two-photon imaging reveals a role of arc in enhancing orientation specificity in visual cortex. Cell *126*,
389–402.

Waung, M.W., Pfeiffer, B.E., Nosyreva, E.D., Ronesi, J.A., and Huber, K.M. (2008). Rapid Translation of
Arc/Arg3.1 Selectively Mediates mGluR-Dependent LTD through Persistent Increases in AMPAR
Endocytosis Rate. Neuron *59*, 84–97.

Weng, F.-J., Garcia, R.I., Lutzu, S., Alviña, K., Zhang, Y., Dushko, M., Ku, T., Zemoura, K., Rich, D., Garcia-
Dominguez, D., et al. (2018). Npas4 Is a Critical Regulator of Learning-Induced Plasticity at Mossy Fiber-
CA3 Synapses during Contextual Memory Formation. Neuron.

Widmer, F.C., and Keller, G.B. (2021). Developmental plasticity in visual cortex is necessary for normal
visuomotor integration and visuomotor skill learning. BioRxiv 2021.06.20.449148.

Xie, H., Liu, Y., Zhu, Y., Ding, X., Yang, Y., and Guan, J.-S. (2014). In vivo imaging of immediate early gene
expression reveals layer-specific memory traces in the mammalian brain. Proceedings of the National
Academy of Sciences *111*, 2788–2793.

Yamada, Y., Hada, Y., Imamura, K., Mataga, N., Watanabe, Y., and Yamamoto, M. (1999). Differential
expression of immediate-early genes, c-fos and zif268, in the visual cortex of young rats: effects of a
noradrenergic neurotoxin on their expression. Neuroscience *92*, 473–484.

Yap, E.-L., and Greenberg, M.E. (2018). Activity-Regulated Transcription: Bridging the Gap between
Neural Activity and Behavior. Neuron *100*, 330–348.

Yasoshima, Y., Sako, N., Senba, E., and Yamamoto, T. (2006). Acute suppression, but not chronic genetic
deficiency, of c-fos gene expression impairs long-term memory in aversive taste learning. Proceedings of
the National Academy of Sciences of the United States of America *103*, 7106–7111.

Yassin, L., Benedetti, B.L., Jouhanneau, J.S., Wen, J.A., Poulet, J.F.A., and Barth, A.L. (2010). An
Embedded Subnetwork of Highly Active Neurons in the Neocortex. Neuron *68*, 1043–1050.

Zmarz, P., and Keller, G.B. (2016). Mismatch Receptive Fields in Mouse Visual Cortex. Neuron *92*, 766–
772.