Host specialization in microparasites transmitted by generalist vectors: insights into the

cellular and immunological mechanisms

Yi-Pin Lin[1,2#]*, Danielle M. Tufts[3#], Alan P. Dupuis, II[1], Matthew Combs[3], Ashley L.

Marcinkiewicz[1], Andrew D. Hirsbrunner[1], Alexander J. Diaz[1], Jessica L. Stout[1], Anna M. Blom[4],

Klemen Strle[1], April D. Davis[1], Laura D. Kramer[1,2], Maria A. Diuk-Wasser[3]*

[1]Division of Infectious Diseases, Wadsworth Center, New York State Department of Health,

Albany, NY, USA, [2]Department of Biomedical Sciences, State University of New York at

Albany, NY, USA, [3]Department of Ecology, Evolution, and Environmental Biology, Columbia

University, New York, NY USA, [4]Division of Medical Protein Chemistry, Department of

Translational Medicine, Lund University, Malmo, Sweden

Short title: Early defense and pathogen adhesiveness confer spirochete host specialization

Key words: Lyme disease, *Borrelia*, Early immune responses, Adhesion

[#]Contributed equally to this work

*correspondence: Yi-Pin Lin, Ph.D.
                 Division of Infectious Disease,
                 Wadsworth Center, New York State Department of Health
                 120 New Scotland Ave., Albany, NY 12047
                 Telephone: 518-402-2233; Fax: 518-473-1326
                 Email: Yi-Pin.Lin@health.ny.gov

*correspondence: Maria A. Diuk-Wasser, Ph.D.
                 Department of Ecology, Evolution, and Environmental Biology,
                 Columbia University
                 1200 Amsterdam Ave., New York, NY 10027
                 Telephone: 212-854-3355
                 Email: mad2256@columbia.edu

ABSTRACT (289 words)

Host specialization is an ecological and evolutionary process by which a pathogen becomes differentially adapted to a subset of hosts, restricting its host range. For parasites transmitted by generalist vectors, host specialization is not expected to evolve because of the decreased survival of those parasites in inadequate hosts. Thus, parasites may develop adaptation strategies, resulting in host specialization. The causative agents of Lyme disease are multiple species of bacteria, *Borrelia burgdorferi* sensu lato species complex (*Bb*sl), and are suitable for examining host specialization as birds and rodents were found to carry different species of these bacteria. Debate exists on whether host specialization occurs among these strains within a particular species of *Bb*sl, such as *B. burgdorferi* sensu stricto (*Bb*ss). Current evidence supports some *Bb*ss strains are widespread in white-footed mice but others are in non-rodent vertebrates, such as birds. To recapitulate specialization in the laboratory and define the mechanisms for host specialization, we introduced different genotypes of *Bb*ss via tick transmission to American robins and white-footed mice, the Lyme disease reservoirs in North America. Among these strains, we found distinct levels of spirochete presence in the bloodstream and tissues and maintenance by these animals in a host-dependent fashion. We showed that the late stage persistence of these strains largely corresponds to bacterial survival at early infection onsets. We also demonstrated that those early survival phenotypes correspond to spirochete adhesiveness, evasion of complement-mediated killing in sera, and/or not triggering high levels of pro-inflammatory cytokines and antibodies. Our findings thus link host competence to *Bb*ss with spirochete genotypic variation of adhesiveness and inducing/escaping host immune responses, illuminating the potential mechanisms that dictate host specialization. Such information will provide a foundation for further investigation into multi-disciplinary processes driving host specialization of microparasites.

AUTHOR SUMMARY (200 words)

Host specialization arises when microparasites adapt to a subset of available hosts, restricting the host ranges they can infect. The mechanisms and selective pressures for the evolution of host specialization remain unclear. The causative agent of Lyme disease (LD), the bacteria species complex of *Borrelia burgdorferi* sensu lato, is adapted to different vertebrates. However, whether such a differential host adaption also applies to each genotype within the same species is under debate. Further, the mechanisms that drive such host specialization are unclear. We thus introduced three genotypes of one LD bacteria species (*B. burgdorferi* sensu stricto) individually via tick bite to American robins and white-footed mice, the most common LD reservoirs in North America. We found that these genotypes differed in the persistent maintenance by those reservoirs and occurred in a host-specific fashion. The ability of those bacteria for long-term maintenance was linked with their capability to attach to cells and a lack of induction of high levels of immune responses at early infection onsets. This work demonstrates the potential mechanisms that dictate host specialization of LD bacteria circulating in natural populations. Such information will pave the road to define the molecular, ecological, and evolutionary determinants that drive host-microparasite interactions.

INTRODUCTION

The range of hosts a parasite can infect is arguably one of the most important properties of a parasite because it can determine, among other things, whether a parasite can survive the extinction of a host species and whether it can become established and spread following its introduction to a new area (1, 2). Host specialization is defined as the ecological and evolutionary process by which a pathogen becomes differentially adapted and thus restricts its host range to a subset of potential hosts. For vector-borne microparasites, such a process is expected to evolve when vectors are specialized to competent hosts, leading to microparasite amplification. However, it is much less obvious how and when host specialization in a vector-borne microparasite can occur and when vectors are host-indiscriminate. Given the significant cost incurred when microparasites are inoculated into incompetent hosts, the evolution of host specialization is only expected to occur when it provides a significant selective advantage for the microparasite to overcome cellular or immunological barriers to infection.

The spirochetes of the *B. burgdorferi* sensu lato (s.l.) species complex, agents of Lyme disease, represent an ideal system to investigate the tradeoffs involved in the evolution of host specialization. *B. burgdorferi* s.l. is maintained in an enzootic cycle between generalist ticks of the *Ixodes ricinus* complex and reservoir hosts, including small and medium-sized mammals, birds, and reptiles (3-5). Despite being transmitted by the same generalist vector, some *B. burgdorferi* s.l. genospecies in Europe are almost exclusively associated with a host taxa (e.g. *B. afzelii* in rodents; *B. garinii* in avian hosts). In contrast, *B. burgdorferi* sensu stricto (hereafter *B. burgdorferi*) in North America is considered a host generalist as it has been isolated from multiple classes of vertebrate animals including mammalian and avian hosts (3, 6-8). However, laboratory infection studies indicate that the fitness of *B. burgdorferi* strains varies in different hosts (9, 10).

In natural populations, weak associations were also found between hosts and particular *B.*

burgdorferi genotypes, defined by polymorphic markers, such as *ospC* or 16S-23S rRNA

intergenic spacer type (RST) (11-13). An intriguing possibility is that the partially and regionally

constrained host associations observed in *B. burgdorferi* genotypes represent an incipient

evolutionary process of host specialization (4, 12, 14-17). That is, *B. burgdorferi* may be on an

evolutionary path to diversify into host specialized genospecies, similar to those within the *B.*

burgdorferi s.l. species complex in Europe. Nonetheless, the molecular processes underlying such

diversification and the extent to which they might drive genome-wide diversification in *B.*

burgdorferi remain under debate.

Multiple-niche polymorphism (MNP), or diversifying selection, has been proposed as a

balancing selection mechanism maintaining the diversity of *B. burgdorferi* genotypes through

adaptation to host 'niches' (11, 12, 18, 19). This model was applied to the maintenance of the

polymorphism in the outer surface protein C (OspC), one of the most diverse Lyme borreliae

antigens that is heavily targeted by the vertebrate immune system (20-22). Alternatively (or

concurrently), the OspC polymorphism could be maintained in a host-independent manner,

through negative frequency-dependent selection. In this form of balancing selection, rare *B.*

burgdorferi OspC genotypes to which few hosts have been exposed would have a selective

advantage over the more common genotypes to which many hosts have mounted an adaptive

immune response; thus maintaining diversity in the population (14, 17, 23-25). Theoretical studies

have examined patterns of OspC diversity predicted by the different proposed eco-evolutionary

mechanisms in different ecological settings (25-28), but a mechanistic understanding of the cellular

and immunological mechanisms mediating strain-host interactions is critical to disentangle the

multiple co-occurring selective pressures.

In this study, we simultaneously assessed cellular and immunological mechanisms that mediate strain-host interactions using three *B. burgdorferi* strains with variable *ospC* genotypes, as well as the North American rodent and avian reservoir hosts, American robins (hereafter robins) and white-footed mice (*Peromyscus leucopus*), respectively. We aimed to determine which cellular and immunological mechanisms mediate fitness variation in strain-host species pairs. Specifically, we identified genotypic differences across *B. burgdorferi* interacting with hosts in (1) the role of complement inhibition and inflammation induction; (2) the role of bacterial adhesion; (3) the role of antibodies in driving late stage persistence in specific hosts and (4) transmission efficiency to larval ticks. By characterizing these mechanisms, we aimed to identify plausible selective pressures that shape the composition of *B. burgdorferi* strain community, its diversity, and population dynamics.

RESULTS

***Borrelia burgdorferi* B31-5A4, 297, and cN40 differed in their adhesiveness to fibroblasts, *ex vivo* cytokine induction, and infection establishment in robins and white-footed mice.** To compare the capability of genotypically distinct *B. burgdorferi* to initiate infection in reservoir hosts, the cloned *B. burgdorferi* strains B31-5A4, 297, and cN40 belonging to different genotypes (Table S1) and displaying similar growth rates *in vitro* (Fig. S1) were used in this study. Bacterial burdens at inoculation sites of skin and in blood were determined at 1 day after each of these strains was intradermally introduced into robins and white-footed mice. We did not detect any of these strains in robin blood (Fig. S2A), in agreement with no hematogenous dissemination at such an early time point (29). However, we found that B31-5A4 and cN40 showed significantly higher spirochete burdens at the initial injection sites, compared to mock-infected robins (Fig. 1). The

injection sites from only two out of four 297-inoculated robins had spirochete burdens higher than

detection limits (10 bacteria per 100ng of DNA from tissues), leading to non-significant

differences in the levels of colonization from mock-infected robins (Fig. 1A). In white-footed mice

infections, we did not observe spirochete burdens higher than detection limits of any of these

strains in the blood of white-footed mice (Fig. S2B). Nonetheless, the burdens of B31-5A4 and

297 were significantly higher than those from mock-infected individuals at the inoculation site

($\sim10^2$ bacteria per 100ng of DNA). Only two of five white-footed mice had cN40 bacterial loads

greater than the detectable threshold limit, resulting in non-significant differences in burdens from

mock-infected mice (Fig. 1B). These findings indicate B31-5A4, 297, and cN40 have the capacity

to establish infection selectively in robins and white-footed mice.

As the burdens of *B. burgdorferi* during infection initiation can be attributed to differences in

spirochete attachment to cells or clearance by host inflammatory responses, we incubated B31-

5A4, 297, and cN40 with fibroblasts isolated from robin or white-footed mouse for 1 h and

determined the levels of bacterial attachment using fluorescent microscopy (Fig. S3). A non-

adhesive *B. burgdorferi* strain B314 was also included as control (Table S1). Compared to B314,

we found that B31-5A4, 297, and cN40 more efficiently bind to the fibroblasts from robins (Fig.

1C) or white-footed mice (Fig. 1D). Whereas B31-5A4 and cN40 bound to robin fibroblasts at

significantly higher levels compared to 297 (Fig. 1C), B31-5A4 and 297 attached to white-footed

mouse fibroblasts at significantly greater levels compared to cN40 (Fig. 1D). To determine

whether B31-5A4, 297, and cN40 trigger differential pro-inflammatory responses in fibroblasts

from different reservoir origins, we measured the expression levels of the genes encoding pro-

inflammatory cytokines, IFNγ and TNF, or the proteins related to these cytokines after incubating

cells with each of the spirochete strains for 24h. The expression levels of housekeeping genes (18S

rRNA and β-actin) were also included as controls. We found that the levels of expression for each

of the genes in any spirochete-treated cells were not significantly different from the mock-treated

cells at 1 to 10 of the spirochetes to cell ratio (Fig. S4). At 1 to 100 of spirochete to cell ratio, the

B31-5A4- or cN40-treated robin fibroblasts expressed similar levels of the tested cytokines to the

cells under mock treatment while 297-incubated cells had significantly greater expression of IFNγ

and TNF-induced protein than mock-treated cells, respectively (Fig. 1F and G). Conversely, while

white-footed mouse fibroblasts incubated with B31-5A4 or 297 expressed those cytokines

indistinguishably from mock-treated cells, cN40-treated cells showed significantly higher

expression of IFNγ and TNF (4.6- and 30.1-fold, respectively) than the cells under mock treatment

(Fig. 1I and J). These results suggest that *B. burgdorferi* B31-5A4, 297, and cN40 differ in their

fibroblast adhesiveness and cytokine induction in a host-dependent fashion.

Ticks acquired different levels of *B. burgdorferi* B31-5A4, 297, and cN40 from robins and

white-footed mice infected with each of these spirochete strains. To compare reservoir host

competence to B31-5A4, 297, and cN40, we assessed spirochete transmission from host to feeding

larvae. Unfed *I. scapularis* nymphs were allowed to feed on robins or white-footed mice until

repletion. We subsequently measured bacterial loads in flat and fed nymphs and found

indistinguishable burdens among these ticks (~10^4 bacteria per nymph), indicating no differences

of the ability for any tested strain to survive in flat or fed nymphs (Fig. S5).

We then placed *I. scapularis* larvae on robins or white-footed mice at different time points (Fig.

2A) to determine spirochete burdens in each larva and the percentage of spirochete-positive larvae

per individual host (defined as percent positivity). The spirochete burdens in uninfected (control)

robin-derived larvae were below the detection limit (10 bacteria per tick), resulting in zero

spirochete positivity at 14, 28, 35, and 56 days post nymph feeding (dpf) (Fig. 2B to F and Table 1). At least 18% of larvae feeding on B31-5A4-, 297-, or cN40-infected robins were spirochete positive at all time points with greater than detection limits of bacterial loads, indicating the ability of robins to persistently maintain and transmit these strains (Fig. 2C to F and Table 1). The percent positivity of fed larvae carrying cN40 was significantly higher than those harboring 297 at all four time points and significantly greater than those carrying B31-5A4 at 28, 35, and 56 dpf (Fig. 2B and Table 1). Moreover, the cN40-infected larvae had significantly higher bacterial loads than the 297-infected larvae at all time points and significantly greater than those in B31-5A4-infected larvae at 28, 35, and 56dpf (Fig. 2C to F). These data suggest that the cN40 is maintained and transmitted more efficiently in robins than B31-5A4 and 297.

Similarly, we found bacterial burdens lower than detection limits in the larvae feeding on uninfected (control) mice, resulting in zero spirochete positivity (Fig. 2G to K, Table 1). At least 9% of the larvae feeding on B31-5A4-, 297-, or cN40-infected mice were positive throughout the experiment (Fig. 2G and Table 1). The bacterial loads from those mice were also statistically different from those derived from uninfected mice (Fig. 2H to K), except that larvae infected with cN40 had similar spirochete burdens to uninfected mouse-derived larvae at 56dpi. These data indicate the ability of white-footed mice to maintain and transmit each of these strains. We found significantly lower spirochete positivity in larvae infected with 297, compared to that in larvae infected with B31-5A4 at 28 and 35 dpf, respectively (p< 0.05, Fig. 2G, I, J and Table 1). Finally, larvae infected with cN40 showed more than significantly lower spirochete positivity (Fig. 2G and Table 1) and had significantly less bacterial burdens than B31-5A4- and 297-infected ticks throughout the experiments (Fig. 2H to K). These data suggest less capability of white-footed mice to maintain and transmit cN40 compared to B31-5A4 or 297.

Borrelia burgdorferi B31-5A4, 297, and cN40 varied in their capability to trigger bacteremia

and persistently colonize robins and white-footed mice. We next determined the ability of B31-

5A4, 297, and cN40 to survive in the bloodstream of robins and white-footed mice. We found that

robins infected with cN40 develop significantly greater levels of bacteremia at 7, 14, and 21dpf

than the uninfected robins, which had spirochete burdens lower than the detection limits (10

bacteria per 100ng DNA, Fig. 3A to C). However, the burdens of that strain in the blood was no

statistically different from those in uninfected robins at 28dpf (Fig. 3D). Robins infected with 297

had spirochete burdens in the blood no statistically different from uninfected robins at 7, 21 and

28 dpf but developed statistically greater bacterial loads at 14dpf (Fig. 3A to D). Though B31-5A4

was capable of inducing bacteremia in robins at 7 and 14dpf at the levels statistically greater than

those in uninfected robins, the burdens of this strain in the blood were not statistically different

from those in uninfected robins at 21 and 28dpf (Fig. 3A to D). These results indicate the ability

of cN40 to induce long-lasting bacteremia in robins, compared to other strains, and 297 displayed

delayed onsets of bacteremia. Similar to uninfected robins, uninfected white-footed mice did not

exhibit bacteremia above the levels of detection throughout the experiment (Fig. 3E to H). At 21

and 28dpf, the spirochetes burdens in the blood from the mice infected with any of tested strains

were no statistically different from those in uninfected mice (Fig. 3E and H). At 7 and 14 dpf, B31-

5A4 and 297 triggered significantly higher levels of bacteremia than uninfected mice (Fig. 3E and

F). However, the majority of mice infected with cN40 did not show detectable bacteremia (3/5 and

5/5 mouse blood had burdens below detection limits at 7 and 14dpf, respectively), resulting in no

significant different bacterial loads from uninfected mice (Fig. 3E and F). These data suggest that

cN40 is less capable of surviving in the white-footed mouse bloodstream compared to B31-5A4

and 297, in contrast to the results derived from robins.

We also evaluated the ability of B31-5A4, 297, and cN40 to colonize robin and white-footed

mouse tissues at 64dpf (Fig. 2A). Robins infected with each of these strains had significantly

greater spirochete burdens in skin than uninfected robins, which exhibited bacterial loads below

the detection limits (Fig. 3I). In addition, spirochetes were detected at significantly greater burdens

in the heart and brain of cN40-infected robins and the heart of robins infected with B31-5A4,

compared to respective tissues from uninfected robins (Fig. 3J and K). Majority of the 297- or

cN40-infected robins (3 out of 4 and 3 out of 5 in 297- and cN40-infected robins, respectively)

had burdens below the detection limits in livers, yielding no significantly different burdens from

those in uninfected robins (Fig. 3L). Similarly, the burdens of B31-5A4 in the livers from

spirochete-infected robins were no significantly different from those from uninfected robins (Fig.

3L). These data showed persistent skin colonization of strains B31-5A4, 297, and cN40 in robin,

but the ability of each strain to colonize other tissues varied (Fig. 3I to L). We also measured the

bacterial burdens at 64dpf in the tissues from white-footed mice infected with each of these strains.

We detected spirochetes in ears, tibiofemoral joints, heart, and bladder from mice infected with

B31-5A4 or 297 ($\sim 10^2$ to 10^3 bacteria per 100ng of DNA from tissues). The bacterial burdens in

tibiofemoral joints, heart, and bladder from B31-5A4- or 297- but not cN40-infected mice were

significantly greater than those in uninfected mice (Fig. 3M to P). These findings demonstrated

the ability of B31-5A4 and 297 but not cN40 to persistently colonize white-footed mice.

Robin but not white-footed mouse complement differentiated the ability of *B. burgdorferi*

B31-5A4, 297, and cN40 to survive in sera. To define the capability of B31-5A4, 297, and cN40

to evade complement-mediated killing, we evaluated the percent survival after incubation of these

strains individually with the sera from uninfected robins or white-footed mice by counting the

number of motile bacteria under microscopes. More than 95% of all tested strains including a

serum sensitive control strain, *B. burgdorferi* B313 (30), survived in heat inactivated robin sera,

in which the heat sensitive components such as complement were not functional (Fig. 4A and

Table S1). Less than 16% of B313 remained motile, verifying the bactericidal activity of these

robin sera (Fig. 4A). More than 85% of B31-5A4 and cN40 were alive, but only 50% of 297 was

motile after incubated with robin sera (Fig. 4A). However, all strains survived at similar levels

(greater than 80% of live spirochetes) in the sera pre-treated with OmCI, which inactivates robin

complement by abolishing the lytic activity towards gram negative bacteria (Fig. 4B) (31). These

results indicate that the 297 is more vulnerable to robin complement-mediated killing than B31-

5A4 and cN40. We also determined the ability of spirochete strains to evade killing by white-

footed mouse complement in the similar fashion. Greater than 95% of B313, B31-5A4, 297, and

cN40 survived in heat inactivated white-footed mouse sera (Fig. 4C). Whereas only 29% of B313

remained motile after treated with these sera, more than 97% of other tested strains were alive in

the untreated white-footed mouse sera (Fig. 4C). More than 93% of tested strains remained motile

in white-footed mouse sera treated with CVF, which inactivates mammal complement (Fig. 4D)

(32). These results indicate that all tested strains evade the killing by white-footed mouse

complement at indistinguishable levels.

Robins and white-footed mice generated different levels of pro-inflammatory cytokines in

response to early infection of *B. burgdorferi* B31-5A4, 297, and cN40. We compared the levels

of pro-inflammatory cytokines, IFNγ and TNFα, in the sera derived from robin- and white-footed

mice at different times during the infection. We found no significant different levels of IFNγ and

TNFα in robins prior to the infection among different infection groups (Fig. 5A and E). At 7dpf,

B31-5A4- or cN40-infected robins were not significantly different in those cytokines compared to

uninfected robins, 297-infected robins produced significantly greater levels of IFNγ and TNFα

than those uninfected birds, respectively (Fig. 5B and F). At 14dpf, the levels of these cytokines

from those robins are no different from those in uninfected robins. In contrast to robins at 7dpf,

cN40-infected mice displayed significantly higher levels of these cytokines than uninfected mice

at this time point while 297-infected mice had no significantly different in the levels of those

cytokines, compared to uninfected mice (Fig. 5J and N). Note that B31-5A4-infected mice had

significantly greater levels of IFNγ (Fig. 5J) but were not significantly different from levels of

TNFα at 7dpf (Fig. 5N), compared to uninfected mice. These data indicate differences of pro-

inflammatory cytokine induction by each of these spirochete strains, particularly at early stages of

robin and white-footed mouse infection.

Robins and white-footed mice developed distinct levels of bactericidal antibodies during

early infection of *B. burgdorferi* B31-5A4, 297, or cN40. We aimed to quantitatively measure

the levels of antibodies induced by the infection of B31-5A4, 297, and cN40. To avoid the

confounding factors of strain-specific antibody recognition observed previously (10, 33), we

mixed these strains and determined the IgG titers against such mixtures in robins infected with

B31-5A4, 297, or cN40. We found that robins infected with any strain develop significantly higher

anti-spirochete IgG titers than uninfected robins at 14, 21, and 28dpf whereas only 297-infected

robins had significantly higher tiers of IgG than uninfected robins at 7dpf (Fig. 6B to E). cN40

triggered significantly greater levels of IgG antibodies compared to the other strains at 21 and 28

dpf whereas 297 induced IgG titers significantly higher than those titers triggered by other strains

at 7dpf (Fig. 6B, D and E). Further, we found that white-footed mice infected with each of these

strains had anti-spirochete IgG titers at more robust levels than uninfected mice at 14, 21, and

28dpf whereas only cN40-infected mice developed significantly higher tiers of IgG than

uninfected white-footed mice at 7dpf (Fig. 6G to J). cN40 induced significantly lower titers of

such IgG at 21 and 28dpf but significantly higher titers at 7dpf, compared to B31-5A4 and 297

(Fig. 6G, I and J). These results indicate that B31-5A4, 297, and cN40 differ in their ability to

induce antibodies, and such differences depend on the hosts and infection onset.

The levels of antibodies against B31-5A4, 297, or cN40 at 7dpf in robins and white-footed mice

negatively corresponded to the ability of these hosts to carry spirochetes (Fig. 2B and G). This

finding raised a hypothesis that such correspondences are mediated by bactericidal activities of

those antibodies. We thus tested this hypothesis by incubating a mixture of B31-5A4, 297, and

cN40 with different dilution rates of sera from robins or white-footed mice infected with each of

these strains at 7dpf. We found that B31-5A4- and cN40-derived robin sera did not kill spirochetes

at any of the tested dilution rates, similar to sera from uninfected robins. However, 297-derived

sera eliminated 50% of spirochetes at the dilution rate of 1:52× (Fig. 6K). Conversely, B31-5A4-

and 297-derived white-footed mouse sera did not eliminate spirochetes, similar to the sera from

uninfected mice. However, cN40-derived sera eliminated 50% of spirochetes at the dilution rate

of 1:32× (Fig. 6L). These finding showed different levels of bactericidal activities from the

antibodies produced by B31-5A4-, 297-, or cN40-infected robins and white-footed mice early

during infection.

DISCUSSION

Theory predicts that generalist vectors should select for generalist pathogens, to minimize the

latter's loss to incompetent hosts. Host-specialized genospecies in *B. burgdorferi* s.l. thus represent

a paradox, with the intriguing possibility of incipient host specialization within genotypes of the

generalist *B. burgdorferi*. We present evidence of molecular mechanisms that differentially

influence the ability of three strains of *B. burgdorferi* to colonize, disseminate to distal tissues,

evade host immune responses, and be transmitted from the host to feeding ticks between two

representative natural reservoir hosts (Fig. 7). We found that cellular and immunological

mechanisms act mostly synergistically, resulting in increased fitness of strain cN40 in robins and

297 in white-footed mice. Contrary to theoretical expectations, strain B31-5A4 was able to

efficiently infect both hosts, with higher fitness in white-footed mice than the 'specialized' strain,

297, and exhibited intermediate fitness in robins. The synergistic nature of those cellular and

immunological mechanisms indicates a strong selective pressure for the evolution of host

specialization, as predicted by the MNP theory. However, the higher overall fitness of B31-5A4

does not support the existence of tradeoffs, with this strain having high fitness in both

representative hosts. This higher fitness advantage is consistent with the high overall prevalence

of the genotype of this strain (RST type 1/OspC type A) circulating in Lyme disease prevalent

regions of North America (10, 13, 34-37).

Multiple mechanisms were tested in this study for their role in contributing to host specialization

of *B. burgdorferi*. At one day after intradermal infection of spirochetes, we found that *B.*

burgdorferi B31-5A4 and cN40 colonized robin inoculation sites (skin tissue) more efficiently

than 297, whereas B31-5A4 and 297 colonized the inoculation sites of white-footed mice at

significantly higher levels than cN40. We also found that these strains differed in their capability

to attach to robin and white-footed mouse skin fibroblasts *in vitro*. These results showed that strain-

to-strain variability of fibroblast adhesion is host-dependent and corresponds to the ability of *B.*

burgdorferi spirochetes to colonize the inoculation sites of these hosts immediately after infection.

This supports prior findings of Lyme borreliae strains varying in their adhesive activities (38),

demonstrating the role of such cellular processes (adhesion) in conferring strain-host associations.

However, the levels of robin fibroblast adhesion of B31-5A4, 297, and cN40 did not perfectly

agree to strain colonization of other tissues (e.g. brain and liver) in robins at 64 days after being

fed on by *B. burgdorferi*-infected nymphs. Additionally, strain 297 colonized robin skin at similar

levels to B31 and cN40 at 64 dpf, but this strain attached to robin fibroblast at lower levels than

other strains. These disagreements between fibroblast adhesion/early colonization of inoculation

sites and tissue colonization at later stages suggest the possibility of adhesion-independent

mechanisms to determine strain-host associations. In fact, the complement system is the first line

of immune defense in vertebrate animal sera and confers Lyme borreliae clearance as soon as

infection begins (39, 40). Our findings of greater percentages of B31-5A4 and cN40 surviving in

robin sera, compared to those of 297, mirror the trends of these spirochete strains being maintained

at earlier infection onsets (e.g. 14 dpf) and early bloodstream survival of *B. burgdorferi* (e.g. 7

dpf). These results imply that the host complement plays a role in determining spirochete

genotype-specific robin competence. In contrast, all three *B. burgdorferi* strains showed similar

levels of survival in the presence of white-footed mouse sera, suggesting a non-complement-

mediated defense mechanism to dominate the association of white-footed mice with these

spirochetes.

In addition to complement, we compared the cytokine production in responding to the presence

of genotypically distinct *B. burgdorferi*. Cytokines are generally triggered shortly after pathogen

invasions, often leading to activation of different pathways and downstream bactericidal responses

as a bottleneck of infection initiation (i.e. recruitment of innate and adaptive immune cells) (41-

45). *Borrelia burgdorferi* strain-to-strain variability in inducing pro-inflammatory cytokines has

been observed in humans, *Mus musculus*, and *P. leucopus* mice or from cells derived from these

animals (46-52). In humans, the levels of multiple cytokines are positively correlated with disease

severity but are independent from spirochete burdens (46, 47, 53-55). We found that the three

tested *B. burgdorferi* strains differed in their ability, in robins and white-footed mouse fibroblasts,

to induce the expression of two pro-inflammatory cytokines and their related proteins, IFNγ, TNFα,

and TNFα-induced proteins. Such differences also matched the variation of these strains to trigger

IFNγ and TNFα during early infection. The ability of these strains to induce those pro-

inflammatory cytokines were negatively associated with larval percent positivity from the

infection of these *B. burgdorferi* strains. These results raised the hypothesis that less robust host

cytokine induction by spirochetes facilitates reservoir competence, and differences in cytokine

induction by genotypically distinct spirochetes shape Lyme borreliae-reservoir associations.

Following the induction of innate immune responses, the adaptive responses including

antibodies are essential in Lyme borreliae clearance by vertebrates (41, 56, 57). Distinct levels of

antibodies were observed during the infection time period by different spirochete strains (10, 58).

We observed at 7 dpf that 297 and cN40 triggered significantly higher titers of anti-*B. burgdorferi*

IgG in robins and white-footed mice, respectively, compared to the other spirochete strains.

Furthermore, *B. burgdorferi* was selectively eliminated by the sera collected at 7dpf from robins

infected with 297 or white-footed mice infected with cN40, suggesting that the antibody response

varies among different spirochete strain-host pairings. Interestingly, a previous study showed

variable IgG levels against lp36- and lp28-1-derived proteins in white-footed mice infected with

different *B. burgdorferi* strains (10). However, in that study, those lp36- and lp28-1-derived

proteins were produced from a single strain of *B. burgdorferi* (*B. burgdorferi* strain B31), leading

to the possibility that allelic-specific recognition is the result of antigenic sequence variation

among spirochete strains (10, 33, 59-61). Thus, the possibility of the antibodies against other

spirochete proteins to confer *B. burgdorferi* clearance may not be completely ruled out. Our

finding of early antibody-mediated bactericidal activities in robin and white-footed mouse sera

highlights the potential role of antibodies in dictating pathogen-host associations. We observed

high titers of antibody responses at late time points (21 and 28 dpf) in robins infected with cN40

and mice infected with B31-5A4- or 297. Nonetheless, cN40 was persistently maintained in robins

whereas B31-5A4 and 297 efficiently survived long-term in white-footed mice. Such observations

are consistent with the evidence of pathogen-specific antibodies present in persistently infected

animals (62-64). These results thus suggest inefficient spirochete elimination by those antibodies

at later stages of infection. That inefficient pathogen killing could be because of the onset-

dependent reduction of highly antigenic spirochete proteins (i.e. downregulation of OspC at late

infection onsets) (65-69), constantly changing features of antigen sequences to evade antibody

responses (i.e. VlsE) (70-72), or the rapid dissemination of spirochetes to the sites less vulnerable

to antibody-mediated clearance (73).

Our results suggest spirochete adhesiveness and early immune responses as the cellular and

immunological mechanisms that differentially confer spirochete fitness in reservoir-strain-host

combinations (Fig. 7). Both mechanisms could play in concert to determine *B. burgdorferi* strain-

to-strain variation in host fitness. Our findings support previous studies which determined that

some polymorphic Lyme borreliae proteins confer these immune response functions in a variant-

specific manner, and such manner is host dependent (30, 59, 62, 63, 74-80). The potential for host

specialization driving genome-wide diversification is illustrated by the recent definition of a new

species of *B. burgdorferi* s.l., *B. bavariensis*, which was previously considered a genotype of the

avian associated species, *B. garinii*. Such a redefinition is based on the variation in chromosomal

housekeeping genes and the association of this species with rodents, unlike other *B. garinii* strains

(81). This reclassification reflects how host adaptation can lead to speciation in Lyme borreliae

(82). Prior field studies along with our current laboratory evidence showing incipient host

specialization raises the possibility of diversification of *B. burgdorferi* in North America into

multiple genospecies. Here, we used a controlled laboratory setup to demonstrate differences in

fitness of *B. burgdorferi* strains in North American reservoir hosts. The molecular mechanisms

supported by our findings provide a potential model of how specific adaptations may lead to

specialization even with a significant cost in lost propagules to incompetent hosts. The identified

mechanisms can guide future empirical and modeling studies to understand the role of host-

pathogen-vector interactions in shaping the microparasite host ranges and their potential for

pathogen spillover into livestock, wildlife or humans.

MATERIALS AND METHODS

Ethics statement. All experiments involving American robins (*Turdus migratorius*) and white-

footed mice (*Peromyscus leucopus*) were performed in strict accordance with all provisions of the

Animal Welfare Act, the Guide for the Care and Use of Laboratory Animals, and the PHS Policy

on Humane Care and Use of Laboratory Animals. Additionally, the protocol was approved by the

Institutional Animal Care and Use Committee (IACUC) of Wadsworth Center, New York State

Department of Health (Protocol docket number 18-412 and 19-451), and Columbia University

(Protocol number AC-AAAY2450), and the City University in New York Advanced Science

Research Center (Protocol number ASRC AUP 2016-20). To mistnet robins, personnel were

approved on scientific collecting permits USFWS Collecting Permit MB035731 and NYSDEC

Permit #1236.

Bird, Mouse, Tick, and Bacterial strains. From June-July, 2019 and July, 2020, American robins

were mist netted around the property of Griffin Laboratory of Wadsworth Center, New York State

Department of Health at Albany, NY. This site was selected because of previously reported low

tick abundance, thus minimizing the probability of previous tick exposure. Sera were collected

from 32 and 17 robins in 2019 and 2020, respectively, using BD Microtainer Capillary Blood

Collector tubes (Fisher Scientific, Hampton, NH, USA) to assess previous infection with *B.*

burgdorferi infection by the methods previously described (31). To confirm seronegative status,

robins were quarantined for two weeks at 25°C on a 12L:12D (light: dark) cycle by housing them

in cages with a wire bottom, under which a water moat was placed. If replete ticks were found

while the birds were in quarantine, quantitative PCR (qPCR) was used to determine the spirochete

burdens in those ticks (see "Quantification of *B. burgdorferi* in infected ticks, tissues, and blood

samples" for more details). The birds with spirochete positive ticks attached were removed from

the experiments. After two weeks of quarantine, robins were subjected to additional serological

examination as described above, and 40 seronegative juvenile (hatch year) robins were considered

non-infectious and used in this study.

White-footed mice were purchased from the *Peromyscus* Genetic Stock Center at the University

of South Carolina (Columbia, SC). Non-sibling mice were bred in-house at Columbia University.

Immunodeficient Fox Chase SCID mice (C.B.17 SCID) were obtained from Charles River (Boston,

MA) and used to generate infected nymphs for each *B. burgdorferi* strain as described in the

"Mouse infection experiments by ticks" section. *Ixodes scapularis* larvae were purchased from the

National Tick Research and Education Center, Oklahoma State University (Stillwater, OK). Mice

and birds were housed individually and maintained at 21 to 24°C on a 14L:10D (light: dark) cycle

and handled humanely. The *B. burgdorferi* strains used in this study were cultivated in BSK-II

completed medium as described in Table 1 (83).

Determination of spirochete growth curves and generation time. *Borrelia burgdorferi* strains

B31-5A4, 297, and cN40 were cultivated in BSK-II complete media at 33°C in the initial

concentration of 5×10^6 ml^{-1}. The concentration of spirochetes was measured prior to incubation

and at 24-, 48-, 72-, 96-, 120-, 144-, and 168-h post incubation using a Nikon Eclipse E600 dark

field microscope (Nikon, Melville, NY). The generation time of each spirochete strain at the

exponential phase was calculated as described previously (84).

Robin, C.B.17 SCID mouse, and white-footed mouse infection by intradermal inoculation.

Four to eight week old male or female white-footed mice, American robins, or C.B.17 SCID mice

were intradermally inoculated, using a 27-gauge needle, with *B. burgdorferi* B31-5A4, 297, cN40,

or BSK-II medium without rabbit sera (1×10^5 bacteria per C.B.17 SCID mouse or 1×10^4 bacteria

per robin or white-footed mouse) as a control (63). The plasmid profiles and the presence of the

shuttle plasmids of each of these *B. burgdorferi* strains were verified prior to infection to ensure

no loss of plasmids, as described previously (85-87). Both robins and white-footed mice were

euthanized at one day post injection. The inoculation site of the skin and blood from robins and

white-footed mice were collected to quantitatively evaluate spirochete burdens as described in the

"Quantification of *B. burgdorferi* in infected ticks, tissues and blood samples" section.

Quantification of *B. burgdorferi* in infected ticks, tissues, and blood samples. The white-footed

mouse- or robin-derived replete nymphs were mixed with glass beads and homogenized by a

Precellys 24 High-Powered Bead Mill Homogenizer (Bertin, Rockville, MD). DNA was extracted

from blood and tissue samples of white-footed mice, robins and homogenized ticks using an EZ-

10 Genomic DNA kit (Biobasic, Amherst, NY). The quantity and quality of DNA for each tissue

sample was assessed by measuring the concentration of DNA and the ratio of the UV absorption

at 280 to 260 using a nanodrop 1000 UV/Vis spectrophotometer (ThermoFisher Scientific,

Waltham, MA). The amount of DNA used in this study was 100 ng for each sample, and the

280:260 ratio was between 1.75 to 1.85, indicating the lack of contamination by RNA or proteins.

qPCR was performed to quantitate bacterial loads. *Borrelia burgdorferi* genomic equivalents were

calculated using an Applied Biosystems 7500 Real-Time PCR system (ThermoFisher Scientific)

in conjunction with PowerUp SYBR Green Master Mix (ThermoFisher Scientific), based on

amplification of the *B. burgdorferi 16S rRNA* gene using primers 16S rRNAfp and 16S rRNArp

(Table S2), as described previously (88). Cycling parameters for SYBR green-based reactions

were 50°C for 2 min, 95°C for 10 min, 45 cycles of 95°C for 15 s, and 60°C for 1 min. The number

of *16S rRNA* copies was calculated by establishing a threshold cycle (Ct) standard curve of a

known number of *16S rRNA* gene copies extracted from *B. burgdorferi* strain B31, and compared

to the Ct values of the experimental samples. To ensure low signals were not simply a function of

the presence of PCR inhibitors in the DNA preparation, five samples of blood, tibiofemoral joints,

bladders of white-footed mice and the skin, brain, and heart of robins in the B31-5A4 experimental

group were applied to qPCR to determine the levels of β-actin from white-footed mice (pActfp

and pActrp) and robins (rActfp and rActrp), respectively (Table S2) (63, 89). Note that the primers

used to determine the expression of the genes encoding robin actin were based on the mRNA

sequences that were translated to protein (Genbank Accession number: PYHW01009720.1) from

a closely related avian host, rufous-bellied thrush (*Turdus rufiventris*) due to the lack of sequence

information of American robin actin. As predicted, we detected 10^7 copies of the actin gene from

100ng of each DNA sample in robins and white-footed mice, ruling out the presence of PCR

inhibitors in these samples.

Isolation of robin fibroblasts. The procedures to isolate robin fibroblasts were described

previously (90). Five 5mm × 5mm sections of skin were removed from the breast of euthanized

robins, washed twice in PBS buffer, and then incubated in a transferring solution until the next

step was ready to be performed. The constituents of the transferring solution included Dulbecco's

modified Eagle medium (DMEM) (Wadsworth media & tissue culture core) with glucose (4.5mg

ml^{-1}) (Sigma-Aldrich, St. Louis, MO), sodium pyruvate (110mg l^{-1}) (Sigma-Aldrich), L-glutamine

(ThermoFisher Scientific), supplemented with 10% heat-inactivated fetal bovine serum

(ThermoFisher Scientific), 2% heat-inactivated chicken serum (Biowest, Riverside, MO, USA),

and antibiotics (100U ml^{-1} of mixture of penicillin and streptomycin) (ThermoFisher Scientific).

The skin was then submerged in 70% ethanol (Sigma-Aldrich) for 30s, minced using sterile

scalpels, and placed in collagenase B at 37°C overnight (ThermoFisher Scientific). We then

applied the mixture of the cells to a 20μm cell strainer and spun down the samples. The pellets

containing fibroblasts were re-suspended in growth media and incubated at 37°C with 5% CO_2.

Growth media had the same components as the transferring solution except it was supplemented

with amphotericin B (ThermoFisher Scientific) to a concentration of 0.25μg/ml. The cells were

harvested by trypsinization using Trypsin (0.25% trypsin in DMEM media, ThermoFisher

Scientific) and used in the spirochete attachment experiment.

Determination of the levels of spirochete attachment to robin and white-footed mouse

fibroblasts. The primary fibroblasts from the neck skin of American robins (see the previous

section) and ears of white-footed mice were acquired commercially (#AG22353, Coriell Institute

for Medical Research, Camden, NJ) and cultivated on cover slips in 24-well plates (2×10^5 cells

per well). *Borrelia burgdorferi* strains B31-5A4, 297, cN40, or B314 were suspended in BSK-II

medium and added to prepared plates (2×10^6 spirochetes per well). The plates were centrifuged

at 106 g for 5 min and then rocked at room temperature for 1 h. After removing unbound bacteria

through washing each well with PBS containing 0.2% BSA, the bound bacteria and cells on the

cover slips were fixed using 100% chilled methanol for 1 h followed by blocking with PBS

containing 0.2% BSA for 1 h. After washing with PBS containing 0.2% BSA, the cover slips were

incubated with a fluorescein isothiocyanate (FITC)-conjugated goat anti-*B. burgdorferi* polyclonal

antibody (Abcam, Cambridge, MA) for 1 h and mounted using ProLong Gold antifade mountant

with DAPI fluorescent stain (ThermoFisher Scientific). The spirochetes (green) and the DNA from

spirochetes and the nuclei of fibroblasts (blue) were then visualized under overlaid FITC and DAPI

filters using an Olympus BX51 fluorescence microscope (Olympus Corporation, Waltham, MA)

(Fig. S3). The number of spirochetes from three fields of view were counted in four independent

events. The results were presented as the average number of spirochetes per 50 fibroblast cells.

Determination of the relative expression levels of the genes encoding IFNγ and TNFα or

TNFα-induced proteins in fibroblasts by quantitative reverse transcription PCR (RT-qPCR).

The procedures for the examination of expression levels in the genes encoding cytokines of

fibroblasts were described previously (52). In brief, the primary fibroblasts from robins and white-

footed mice were cultivated in 24-well plates (2×10^5 cell per well). When the cells were greater

than 80% confluent, *B. burgdorferi* strains B31-5A4, 297, or cN40 suspended in BSK-II medium

were added to corresponding wells on each plate (2×10^6 and 2×10^7 spirochetes per well for the

cell to spirochete ratio (MOI) of 1:10 and 1:100, respectively) for 24h. The cells incubated with

BSK-II medium without bacteria (mock-treated cells) were included as a control.

After incubation, the supernatant was removed, and the cells were washed with PBS buffer.

These fibroblasts were then suspended in Trizol (ThermoFisher Scientific) at room temperature

for 1 h to inactivate RNase. The procedure of RNA extraction was performed using Direct-Zol

RNA MiniPrep Plus Kit (Zymo Research, Irvine, CA) as previously described (84), and DNA was

removed using RQ1 RNase-Free DNase (Promega, Madison, WI). We then synthesized cDNA

using these RNA samples (1 µg per sample) by qScript cDNA SuperMix (Quanta Bioscience,

Beverly, MA). The expression levels of the house keeping genes encoding the 18S rRNA gene

from robins (Genbank Accession number: M59402.1) or white-footed mice (Genbank Accession

number: AY591913.1) were included as controls. The primers used to quantitate the expression of

the genes encoding white-footed mouse IFNγ, TNF, and 18S rRNA, and robin IFNγ, TNFα-

induced proteins, and 18S rRNA are listed in Table S2 (89). Note, because of the lack of sequence

information for robin cytokines, the primers to determine the expression of these cytokines were

based on the mRNA sequences of IFNγ (Genbank Accession number: PYHW01010552.1) and

TNFα-induced proteins (Genbank Accession number: PYHW01009717.1) from rufous-bellied

thrush (*Turdus rufiventris*). The quantity and quality of cDNA for each sample was evaluated by

obtaining the concentration of DNA and the ratio of the UV absorption at 260 and 280 using a

Nanodrop 1000 UV/Vis spectrophotometer (ThermoFisher, Waltham, MA). The resulting ratio

between 1.75 and 1.85, indicated the lack of RNA or protein contamination. Samples were applied

to an Applied Biosystems 7500 Real-Time PCR System (ThermoFisher) in conjunction with

PowerUp SYBR Green Master Mix (ThermoFisher) to detect the expression levels of the above-

mentioned genes. The cycling parameters were 50°C for 2 min, 95°C for 10 min, and 45 cycles of

95°C for 15 s, and 49°C for 1 min, and the resulting values of threshold cycles (Ct) were

determined from duplicate experiments in three independent events. The relative expression of the

genes encoding IFNγ, TNF, TNFα-induced proteins, or 18S rRNA was presented by normalizing

the Ct-derived from each of these cytokines to that of β-actin from respective animals through the

following equation (Equation 1).

$$\textit{cytokine expression relative to actin} = 2^{-(Ct(cytokines)-Ct(actin))} \text{ (Equation 1)}$$

Generation of *B. burgdorferi*-infected nymphal ticks. Four-week-old male and female C.B.17

SCID mice were injected with a concentration of 1×10^5 of either *B. burgdorferi* strain B31-5A4,

297, or cN40 via subcutaneous injection. The plasmid profiles and the presence of the shuttle

vector of each of these *B. burgdorferi* strains were verified prior to injection to ensure no loss of

plasmids, as previously described (85-87). SCID mice inoculated with BSK-II media without

rabbit sera were used to generate uninfected nymphs. A 3 mm ear biopsy was collected 7 days post

injection from each mouse inoculated with a *B. burgdorferi* strain and DNA was extracted using a

DNeasy Blood and Tissue kit (Qiagen, Germantown, MD). DNA from ear tissue was subjected to

qPCR analysis to verify infection in each host (30). At 14 days post injection, approximately 200

larvae were placed in the ears of anesthetized mice using a paint brush. Mice were anesthetized

with isoflurane for 45 min to allow larvae to attach before being housed individually in water bath

cages, larvae were allowed to feed to repletion, as described previously (30, 91). Engorged larvae

were collected, cleaned with 10% bleach and 70% ethanol solutions, and stored in an incubator at

21°C, 95% relative humidity, and a 14L:10D light: dark cycle. Larvae molted into the nymphal

life stage in approximately 4-6 weeks and were used in the subsequent experiments with white-

footed mice and robins.

Robin and white-footed mouse infection by nymphs. The timeline of experimental procedures

is provided in Fig. 2A. Basically, unfed nymphs carrying B31-5A4, 297, cN40 or unfed, uninfected

nymphs were placed in the ear canals of each mouse ($n = 5$ mice per group, 5 nymphs in each ear).

Mice were maintained under anesthesia for 60 min to allow nymphs to attach before being placed

individually in water bath cages. Approximately 100 to 150 xenodiagnostic larvae were placed on

each mouse at 14, 28, 35, and 56 days post nymph feeding using the aforementioned procedure.

To permit flat nymphs to feed on robins, the birds were placed into a PVC pipe (~2.5-3.0 inches

in diameter; 10 inches in length) as described (92). After unfed nymphs carrying B31-5A4, 297,

cN40 or unfed, uninfected nymphs were placed on these robins ($n = 5$ robins per group, 10 nymphs

per bird), the PVC pipes were covered by a fine mosquito net ('no-see-um' netting, Skeeta,

Bradenton, FL) secured with a rubber band. The birds were then moved into wire bottom cages

with a water moat. These tick-infested robins were kept in a dark room for 1 h to minimize

grooming and allow ticks to attach before they were released and returned to their cages. To allow

xenodiagnostic larvae to feed on robins, approximately 100 to 200 naïve larvae were placed on

robins restrained in the PVC pipes as described above at 14, 28, 35, and 56 days post nymph

feeding.

Blood samples were collected from each animal (robins and white-footed mice) prior to nymph

feeding and at 7, 14, 21, and 28 days post nymph feeding. The skin, heart, liver, and brain from

robins and the ears, tibiofemoral joints, heart, and bladder from white-footed mice were obtained

at 64 days post nymphs feeding. The blood, tissues, and replete nymphs collected from each animal

were analyzed via qPCR to determine spirochete burdens as described in the section

"Quantification of *B. burgdorferi* in infected ticks, tissues, and blood samples".

Serum resistance assays. Serum resistance of *B. burgdorferi* strains was determined as described

previously with modifications (30, 84). The mid-log phase of *B. burgdorferi* strains B31-5A4, 297,

and cN40 as well as a high passage, serum sensitive strain B313 (negative control) were cultivated

in triplicate. The resulting spirochete culture was diluted to a final concentration of 5×10^6 bacteria

per milliliter into BSK-II medium without rabbit sera. The cell suspensions were mixed with sera

collected from naïve white-footed mice or robins (60% spirochetes and 40% sera) in the presence

or absence of 2 µM of cobra venom factor (CVF) or recombinant *Ornithordorus moubata*

complement inhibitor (OmCI). The generation of recombinant OmCI has been described

previously (31). The sera were incubated at 65°C for 2 h (heat-inactivated sera) and included as a

control. At 0 and 4 h following incubation with sera, the number of motile spirochetes was

measured under dark field microscopy. The percent survival of *B. burgdorferi* was calculated by

the normalization of motile spirochetes at 4 h post incubation to that immediately after incubation

with sera.

Determination of the relative levels of IFNγ and TNFα in sera by ELISA. Due to the lack of

commercially available ELISA kits to detect robin cytokines, the kits for chicken IFNγ

(ThermoFisher) and TNFα (Genorise, Glen Mills, PA) were used to measure the levels of those

cytokines in robins based on the absorption values derived from ELISA. Similarly, the ELISA kits

to determine the levels of IFNγ and TNFα from house mouse (*Mus muscuslus*) (Tonbo Bioscience,

San Diego, CA) were utilized to detect those cytokines in white-footed mice. We observed the

antibodies in the ELISA kits cross reacted with IFNγ and TNFα from robins and white-footed mice.

However, the recombinant proteins provided in these kits to generate standard curves were based

on the sequences from chicken and house mouse IFNγ and TNFα, potentially leading to inaccuracy

of quantification for those cytokines by normalizing the results obtained from robin and white-

footed mouse samples to standard curves. Therefore, we chose to present reservoir animal

cytokines as relative levels (arbitrary unit; A.U.), similar to a previous study (93). In brief, the

monoclonal capture antibodies that recognized chicken or house mouse IFNγ or TNFα were coated

on microtiter plate wells. After blocked by 5% PBS-BSA, different dilutions (1:100×, 1:300×, or

1:900×) of sera from robins and white-footed mice at 0, 7, 14, or 21 days post nymph feeding were

added to the wells. Wells were washed using PBST buffer (PBS with 0.5% of Tween 20) and the

detection of antibodies against chicken or house mouse IFNγ or TNFα were incubated for 1 h at

room temperature. The wells were then washed with PBST buffer and subsequently mixed with a

tetramethyl benzidine solution (ThermoFisher). The absorbance level was detected at 620nm for

10 cycles of 60 s kinetic intervals with 10 s of shaking duration in a Sunrise absorbance ELISA

plate reader (Tecan, Männedorf, Switzerland). We then obtained the greatest maximum slope of

optical density/min per sample multiplied by respective serum dilution factors to represent the

relative levels of cytokines shown as arbitrary units.

Determination of the titers of the antibodies against spirochetes. The titers of IgG against

spirochetes were measured as described previously with modifications (94). Basically, microtiter

plate wells were coated with a mixture of *B. burgdorferi* B31, 297, and cN40 (1×10^6 spirochetes

per strain in a well). After blocking with 5% PBS-BSA, the sera from white-footed mice or robins

collected at 7, 14, 21, 28, and 56 days post nymph feeding were diluted in 50µl of PBS (1:100×,

1:300×, or 1:900×) and then added to the wells. After the samples were washed with PBST buffer,

the microtiter wells were incubated with antibodies that recognize the Fc region of IgG from *P.*

leucopus (1:1,000×, Serocare, Inc, Milford, MA), or wild bird (1:10,000×, Bethyl laboratory,

Montgomery, TX). After the addition of these antibodies, tetramethyl benzidine solution

(ThermoFisher) was added, and the absorbance was detected at 620nm for 10 cycles of 60 s kinetic

intervals with 10 s of shaking duration in a Sunrise absorbance ELISA plate reader (Tecan,

Männedorf, Switzerland). The values of greatest maximum slopes of optical density/min/sample

were multiplied by respective serum dilution factors as shown as arbitrary units, representing

antibody titers.

Borreliacidal assays. Sera collected from robins and white-footed mice at 7 days post tick feeding

were used to measure the bactericidal activity against *B. burgdorferi* as described previously (95).

The complement in these serum samples was heat-inactivated by incubating these samples at 55°C

for 2 h. The heat inactivated sera were serially diluted in BSK-II media without rabbit sera

followed by being mixed with complement-preserved sera from guinea pig (Sigma-Aldrich) or

chicken (Biowest, Riverside). Heat-inactivated guinea pig and chicken sera were included as

controls. *Borrelia burgdorferi* strains B31-5A4, 297, or cN40 (1×10^7 spirochetes per strain) were

then mixed prior to their addition to the reaction. At 24 h after incubation, the surviving spirochetes

were quantified by counting the motile bacteria under dark-field microscopy and presented as the

proportion of serum-treated to untreated spirochetes. We also calculated the 50% borreliacidal titer

(BA), which represents the serum dilution rate that eradicates 50% of spirochetes, using dose-

response stimulation fitting in GraphPad Prism 7.

Statistical analyses. Because the data were not normally distributed, we used a Kruskal-Wallis test followed by a two-stage step-up method with a Benjamini, Krieger, and Yekutieli correction for all comparisons (96). Geometric means of duplicated individual samples (qPCR) were used in our calculations. Spirochete burdens among the three *B. burgdorferi* strains (B31-5A4, 297, and cN40) and uninfected samples in robins, white-footed mice, or ticks were compared using normalized qPCR quantity values for each individual. Cell adhesion variation among the *B. burgdorferi* strains were analyzed using the number of spirochetes per 50 fibroblast cells observed under fluorescent microscopy while the variation in expression levels of the genes encoding IFN-γ, TNF, or TNFα-induced protein were first normalized by the gene encoding actin and then significant differences between the constitutively expressed gene, actin, and 18S rRNA were determined using RT-qPCR quantity values. However, when comparing the differences between levels of pro-inflammatory cytokines at early stages of host infection, we used quantitative ELISA values. Spirochete survival in treated and untreated host sera was assessed using the number of mobile spirochetes at 4 h post incubation. We used quantitative ELISA values to determine the significant differences among the *B. burgdorferi* strains and levels of IgG antibody production in robins and white-footed mice. All analyses and figure generation were completed using GraphPad Prism 7 software.

ACKNOWLEDGEMENTS

We thank Thomas Hart and Tristan Nowak for graph editing, and Jen Owen and Jean Tsao for technical support in the protocol of tick infection in robins. We also thank Thomas Hart, Patricia Lederman, Tristan Nowak, and Frank Blaisdell for the husbandry work to maintain tick colony

and robins. We are grateful for John Leong and George Chaconas for sharing the *B. burgdorferi*

strains B31-5A4, 297, and cN40, Sudha Chaturvedi allowing us to use her Bead Homogenizer,

Susan Madison-Antenucci allowing us to use her fluorescence microscope, and Noel Espina for

the assistance with that microscope. We thank Wadsworth Media & Tissue Culture Core for

preparation of *Borrelia* and American robin and white-footed mouse fibroblast culture medium.

This work was supported by NSF IOS-1755370 (DMT, MC and MDW), NSF-IOS1755286 (YL,

ADII, ALM, JS, and LDK), New York State Department of Health Wadsworth Center Start-Up

Grant (ALM and YL). The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript. The authors declare that the research was

conducted in the absence of any commercial or financial relationships that could be construed as

a potential conflict of interest.

REFERENCES

1.   Poulin R. KBR, Morand S. Micromammals and Macroparasites. Tokyo: Springer; 2006.

233-56 p.

2.   Futuyma DJ, Moreno, G. The evolution of ecological specialization. Ann Rev Ecol Syst

1988;19:207-33.

3.   Tufts DM, Hart TM, Chen GF, Kolokotronis SO, Diuk-Wasser MA, Lin YP. Outer surface

protein polymorphisms linked to host-spirochete association in Lyme borreliae. Molecular

microbiology. 2019.

4.   Lin YP, Diuk-Wasser MA, Stevenson B, Kraiczy P. Complement Evasion Contributes to

Lyme Borreliae-Host Associations. Trends in parasitology. 2020;36(7):634-45.

5.   O'Keeffe KR, Oppler ZJ, Brisson D. Evolutionary ecology of Lyme *Borrelia*. Infection,

genetics and evolution : journal of molecular epidemiology and evolutionary genetics in infectious

diseases. 2020:104570.

6.   Kurtenbach K, De Michelis S, Etti S, Schafer SM, Sewell HS, Brade V, et al. Host

association of *Borrelia burgdorferi* sensu lato--the key role of host complement. Trends in

microbiology. 2002;10(2):74-9.

7.   Kurtenbach K, Peacey M, Rijpkema SG, Hoodless AN, Nuttall PA, Randolph SE.

Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and

small rodents in England. Applied and environmental microbiology. 1998;64(4):1169-74.

8.   Kurtenbach K, Schafer SM, Sewell HS, Peacey M, Hoodless A, Nuttall PA, et al.

Differential survival of Lyme borreliosis spirochetes in ticks that feed on birds. Infection and

immunity. 2002;70(10):5893-5.

9.   Hanincova K, Ogden NH, Diuk-Wasser M, Pappas CJ, Iyer R, Fish D, et al. Fitness

variation of *Borrelia burgdorferi* sensu stricto strains in mice. Applied and environmental

microbiology. 2008;74(1):153-7.

10.   Baum E, Hue F, Barbour AG. Experimental infections of the reservoir species *Peromyscus*

leucopus with diverse strains of Borrelia burgdorferi, a Lyme disease agent. mBio.

2012;3(6):e00434-12.

11.   Hanincova K, Kurtenbach K, Diuk-Wasser M, Brei B, Fish D. Epidemic spread of Lyme

borreliosis, northeastern United States. Emerging infectious diseases. 2006;12(4):604-11.

12.   Brisson D, Dykhuizen DE. ospC diversity in *Borrelia burgdorferi*: different hosts are

different niches. Genetics. 2004;168(2):713-22.

13. Vuong HB, Canham CD, Fonseca DM, Brisson D, Morin PJ, Smouse PE, et al. Occurrence

and transmission efficiencies of *Borrelia burgdorferi ospC* types in avian and mammalian wildlife.

Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in

infectious diseases. 2014;27:594-600.

14. Kurtenbach K, Hanincova K, Tsao JI, Margos G, Fish D, Ogden NH. Fundamental

processes in the evolutionary ecology of Lyme borreliosis. Nature reviews Microbiology.

2006;4(9):660-9.

15. Haven J, Magori K, Park AW. Ecological and inhost factors promoting distinct parasite

life-history strategies in Lyme borreliosis. Epidemics. 2012;4(3):152-7.

16. Qiu WG, Bosler EM, Campbell JR, Ugine GD, Wang IN, Luft BJ, et al. A population

genetic study of *Borrelia burgdorferi* sensu stricto from eastern Long Island, New York, suggested

frequency-dependent selection, gene flow and host adaptation. Hereditas. 1997;127(3):203-16.

17. Qiu WG, Dykhuizen DE, Acosta MS, Luft BJ. Geographic uniformity of the Lyme disease

spirochete (*Borrelia burgdorferi*) and its shared history with tick vector (*Ixodes scapularis*) in the

Northeastern United States. Genetics. 2002;160(3):833-49.

18. Brisson D, Dykhuizen DE, Ostfeld RS. Conspicuous impacts of inconspicuous hosts on

the Lyme disease epidemic. Proc Biol Sci. 2008;275(1631):227-35.

19. Brisson D, Baxamusa N, Schwartz I, Wormser GP. Biodiversity of *Borrelia burgdorferi*

strains in tissues of Lyme disease patients. PloS one. 2011;6(8):e22926.

20. Xu Q, Seemanapalli SV, McShan K, Liang FT. Constitutive expression of outer surface

protein C diminishes the ability of *Borrelia burgdorferi* to evade specific humoral immunity.

Infection and immunity. 2006;74(9):5177-84.

21.   Tilly K, Krum JG, Bestor A, Jewett MW, Grimm D, Bueschel D, et al. *Borrelia burgdorferi*

OspC protein required exclusively in a crucial early stage of mammalian infection. Infection and

immunity. 2006;74(6):3554-64.

22.   Liang FT, Yan J, Mbow ML, Sviat SL, Gilmore RD, Mamula M, et al. *Borrelia burgdorferi*

changes its surface antigenic expression in response to host immune responses. Infection and

immunity. 2004;72(10):5759-67.

23.   Andersson M, Scherman K, Raberg L. Multiple-strain infections of *Borrelia afzelii*: a role

for within-host interactions in the maintenance of antigenic diversity? Am Nat. 2013;181(4):545-

54.

24.   Brisson D, Dykhuizen DE. A modest model explains the distribution and abundance of

Borrelia burgdorferi strains. The American journal of tropical medicine and hygiene.

2006;74(4):615-22.

25.   Haven J, Vargas LC, Mongodin EF, Xue V, Hernandez Y, Pagan P, et al. Pervasive

recombination and sympatric genome diversification driven by frequency-dependent selection in

Borrelia burgdorferi, the Lyme disease bacterium. Genetics. 2011;189(3):951-66.

26.   Woolhouse ME, Taylor LH, Haydon DT. Population biology of multihost pathogens.

Science. 2001;292(5519):1109-12.

27.   Gog JR, Grenfell BT. Dynamics and selection of many-strain pathogens. Proceedings of

the National Academy of Sciences of the United States of America. 2002;99(26):17209-14.

28.   Barrett LG, Thrall PH, Burdon JJ, Linde CC. Life history determines genetic structure and

evolutionary potential of host-parasite interactions. Trends Ecol Evol. 2008;23(12):678-85.

29.   Barthold SW, Persing DH, Armstrong AL, Peeples RA. Kinetics of *Borrelia burgdorferi*

dissemination and evolution of disease after intradermal inoculation of mice. The American

journal of pathology. 1991;139(2):263-73.

30.   Hart T, Nguyen NTT, Nowak NA, Zhang F, Linhardt RJ, Diuk-Wasser M, et al.

Polymorphic factor H-binding activity of CspA protects Lyme borreliae from the host complement

in feeding ticks to facilitate tick-to-host transmission. PLoS pathogens. 2018;14(5):e1007106.

31.   Frye AM, Hart TM, Tufts DM, Ram S, Diuk-Wasser MA, Kraiczy P, et al. A soft tick

Ornithodoros moubata salivary protein OmCI is a potent inhibitor to prevent avian complement

activation. Ticks and tick-borne diseases. 2020;11(2):101354.

32.   Finnie JA, Stewart RB, Aston WP. A comparison of cobra venom factor-induced depletion

of serum C3 in eight different strains of mice. Dev Comp Immunol. 1981;5(4):697-701.

33.   Ivanova L, Christova I, Neves V, Aroso M, Meirelles L, Brisson D, et al. Comprehensive

seroprofiling of sixteen *B. burgdorferi* OspC: implications for Lyme disease diagnostics design.

Clinical immunology. 2009;132(3):393-400.

34.   States SL, Brinkerhoff RJ, Carpi G, Steeves TK, Folsom-O'Keefe C, DeVeaux M, et al.

Lyme disease risk not amplified in a species-poor vertebrate community: similar *Borrelia*

burgdorferi tick infection prevalence and OspC genotype frequencies. Infection, genetics and

evolution : journal of molecular epidemiology and evolutionary genetics in infectious diseases.

2014;27:566-75.

35.   Wormser GP, Brisson D, Liveris D, Hanincova K, Sandigursky S, Nowakowski J, et al.

Borrelia burgdorferi genotype predicts the capacity for hematogenous dissemination during early

Lyme disease. The Journal of infectious diseases. 2008;198(9):1358-64.

36. Jones KL, Glickstein LJ, Damle N, Sikand VK, McHugh G, Steere AC. *Borrelia*

burgdorferi genetic markers and disseminated disease in patients with early Lyme disease. Journal

of clinical microbiology. 2006;44(12):4407-13.

37. Wang G, Ojaimi C, Wu H, Saksenberg V, Iyer R, Liveris D, et al. Disease severity in a

murine model of lyme borreliosis is associated with the genotype of the infecting *Borrelia*

burgdorferi sensu stricto strain. The Journal of infectious diseases. 2002;186(6):782-91.

38. Parveen N, Robbins D, Leong JM. Strain variation in glycosaminoglycan recognition

influences cell-type-specific binding by lyme disease spirochetes. Infection and immunity.

1999;67(4):1743-9.

39. Zipfel PF, Skerka C. Complement regulators and inhibitory proteins. Nature reviews

Immunology. 2009;9(10):729-40.

40. Sjoberg AP, Trouw LA, Blom AM. Complement activation and inhibition: a delicate

balance. Trends in immunology. 2009;30(2):83-90.

41. Tracy KE, Baumgarth N. Borrelia burgdorferi Manipulates Innate and Adaptive Immunity

to Establish Persistence in Rodent Reservoir Hosts. Frontiers in immunology. 2017;8:116.

42. Hyde JA. Borrelia burgdorferi Keeps Moving and Carries on: A Review of Borrelial

Dissemination and Invasion. Frontiers in immunology. 2017;8:114.

43. Liu N, Montgomery RR, Barthold SW, Bockenstedt LK. Myeloid differentiation antigen

88 deficiency impairs pathogen clearance but does not alter inflammation in *Borrelia burgdorferi-*

infected mice. Infection and immunity. 2004;72(6):3195-203.

44. Bolz DD, Sundsbak RS, Ma Y, Akira S, Kirschning CJ, Zachary JF, et al. MyD88 plays a

unique role in host defense but not arthritis development in Lyme disease. Journal of immunology.

2004;173(3):2003-10.

45.   Troy EB, Lin T, Gao L, Lazinski DW, Camilli A, Norris SJ, et al. Understanding barriers

to *Borrelia burgdorferi* dissemination during infection using massively parallel sequencing.

Infection and immunity. 2013;81(7):2347-57.

46.   Strle K, Jones KL, Drouin EE, Li X, Steere AC. *Borrelia burgdorferi* RST1 (OspC type

A) genotype is associated with greater inflammation and more severe Lyme disease. The American

journal of pathology. 2011;178(6):2726-39.

47.   Cerar T, Strle F, Stupica D, Ruzic-Sabljic E, McHugh G, Steere AC, et al. Differences in

Genotype, Clinical Features, and Inflammatory Potential of *Borrelia burgdorferi* sensu stricto

Strains from Europe and the United States. Emerging infectious diseases. 2016;22(5):818-27.

48.   Petzke MM, Iyer R, Love AC, Spieler Z, Brooks A, Schwartz I. *Borrelia burgdorferi*

induces a type I interferon response during early stages of disseminated infection in mice. BMC

microbiology. 2016;16:29.

49.   Mason LM, Herkes EA, Krupna-Gaylord MA, Oei A, van der Poll T, Wormser GP, et al.

Borrelia burgdorferi clinical isolates induce human innate immune responses that are not

dependent on genotype. Immunobiology. 2015;220(10):1141-50.

50.   Wang G, Petzke MM, Iyer R, Wu H, Schwartz I. Pattern of proinflammatory cytokine

induction in RAW264.7 mouse macrophages is identical for virulent and attenuated *Borrelia*

burgdorferi. Journal of immunology. 2008;180(12):8306-15.

51.   Kern A, Schnell G, Bernard Q, Boeuf A, Jaulhac B, Collin E, et al. Heterogeneity of

Borrelia burgdorferi Sensu Stricto Population and Its Involvement in *Borrelia* Pathogenicity:

Study on Murine Model with Specific Emphasis on the Skin Interface. PloS one.

2015;10(7):e0133195.

52.   Schramm F, Kern A, Barthel C, Nadaud S, Meyer N, Jaulhac B, et al. Microarray analyses

of inflammation response of human dermal fibroblasts to different strains of *Borrelia burgdorferi*

sensu stricto. PloS one. 2012;7(6):e40046.

53.   Li X, McHugh GA, Damle N, Sikand VK, Glickstein L, Steere AC. Burden and viability

of *Borrelia burgdorferi* in skin and joints of patients with erythema migrans or lyme arthritis.

Arthritis and rheumatism. 2011;63(8):2238-47.

54.   Jones KL, McHugh GA, Glickstein LJ, Steere AC. Analysis of *Borrelia burgdorferi*

genotypes in patients with Lyme arthritis: High frequency of ribosomal RNA intergenic spacer

type 1 strains in antibiotic-refractory arthritis. Arthritis and rheumatism. 2009;60(7):2174-82.

55.   Dykhuizen DE, Brisson D, Sandigursky S, Wormser GP, Nowakowski J, Nadelman RB,

et al. The propensity of different *Borrelia burgdorferi* sensu stricto genotypes to cause

disseminated infections in humans. The American journal of tropical medicine and hygiene.

2008;78(5):806-10.

56.   Barthold SW, Bockenstedt LK. Passive immunizing activity of sera from mice infected

with *Borrelia burgdorferi*. Infection and immunity. 1993;61(11):4696-702.

57.   Fikrig E, Bockenstedt LK, Barthold SW, Chen M, Tao H, Ali-Salaam P, et al. Sera from

patients with chronic Lyme disease protect mice from Lyme borreliosis. The Journal of infectious

diseases. 1994;169(3):568-74.

58.   Wang G, Ojaimi C, Iyer R, Saksenberg V, McClain SA, Wormser GP, et al. Impact of

genotypic variation of *Borrelia burgdorferi* sensu stricto on kinetics of dissemination and severity

of disease in C3H/HeJ mice. Infection and immunity. 2001;69(7):4303-12.

59.   Benoit VM, Fischer JR, Lin YP, Parveen N, Leong JM. Allelic variation of the Lyme

disease spirochete adhesin DbpA influences spirochetal binding to decorin, dermatan sulfate, and

mammalian cells. Infection and immunity. 2011;79(9):3501-9.

60.   Roberts WC, Mullikin BA, Lathigra R, Hanson MS. Molecular analysis of sequence

heterogeneity among genes encoding decorin binding proteins A and B of *Borrelia burgdorferi*

sensu lato. Infection and immunity. 1998;66(11):5275-85.

61.   Wilske B, Preac-Mursic V, Jauris S, Hofmann A, Pradel I, Soutschek E, et al.

Immunological and molecular polymorphisms of OspC, an immunodominant major outer surface

protein of Borrelia burgdorferi. Infection and immunity. 1993;61(5):2182-91.

62.   Lin YP, Tan X, Caine JA, Castellanos M, Chaconas G, Coburn J, et al. Strain-specific joint

invasion and colonization by Lyme disease spirochetes is promoted by outer surface protein C.

PLoS pathogens. 2020;16(5):e1008516.

63.   Lin YP, Benoit V, Yang X, Martinez-Herranz R, Pal U, Leong JM. Strain-specific variation

of the decorin-binding adhesin DbpA influences the tissue tropism of the lyme disease spirochete.

PLoS pathogens. 2014;10(7):e1004238.

64.   Isogai E, Tanaka S, Braga IS, 3rd, Itakura C, Isogai H, Kimura K, et al. Experimental

Borrelia garinii infection of Japanese quail. Infection and immunity. 1994;62(8):3580-2.

65.   Schwan TG, Piesman J, Golde WT, Dolan MC, Rosa PA. Induction of an outer surface

protein on *Borrelia burgdorferi* during tick feeding. Proceedings of the National Academy of

Sciences of the United States of America. 1995;92(7):2909-13.

66.   Hodzic E, Feng S, Freet KJ, Borjesson DL, Barthold SW. *Borrelia burgdorferi* population

kinetics and selected gene expression at the host-vector interface. Infection and immunity.

2002;70(7):3382-8.

67.   Hodzic E, Feng S, Freet KJ, Barthold SW. *Borrelia burgdorferi* population dynamics and

prototype gene expression during infection of immunocompetent and immunodeficient mice.

Infection and immunity. 2003;71(9):5042-55.

68.   Bykowski T, Woodman ME, Cooley AE, Brissette CA, Brade V, Wallich R, et al.

Coordinated expression of *Borrelia burgdorferi* complement regulator-acquiring surface proteins

during the Lyme disease spirochete's mammal-tick infection cycle. Infection and immunity.

2007;75(9):4227-36.

69.   Iyer R, Caimano MJ, Luthra A, Axline D, Jr., Corona A, Iacobas DA, et al. Stage-specific

global alterations in the transcriptomes of Lyme disease spirochetes during tick feeding and

following mammalian host adaptation. Molecular microbiology. 2015;95(3):509-38.

70.   Zhang JR, Hardham JM, Barbour AG, Norris SJ. Antigenic variation in Lyme disease

borreliae by promiscuous recombination of VMP-like sequence cassettes. Cell. 1997;89(2):275-

85.

71.   Verhey TB, Castellanos M, Chaconas G. Analysis of recombinational switching at the

antigenic variation locus of the Lyme spirochete using a novel PacBio sequencing pipeline.

Molecular microbiology. 2018;107(1):104-15.

72.   Coutte L, Botkin DJ, Gao L, Norris SJ. Detailed analysis of sequence changes occurring

during *vlsE* antigenic variation in the mouse model of *Borrelia burgdorferi* infection. PLoS

pathogens. 2009;5(2):e1000293.

73.   Shih CM, Telford SR, 3rd, Pollack RJ, Spielman A. Rapid dissemination by the agent of

Lyme disease in hosts that permit fulminating infection. Infection and immunity. 1993;61(6):2396-

9.

74.   Caine JA, Lin YP, Kessler JR, Sato H, Leong JM, Coburn J. *Borrelia burgdorferi* outer

surface protein C (OspC) binds complement component C4b and confers bloodstream survival.

Cellular microbiology. 2017.

75.   Lagal V, Portnoi D, Faure G, Postic D, Baranton G. *Borrelia burgdorferi* sensu stricto

invasiveness is correlated with OspC-plasminogen affinity. Microbes and infection / Institut

Pasteur. 2006;8(3):645-52.

76.   Rogers EA, Marconi RT. Delineation of species-specific binding properties of the CspZ

protein (BBH06) of Lyme disease spirochetes: evidence for new contributions to the pathogenesis

of *Borrelia* spp. Infection and immunity. 2007;75(11):5272-81.

77.   Wallich R, Pattathu J, Kitiratschky V, Brenner C, Zipfel PF, Brade V, et al. Identification

and functional characterization of complement regulator-acquiring surface protein 1 of the Lyme

disease spirochetes *Borrelia afzelii* and *Borrelia garinii*. Infection and immunity.

2005;73(4):2351-9.

78.   Hammerschmidt C, Koenigs A, Siegel C, Hallstrom T, Skerka C, Wallich R, et al. Versatile

roles of CspA orthologs in complement inactivation of serum-resistant Lyme disease spirochetes.

Infection and immunity. 2014;82(1):380-92.

79.   Alitalo A, Meri T, Comstedt P, Jeffery L, Tornberg J, Strandin T, et al. Expression of

complement factor H binding immunoevasion proteins in *Borrelia garinii* isolated from patients

with neuroborreliosis. European journal of immunology. 2005;35(10):3043-53.

80.   Xie J, Zhi H, Garrigues RJ, Keightley A, Garcia BL, Skare JT. Structural determination of

the complement inhibitory domain of *Borrelia burgdorferi* BBK32 provides insight into classical

pathway complement evasion by Lyme disease spirochetes. PLoS pathogens.

2019;15(3):e1007659.

81.   Margos G, Vollmer SA, Cornet M, Garnier M, Fingerle V, Wilske B, et al. A new *Borrelia*

species defined by multilocus sequence analysis of housekeeping genes. Applied and

environmental microbiology. 2009;75(16):5410-6.

82.   Norte AC, Margos G, Becker NS, Albino Ramos J, Nuncio MS, Fingerle V, et al. Host

dispersal shapes the population structure of a tick-borne bacterial pathogen. Mol Ecol. 2019.

83.   Barbour AG. Isolation and cultivation of Lyme disease spirochetes. The Yale journal of

biology and medicine. 1984;57(4):521-5.

84.   Marcinkiewicz AL, Dupuis AP, 2nd, Zamba-Campero M, Nowak N, Kraiczy P, Ram S, et

al. Blood treatment of Lyme borreliae demonstrates the mechanism of CspZ-mediated complement

evasion to promote systemic infection in vertebrate hosts. Cellular microbiology.

2019;21(2):e12998.

85.   Purser JE, Norris SJ. Correlation between plasmid content and infectivity in *Borrelia*

burgdorferi. Proceedings of the National Academy of Sciences of the United States of America.

2000;97(25):13865-70.

86.   Blevins JS, Hagman KE, Norgard MV. Assessment of decorin-binding protein A to the

infectivity of *Borrelia burgdorferi* in the murine models of needle and tick infection. BMC

microbiology. 2008;8:82.

87.   Chan K, Casjens S, Parveen N. Detection of established virulence genes and plasmids to

differentiate *Borrelia burgdorferi* strains. Infection and immunity. 2012;80(4):1519-29.

88.   Marcinkiewicz AL, Lieknina I, Yang X, Lederman PL, Hart TM, Yates J, et al. The Factor

H-Binding Site of CspZ as a Protective Target against Multistrain, Tick-Transmitted Lyme

Disease. Infection and immunity. 2020;88(5).

89.   Izuogu AO, McNally KL, Harris SE, Youseff BH, Presloid JB, Burlak C, et al. Interferon

signaling in *Peromyscus leucopus* confers a potent and specific restriction to vector-borne

flaviviruses. PloS one. 2017;12(6):e0179781.

90.   Calhoon EA, Miller, M. K., Jimenez, A. G., Harper, J. M., Williams, J. B. . Changes in

cultured dermal fibroblasts during early passages across five wild bird species. Canadian Journal

of Zoology. 2013;91:653-9.

91.   Kern A, Zhou CW, Jia F, Xu Q, Hu LT. Live-vaccinia virus encapsulation in pH-sensitive

polymer increases safety of a reservoir-targeted Lyme disease vaccine by targeting gastrointestinal

release. Vaccine. 2016;34(38):4507-13.

92.   Johnston E, Tsao JI, Munoz JD, Owen J. *Anaplasma phagocytophilum* infection in

American robins and gray catbirds: an assessment of reservoir competence and disease in captive

wildlife. Journal of medical entomology. 2013;50(1):163-70.

93.   Varkey R, Du Q, Karnell JL, Xiao X, Casey KA, Woods R, et al. Discovery and

characterization of potent IL-21 neutralizing antibodies via a novel alternating antigen

immunization and humanization strategy. PloS one. 2019;14(1):e0211236.

94.   Lin YP, Yu Y, Marcinkiewicz AL, Lederman P, Hart TM, Zhang F, et al. Non-

anticoagulant Heparin as a Pre-exposure Prophylaxis Prevents Lyme Disease Infection. ACS

Infect Dis. 2020;6(3):503-14.

95.   Marcinkiewicz AL, Lieknina I, Kotelovica S, Yang X, Kraiczy P, Pal U, et al. Eliminating

Factor H-Binding Activity of Borrelia burgdorferi CspZ Combined with Virus-Like Particle

Conjugation Enhances Its Efficacy as a Lyme Disease Vaccine. Frontiers in immunology.

2018;9:181.

96. Benjamini YK, A. M. Yekutieli, D. . Adaptive linear step-up procedures that control the
false discovery rate. Biometrika. 2006;93:491-507.

97. Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. Lyme disease-
a tick-borne spirochetosis? Science. 1982;216(4552):1317-9.

98. Leong JM, Moitoso de Vargas L, Isberg RR. Binding of cultured mammalian cells to
immobilized bacteria. Infection and immunity. 1992;60(2):683-6.

99. Steere AC, Grodzicki RL, Kornblatt AN, Craft JE, Barbour AG, Burgdorfer W, et al. The
spirochetal etiology of Lyme disease. The New England journal of medicine. 1983;308(13):733-
40.

100. Barthold SW, de Souza MS, Janotka JL, Smith AL, Persing DH. Chronic Lyme borreliosis
in the laboratory mouse. The American journal of pathology. 1993;143(3):959-71.

101. Barthold SW, Beck DS, Hansen GM, Terwilliger GA, Moody KD. Lyme borreliosis in
selected strains and ages of laboratory mice. The Journal of infectious diseases. 1990;162(1):133-
8.

102. Sadziene A, Wilske B, Ferdows MS, Barbour AG. The cryptic *ospC* gene of *Borrelia*
burgdorferi B31 is located on a circular plasmid. Infection and immunity. 1993;61(5):2192-5.

103. Lin T, Oliver JH, Jr., Gao L. Genetic diversity of the outer surface protein C gene of
southern *Borrelia* isolates and its possible epidemiological, clinical, and pathogenetic
implications. Journal of clinical microbiology. 2002;40(7):2572-83.

104. Wang IN, Dykhuizen DE, Qiu W, Dunn JJ, Bosler EM, Luft BJ. Genetic diversity of *ospC*
in a local population of *Borrelia burgdorferi* sensu stricto. Genetics. 1999;151(1):15-30.

105.   Chan K, Awan M, Barthold SW, Parveen N. Comparative molecular analyses of *Borrelia burgdorferi* sensu stricto strains B31 and N40D10/E9 and determination of their pathogenicity. BMC microbiology. 2012;12:157.

106.   Barbour AG, Travinsky B. Evolution and distribution of the ospC Gene, a transferable serotype determinant of *Borrelia burgdorferi*. mBio. 2010;1(4).

TABLES

Table 1. The number of positive xenodiagnostic larval ticks collected from robins and white-footed mice.

dpf[a]	Positive xenodiagnostic larval ticks acquired from			
	Mice fed on by naïve nymphs	Mice fed on by nymphs carrying *B. burgdorferi*		
		B31-5A4	297	cN40
American robins				
14	0/44	39/44[b]	19/44[b,c]	41/55[b]
28	0/44	18/33[b,c]	11/44[b,c]	46/55[b]
35	0/44	11/33[b,c]	12/44[b,c]	34/55[b]
56	0/44	8/33[b,c]	7/44[b,c]	30/55[b]
White-footed mice				
14	0/55	47/55[b,c]	47/55[b,c]	16/55[b]
28	0/33	50/55[b,c]	40/55[b,c]	8/55[b]
35	0/33	51/55[b,c]	40/55[b,c]	9/55[b]
56	0/33	38/55[b,c]	35/55[b,c]	5/55[b]

[a]Days post nymph feeding.
[b]Difference compared with larvae from the mice fed on by naïve nymphs by two-tailed Fisher test.
[c]Difference compared with larvae from the mice fed on by nymphs carrying strain cN40 by two-tailed Fisher test.

 FIGURE LEGENDS

Figure 1. *Borrelia burgdorferi* **B31-5A4, 297, and cN40 differed in early colonization in robins and white-footed mice and adhesion and cytokine triggering in these animals-derived fibroblasts. (A and B): (A)** American robins and **(B)** white-footed (WF) mice were intradermally inoculated with 10^4 *B. burgdorferi* B31-5A4, 297, or cN40, or with BSK-II medium without rabbit sera as mock infection ("Mock"). The inoculation site of skin from these robins and mice were collected at 1-day post infection (1dpi) to determine bacterial burdens by qPCR. The bacterial loads in the tissues or blood from robins or mice were normalized to 100 ng total DNA. Shown are the geometric mean ± geometric standard deviation of bacterial burdens in those tissues from four robins or mice per group. Significant differences (P < 0.05) in the spirochete burdens (normalized qPCR quantity values) relative to the mock infected group are indicated (''*''). **(C and D):** *B. burgdorferi* B31-5A4, 297, cN40, or B314 (negative control) (2×10^6 spirochetes) were incubated with fibroblasts (2×10^5 cells) from **(C)** American robins and **(D)** white-footed (WF) mice for 1 h. We mixed those cells with FITC-conjugated goat anti-*B. burgdorferi* polyclonal antibodies and visualized the spirochetes after fixation. Additionally, DAPI was incubated with these cells to localize the nuclei from fibroblasts. The levels of spirochete attachment were evaluated by counting the number of bacteria per 50 cells under fluorescence microscopy described in the section "Materials and Methods." Each bar represents the mean of four independent determinations ± standard deviation. Asterisks indicate significant differences (P < 0.05) in spirochetal attachment relative using normalized qPCR quantity values to strain B314. **(E to J):** Fibroblasts (2×10^5 cells) from **(E to G)** American robins and **(H to J)** white-footed (WF) mice were incubated for 24h with *B. burgdorferi* B31-5A4, 297, or cN40 (2×10^7 of spirochetes for spirochete to cell ration (MOI) at 1:100). Cell media-treated fibroblasts were included as control

("Mock"). After the RNA was extracted from these cells, the expression levels of the genes

encoding IFN-γ, TNF, or TNFα-induced protein and the constitutively expressed gene, actin, and

18S rRNA, from robins and white-footed mice were determined using quantitative reverse

transcription polymerase chain reaction. The expression levels of the genes encoding **(E and H)**

18S rRNA, **(F and I)** IFNγ, **(G)** TNFα-induced protein, and **(J)** TNF are presented by normalizing

to the expression levels of the gene encoding actin. Each bar represents the mean of three

independent determinations ± SD. The asterisk ("*") indicates significant differences ($P < 0.05$;)

in the normalized expression levels of the gene encoding IFN-γ ,TNF, and TNFα-induced protein

in fibroblasts treated with indicated spirochete strains relative to those in mock-treated cells.

1099

Figure 2. *Borrelia burgdorferi* B31-5A4, 297, and cN40 displayed strain-to-strain variation of

xenodiagnostics acquisition from robins and white-footed mice. (A) The experimental timeline

of infection. **(B to K):** *Ixodes scapularis* nymphs carrying *B. burgdorferi* B31-5A4, 297, or cN40,

or naïve nymphs (Uninfect.) were allowed to feed to repletion on **(B to F)** 3, 4, 5, or 4 American

robins, respectively, or **(G to K)** 5 white-footed (WF) mice per group. Approximately 100 larval

ticks were placed on each robin at the time points indicated in Fig. 2A to feed till repletion. qPCR

was used to determine spirochete burdens derived from 55 larvae feeding on *B. burgdorferi*-

infected white-footed mice or 33, 44, 55, and 44 larvae feeding on robins fed on by nymphs

carrying B31-5A4, 297, or cN40, or uninfected nymphs, respectively. **(B and F)** The larvae were

considered xenodiagnostic positive if their spirochete burdens were greater than the threshold, the

mean plus three-fold standard deviation of spirochete burdens in the uninfected group. Shown are

the means ± SEM of percent positive larvae. **(C to F and H to K)** Shown are the geometric means

± geometric standard deviation of bacterial burdens in larvae that are allowed to feed on robins or

white-footed (WF) mice at **(C and H)** 14, **(D and I)** 28, **(E and J)** 35, and **(F and K)** 56 days post nymph feeding (dpf). Significant differences (p < 0.05) in the spirochete burdens relative to larvae feeding on naïve robins or white-footed mice (''*'') or between different groups ("#") are indicated.

Figure 3. *Borrelia burgdorferi* B31-5A4, 297, and cN40 differed in their ability to induce bacteremia and colonize tissues in robins and white-footed mice. *Ixodes scapularis* nymphs carrying *B. burgdorferi* B31-5A4, 297, or cN40, or naïve nymphs (Uninfect.) were allowed to feed to repletion on **(A to D and I to L)** 3, 4, 5, and 4 American robins, respectively, or **(E to H and M to P)** 5 white-footed (WF) mice per group. The bacterial loads in the blood at **(A and E)** 7, **(B and F)** 14, **(C and G)** 21, and **(D and H)** 28 days post nymph feeding (dpf) and in **(I)** skin, **(J)** heart, **(K)** brain, and **(L)** liver of robins and **(M)** ears, **(N)** tibiofemoral joints (Tibio.), **(O)** heart, **(P)** bladder of white-footed mice at 64 days post nymph feeding (dpf) were determined by qPCR. The bacterial loads in blood were normalized to 100 ng total DNA. Shown are the geometric mean ± geometric standard deviation of indicated number of robins or white-footed mice. Significant differences (p < 0.05) in the spirochete burdens relative to robins or white-footed mice fed on by naïve nymphs (''*'') are indicated.

Figure 4. *Borrelia burgdorferi* B31-5A4, 297, and cN40 varied in their ability to survive in robin but not white-footed mouse sera. A high passage, non-infectious, serum sensitive *B. burgdorferi* strain B313 ("B313") or *B. burgdorferi* B31-5A4, 297, or cN40 were incubated for 4 h with the sera from American robins at a final concentration of 40% in the **(A)** absence or **(B)** presence of 2 μM of OmCI. Each of these spirochete strains was also incubated for 4 h with the

sera from white-footed (WF) mice at a final concentration of 40% in the **(C)** absence or **(D)**

presence of 2 µM of CVF. The above-mentioned sera were also heat-inactivated and included as

controls. The number of motile spirochetes was assessed microscopically. The percentage of

survival for *B. burgdorferi* was calculated using the number of mobile spirochetes at 4 h post

incubation normalized to that prior to the incubation with serum. The assays were performed at

three independent occasions; within each experiment, samples were run in triplicate, and the

survival percentage for each experiment was calculated by averaging the results from triplicate

runs. The result shown here are the average ± standard deviation of the survival percentage from

three independent experiments. Significant differences ($P < 0.05$) of the percent survival of

spirochetes between groups are indicated (#).

Figure 5. *Borrelia burgdorferi* B31-5A4, 297, and cN40 triggered different levels of pro-inflammatory cytokines at early stages of robin and white-footed mice infection. *Ixodes scapularis* nymphs infected with *B. burgdorferi* B31-5A4, 297, cN40, or naïve nymphs (Uninfect.) were allowed to feed to repletion on **(A to H)** American robins or **(I to P)** white-footed (WF) mice. The sera were obtained **(A, E, I, and M)** prior to tick feeding or at **(B, F, J, and N)** 7-, **(C, G, K, and O)** 14-, or **(D, H, L, and P)** 21-days post nymph feeding ("dpf"). The levels of IFNγ **(A to D and I to L)** and TNFα **(E to H and M to P)** were determined using quantitative ELISA. Shown are the geometric mean ± geometric standard deviation of five white-footed mice per group or robins (3, 4, 5, and 4 for the strains B31-5A4-, 297-, cN40-infected robins or uninfected robins per group, respectively). Significant differences ($p < 0.05$) in the cytokine levels relative to robins or white-footed mice fed on by naïve nymphs ("*") are indicated.

Figure 6. *Borrelia burgdorferi* **B31-5A4, 297, and cN40 differed in the ability to induce antibodies against spirochetes in robin and white-footed mice during infection.** *Ixodes scapularis* nymphs infected with *B. burgdorferi* B31-5A4, 297, or cN40, or naïve nymphs (Uninfect.) were allowed to feed to repletion on **(A to E)** American robins or **(F to J)** white-footed (WF) mice. The sera were obtained at **(A and F)** 0-, **(B and G)** 7-, **(C and H)** 14-, **(D and I)** 21-, or **(E and J)** 28-days post nymph feeding ("dpf"). The levels of IgG against the mixture of *B. burgdorferi* strains B31-5A4, 297, and cN40 (1×10^6 spirochetes per strain in each microtiter well) were determined using quantitative ELISA. Shown are the geometric mean ± geometric standard deviation of five white-footed mice or robins per group (3, 4, 5, and 4 for the strains B31-5A4-, 297-, cN40-infected robins or uninfected robins per group, respectively). Significant differences ($p < 0.05$) in the antibody titers relative to robins or white-footed mice fed on by naive nymphs ("*") are indicated. Sera from the **(K)** American robins and **(L)** white-footed (WF) mice at 7 days after fed on by *I. scapularis* nymphs carrying *B. burgdorferi* strains B31-5A4, 297, or cN40, or naïve nymphs (Uninfect.) were collected. These sera were serially diluted as indicated and mixed with **(K)** chicken or **(L)** guinea pig complement and the mixture of the strains B31-5A4, 297, and cN40 (1×10^7 spirochetes per strain in each reaction). After incubation for 24 h, surviving spirochetes were quantified from three fields of view for each sample microscopically. The experiment was performed at three independent occasions. The survival percentage was derived from the proportion of serum-treated to untreated spirochetes. Data shown are the mean ± standard error of the mean (SEM) of the survival percentage from three replicates in one representative experiment. The 50% borreliacidal dilution of each serum sample (50% BA), representing the dilution rate that effectively killed 50% of spirochetes, was obtained from curve-fitting and extrapolation of data shown in the table above respective panels. The 50% BA of some strain-sera

1182 pairs could not be determined because no robust killing was observed, resulting in curves that did

1183 not fit (N.F.).

1184

Figure 7. The schematic diagram showing the model supported by this study. Upon *B.*

burgdorferi transmission from ticks to hosts, multiple cellular and immunological mechanisms

may play in concert to determine the fitness of spirochetes in hosts. These mechanisms include the

ability of *B. burgdorferi* to attach to cells or tissues (spirochete adhesiveness) and escape from host

immune clearance (early host defense), such as complement, cytokine-mediated responses, and

antibodies. Both mechanisms applied on spirochetes would lead to host specialization, resulting in

the association of *B. burgdorferi* with particular hosts (i.e. mammal or avian specialists).

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

Fig. 1

Fig. 2

A

Days post initial nymph feeding (dpf)

0	7	14	21	28	35	56	64
Place nymphs	Serum & blood collection	Xenodiag. larvae Serum & blood collection	Serum & Blood collection	Xenodiag. larvae Serum & blood collection	Xenodiag. larvae	Xenodiag. larvae	Tissue collection

Fig. 3

A Blood 7dpf **B** 14dpf **C** 21dpf **D** 28dpf

E 7dpf **F** 14dpf **G** 21dpf **H** 28dpf

I Skin 56dpf **J** Heart 56dpf **K** Brain 56dpf **L** Liver 56dpf

M Ears 56dpf **N** Tibio. 56dpf **O** Heart 56dpf **P** Bladder 56dpf

Robin

WF mouse

Robin

WF mouse

Spirochete burdens (Bact. per 100ngDNA)

Unifect. B31-5A4 297 cN40

Fig. 4

Fig. 5

Fig. 6

Fig. 7

1) **Spirochete adhesiveness**
2) **Early Host defense:**
   Complement
   Cytokine-mediated response
   Antibodies

Host specialization

Avian specialist

Mammalian specialist