SCNIC: Sparse Correlation Network Investigation for Compositional Data

Michael Shaffer[1,2,†], Kumar Thurimella[1,3,†] , John D. Sterrett[4] and Catherine A. Lozupone[1,*]

1 Department of Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA 80045

2 Current: Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO 80523, USA

3 Current: Department of Chemical Engineering and Biotechnology, University of Cambridge, Cambridge, UK

4 Department of Integrative Physiology, University of Colorado, Boulder, CO, USA 80309

†*These authors contributed equally to this work*

**To whom correspondence should be addressed*

ABSTRACT

Background

Microbiome studies are often limited by a lack of statistical power due to small sample sizes and a large number of features. This problem is exacerbated in correlative studies of multi-omic datasets. Statistical power can be increased by finding and summarizing modules of correlated observations, which is one dimensionality reduction method. Additionally, modules provide biological insight as correlated groups of microbes can have relationships among themselves.

Results

To address these challenges, we developed SCNIC: Sparse Cooccurrence Network Investigation for compositional data. SCNIC is open-source software that can generate correlation networks and detect and summarize modules of highly correlated features. Modules can be formed using either the Louvain Modularity Maximization (LMM) algorithm or a Shared Minimum Distance algorithm (SMD) that we newly describe here and relate to LMM using simulated data. We applied SCNIC to two published datasets and we achieved increased statistical power and identified microbes that not only differed across groups, but also correlated strongly with each other, suggesting shared environmental drivers or cooperative relationships among them.

Conclusions

SCNIC provides an easy way to generate correlation networks, identify modules of correlated features and summarize them for downstream statistical analysis. Although SCNIC was designed considering properties of microbiome data, such as compositionality and sparsity, it can be applied to a variety of data types including metabolomics data and used to integrate multiple data types. SCNIC allows for the identification of functional microbial relationships at scale while increasing statistical power through feature reduction.

BACKGROUND

Microbial communities play important roles in environmental and human health systems and can often reach great complexity. In these rich ecosystems, microbes interact with each other, forming relationships based on predator-prey dynamics [1], competition for resources [2], cross-feeding of small compounds, [3] and other factors. Identifying correlated pairs of microbes can suggest potential interactions or shared environmental preferences. Accordingly, studies have identified complex networks of co-occurring microbes in a variety of different environments ranging from the human mouth and gut [4] to soil [5] and stream ecosystems [6].

To detect correlations between microbes, a variety of methods have been developed. While traditional correlation metrics are used by some [7-9], newer methods have been developed that take into account the properties of 16S rRNA sequencing data [10-12]. A recent review tested these methods on a variety of models and identified some methods that performed better than others in ways that can depend on underlying data characteristics [13]. Although these tools are useful for finding pairwise relationships between organisms, less attention has been given toward developing methods for finding correlations among groups of microbes.

One way to explore complex interactions is to form networks in which correlated organisms are joined with an edge, and highly correlated sets of microbes are defined. Here, we refer to these sets as modules, which are synonymous to clusters or groups. There are two primary benefits of finding modules of correlated microbes. First, the combination of microbes in a module could be further explored to understand microbial interactions, such as cross-feeding relationships, or shared environmental niches [5, 14-16]. Second, considering correlation structure among microbes can aid in statistical analysis aimed at uncovering relationships between microbes and other environmental factors. Specifically, by eliminating or summarizing

highly correlated features, dependence between features is decreased. Feature reduction will

increase accuracy of methods that assume the independence of features such as false discovery

rate technique (FDR) measurements like the Benjamini-Hochberg Correction [17], and statistical

power is increased by reducing the number of feature comparisons.

One workflow for considering groups of correlated microbes in downstream statistical

analyses requires three steps: first, correlations between microbes must be measured and used to

form a network; second, modules must be identified; and third, abundance of the microbes in

modules must be summarized for use in subsequent statistical analyses. One software tool that

has implemented this workflow, developed for application to gene expression data, is weighted

gene correlation network analysis (WGCNA)[18]. WGCNA builds correlation networks based

on a correlation coefficient (such as Pearson, Spearman, or biweight midcorrelation [19]), and

detects modules as subtrees in a hierarchical cluster of features [20]. Modules are summarized by

setting module abundance to that of network hubs or an eigenvector of the abundance of all

module members [18].

Several groups have used WGCNA to find correlations within 16S rRNA sequencing

data [21-24], but this approach may not be appropriate for several reasons [25]. First, the

correlation metrics implemented in WGCNA do not account for sparsity and compositionality.

Most sequencing-based microbiome datasets are sparse (i.e. there are many zeros) and

compositional, meaning they only carry information on relative abundances of taxa instead of

absolute abundances, which can lead to the detection of spurious correlations if proper statistical

methods are not used [26]. Thus, the use of WGCNA for compositional data may be leading to

the detection of spurious edges in microbiome networks. Second, the primary method WGCNA

uses to pick modules assumes the correlation network will have a scale-free topology that may

not be relevant to microbiome data [27]. Third, summarizing modules through identifying hub

taxa works well in gene expression where a single transcription factor can control the expression

of many genes, but may not be appropriate in microbial communities. Both the hub and

eigenvector approaches to module summarization do not allow for output tables that maintain the

total counts of microbial abundance per sample. Therefore, the hub and eigenvector approaches

cannot be used with tools developed for microbiome data analysis that make assumptions based

on total sample counts, such as ANCOM [28] or metagenomeSeq [29].

95   Optimal methods for identifying and summarizing modules of correlated features in 16S

rRNA sequencing data have not been deeply explored. One study [25] recommended an

ensemble approach for correlation detection, and the Louvain modularity maximization (LMM)

method [30] to identify modules. LULU is a tool that follows a binning approach towards OTUs

that co-occur, but only does so if they are highly phylogenetically related [31]. Another tool,

CoNet, uses an ensemble approach to build and visualize networks [32]. However, no

implementation of module summarization was made available for downstream statistical

analysis.

103   To address these gaps, we have developed a tool for sparse, compositional correlation

network investigation for compositional data (SCNIC), which uses methods optimized for

microbiome data analysis. SCNIC is available as standalone Python software, via Bioconda [33]

and the package installer for Python (pip), and as a QIIME 2 plugin [34]. The source code for

SCNIC and the QIIME 2 plugin is freely available on GitHub

(https://github.com/lozuponelab/SCNIC, https://github.com/lozuponelab/q2-SCNIC) under the

BSD-3-Clause License.

MATERIALS AND METHODS

The SCNIC method

SCNIC takes a feature table containing counts of each feature in all samples as input and

performs three steps: 1) a correlation network is built, 2) modules are detected in the network

and 3) feature counts within a module are summed into a new single feature (identified as

"module-*x*" where x is whole numbered consecutively starting at zero)(Figure 1). The modules

are ordered based on size, where the lower numbered modules have a larger number of members

compared to higher numbered modules. To summarize modules, SCNIC uses a sum of count

data from all features in a module. There is no maximum or minimum size constraint on module

size when modules are created. The newly generated modules are included in a new feature table

alongside all features not grouped into a module. This maintains the total counts per sample,

allowing for downstream analyses with tools that have assumptions related to total sample

counts. SCNIC produces a graph modeling language (GML) format [35] file compatible with

Cytoscape [36] for network visualization in which the edges in the correlation network represent

the positive correlations which are stronger than a user specified R-value cutoff (between 0 and

1), a file describing which features compose each defined module, and a feature table in the

Biological Observation Matrix (BIOM) [37] (Figure 1).

SCNIC allows users to choose between multiple methods for detecting correlations and

of defining modules of co-occurring microbes. For correlations, SCNIC can implement

traditional correlation metrics (including Pearson's *r*, Spearman's ρ and Kendall's τ) or the

compositionality- and sparsity-aware correlation metric from SparCC [38, 39] to correct for

aspects of microbiome data. SparCC has been shown to perform well in detecting correlations

compared to other correlation measures [13]. Specifically, SparCC performs well in communities

with an inverse Simpson index above 13 (which would be indicative of a high number of

successful species, a complex food web, and many ecological niches, as would be seen in many

high biomass microbial communities such as gut or soil microbiomes) [39,40], and it thus was

chosen as the default metric.

To define modules of co-correlated features, we implement two methods: 1) Louvain

modularity maximization (LMM) and 2) a novel shared minimum distance (SMD) module

detection algorithm; unlike WGCNA, neither of these algorithms make assumptions about

network topology. LMM was previously proposed as a method for clustering correlation

networks of microbes into modules [30]. LMM works by first assigning one module per feature.

Each pair of adjacent modules are joined and the change in modularity (defined by the number of

edges within the module compared to outside) is calculated for each module. The pair which

increases the mean modularity of the network the most is then joined. This process is repeated

until the modularity of the network is not increased. LMM uses two parameters provided by the

user: The first parameter, R-value, defines the minimum correlation coefficient for defining an

edge between features. The second parameter, gamma (also referred to as resolution), controls

the size of modules detected, with large gamma values yielding larger modules.

WGCNA and LMM have a potential weakness in that modules can contain pairs of taxa

that are not strongly correlated (e.g. if they are several steps away from each other in the

network). To address this weakness, we also implement the SMD method to ensure that

correlations between all pairs of features in the module have an R-value greater than the user

provided minimum (Figure 2). Specifically, the SMD method defines modules by first applying

complete linkage hierarchical clustering to correlation coefficients to make a tree of features.

Next, SMD defines modules as subtrees where correlations between all pairs of tips have an R-

value above the specified value. SMD has been set as the default method in SCNIC because of

the desirable property of only producing modules where all features are correlated over a user-

specified threshold.

160   A large proportion of microbiome studies sample highly uneven communities which

leads to strong compositionality-driven artifacts [26, 40, 41]. Because of this, we use SparCC,

specifically the implementation of FastSpar [39], as the default correlation measure. SparCC was

used as the correlation metric based on analysis that suggested a high precision in the number of

correct edges recovered when correlations were calculated in synthetic data [13]. SCNIC

additionally includes the option of using Pearson's r, Spearman's ρ and Kendall's τ to evaluate

non-compositional or dense data types.

Evaluating the SMD algorithm using simulated data

169   Since SMD has not been applied to microbiome module detection before, we compared

SMD to LMM using simulated data. In order to evaluate the performance of SMD for module

detection under different parameter settings and compare it to LMM, we simulated a wide range

of networks. The simulations had networks with similar characteristics to those seen in networks

generated from microbiome datasets. These included networks with power law degree

distributions (N=175) with values of α, the exponent term of the power law formula ($y = kX^{-a}$),

varying between 1.8 and 2.6, as well as networks with regular degree distributions (N=200) with

p, the probability of one node being connected to another, varying from 0.001 to 0.2. The power

law and regular degree distribution networks were created using the NetworkX v2.6.3

implementations of configuration_model and erdos_renyi_graph, respectively, and all had a size

of 500. The networks with power law degree distributions had modularity values between 0.2

and 0.9, with higher α corresponding to higher modularity, and the networks with regular degree

distributions had modularity values between 0.07 and 0.98, with lower p corresponding to higher

modularity. Higher modularity scores indicate many connections within modules and fewer

connections between modules. We then calculated SMD and LMM partitions (with LMM

gamma = 1) of each network and compared the homogeneity between the two partitions.

Because SMD modules are smaller than LMM modules, we used the homogeneity metric

described by Rosenberg and Hirschberg [42] (implemented via Scikit-learn v0.24.2) to assess

whether nodes partitioned together by SMD are a subset of the module partitioned by LMM. A

score of 1 represents that all nodes in SMD modules represent sub-modules of LMM-partitioned

modules, whereas a score of 0 represents that no two nodes that were classified by SMD into the

same module were partitioned into a module together by the LMM method.

Demonstrating the use of SCNIC

We demonstrate the use of SCNIC with two example datasets. These are 1) a study that

used 16S rRNA sequencing of fecal material to compare microbiome composition in individuals

with and without HIV and in men who have sex with men (MSM) who were at a high risk of

contracting HIV [43], and 2) a dataset analyzing the microbiome of water samples at various

depths in two of the Great Lakes. We chose these two datasets so that we could evaluate

performance using datasets from both host-associated and free-living microbiomes. We also used

the Great Lakes dataset to compare module size and modularity between SMD and LMM

selected modules.

HIV dataset:

The HIV data set was retrieved from NCBI SRA accession number SRP068240, and

samples from the BCN0 cohort were used for these analyses. Reads were error corrected, quality

trimmed, and primers were removed using default parameters in BBTools [44]. DADA2 [45]

was used to define amplicon sequence variants (ASVs) with reads trimmed from the left by 30

base pairs and truncated at 269. ASVs were binned into operational taxonomic units (OTUs)

using USEARCH [46] at 99% identity using QIIME 1 [47]. A phylogenetic tree was made using

a single representative sequence from each OTU and the SEPP protocol [48, 49] using QIIME 2

[34]. We evaluated the average phylogenetic distance between OTUs in the same module using

the *distance* method of Biopython [50, 51]. Taxonomy was assigned using the Naive Bayes

QIIME 2 feature classifier, version gg-13-8-99-515-806-nb-classifier.qza.

The original study describing these data showed a strong divergence in gut microbiome

composition in MSM compared to non-MSM independent of HIV infection status and more

subtle differences associated with HIV infection when controlling for MSM behavior. The goal

of our analysis was to evaluate whether comparing gut microbiome composition between HIV

negative MSM and non-MSM with SCNIC modules provide additional significant taxa

compared to without, and additional insights as to which taxa that differ with MSM also are in

turn demonstrating co-correlated structure with each other. Co-correlation of microbes may

indicate that they are a part of a broader community type, interact with each other, or have shared

environmental drivers of their prevalence. A further goal of this analysis is to examine the effects

of using different R-value thresholds on the results. The SMD method was specifically used with

SparCC R-value thresholds between 0.20 and 1.0, with 0.05 increments.

Great Lakes dataset

The Great Lakes dataset was previously published as part of the Earth Microbiome

Project [52]. This study evaluated patterns of microbial relative abundance across depths in Lake

Michigan (N=16) and Lake Superior (N=33), with depth of samples collected ranging from 5 to

3654 meters. The study additionally recorded data on pH and salinity. The Great Lakes data set

was retrieved from QIITA accession number 1041 [53]. ASVs were found using DADA2 with a

left trim of 30 and a truncation length of 135. OTUs were subsequently picked on the ASVs

using VSEARCH [54] with a 99% identity threshold, resulting in 3,871 OTUs. These steps were

done with QIIME 2 [34]. SCNIC was applied with the SMD method and .2, .4 and .65 R-value

thresholds.

Comparison of SMD to LMM using the Great Lakes dataset

To identify differences in module structure from SMD versus LMM partitions, we

assessed the module size and modularity of 221 separately partitioned networks from the Great

Lakes dataset using varying parameters for SCNIC. The parameters included SCNIC R

thresholds ranging from 0.1 to 0.7 and gamma ranging from 0.15 to 0.9 for LMM.

Evaluating effects of applying SCNIC to discern microbes that differ between groups in the HIV

and Great Lakes datasets

OTUs/modules that differed with MSM status (HIV study) and between lakes (Great

Lakes study) were identified using ANCOM [28] for each feature. For the first study, we focused

on evaluating differences in the microbiome between MSM and non-MSM without confounding

by HIV infection status, by only using samples from HIV negative individuals. We chose

ANCOM because it is also a tool designed specifically for working with compositional

microbiome data. ANCOM was applied to the original feature table where SCNIC was not

applied, as well as to feature tables output from SCNIC using SparCC at different R-value

thresholds with the SMD algorithm.

While applying SparCC, SCNIC uses the recommended practice of the SparCC

manuscript of filtering based on average relative abundance across samples [38] .The SparCC

manuscript suggests this filter because removing features with high abundances, even in a few

samples, will upset the ability of the method to control for the number of reads per sample in its

compositionality adjustment. Because this method can retain OTUs that are highly abundant in

only a single sample, we removed features that had 0 values in more than 5% (~ 29/146) of

samples before applying ANCOM but after applying SparCC. Significant differences between

groups were determined as those above the W-value threshold determined by ANCOM.

Time and memory usage by SCNIC

To evaluate the memory resources needed by SCNIC, we ran the SCNIC modules step

locally on a 2015 MacBook Pro with 16 GB RAM with a 2.5 GHz Quad-Core Intel Core i7

processor for both the Great Lakes dataset and an integrated microbiome-metabolome dataset

with 1,301 features, which will be referred to as the GT dataset. The runtime was recorded across

3 runs per method (SMD vs LMM) for each dataset using GNU Time, and memory was profiled

using memory-profiler 0.60.0. The "within" step, which calculates correlations between features

and creates the correlation network was not tested because it depends greatly on the correlation

metric used, and the runtime and memory usage of FastSpar (likely the most computationally

intensive correlation metric to be used in this step) have already been profiled[39]. The modules

step only utilizes a correlation matrix and as such does not scale with the number of samples,

only the number of features, except when the values of the count table are being summed, which

is a generally trivial calculation compared to the module generation step.

RESULTS

Comparison of LMM to SMD on real and simulated data

In order to evaluate the relationship between modules detected with SMD versus LMM,

we chose modules on the Great Lakes dataset using SMD at R-value thresholds ranging from

0.05 to 0.7 and with LMM at the same R-value thresholds and gamma values ranging from 0.15

to 0.9. We found that with both LMM and SMD, modularity increased with increasing R-value

thresholds. However, SMD produced less modular partitions and smaller modules than LMM,

even when LMM was applied with very low values for the gamma parameter that controls

module size (Figure 3 A,B).

In order to determine whether SMD produced related modules to LMM (e.g. since SMD

modules are smaller, whether they represent sub-graphs of the larger LMM modules), we

calculated a homogeneity score (described in methods section) between SMD and LMM

modules in simulated networks. All networks contained 500 nodes. Modularity was calculated

for the LMM partitions, and the homogeneity of SMD and LMM partitions was calculated. From

our simulated networks, the homogeneity between SMD and LMM module partitions was

between 0.55 and 0.87 (Figure 3 C,D). Notably, we found that for networks simulated with both

power law and regular node degree distributions, as modularity of LMM partitions increased, the

homogeneity between SMD- and LMM-partitioned modules increased (power law network

Pearson $R = 0.87$, $p < 0.001$; regular network Pearson $R = 0.93$, $p < 0.001$). Thus, when network

modularity is high (i.e. there is a high number of edges within the module compared to between

modules), SMD partitions tend to be sub-partitions of LMM partitions.

R-value thresholds influence module size and phylogenetic relatedness of OTUs binned into a

module

A key parameter in SCNIC is the R-value threshold used to pick modules. Use of a high

R-value threshold would be expected to bin only very tightly correlated microbes with strong

relationships, while less stringent thresholds may identify community-level patterns representing

more loosely connected microbial pairs. To illustrate this concept, we binned OTUs into modules

using the SMD method at R-value thresholds between 0.2 and 1.0 using the HIV dataset. As

expected, at lower R-value thresholds, more OTUs were binned into modules and lower numbers

of modules of smaller average size were formed as the threshold increased (Figure 4A). To

illustrate the effects of R-values thresholds on the nature of the identified modules, we compare

SCNIC outputs using R-value thresholds of 0.2, 0.4, and 0.65. As shown in Figure 4, which

visualizes modules in Cytoscape using SCNIC output files, the R-value threshold influences the

size and connectivity of the network. We also illustrate the effect of using different thresholds by

examining the correlations between OTUs that are included in the first module output by SCNIC,

which is the largest module (module-*0*) (Figure 5). All OTUs in module-*0* are positively

correlated with each other, since SCNIC only captures positive correlations.

Microbes co-occurring in the same environmental niche have previously been observed

to be phylogenetically closer on average[4]. This is likely because phylogenetic relatedness has

been correlated with functional relatedness, such as through having more shared genome content,

leading towards success in similar environments [55]. We show that increasing the R-value

threshold results in modules that contain OTUs that are more phylogenetically similar on average

(Figure 5).

Use of SCNIC influences the detection of OTUs that differ between MSM and non-MSM

We next evaluated the effects of applying SCNIC with default SparCC and SMD

parameters and varying R-value thresholds on downstream statistical analysis results. To

investigate differential abundance based on MSM status in the HIV dataset we used

ANCOM[28]. After removing taxa that were not present in at least 5% of the samples the OTU

table had 317 samples and 639 OTUs. We found that 12 OTUs were significantly different

between MSM and non-MSM without using SCNIC. Using SCNIC at R-values of 0.2, 0.4, and

0.65 and running ANCOM on the filtered output feature table, we found that most of the

significant features were modules at an R-value of 0.2 and 0.4 but not 0.65 (e.g. 14 of the 15

significant features were modules at R=0.2) (Table 1). This was the case even though the vast

majority of OTUs were not a part of modules at the 0.4 R-value threshold (Figure 4A). The

majority of 12 of the OTUs that were significant without running SCNIC, were grouped into

modules with each other and with OTUs that were not individually significant without running

SCNIC. These significant modules contained 74, 26, and 1 new OTU at R-values of 0.2, 0.4 and

0.65 respectively. Using SCNIC also resulted in the identification of 1, 5 and 25 (at R-values of

0.2, 0.4 and 0.65) OTUs that were individually significant that were not significant without

running SCNIC, with no OTUs that were individually significant losing significance because

they were binned in a module, indicating an increase in statistical power resulting from running a

test like ANCOM that controls the FDR.

Considering correlation structure of significant features can help in understanding the

broader community context of bacteria that differ with MSM status. In module-*0* for each of the

R-values, which significantly differed by MSM status in all cases, *Prevotella* was the dominant

genus (Figure 5). At an R-value of 0.65, all OTUs in module-*0* were assigned to the genus

Prevotella (Figure 5C). However, at an R-value of 0.4 module-*0* included seven *Prevotella*

OTUs, one *Dialister*, and an unidentified member of the *Paraprevotellaceae* family. At the R-

value of 0.2, *Prevotella* accounted for 13 of the 25 OTUs and 11 of the 12 pre-SCNIC significant

OTUs were all found in this module. This suggests that individual OTUs that differ with MSM

status may in some cases be a part of a consortium of diverse members that collectively display

features that may contribute to differences in microbiome function.

To further explore this concept, we investigated the results generated with an R-value of

0.4, as the significant features maintain a strong level of correlation while being phylogenetically

diverse. When running ANCOM on this feature table, we found that these individually

significant OTUs tended to be joined into modules with other highly co-correlated microbes and

that these modules significantly differed with MSM (Figure 6). Of particular note, we observe

that the modules and taxa that are significantly related to MSM do not all correlate with each

other. At the R-value of 0.4, module-*36* contains two taxa, *Erysipelotrichaceae* and *Clostridium*

that are negatively correlated with the other significant taxa and modules (Figure 6). Module-*2*

contains *Eubacterium, Catenibacterium* and *Prevotella* which are phylogenetically heterogenous

but mutually co-occurring. A follow up experiment, which leverages insights that SCNIC

generates, may combine different strains of microbes to assemble a community type to test for

functional correlates of disease.

Use of SCNIC results in the detection of lake associated taxa

To test consistency in patterns across different datasets, we also tested the effects of

applying SCNIC with default SparCC and SMD parameters and varying R-value thresholds on

results from the Great Lakes dataset. Specifically, we identified features that significantly

differed between Lake Michigan (N=16) and Lake Superior (N=33) using ANCOM[28].

We began with a table of 3,871 OTUs, and 725 of these remained after removing OTUs

not present in at least 5% of the samples. We found that 168 OTUs were significantly different

between lakes without using SCNIC using ANCOM. Using SCNIC at R-values of 0.2, 0.4, and

0.65 and running ANCOM on the filtered output OTU table, we found that most significant

features were modules at an R threshold of 0.4 but not 0.2 or 0.65 (Table 1). Use of SCNIC

resulted in the detection of individual OTUs that were now significant that were not before (3

and 13 for R-value thresholds of 0.4 and 0.65 respectively). Application of SCNIC also

identified many additional OTUs that become of interest because they were now part of

significant modules (139, 64, and 12 OTUs at 0.2, 0.4, and 0.65 respectively; Table 1). However,

unlike for the HIV dataset, several OTUs that were individually significant were no longer

significant with ANCOM after applying SCNIC and this effect was the most pronounced with

lower R-value thresholds (64, 14, and 6 OTUs that were significant with SCNIC were no longer

significant after applying SCNIC at 0.2, 0.4, and 0.65 R-value thresholds respectively)(Table 1).

This is likely because microbes that differed between lakes were binned with loosely correlated

microbes that did not, leading to a loss of signal. Thus, in this case, only SCNIC with a moderate

to high R-value threshold appeared to balance the benefit of the increased power and

disadvantages of loss of signal from binning loosely correlated features.

Time and memory resources used by SCNIC

SCNIC's module generation step can be run locally on a laptop computer. For both the

Great Lakes and GT datasets, which had 764 and 1,301 features respectively, SCNIC's module

generation step ran in less than 2 minutes when using the SMD method and less than 30 seconds

when using the Louvain method. The Great Lakes dataset used less than 200 mebibytes (MiB)

memory, and the GT dataset required a maximum of 300 MiB memory. Supplemental table 1

shows time and memory usage during SCNIC.

DISCUSSION

SCNIC provides a method to measure correlations, find and visualize modules of

correlated features, and summarize modules by summing their counts for use in downstream

statistical analysis as one method for dimensionality reduction. Using SCNIC with the SMD

algorithm for module detection aids in feature reduction in 16S rRNA sequencing data while

ensuring a minimum strength of association within modules. As expected, our workflow

identified modules in which OTUs tended to be phylogenetically related, especially at relatively

high values of R. Using SCNIC, we overall achieved increased statistical power from performing

less comparisons, but use of low R-value thresholds had the potential to lead to loss of

significance by binning loosely correlated features. In this analysis, we used OTUs as features;

however, other microbiome features can be used with SCNIC, such as ASVs, genera, or species

defined with a taxonomic classifier, as well as other data types such as metabolome data. SCNIC

has also been used in previously published work to perform feature reduction prior to random

forest analysis with the microbiome and diverse other data types [56].

SCNIC complements existing methods because these either: 1) form correlation networks

of microbes for visualization but do not have functionality for selecting and summarizing

modules for downstream statistical analysis[32], 2) can select and summarize modules for

downstream statistical analysis but are designed for gene expression and not microbiome

data[18], only summarize features if they are phylogenetically related[31], or suggest methods

for finding modules of correlated microbes but do not provide a convenient implementation[30].

SCNIC is available both as a stand-alone application and as a QIIME 2 plugin for easy

integration with existing microbiome workflows.

SCNIC implements both the LMM algorithm, which had been previously recommended

for selecting modules of correlated microbes [25, 57], and a novel SMD algorithm. The

advantage of the SMD algorithm is that all pairs of features in the module have an R-value

greater than the user-provided minimum threshold. Using real and simulated data, we showed

that SMD produced smaller modules that generally represent sub-graphs of the larger LMM

modules. Since the use of lower R-value thresholds similarly produced larger modules including

more weakly correlated modules, we speculate that use of LMM might result in a similar trend of

identifying more OTUs within significant modules, but with the disadvantage of individually

significant OTUs being lost because they are combined with loosely correlated microbes that are

not related to the outcome being tested.

We illustrate here that varying the R-value threshold input by the user has a great impact

on the results. However, we have avoided giving specific R-value threshold recommendations

here, because optimal R-values may vary across datasets and data types. Using higher R-values

thresholds was more likely to identify highly phylogenetically related microbes that likely share

overlapping functionality, and in principle could also identify diverse organisms with

overlapping niches or highly complementary metabolic functions. Using a lower R-value

threshold bins a broader community of more loosely correlated features with the risk of bringing

together features which should not be grouped and loosing significance of OTUs – as was

illustrated in the Great Lakes dataset analysis conducted here. By summarizing correlated

features, SCNIC mitigates overcorrection in multiple test adjustments by reducing the number of

taxa and false discovery rate for downstream analysis. When these organisms are grouped into a

broader module that is truly independent from other modules, any penalties on two highly similar

features may be avoided in statistical analysis.

The results of our HIV dataset analysis confirm original findings, as well as those of

another study[58], but included many new significantly associated taxa. SCNIC also assists in

interpretation of microbiome data by identifying correlations among these taxa. Our results

recapitulated those of the original publication of these data and previous HIV microbiome

studies that all found enrichment of *Prevotella* with MSM status [43, 58-60]. However, our

analyses provide additional insight by identifying correlations between differentiating taxa. For

instance, in module-*0*, which was more abundant in MSM samples, OTUs assigned

taxonomically to the *Prevotella* genus are correlated with two OTUs identified as *Eubacterium*

biforme (which has recently been renamed *Holdemanella biformis [61]*). *Prevotella copri* has

previously been associated with increased inflammation [59] while *in vitro* stimulations of

human immune cells have found that *P. copri* did not induce particularly high levels of

inflammation but *E. biforme* did [60]. This strong correlation between *P. copri* and *E. biforme* in

MSM could explain the increased inflammation seen in individuals with higher levels of *P.*

copri, with *E. biforme* being the true driver. Indeed, MSM status has previously been associated

with increased inflammation [62, 63]. With the use of SCNIC, this correlation highlighted a

route of mechanistic understanding which could be functionally followed up on in further

experimental studies.

SCNIC detected multiple significant modules, of which none of the OTUs within were

significant when analyzed separately. Module-*20*, which was associated with MSM status, is the

fourth most significant feature at R-value of 0.2, and is made up of *Acidaminococcus*,

Megasphaera, and *Mitsuokella* species. These are all from the *Veillonellaceae* family which is

likely the explanation for their correlation. Members of the *Veillonellaceae* family have been

linked with inflammation [64].

By increasing statistical power and providing context for the relationships between

significant taxa, SCNIC modules open new opportunities for analysis. When a module is

associated with a variable of interest, the correlations within the module may imply functional

relationships. These can be further investigated with *in vitro* and *in vivo* experiments. Studies

which aim to test hypotheses generated from correlative analysis will commonly use a single

significantly associated microbes. This often does not adequately represent *in vivo* systems

because microbes in isolation often do not affect a disease state or their environment. SCNIC can

enhance these confirmatory studies by identifying groups of microbes that may grow better than

individual microbes and may better elicit relevant phenotypes than when grown separately.

ACKNOWLEDGEMENTS

We would like to thank Elmar Pruesse for input on the design of SCNIC. We thank

Jennifer Fouquier, Abigail Armstrong and Casey Martin for beta testing SCNIC. We thank

Jennifer Fouquier for reading and commenting on the manuscript. Funding for KT came from the

University of Colorado School of Medicine Research Track. Funding for MS came from NIH

NLM 4 T15 LM009451-10.

REFERENCES

1. Corno, G., J. Villiger, and J. Pernthaler, *Coaggregation in a microbial predator–prey system affects competition and trophic transfer efficiency.* Ecology, 2013. **94**(4): p. 870-881.

2. Burkepile, D.E., et al., *Chemically mediated competition between microbes and animals: microbes as consumers in food webs.* Ecology, 2006. **87**(11): p. 2821-2831.

3. LaSarre, B., et al., *Microbial mutualism dynamics governed by dose-dependent toxicity of cross-fed nutrients.* The ISME journal, 2017. **11**(2): p. 337-348.

4. Faust, K., et al., *Microbial co-occurrence relationships in the human microbiome.* PLoS Comput. Biol., 2012. **8**(7): p. e1002606.

5. Barberán, A., et al., *Using network analysis to explore co-occurrence patterns in soil microbial communities.* The ISME journal, 2012. **6**(2): p. 343-351.

6. Widder, S., et al., *Fluvial network organization imprints on microbial co-occurrence networks.* Proc. Natl. Acad. Sci. U. S. A., 2014. **111**(35): p. 12799-12804.

7. Bray, J.R. and J.T. Curtis, *An Ordination of the Upland Forest Communities of Southern Wisconsin.* Ecological Monographs, 1957. **27**(4): p. 325-349.

8. Pearson, K., *DETERMINATION OF THE COEFFICIENT OF CORRELATION.* Science, 1909. **30**(757): p. 23-25.

9. Spearman, C., *Measurement of association, Part II. Correction of 'systematic deviations'.* Am. J. Psychol, 1904. **15**: p. 88-101.

10. Haas, B.J., et al., *Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons.* Genome Research, 2011. **21**(3): p. 494-504.

11. Huse, S.M., et al., *Ironing out the wrinkles in the rare biosphere through improved OTU clustering.* Environmental Microbiology, 2010. **12**(7): p. 1889-1898.

12. Kuczynski, J., et al., *Microbial community resemblance methods differ in their ability to*

detect biologically relevant patterns. Nat. Methods, 2010. **7**(10): p. 813-819.

13. Weiss, S., et al., *Correlation detection strategies in microbial data sets vary widely in*

sensitivity and precision. ISME J., 2016. **10**(7): p. 1669-1681.

14. Ban, Y., L. An, and H. Jiang, *Investigating microbial co-occurrence patterns based on*

metagenomic compositional data. Bioinformatics, 2015. **31**(20): p. 3322-3329.

15. Dugas, L.R., et al., *Decreased microbial co-occurrence network stability and SCFA receptor*

level correlates with obesity in African-origin women. Sci. Rep., 2018. **8**(1): p. 17135.

16. Lozupone, C., et al., *Identifying genomic and metabolic features that can underlie early*

successional and opportunistic lifestyles of human gut symbionts. Genome Research, 2012.

22(10): p. 1974-1984.

17. Benjamini, Y. and Y. Hochberg, *On the Adaptive Control of the False Discovery Rate in*

Multiple Testing With Independent Statistics. J. Educ. Behav. Stat., 2000. **25**(1): p. 60-83.

18. Langfelder, P. and S. Horvath, *WGCNA: an R package for weighted correlation network*

analysis. BMC Bioinformatics, 2008. **9**: p. 559.

19. Wilcox, R.R., *Introduction to Robust Estimation and Hypothesis Testing*. 2011: Academic

Press.

20. Barabási, A.-L. and R. Albert, *Emergence of Scaling in Random Networks.* Science, 1999.

286(5439): p. 509-512.

21. Castillo, J.D., J.M. Vivanco, and D.K. Manter, *Bacterial Microbiome and Nematode*

Occurrence in Different Potato Agricultural Soils. Microb. Ecol., 2017. **74**(4): p. 888-900.

22. Tong, M., et al., *Reprograming of gut microbiome energy metabolism by the FUT2 Crohn's*

disease risk polymorphism. The ISME Journal, 2014. **8**(11): p. 2193-2206.

23. Yin, J., et al., *Lysine Restriction Affects Feed Intake and Amino Acid Metabolism via Gut*

Microbiome in Piglets. Cell. Physiol. Biochem., 2017. **44**(5): p. 1749-1761.

24. Younge, N., Q. Yang, and P.C. Seed, *Enteral High Fat-Polyunsaturated Fatty Acid Blend*

Alters the Pathogen Composition of the Intestinal Microbiome in Premature Infants with an

Enterostomy. J. Pediatr., 2017. **181**: p. 93-101.e6.

25. Jackson, M.A., et al., *Detection of stable community structures within gut microbiota co-*

occurrence networks from different human populations. PeerJ, 2018. **6**: p. e4303.

26. Gloor, G.B., et al., *Microbiome Datasets Are Compositional: And This Is Not Optional.*

Frontiers in Microbiology, 2017. **8**.

27. Broido, A.D. and A. Clauset, *Scale-free networks are rare.* Nat. Commun., 2019. **10**(1): p.

1017.

28. Mandal, S., et al., *Analysis of composition of microbiomes: a novel method for studying*

microbial composition. Microb. Ecol. Health Dis., 2015. **26**: p. 27663.

29. Paulson, J.N., et al., *Differential abundance analysis for microbial marker-gene surveys.*

Nat. Methods, 2013. **10**(12): p. 1200-1202.

30. Blondel, V.D., et al., *Fast unfolding of communities in large networks.* Journal of Statistical

Mechanics: Theory and Experiment, 2008. **2008**(10): p. P10008.

31. Frøslev, T.G., et al., *Algorithm for post-clustering curation of DNA amplicon data yields*

reliable biodiversity estimates. Nature Communications, 2017. **8**(1): p. 1188.

32. Faust, K. and J. Raes, *CoNet app: inference of biological association networks using*

Cytoscape. F1000Research, 2016. **5**: p. 1519.

33. Grüning, B., et al., *Bioconda: sustainable and comprehensive software distribution for the*

life sciences. Nature Methods, 2018. **15**(7): p. 475-476.

34. Bolyen, E., et al., *Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.* Nature Biotechnology, 2019. **37**(8): p. 852-857.

35. Himsolt, M., *GML: A portable graph file format.* 1997, Technical report, Universitat Passau.

36. Shannon, P., et al., *Cytoscape: a software environment for integrated models of biomolecular interaction networks.* Genome Res., 2003. **13**(11): p. 2498-2504.

37. McDonald, D., et al., *The Biological Observation Matrix (BIOM) format or: how I learned to stop worrying and love the ome-ome.* Gigascience, 2012. **1**(1): p. 7.

38. Friedman, J. and E.J. Alm, *Inferring correlation networks from genomic survey data.* PLoS Comput. Biol., 2012. **8**(9): p. e1002687.

39. Watts, S.C., et al., *FastSpar: rapid and scalable correlation estimation for compositional data.* Bioinformatics, 2019. **35**(6): p. 1064-1066.

40. Fernandes, A.D., et al., *Unifying the analysis of high-throughput sequencing datasets: characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by compositional data analysis.* Microbiome, 2014. **2**(1): p. 15.

41. Tsilimigras, M.C.B. and A.A. Fodor, *Compositional data analysis of the microbiome: fundamentals, tools, and challenges.* Annals of Epidemiology, 2016. **26**(5): p. 330-335.

42. Rosenberg, A. and J. Hirschberg. *V-Measure: A Conditional Entropy-Based External Cluster Evaluation Measure.* 2007. Prague, Czech Republic: Association for Computational Linguistics.

43. Noguera-Julian, M., et al., *Gut Microbiota Linked to Sexual Preference and HIV Infection.* EBioMedicine, 2016. **5**: p. 135-146.

44. Bushnell, B., J. Rood, and E. Singer, *BBMerge – Accurate paired shotgun read merging via overlap.* PLOS ONE, 2017. **12**(10): p. e0185056.

45. Callahan, B.J., et al., *DADA2: High-resolution sample inference from Illumina amplicon*

data. Nat Methods, 2016. **13**(7): p. 581-3.

46. Edgar, R.C., *Search and clustering orders of magnitude faster than BLAST.* Bioinformatics,

2010. **26**(19): p. 2460-2461.

47. Caporaso, J.G., et al., *QIIME allows analysis of high-throughput community sequencing*

data. Nat. Methods, 2010. **7**(5): p. 335-336.

48. Janssen, S., et al., *Phylogenetic Placement of Exact Amplicon Sequences Improves*

Associations with Clinical Information. mSystems, 2018. **3**(3).

49. Mirarab, S., N. Nguyen, and T. Warnow, *SEPP: SATé-enabled phylogenetic placement.* Pac.

Symp. Biocomput., 2012: p. 247-258.

50. Cock, P.J.A., et al., *Biopython: freely available Python tools for computational molecular*

biology and bioinformatics. Bioinformatics, 2009. **25**(11): p. 1422-1423.

51. Talevich, E., et al., *Bio.Phylo: A unified toolkit for processing, analyzing and visualizing*

phylogenetic trees in Biopython. BMC Bioinformatics, 2012. **13**(1): p. 209.

52. Thompson, L.R., et al., *A communal catalogue reveals Earth's multiscale microbial*

diversity. Nature, 2017. **551**(7681): p. 457-463.

53. Gonzalez, A., et al., *Qiita: rapid, web-enabled microbiome meta-analysis.* Nat. Methods,

2018. **15**(10): p. 796-798.

54. Rognes, T., et al., *VSEARCH: a versatile open source tool for metagenomics.* PeerJ, 2016. **4**:

p. e2584.

55. Zaneveld, J.R., et al., *Ribosomal RNA diversity predicts genome diversity in gut bacteria and*

their relatives. Nucleic Acids Research, 2010. **38**(12): p. 3869-3879.

56. Armstrong, A.J.S., et al., *Systems Analysis of Gut Microbiome Influence on Metabolic*

Disease in HIV-Positive and High-Risk Populations. mSystems, 2021. **6**(3).

57. Baldassano, S.N. and D.S. Bassett, *Topological distortion and reorganized modular*

structure of gut microbial co-occurrence networks in inflammatory bowel disease. Sci Rep,

2016. **6**: p. 26087.

58. Armstrong, A.J.S., et al., *An exploration of Prevotella-rich microbiomes in HIV and men*

who have sex with men. Microbiome, 2018. **6**(1): p. 198.

59. Dillon, S.M., et al., *An altered intestinal mucosal microbiome in HIV-1 infection is*

associated with mucosal and systemic immune activation and endotoxemia. Mucosal

Immunology, 2014. **7**(4): p. 983-994.

60. Lozupone, C.A., et al., *Alterations in the gut microbiota associated with HIV-1 infection.*

Cell Host Microbe, 2013. **14**(3): p. 329-339.

61. De Maesschalck, C., et al., *Faecalicoccus acidiformans gen. nov., sp. nov., isolated from the*

chicken caecum, and reclassification of Streptococcus pleomorphus (Barnes et al. 1977),

Eubacterium biforme (Eggerth 1935) and Eubacterium cylindroides (Cato et al. 1974) as

Faecalicoccus pleomorphus comb. nov., Holdemanella biformis gen. nov., comb. nov. and

Faecalitalea cylindroides gen. nov., comb. nov., respectively, within the family

Erysipelotrichaceae. International Journal of Systematic and Evolutionary Microbiology,

2014. **64**(Pt_11): p. 3877-3884.

62. Gianella, S., et al., *Associations between virologic and immunologic dynamics in blood and*

in the male genital tract. J. Virol., 2012. **86**(3): p. 1307-1315.

63. Palmer, C.D., et al., *Enhanced immune activation linked to endotoxemia in HIV-1*

seronegative MSM. AIDS, 2014. **28**(14): p. 2162-2166.

64. Bajaj, J.S., et al., *Linkage of gut microbiome with cognition in hepatic encephalopathy.*

American Journal of Physiology-Gastrointestinal and Liver Physiology, 2012. **302**(1): p.

G168-G175.

DATA ACCESSIBILITY

The Noguera-Julian et al. data set is available from NCBI SRA accession number

SRP068240. The lakes dataset is available from QIITA accession number 1041.

AUTHOR CONTRIBUTIONS

M.S. coded the initial implementation of SCNIC and made major contributions to its

conceptualization and design. K.T. improved the SCNIC implementation and performed the HIV

case study. J.S. performed the comparisons of SMD to LMM with real and simulated data. C.L.

conceptualized SCNIC and guided its implementation and design. K.T., M.S., J.S. and C.L. all

wrote the manuscript together.

A)

635
636 **B)**

637
638

639
640
Figure 1: SCNIC Schematic and Data Flow

A. The basic process of SCNIC involves first identifying pairwise correlations between species
and using them to build a correlation network. Modules of correlated features are identified and
then summarized for downstream statistical analysis, or multi-omic analysis between modules of
microbes and other feature types. B. The input to SCNIC comes in the form of a count table in
BIOM format. The first step takes the table and generates a correlation table and network. The
table is in a tab delimited format and the network is in GML format and can be used to visualize
the network in Cytoscape. Modules are detected and summarized in the final step which
generates a module membership file indicating which features are in each module. The collapsed
BIOM table contains the same total counts per sample as the original table, but with less features.
All features not included in modules are retained with their original counts and all modules have
a total count per sample of the sum of the counts of all features in that module.

653
654
Figure 2: SMD Algorithm for Defining Modules

656
657 SMD defines a module as a group where all features have a correlation above a given threshold.
658 To do this SMD first uses complete linkage hierarchical clustering on correlation coefficients to
659 create a tree. Each module is defined as a subtree where the correlation coefficients between all
660 tips are greater than the threshold.

661
662
Figure 3: Comparison of the SMD and LMM Algorithms for Module Selection
Panels A and B show a comparison of module size and modularity between SMD and LMM
selected modules on the Great Lakes data at R thresholds ranging from 0.1 to 0.7 and gamma
ranging from 0.15 and 0.9 for LMM. Panel A compares average module size and Panel B shows
modularity across these parameters. In Panels C and D the homogeneity of LMM versus SMD
selected module was calculated using simulated networks formed using both Power-Law (Panel
C) and Regular (Panel D) degree distributions. Nodes are colored by the α (exponent) parameter
for power law degree distributions in Panel C and by p (probability of one node being connected
to any other node) in regular degree distributions in Panel D. A homogeneity of 1 denotes that all
SMD modules are sub-modules of the LMM modules, whereas a homogeneity of 0 denotes that
no two nodes grouped into a SMD module were partitioned into the same module by LMM.

Figure 4: SCNIC Feature Reduction and Visualization of SCNIC Networks

A) We used SCNIC to select modules using the OTU table from the HIV dataset, the SMD
module selection algorithm, and SparCC R-values ranging from 0.2 to 1.0, in increments of 0.05.
The R-value is plotted against the number of features in the resulting BIOM table produced by
SCNIC. As the R-value increases the number of modules decreases and the number of single
features (modules + OTUs not included in modules) increases. After the R-value of 0.65, the
number of features in the resulting file remained the same at 4351 features, because there were
no modules that were created past a SparCC R of 0.65.The Cytoscape output allows for easy
exploration and visualization of relationships between different OTUS/taxa in an interactive
interface. B) R = 0.65 C) R = 0.4 D) R = 0.2. As the R-value increases, the size of the network
decreases as SCNIC does not include singletons (features with no significant positive
correlations) in the resulting network file. Correlation network where edges are correlations with
a R-value greater than the threshold set. Nodes are OTUs and node color represents module
membership (i.e. module-*0* is pink in Panel B).
689

690
691 **Figure 5: Module-*0* Across Different R-Values**
692
Module-*0* expanded to view taxonomy and correlations amongst them at R-values of 0.2 (A), 0.4
(B), and 0.65 (C). The heatmap in the lower triangle corresponds to the correlation found by
SparCC colored on a light red (low correlation) to dark red (high correlation) spectrum as
defined in the color bar on the right. The heatmap in the upper triangle represents the
phylogenetic distance between organism pairs colored on a yellow (small phylogenetic distance)
to dark blue (high phylogenetic distance) spectrum as defined in the color bar on the right.
As the R-value increases, the species in module-*0* become more phylogenetically similar.
Module-*0* has 11, 5 and 2 of the significant Pre-SCNIC OTUs at R-values of 0.2, 0.4 and 0.65,
and are highlighted in a yellow border.

Figure 6: All Significant Features at R-value 0.4 Found by ANCOM

Each of the borders in the y-axis represents the different modules, with the module number
bolded. The Pre-SCNIC OTUs that were significant are highlighted in a yellow border. The
heatmap in the lower triangle corresponds to the correlation found by SparCC colored on a light
red (low correlation) to dark red (high correlation) spectrum as defined in the color bar on the
right. The heatmap in the upper triangle represents the phylogenetic distance between organism
pairs colored on a yellow (small phylogenetic distance) to dark blue (high phylogenetic distance)
spectrum as defined in the color bar on the right.

712   HIV dataset

R-Value	New OTUs in Sig. Modules	New Significant OTUs	Lost Significant OTUs	# of Significant Modules	Total Significant Features
0.2	74	1	0	14	15
0.4	26	5	0	11	17
0.65	1	25	0	2	35

714   Great Lakes dataset

R-Value	New OTUs in Sig. Modules	New Significant OTUs	Lost Significant OTUs	# of Significant Modules	Total Significant Features
0.2	139	0	64	1	25
0.4	64	3	14	29	33
0.65	12	13	6	24	74
R-Value	New OTUs in Sig. Modules	New Significant OTUs	Lost Significant OTUs	# of Significant Modules	Total Significant Features
0.2	74	1	0	14	15
0.4	26	5	0	11	17
0.65	1	25	0	2	35

716   **Table 1: Significant SCNIC Modules and Features Across R-Values in the HIV and Great**

717   **Lakes Datasets**

718   ANCOM analysis of the HIV and Great Lakes datasets after using SCNIC at R-Value thresholds
719   at 0.2, 0.4, and 0.65. MSM was used as the categorical variable for differential abundance in the
720   HIV analysis and the Great Lakes analysis tested for taxa that differed between Lakes.
721
722
723

Dataset	Features	Time 1	Time 2	Time 3	Max Memory
Lakes (SMD)	764	0 m 31 s	0 m 30 s	0 m 32 s	190 MiB
Lakes (Louvain)	764	0 m 7 s	0 m 7 s	0 m 7 s	190 MiB
GT (SMD)	1,301	1 m 28 s	1 m 27 s	1 m 27 s	302 MiB
GT (SMD + table)	1,301	1 m 45 s	1 m 32 s	1 m 52 s	301 MiB
GT (Louvain)	1,301	0 m 20 s	0 m 18 s	0 m 16 s	299 MiB
GT (Louvain + table)	1,301	0 m 15 s	0 m 15 s	0 m 16 s	301 MiB

Supplemental Table 1: Runtime and memory usage by SCNIC module generation step

Modules were calculated based on correlation matrices for the Great Lakes dataset and an
integrated metabolomics-microbiome (GT) dataset. Runtime was profiled using GNU Time
across 3 runs per dataset and method, and maximum memory usage in mebibytes (MiB) was
profiled using memory-profiler. A count table to sum module relative abundances was passed to
SCNIC for the GT dataset when denoted in the leftmost column.