In Vitro Activity of Itraconazole Against SARS-CoV-2

Ellen Van Damme PhD[1], Sandra De Meyer PhD[1], Denisa Bojkova MSc[2], Sandra Ciesek MD[2,4],

Jindrich Cinatl PhD[2], Steven De Jonghe PhD[3], Dirk Jochmans PhD[3], Pieter Leyssen PhD[3],

Christophe Buyck MEng[1], Johan Neyts PhD[3], Marnix Van Loock PhD[1]*

[1]Janssen Pharmaceutica NV, Beerse, Belgium

[2]Institute of Medical Virology, University Hospital Frankfurt, Goethe University, Frankfurt am

Main, Germany

[3]KU Leuven, Department of Microbiology, Immunology and Transplantation, Rega Institute,

Laboratory of Virology and Chemotherapy, Leuven, Belgium

[4]German Center for Infection Research (DZIF), Frankfurt, Germany

*Corresponding Author: mvloock@its.jnj.com

15
Address for Correspondence

Marnix Van Loock, PhD,

Infectious Diseases & Vaccines Discovery

Janssen Pharmaceutica NV,

Turnhoutseweg 30,

2340 Beerse

Belgium

mvloock@its.jnj.com

Highlights

• Itraconazole exerted *in vitro* low micromolar activity against SARS-CoV-2 (EC_{50} = 2.3 µM)

• Remdesivir demonstrated potent antiviral activity, confirming validity of the assay

• Itraconazole has since shown no efficacy in a clinical study in hospitalized COVID-19

patients

Abstract

31
Background: As long as there is no vaccine available, having access to inhibitors of SARS-

CoV-2 will be of utmost importance. Antivirals against coronaviruses do not exist, hence global

drug re-purposing efforts have been carried out to identify agents that may provide clinical

benefit to patients with COVID-19. Itraconazole, an antifungal agent, has been reported to have

potential activity against animal coronaviruses.

Methods: Using cell-based phenotypic assays, the *in vitro* antiviral activity of itraconazole and

17-OH itraconazole was assessed against clinical isolates from a German and Belgian patient

infected with SARS-CoV-2.

Results: Itraconazole demonstrated antiviral activity in human Caco-2 cells (EC_{50} = 2.3 µM;

MTT assay). Similarly, its primary metabolite, 17-OH itraconazole, showed inhibition of SARS-

CoV-2 activity (EC_{50} = 3.6 µM). Remdesivir inhibited viral replication with an EC_{50} = 0.4 µM.

Itraconazole and 17-OH itraconazole resulted in a viral yield reduction *in vitro* of approximately

$2\text{-}log_{10}$ and approximately $1\text{-}log_{10}$, as measured in both Caco-2 cells and VeroE6-eGFP cells,

respectively. The viral yield reduction brought about by remdesivir or GS-441524 (parent

nucleoside of the antiviral prodrug remdesivir; positive control) was more pronounced, with an

approximately $3 \, log_{10}$ drop and $>4 \, log_{10}$ drop in Caco-2 cells and VeroE6-eGFP cells,

respectively.

Discussion: Itraconazole and 17-OH itraconazole exert *in vitro* low micromolar activity against

SARS-CoV-2. Despite the *in vitro* antiviral activity, itraconazole did not result in a beneficial

effect in hospitalized COVID-19 patients in a clinical study (EudraCT Number: 2020-001243-15).

Introduction

The rapid spread of the severe acute respiratory syndrome-associated coronavirus-2 (SARS-CoV-2) disease (COVID-19) has been declared a global pandemic.[1,2] SARS-CoV-2 is a beta-coronavirus, which are enveloped viruses containing single-strand, positive-sense RNA.[3] Since COVID-19 emerged in humans in late December 2019,[4] aside from a significant economic loss, more than a million death have been reported worldwide, driving urgently the need to identify potential vaccines and therapies.

Given that therapeutic options for antiviral treatment of SARS-CoV-2 remain limited, research has initially focused on re-purposing available drugs that have demonstrated antiviral activity against coronaviruses. As part of such an initiative, itraconazole was identified as a potential candidate. Itraconazole has been described previously to have activity in an *in vitro* screen using a luciferase reporter-expressing recombinant murine betacoronavirus,[5] as well as against a feline alphacoronavirus that causes feline infectious peritonitis (FIP).[6] Furthermore, *in vitro* and *in vivo* activity has been described against other respiratory viruses, such as influenza A and human rhinovirus.[7,8]

Itraconazole is a member of the triazole group of broad-spectrum antifungals,[9] with a well-established efficacy and safety profile.[8–12] The primary mechanism of its antifungal action is the inhibition of ergosterol biosynthesis, by acting on the oxysterol-binding protein (OSBP).[13–16] Since ergosterol is closely related to cholesterol, it has been suggested that also in mammalian cells, itraconazole acts as a cholesterol trafficking inhibitor resulting in disruption of the cholesterol-enriched membranes and thus inhibition of virus replication.[6,8,17] However, a role in interferon priming has also been suggested as a contributing factor to the antiviral activity.[18]

In an attempt to identify therapeutic options for treating COVID-19 patients, the *in vitro* antiviral

activities of itraconazole, and its metabolite 17-OH itraconazole were investigated in Caco-2 and

VeroE6-eGFP cells infected with SARS-CoV-2, isolated from COVID-19 patients.

Methods

Cell culture

Caco-2 cells (human colon carcinoma cell line; obtained from the Deutsche Sammlung von

Mikroorganismen und Zellkulturen, Braunschweig, Germany) were cultured in Minimal Essential

Medium (MEM) supplemented with 10% fetal bovine serum (FBS) with penicillin (100 IU/mL)

and streptomycin (100 µg/mL) at 37°C in a 5% CO_2 atmosphere. VeroE6-eGFP (African green

monkey kidney cell line; provided by Dr. K. Andries J&JPRD; Beerse, Belgium) were cultured in

Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS, 0.075% sodium

bicarbonate, penicillin and streptomycin (100 µg/mL) at 37°C in a 5% CO_2 atmosphere. All cell

culture reagents were obtained from Sigma-Aldrich (Hamburg, Germany).

SARS-CoV-2 Preparation

SARS-CoV-2-FFM1 (strain hCoV-19/Germany/FrankfurtFFM1/2020) was isolated from a

German human-case sample and cultured in Caco-2 cells, as previously described.[1,19]

SARS-CoV-2-Germany stocks were passaged twice in Caco-2 cells prior to storage (–80°C).

SARS-CoV-2-Belgium (strain BetaCov/Belgium/GHB-03021/2020) was recovered from a

nasopharyngeal swab taken from a patient returning from Wuhan, China. SARS-CoV-2 Belgium

stocks were passaged six times in VeroE6-eGFP cells prior to storage (-80°C).

Assessment of Antiviral Activity

Itraconazole and its metabolite, 17-OH itraconazole, and remdesivir were synthesized at

Johnson and Johnson. GS-441524, the parent nucleoside of remdesivir, used in studies at KU

Leuven was obtained from MedChemExpress (NJ, USA). Antiviral activity was assessed by

inhibition of virus-induced cytopathogenic effect (CPE) as described previously.[20] In brief,

confluent layers of Caco-2 cells cultured for 72 hours on 96 multi-well plates (50,000 cells/well)

were challenged with SARS-CoV-2 FFM1 at a multiplicity of infection of 0.01. The virus was

added together with the compound under investigation and incubated in MEM supplemented

with 1% FBS. Itraconazole diluted in MEM without FBS in 4-fold dilutions was added to a

concentration range of 0.01 µM to 50 µM; 17-OH itraconazole diluted in MEM without FBS was

added in 4-fold dilutions to a concentration range of 0.02 µM to 100 µM and remdesivir diluted in

MEM without FBS in 4-fold dilutions was added to a concentration range of 0.02 µM to 100 µM.

Cells were incubated for 48 hours; the CPE was then visually scored by two independent

laboratory technicians. In addition, CPE was also assessed using a 3-(4,5-dimethylthiazol-2-yl)-

2,5-diphenyltetrazolium bromide (MTT) assay, performed according to the manufacturer's

instructions. Optical densities were measured at 560/620 nm in a Multiskan Reader (MCC/340

Labsystems). Two series of three independent experiments (or one series of three experiments

for 17OH-itraconazole), each containing three replicates, were performed. Data were analyzed

by four-parameter curve-fitting from a dose-response curve using GraphPad Prism (version

7.00) to calculate the EC_{50} (concentration of the compound that inhibited 50% of the infection)

based on visual CPE scoring or the MTT assay.

Assessment of Cell Viability

Cell viability in Caco-2 cells was measured following administration of each of the compounds or

metabolites under investigation over the range of concentrations in the absence of virus using

the Rotitest Vital (Roth, Karlsruhe, Germany) test according to manufacturer's instructions, as

previously described.[21] All assays were performed three times independently in triplicate, this

was performed twice (once for 17OH-itraconazole). Data were analyzed by four-parameter

curve-fitting from a dose-response curve using GraphPad Prism (version 7.00) to calculate the

CC_{50} (cytotoxic concentration of the compound that reduced cell viability to 50%).

Viral RNA Yield Reduction Assay

Antiviral activity was assessed by inhibition of viral yield in Caco-2 cells. Confluent layers of

Caco-2 cells cultured for 72 hours on 96 multi-well plates (50,000 cells/well) were challenged with

SARS-CoV-2-FFM1 at a multiplicity of infection of 0.01. The virus was added together with the

compound under investigation and incubated in MEM supplemented with 1% FBS. Itraconazole

diluted in MEM without FBS was added in 4-fold dilutions to a concentration range of 0.01 µM to

50 µM; 17-OH itraconazole diluted in MEM without FBS was added in 4-fold dilutions to a

concentration range of 0.02 µM to 100 µM and remdesivir diluted in MEM without FBS was added

in 4-fold dilutions to a concentration range of 0.02 µM to 100 µM. Two series of three independent

experiments, each containing two replicates, were performed.

SARS-CoV-2 RNA from cell culture supernatant samples was isolated using AVL buffer and the

QIAamp Viral RNA Kit (Qiagen) according to the manufacturer's instructions. Absorbance-based

quantification of the RNA yield was performed using the Genesys 10S UV-Vis Spectrophotometer

(Thermo Scientific). RNA was subjected to OneStep qRT-PCR analysis using the Luna Universal

One-Step RT-qPCR Kit (New England Biolabs) and a CFX96 Real-Time System, C1000 Touch

Thermal Cycler (BioRad). Primers were adapted from the WHO protocol targeting the open

reading frame for RNA-dependent RNA polymerase (RdRp): RdRP_SARSr-F2 (GTG ARA TGG

TCA TGT GTG GCG G) and RdRP_SARSr-R1 (CAR ATG TTA AAS ACA CTA TTA GCA TA)

using 0.4 µM per reaction. Standard curves were created using plasmid DNA (pEX-A128-RdRP)

harboring the corresponding amplicon regions for RdRP target sequence according to GenBank

Accession number NC_045512.

Antiviral activity was also assessed by reduction of viral yield in VeroE6-eGFP cells. VeroE6-

eGFP cells were cultured for 24 hours on 96 multi-well plates (10,000 cells/well) in the absence

or presence of compound and infected with SARS-CoV-2-Belgium at a multiplicity of infection of

1 Tissue Culture Infectious Dose ($TCID_{50}$)/cell, in 200 µL assay medium (DMEM supplemented

with 2% FBS and 0.075% sodium bicarbonate). After 2 hours at 37°C the cells were washed

once and further cultured in 200 µL assay medium containing the same compound

concentrations (37°C, 5% CO_2) for another 48 hours. The viral RNA (vRNA) in the culture

supernatant (SN) was extracted using the NucleoSpin kit (Macherey-Nagel), according to the

manufacturer's instructions and quantified by real-time quantitative polymerase chain reaction

(RT-qPCR) performed on a LightCycler96 platform (Roche) using the iTaq Universal Probes

One-Step RT-qPCR kit (BioRad) with primers and probes specific for SARS-CoV-2. A standard

curve was created from serial dilutions of a virus stock with known titer and used to correlate the

cycle threshold (Ct) values of the experimental samples with absolute virus quantities. One

independent experiment, containing two replicates (for itraconazole) or three replicates (17OH-

Itraconazole), were performed.

Results

In Vitro **Antiviral Activity in Caco-2 cells**

Independent experiments with triplicate measurements were performed with itraconazole (n=6),

17-OH itraconazole (n=3) and remdesivir (n=6). In Caco-2 cells, itraconazole resulted in a dose-

dependent inhibition of SARS-CoV-2-FFM1 measured by visual scoring of inhibition of CPE with

an EC_{50} = 1.5 µM (Figure 1). Similar findings were obtained using the MTT assay (EC_{50} = 2.3

µM). The 17-OH itraconazole metabolite reduced SARS-CoV-2 induced CPE with an $EC_{50\ visual}$

of 1.2 µM and $EC_{50\ MTT}$ = 3.6 µM. The positive control, remdesivir, resulted in EC_{50} values of 0.3

and 0.4 µM in the CPE and MTT assays in Caco-2 cells, respectively.

Minimal cytotoxicity was seen in Caco-2 cells with both itraconazole (CC_{50} >50 µM) and 17-OH

itraconazole (CC_{50} >100 µM) (Figure 1). For remdesivir CC_{50} values >100 µM were observed in

Caco-2 cells.

Figure 1. Effect of either itraconazole, 17-OH itraconazole or remdesivir on SARS-CoV-2

replication in CPE assays and on viability of Caco-2 cells.

Mean percent inhibition for each readout across two series of three independent experiments

(itraconazole, remdesivir) or three independent experiments (17-OH itraconazole) with triplicate

measurements are plotted. The error bars represent the standard deviation. Orange represents

CPE visual read-out; purple represents MTT assay; and green represents cytotoxicity.

A)

197
198
199   EC_{50} by visual scoring of inhibition of CPE = 1.5 μM; EC_{50} by MTT assay = 2.3 μM
200
201   **B)**
202

203
204
205   EC_{50} by visual scoring of inhibition of CPE = 1.2 μM; EC_{50} by MTT assay = 3.6 μM

C)

208
209
210
EC_{50} by visual scoring of inhibition of CPE = 0.3 µM; EC_{50} by MTT assay = 0.4 µM

***In Vitro* Viral RNA Yield Reduction**

The *in vitro* effect of itraconazole and remdesivir (Caco-2 cells) orGS-441522 (VeroE6-eGFP

cells) on SARS-CoV-2 RNA yield reduction was assessed in Caco-2 cells and VeroE6-eGFP

cells. A concentration of 6.25 µM and 3.1 µM of itraconazole and its main metabolite 17-OH

itraconazole, respectively, resulted in an approximately 2-$\log_{10}$ reduction in viral RNA levels (as

a measure of the number of virus particles in the culture SN) in Caco-2 cells. This reduction was

observed at all higher doses, in the absence of toxicity. However, remdesivir proved more

potent (a reduction of ~3 $\log_{10}$ at 6.25 µM) (Figure 2).

Fig 2. Effect of itraconazole, 17-OH itraconazole or remdesivir on SARS-CoV-2 vRNA

yield and viability in Caco-2 cells. A) Mean differences in viral RNA in the supernatant

between untreated cultures and treated cultures at 48 hours post-infection with SARS-CoV-2-

FFM1 of three independent experiments each containing two replicates is shown. Error bars

represent the standard deviation. B) Mean viability of the cells, based on three independent

experiments each containing three replicates is shown. Error bars represent the standard

deviation.

230
231   **A)**

232
B)

In VeroE6-eGFP cells, a concentration of 1 µM itraconazole resulted in a ~1-$\log_{10}$ reduction in

viral RNA levels (Figure 3). This reduction was also observed at higher concentrations. As in

Caco-2 cells, GS-441524 proved more potent in reducing viral RNA than itraconazole. At a

concentration of 3.7 µM, GS-441524 reduced viral RNA load by >4 $\log_{10}$ to undetectable levels.

At concentrations of 10 µM and higher, 17-OH itraconazole resulted in limited activity, but

VeroE6-eGFP cell viability was markedly affected (Figure 3).

Fig 3. Effect of itraconazole, 17-OH itraconazole or GS-441524 on SARS-CoV-2 vRNA

yield and viability in VeroE6-eGFP cells. A) and B) Mean differences in viral RNA in the

supernatant between untreated cultures and treated cultures at 48 hours post-infection with

SARS-CoV-2-Belgium of one independent experiment containing 2 replicates (A) or 3 replicates

(B) is shown. Error bars represent the standard deviation. C) and D) Mean viability of the cells,

based on MTT readout of uninfected cells. Error bars represent the standard deviation.

A)   **B)**

256   **C)**   **D)**

Discussion

While a number of vaccines are in development, for the moment there are no antivirals for the prevention of COVID-19 and only a limited number of antivirals for the treatment of COVID-19.[22] Given the great unmet need to identify potential treatments for COVID-19, and the fact that a *de novo* drug development project will take many years to bring a drug to patients, efforts have been expedited into the *in vitro* screening of compounds already on the market, or in late-stage development. In such an effort, itraconazole was identified, with single digit micromolar activity in Caco-2 cells. Itraconazole was first approved (in the US) in 1992 as an oral treatment for a number of fungal infections in immunocompromised and non-immunocompromised patients, with a well-established safety profile.[11]

These experiments demonstrated that itraconazole inhibits *in vitro* SARS-CoV-2 replication in Caco-2 cells. The EC_{50} values for inhibition of virus-induced CPE formation is in the low micromolar range and in that respect, is only around 5-fold higher than that of remdesivir. To obtain further evidence for the antiviral potency of the drug(s), the effect on viral RNA yield in culture was assessed in two cell lines. Itraconazole results in a marked inhibition of virus yield; however, itraconazole was less potent than remdesivir. Our findings are in line with recent data showing that itraconazole inhibited SARS-CoV-2 replication in a dose-dependent manner in both Calu-3 and Vero E6 cell lines (with EC_{50}s of 0.43 µM and 0.39 µM respectively).[23] In addition, *in vitro* studies have also demonstrated activity of itraconazole against murine coronavirus (recombinant murine coronavirus, mouse hepatitis virus, EC_{50} = 7.9 µM)[5] and type I (but not type II) feline coronavirus (EC_{50} across three strains tested = 0.146–0.597 µM).[6] Itraconazole has documented activity against a number of enteroviruses, including rhinovirus.[8,15] Further, at a dose of 5.7 mg/kg, itraconazole resulted in improved mortality and a lower viral load in mice that had been infected with human influenza (strain PR8M) compared with the control group.[7]

As an antifungal agent, itraconazole inhibits fungal ergosterol biosynthesis, through inhibition of

a cytochrome P450 enzyme, lanosterol-14-alpha-demethylase.[13–16] It is also thought to interfere

with cholesterol homeostasis and *de novo* synthesis in the host's cells through inhibition of this

same enzyme, thereby influencing the virus/host cell interaction.[17,24–26] The antiviral effect of

itraconazole may be, at least in part, based on such a mechanism. Itraconazole impairs

cholesterol trafficking and blocks late endosomal/lysosomal export of cholesterol to the plasma

membrane through NPC1 leading to cholesterol accumulation in sub-cellular

compartments.[8,16,24–27] In turn, elevated cholesterol levels in the endosomal membrane impede

fusion of the viral lipid envelope and prevent viral genome transfer into the host cell cytosol.[7,28] It

has been postulated that such changes in cellular cholesterol are linked with the host's immune

response.[29] Disruption of cholesterol biosynthesis has been shown to upregulate type I

interferons and thereby accelerate the virus-induced interferon-mediated host cell response.[7,29]

It has also been suggested that itraconazole may have antiviral effects through proteins other

than NPC intracellular cholesterol transporter 1 (NPC1).[6] In enteroviruses, itraconazole has

been shown to inhibit viral RNA replication by oxysterol-binding protein (OSBP), which is

responsible for trafficking of cholesterol and phosphatidylinositol-4-phosphate between

membranes, thus also affecting the membranes of the replication complex.[15,16] Since

coronaviruses result in the formation of intracellular viral replicative organelles within the

endoplasmic reticulum that drive the viral replication cycle,[30–32] it is possible that inhibition of

OSBP may cause detrimental changes to replicative organelle membranes.[15,16,32,33] Although it

remains to be confirmed how itraconazole may inhibit the replication cycle of coronaviruses, it is

conceivable that it acts through multiple mechanisms.[7,17]

As a lipophilic compound, itraconazole has high bioavailability and extensive distribution

throughout the lung, kidney, epidermis and brain.[9] Itraconazole is predominantly metabolized by

the cytochrome P450 3A4 isoenzyme system, forming many metabolites.[34,35] The main

metabolite produced is 17-OH itraconazole, which also has considerable antifungal activity.[34]

Following administration of itraconazole 200mg bid, itraconazole and its 17-OH hydroxy-

metabolite have been shown to reach mean maximal plasma concentrations of 2282 and 3488

ng/mL (~ 3.2 µM and ~4.8 µM) and a C_{trough} of 1855 and 3349 ng/ml (~2.6 µM and ~4.7 µM),

respectively, which is close to the EC_{50} for inhibition of virus-induced CPE. Both the parent drug

and metabolite have long half-lives (64 and 56 hours, respectively).[11]

In conclusion, itraconazole and its metabolite, 17-OH itraconazole show *in vitro* activity against

SARS-CoV-2. Based on these *in vitro* findings, a clinical study in hospitalized COVID-19

patients was initiated by UZ Leuven in March 2020.[36] However, no efficacy of the drug was

observed in this particular study population; other studies in non-hospitalized patients were not

conducted.

Acknowledgments

This project has received funding from the European Union's Horizon 2020 research and

innovation program under grant agreement no.: 101003627. Part of this research work was

performed using the 'Caps-It' research infrastructure (project ZW13-02) that was financially

supported by the Hercules Foundation and Rega Foundation, KU, Leuven. This work has been

funded in part with Federal funds from the Office of the Assistant Secretary for Preparedness

and Response, Biomedical Advanced Research and Development Authority, under OTA No.

HHSO100201800012C. Medical writing supporting for the development of this manuscript was

provided by Patrick Hoggard of Zoetic Science, an Ashfield company, part of UDG Healthcare

plc; this support was funded by Janssen Pharmaceuticals. We thank Lena Stegmann, Winston

Chiu, Piet Maes, Robbert Boudewijns, Nguyen Dan Thuc Do, Xin Zhang, Jana Van Dycke,

Rana Abdelnabi, Laura Thijs, Peggy Geluykens, Doortje Borrenberghs, and Christel Van den

Eynde for technical support for antiviral assays.

Disclaimers

Sandra De Meyer, Ellen Van Damme, Christophe Buyck and Marnix Van Loock are employees

and may be stock owners of Johnson & Johnson. Sandra Ciesek received research funding

from Janssen for this research. Steven De Jonghe, Dirk Jochmans, Pieter Leyssen and Johan

Neyts are employees from KU Leuven and received funding from Janssen for this research.

The data sharing policy of Janssen Pharmaceutical Companies of Johnson & Johnson is

available at https://www.janssen.com/clinical-trials/transparency. As noted on this site, requests

for access to the study data can be submitted through Yale Open Data Access (YODA) Project

site at http://yoda.yale.edu.

References

1.   Hoehl S, Rabenau H, Berger A, Kortenbusch M, Cinatl J, Bojkova D, et al. Evidence of

SARS-CoV-2 infection in returning travelers from Wuhan, China. N Engl J Med.

2020;328:1278–80.

2.   Lu R, Zhao X, Juan L, Niu P, Yang B, Wu H, et al. Genomic characterisation and

epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding.

Lancet. 2020a;395:565–74.

3.   Su S, Wong G, Shi W, Liu J, Lai ACK, Zhou J, et al. Epidemiology, Genetic Recombination,

and Pathogenesis of Coronaviruses. Trends Microbiol. 2016;24:490–52.

4.   Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients

with pneumonia in China, 2019. New Engl J Med. 2020;382:727–33.

5.   Cao J, Forrest CJ, Zhang X. A screen of the NIH Clinical Collection small molecule library

identifies potential anti-coronavirus drugs. Antiviral Res. 2015;114:1–10.

6.   Takano T, Akiyama M, Doki T, Hohdatsu T. Antiviral activity of itraconazole against type 1

feline coronavirus infection. Vet Res. 2019;50:5.

7.   Schloer S, Goretzko J, Kühnl A, Brunotte L, Luwdig S, Rescher U. The clinically licensed

antifungal drug itraconazole inhibits influenza virus *in vitro* and *in vivo*. Emerg Microbes

Infect. 2019;8:80–93.

8.   Shim A, Song JH, Kwon BE, Lee JJ, Ahn JH, Kim YJ. Therapeutic and prophylactic activity

of itraconazole against human rhinovirus infection in a murine model. Sci Rep.

2016;6:23110.

9.   Saag MS, Dismukes WE. Azole antifungal agents: emphasis on new thiazoles. Antimicrob

Agents Chemother. 1988;32:1–8.

10. Kim J, Tang JY, Gong R, Kim J, Lee JJ, Clemons KV, et al. Itraconazole, a commonly used

antifungal that inhibits Hedgehog pathway activity and cancer growth. Cancer Cell.

2010;17:388–99.

11. Sporanox® Summary of Product Characteristics. Ortho-McNeil-Janssen Pharmaceuticals,

Inc. 2012. Available at:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2012/020083s048s049s050lbl.pdf

(Last accessed: November 2020).

12. De Doncker P, Pande S, Richarz U, Garodia N. Itraconazole: What clinicians should know?

Indian J Drugs Dermatol. 2017;3:4–10.

13. Da Silva Ferreira ME, Colombo AL, Paulsen I, Ren Q, Wortman J, Huang J, et al. The

ergosterol biosynthesis pathway, transporter genes, and azole resistance in Aspergillus

fumigatus. Med Mycol. 2005;43(Suppl 1):S313–S319.

14. Lepesheva GI, Waterman MR. Sterol 14α-demethylase cytochrome P450 (CYP51), a P450

in all biological kingdoms. Biochim Biophys Acta. 2007;1770:467–77.

15. Strating JR, van der Linden L, Albulescu L, Bigay J, Arita M, Delang L, et al. Itraconazole

inhibits enterovirus replication by targeting the oxysterol-binding protein. Cell Rep.

2015;10:600–15.

16. Bauer L, Ferla SA, Bhat S, Pasunooti KK, Shi WQ, Albulescu L, et al. Structure-activity

relationship study of itraconazole, a broad-range inhibitor of picornavirus replication that

targets oxysterol-binding protein (OSBP). Antiviral Res. 2018; 156:55–63.

17. Takano T, Endoh M, Fukatsu H, Sakurada H, Doki T, Hohdatsu T. The cholesterol transport

inhibitor U18666A inhibits type 1 feline coronavirus infection. Antiviral Res. 2017;145:96–

102.

18. Schloer S, Goretzko J, Kühnl A, Brunotte L, Ludwig S, Rescher U. The clinically licensed

antifungal drug itraconazole inhibits influenza virus in vitro and in vivo. Emerg Microbes

Infect. 2019;8:80–93.

19. Ivens T, Van den Eynde C, Van Acker K, Nijs E, Dams G, Bettens E, et al. Development of

a homogeneous screening assay for automated detection of antiviral agents active against

severe acute respiratory syndrome-associated coronavirus. J Virol Methods. 2005t;129:56–

63.

20. De Meyer S, Bojkova D, Cinatl J, Van Damme E, Buyck C, Van Loock M, et al. Lack of

antiviral activity of darunavir against SARS-CoV-2. Int J Infect Dis. 2020;97:7-10.

21. Bojkova D, Klann K, Koch B, Widera M, Krause D, Ciesek S, et al. Proteomics of SARS-

CoV-2-infected host cell reveals therapy targets. Nature. 2020;583:469-472.

22. Lu H. Drug treatment options for the 2019-new coronavirus (2019-nCoV). Biosci trends.

2020b;14:69–71.

23. Schloer S, Brunotte L, Mecate-Zambrano A, Zheng S, Tang S, Ludwig S, et al. Drug

synergy of combinatory treatment with remdesivir and the repurposed drugs fluoxetine and

itraconazole effectively impairs SARS-CoV-2 infection in vitro. bioRxiv:

2020.2010.2016.342410.

24. Shoemaker CJ, Schornberg KL, Delos SE, Scully C, Pajouhesh H, Olinger GG, et al.

Multiple cationic amphiphiles induce a Niemann-Pick C phenotype and inhibit Ebola virus

entry and infection. PLoS One. 2013;8:e56265.

25. Trinh MW, Lu F, Li X, Das A, Liang Q, De Brabander JK, et al. Proc Natl Acad Sci USA.

2017;114:89–94.

26. Amini-Bavil-Olyaee S, Choi YJ, Lee JH, Shi M, Huang IC, Farzan M, et al. The antiviral

effector IFITM3 disrupts intracellular cholesterol homeostasis to block viral entry. Cell Host

Microbe. 2013;13:452–64.

27. Head SA, Shi WQ, Yang EJ, Nacev BA, Hong SY, Pasunooti KK, et al. Simultaneous

targeting of NPC1 and VDAC1 by itraconazole leads to synergistic inhibition of mTOR

signaling and angiogenesis. ACS Chem Biol. 2017;12:174–82.

28. Kühnl A, Musiol A, Heitzig N, Johnson DE, Ehrhardt C, Grewal T, et al. Late

endosomal/lysosomal cholesterol accumulation is a host cell-protective mechanism

inhibiting endosomal escape of influenza A virus. MBio. 2018;9:e01345–18.

29. York AG, Williams KJ, Argus JP, Zhou QD, Brar G, Vergnes L, et al. Limiting cholesterol

biosynthetic flux spontaneously engages type I IFN signaling. Cell. 2015;163:1716–29.

30. Risco C, de Castro IF Sanz-Sánchez L, Narayan K, Grandinetti G, Subramaniam S. Three-

dimensional imaging of viral infections. Annu Rev Virol. 2014;1:453–73.

31. Romero-Brey I, Bartenschlager R. Membranous replication factories induced by plus-strand

RNA viruses. Viruses. 2014;6:2826–57.

32. Neuman BW, Angelini MM, Buchmeier MJ. Does form meet function in the coronavirus

replicative organelle? Trends Microbiol. 2014;22:642–7.

33. Takano T, Wakama Y, Doki T. Endocytic pathway of feline coronavirus for cell entry:

differences in serotype-dependent viral entry pathway. Pathogens. 2019;8:300.

34. Peng C, Shi W, Lutz JD, Kunze KL, Liu JO, Nelson WL, et al. Stereospecific metabolism of

itraconazole by CYP3A4: Dioxoalane ring scission of azole antifungals. Drug Metab Dispos.

2012;40:426–35.

35. Prentice AG, Glasmacher A. Making sense of itraconazole pharmockinetics. J Antimicrob

Chemother. 2005;56(Suppl 1):i17–i22.

36. EUDRACT. Covid-19: A randomized, open-label, adaptive, proof-of- concept clinical trial of

new antiviral drug candidates against SARS-CoV-2. 2020-001243-15

https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-001243-15/BE (Last accessed:

November 2020).