Predictive attenuation of touch and tactile gating are distinct perceptual phenomena

Konstantina Kilteni[1*] and H. Henrik Ehrsson[1]

[1]Department of Neuroscience, Karolinska Institutet, Solnavägen 9, 17165 Stockholm, Sweden
[*]Correspondence and requests for materials should be addressed to konstantina.kilteni@ki.se

In recent decades, research on somatosensory perception has led to two important observations. First, self-generated touches that are predicted by voluntary movements become attenuated compared to externally generated touches of the same intensity (attenuation). Second, externally generated touches feel weaker and are more difficult to detect during movement compared to rest (gating). Researchers today often consider gating and attenuation to be the same suppression process; however, this assumption is unwarranted because, despite more than forty years of research, no study has combined them in a single paradigm. We quantified how people perceive self-generated and externally generated touches during movement and rest. We demonstrate that whereas voluntary movement gates the precision of both self-generated and externally generated touch, the amplitude of self-generated touch is selectively attenuated compared to externally generated touch. We further show that attenuation and gating neither interact nor correlate, and we conclude that they represent distinct perceptual phenomena.

Let's imagine you are at your doctor for a medical examination. Upon her request, you apply pressure with your index finger on your leg to indicate where exactly you feel the pain. The pressure you feel on your leg and the tip of your finger is the feedback from your voluntary finger movement, and it is called *somatosensory reafference*. To reproduce and confirm your sensations, imagine that the doctor now applies pressure with her index finger at the same spot on your leg. This pressure is generated by the doctor and not from you, and it is called *somatosensory exafference*. Now imagine that the doctor asks you to first keep your leg relaxed and then flex and extend it while you or her continuously apply pressure on your leg. You can therefore experience your reafferent or her exafferent touches on your leg, while it is moving or resting. Distinguishing between these four conditions is fundamental for your sensorimotor control; your nervous system must know both the source of the touch and the state of your limb to make appropriate use of the sensory feedback. A cutaneous mechanoreceptor in your peripheral nervous system though, cannot distinguish whether a touch is reafferent or exafferent, and thus this distinction must be made centrally, where tactile signals from the skin, sensory information from muscles and joints, and information from motor commands are available. How, then, does the central nervous system classify somatosensory signals during movement?

Several experimental studies in humans have shown that the brain *attenuates* somatosensory reafference compared to exafference. In behavioral research, this refers to participants perceiving self-generated strokes, forces, or taps as weaker than external equivalents of the same intensity [1–10]. This somatosensory attenuation has been related to reduced activity in the secondary somatosensory cortex [11–13] and the cerebellum [11,13,14] and an increased connectivity between the two areas [13,15] during self-generated compared to externally generated touches. Somatosensory attenuation was found in 98% or 315 out of 322 people across a wide age range [16], and it is considered one of the reasons why we cannot tickle ourselves [17–19].

Sensory attenuation is not exclusive to humans; similar strategies are used in other species across the animal kingdom (for reviews, see [20–24]). For example, during self-chirping, the cricket's central auditory processing is inhibited (both presynaptically and postsynaptically) in phase with the insect's chirps to prevent desensitization of its auditory system while maintaining sensitivity to external sounds [25,26]. In mice, auditory cortical responses in response to self-generated sounds are attenuated, and this attenuation is present only for the tone frequencies the animal has associated with its locomotion and absent when the same sounds are externally produced [27]. A weakly electric fish [28] is able to respond exclusively to externally generated electrical discharges by attenuating its predicted electrosensory reafference [21,29]. In primates, activity in the vestibular nucleus in response to vestibular reafference is attenuated during active head movements compared to passive head movements, allowing the animal to maintain its head and body posture and activate the vestibular-related reflexes when appropriate [20,21,30–32].

At the same time, another branch of experimental research has shown that somatosensory sensitivity in response to externally generated stimuli is *gated* during and before a voluntary movement. In human psychophysical research, this phenomenon of *movement-related tactile gating* or *tactile suppression* manifests as an elevation in the detection threshold [33–42] and a decrease in the subjective intensity of externally generated stimuli [35,37,43] when the stimulated body part moves compared to when it is at rest. Several electrophysiological studies have shown this gating as an inhibition of somatosensory evoked potentials during active movement compared to rest, at subcortical and cortical sites along the somatosensory pathway [43–48]. Similar to somatosensory attenuation, tactile gating is a biologically preserved mechanism that is seen across different species [49]. For example, responses recorded at the cat's medial lemniscus evoked by nerve stimulation are suppressed prior to and during limb movements [50]. Similarly, the transmission of cutaneous afferent signals to the primary somatosensory cortex is suppressed in rats during movement compared to rest [51]. In monkeys, the gating of cutaneous afferent input during active movement has been observed both in the primary somatosensory cortex [52–54] and the spinal cord [54,55].

Somatosensory attenuation and tactile gating share two important conceptual similarities. First, they both refer to a modulation —either in terms of magnitude or precision— in the perception of cutaneous stimuli during movement. Second, they have been assigned the same functional role [39]: to reduce the flow of afferent information that can be predicted from the motor command, so that detection of external input that can be biologically important, such as touches caused by predators[17,24,56], or input that is task-relevant for the upcoming or ongoing movement [45,47,57] is facilitated.

Importantly, however, the two phenomena present one striking difference. Somatosensory attenuation relates to somatosensory reafference, that is, touches caused *by* our voluntary movement. In contrast, gating relates to somatosensory exafference, that is, external touches occurring *during* our voluntary movement. Nevertheless, somatosensory research, including very recent studies and review articles, e.g., [41,42,58–64], often treats the two phenomena as a single generalized suppression strategy of the brain and uses the terms attenuation and gating interchangeably. If the two phenomena are indeed different, this false equivalence is detrimental for understanding human sensorimotor control. First, it cannot advance our understanding at the computational level because researchers try to explain gating (e.g., [41]) and attenuation (e.g., [65]) using the computational processes proposed for attenuation and gating, respectively. Similarly, at the neurobiological level, researchers intermix neural correlates of gating and attenuation (e.g., [62,63]) because they assume that these measure the same single

100 phenomenon. Second, this confusion becomes particularly disadvantageous in clinical studies
101 using gating and attenuation when interpreting findings of sensorimotor deficits in patients with
102 schizophrenia [66–68], functional movement disorders [69] or Parkinson's disease [45,70,71].
103
Are these phenomena the same, or do they represent two fundamentally distinct processes?
Does the brain treat all sensory stimuli similarly during movement, independent of whether
they are reafferent or exafferent? To the best of our knowledge, there has been no previous
attempt to simultaneously test attenuation and gating with the same stimulus and psychophysics
task. Here, in a single experimental design, we investigated the perception of touches applied
on the left hand while manipulating whether the left arm was in movement or resting (left limb
state). We additionally manipulated whether the touches were reafferent, generated by the right
hand, or exafferent, generated by an external source (origin of touch). We reasoned that if the
two phenomena are the same, then they should influence somatosensory perception in similar
ways. Our results demonstrate that this is not the case: voluntary movement reduces the
somatosensory precision of both sensory reafference and exafference (gating), while the
perceived amplitude of sensory reafference is selectively attenuated compared to exafference
(attenuation). Notably, the two phenomena do not correlate with each other or interact when
present simultaneously. Thus, collectively, our results show that gating and attenuation are two
separate processes that can be experimentally dissociated.
119
120
Results
Participants rested their left hands palm up with their index fingers placed on a molded support,
and their right hands were palm down on top of a set of sponges (**Fig. 1a-d**). In all conditions,
they performed a force-discrimination task [3,4,10,67]: in each trial, a motor delivered two taps (the
test tap and the *comparison* tap) on the pulp of their left index finger, and they were asked to
verbally indicate which tap felt stronger (**Fig. 1e-h**). While the *test* tap had a fixed intensity (2
N), the *comparison* tap randomly changed its intensity on every trial (1, 1.5, 1.75, 2, 2.25, 2.5,
or 3 N). An auditory 'go' signal indicated the trial's onset and the onset of the response period.
129
In all conditions, our experimental manipulation exclusively concerned the *test tap*; the
comparison tap was always externally triggered and delivered on the relaxed left arm, serving,
therefore, as a reference stimulus. In a factorial design, we controlled for whether the left arm
moved **(Fig. 1c, d)** or remained relaxed **(Fig. 1a, b)** during the *test* tap, and whether the *test* tap
was produced by the right hand (self-generated) **(Fig. 1b, d)** or not (externally generated) **(Fig.
1a, c)**. This design resulted in four experimental conditions, the order of which was fully
counterbalanced (**Fig. 1a-d**).
137
In the *baseline* condition (**Fig. 1a**), participants did not move their limbs but passively received
the *test* and the *comparison* taps on the left index. This control condition assesses the
participants' somatosensory perception in the absence of any movement [2,4,10]. In the *attenuation*
condition (**Fig. 1b**), participants actively tapped with their right index finger a force sensor
placed on top of their left index finger. The tap of their right index finger on the force sensor
triggered the *test* tap on their left index finger. This classic condition assesses the perception of
a self-generated tap on a passive limb [2,4,10]. In the *gating* condition (**Fig. 1c**), participants were
asked to continuously move their left arm forward, sliding the experimental setup with the
motor and the force sensors between a start and an end position (distance 25 cm) at a
comfortable velocity of approximately 20 cm/sec (**Fig. 1i-j, Supplementary Fig. S1, S2**).
During this movement, participants received the *test* tap on their left index finger. This is a
gating condition because it assesses the perception of an externally generated tap on a moving

limb[36–38,40]. Finally, in the *attenuation&gating* condition (**Fig. 1d**), the participants performed the same movement with their left arm, but they were additionally asked to actively tap with their right index finger on the force sensor that triggered the *test* tap on their left index finger during the movement. The force sensor was attached to the experimental setup and moved together with the left hand. This condition combines the gating and attenuation phenomena since it assesses the perception of a self-generated tap on a moving limb.

Fig. 1. Experimental conditions. Two factors were manipulated in the experiment, resulting in four experimental conditions: whether the left arm was in rest (**a**, **b**) or in movement while receiving the *test* tap (**c**, **d**) (state of the left limb) and whether the *test* tap was externally triggered by the motor (**a**, **c**) or self-triggered by the participants' right hand (**b**, **d**) (origin of touch). In all four conditions (**e-h**), the participants received the two taps (*test* and *comparison* tap) on the pulp of their left index fingers from the motor (black stars), and they had to verbally indicate which was stronger: the first or the second tap. In the *attenuation* and *attenuation&gating* conditions, the participants self-triggered the test tap on their left index finger by moving their right arm to tap a sensor with their right index finger (blue stars). **(i)** In the *gating* and *attenuation&gating* conditions (**c** and **d**), the participants extended their left arm from a start position to an end position, sliding the experimental setup along two drawer runners. During this movement, participants experienced the *test* tap. A motion sensor recorded the position of the platform in time. **(f)** Example position traces recorded by the motion sensor for the movements of one participant during the *attenuation&gating* condition. **(g)** Responses of a representative participant and fitted logistic

models for the four experimental conditions. Data points have been horizontally jittered to avoid overlapping.

The participant's responses in each condition were fitted with a generalized linear model (**Fig. 1k**, **Supplementary Fig. S3**), and two parameters of interest were extracted: the point of subjective equality (PSE), which represents the intensity at which the test tap felt as strong as the comparison tap ($p = 0.5$), and the just noticeable difference (JND), which reflects the participants' sensitivity in force discrimination. We hypothesized that the two phenomena are different and, thus, that they would affect the PSE and JND differently. Specifically, we expected a decrease in the perceived magnitude (lower PSE) for the conditions with sensory reafference (*attenuation* and *attenuation&gating* conditions), with no effects on the sensory precision of the participants (JND) when they received the reafference on a still limb (*attenuation* condition). In contrast, we predicted a decrease in the sensory precision (higher JND) of both sensory reafference and exafference for the conditions where the limb that receives the touches moves (*gating*, *attenuation&gating*) and small, if any, effects in the PSE. In other words, we expected that the attenuation phenomenon affects the PSE and not the JND, and the gating phenomenon mainly affects the JND. Our hypotheses were supported by the data (**Fig. 2a-g**).

Fig. 2. Experimental results. (a, b) Bar graphs show the PSEs and JNDs (mean ± SEM) for each condition. A lower PSE value indicates a lower perceived magnitude, while a higher JND value indicates a lower somatosensory sensory precision. Reafferent touches produced by the movement of the right arm (*attenuation* and *attenuation&gating* conditions) were associated with a significant drop in the PSEs, while the movement of the left arm that receives the touches (*gating* and *attenuation&gating*) produces a significant increase in the JNDs. **(c)** Group psychometric functions for each condition generated using the mean PSE and the mean JND across participants. The leftward shift of the curves

for the *attenuation* and *attenuation&gating* conditions illustrates that somatosensory reafference is
perceived as weaker than exafference. The flattening of the curves for the *gating* and *attenuation&gating*
conditions illustrates the worsening in somatosensory precision of both reafference and exafference on
a moving limb. **(d, e)** Line plots illustrate the decreases in PSEs when experiencing reafferent touches
compared to exafferent touches, when the left arm is still **(d)** and when the left arm moves **(e)**. **(f, g)**
Line plots illustrate the increases in JNDs when receiving the touches on a moving limb compared to
rest, when the touches are exafferent **(f)** and when the touches are reafferent **(g)**.
205
We performed a repeated-measures ANOVA on the PSEs with the origin of the touch
(reafference *vs* exafference) and the state of the left limb (movement *vs* rest) as factors. This
revealed a significant main effect of the origin of the touch ($F(1, 23) = 36.10$, $p < 0.001$, $\eta_p^2 =$
0.611), a significant main effect of the left limb state ($F(1, 23) = 13.91$, $p = 0.001$, $\eta_p^2 = 0.377$)
and a nonsignificant interaction ($F(1, 23) = 0.26$, $p = 0.615$, $\eta_p^2 = 0.011$) (**Fig. 2a**). A Bayesian
repeated-measures ANOVA further concluded against the interaction term, by supporting the
model without the interaction term compared to the full factorial ($BF_{M1} / BF_{M2} = 3.631$). The
attenuation condition produced a significant drop in the PSE compared to the *baseline* condition
($n = 24$, $t(23) = -5.908$, $p < 0.001$, *Cohen's d* $= -1.206$, $CI^{95} = [-0.332, -0.160]$, $BF_{01} < 0.0003$)
(**Fig. 2d**). This replicates previous attenuation findings indicating that a self-generated tap on a
passive limb feels weaker than a tap of the same intensity but of an external origin [2,4–7,10,13,16].
Similarly, the PSE in the *attenuation&gating* condition was significantly lower than that in the
gating condition ($n = 24$, $t(23) = -5.032$, $p < 0.001$, *Cohen's d* $= -1.027$, $CI^{95} = [-0.377, -0.157]$,
$BF_{01} = 0.002$) (**Fig. 2e**), extending the previous conclusion to when the receiving limb is
moving. Together, these two contrasts show that reafferent (self-generated) touches feel weaker
than exafferent touches, both when the receiving hand is in movement or rest.
222
The *gating* and *attenuation&gating* conditions also produced a significant drop in the PSE
compared to the *baseline* condition ($n = 24$, $t(23) = -2.409$, $p = 0.024$, *Cohen's d* $= -0.492$, CI^{95}
$= [-0.141, -0.011]$) and the *attenuation* condition ($n = 24$, $V = 55$, $p = 0.005$, $rrb = -0.633$, CI^{95}
$= [-0.161, -0.022]$), respectively. However, these decreases were quite modest ($\cong 30\%$ of the
weakening produced by the *attenuation* condition) and supported only by anecdotal evidence
from Bayesian statistics ($BF_{01} = 0.433$ and $BF_{01} = 0.738$, respectively). Together, these
contrasts suggest that exafferent touches can feel slightly weaker on a moving limb than on a
passive limb, in agreement with previous findings for tactile gating [35,37,43]. Nevertheless, when
compared with tactile reafference, the perceived magnitude of tactile exafference is not strongly
decreased.
233
When testing for the effects of the conditions on the somatosensory precision of the participants
(JND), there was a significant main effect of the state of the left limb ($F(1, 23) = 17.1$, $p <$
0.001, $\eta_p^2 = 0.426$), but there was neither a significant main effect of the origin of touch ($F(1,$
$23) = 0.52$, $p = 0.478$, $\eta_p^2 = 0.022$) nor a significant interaction ($F(1, 23) = 0.06$, $p = 0.809$, η_p^2
$= 0.003$) (**Fig. 2b**). As with PSEs, the absence of interaction for the JNDs was further supported
with a Bayesian repeated-measures ANOVA that provided evidence against the interaction term
($BF_{M1} / BF_{M2} = 3.522$). The *attenuation* condition did not produce any change in the JND with
respect to the *baseline* condition ($n = 24$, $t(23) = 0.331$, $p = 0.744$, *Cohen's d* $= 0.068$, $CI^{95} = [-$
$0.034, 0.047]$), and this was substantially confirmed with a Bayesian analysis ($BF_{01} = 4.432$).
In alignment, no significant differences in the JND were detected between the *gating* and the
attenuation&gating conditions ($n = 24$, $t(23) = 0.72$, $p = 0.481$, *Cohen's d* $= 0.146$, $CI^{95} = [-$
$0.024, 0.05]$), and this was again confirmed with a Bayesian analysis ($BF_{01} = 3.691$). Together,
these two contrasts demonstrate that receiving sensory reafference *per se* is not accompanied
by a worsening of sensory precision on the receiving limb.
248

In contrast, moving the limb while receiving an external touch (*gating* condition) produces a
significant increase in the JND compared to the *baseline* condition ($n = 24$, $t(23) = 3.134$, $p =$
0.005, *Cohen's d* $= 0.640$, $CI^{95} = [0.019, 0.093]$, $BF_{01} = 0.108$) (**Fig. 2f**). This was further
confirmed by a significant increase in JND between the *attenuation* and the *attenuation&gating*
conditions ($n = 24$, $t(23) = 2.984$, $p = 0.007$, *Cohen's d* $= 0.609$, $CI^{95} = [0.019, 0.106]$, $BF_{01} =$
0.146) (**Fig. 2g**). Together, these two contrasts indicate that voluntary movement *per se*
deteriorates the precision with which reafferent and exafferent stimuli are perceived on the
moving limb.
257
Together, our results indicate that predicting the sensory consequences of a voluntary
movement produces a decrease in the perceived magnitude of sensory reafference (PSE)
without a concomitant worsening in somatosensory precision (JND). In contrast, a voluntary
movement leads to a decrease in somatosensory precision (JND) for both sensory reafference
and exafference. These effects are observed in the group psychometric fits (**Fig. 2c**). To further
illustrate that somatosensory attenuation affects the amplitude (PSE), while tactile gating
affects the precision (JND) and not *vice versa*, we tested whether we could predict better the
participants' performance in the *attenuation&gating* condition when using the PSE from the
attenuation condition and the JND from the *gating* condition than the PSE from the *gating*
condition and the JND from the *attenuation* condition. Indeed, the first model was significantly
better: $n = 24$, $V = 39$, $p < 0.001$, $rrb = -0.74$, $CI^{95} = [-68.980, -12.736]$, $BF_{01} = 0.443$
(**Supplementary Fig. S4, Fig. 3a-b**).
270
In the abovementioned ANOVAs, there were no significant interactions between the two factors
(the origin of touch and the state of the limb), neither for the PSEs nor for the JNDs, according
to both frequentist and Bayesian analyses. This indicates that the effect of the left limb state
was not influenced by the effect of the origin of touch, neither for the PSE nor for the JND.
Instead, the two effects just added up when simultaneously present. To further test whether the
effects produced by each phenomenon relate to each other at all, we performed a correlation
analysis across the PSEs and JNDs. No significant correlations were detected between any of
the PSEs and any of the JNDs (all *p*-values > 0.225, $BF_{01} = [1.971, 3.950]$). The only significant
correlation found within PSEs was between the PSE in the *attenuation* condition and the PSE
in the *attenuation&gating* condition ($t(22) = 4.89$, $r = 0.722$, $p < 0.001$, $CI^{95} = [0.449, 0.871]$,
$BF_{01} = 0.002$) (**Fig. 3c**). That is, the weaker the participants perceived the magnitude of their
self-generated touch during rest, the weaker the magnitude of their self-generated touch during
movement felt. Given that the PSEs significantly dropped in these two conditions and that these
decreases correlated with each other, this result provides further support that their common
experimental denominator, i.e., the reafferent nature of the touch, was responsible for the drop
in the PSE and thus for the attenuation phenomenon. In contrast, the JND in the *gating* condition
was significantly correlated only with the *attenuation&gating* condition ($t(22) = 3.47$, $r = 0.595$,
$p = 0.008$, $CI^{95} = [0.252, 0.805]$, $BF_{01} = 0.047$) (**Fig. 3d**). This specific correlation means that
the worse the somatosensory precision of an external touch when participants moved their
receiving hand, the worse the sensory precision for a self-generated touch during the same
movement of the receiving hand. Given that the JNDs significantly increased only in these two
conditions and that these increases correlated with each other, this result provides evidence that
their common experimental denominator, i.e., the left limb's movement that receives the
touches, was responsible for the increase in the JND.
295

Fig. 3. Model predictions and scatterplots for PSEs and JNDs. (a) Average participants' responses in the *attenuation&gating* condition (yellow) and average predicted responses using the parameters from the *attenuation* and *gating* conditions (blue). The responses depicted in dark blue used the PSE of the *attenuation* condition and the JND from the *gating* condition, while the responses depicted in light blue used the PSE of the *gating* condition and the JND of the *attenuation* condition. Error bars and ribbons represent 95% confidence intervals. **(b)** For each participant, we estimated the Bayesian Information Criterion (BIC) of the fitted logistic model in the *attenuation&gating* condition and the two models with fixed parameters from the *attenuation* and *gating* conditions. The model using the PSE of the *attenuation* condition and the JND of the *gating* condition was a significantly better model than the one using the PSE of the *gating* condition and the JND from the *attenuation* condition. The upper right panel represents the same data after excluding one participant corresponding to the outlier observation, for illustration purposes. The exclusion of the outlier did not change the statistical results. **(c)** The participants' PSEs in the *attenuation* condition were significantly correlated with those in the *attenuation&gating* condition. **(d)** The participants' JNDs in the *gating* condition were significantly correlated with those in the *attenuation&gating* condition. There were no significant correlations between PSEs and JNDs.

A common finding in tactile gating studies is that the gating effects are stronger with higher movement velocities [33,36,40,41]; that is, the faster the limb movements are, the worst the

perception of the moving limb. Therefore, one could hypothesize that any differences observed
between the *gating* and the *attenuation&gating* conditions might be due to differences in the
velocity of the participants' movements. Since there were no significant JND differences in
these two conditions (**Fig. 2**), which was further supported by a Bayesian analysis, this concern
can be ruled out. However, one can argue that the PSE in the *attenuation&gating* condition was
lower than the PSE in the *attenuation* condition because the participants moved faster in the
attenuation&gating condition and not because of the reafferent nature of the touch. This
concern can also be ruled out since, rather than faster, we observed that the participants moved
slightly slower in the *attenuation&gating* condition (20.3 ± 0.003 cm/s) than in the *gating*
condition (23.3 ± 0.003 cm/s): (peak trial velocity; $t(23) = -4.062$, $p < 0.001$, *Cohen's d* = -
0.829, $CI^{95} = [-0.001, -0.004]$, $BF_{01} = 0.015$). Although the total distances the participants ran
with their left arm were comparable, participants in the *attenuation&gating* condition moved
slower because they had to coordinate both their arms to tap the sensor that the left arm moves
with the right hand (**Fig. 1d**). This difference was further confirmed when looking at the peak
velocities during the *test* tap between the two conditions (*gating*: 22.2 ± 0.003 cm/s;
attenuation&gating: 19 ± 0.003 cm/s) (**Supplementary Fig. S5**). Therefore, these kinematic
differences cannot explain the significant decrease in the PSEs in the *attenuation&gating*
condition compared to the *gating* condition, but they actually underscore the importance of
reafference in somatosensory attenuation.

336
337

Discussion

The present study contrasted the conditions of attenuation and gating in a single experimental
paradigm to investigate the relationship between the two phenomena. To this end, we
independently manipulated the origin of the touch (reafference *vs* exafference) and the state of
the receiving limb (movement *vs* rest), taking on all four possible combinations of these levels.
We replicated the classic phenomenon of predictive attenuation of touch [1,2,16,3–10] by showing
that somatosensory reafference feels weaker than somatosensory exafference. Importantly,
however, this decrease in the perceived amplitude (PSE) was not accompanied by a
concomitant worsening in somatosensory precision (JND). Participants had the same
discrimination capacity for both reafferent and exafferent touches applied on their passive limb,
a conclusion supported by Bayesian statistics. Nevertheless, when the limb that receives the
touches moved, this voluntary movement *per se* led to a decrease in somatosensory precision
(JND) for both reafferent and exafferent touches, replicating the classic tactile gating
phenomenon [33–36,39–42]. The two effects did not correlate and did not interact but summed up
when present together.

The main conclusion of the present study is that the predictive attenuation of touch and tactile
gating are two distinct perceptual phenomena. Our findings can help conciliate several previous
observations on gating and attenuation that have been studied in isolation during recent decades.
First, attenuation is observed not only on the active limb [8] but also on a passive limb (the
contralateral hand in the present study), as long as the contact between the body parts is
predicted by the voluntary movement [2–7,9,10,13,16]. In contrast, there is abundant evidence that
external touches applied to the limb contralateral to the one that moves are not gated [34,43,45,72,73].
Second, a touch that results from a passive movement [10] or touches that are simultaneously
presented in both hands (double touch) [67] are not attenuated. In contrast, gating effects have
been repeatedly demonstrated for passive movements, both electrophysiologically [45] and
behaviorally [34,38,39]. Third, self-generated tactile signals are attenuated as long as they are
presented at the timing predicted by the action [1,2,4]; even a 100 ms delay between the movement

and its tactile feedback greatly reduces the attenuation of the latter. In contrast, externally generated stimuli are gated when presented at any (unpredicted) timing during the movement [45], and importantly, even at hundreds of milliseconds before the movement onset [36]. Fourth, whereas gating shows no specificity for the type of motor activity and manifests both during isotonic [33–35,43] and isometric [35,73] contractions, attenuation is motor command-specific; a consistent but arbitrary and unnatural mapping between the motor command and the touch, for example, moving a joystick with one hand to produce touch on the other, does not produce attenuation [5,6,69]. Together, when seen in light of our findings, these results are not contradictory; instead, they refer to different perceptual phenomena.

Motor control relies on integrating afferent sensory information with efferent motor signals [74]. Distinguishing between gating and attenuation is fundamentally important for motor control theories because this can indicate a different integration or weighting mechanism of the motor and sensory information depending on the context. One well-established computational framework inspired by engineering approaches poses that the brain produces the motor command through an inverse model [75] or controller [76]. A copy of the motor command — termed 'efference copy' — is used by a forward model to predict the expected sensory feedback of the movement, which is then combined with the actual sensory input to estimate the state of the body [56,75–77]. With respect to the attenuation of sensory reafference, it has been proposed that the prediction signal of the forward model is used to 'cancel' the sensory reafference [17,56,78–80]. In other words, central motor processes play a major role in somatosensory attenuation compared to the actual sensory feedback. Support for this comes from studies showing that conditions that present highly predictable touches but in the absence of movement do not yield attenuation [2,10] (see also [81–83] for similar conclusions). The dependence of attenuation on action prediction was further shown when participants attenuated touches applied on one hand that were predicted by their other hand's movement, even when the two hands unexpectedly failed to make contact [3]. In further agreement, neuroimaging studies on somatosensory attenuation consistently report activation of the cerebellum [11–14,84], a structure that is associated with motor prediction [74,85–88].

In contrast, this computational account that relies on action prediction, efference copy, and internal forward models is inapplicable for tactile gating, since gated touches can be of exafferent nature and occur at any (unpredictable) time during movement, even before movement onset [36–38,47]. In other words, there is no information that the brain can use to predict the exafferent touches with the forward model because there is no causal relationship between the motor command and sensory input. This observation agrees with the proposal that peripheral afferent signals from muscle spindles and joint afferents play the major role in gating [36,45], and the fact that gating effects have also been observed during passive movement, without significant differences from active movements [34,38]. Then, if efference copy is not the basis for gating, how can the gating effects be computationally explained?

The alternative computational framework of active inference has been proposed to explain somatosensory attenuation[59]; however, based on an assumed equivalence between attenuation and gating. The active inference approach refutes the necessity of an efference copy, and emphasizes the importance of a generative model and reflex arcs in the place of forward and inverse models, and controllers [89,90]. According to the active inference account, the brain predicts the sensory input that would be expected from a specific action, and the body moves in order to fulfill these sensory predictions. Motor commands are thus conceptually replaced by proprioceptive predictions, and action occurs as a way to minimize the proprioceptive prediction errors when the movement has not been yet executed [91]. A major role is assigned to

the precision (i.e., reliability) that weights these sensory prediction errors depending on the context, and can be manipulated through attention allocation. Within this computational architecture, attenuation of somatosensory reafference is seen as a reduction in the precision of somatosensory evidence during movement in order to allow the expression of proprioceptive predictions that trigger the movement [59]. In other words, the agent attends away from somatosensory input in order to execute the movement. However, this proposal does not address the attenuation of sensory reafference with respect to exafference, since the agent should theoretically attend away from *all* somatosensory input, independently of their source. Moreover, the proposal cannot explain why increasing the precision of the proprioceptive prediction errors on the hand that is to move would reduce the precision of somatosensory prediction errors on the contralateral limb that is not meant to move and thus, there should not exist any proprioceptive predictions about it. In contrast, the active inference account can sufficiently explain the tactile gating effect; i.e., the reduction in the precision of somatosensory input on the moving limb during the movement.

A recent study [64] suggested that action prediction does not produce somatosensory attenuation but enhancement and argued that the attenuation observed in all earlier experiments is actually due to nonpredictive gating processes. This claim cannot be supported based on several observations and arguments. First, the present results showed that the predictive attenuation of touch and gating are different phenomena; if sensory attenuation was the same as tactile gating, then *all* stimuli applied on the moving limb would be attenuated. In contrast, we show that sensory reafference is selectively attenuated compared to sensory exafference, both in a passive and a moving limb. Similar results were shown in [79], where participants attenuated only their reafferent touches and not exafferent touches presented at the same limb simultaneously. Second, Thomas and colleagues [64] provided participants with an arbitrary mapping between the movement of one hand and sensory feedback on the other hand. Using a very similar setup, a study from 2006 [3] found neither attenuation, which is consistent with the predictive account of attenuation, nor enhancement effects, which is inconsistent with those from Thomas and colleagues [64]. Third, it is essential to keep in mind that attenuation has also been shown on an anesthetized limb [8] and for imagined movements [9], further emphasizing the contribution of motor prediction in relation to peripheral feedback for somatosensory attenuation. Finally, it has been shown that when reaching with one arm towards the other, the perceived intensity of an external tactile stimulus applied to the passive target limb is enhanced [92]. Although we believe that future experimentation is needed to understand and replicate the reported enhancement effects [64], we speculate that, as in [92], increased spatial attention or other kinematic differences play a significant role in the somatosensory enhancement reported by Thomas and colleagues [64].

We conclude that the human brain uses two different basic processes to suppress reafferent and exafferent information during movement and rest. This separation of attenuation and gating can explain why although we cannot tickle ourselves because we attenuate our self-tickles, we do tense our muscles when being tickled by others to decrease our sensitivity to the external tickles.

Materials and Methods

Participants
After providing written informed consent, twenty-four participants (12 women and 12 men, 22 right-handed, 1 ambidextrous, 1 left-handed) aged 21-40 years participated. Handedness was assessed using the Edinburgh Handedness Inventory [93]. The sample size was set to twenty-four (24) before data collection commenced based on our previous studies using the same methods

467 [4,10] while ensuring a counterbalanced order of conditions. The Swedish Ethical Review
468 Authority (https://etikprovningsmyndigheten.se/) approved the study (no. 2016/445- 31/2,
469 amendment 2019-04536). All participants provided their written informed consent.
470
General Procedure
Participants sat comfortably on a chair with their arms placed on a table. Their left hands rested
palm up, with their index fingers placed on a molded support. The right arms rested palm down
on top of a set of sponges. In each trial, a motor (Maxon EC Motor EC 90 flat; Switzerland)
delivered two taps (the *test* tap and the *comparison* tap) on the pulp of their left index finger
through a cylindrical probe (25 mm height) with a flat aluminum surface (20 mm diameter)
attached to the motor's lever. A force sensor (FSG15N1A, Honeywell Inc.; diameter, 5 mm;
minimum resolution, 0.01 N; response time, 1 ms; measurement range, 0–15 N) within the
probe recorded the forces applied on the left index finger. Following the presentation of the two
taps, participants had to verbally indicate which tap felt stronger: the first or the second.
481
A wooden surface was placed under the motor and the sensors. This surface was placed on top
of two commercially available drawer runners (IKEA, https://www.ikea.com/us/en/p/besta-
drawer-runner-soft-closing-40348715/). One side of the runners was attached to the table with
Velcro material, and the other side was attached to the bottom side of the surface. With this
configuration, the surface – with the motor, the sensors and the participants' hands – could be
moved forward and backward.
488
489 In all conditions, the *comparison* tap was delivered on the left index finger with a random delay
490 of 800-1500 ms from the *test* tap. In the *attenuation* and *attenuation&gating* conditions (**Fig.
491 1b, d**), the tap of the participants' right index fingers on the force sensor triggered the *test* tap
492 on their left index finger with an intrinsic delay of ≈ 36 ms.
493
In the *gating* and *attenuation&gating* conditions (**Fig. 1c, d**), participants were asked to extend
their elbow upon the auditory 'go' cue. The extension of the elbow moved the platform forward
on the table (**Fig. 1f**). A green tape on the table (**Fig. 1i**) indicated the start position of the
platform, while a red tape indicated its end position. The participants were asked to move the
platform from the start position to the end position (distance = 25 cm). During the movement
of the left arm, the participants received the *test* tap on their left index finger. Before the
condition started, we emphasized to the subjects that their task was to pay attention to the force
that they would receive during the movement, rather than covering exactly the distance between
the lines. Moreover, the participants were trained to perform the movement in approximately
1000-1500 ms after the 'go' cue and then stop. In the *gating* condition, the *test* tap was applied
800 ms after the 'go' cue so that its delivery was during the movement. Similarly, in the
attenuation&gating condition, the participants triggered the *test* tap during the movement. The
comparison tap was applied 800-1500 ms after the test tap to ensure that the participants had
stopped moving. Once the participants gave a response, they returned the platform to the start
position.
509
510 A motion tracking sensor (6DOF Polhemus Fastrak, USA, weight = 9.1 g, dimensions = 2.29
511 cm x 2.82 cm x 1.52 cm) was placed on top of the platform to record the motion of the platform
512 due to the movement of the participants' left arm. The sensor recorded the x, y and z positions
513 at a sampling rate of approximately 120 Hz.
514
Each condition included 70 trials. The test tap was set to 2 N, while the intensity of the
comparison tap was systematically varied among seven different force levels (1, 1.5, 1.75, 2,

2.25, 2.5 or 3 N). Each tap lasted for 100 ms. On every trial, participants verbally indicated
which tap on their left index finger felt stronger: the first (*test*) or the second (*comparison*).
Participants were told not to try to balance their responses (50% first and 50% second), and
they were further instructed to make their best guess if the intensity of the two taps felt similar.

In addition, participants were administered white noise through a pair of headphones to
preclude any sounds created by the motor to serve as a cue for the task. The loudness of the
white noise was adjusted so that participants could clearly hear the auditory cues of the trial. In
all conditions, the view of the pulp of the left index finger was occluded. Participants were
asked to fixate on a cross placed on a wall 2 m opposite them, but they were allowed to look at
the force sensor to guide the movement of the right index finger when needed (**Fig. 1b, d**). No
feedback was provided to the participants about their responses.

Force discrimination analysis
In each condition, the participants' responses were fitted with a generalized linear model using
a *logit* link function (Equation 1):

$$533 \qquad p = \frac{e^{\beta 0 + \beta 1 x}}{1 + e^{\beta 0 + \beta 1 x}} \text{ (Equation 1)}$$

We extracted two parameters of interest: the PSE, which represents the intensity at which the
test tap felt as strong as the comparison tap ($p = 0.5$) and which quantifies the perceived
intensity, and the JND, which reflects the participants' discrimination capacity. Before fitting
the responses, the values of the applied comparison taps were binned to the closest value with
respect to their theoretical values (1, 1.5, 1.75, 2, 2.25, 2.5 or 3 N).

Kinematic analysis.
Both position and velocity data were smoothed with a moving average filter in MATLAB
2018a. Velocity was calculated as the first derivative of position. To calculate the distance
participants ran in every trial of each condition, we calculated the minimum and the maximum
position of the platform during the entire trial duration. The peak trial velocity was defined as
the peak velocity of the entire trial. The peak tap velocity was defined as the peak velocity
during the period that the *test* tap was applied.

Rejection of trials
After data collection, one hundred seventy-three (173) trials out of 6720 (2.57%) were rejected.
First, in thirty-four trials (34), the intensity of the *test* tap (2 N) was not applied accurately (*test*
tap < 1.85 N or *test* tap > 2.15 N), and in sixteen (16) trials, the responses were missing. Second,
we rejected one hundred seventeen (117) trials in total from the *gating* and *attenuation&gating*
conditions because participants either did not move their left arm (or moved it too slowly)
during the *test* tap (mean velocity < 10 cm/s) or they moved it during the *comparison* tap (mean
velocity > 5 cm/s). The thresholds were based on a previous study [40] showing no gating effects
for velocities smaller than 5 cm/s. The analysis was therefore performed with 6547 trials in
total.

Statistical analysis
We used R [94] and JASP [95] to analyze our data. The data normality was assessed using the
Shapiro-Wilk test. Depending on the data normality, we then performed planned comparisons
using either a paired t-test or a Wilcoxon signed-rank test. We report 95% confidence intervals
(CI^{95}) for each statistical test. Effect sizes are given by the partial eta-squared (η_P^2) for the
ANOVAs, by the Cohen's *d* for t-tests or by the matched rank biserial correlation *rrb* for the
Wilcoxon signed-rank tests. In addition, a Bayesian factor analysis using default Cauchy priors

with a scale of 0.707 was carried out for all statistical tests to provide information about the
level of support for the null hypothesis compared to the alternative hypothesis (BF_{01}) given the
data. Finally, correlations were performed using Pearson's coefficient r given that the data were
normally distributed. All tests were two-tailed.
571
Corrections for multiple comparisons
Since our PSE and JND comparisons were planned, we did not apply corrections for multiple
comparisons. However, all results remained exactly the same when applying corrections for
false discovery rate (FDR) [96]. In the correlation analyses, we did apply corrections for multiple
comparisons (FDR) since, although we expected correlations between the PSEs and between
the JNDs, we had no *a priori* hypotheses for correlations between PSEs and JNDs.
578

Acknowledgements

580 We thank L. Miller, M. Hauser and M. Chancel for their feedback on an earlier version of the
581 manuscript. K.K. was supported by the Swedish Research Council (VR Starting Grant 2019-
582 01909 granted to K.K.). This work was supported by the Swedish Research Council, Torsten
583 Söderbergs Stiftelse, and Göran Gustafssons Stiftelse. We thank Kanaka Raghavan for help
584 with the illustrations.
585

Conflicts of interest/Competing interests

587 The authors declare no competing financial interests.
588

Authors' contributions

590 K.K. and H.H.E. conceived and designed the experiment. K.K. collected the data and conducted
591 the statistical analysis. K.K. and H.H.E. wrote the manuscript.
592

References

1. Blakemore, S. J., Frith, C. D. & Wolpert, D. M. Spatio-temporal prediction modulates
the perception of self-produced stimuli. *J. Cogn. Neurosci.* **11**, 551–559 (1999).
2. Bays, P. M., Wolpert, D. M. & Flanagan, J. R. Perception of the consequences of self-
action is temporally tuned and event driven. *Curr. Biol.* **15**, 1125–1128 (2005).
3. Bays, P. M., Flanagan, J. R. & Wolpert, D. M. Attenuation of self-generated tactile
sensations is predictive, not postdictive. *PLoS Biol.* **4**, 281–284 (2006).
4. Kilteni, K., Houborg, C. & Ehrsson, H. H. Rapid learning and unlearning of sensory
delays in self-touch. *Elife* 1–17 (2019). doi:10.7554/eLife.42888.001
5. Shergill, S. S., Bays, P. M., Frith, C. D. & Wolpert, D. M. Two eyes for an eye: the
neuroscience of force escalation. *Science* **301**, 187 (2003).
6. Kilteni, K. & Ehrsson, H. H. Sensorimotor predictions and tool use: Hand-held tools
attenuate self-touch. *Cognition* **165**, 1–9 (2017).
7. Kilteni, K. & Ehrsson, H. H. Body ownership determines the attenuation of self-
generated tactile sensations. *Proc. Natl. Acad. Sci.* 201703347 (2017).
doi:10.1073/PNAS.1703347114
8. Walsh, L. D., Taylor, J. L. & Gandevia, S. C. Overestimation of force during matching
of externally generated forces. *J. Physiol.* **589**, 547–557 (2011).
9. Kilteni, K., Andersson, B. J., Houborg, C. & Ehrsson, H. H. Motor imagery involves
predicting the sensory consequences of the imagined movement. *Nat. Commun.* **9**,
1617 (2018).
10. Kilteni, K., Engeler, P. & Ehrsson, H. H. Efference Copy Is Necessary for the
Attenuation of Self-Generated Touch. *iScience* (2020). doi:10.1016/j.isci.2020.100843
11. Blakemore, S. J., Wolpert, D. M. & Frith, C. D. Central cancellation of self-produced

617		tickle sensation. *Nat. Neurosci.* **1**, 635–640 (1998).
618	12.	Shergill, S. S. *et al.* Modulation of somatosensory processing by action. *Neuroimage*
619		**70**, 356–362 (2013).
620	13.	Kilteni, K. & Ehrsson, H. H. Functional Connectivity between the Cerebellum and
621		Somatosensory Areas Implements the Attenuation of Self-Generated Touch. *J.*
622		*Neurosci.* **40**, 894 LP – 906 (2020).
623	14.	Blakemore, S. J., Frith, C. D. & Wolpert, D. M. The cerebellum is involved in
624		predicting the sensory consequences of action. *Neuroreport* **12**, 1879–1884 (2001).
625	15.	Blakemore, S. J., Wolpert, D. M. & Frith, C. D. The cerebellum contributes to
626		somatosensory cortical activity during self-produced tactile stimulation. *Neuroimage*
627		**10**, 448–459 (1999).
628	16.	Wolpe, N. *et al.* Ageing increases reliance on sensorimotor prediction through
629		structural and functional differences in frontostriatal circuits. *Nat. Commun.* **7**, (2016).
630	17.	Blakemore, S. J., Wolpert, D. M. & Frith, C. Why can't you tickle yourself?
631		*Neuroreport* **11**, R11–R16 (2000).
632	18.	Leavens, D. A. & Bard, K. A. Tickling. *Current Biology* (2016).
633		doi:10.1016/j.cub.2015.06.014
634	19.	Weiskrantz, L., Elliott, J. & Darlington, C. Preliminary observations on tickling
635		oneself. *Nature* **230**, 598–599 (1971).
636	20.	Crapse, T. B. & Sommer, M. A. Corollary discharge across the animal kingdom.
637		*Nature Reviews Neuroscience* **9**, 587–600 (2008).
638	21.	Cullen, K. E. Sensory signals during active versus passive movement. *Curr. Opin.*
639		*Neurobiol.* **14**, 698–706 (2004).
640	22.	Straka, H., Simmers, J. & Chagnaud, B. P. A New Perspective on Predictive Motor
641		Signaling. *Curr. Biol.* **28**, R232–R243 (2018).
642	23.	Schneider, D. M. & Mooney, R. How movement modulates hearing. *Annual Review of*
643		*Neuroscience* (2018). doi:10.1146/annurev-neuro-072116-031215
644	24.	Brooks, J. X. & Cullen, K. E. Predictive Sensing: The Role of Motor Signals in
645		Sensory Processing. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*
646		(2019). doi:10.1016/j.bpsc.2019.06.003
647	25.	Poulet, J. F. A. & Hedwig, B. The cellular basis of a corollary discharge. *Science (80-.*
648		*).* (2006). doi:10.1126/science.1120847
649	26.	Poulet, J. F. A. & Hedwig, B. Corollary discharge inhibition of ascending auditory
650		neurons in the stridulating cricket. *J. Neurosci.* (2003). doi:10.1523/jneurosci.23-11-
651		04717.2003
652	27.	Schneider, D. M., Sundararajan, J. & Mooney, R. A cortical filter that learns to
653		suppress the acoustic consequences of movement. *Nature* (2018). doi:10.1038/s41586-
654		018-0520-5
655	28.	Fukutomi, M. & Carlson, B. A. A History of Corollary Discharge: Contributions of
656		Mormyrid Weakly Electric Fish. *Frontiers in Integrative Neuroscience* (2020).
657		doi:10.3389/fnint.2020.00042
658	29.	Sawtell, N. B. Neural Mechanisms for Predicting the Sensory Consequences of
659		Behavior: Insights from Electrosensory Systems. *Annual Review of Physiology* (2017).
660		doi:10.1146/annurev-physiol-021115-105003
661	30.	Cullen, K. E. The vestibular system: Multimodal integration and encoding of self-
662		motion for motor control. *Trends in Neurosciences* (2012).
663		doi:10.1016/j.tins.2011.12.001
664	31.	Brooks, J. X., Carriot, J. & Cullen, K. E. Learning to expect the unexpected: Rapid
665		updating in primate cerebellum during voluntary self-motion. *Nat. Neurosci.* **18**, 1310–
666		1317 (2015).

667   32.   Roy, J. E. & Cullen, K. E. Dissociating Self-Generated from Passively Applied Head
668   Motion: Neural Mechanisms in the Vestibular Nuclei. *J. Neurosci.* (2004).
669   doi:10.1523/JNEUROSCI.3988-03.2004
670   33.   Angel, R. W. & Malenka, R. C. Velocity-dependent suppression of cutaneous
671   sensitivity during movement. *Exp. Neurol.* (1982). doi:10.1016/0014-4886(82)90244-8
672   34.   Chapman, C. E., Bushnell, M. C., Miron, D., Duncan, G. H. & Lund, J. P. Sensory
673   perception during movement in man. *Exp. Brain Res.* **68**, 516–524 (1987).
674   35.   Post, L. J., Zompa, I. C. & Chapman, C. E. Perception of vibrotactile stimuli during
675   motor activity in human subjects. *Exp. Brain Res.* **100**, 107–120 (1994).
676   36.   Williams, S. R., Shenasa, J. & Chapman, C. E. Time course and magnitude of
677   movement-related gating of tactile detection in humans. I. Importance of stimulus
678   location. *J. Neurophysiol.* **79**, 947–963 (1998).
679   37.   Williams, S. R. & Chapman, C. E. Time course and magnitude of movement-related
680   gating of tactile detection in humans. II. Effects of stimulus intensity. *J. Neurophysiol.*
681   (2000). doi:10.1152/jn.2000.84.2.863
682   38.   Williams, S. R. & Chapman, C. E. Time course and magnitude of movement-related
683   gating of tactile detection in humans. III. Effect of motor tasks. *J. Neurophysiol.* **88**,
684   1968–1979 (2002).
685   39.   Chapman, C. E. & Beauchamp, E. Differential controls over tactile detection in
686   humans by motor commands and peripheral reafference. *J. Neurophysiol.* **96**, 1664–
687   1675 (2006).
688   40.   Cybulska-Klosowicz, A., Meftah, E. M., Raby, M., Lemieux, M. L. & Chapman, C. E.
689   A critical speed for gating of tactile detection during voluntary movement. *Exp. Brain*
690   *Res.* **210**, 291–301 (2011).
691   41.   Gertz, H., Voudouris, D. & Fiehler, K. Reach-relevant somatosensory signals modulate
692   tactile suppression. *J. Neurophysiol.* **117**, 2262–2268 (2017).
693   42.   Fraser, L. E. & Fiehler, K. Predicted reach consequences drive time course of tactile
694   suppression. *Behav. Brain Res.* **350**, 54–64 (2018).
695   43.   Papakostopoulos, D., Cooper, R. & Crow, H. J. Inhibition of cortical evoked potentials
696   and sensation by self-initiated movement in man. *Nature* **258**, 321–324 (1975).
697   44.   Starr, A. & Cohen, L. G. 'Gating' of somatosensory evoked potentials begins before
698   the onset of voluntary movement in man. *Brain Res.* **348**, 183–186 (1985).
699   45.   Rushton, D. N., Roghwell, J. C. & Craggs, M. D. Gating of somatosensory evoked
700   potentials during different kinds of movement in man. *Brain* **104**, 465–491 (1981).
701   46.   Giblin, D. R. SOMATOSENSORY EVOKED POTENTIALS IN HEALTHY
702   SUBJECTS AND IN PATIENTS WITH LESIONS OF THE NERVOUS SYSTEM.
703   *Ann. N. Y. Acad. Sci.* **112**, 93–142 (1964).
704   47.   Chapman, C. E. Active versus passive touch: Factors influencing the transmission of
705   somatosensory signals to primary sornatosensory cortex. *Can. J. Physiol. Pharmacol.*
706   **72**, 558–570 (1994).
707   48.   Lei, Y., Ozdemir, R. A. & Perez, M. A. Gating of sensory input at subcortical and
708   cortical levels during grasping in humans. *J. Neurosci.* (2018).
709   doi:10.1523/JNEUROSCI.0545-18.2018
710   49.   Azim, E. & Seki, K. Gain control in the sensorimotor system. *Curr. Opin. Physiol.* **8**,
711   177–187 (2019).
712   50.   Ghez, C. & Lenzi, G. L. Modulation of sensory transmission in cat lemniscal system
713   during voluntary movement. *Pflügers Arch. Eur. J. Physiol.* **323**, 273–278 (1971).
714   51.   Chapin, J. K. & Woodward, D. J. Modulation of sensory responsiveness of single
715   somatosensory cortical cells during movement and arousal behaviors. *Exp. Neurol.*
716   (1981). doi:10.1016/0014-4886(81)90135-7

52. Jiang, W., Chapman, C. E. & Lamarre, Y. Modulation of the cutaneous responsiveness
of neurones in the primary somatosensory cortex during conditioned arm movements in
the monkey. *Exp. Brain Res.* (1991). doi:10.1007/BF00231455
53. Jiang, W., Lamarre, Y. & Chapman, C. E. Modulation of cutaneous cortical evoked
potentials during isometric and isotonic contractions in the monkey. *Brain Res.* (1990).
doi:10.1016/0006-8993(90)90010-9
54. Seki, K. & Fetz, E. E. Gating of sensory input at spinal and cortical levels during
preparation and execution of voluntary movement. *J. Neurosci.* **32**, 890–902 (2012).
55. Seki, K., Perlmutter, S. I. & Fetz, E. E. Sensory input to primate spinal cord is
presynaptically inhibited during voluntary movement. *Nat. Neurosci.* **6**, 1309–1316
(2003).
56. McNamee, D. & Wolpert, D. M. Internal Models in Biological Control. *Annu. Rev.*
Control. Robot. Auton. Syst. **2**, 339–364 (2019).
57. Collins, D. F., Cameron, T., Gillard, D. M. & Prochazka, A. Muscular sense is
attenuated when humans move. *J. Physiol.* **508**, 635–643 (1998).
58. Juravle, G., Binsted, G. & Spence, C. Tactile suppression in goal-directed movement.
Psychon. Bull. Rev. (2017). doi:10.3758/s13423-016-1203-6
59. Brown, H., Adams, R. a., Parees, I., Edwards, M. & Friston, K. Active inference,
sensory attenuation and illusions. *Cogn. Process.* **14**, 411–427 (2013).
60. Limanowski, J. *et al.* Action-Dependent Processing of Touch in the Human Parietal
Operculum and Posterior Insula. *Cereb. Cortex* (2020). doi:10.1093/cercor/bhz111
61. Saradjian, A. H. Sensory modulation of movement, posture and locomotion.
Neurophysiol. Clin. **45**, 255–267 (2015).
62. Palmer, C. E., Davare, M. & Kilner, J. M. Physiological and Perceptual Sensory
Attenuation Have Different Underlying Neurophysiological Correlates. *J. Neurosci.* **36**,
10803–10812 (2016).
63. Boehme, R., Hauser, S., Gerling, G. J., Heilig, M. & Olausson, H. Distinction of self-
produced touch and social touch at cortical and spinal cord levels. *Proc. Natl. Acad.*
Sci. 201816278 (2019). doi:10.1073/PNAS.1816278116
64. Thomas, E. R., Yon, D., Lange, F. P. de & Press, C. Action enhances predicted touch.
bioRxiv (2020). doi:10.1101/2020.03.26.007559
65. Voss, M., Ingram, J. N., Wolpert, D. M. & Haggard, P. Mere expectation to move
causes attenuation of sensory signals. *PLoS One* **3**, 2–6 (2008).
66. Blakemore, S. J., Smith, J., Steel, R., Johnstone, C. E. & Frith, C. D. The perception of
self-produced sensory stimuli in patients with auditory hallucinations and passivity
experiences: evidence for a breakdown in self-monitoring. *Psychol. Med.* **30**, 1131–
1139 (2000).
67. Shergill, S. S., Samson, G., Bays, P. M., Frith, C. D. & Wolpert, D. M. Evidence for
sensory prediction deficits in schizophrenia. *Am. J. Psychiatry* **162**, 2384–2386 (2005).
68. Shergill, S. S. *et al.* Functional magnetic resonance imaging of impaired sensory
prediction in schizophrenia. *JAMA psychiatry* **71**, 28–35 (2014).
69. Parees, I. *et al.* Loss of sensory attenuation in patients with functional (psychogenic)
movement disorders. *Brain* **137**, 2916–2921 (2014).
70. Wolpe, N. *et al.* Sensory attenuation in Parkinson's disease is related to disease
severity and dopamine dose. *Sci. Rep.* (2018). doi:10.1038/s41598-018-33678-3
71. Macerollo, A. *et al.* Dopaminergic Modulation of Sensory Attenuation in Parkinson's
Disease: Is There an Underlying Modulation of Beta Power? *Front. Neurol.* **10**, 1–14
(2019).
72. Cohen, L. G. & Starr, A. Localization, timing and specificity of gating of
somatosensory evoked potentials during active movement in man. *Brain* (1987).

doi:10.1093/brain/110.2.451

73. Pertovaara, A., Kemppainen, P. & Leppänen, H. Lowered cutaneous sensitivity to
nonpainful electrical stimulation during isometric exercise in humans. *Exp. Brain Res.*
89, 447–452 (1992).

74. Shadmehr, R. & Krakauer, J. W. A computational neuroanatomy for motor control.
Exp. Brain Res. **185**, 359–381 (2008).

75. Kawato, M. Internal models for motor control and trajectory planning. *Curr Opin*
Neurobiol **9**, 718–727 (1999).

76. Todorov, E. Optimality principles in sensorimotor control. *Nat. Neurosci.* **7**, 907–15
(2004).

77. Wolpert, D. M. & Ghahramani, Z. Computational principles of movement
neuroscience. *Nat. Neurosci.* **3**, 1212–1217 (2000).

78. Bays, P. M. & Wolpert, D. M. Computational principles of sensorimotor control that
minimize uncertainty and variability. *J. Physiol.* **578**, 387–396 (2007).

79. Bays, P. M. & Wolpert, D. M. Predictive attenuation in the perception of touch. in
Sensorimotor Foundations of Higher Cognition (eds. Haggard, E. P., Rosetti, Y. &
Kawato, M.) **22**, 339–358 (Oxford University Press, 2008).

80. Wolpert, D. M. & Flanagan, J. R. Motor prediction. *Curr. Biol.* **11**, R729–R732 (2001).

81. Diedrichsen, J., Verstynen, T., Hon, A., Lehman, S. L. & Ivry, R. B. Anticipatory
adjustments in the unloading task: Is an efference copy necessary for learning? *Exp.*
Brain Res. **148**, 272–276 (2003).

82. Klaffehn, A. L., Baess, P., Kunde, W. & Pfister, R. Sensory attenuation prevails when
controlling for temporal predictability of self- and externally generated tones.
Neuropsychologia (2019). doi:10.1016/j.neuropsychologia.2019.107145

83. Bäß, P., Jacobsen, T. & Schröger, E. Suppression of the auditory N1 event-related
potential component with unpredictable self-initiated tones: Evidence for internal
forward models with dynamic stimulation. *Int. J. Psychophysiol.* (2008).
doi:10.1016/j.ijpsycho.2008.06.005

84. Blakemore, S. J., Wolpert, D. M. & Frith, C. D. The cerebellum contributes to
somatosensory cortical activity during self-produced tactile stimulation. *Neuroimage*
10, 448–459 (1999).

85. Miall, R. C. & Wolpert, D. M. Forward models for physiological motor control. *Neural*
Networks **9**, 1265–1279 (1996).

86. Wolpert, D. M., Miall, R. C. & Kawato, M. Internal models in the cerebellum. *Trends*
in Cognitive Sciences **2**, 338–347 (1998).

87. Shadmehr, R., Smith, M. a & Krakauer, J. W. Error correction, sensory prediction, and
adaptation in motor control. *Annu. Rev. Neurosci.* **33**, 89–108 (2010).

88. Therrien, A. S. & Bastian, A. J. The cerebellum as a movement sensor. *Neurosci. Lett.*
0–1 (2018). doi:10.1016/j.neulet.2018.06.055

89. Adams, R. a., Shipp, S. & Friston, K. J. Predictions not commands: Active inference in
the motor system. *Brain Struct. Funct.* **218**, 611–643 (2013).

90. Friston, K. The free-energy principle: a unified brain theory? *Nat. Rev. Neurosci.* **11**,
127–38 (2010).

91. Pickering, M. J. & Clark, A. Getting ahead: Forward models and their place in
cognitive architecture. *Trends Cogn. Sci.* **18**, 451–456 (2014).

92. Voudouris, D. & Fiehler, K. Enhancement and suppression of tactile signals during
reaching. *J. Exp. Psychol. Hum. Percept. Perform.* (2017). doi:10.1037/xhp0000373

93. Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh inventory.
Neuropsychologia **9**, 97–113 (1971).

94. Core Team, R. R: A language and environment for statistical computing. *R Foundation*

817   *for Statistical Computing* (2018). doi:10.1007/978-3-540-74686-7

818   95.   JASP & JASP Team. JASP. *[Computer software]* (2019).

819   96.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

820   Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* (1995).

821   doi:10.1111/j.2517-6161.1995.tb02031.x