Predictive attenuation of touch and tactile gating are distinct perceptual phenomena

Konstantina Kilteni[1][*] and H. Henrik Ehrsson[1]

[1]Department of Neuroscience, Karolinska Institutet, Solnavägen 9, 17165 Stockholm, Sweden
[*]Correspondence and requests for materials should be addressed to konstantina.kilteni@ki.se

In recent decades, research on somatosensory perception has led to two important observations. First, self-generated touches that are predicted by voluntary movements become attenuated compared to externally generated touches of the same intensity (attenuation). Second, externally generated touches feel weaker and are more difficult to detect during movement than at rest (gating). Researchers today often consider gating and attenuation to be the same suppression process; however, this assumption is unwarranted because, despite more than forty years of research, no study has combined them in a single paradigm. We quantified how people perceive self-generated and externally generated touches during movement and rest. We demonstrate that whereas voluntary movement gates the precision of both self-generated and externally generated touch, the amplitude of self-generated touch is selectively attenuated compared to externally generated touch. We further show that attenuation and gating do not interact and are not correlated, and we conclude that they represent distinct perceptual phenomena.

Let us imagine you are at your doctor for a medical examination. Upon her request, you apply pressure with your index finger to your leg to indicate where exactly you feel the pain. The pressure you feel on your leg and the tip of your finger is the feedback from your voluntary finger movement, and it is called *somatosensory reafference*. To reproduce and confirm your sensations, imagine that the doctor now applies pressure with her index finger to the same spot on your leg. This pressure is generated by the doctor and not by you, and it is called *somatosensory exafference*. Now imagine that the doctor asks you to first keep your leg relaxed and then flex and extend it while you or her continuously apply pressure to your leg. You can therefore experience your reafferent or her exafferent touches on your leg while it is moving or resting. Distinguishing between these four conditions is fundamental for your sensorimotor control; your nervous system must know both the source of the touch and the state of your limb to make appropriate use of the sensory feedback. A cutaneous mechanoreceptor in your peripheral nervous system, however, cannot distinguish whether a touch is reafferent or exafferent, and thus, this distinction must be made centrally, where tactile signals from the skin, sensory information from muscles and joints, and information from motor commands are available. How, then, does the central nervous system classify somatosensory signals during movement?

Several experimental studies in humans have shown that the brain *attenuates* somatosensory reafference compared to exafference. In behavioral research, this refers to participants perceiving self-generated strokes, forces, or taps as weaker than external equivalents of the same intensity [1–10]. This somatosensory attenuation has been related to reduced activity in the secondary somatosensory cortex [11–13] and the cerebellum [11,13,14] and increased connectivity between the two areas [13,15] during self-generated touches compared to externally generated touches. Somatosensory attenuation was found in 98% or 315 out of 322 people across a wide age range [16], and it is considered one of the reasons why we cannot tickle ourselves [17–19].

Sensory attenuation is not exclusive to humans; similar strategies are used by other species across the animal kingdom (for reviews, see [20–24]). For example, during self-chirping, the cricket's central auditory processing is inhibited (both presynaptically and postsynaptically) in phase with the insect's chirps to prevent desensitization of its auditory system while maintaining sensitivity to external sounds [25,26]. In mice, auditory cortical responses in response to self-generated sounds are attenuated, and this attenuation is present only for the tone frequencies the animal has associated with its locomotion and absent when the same sounds are externally produced [27]. A weakly electric fish [28] is able to respond exclusively to externally generated electrical discharges by attenuating its predicted electrosensory reafference [21,29]. In primates, activity in the vestibular nucleus in response to vestibular reafference is attenuated during active head movements compared to passive head movements, allowing the animal to maintain its head and body posture and activate vestibular-related reflexes when appropriate [20,21,30–32].

At the same time, another branch of experimental research has shown that somatosensory sensitivity in response to externally generated stimuli is *gated* during and before a voluntary movement. In human psychophysical research, this phenomenon of *movement-related tactile gating* or *tactile suppression* manifests as an elevation of the detection threshold [33–38], a decrease in the detection rate [33–35,39–43,] a decrease in the detection precision [36,38,44,45], and a decrease in the subjective intensity of externally generated stimuli [35,40,46] when the stimulated body part moves compared to when it is at rest. Several electrophysiological studies have shown that this gating reflects inhibition of somatosensory evoked potentials during active movement compared to rest at subcortical and cortical sites along the somatosensory pathway [46–51]. Similar to somatosensory attenuation, tactile gating is a biologically preserved mechanism that is seen across different species [52]. For example, responses recorded at the cat's medial lemniscus evoked by nerve stimulation are suppressed prior to and during limb movements [53]. Similarly, the transmission of cutaneous afferent signals to the primary somatosensory cortex is suppressed in rats during movement compared to rest [54]. In monkeys, the gating of cutaneous afferent input during active movement has been observed in both the primary somatosensory cortex [55–57] and the spinal cord [57,58].

Somatosensory attenuation and tactile gating share two important conceptual similarities. First, they both refer to modulation, either in terms of magnitude or precision, of the perception of cutaneous stimuli during movement. Second, they have been assigned the same functional role [41]: to reduce the flow of afferent information that can be predicted from the motor command so that detection of external input that can be biologically important, such as touches caused by predators [17,24,59], or input that is task-relevant for the upcoming or ongoing movement [48,50,60] is facilitated.

Importantly, however, the two phenomena present one striking difference. Somatosensory attenuation relates to somatosensory reafference, that is, touches caused *by* our voluntary movement. In contrast, gating relates to somatosensory exafference, that is, external touches occurring *during* our voluntary movement. Nevertheless, somatosensory research, including very recent studies and review articles, e.g., [36,37,61–67], often treats the two phenomena as a single generalized suppression strategy of the brain and uses the terms attenuation and gating interchangeably. If the two phenomena are indeed different, this false equivalence is detrimental for understanding human sensorimotor control. First, it prevents advancements in our understanding at the computational level because researchers try to explain gating (e.g., [36]) and attenuation (e.g., [68]) using the computational processes proposed for attenuation and gating,

respectively. Similarly, at the neurobiological level, researchers intermix neural correlates of gating and attenuation (e.g., [65,66]) because they assume that these measure the same single phenomenon. Second, this confusion becomes particularly disadvantageous in clinical studies using gating and attenuation when interpreting findings of sensorimotor deficits in patients with schizophrenia [69–71], functional movement disorders [72] or Parkinson's disease [48,73,74].

Are these phenomena the same, or do they represent two fundamentally distinct processes? Does the brain treat all sensory stimuli similarly during movement, independent of whether they are reafferent or exafferent? To the best of our knowledge, there has been no previous attempt to simultaneously test attenuation and gating with the same stimulus and psychophysics task. Here, in a single experimental design, we investigated the perception of touches applied on the left hand while manipulating whether the left arm was in movement or resting (left limb state). We additionally manipulated whether the touches were reafferent, generated by the right hand, or exafferent, generated by an external source (origin of touch). We reasoned that if the two phenomena are the same, they should influence somatosensory perception in similar ways. Our results demonstrate that this is not the case: voluntary movement reduces the somatosensory precision of both sensory reafference and exafference (gating), while the perceived amplitude of sensory reafference is selectively attenuated compared to that of exafference (attenuation). Notably, the two phenomena do not correlate with each other or interact when present simultaneously. Thus, collectively, our results show that gating and attenuation are two separate processes that can be experimentally dissociated.

Results

Participants rested their left hands palm up with their index fingers on a molded support, and their right hands were placed palm down on top of a set of sponges (**Fig. 1a-d**). In all conditions, they performed a force-discrimination task [3,4,10,70]: in each trial, a motor delivered two taps (the *test* tap and the *comparison* tap) on the pulp of their left index finger, and they were asked to verbally indicate which tap felt stronger (**Fig. 1e-h**). While the *test* tap had a fixed intensity (2 N), the intensity of the *comparison* tap randomly changed in every trial (1, 1.5, 1.75, 2, 2.25, 2.5, or 3 N). An auditory 'go' signal indicated the trial's onset and the onset of the response period.

In all conditions, our experimental manipulation exclusively concerned the *test tap*; the *comparison* tap was always externally triggered and delivered on the relaxed left arm, serving, therefore, as a reference stimulus. In a factorial design, we controlled for whether the left arm moved (**Fig. 1c, d**) or remained relaxed (**Fig. 1a, b**) during the *test* tap and whether the *test* tap was produced by the right hand (self-generated) (**Fig. 1b, d**) or not (externally generated) (**Fig. 1a, c**). This design resulted in four experimental conditions, the order of which was fully counterbalanced (**Fig. 1a-d**).

In the *baseline* condition (**Fig. 1a**), participants did not move their limbs but passively received the *test* and the *comparison* taps on the left index. This control condition was used to assess the participants' somatosensory perception in the absence of any movement [2,4,10]. In the *attenuation* condition (**Fig. 1b**), participants actively tapped with their right index finger a force sensor placed on top of their left index finger. The tap of their right index finger on the force sensor triggered the *test* tap on their left index finger. This classic condition was used to assess the perception of a self-generated tap on a passive limb [2,4,10]. In the *gating* condition (**Fig. 1c**), participants were asked to continuously move their left arm forward, sliding the experimental setup with the motor and the force sensors between a start and an end position (distance 25 cm)

at a comfortable velocity of approximately 20 cm/sec (**Fig. 1i-j, Supplementary Fig. S1, S2**). During this movement, participants received the *test* tap on their left index finger. This is a gating condition because it assesses the perception of an externally generated tap on a moving limb[39,40,42,75]. Finally, in the *attenuation&gating* condition (**Fig. 1d**), the participants performed the same movement with their left arm, but they were additionally asked to actively tap with their right index finger on the force sensor that triggered the *test* tap on their left index finger during the movement. The force sensor was attached to the experimental setup and moved together with the left hand. This condition combines the gating and attenuation phenomena since it is used to assess the perception of a self-generated tap on a moving limb.

Fig. 1. Experimental conditions. Two factors were manipulated in the experiment, resulting in four experimental conditions: whether the left arm was in rest (**a**, **b**) or in movement while receiving the *test* tap (**c**, **d**) (state of the left limb) and whether the *test* tap was externally triggered by the motor (**a**, **c**) or self-triggered by the participants' right hand (**b**, **d**) (origin of touch). In all four conditions (**e-h**), the participants received the two taps (*test* and *comparison* tap) on the pulp of their left index fingers from the motor (black stars), and they had to verbally indicate which was stronger: the first or the second tap. In the *attenuation* and *attenuation&gating* conditions, the participants self-triggered the test tap on their left index finger by moving their right arm to tap a sensor with their right index finger (blue stars). **(i)** In the *gating* and *attenuation&gating* conditions (**c** and **d**), the participants extended their left arm from a starting position to an ending position, sliding the experimental setup along two drawer runners.

171 During this movement, participants experienced the *test* tap. A motion sensor recorded the position of
172 the platform in time. **(f)** Example position traces recorded by the motion sensor for the movements of
173 one participant during the *attenuation&gating* condition. **(g)** Responses of a representative participant
174 and fitted logistic models for the four experimental conditions. The data points were horizontally jittered
175 to avoid overlapping.
176
The participant's responses in each condition were fitted with a generalized linear model (**Fig.**
1k, **Supplementary Fig. S3**), and two parameters of interest were extracted: the point of
subjective equality (PSE), which represents the intensity at which the test tap felt as strong as
the comparison tap ($p = 0.5$), and the just noticeable difference (JND), which reflects the
participants' sensitivity in force discrimination. A lower PSE in an experimental condition
indicates that the test tap felt weaker in that condition. A higher JND in an experimental
condition indicates that the discrimination sensitivity was lower in that condition (i.e., a larger
difference in the force intensities needed to be detected). We hypothesized that the two
phenomena are different and, thus, that they affect the PSE and JND differently. Specifically,
we expected a decrease in the perceived magnitude (lower PSE) for conditions with sensory
reafference (*attenuation* and *attenuation&gating* conditions), with no effects on the sensory
precision of the participants (JND) when they received reafference on a still limb (*attenuation*
condition). That is, a self-generated touch will feel weaker than an externally generated touch,
but there will be no effect on somatosensory precision. In contrast, we predicted a decrease in
the somatosensory precision (higher JND) of both sensory reafference and exafference for
conditions where the limb that receives the touches moves (*gating*, *attenuation&gating*). That
is, the precision with which a touch (either self-generated or externally generated) is perceived
is lower on a moving limb because of the additional kinaesthetic, proprioceptive, tactile signals
from the movement. A small, if any, effect in the PSE was expected for the *gating* condition.
We hypothesized that the attenuation phenomenon mainly affects the PSE and not the JND, and
the gating phenomenon mainly affects the JND and not the PSE. Our hypotheses were
supported by the data (**Fig. 2a-g**).
199

Fig. 2. Experimental results. (a, b) Bar graphs showing the PSEs and JNDs (mean ± SEM) for each condition. A lower PSE value indicates a lower perceived magnitude, while a higher JND value indicates lower somatosensory sensory precision. Reafferent touches produced by the movement of the right arm (*attenuation* and *attenuation&gating* conditions) were associated with a significant drop in the PSEs, while movement of the left arm that receives the touches (*gating* and *attenuation&gating*) results in a significant increase in the JNDs. A small decrease in PSE was also observed in the *gating* and *attenuation&gating* conditions compared to the *baseline* and *attenuation* conditions, respectively. (c) Group psychometric functions for each condition generated using the mean PSE and the mean JND across participants. The leftward shift of the curves for the *attenuation* and *attenuation&gating* conditions illustrates that somatosensory reafference is perceived as weaker than the exafference. The flattening of the curves for the *gating* and *attenuation&gating* conditions illustrates the worsening in somatosensory precision of both the reafference and exafference on a moving limb. (d, e) Line plots illustrating the decreases in PSEs when experiencing reafferent touches compared to exafferent touches when the left arm is still (d) and when the left arm moves (e). (f, g) Line plots illustrating the increases in JNDs when receiving touches on a moving limb compared to rest when touches are exafferent (f) and when touches are reafferent (g).

We performed repeated-measures ANOVA on the PSEs with the origin of the touch (reafference *vs* exafference) and the state of the left limb (movement *vs* rest) as factors. This revealed a significant main effect of the origin of the touch ($F(1, 23) = 36.10$, $p < 0.001$, $\eta_p^2 = 0.611$), a significant main effect of the left limb state ($F(1, 23) = 13.91$, $p = 0.001$, $\eta_p^2 = 0.377$) and a nonsignificant interaction ($F(1, 23) = 0.26$, $p = 0.615$, $\eta_p^2 = 0.011$) (Fig. 2a). A Bayesian repeated-measures ANOVA further concluded against the interaction term by supporting the model without the interaction term compared to the full factorial ($BF_{M1} / BF_{M2} = 3.631$). The *attenuation* condition produced a significant drop in the PSE compared to the *baseline* condition ($n = 24$, $t(23) = -5.908$, $p < 0.001$, *Cohen's d* $= -1.206$, $CI^{95} = [-0.332, -0.160]$, $BF_{01} < 0.0003$)

(**Fig. 2d**). This replicates previous attenuation findings indicating that a self-generated tap on a
passive limb feels weaker than a tap of the same intensity but of an external origin [2,4–7,10,13,16].
Similarly, the PSE in the *attenuation&gating* condition was significantly lower than that in the
gating condition ($n = 24$, $t(23) = -5.032$, $p < 0.001$, *Cohen's d* $= -1.027$, $CI^{95} = [-0.377, -0.157]$,
$BF_{01} = 0.002$) (**Fig. 2e**), extending the previous conclusion to when the receiving limb is
moving. Together, these two contrasts show that reafferent (self-generated) touches feel weaker
than exafferent touches, both when the receiving hand is in movement and when it is at rest.
234
The *gating* and *attenuation&gating* conditions also resulted in a significant drop in the PSE
compared to that in the *baseline* condition ($n = 24$, $t(23) = -2.409$, $p = 0.024$, *Cohen's d* $= -$
0.492, $CI^{95} = [-0.141, -0.011]$) and the *attenuation* condition ($n = 24$, $V = 55$, $p = 0.005$, $rrb = -$
0.633, $CI^{95} = [-0.161, -0.022]$), respectively. However, these decreases were quite modest ($\cong$
30% of the weakening produced by the *attenuation* condition) and supported only by anecdotal
evidence from Bayesian statistics ($BF_{01} = 0.433$ and $BF_{01} = 0.738$, respectively). Together,
these contrasts suggest that exafferent touches can feel slightly weaker on a moving limb than
on a passive limb, in agreement with previous findings regarding tactile gating [35,40,46].
Nevertheless, when compared with tactile reafference, the perceived magnitude of tactile
exafference is not strongly decreased.
245
When testing for the effects of the conditions on the somatosensory precision of the participants
(JND), there was a significant main effect of the state of the left limb ($F(1, 23) = 17.1$, $p <$
0.001, $\eta_p^2 = 0.426$), but there was neither a significant main effect of the origin of touch ($F(1,$
$23) = 0.52$, $p = 0.478$, $\eta_p^2 = 0.022$) nor a significant interaction ($F(1, 23) = 0.06$, $p = 0.809$, η_p^2
$= 0.003$) (**Fig. 2b**). As with PSEs, the absence of interaction for the JNDs was further supported
by Bayesian repeated-measures ANOVA, which provided evidence against the interaction term
($BF_{M1} / BF_{M2} = 3.522$). The *attenuation* condition did not result in any change in the JND with
respect to that in the *baseline* condition ($n = 24$, $t(23) = 0.331$, $p = 0.744$, *Cohen's d* $= 0.068$,
$CI^{95} = [-0.034, 0.047]$), and this was substantially confirmed by Bayesian analysis ($BF_{01} =$
4.432). In alignment, no significant differences in the JND were detected between the *gating*
and the *attenuation&gating* conditions ($n = 24$, $t(23) = 0.72$, $p = 0.481$, *Cohen's d* $= 0.146$, CI^{95}
$= [-0.024, 0.05]$), which was again confirmed by Bayesian analysis ($BF_{01} = 3.691$). Together,
these two contrasts demonstrate that receiving sensory reafference *per se* is not accompanied
by worsening of sensory precision on the receiving limb.
260
In contrast, moving the limb while receiving an external touch (*gating* condition) resulted in a
significant increase in the JND compared to that in the *baseline* condition ($n = 24$, $t(23) = 3.134$,
$p = 0.005$, *Cohen's d* $= 0.640$, $CI^{95} = [0.019, 0.093]$, $BF_{01} = 0.108$) (**Fig. 2f**). This was further
confirmed by a significant increase in JND in the *attenuation&gating* condition compared with
the *attenuation* condition ($n = 24$, $t(23) = 2.984$, $p = 0.007$, *Cohen's d* $= 0.609$, $CI^{95} = [0.019,$
$0.106]$, $BF_{01} = 0.146$) (**Fig. 2g**). Together, these two differences indicate that voluntary
movement *per se* deteriorates the precision with which reafferent and exafferent stimuli are
perceived on the moving limb.
269
Together, our results indicate that predicting the sensory consequences of a voluntary
movement produces a decrease in the perceived magnitude of sensory reafference (PSE)
without a concomitant worsening of somatosensory precision (JND). In contrast, voluntary
movement leads to a decrease in somatosensory precision (JND) for both sensory reafference
and exafference. These effects are observed in the group psychometric fits (**Fig. 2c**). To further
illustrate that somatosensory attenuation affects the amplitude (PSE), while tactile gating
affects the precision (JND) and not *vice versa*, we tested whether we could better predict the

participants' performance in the *attenuation&gating* condition when using the PSE from the *attenuation* condition and the JND from the *gating* condition than the PSE from the *gating* condition and the JND from the *attenuation* condition. Indeed, the first model was significantly better: $n = 24$, $V = 39$, $p < 0.001$, $rrb = -0.74$, $CI^{95} = [-68.980, -12.736]$, $BF_{01} = 0.443$ (**Supplementary Fig. S4, Fig. 3a-b**).

In the abovementioned ANOVAs, there were no significant interactions between the two factors (the origin of touch and the state of the limb), neither for the PSEs nor for the JNDs, according to both frequentist and Bayesian analyses. This indicates that the effect of the left limb state was not influenced by the effect of the origin of touch, neither for the PSE nor for the JND. Instead, the two effects were summed when simultaneously present, and there were no superadditive effects (interactions), which is in line with the proposal that they are two independent processes. To further test whether the effects produced by each phenomenon relate to each other at all, we performed a correlation analysis of the PSEs and JNDs. No significant correlations were detected between any of the PSEs and any of the JNDs (all *p*-values > 0.225, $BF_{01} = [1.971, 3.950]$). The only significant correlation found for PSEs was between the PSE in the *attenuation* condition and the PSE in the *attenuation&gating* condition ($t(22) = 4.89$, $r = 0.722$, $p < 0.001$, $CI^{95} = [0.449, 0.871]$, $BF_{01} = 0.002$) (**Fig. 3c**). That is, the weaker the participants perceived the magnitude of their self-generated touch during rest, the weaker the magnitude of their self-generated touch during movement felt. Given that the PSEs significantly dropped in these two conditions and that these decreases correlated with each other, this result provides further support that their common experimental denominator, i.e., the reafferent nature of the touch, was responsible for the drop in the PSE and thus for the attenuation phenomenon. In contrast, the JND in the *gating* condition was significantly correlated only with the JND in the *attenuation&gating* condition ($t(22) = 3.47$, $r = 0.595$, $p = 0.008$, $CI^{95} = [0.252, 0.805]$, $BF_{01} = 0.047$) (**Fig. 3d**). This specific correlation means that the worse the somatosensory precision of an external touch when participants moved their receiving hand, the worse the sensory precision for a self-generated touch during the same movement of the receiving hand. Given that the JNDs significantly increased only in these two conditions and that these increases correlated with each other, this result provides evidence that their common experimental denominator, i.e., the left limb's movement that receives the touches, was responsible for the increase in the JND.

Fig. 3. Model predictions and scatterplots for PSEs and JNDs. (a) Average participants' responses in the *attenuation&gating* condition (yellow) and average predicted responses using the parameters from the *attenuation* and *gating* conditions (blue). The responses depicted in dark blue were for the PSE of the *attenuation* condition and the JND from the *gating* condition, while the responses depicted in light blue were for the PSE of the *gating* condition and the JND of the *attenuation* condition. The error bars and ribbons represent 95% confidence intervals. **(b)** For each participant, we estimated the Bayesian information criterion (BIC) of the fitted logistic model in the *attenuation&gating* condition and the two models with fixed parameters from the *attenuation* and *gating* conditions. The model using the PSE of the *attenuation* condition and the JND of the *gating* condition was a significantly better model than the one using the PSE of the *gating* condition and the JND from the *attenuation* condition. The upper right panel represents the same data after excluding one participant corresponding to the outlier for illustration purposes. The exclusion of the outlier did not change the statistical results. **(c)** The participants' PSEs in the *attenuation* condition were significantly correlated with those in the *attenuation&gating* condition. **(d)** The participants' JNDs in the *gating* condition were significantly correlated with those in the *attenuation&gating* condition. There were no significant correlations between PSEs and JNDs.

A common finding in tactile gating studies is that the gating effects, both behavioral and electrophysiological, are stronger with higher movement velocities [33,36,39,42,76]; that is, the faster the limb movements are, the worst the perception of the moving limb. Therefore, one could

hypothesize that any differences observed between the *gating* and *attenuation&gating* conditions might be due to differences in the velocity of the participants' movements. Since there were no significant JND differences in these two conditions (**Fig. 2**), which was further supported by Bayesian analysis, this concern can be ruled out. However, one can argue that the PSE in the *attenuation&gating* condition was lower than the PSE in the *attenuation* condition because the participants moved faster in the *attenuation&gating* condition and not because of the reafferent nature of the touch. This concern can also be ruled out since, rather than faster, we observed that the participants moved slightly slower in the *attenuation&gating* condition (20.3 ± 0.003 cm/s) than in the *gating* condition (23.3 ± 0.003 cm/s): (peak trial velocity; $t(23) = -4.062$, $p < 0.001$, *Cohen's d* = -0.829, $CI^{95} = [-0.001, -0.004]$, $BF_{01} = 0.015$). Although the total distances the participants ran with their left arm were comparable, participants in the *attenuation&gating* condition moved slower because they had to coordinate both their arms to tap the sensor that the left arm moves with the right hand (**Fig. 1d**). This difference was further confirmed when looking at the peak velocities during the *test* tap between the two conditions (*gating*: 22.2 ± 0.003 cm/s; *attenuation&gating*: 19 ± 0.003 cm/s) (**Supplementary Fig. S5**). Therefore, these kinematic differences cannot explain the significant decrease in the PSEs in the *attenuation&gating* condition compared to the *gating* condition, but they actually underscore the importance of reafference in somatosensory attenuation.

Discussion

The present study contrasted the conditions of attenuation and gating in a single experimental paradigm to investigate the relationship between the two phenomena. To this end, we independently manipulated the origin of the touch (reafference *vs* exafference) and the state of the receiving limb (movement *vs* rest), studying all four possible combinations of these levels. We replicated the classic phenomenon of predictive attenuation of touch [1,2,16,3–10] by showing that somatosensory reafference feels weaker than somatosensory exafference. Importantly, however, this decrease in the perceived amplitude (PSE) was not accompanied by a concomitant worsening of somatosensory precision (JND). That is, participants had the same discrimination capacity (JND) for both reafferent and exafferent touches applied on their passive limb, a conclusion supported by Bayesian statistics. Nevertheless, when the limb that received the touches moves, this voluntary movement *per se* led to a decrease in somatosensory precision (JND) for both reafferent and exafferent touches, replicating the classic tactile gating phenomenon [33–37,39,41,42]. The two effects did not correlate and did not interact, but were summed when present together.

The main conclusion of the present study is that the predictive attenuation of touch and tactile gating are two distinct perceptual phenomena. Our findings can help conciliate several previous observations on gating and attenuation that have been studied in isolation during recent decades. First, attenuation is observed not only on the active limb [8] but also on a passive limb (the contralateral hand in the present study), as long as the contact between the body parts is predicted by voluntary movement [2–7,9,10,13,16]. In contrast, there is abundant evidence that external touches applied to the limb contralateral to the limb that moves are not gated [34,45,46,48,77,78]. Second, a touch that results from a passive movement [10] or touches that are simultaneously presented in both hands (double touch) [70] are not attenuated. In contrast, gating effects have been repeatedly demonstrated for passive movements, both electrophysiologically [48] and behaviorally [34,41,75]. Third, self-generated tactile signals are attenuated as long as they are presented at the timing predicted by the action [1,2,4]; even a 100 ms delay between the movement and its tactile feedback greatly reduces the attenuation of the latter. In contrast, externally

generated stimuli are gated with less temporal sensitivity; for example, gating is observed for stimuli presented at several (unpredicted) times during the movement [48], at the movement onset [43], and importantly, even at hundreds of milliseconds before the movement onset [39,41,43,45]. Fourth, whereas gating shows no specificity for the type of motor activity and manifests both during isotonic [33–35,46] and isometric [35,78] contractions, attenuation is motor command-specific; a consistent but arbitrary and unnatural mapping between the motor command and the touch, for example, moving a joystick with one hand to produce touch on the other, does not produce attenuation [5,6,72]. These results are not contradictory in light of our findings; instead, they refer to different perceptual phenomena.

Our study is the first to behaviorally dissociate the predictive attenuation of touch and tactile gating. A previous study [79] found that the perceived intensity of an electrical stimulus applied to the tip of the relaxed left index finger does not change if this finger receives a self-produced force generated by the right index finger. However, in that study, the perception of touch was tested on a resting limb only, which is already known not to elicit gating effects [34,45,46,48,77,78]. Therefore, the two phenomena were not disentangled because they were not tested on the same limb as in the current study. Similarly, another study [65] measured electrophysiological responses (somatosensory evoked potentials) to external electrical stimulation on the right and the left wrist, while the participants produced a self-generated force with their right index finger on their left index finger. The authors did not measure the perception of external touches during movement (gating) but only the perception of self-generated touches during rest (attenuation) and could thus not dissociate the two phenomena behaviorally. Moreover, the authors observed modulation of the late electrophysiological responses depending on whether the participants produced a self-generated force, and they suggested that the attenuation and gating might share the same mechanism: that of reduced sensory precision. However, the present study demonstrates that sensory precision is reduced during gating and not during attenuation, suggesting that a common mechanism governing both phenomena is unlikely.

Motor control relies on integrating afferent sensory information with efferent motor signals [79]. Distinguishing between gating and attenuation is fundamentally important for motor control theories because it can indicate a different integration or weighting mechanism of the motor and sensory information depending on the context. One well-established computational framework inspired by engineering approaches poses that the brain produces motor commands through an inverse model [80] or controller [81]. A copy of the motor command, termed 'efference copy', is used by a forward model to predict the expected sensory feedback of the movement, which is then combined with the actual sensory input to estimate the state of the body [59,80–82]. With respect to the attenuation of sensory reafference, it has been proposed that the prediction signal of the forward model is used to 'cancel' the sensory reafference [17,59,83–85]. In other words, central motor processes play a more important role in somatosensory attenuation than actual sensory feedback. Support for this comes from studies showing that conditions that present highly predictable touches but in the absence of movement do not yield attenuation [2,10] (see also [86–88] for similar conclusions). The dependence of attenuation on action prediction was further shown when participants attenuated the touches applied to one hand that were predicted by their other hand's movement, even when the two hands unexpectedly failed to make contact [3]. In further agreement, neuroimaging studies on somatosensory attenuation consistently report activation of the cerebellum [11–15], a structure that is associated with motor prediction [79,89–92]. Our findings of a reduced perceived magnitude of somatosensory reafference compared to exafference (i.e., lower PSEs in the *attenuation* and *attenuation&gating* conditions compared to the *baseline* and *gating* conditions, respectively) support this forward model mechanism for the attenuation of self-generated input.

431
In contrast, this computational account that relies on action prediction, efference copy, and
internal forward models is inapplicable for tactile gating since gated touches can be of
exafferent nature and occur at any (unpredictable) time during movement, even before
movement onset [39,40,50,75]. In other words, there is no information that the brain can use to predict
exafferent touches with the forward model because there is no causal relationship between the
motor command and sensory input. This observation agrees with the proposal that peripheral
afferent signals from muscle spindles and joint afferents play the major role in gating [39,48] and
that gating effects have also been observed during passive movement, without significant
differences from active movements [34,75]. Then, if efference copy is not the basis for gating, how
can the gating effects be computationally explained?
442
The alternative computational framework of active inference has been proposed to explain
somatosensory attenuation [62]; however, this was based on an assumed equivalence between
attenuation and gating. The active inference approach refutes the necessity of an efference copy
and emphasizes the importance of a generative model and reflex arcs in the place of forward
and inverse models and controllers [93,94]. According to the active inference account, the brain
predicts the sensory input that would be expected from a specific action, and the body moves
to fulfill these sensory predictions. Motor commands are thus conceptually replaced by
proprioceptive predictions, and action occurs as a way to minimize the proprioceptive
prediction errors when the movement has not yet been executed [95]. A major role is assigned to
the precision (i.e., reliability), which weights these sensory prediction errors depending on the
context and can be manipulated through attention allocation. Within this computational
architecture, attenuation of somatosensory reafference is seen as a reduction in the precision of
somatosensory evidence during movement to allow the expression of proprioceptive
predictions that trigger the movement [62]. In other words, the agent attends away from
somatosensory input to execute the movement. However, this proposal does not address the
attenuation of sensory reafference with respect to exafference since the agent should
theoretically attend away from *all* somatosensory inputs independent of their source. Moreover,
the proposal cannot explain why increasing the precision of the proprioceptive prediction errors
on the hand that is to move reduces the precision of somatosensory prediction errors on the
contralateral limb that is not meant to move and thus, there should not exist any proprioceptive
predictions about it. In contrast, the active inference account can sufficiently explain the tactile
gating effect, i.e., the reduction in the precision of somatosensory input on the moving limb
during movement. Our findings of a reduced precision of somatosensory input on the moving
limb (i.e., higher JNDs in the *gating* and *attenuation&gating* conditions compared to the
baseline and *attenuation* conditions, respectively) are compatible with this active inference
mechanism for the gating of sensory input during movement.
469
Another mechanism that has been proposed to explain tactile gating is a backward masking or
postdictive mechanism [68] that is not necessarily dependent on motor signals. Accordingly, when
moving a limb, the sensation from the muscles, joints, and skin of the moving limb *masks* the
externally generated touches that are applied on this limb. These sensations could affect the
perception of earlier stimuli in a postdictive manner [75], and this could explain the gating effects
observed for external touches applied even before the movement onset [41]. Our gating findings
of a reduced precision of somatosensory input on the moving limb are compatible with this
backward masking mechanism. Furthermore, in the context of a passive movement (i.e., in the
absence of motor commands), the sensations from the muscles, joints, and skin of the passively
moving limb could mask the externally generated touches on the same limb, and this

mechanism could account for the absence of gating differences between active and passive
movements [34,75].
482
Finally, a recent study [67] suggested that action prediction results in somatosensory enhancement
rather than attenuation and argued that the attenuation observed in all earlier experiments is
actually due to nonpredictive gating processes. This claim cannot be supported based on several
observations and arguments. First, the present results showed that the predictive attenuation of
touch and gating are different phenomena; if sensory attenuation was the same as tactile gating,
then *all* stimuli applied on the moving limb would be attenuated. In contrast, we show that
sensory reafference is selectively attenuated compared to sensory exafference, both in passive
and moving limbs. Similar results were shown in [84], where participants attenuated only their
reafferent touches and not exafferent touches presented on the same limb simultaneously.
Second, Thomas and colleagues [67] provided participants with an arbitrary mapping between the
movement of one hand and sensory feedback on the other hand. Using a very similar setup, a
study from 2006 [3] found neither attenuation, which is consistent with the predictive account of
attenuation, nor enhancement effects, which is inconsistent with the findings of Thomas and
colleagues [67]. Third, it is essential to keep in mind that attenuation has also been shown on an
anesthetized limb [8] and for imagined movements [9], further emphasizing the contribution of
motor prediction in relation to peripheral feedback for somatosensory attenuation. Finally, it
has been shown that when reaching with one arm towards the other, the perceived intensity of
an external tactile stimulus applied to the passive target limb is enhanced [44]. Although we
believe that future experimentation is needed to understand and replicate the reported
enhancement effects [67], we speculate that, as in [44], increased spatial attention or other kinematic
differences play a significant role in the somatosensory enhancement reported by Thomas and
colleagues [67].
505
506   We conclude that the human brain uses two different basic processes to suppress reafferent and
507   exafferent information during movement and rest. This separation of attenuation and gating can
508   explain why although we cannot tickle ourselves because we attenuate our self-tickles, we do
509   tense our muscles when being tickled by others to decrease our sensitivity to external tickles.
510

Materials and Methods

512
Participants
After providing written informed consent, twenty-four participants (12 women and 12 men, 22
right-handed, 1 ambidextrous, 1 left-handed) aged 21-40 years participated. Handedness was
assessed using the Edinburgh Handedness Inventory [96]. The sample size was set to twenty-four
(24) before data collection commenced based on our previous studies using the same methods
[4,10] while ensuring a counterbalanced order of conditions. Three participants were excluded
because of technical issues with the kinematic recordings and replaced by three new participants
to reach the target sample size. The Swedish Ethical Review Authority
(https://etikprovningsmyndigheten.se/) approved the study (no. 2016/445- 31/2, amendment
2019-04536). All participants provided their written informed consent.
523
General Procedure
Participants sat comfortably on a chair with their arms placed on a table. Their left hands rested
palm up, with their index fingers placed on a molded support. The right arms rested palm down
on top of a set of sponges. In each trial, a motor (Maxon EC Motor EC 90 flat; Switzerland)
delivered two taps (the *test* tap and the *comparison* tap) on the pulp of their left index finger
through a cylindrical probe (25 mm height) with a flat aluminum surface (20 mm diameter)

attached to the motor's lever. A force sensor (FSG15N1A, Honeywell Inc.; diameter, 5 mm; minimum resolution, 0.01 N; response time, 1 ms; measurement range, 0–15 N) within the probe recorded the forces applied on the left index finger. Following the presentation of the two taps, participants had to verbally indicate which tap felt stronger: the first or the second. A second identical force sensor within an identical cylindrical probe was placed on top of, but not in contact with, the probe of the left index finger (**Fig. 1**).

A wooden surface was placed under the motor and the sensors. This surface was placed on top of two commercially available drawer runners (IKEA, https://www.ikea.com/us/en/p/besta-drawer-runner-soft-closing-40348715/). One side of the runners was attached to the table with Velcro, and the other side was attached to the bottom side of the surface. With this configuration, the surface, with the motor, the sensors and the participants' hands, could be moved forward and backward.

In all conditions, the *comparison* tap was delivered on the left index finger with a random delay of 800-1500 ms from the *test* tap. In the *attenuation* and *attenuation&gating* conditions (**Fig. 1b, d**), the tap of the participants' right index fingers on the force sensor triggered the *test* tap on their left index finger with an intrinsic delay of ≈ 36 ms.

In the *gating* and *attenuation&gating* conditions (**Fig. 1c, d**), participants were asked to extend their elbow upon an auditory 'go' cue. The extension of the elbow moved the platform forward on the table (**Fig. 1f**). A piece of green tape on the table (**Fig. 1i**) indicated the start position of the platform, while a piece of red tape indicated the end position. The participants were asked to move the platform from the start position to the end position (distance = 25 cm). During the movement of the left arm, the participants received the *test* tap on their left index finger. Before the condition started, we emphasized to the subjects that their task was to pay attention to the force that they would receive during the movement rather than covering exactly the distance between the lines. Moreover, the participants were trained to perform the movement in approximately 1000-1500 ms after the 'go' cue and then stop. In the *gating* condition, the *test* tap was applied 800 ms after the 'go' cue so that its delivery was during the movement. Similarly, in the *attenuation&gating* condition, the participants triggered the *test* tap during the movement. The *comparison* tap was applied 800-1500 ms after the test tap to ensure that the participants had stopped moving. Once the participants gave a response, they returned the platform to the start position.

A motion tracking sensor (6DOF Polhemus Fastrak, USA, weight = 9.1 g, dimensions = 2.29 cm x 2.82 cm x 1.52 cm) was placed on top of the platform to record the motion of the platform due to the movement of the participants' left arm. The sensor recorded the x, y and z positions at a sampling rate of approximately 120 Hz.

Each condition included 70 trials. The test tap was set to 2 N, while the intensity of the comparison tap was systematically varied among seven different force levels (1, 1.5, 1.75, 2, 2.25, 2.5 or 3 N). Each tap lasted for 100 ms. In every trial, participants verbally indicated which tap on their left index finger felt stronger: the first (*test*) or the second (*comparison*). Participants were told not to try to balance their responses (50% first and 50% second), and they were further instructed to make their best guess if the intensity of the two taps felt similar.

In addition, participants were administered white noise through a pair of headphones to preclude any sounds created by the motor to serve as a cue for the task. The loudness of the white noise was adjusted so that participants could clearly hear the auditory cues of the trial. In

580   all conditions, the view of the pulp of the left index finger was occluded. Participants were
581   asked to fixate on a cross placed on a wall 2 m opposite them, but they were allowed to look at
582   the force sensor to guide the movement of the right index finger when needed (**Fig. 1b, d**). No
583   feedback was provided to the participants about their responses.
584
585   *Force discrimination analysis*
586   In each condition, the participants' responses were fitted with a generalized linear model using
587   a *logit* link function (Equation 1):

$$p = \frac{e^{\beta_0 + \beta_1 x}}{1 + e^{\beta_0 + \beta_1 x}} \text{ (Equation 1)}$$

589
590   We extracted two parameters of interest: the PSE, which represents the intensity at which the
591   test tap felt as strong as the comparison tap ($p = 0.5$) and which quantifies the perceived
592   intensity, and the JND, which reflects the participants' discrimination capacity. Before fitting
593   the responses, the values of the applied comparison taps were binned to the closest value with
594   respect to their theoretical values (1, 1.5, 1.75, 2, 2.25, 2.5 or 3 N).
595
Kinematic analysis.
Both position and velocity data were smoothed with a moving average filter in MATLAB
2018a. Velocity was calculated as the first derivative of position. To calculate the distance
participants ran in every trial of each condition, we calculated the minimum and the maximum
position of the platform during the entire trial duration. The peak trial velocity was defined as
the peak velocity of the entire trial. The peak tap velocity was defined as the peak velocity
during the period that the *test* tap was applied.
603
Rejection of trials
After data collection, one hundred seventy-three (173) trials out of 6720 (2.57%) were rejected.
First, in thirty-four trials (34), the intensity of the *test* tap (2 N) was not applied accurately (*test*
tap < 1.85 N or *test* tap > 2.15 N), and in sixteen (16) trials, the responses were missing. Second,
we rejected one hundred seventeen (117) trials in total from the *gating* and *attenuation&gating*
conditions because participants either did not move their left arm (or moved it too slowly)
during the *test* tap (mean velocity < 10 cm/s) or they moved it during the *comparison* tap (mean
velocity > 5 cm/s). The thresholds were based on a previous study [42] showing no gating effects
for velocities smaller than 5 cm/s. The analysis was therefore performed with 6547 trials in
total.
614
Statistical analysis
We used R [97] and JASP [98] to analyze our data. The data normality was assessed using the
Shapiro-Wilk test. Depending on the data normality, we then performed planned comparisons
using either a paired t-test or a Wilcoxon signed-rank test. We report 95% confidence intervals
(CI^{95}) for each statistical test. Effect sizes are given by the partial eta-squared (η_P^2) for the
ANOVAs, by Cohen's *d* for t-tests or by the matched rank biserial correlation *rrb* for the
Wilcoxon signed-rank tests. In addition, a Bayesian factor analysis using default Cauchy priors
with a scale of 0.707 was carried out for all statistical tests to provide information about the
level of support for the null hypothesis compared to the alternative hypothesis (BF_{01}) given the
data. Finally, correlations were performed using Pearson's coefficient *r* given that the data were
normally distributed. All tests were two-tailed.
626
627   *Corrections for multiple comparisons*
628   Since our PSE and JND comparisons were planned, we did not apply corrections for multiple
629   comparisons. However, all results remained exactly the same when applying corrections for the

false discovery rate (FDR) [99]. In the correlation analyses, we did apply corrections for multiple comparisons (FDR) since, although we expected correlations between the PSEs and between the JNDs, we had no *a priori* hypotheses for correlations between PSEs and JNDs.

Acknowledgments

We thank M. Chancel, M. Hauser and L. Miller for their feedback on an earlier version of the manuscript. K.K. was supported by the Swedish Research Council (VR Starting Grant 2019-01909 granted to K.K.). H.H.E. and experimental costs were supported by the Swedish Research Council, Torsten Söderbergs Stiftelse, and Göran Gustafssons Stiftelse. We thank Kanaka Raghavan for help with the illustrations.

Data availability

We do not have an ethical permit to make the data publicly available. However, the data that support the findings of this study are available from the corresponding author upon reasonable request.

Conflicts of interest/competing interests

The authors declare no competing financial interests.

Authors' contributions

K.K. and H.H.E. conceived and designed the experiment. K.K. collected the data and conducted the statistical analysis. K.K. and H.H.E. wrote the manuscript.

References

1. Blakemore, S. J., Frith, C. D. & Wolpert, D. M. Spatio-temporal prediction modulates the perception of self-produced stimuli. *J. Cogn. Neurosci.* **11**, 551–559 (1999).
2. Bays, P. M., Wolpert, D. M. & Flanagan, J. R. Perception of the consequences of self-action is temporally tuned and event driven. *Curr. Biol.* **15**, 1125–1128 (2005).
3. Bays, P. M., Flanagan, J. R. & Wolpert, D. M. Attenuation of self-generated tactile sensations is predictive, not postdictive. *PLoS Biol.* **4**, 281–284 (2006).
4. Kilteni, K., Houborg, C. & Ehrsson, H. H. Rapid learning and unlearning of predicted sensory delays in self-generated touch. *Elife* **8**, 1–17 (2019).
5. Shergill, S. S., Bays, P. M., Frith, C. D. & Wolpert, D. M. Two eyes for an eye: the neuroscience of force escalation. *Science* **301**, 187 (2003).
6. Kilteni, K. & Ehrsson, H. H. Sensorimotor predictions and tool use: Hand-held tools attenuate self-touch. *Cognition* **165**, 1–9 (2017).
7. Kilteni, K. & Ehrsson, H. H. Body ownership determines the attenuation of self-generated tactile sensations. *Proc. Natl. Acad. Sci.* **114**, 8426–8431 (2017).
8. Walsh, L. D., Taylor, J. L. & Gandevia, S. C. Overestimation of force during matching of externally generated forces. *J. Physiol.* **589**, 547–557 (2011).
9. Kilteni, K., Andersson, B. J., Houborg, C. & Ehrsson, H. H. Motor imagery involves predicting the sensory consequences of the imagined movement. *Nat. Commun.* **9**, 1617 (2018).
10. Kilteni, K., Engeler, P. & Ehrsson, H. H. Efference Copy Is Necessary for the Attenuation of Self-Generated Touch. *iScience* **23**, 100843 (2020).
11. Blakemore, S.-J., Wolpert, D. M. & Frith, C. D. Central cancellation of self-produced tickle sensation. *Nat. Neurosci.* **1**, 635–640 (1998).
12. Shergill, S. S. *et al.* Modulation of somatosensory processing by action. *Neuroimage* **70**, 356–362 (2013).
13. Kilteni, K. & Ehrsson, H. H. Functional Connectivity between the Cerebellum and

Somatosensory Areas Implements the Attenuation of Self-Generated Touch. *J.*
Neurosci. **40**, 894–906 (2020).

14. Blakemore, S. J., Frith, C. D. & Wolpert, D. M. The cerebellum is involved in
predicting the sensory consequences of action. *Neuroreport* **12**, 1879–1884 (2001).

15. Blakemore, S. J., Wolpert, D. M. & Frith, C. D. The cerebellum contributes to
somatosensory cortical activity during self-produced tactile stimulation. *Neuroimage*
10, 448–459 (1999).

16. Wolpe, N. *et al.* Ageing increases reliance on sensorimotor prediction through
structural and functional differences in frontostriatal circuits. *Nat. Commun.* **7**, 13034
(2016).

17. Blakemore, S.-J., Wolpert, D. & Frith, C. Why can't you tickle yourself? *Neuroreport*
11, R11–R16 (2000).

18. Leavens, D. A. & Bard, K. A. Tickling. *Curr. Biol.* **26**, R91–R93 (2016).

19. Weiskrantz, L., Elliott, J. & Darlington, C. Preliminary observations on tickling
oneself. *Nature* **230**, 598–599 (1971).

20. Crapse, T. B. & Sommer, M. A. Corollary discharge across the animal kingdom. *Nat.*
Rev. Neurosci. **9**, 587–600 (2008).

21. Cullen, K. E. Sensory signals during active versus passive movement. *Curr. Opin.*
Neurobiol. **14**, 698–706 (2004).

22. Straka, H., Simmers, J. & Chagnaud, B. P. A New Perspective on Predictive Motor
Signaling. *Curr. Biol.* **28**, R232–R243 (2018).

23. Schneider, D. M. & Mooney, R. How Movement Modulates Hearing. *Annu. Rev.*
Neurosci. **41**, 553–572 (2018).

24. Brooks, J. X. & Cullen, K. E. Predictive Sensing: The Role of Motor Signals in
Sensory Processing. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* **4**, 842–850
(2019).

25. Poulet, J. F. A. A. & Hedwig, B. The Cellular Basis of a Corollary Discharge. *Science*
(80-.). **311**, 518–522 (2006).

26. Poulet, J. F. A. & Hedwig, B. Corollary Discharge Inhibition of Ascending Auditory
Neurons in the Stridulating Cricket. *J. Neurosci.* **23**, 4717–4725 (2003).

27. Schneider, D. M., Sundararajan, J. & Mooney, R. A cortical filter that learns to
suppress the acoustic consequences of movement. *Nature* **561**, 391–395 (2018).

28. Fukutomi, M. & Carlson, B. A. A History of Corollary Discharge: Contributions of
Mormyrid Weakly Electric Fish. *Front. Integr. Neurosci.* **14**, (2020).

29. Sawtell, N. B. Neural Mechanisms for Predicting the Sensory Consequences of
Behavior: Insights from Electrosensory Systems. *Annu. Rev. Physiol.* **79**, 381–399
(2017).

30. Cullen, K. E. The vestibular system: multimodal integration and encoding of self-
motion for motor control. *Trends Neurosci.* **35**, 185–196 (2012).

31. Brooks, J. X., Carriot, J. & Cullen, K. E. Learning to expect the unexpected: rapid
updating in primate cerebellum during voluntary self-motion. *Nat. Neurosci.* **18**, 1310–
1317 (2015).

32. Roy, J. E. Dissociating Self-Generated from Passively Applied Head Motion: Neural
Mechanisms in the Vestibular Nuclei. *J. Neurosci.* **24**, 2102–2111 (2004).

33. Angel, R. W. & Malenka, R. C. Velocity-dependent suppression of cutaneous
sensitivity during movement. *Exp. Neurol.* (1982). doi:10.1016/0014-4886(82)90244-8

34. Chapman, C. E., Bushnell, M. C., Miron, D., Duncan, G. H. & Lund, J. P. Sensory
perception during movement in man. *Exp. Brain Res.* **68**, 516–524 (1987).

35. Post, L. J., Zompa, I. C. & Chapman, C. E. Perception of vibrotactile stimuli during
motor activity in human subjects. *Exp. Brain Res.* **100**, 107–120 (1994).

36. Gertz, H., Voudouris, D. & Fiehler, K. Reach-relevant somatosensory signals modulate
tactile suppression. *J. Neurophysiol.* **117**, 2262–2268 (2017).
37. Fraser, L. E. & Fiehler, K. Predicted reach consequences drive time course of tactile
suppression. *Behav. Brain Res.* **350**, 54–64 (2018).
38. Voudouris, D., Broda, M. D. & Fiehler, K. Anticipatory grasping control modulates
somatosensory perception. *J. Vis.* **19**, 1–10 (2019).
39. Williams, S. R., Shenasa, J. & Chapman, C. E. Time course and magnitude of
movement-related gating of tactile detection in humans. I. Importance of stimulus
location. *J. Neurophysiol.* **79**, 947–963 (1998).
40. Williams, S. R. & Chapman, C. E. Time course and magnitude of movement-related
gating of tactile detection in humans. II. Effects of stimulus intensity. *J. Neurophysiol.*
(2000). doi:10.1152/jn.2000.84.2.863
41. Chapman, C. E. & Beauchamp, E. Differential controls over tactile detection in
humans by motor commands and peripheral reafference. *J. Neurophysiol.* **96**, 1664–
1675 (2006).
42. Cybulska-Klosowicz, A., Meftah, E. M., Raby, M., Lemieux, M. L. & Chapman, C. E.
A critical speed for gating of tactile detection during voluntary movement. *Exp. Brain*
Res. **210**, 291–301 (2011).
43. Colino, F. L. & Binsted, G. Time Course of Tactile Gating in a Reach-to-Grasp and
Lift Task. *J. Mot. Behav.* **48**, 390–400 (2016).
44. Voudouris, D. & Fiehler, K. Enhancement and suppression of tactile signals during
reaching. *J. Exp. Psychol. Hum. Percept. Perform.* (2017). doi:10.1037/xhp0000373
45. Colino, F. L., Buckingham, G., Cheng, D. T., van Donkelaar, P. & Binsted, G. Tactile
gating in a reaching and grasping task. *Physiol. Rep.* **2**, 1–11 (2014).
46. Papakostopoulos, D., Cooper, R. & Crow, H. J. Inhibition of cortical evoked potentials
and sensation by self-initiated movement in man. *Nature* **258**, 321–324 (1975).
47. Starr, A. & Cohen, L. G. 'Gating' of somatosensory evoked potentials begins before
the onset of voluntary movement in man. *Brain Res.* **348**, 183–186 (1985).
48. Rushton, D. N., Roghwell, J. C. & Craggs, M. D. Gating of somatosensory evoked
potentials during different kinds of movement in man. *Brain* **104**, 465–491 (1981).
49. Giblin, D. R. SOMATOSENSORY EVOKED POTENTIALS IN HEALTHY
SUBJECTS AND IN PATIENTS WITH LESIONS OF THE NERVOUS SYSTEM.
Ann. N. Y. Acad. Sci. **112**, 93–142 (1964).
50. Chapman, C. E. Active versus passive touch: Factors influencing the transmission of
somatosensory signals to primary sornatosensory cortex. *Can. J. Physiol. Pharmacol.*
72, 558–570 (1994).
51. Lei, Y., Ozdemir, R. A. & Perez, M. A. Gating of sensory input at subcortical and
cortical levels during grasping in humans. *J. Neurosci.* (2018).
doi:10.1523/JNEUROSCI.0545-18.2018
52. Azim, E. & Seki, K. Gain control in the sensorimotor system. *Curr. Opin. Physiol.* **8**,
177–187 (2019).
53. Ghez, C. & Lenzi, G. L. Modulation of sensory transmission in cat lemniscal system
during voluntary movement. *Pflügers Arch. Eur. J. Physiol.* **323**, 273–278 (1971).
54. Chapin, J. K. & Woodward, D. J. Modulation of sensory responsiveness of single
somatosensory cortical cells during movement and arousal behaviors. *Exp. Neurol.* **72**,
164–178 (1981).
55. Jiang, W., Chapman, C. E. & Lamarre, Y. Modulation of the cutaneous responsiveness
of neurones in the primary somatosensory cortex during conditioned arm movements in
the monkey. *Exp. Brain Res.* (1991). doi:10.1007/BF00231455
56. Jiang, W., Lamarre, Y. & Chapman, C. E. Modulation of cutaneous cortical evoked

potentials during isometric and isotonic contractions in the monkey. *Brain Res.* (1990).
doi:10.1016/0006-8993(90)90010-9

57.　Seki, K. & Fetz, E. E. Gating of sensory input at spinal and cortical levels during
preparation and execution of voluntary movement. *J. Neurosci.* **32**, 890–902 (2012).

58.　Seki, K., Perlmutter, S. I. & Fetz, E. E. Sensory input to primate spinal cord is
presynaptically inhibited during voluntary movement. *Nat. Neurosci.* **6**, 1309–1316
(2003).

59.　McNamee, D. & Wolpert, D. M. Internal Models in Biological Control. *Annu. Rev.*
Control. Robot. Auton. Syst. **2**, 339–364 (2019).

60.　Collins, D. F., Cameron, T., Gillard, D. M. & Prochazka, A. Muscular sense is
attenuated when humans move. *J. Physiol.* **508**, 635–643 (1998).

61.　Juravle, G., Binsted, G. & Spence, C. Tactile suppression in goal-directed movement.
Psychon. Bull. Rev. (2017). doi:10.3758/s13423-016-1203-6

62.　Brown, H., Adams, R. a., Parees, I., Edwards, M. & Friston, K. Active inference,
sensory attenuation and illusions. *Cogn. Process.* **14**, 411–427 (2013).

63.　Limanowski, J. *et al.* Action-Dependent Processing of Touch in the Human Parietal
Operculum and Posterior Insula. *Cereb. Cortex* **30**, 607–617 (2020).

64.　Saradjian, A. H. Sensory modulation of movement, posture and locomotion.
Neurophysiol. Clin. **45**, 255–267 (2015).

65.　Palmer, C. E., Davare, M. & Kilner, J. M. Physiological and Perceptual Sensory
Attenuation Have Different Underlying Neurophysiological Correlates. *J. Neurosci.* **36**,
10803–10812 (2016).

66.　Boehme, R., Hauser, S., Gerling, G. J., Heilig, M. & Olausson, H. Distinction of self-
produced touch and social touch at cortical and spinal cord levels. *Proc. Natl. Acad.*
Sci. **116**, 2290–2299 (2019).

67.　Thomas, E. R., Yon, D., Lange, F. P. de & Press, C. Action enhances predicted touch.
bioRxiv (2020). doi:10.1101/2020.03.26.007559

68.　Voss, M., Ingram, J. N., Wolpert, D. M. & Haggard, P. Mere expectation to move
causes attenuation of sensory signals. *PLoS One* **3**, 2–6 (2008).

69.　Blakemore, S.-J., Smith, J., Steel, R., Johnstone, E. C. & Frith, C. D. The perception of
self-produced sensory stimuli in patients with auditory hallucinations and passivity
experiences: evidence for a breakdown in self-monitoring. *Psychol. Med.* **30**, 1131–
1139 (2000).

70.　Shergill, S. S., Samson, G., Bays, P. M., Frith, C. D. & Wolpert, D. M. Evidence for
sensory prediction deficits in schizophrenia. *Am. J. Psychiatry* **162**, 2384–2386 (2005).

71.　Shergill, S. S. *et al.* Functional Magnetic Resonance Imaging of Impaired Sensory
Prediction in Schizophrenia. *JAMA Psychiatry* **71**, 28 (2014).

72.　Parees, I. *et al.* Loss of sensory attenuation in patients with functional (psychogenic)
movement disorders. *Brain* **137**, 2916–2921 (2014).

73.　Wolpe, N. *et al.* Sensory attenuation in Parkinson's disease is related to disease
severity and dopamine dose. *Sci. Rep.* **8**, 15643 (2018).

74.　Macerollo, A. *et al.* Dopaminergic Modulation of Sensory Attenuation in Parkinson's
Disease: Is There an Underlying Modulation of Beta Power? *Front. Neurol.* **10**, 1–14
(2019).

75.　Williams, S. R. & Chapman, C. E. Time course and magnitude of movement-related
gating of tactile detection in humans. III. Effect of motor tasks. *J. Neurophysiol.* **88**,
1968–1979 (2002).

76.　Rauch, R., Angel, R. W. & Boylls, C. C. Velocity-dependent suppression of
somatosensory evoked potentials during movement. *Electroencephalogr. Clin.*
Neurophysiol. Evoked Potentials **62**, 421–425 (1985).

77.   Cohen, L. G. & Starr, A. Localization, timing and specificity of gating of
831   somatosensory evoked potentials during active movement in man. *Brain* (1987).
832   doi:10.1093/brain/110.2.451

833   78.   Pertovaara, A., Kemppainen, P. & Leppänen, H. Lowered cutaneous sensitivity to
834   nonpainful electrical stimulation during isometric exercise in humans. *Exp. Brain Res.*
835   **89**, 447–452 (1992).

836   79.   Shadmehr, R., Krakauer, J. W., Neuroanatomy, A. C. & Motor, F. O. R. A
837   computational neuroanatomy for motor control. *Exp. Brain Res.* **185**, 359–381 (2008).

838   80.   Kawato, M. Internal models for motor control and trajectory planning. *Curr. Opin.*
839   *Neurobiol.* **9**, 718–727 (1999).

840   81.   Todorov, E. Optimality principles in sensorimotor control. *Nat. Neurosci.* **7**, 907–15
841   (2004).

842   82.   Wolpert, D. M. & Ghahramani, Z. Computational principles of movement
843   neuroscience. *Nat. Neurosci.* **3**, 1212–1217 (2000).

844   83.   Bays, P. M. & Wolpert, D. M. Computational principles of sensorimotor control that
845   minimize uncertainty and variability. *J. Physiol.* **578**, 387–396 (2007).

846   84.   Bays, P. M. & Wolpert, D. M. Predictive attenuation in the perception of touch. in
847   *Sensorimotor Foundations of Higher Cognition* (eds. Haggard, E. P., Rosetti, Y. &
848   Kawato, M.) **22**, 339–358 (Oxford University Press, 2008).

849   85.   Wolpert, D. M. & Flanagan, J. R. Motor prediction. *Curr. Biol.* **11**, R729–R732 (2001).

850   86.   Diedrichsen, J., Verstynen, T., Hon, A., Lehman, S. L. & Ivry, R. B. Anticipatory
851   adjustments in the unloading task: Is an efference copy necessary for learning? *Exp.*
852   *Brain Res.* **148**, 272–276 (2003).

853   87.   Klaffehn, A. L., Baess, P., Kunde, W. & Pfister, R. Sensory attenuation prevails when
854   controlling for temporal predictability of self- and externally generated tones.
855   *Neuropsychologia* **132**, 107145 (2019).

856   88.   Bäß, P., Jacobsen, T. & Schröger, E. Suppression of the auditory N1 event-related
857   potential component with unpredictable self-initiated tones: Evidence for internal
858   forward models with dynamic stimulation. *Int. J. Psychophysiol.* (2008).
859   doi:10.1016/j.ijpsycho.2008.06.005

860   89.   Miall, R. C. & Wolpert, D. M. Forward models for physiological motor control. *Neural*
861   *Networks* **9**, 1265–1279 (1996).

862   90.   Wolpert, D. M., Miall, R. C. & Kawato, M. Internal models in the cerebellum. *Trends*
863   *in Cognitive Sciences* **2**, 338–347 (1998).

864   91.   Shadmehr, R., Smith, M. a & Krakauer, J. W. Error correction, sensory prediction, and
865   adaptation in motor control. *Annu. Rev. Neurosci.* **33**, 89–108 (2010).

866   92.   Therrien, A. S. & Bastian, A. J. The cerebellum as a movement sensor. *Neurosci. Lett.*
867   **688**, 0–1 (2019).

868   93.   Adams, R. a., Shipp, S. & Friston, K. J. Predictions not commands: Active inference in
869   the motor system. *Brain Struct. Funct.* **218**, 611–643 (2013).

870   94.   Friston, K. *The free-energy principle: A unified brain theory? Nature Reviews*
871   *Neuroscience* **11**, 127–38 (2010).

872   95.   Pickering, M. J. & Clark, A. Getting ahead: Forward models and their place in
873   cognitive architecture. *Trends Cogn. Sci.* **18**, 451–456 (2014).

874   96.   Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh inventory.
875   *Neuropsychologia* **9**, 97–113 (1971).

876   97.   Core Team, R. R: A language and environment for statistical computing. *R Foundation*
877   *for Statistical Computing* (2018). doi:10.1007/978-3-540-74686-7

878   98.   JASP & JASP Team. JASP. *[Computer software]* (2019).

879   99.   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

880   Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* (1995).
881   doi:10.1111/j.2517-6161.1995.tb02031.x
882