The BEACH Domain-Containing Protein SPIRRIG Modulates Actin-Dependent

Root Hair Development in Coordination with the WAVE/SCAR and ARP2/3

Complexes

Sabrina Chin[a], Taegun Kwon[a], Bibi Rafeiza Khan[a, 2], J. Alan Sparks[a], Eileen L.

Mallery[b], Daniel B. Szymanski[b,c] and Elison B. Blancaflor[a, 1]

[a] Noble Research Institute LLC, 2510 Sam Noble Parkway, Ardmore, Oklahoma 73401,

[b] Department of Botany and Plant Pathology, [c] Dept. of Biological Sciences, Purdue

University, West Lafayette, Indiana 47907

ORCID IDs: 0000-0003-3897-3812 (S.C.); 0000-0002-0205-8535 (T.K.); 0000-0002-

7661-0142 (B.R.K.); 0000-0002-4492-9810 (J.A.S.); 0000-0002-0363-5920 (E.L.M.),

0000-0001-8255-424X (D.B.S.); 0000-0001-6115-9670 (E.B.B.)

[1] Address correspondence to eblancaflor@noble.org

[2] Current address: Department of Biology, University of Guyana, Turkeyen Campus, 41

Georgetown, Guyana

Short Title: Actin-mediated root hair development

The author responsible for distribution of materials integral to the findings presented in

this article in accordance with the policy described in the Instructions for Authors are:

Daniel B. Szymanski (szymandb@purdue.edu) and Elison B. Blancaflor

(eblancaflor@noble.org).

Abstract

Root hairs are single cell protrusions that enable roots to optimize nutrient and water acquisition. They attain their tubular shapes by confining growth to the cell apex, a process called tip growth. The actin cytoskeleton and endomembrane system are essential for tip growth; however, little is known about how these cellular components coordinate their activities during this process. Here, we show that SPIRRIG (SPI), a BEACH domain-containing protein involved in membrane trafficking, and BRK1 and SCAR2, subunits of the WAVE/SCAR (W/SCR) and actin related protein (ARP)2/3 activation complexes, display polarized localizations to root hairs at distinct developmental stages. SPI accumulates at the root hair apex via post-Golgi vesicles and positively regulates tip growth by maintaining tip-focused vesicle secretion and filamentous-actin integrity. BRK1 and SCAR2 on the other hand, mark the root hair initiation domain to specify the position of root hair emergence. Live cell microscopy revealed that BRK1 depletion coincided with SPI accumulation as root hairs transitioned from initiation to tip growth. Furthermore, double mutant studies showed that *SPI* genetically interacts with *BRK1* and *ARP2/3*. Taken together, our work uncovers a role for SPI in facilitating actin-dependent root hair development through pathways that intersect with the W/SCR and ARP2/3 complexes.

INTRODUCTION

Root hairs are single cell tubular projections that emerge from root epidermal

cells. They increase the effective surface area of the root system by extending laterally

into soil pores, thus enabling the increased access to nutrients and water (Carminati et

al., 2017, Ruiz et al., 2020). Root hairs have been studied extensively by plant biologists

for decades because they serve as excellent models to unravel mechanisms by which

cell size and shape in plants is regulated (Grierson et al., 2014). To attain their

cylindrical shapes, root hairs undergo tip growth, a process in which expansion of the

cell is confined to its apical domain. Tip growth involves a balance between the directed

delivery of post-Golgi vesicles carrying protein complexes and cell wall building blocks

to the cell apex, and localized cell wall loosening and recycling of excess membranes.

Besides root hairs, tip growth is exhibited by other cell types such as pollen tubes,

fungal hyphae, and rhizoids of mosses, liverworts and algae (Bascom et al., 2018a).

Root epidermal cells called trichoblasts are the cell types that form root hairs.

Work in *Arabidopsis thaliana* has shown that trichoblasts are specified to become root

hair-forming cells early during root development through the patterned assembly of

protein complexes of transcriptional activators and repressors in different cell files

(Schiefelbein et al., 2014, Shibata et al., 2018). Upon establishing their identity,

trichoblasts undergo two developmental stages that lead to root hair outgrowth. The first

stage is the establishment of a root hair initiation domain (RHID) that eventually leads to

a conspicuous root hair bulge at the basal (root-tip oriented) end of the trichoblast

(Grierson et al., 2014, Nakamura and Grebe, 2018). Several proteins accumulate at the

RHID, most prominently, the Rho of Plants (ROP)/RAC small GTPases and their

Guanine Nucleotide Exchange Factor (GEF) activators (Denninger et al., 2019). The

second stage is tip growth. In addition to small GTPases, the cytoskeleton, the

endomembrane trafficking machinery, cytoplasmic calcium, phosphoinositide lipids,

hormones (e.g. auxin and ethylene) and reactive oxygen species are other major

players in signaling pathways that modulate root hair development (Bascom et al.,

2018a, Nakamura and Grebe, 2018).

The filamentous-actin (F-actin) and microtubule cytoskeletons orchestrate root

hair development. The actin cytoskeleton in particular has been studied widely during tip

growth as it functions as tracks for the traffic of cellular cargo to the cell apex (Bascom

et al., 2018a, Stephan, 2017, Szymanski and Staiger, 2018). Many insights into the role

of the actin cytoskeleton in tip growth have been demonstrated through work with

Arabidopsis mutants and pharmacological approaches involving the use of chemicals

that disrupt F-actin. For example, the actin-disrupting compound, latrunculin B (LatB),

inhibits tip growth, while also inducing the formation of root hairs and pollen tubes with

irregular shapes (Bibikova et al., 1999, Gibbon et al., 1999). In this regard, *Arabidopsis*

mutants with defects in the root hair-expressed vegetative *ACTIN2* (*ACT2*) gene are

characterized by root hairs that mirror those treated with LatB (Ringli et al., 2002, Yoo et

al., 2012). Furthermore, *ACT2* and *ACT7* mutants display an apical (shoot-ward) shift in

the position of the RHID when compared to wild type (Kiefer et al., 2015).

To fulfill its cellular functions, the actin cytoskeleton is organized into higher order

networks that correspond to the growth strategy of the cell. This is evident in tip-growing

cells, whereby the base and shank of the cell consists mostly of thick, longitudinal F-

actin bundles, while the apex contains actin fringes, rings, patches or a fine meshwork

depending on the plant species or cell type (Stephan, 2017). To sustain tip growth, the

integrity and organization of these tip-focused F-actin arrays must be maintained, a task

that is facilitated by a diverse collection of actin-binding proteins and actin nucleators (Li

et al., 2015, Paez-Garcia, 2018, Szymanski and Staiger, 2018).

The actin-related protein (ARP2/3) complex and its activator, suppressor of

cAMP receptor (SCAR)-WASP family verprolin homologous (WAVE) complex (W/SRC-

ARP2/3), is the best characterized actin nucleator in plants (Deeks and Hussey, 2005,

Szymanski, 2005, Yanagisawa et al., 2013). The ARP2/3 complex consists of seven

subunits that include the proteins ARP2, ARP3, ARPC1, ARPC2, ARPC3, ARPC4, and

ARPC5, which were first identified in *Acanthamoeba* (Machesky et al., 1994). It was

shown that the membrane-associated plant ARP2/3 complex has the same assembly

mechanisms as the non-plant complexes (Kotchoni et al., 2009). Upon conversion from

an inactive open conformation to an inactive closed conformation, the ARP2/3 complex

promotes F-actin nucleation from the sides of existing filaments by forming a surface

that mimics stable actin dimers (Blanchoin et al., 2000, Robinson et al., 2001, Rodal et

al., 2005). Activation of ARP2/3 for efficient nucleation of F-actin requires the W/SCAR

nucleation promoting factor (NPF) complex. In addition to WAVE/SCAR, the NPF

consists of the proteins SRA1, NAP1, ABIL and HSPC300/BRICK1 (BRK1) (Deeks et

al., 2004, Djakovic, 2006, Frank and Smith, 2002, Jörgens et al., 2010, Le et al., 2006,

Saedler et al., 2004). Studies of *Arabidopsis* trichome development have been most

instrumental in revealing insights into the function of W/SCR-ARP2/3 in plants. Plant

counterparts to the mammalian W/SCR-ARP2/3 subunits were first uncovered through

the cloning of disrupted genes in a set of *DISTORTED* (*DIS*) *Arabidopsis* trichome

mutants (Hülskamp et al., 1994, Mathur et al., 2003, Le et al., 2003, El-Assal Sel et al.,

2004, Basu et al., 2004, Deeks et al., 2004, Zhang et al., 2005). W/SCR is the only

known NPF for ARP2/3, and *BRK1* and *SCAR* mutants display null *arp2/3* trichome

phenotypes (Le et al., 2006, Zhang et al., 2008). NPF activity of W/SCR to initiate an

ARP2/3 actin nucleating response is also known to be positively regulated by

ROP/RAC-GTP effector complexes (Basu et al., 2004, Uhrig et al., 2007). In

Arabidopsis trichomes, active ROP/RAC-GTPase is coupled to W/SCR by the DOCK

(dedicator of cytokinesis)-family GEF, SPIKE1 (SPK1) (Basu et al., 2008) (Yanagisawa

et al., 2018).

In addition to trichome development, the W/SRC-ARP2/3 actin filament-

nucleating module has been implicated in specifying leaf pavement cell shape, light and

auxin-dependent root growth, stomatal gating, gravitropism, responses to salinity stress,

and plant immunity (Li et al., 2003, Basu et al., 2005, Zhang et al., 2005, Dyachok et al.,

2011, Zhao et al., 2013, Li et al., 2014, Zou et al., 2016, Isner et al., 2017, Badet et al.,

2019, Pratap Sahi et al., 2018). The W/SRC-ARP2/3 complex has also been shown to

play an essential role in some tip growing cells such as the protonemal cells of the moss

Physcomitrella patens. This was shown when RNA interference (RNAi) of transcripts

encoding selected subunits of the moss W/SRC-ARP2/3 complex resulted in substantial

protonemal tip growth defects (Harries et al., 2005, Perroud and Quatrano, 2006, Finka

et al., 2008). The importance of the W/SRC-ARP2/3 complex was further demonstrated

by Perroud and Quatrano (2006, 2008) who showed that two of its components, BRK1

and ARPC4, accumulated in tips of moss protonemal cells. By contrast, *Arabidopsis*

plants with mutations in components of the W/SRC-ARP2/3 complex displayed minimal

or no defects in root hair and pollen tube growth (Li et al., 2003, Mathur et al., 2003),

and none of the subunits have been conclusively shown to exhibit clear polar

localization in these cell types. Therefore, the extent by which the W/SRC-ARP2/3

complex functions in tip growing cells of higher plants remains to be determined.

SPIRRIG (SPI) is one of eight genes that have been considered to be a member

of the DIS group, but compared to other DIS mutants, trichome phenotypes of spi are

less severe and the mutant does not display early stage cell swelling that is diagnostic

of the DIS group (Schwab et al., 2003). SPI was shown to encode a 3571 long amino

acid protein with N-terminally located armadillo (ARM) and concanavalin A (ConA)-like

lectin domains and C-terminally-located pleckstrin homology (PH), beige and Chediak

Higashi (BEACH) and WD40 repeat domains (Saedler et al., 2009). BEACH domain-

containing proteins are highly conserved in eukaryotes and are known to function in

membrane dynamics, vesicle transport, apoptosis and receptor signaling. This family of

proteins are of clinical importance as they have been implicated in a variety of human

disorders such as cancer, autoimmunity syndrome and autism (Cullinane et al., 2013).

In addition to mild trichome defects, Arabidopsis SPI mutants have short root

hairs characterized by fragmented vacuoles suggesting that SPI, like other eukaryotic

BEACH domain-containing proteins, functions in membrane trafficking (Saedler et al.,

2009). Additionally, an observation by Steffens et al. (2017) that showed SPI physically

interacting with proteins involved in endosomal sorting reinforces its role in membrane

remodeling. SPI was also demonstrated to associate with mRNA processing bodies (P-

bodies) suggesting a novel role for SPI in post-transcriptional regulation (Steffens et al.,

2015). Because SPI is one of the DIS genes, it was suggested that SPI might be

involved in actin-mediated cell developmental processes (Saedler et al., 2009), and

perhaps function in coordination with W/SCR and ARP2/3 complexes. However, SPI is

not a known W/SCR or ARP2/3 subunit; therefore, its relationship to ARP2/3 function is

unclear. Moreover, the uncertainty with regard to SPI function is confounded by the fact

that its subcellular localization in root hairs, which exhibit the most profound phenotype

when SPI is mutated, remains unknown. In this paper, we have addressed these

questions by showing that a functional SPI-fluorescent protein fusion accumulated at

the tips of rapidly growing root hairs. A SPI-tagged fluorescent protein fusion was not

detected in the RHID and in tips of non-growing mature root hairs. We further show that

the W/SCR complex subunits, BRK1 and SCAR2, localized to the RHID and exhibited

temporal dynamics opposite to that of SPI (i.e. functional fluorescent protein-tagged

BRK1 and SCAR2 declined as rapid root hair tip growth commenced). Double mutant

studies revealed that *SPI* and *W/SCR-ARP2/3* genetically interact suggesting that one

function of SPI is to modulate actin-dependent root hair development, in part through

pathways that overlap with W/SCR and ARP2/3 complexes.

RESULTS

Isolation of a New *SPIRRIG* Mutant Allele

We previously described a forward genetic screen that led to the isolation of three non-allelic recessive *Arabidopsis* mutants that were hypersensitive to the growth inhibitory effects of LatB. The *hypersensitive to LatB1* (*hlb1*) and *hlb3* mutants have been described previously (Sparks et al., 2016, Sun et al., 2019). *hlb1* was disrupted in a gene encoding a *trans*-Golgi Network-localized tetratricopeptide repeat protein involved in actin-mediated membrane recycling (Sparks et al., 2016), whereas the genetic lesion in *hlb3* was found to encode the class II actin nucleator formin (Sun et al., 2019). Here, we report on *hlb2*, the third of these recessive mutants. Like *hlb1* and *hlb3*, *hlb2* primary root growth was more severely inhibited by LatB when compared to wild type. In the absence of LatB or at low (i.e. 25 nM) LatB concentrations, the primary root length of *hlb2* was similar to wild type. Differences in root length between wild type and *hlb2* became apparent when seedlings were grown on a concentration of 50 nM LatB and higher (Supplemental Figure 1A and B).

Through map-based cloning, we found that the mutation in *hlb2* was confined to a region between the *AT1G02740* and *AT1G03410* loci. Nucleotide sequencing revealed that *hlb2* had a 10 base pair deletion (Chr1 position: 720,152 - 720,161) in exon 14 of the *AT1G03060* gene. This 10-base pair deletion led to an open-reading frame shift at the Asp^{1526} codon resulting in a truncated protein (Supplemental Figure 2A). *AT1G03060* encodes the BEACH domain-containing protein, SPI (Saedler et al., 2009) (Supplemental Figure 2A). Because the first *spi* mutant alleles were reported to have short root hairs (Saedler et al., 2009), we examined root hairs of *hlb2*. We found that *hlb2* had root hairs that were about 80% shorter than wild type with some root hairs forming only small bulges (Supplemental Figure 2B and C). Moreover, *hlb2* had mild trichome defects that were reminiscent of the phenotypes of previously isolated *spi* mutant alleles (Supplemental Figure 2D; Saedler et al., 2009). To further verify if *HLB2* is *SPI*, we obtained a mutant from the publicly available SALK collection (SALK_065311), which had a T-DNA insertion in the 10[th] exon of the *SPI* gene

(Supplemental Figure 2A; Alonso et al. (2003)). The SALK_065311 line we obtained is

the same as *spi-3* mutant allele reported previously in Steffens et al. (2015). In addition

to having similar root hair and trichome defects as *hlb2*, primary roots of *spi-3* were

hypersensitive to LatB (Supplemental Figure 2E), and a cross between *hlb2* and

SALK_065311 failed to complement each other in the F1 hybrid. Taken together, these

results indicate that *hlb2* is a new *spi* mutant allele. Based on earlier nomenclature

(Steffens et al., 2015), we renamed *hlb2* to *spi-5* (Supplemental Figure 2A). The *spi-5*

mutant allele was used for all subsequent experiments.

SPIRRIG Localizes to the Tips of Rapidly Elongating Root Hairs

SPI fluorescent protein fusions have been previously demonstrated to associate with P-

bodies and endosomes (Steffens et al., 2015, Steffens et al., 2017). However, these

constructs have not been shown to complement the defective root hair and trichome

phenotypes of *spi*. Because of the large size of SPI, we were unable to generate native

promoter-driven fluorescent protein fusions to the full length *SPI* complementary DNA or

genomic DNA. To overcome this problem, SPI was tagged with the 3x-Yellow

Fluorescent Protein (<u>YFP</u>) for <u>e</u>nergy <u>t</u>ransfer (YPet; Nguyen and Daugherty (2005))

using a method called recombineering. This method involves tagging the gene of

interest in the context of transformation-competent bacterial artificial chromosomes to

ensure that all regulatory sequences are included in the fluorescent protein fusion

(Brumos et al., 2020, Zhou et al., 2011).

Once *spi* was transformed with a recombineered *SPI-YPet* construct, primary

root hypersensitivity to LatB, short root hairs and defective trichome phenotypes were

rescued, indicating that the construct was functional (Figure 1A; Supplemental Figure

3). Transgenic complementation of *spi* with *SPI-YPet* provided additional evidence that

HLB2 is *SPI*. Confocal microscopy revealed that SPI-YPet signal was strongest in the

tips of rapidly elongating root hairs (Figure 1B and C; Supplemental Movie 1). SPI-YPet

fluorescence was not detected or weak at the RHID and during early root hair bulge

formation, but intensified as root hairs transitioned to tip growth (Figure 1C;

Supplemental Movie 2). SPI-YPet signal dissipated as root hairs matured and tip growth

rate declined (Figure 1B and C; Supplemental Movie 3). A linear regression analysis

showed that the intensity of SPI-YPet fluorescence in root hair tips is significantly

positively correlated to rapid root hair growth (Figure 1D). These results indicate that the

SPI protein has functions related to root hair elongation and consistent with the short

root hair phenotypes of *spi*.

SPIRRIG is Transported to the Root Hair Tip via Post-Golgi Vesicles

The prominent SPI-YPet signal at the tips of elongating root hairs is reminiscent of the

localization patterns of post-Golgi markers such as RAB small GTPases, which are

known to function in tip-directed secretion (Preuss et al., 2004). We therefore

hypothesized that SPI is trafficked to the tips of root hairs via post-Golgi vesicles. To

test this hypothesis, seedlings expressing *SPI-YPet* were treated with brefeldin A (BFA).

BFA is a fungal toxin that is routinely used to investigate endomembrane dynamics

because it prevents vesicle formation for exocytosis by inhibiting ADP ribosylation factor

guanine nucleotide exchange factors (ARF-GEFs), while still enabling endocytosis and

some retrograde pathways to continue (Baluška, 2002, Doyle et al., 2015).

Consequently, BFA treatment causes the formation of endosomal agglomerations called

BFA bodies. Treatment of seedlings expressing *SPI-YPet* with 50 μM BFA induced the

formation of SPI-YPet agglomerates in root hairs (Figures 2A and B). The formation of

SPI-YPet fluorescent agglomerates was also observed in the root elongation zone

(Figure 2C). In the absence of BFA, SPI-YPet signal in the elongation zone was barely

detectable. However, upon exposure to BFA, bright fluorescent agglomerates of various

sizes became visible (Figure 2C). These results indicate that SPI-YPet is associated

with endomembranes and is trafficked to the root hair tips via BFA-sensitive post-Golgi

vesicles.

Inhibited root hair growth of *spi* and the formation of SPI-YPet agglomerates after

BFA treatment, suggest that *spi* may be defective in tip-directed protein secretion and

partly explain the short root hair phenotypes of *spi*. To address this question, we

expressed the secreted (SEC)-red fluorescent protein (RFP) in *spi* (Faso et al., 2009).

Similar to previous reports, rapidly elongating wild-type root hairs had a tip-focused

gradient of SEC-RFP (Sparks et al., 2016) (Figure 2D). By contrast, slow-growing root

hairs of *spi* lacked these tip-focused SEC-RFP gradients (Figure 2E), thereby indicating

defects in tip-directed bulk flow exocytosis.

SPIRRIG Maintains Root Hair Tip-Focused F-actin

In a study of other *spi* mutant alleles, the similarities in phenotypes between *spi* and

w/scr-arp2/3 suggests that SPI could function in actin-dependent cellular processes

(Saedler et al., 2009). However, because trichome defects of *spi* were mild compared to

other *w/scr-arp2/3* mutants, no obvious actin phenotypes were observed in *spi*

trichomes (Schwab et al., 2003). To clarify the relationship between SPI and actin, we

focused on investigating actin organization in root hairs since they displayed the most

obvious growth defects in *SPI*-altered plants.

To study actin organization, we expressed the live F-actin reporter, *UBQ10:*

mGFP-Lifeact, in *spi* (Vidali et al., 2010). This particular F-actin reporter was selected

because it prominently labels the tip-focused F-actin meshwork typically observed in

root hairs that are rapidly growing (Sparks et al., 2016). In wild type, fine F-actin

networks were observed in the root hair bulge that had a weaker signal compared to the

thicker actin bundles in other regions of the trichoblast (Figure 3A). As the root hair

bulge expanded and the root hair transitioned to rapid tip growth, the tip-focused F-actin

meshwork, which consisted of short filaments and dynamic puncta, became more

conspicuous (Figures 3B-E; Supplemental Movie 4; Sparks et al. (2016)). When wild-

type root hairs stopped elongating, the tip-focused F-actin meshwork was replaced with

thick F-actin cables (Figure 3F).

F-actin organization in the tips of *spi* root hairs was different from that of wild

type. As noted, some root hairs of *spi* were only able to form small bulges due to

premature termination of tip-growth (Supplemental Figure 2). In *spi* root hairs, the

distinct F-actin meshwork observed in wild-type root hairs was unable to form. Instead,

F-actin in these *spi* root hair bulges contained thick F-actin cables that resembled those

of wild-type root hairs that had terminated growth (Figure 3G and H). However, the thick

F-actin cables in non-growing *spi*root hair bulges were unstable as they dissipated

(Figure 3I) and reformed again at a later time (Figure 3J). In the small population of *spi*

root hairs that were able to undergo tip growth, a few exhibited the tip-focused F-actin

meshwork that resembled those observed in elongating wild-type root hairs (Figure 3K).

However, most of these slow growing *spi* root hairs lacked the tip-focused F-actin

meshwork (Figure 3L and M; Supplemental Movie 4) or had thick F-actin bundles

protruding to the tip, a feature that was reminiscent of non-growing, mature wild-type

root hairs (Figure 3N). The disruption of the tip-focused F-actin meshwork in *spi* was

quantified by measuring F-actin fluorescence from computer reconstructed transverse

sections of the root hair tip and by taking the ratio of the average fluorescence over

background signal. A higher ratio indicates a higher signal of tip-focused F-actin (Figure

3O). Our analysis showed that the fluorescence ratio in *spi* root hairs was significantly

reduced compared to wild-type root hairs, supporting visual observations that the tip-

focused F-actin meshwork in *spi* root hairs is disrupted (Figure 3P).

We next generated plants expressing both *SPI-YPet* and *mRuby-Lifeact* so we

could simultaneously observe SPI and F-actin in growing root hairs. In elongating root

hairs of dual labeled seedlings, SPI-YPet and the mRuby-labeled F-actin meshwork

overlapped at the tip (Figure 3Q; Supplemental Movie 5). This provides support that root

hair tip-localized SPI is associated with the tip-focused F-actin meshwork, and as such

is involved in sustaining normal root hair elongation in coordination with actin.

BRK1 and SCAR2 are Molecular Determinants of the Root Hair Initiation Domain

The identification of *SPI* as one of the genes in the *DIS* group that also included genes

encoding subunits of the W/SRC-ARP2/3 complex (Saedler et al., 2009) raises the

possibility that SPI might function in root hair developmental pathways mediated by

W/SRC-ARP2/3. Furthermore, the observation that SPI-YPET accumulation at the root

hair tip (Figure 1) mirrors the enrichment of BRK1-YFP and APRC4-GFP at apex of

Physcomitrella protonemal cells (Perroud and Quatrano, 2006, Perroud and Quatrano,

2008) raises the possibility that W/SRC-ARP2/3 maybe a root hair tip-localized

complex. In an attempt to link SPI with the W/SCR-ARP2/3 complex, we imaged roots

of *brk1* complemented with *BRK1promoter:BRK1-YFP* (from here on referred to as

BRK1-YFP) (Figures 4A-C) (Dyachok et al., 2008). Unlike SPI-YPet, we did not

observe a BRK1-YFP fluorescence gradient in rapidly elongating root hairs. However,

closer examination of trichoblasts revealed prominent BRK1-YFP signal at the plasma

membrane of the RHID that mirrored other known early root hair initiation site markers

such as ROP (Figure 4A). Unlike ROP, which had a persistent plasma membrane

localization throughout root hair development (Jones et al., 2002, Molendijk et al.,

2001), BRK1-YFP signal was transient and dissipated as root hairs transitioned to rapid

tip growth (Figure 4B; Supplemental Movie 6). This observation was in contrast to the

intensification of SPI-YPet fluorescence as root hairs proceeded with rapid tip growth

(see Figure 1). In agreement with our visual observations, a linear regression analysis

showed that BRK1-YFP fluorescence is inversely proportional to root hair growth rate

(Figure 4C).

Given that BRK1 stabilizes the entire family of SCAR proteins and is required for

functional W/SRC assembly (Le et al., 2006), we investigated if the SCAR protein

localized to the RHID, similarly to BRK1. To address this question, we imaged a

recombineered SCAR2-mCherry fusion expressed in the *scar1 scar2 scar3 scar4*

(*scar1234*) quadruple mutant. We found that like BRK1-YFP, SCAR2-mCherry marked

the RHID and dissipated when rapid root hair tip growth commenced (Figure 4D).

The accumulation of BRK1-YFP and SCAR2-mCherry at the RHID led us to

hypothesize that *brk1* and *scar1234* might have defects in root hair initiation. One

parameter that has been studied extensively as an indicator of root hair initiation defects

is planar polarity, which is a measure of root hair position along the length of the

trichoblast (Nakamura and Grebe, 2018). We found that root hair position of *brk1* and

scar1234 shifted apically (i.e. toward the shoot) when compared to wild type (Figure

4F). The planar polarity defects of *brk1* was rescued by expressing *BRK1-YFP* in *brk1*

while partial complementation of *scar1234* was achieved by expressing *SCAR2-*

mCherry in *scar1234* (Figures 4E and F). Taken together, our results revealed that

BRK1 and SCAR2 are new molecular determinants of the RHID and are required for

specifying the position of root hair emergence.

Given that ARP2/3 is the known target of activated W/SRC, we hypothesized that

ARP2/3 might also localize to the RHID. To test this hypothesis, we imaged a line

expressing ARPC5-GFP, one of the subunits of the ARP2/3 complex (Yanagisawa et

al., 2015). This line has been confirmed to complement the *arpc5* trichome defects. As

reported previously, ARPC5-GFP labeled the plasma membrane of initiating trichome

branches and the apex of elongating trichome branches (Yanagisawa et al. (2015);

Supplemental Figure 4A and B). Nevertheless, when observed in root hairs, the

ARPC5-GFP signal was distributed uniformly at various stages of development with no

distinct accumulation at the RHID or root hair apex (Supplemental Figure 4C and D).

Although there was no clear polarized localization of ARPC5-GFP in root hairs, it is

possible that ARP2/3 is locally activated in RHID or root hair apex, but that active pool is

obscured by a large soluble pool of the complex in this cell type.

SPI is Required for the Depletion of BRK1 as Root Hairs Transition to Tip Growth

Live cell microscopy of BRK1-YFP and SPI-YPet revealed contrasting spatial and

temporal dynamics, with the former intensifying and the latter dissipating as root hairs

elongated (Figure 1 and 4). To observe BRK1 and SPI simultaneously within the same

root hair, we generated plants expressing both *SPI-YPet* and *BRK1-mRuby3*. Similar to

BRK1-YFP, BRK1-mRuby3 labeled the RHID and dissipated as the root hair bulge

expanded (Figure 5A). Within the same root hair cell, SPI-YPet fluorescence at the tip

intensified as BRK1-mRuby3 dissipated (Figure 5A; Supplemental Movie 7). The

depletion of BRK1 coinciding with the accumulation of SPI suggests that these proteins

may be antagonistic to each other. To address this question, we expressed *BRK1-YFP*

in *spi* and *SPI-YPet* in *brk1*. We found that unlike BRK1-YFP in the complemented *brk1*

background (Figure 4B), BRK1-YFP signal in *spi* persisted throughout the entire

imaging time course. In several cases, BRK1-YFP remained visible in non-growing root

hair bulges of *spi* for more than 60 min (Figure 5B). The persistence of BRK1-YFP

signal was also observed in short root hair outgrowths of *spi* (Figures 5C; Supplemental

Movie 8). By contrast, SPI-YPet in *brk1* exhibited similar dynamics as SPI-YPet in the

complemented *spi* (Figure 5D). In *brk1*, SPI-YPet signal was weak during early root hair

bulge formation and intensified as root hair tip growth accelerated (Figure 5D). Taken

together, these results indicate that SPI is antagonistic to BRK1 during the transition

from root hair initiation to rapid tip growth.

***SPI* Genetically Interacts with *W/SRC-ARP2/3* in Root Hairs**

The phenotypic overlap between *SPI* and *W/SCR-*and *ARP2/3* mutants suggest that

SPI participates in similar biological processes (Saedler et al., 2009). The polarized

localization of SPI, BRK1 and SCAR2 in root hairs observed here also indicate potential

functional relationships between SPI and W/SCR-ARP2/3 during root hair development.

However, it remains to be determined whether *SPI* genetically interacts with *W/SRC-*

ARP2/3 pathway components. To address this question, we generated double *spi brk1*

mutants and compared their root hair phenotypes with single mutants. We first

investigated the genetic relationship between *SPI* and *BRK1* based on length of mature

root hairs. We found that at low magnification, root hair length of *brk1* appeared

morphologically similar to wild type (Figure 6A). Quantification of root hair length,

however, uncovered a small but significant reduction in root hair length in *brk1* when

compared to wild type (Figure 6B). On the other hand, *spi brk1* had short root hairs that

were similar to *spi* (Figure 6A and B). We next examined whether *SPI* and *BRK1*

genetically interact in specifying root hair planar polarity. We found that *spi* displayed a

more shoot ward (apical)-shifted root hair position when compared to wild type and

brk1. Quantification of root hair planar polarity showed that that root hair position of *spi*

was identical to *spi brk1* (Figure 6C). Taken together, our results indicate that *SPI* is

epistatic to *BRK1* for root hair planar polarity and tip growth.

Epistasis between *SPI* and *BRK1* prompted us to expand our genetic interaction

studies to other genes in the *ARP2/3* pathway. We therefore generated *spi arp2*, *spi*

arp3 and *spi arpc5* double mutants and analyzed their root hair phenotypes. Like *brk1*,

root hairs of *arp2, arp3* and *arpc5* appeared to be morphologically similar to wild type

(Figure 7A; Supplemental Figure 5). Quantification of root hair length, however,

revealed slightly shorter root hairs in the single *arp2/3* mutants when compared to wild

type (Figure 7C), an observation consistent with the mild tip growth defects reported for

these mutants in the presence of the GFP:TALIN live cell probe for actin (Mathur et al.,

2003). On the other hand, *spi arp2*, *spi arp3* and *spi arpc5* appeared morphologically

hairless when viewed at low magnification (Figure 7; Supplemental Figure 5). This

observation was in contrast to *spi brk1* and *spi,* which had a number of short, but visible

root hair out growths spaced between hairless regions (Figures 6A, 7A, and 7BB;

Supplemental Figure 5). When viewed at high magnification we found that most of the

trichoblasts of *spi arp2*, *spi arp3* and *spi arpc5* had a high number of root hair bulges

that aborted tip growth soon after initiation (Figure 7B; Supplemental Figure 5).

Quantification of root hair length revealed that the joint contribution to the root hair

length phenotype made by combining *SPI* and *ARP2/3* complex mutations was greater

than an additive effect of *SPI* or *ARP* single mutations (Figure 7C). The enhancement of

the root hair length defect of *spi* by *arp2/3* single mutants indicate a synergistic

interaction between *SPI* and *ARP2/3* with regard to root hair tip growth.

439   We next investigated if *SPI* and *ARP2/3* genetically interact in specifying the

position of the RHID. We found a mild but statistically significant shoot ward shift in the

position of the RHID in *arp2*, *arp3* and *arpc5* when compared to wild type. The extent to

which the position of the RHID shifted towards the shoot in single *arp2/3* complex

mutants was less than that of *spi* and *brk1* (compare Figure 6C and 7D). On the other

hand, the position of the RHID in *spi arp2/3* was more similar to single *arp2/3* indicating

that *ARP2/3* is epistatic to *SPI* for root hair planar polarity (Figure 7C and D). In many

cases we observed that the aborted root hair bulges in *spi arp2/3* emerged very close to

the basal (root ward) end wall of the trichoblast (Figure 7C). The synergistic and

epistatic relationship between *SPI* and *ARP2/3,* for root hair length and planar polarity,

respectively, indicate that *SPI* also genetically interacts with *ARP2/3* during root hair

development. However, the nature of this interaction appears to differ from that of

BRK1.

452   Genetic interaction between *SPI* and *W/SRC-ARP2/3* was also analyzed for

other *spi* phenotypes such as primary root hypersensitivity to LatB (Supplemental

Figure 1). For LatB sensitivity, we found that *arpc5* primary roots were inhibited to the

same extent as wild type when grown on LatB (Supplemental Figure 6B). Although

primary roots of *arp2*, *arp3* and *brk1* exhibited some hypersensitivity to LatB, the

response of these mutants to the compound was not as severe as *spi* and leaned more

toward wild-type sensitivity. By contrast, the hypersensitivity of primary roots of *spi arp2*,

spi arp3, *spi arpc5* and *spi brk1* to LatB was more similar to *spi* (Supplemental Figure

6A and B) suggesting genetic interactions also occurs between *SPI* and *W/SCR-*

ARP2/3 for this phenotype.

462   Genetic interaction between *SPI* and *W/SRC-ARP2/3* was also observed in

trichomes (Li et al., 2003, Mathur et al., 2003, Schwab et al., 2003). Consistent with

previous reports, *arp2*, *arp3*, *arpc5* and *brk1* had severe trichome shape defects when

compared to wild type, while those of *spi* were mild when viewed with a scanning

electron microscope. One particular trichome defect is branch length, in which *arp2*,

arp3, *arpc5* and *brk1* trichomes have distinctively short branches when compared to

wild type (Supplemental Figure 7). On the other hand, trichomes of *spi* were wavy and

crooked, but not as short as those of *arp2*, *arp3*, *arpc5* and *brk1*. We found that

trichomes of all the double mutants resembled those of *spi* (Supplemental Figure 7A).

Quantification of branch length of all single and double mutants was consistent with

visual observations, especially in showing that *spi brk1* and *spi arp2/3* resembled the

branch length of *spi* (Supplemental Figures 7A and C). Our observations on trichome

branch length provide additional evidence that *SPI* genetically interacts with *W/SRC-*

ARP2/3.

DISCUSSION

Our work uncovers new insights underlying actin-mediated root hair development. A major result from our studies is the revelation that SPI is a root hair tip-localized protein. Although SPI fused to fluorescent proteins was reported to localize to endosomes and P-bodies, such studies have been limited to transient expression assays in biolistically-bombarded leaves (Steffens et al., 2015; Steffens et al., 2017). As such, a mechanistic link between reported SPI subcellular localization patterns and cell growth phenotypes (i.e. short root hairs and distorted trichome shapes) described in the original *spi* mutant alleles has never been demonstrated. The large size of SPI and the possibility that it is targeted to discrete cellular domains could have hindered transgenic complementation and subsequent *in planta* localization efforts. Through our work, we have addressed this knowledge gap and now show that a SPI-YPet fusion generated through recombineering has complemented *spi* (Brumos et al., 2020, Zhou et al., 2011). In doing so, we provide compelling evidence that SPI is a root hair tip–enriched BEACH domain-containing protein.

Our studies indicate that SPI mediates root hair tip growth by maintaining the tip-focused fine F-actin meshwork, although the precise mechanism by which this is accomplished is unknown. The root hair tip-focused F-actin meshwork is the functional equivalent of domain-specific actin structures found in other tip-growing cells such as cortical actin fringes in pollen tubes and actin spots in moss protonemata (Bascom et al., 2018a). Although the tip-focused F-actin meshwork was able to form in a population of *spi* root hairs, it often dissipated or was replaced by thick actin cables, a feature of mature wild-type root hairs that are terminating tip growth. The root hair phenotypes and corresponding depletion of tip-focused F-actin in *spi*is reminiscent of studies of BEACH domain-containing proteins in mammalian cells, particularly in neurons, which have often been compared to tip-growing plant cells (Baluška, 2010). For instance, neuronal dendritic spines are cytoplasmic protrusions that modulate excitatory synaptic transmission in the mammalian nervous system. Actin is a major component of dendritic spines and it plays a central role in driving the dynamic shape changes and secretory activities of these cytoplasmic protrusions during synaptic signaling (Cingolani and

Goda, 2008). In animals, neurobeachin (Nbea) is a BEACH domain-containing protein

that has a similar domain architecture as SPI. Cultured neurons of *Nbea* mice

knockouts had fewer dendritic spine protrusions and depleted F-actin at the synapse

(Niesmann et al., 2011). Another BEACH domain-containing protein called FAN was

shown to be crucial for the formation of filopodia, actin-rich plasma membrane

extensions that enable motile cells to probe their environment (Mattila and Lappalainen,

2008). FAN deficient fibroblasts had reduced filopodia formation and were unable to

reorganize their actin cytoskeleton in response to upstream activation by tumor necrosis

factors (Haubert et al., 2007). Taken together, these results indicate that actin-mediated

regulation of polarized cell growth by BEACH domain-containing proteins is likely to be

conserved across animals and plants.

The formation of SPI-YPet agglomerates in response to BFA indicates that SPI is

associated with post-Golgi vesicles. This result is consistent with observations made

with BEACH domain-containing proteins in mammals. For example, Nbea was shown to

localize to vesicular endomembranes adjacent to the *trans*-Golgi network, and like SPI,

the distribution of Nbea-positive post-Golgi vesicles was altered by BFA (Wang et al.,

2000). Furthermore, the *Caenorhabditis elegans* Nbea homolog, SEL-2, was

demonstrated to function in endomembrane traffic in polarized epithelial cells based on

the finding that *sel-2* mutants mistarget proteins normally found in the apical cell surface

to the basolateral surface (de Souza et al., 2007). Here, the protein secretory marker

SEC-RFP, which is typically trafficked to the tips of elongating wild-type root hairs

(Sparks et al., 2016), was uniformly distributed in *spi* root hairs. The absence of tip-

directed SEC-RFP gradients in *spi* root hairs shows that, as in mammals, plant BEACH

domain-containing proteins are required for proper targeting of molecular cargo to

points of polarized cell growth.

Another significant result from our studies is the discovery that BRK1 and SCAR2

are molecular determinants of the RHID. The localization of BRK1 and SCAR2 to the

RHID and their dissipation as root hair tip growth commenced, contrasts with findings in

Physcomitrella in which a functional BRK1-YFP distinctly labeled the tips of caulonemal

cells (Perroud and Quatrano, 2008). Like BRK1-YFP, an ARPC4-GFP construct labeled

the tips of caulonemal cells (Perroud and Quatrano, 2008) whereas no clear tip-focused

labeling of ARPC5-GFP was observed in root hairs. The observation that BRK1-YFP

prominently localized to the RHID, but dissipated during active root hair tip growth was

surprising given that the *Arabidopsis BRK1* complemented the defective filamentous

growth of *Physcomitrella BRK1* knockouts (Perroud and Quatrano, 2008). This suggest

that seed plants might have evolved a specialized function for the W/SCR-ARP2/3

complex in which it acts predominantly at the site of root hair emergence and early

bulge formation with only a minor role in driving actin-dependent tip growth. The fact

that *Physcomitrella BRK1 and ARPC4* mutants have clear tip growth defects, while

Arabidopsis W/SCR-ARP2/3 mutants have only weak to no tip growth abnormalities,

supports this possibility (Le et al., 2003, Mathur et al., 2003, Perroud and Quatrano,

2006, Perroud and Quatrano, 2008). By contrast, knockouts of an *Arabidopsis SPI*

orthologue in the liverwort, *Marchantia polymorpha*, led to short rhizoids indicating

conserved functions of SPI across land plants in regulating tip growth (Honkanen et al.,

2016). In other single cell types such as diffusely-growing trichomes, W/SCR-ARP2/3

subunits likely play a more prominent role than SPI based on their more severe

trichome phenotypes when expression of their encoding genes are suppressed.

However, the epistatic relationships between *SPI* and *W/SCR-ARP2/3* for trichome

morphology and primary root hypersensitivity to LatB as reported here continue to point

to functional relationships between these genes in diffusely-growing cells. The

observation that phenotypes of *spi w/scr-arp2/3* tilt more toward *spi* indicate that SPI

might negatively regulate an actin pathway that is parallel to ARP2/3.

The weakening of BRK1-mRuby3 fluorescence coinciding with SPI-YPet

accumulation, and the persistence of SPI-YPet signal in *spi* root hair tips, provides

indirect evidence that SPI might play a role in mediating BRK1 stability or localized

clustering at the plasma membrane of the RHID. Because *spi brk1* root hair phenotypes

resemble *spi*, ectopic expression of BRK1 does not appear to contribute to the *spi*

phenotypes. Although, it is unknown why BRK1-YFP signal persists in *spi,* it is tempting

to speculate that SPI might modulate BRK1 via protein degradation pathways. This

possibility is supported by studies in mammals pointing to a role for BEACH domain-

containing proteins in protein degradation via the ubiquitination pathway. In mouse

models for example, the BEACH domain-containing protein, WDR81, was shown to be

essential for removal of autophagy-dependent ubiquitinated proteins (Liu et al., 2017).

In this regard, it is worth noting that the W/SCR-ARP2/3 complex was demonstrated to

function in stress-induced autophagy (Wang et al., 2016) and proteasome inhibitors

stabilized SCAR during dark-induced primary root growth inhibition (Dyachok et al.,

2011). It is possible that SPI-mediated proteolytic pathways and BRK1 operate

antagonistically to specify the levels of W/SCR at the RHID that enables the transition to

actin-dependent rapid tip growth. However, such as scenario is complicated by the

observation that the formation of tip-directed SPI-YPet does not appear to require

BRK1. This suggest that SPI's appearance at the root hair tip is regulated by other

factors besides W/SCR.

The functional links between SPI and W/SCR-ARP2/3 during root hair

development as suggested by live cell imaging is supported by the double mutant

studies reported here. We found that *SPI* is epistatic to *BRK1* for RHID and root hair tip

growth. For *SPI* and *ARP2/3*, synergistic and epistatic genetic interactions were

observed for root hair tip growth and root hair planar polarity, respectively. Synergistic

genetic interactions can arise when converging pathways are disrupted (Pérez-Pérez et

al., 2009), and the severe root hair phenotypes of *spi arp2/3* double mutants may reflect

parallel pathways that are involved in actin-dependent root hair tip growth. The

detection of synergistic genetic interactions in root hairs with *ARP2/3* mutants, but not

with *brk1* is unexpected as *brk1* is null for W/SRC-dependent ARP2/3 activation (Le et

al., 2006), and W/SRC is the only known ARP2/3 activator in *Arabidopsis* (Zhang et al.,

2008). These genetic data may reflect the existence of an unknown ARP2/3 activator or

an NPF-independent function of ARP2/3. Future studies will probe deeper into the

regulatory relationships between *SPI* and *W/SCR-ARP2/3* that underlie these

unexpected genetic interactions.

In summary, our work provides new data that contribute to our understanding of

actin-mediated root hair development. A crucial result from our work is the discovery

that SPI and the W/SCR subunits, BRK1 and SCAR2, exhibit polarized localization

patterns in root hairs that point to potential functional relationships among these

proteins during root hair development. Based on our cell biological studies, we propose

a model for which new hypotheses can be tested (Figure 8). We propose that a RHID-

localized ROP/RAC-GTPase (Jones et al., 2002, Molendijk et al., 2001) activates

W/SCR during root hair initiation in a manner that mirrors that of the DOCK family GEF,

SPK1, which was shown to occur during trichome morphogenesis (Basu et al., 2008).

Although the precise ROP-based machinery that facilitates this process during root hair

development has yet to be determined, ROP-GEF3 could be the functional equivalent of

SPK1 given that it localizes to the RHID (Denninger et al., 2019) in a manner similar to

BRK1 and SCAR2. Although epistatic and synergistic genetic interactions presented in

our double mutant studies support the occurrence of overlapping pathways between

SPI and *W/SCR-ARP2/3*, the precise nature of such relationships remain unclear. One

plausible scenario is that SPI negatively regulates an actin pathway that is parallel to

W/SCR-ARP2/3. The persistence of BRK1-YFP signal in *spi* suggests that one

mechanism for negative regulation could occur at a node in which SPI and W/SCR

converge. For the future, it will be important to determine whether SPI physically

interacts with actin or subunits of the W/SCR-ARP2/3 complex to better explain the

functional links between SPI and W/SCR-ARP2/3.

METHODS

Forward-Genetic Screening and Map-Based Cloning

The *hlb* mutants from which the *spi-5* mutant allele was identified were isolated from a population of Transfer-DNA (T-DNA) seed (Arabidopsis Biological Research Center stock CS31100). Plants from which the seeds were derived from were transformed with the activation tagging pSKI015 plasmid (Weigel et al. 2000; Sparks et al., 2016; Sun et al., 2019). Briefly, mutagenized seeds were surface sterilized in 95 % (v/v) ethanol and 20% bleach (v/v), followed by three washes in sterile deionized water. A solution of 0.5 x Murashige and Skoog (MS) basal salt medium with vitamins (PhytoTech Labs, USA) and 1 % (w/v) sucrose was prepared, and the pH of the solution was adjusted to 5.7. After adding 0.5% agar (w/v) (Sigma-Aldrich), the solution was autoclaved and allowed to cool to room temperature. Upon reaching 55ºC, a stock solution of 10 mM LatB (CalBiochem-EMD Chemicals) in 100% Dimethyl Sulfoxide (DMSO) was added to make a final LatB concentration of 100 nM. Sterilized seeds were suspended in the MS-agar-LatB medium and gently swirled to evenly distribute the seed. The seed-MS-agar-LatB mixture was poured to a thickness of 2 mm on the base of 10 cm x 10 cm Petri dishes. After keeping plates at 4ºC for 48 h, they were positioned vertically in a Conviron set to 24°C with a 14 hours light (120 µmol m^{-2}s^{-1})/ 10 hours dark cycle. Six days after transfer to the Conviron growth chamber, seedlings that exhibited severe growth inhibition were transplanted to LatB-free media and grown to maturity.

For LatB hypersensitivity assays, seed from selected plants and the *hlb2* mutant were planted on the surface of a 3 mm layer of polymerized 0.5 x MS - 1% agar growth media in gridded square 10 cm × 10 cm Petri plates and grown in the same Conviron used for screening. Four-day-old seedlings with primary roots that were about 1 cm long were selected. Selected seedlings were transplanted to a new set of square Petri dishes containing MS medium supplemented with 50 nM LatB or MS supplemented with the appropriate volume of LatB solvent (i.e. DMSO). During transplant, the tip of the root was positioned at the grid line of the Petri dish and maintained in a vertical orientation. Four days after transplanting, images of the roots were captured with a Nikon Insight

digital camera mounted on a copy stand. Primary root length was expressed as the

distance between the position of the root tip 4 days after transplant and the grid line

where the root tip was positioned during transplant (Supplemental Figure 1).

To identify the *HLB2* gene, homozygous *hlb2* (Col-0 ecotype) was out-crossed to

the Landsberg *erecta* ecotype to generate seeds for map-based cloning because

attempts to identify the responsible mutation for *hlb2* phenotype using TAIL-PCR had

been unsuccessful. Segregating F2 seedlings were surface sterilized as described

above and grown for three days. These seedlings were then transferred to MS media

containing 50 nM LatB and root lengths were marked on the plates to track root growth.

Transferred seedlings were grown vertically for an additional four days in the growth

chamber and seedlings that showed root hypersensitivity to 50 nM LatB were selected

for mapping. Briefly, DNA was extracted (Edwards et al., 1991) from about 2600 LatB

sensitive seedlings and cloned with simple sequence length polymorphism (SSLP) and

cleavage of amplified polymorphic site (CAPS) markers to chromosome one between

AT1G02740 and *AT1G03410* loci (Lukowitz et al., 2000), which spanned 248 kb with 77

annotated genes. Primers were then designed to several candidate genes based on

the sequenced 248 kb region. Nucleotide sequencing revealed that *hlb2* mutation had a

10 base pair deletion (Chr1 position 720,152- 720,161) in exon 14 of the *AT1G03060*

gene. In addition, we obtained a T-DNA insertional mutant (SALK_065311) from the

ABRC with predicted insertion in *SPI*. After genotyping, SALK_065311 was subjected to

similar LatB hypersensitivity and growth assays as *hlb2* as aforementioned. Allelism

was determined by examining the F1 progeny from a cross between *hlb2* ×

SALK_065311. Following the nomenclature of Steffens et al. (2015), SALK_065311 and

hlb2 were named *spi-3* and *spi-5*, respectively.

Generation of Fluorescent Protein-Tagged Constructs and Plant Lines

The SPI protein was tagged at the C-terminus with the fluorescent protein 3x-YPet by

the recombineering method (Zhou et al., 2011). *Agrobacterium tumefaciens* UIA143

pMP90 harboring a SPI-YPet recombineering construct was used to transform the *spi-5*

mutant by the floral dip method (Clough and Bent, 1998). Transgenic lines were isolated

based on resistance to kanamycin and restoration of wild-type root length. Presence of

the SPI-YPet transgene was amplified with primers (5'-

ATTCCACAAGCAACCAGTCAC-3' and 5'-AACAGAGTTGAGAGTGGCTCG-3') and

sequenced to confirm the correct configuration. Final validation of *SPI-YPet* expression

was accomplished by screening selected lines under the confocal microscope for YPet

fluorescence and complementation of the short root hair phenotype.

SCAR2-mCherry lines were also generated by recombineering in which an

mCherry tag was fused to an internal region of SCAR2 (Sharan et al., 2009). The

mCherry tag was inserted after amino acid 585 in SCAR2 to generate SCAR2-

i2mCherry. Recombineering primers were designed as follows: the forward primer

contained 50 SCAR2 nucleotides upstream from the insertion site of SCAR2 and 18

nucleotides of the 5' end of the mCherry cassette, which had a 5x Glycine, 1 Alanine

linker. The reverse primer contained 50 nucleotides after the insert site of SCAR2 and

24 nucleotides of the 3' end of the mCherry cassette. The JAtY75L14 clone was

transformed into the recombineering competent *E.coli* strain SW105 and recombined

with the SCAR2-i2mCherry recombineering cassettes. A flippase recombination

reaction removed the ampicillin resistance marker. The sequence between the two test

primers was verified by DNA sequencing. The clones were transformed into

Agrobacterium and then into *scar1234* plants (Dyachok et al., 2008). Because both

scar1234 plants and the recombineering clones were Basta resistant, plants on MS

plates were screened by eye for rescued trichome phenotypes. Rescued plants were

genotyped with the recombineering test primers and screened for fluorescence with a

confocal microscope.

The *BRK1promoter: BRK1-YFP* introduced into the *brk1-1* mutant and *ARPC5-*

GFP are described in Dyachok et al. (2008) and Yanagisawa et al. (2015), respectively.

For the *BRK1-mRuby3* construct, the fluorescent protein 3x-mRUBY3 was tagged with

a C-terminal linker (10 Alanine, Glycine) using Thermo Fisher Scientific GeneArt to

include BamH1 and XbaI sites at its 5' and 3' ends, respectively. Codon optimization for

Arabidopsis was performed on the 3x-mRUBY3 and internal linkers prior to synthesis.

The 3X-mRuby3 fragment was inserted as a BamH1/XbaI into the plasmid

pBRK:YFPpEZRK (Dyachok et al., 2008).The *Agrobacterium* floral dip method (Clough

and Bent, 1998) was used to transform *brk1-2* plants, and transformation was confirmed

when this construct fully restored defective *brk1-2* trichomes.

Plants expressing *BRK1-YFP* were crossed with *spi-5* to generate *BRK1-YFP* in

spi-5 lines. In parallel, plants expressing *SPI-YPet* were crossed with *brk1* to obtain SPI-

YPet in *brk1* lines. *Spi-5* was directly transformed with the *UBQ10:mGFP-Lifeact*

construct by the floral dip method (Clough and Bent, 1998). *Spi-5* expressing SEC-RFP

was generated by crossing *spi-5* with *SEC7-RFP*-expressing wild-type plants and

progeny in subsequent generations that exhibited fluorescence and the *spi* root hair

phenotypes were selected for analysis.

Generation of Double Mutants and Dual Fluorescent Protein-Labeled Plant Lines

Arp2 (SALK_077920), *arp3* (SALK_010045), *arpc5* (SALK_123936) and *brk1-1*

(Djakovic, 2006, Le et al., 2006, Li et al., 2003) were crossed with *spi-5* to generate

double mutants. Single mutants were obtained from the ABRC and genotyped to verify

zygosity. Progeny from crosses were genotyped for the presence of the double mutation

by PCR-based approaches using the primers: ARP2LP 5'-

CGCTTTTTACTCCGGTAAACC-3' ARP2RP 5'-CATGCAGTAGAAGGAAGCTGG-3'

ARP3LP 5'-AATTGCTGGCAAAGATGTCAC-3' ARP3RP 5'-

AGCTCTTCGTGTGTCAATTGG-3' ARPC5LP 5'-TGCAAATTCCACCAAATCTTC-3'

ARPC5RP 5'-CTTTTGAAATGGTCAAATGGC-3'. The *spi-5* mutation in the double

mutants was confirmed using primers SPIF 5'-CAGTTACGAGGATCTTAGTCTTTC-3'

and SPIR 5'-CCAGTCCAGAAGGACATGTTC-3'. For *brk1-1*, which had a single base

pair deletion (Djakovic, 2006), PCR products were amplified using BRK1F 5′-

TTGAATCGAAAAAGGGCGAAT-3′ and BRK1R 5′-CGAAATCGAGAAAACTCCAAAA-3′

and the resulting fragment was sequenced to ensure deletion using the BigDye

Terminator chemistry (v3.1, Thermo Fisher Scientific) on an ABI3730XL DNA

sequencer (Thermo Fisher Scientific) following the manufacturer's instruction.

To generate plant lines co-expressing *SPI-YPet* and *BRK1-mRuby3, SPI-YPet* in

spi-5 was crossed with *BRK1-mRuby3* in *brk1-2*. Progeny from subsequent generations

that exhibited yellow and red fluorescence and rescued root hair and trichome

phenotypes were selected for analysis. For lines expressing *SPI-YPet* and *mRuby3-*

Lifeact (Bascom et al., 2018b), *SPI-YPet* in *spi-5* was directly transformed with a

UBQ10:Lifeact-mRuby3 construct. Seedlings that showed both YPet and mRuby3

fluorescence were selected under the confocal microscope.

Evaluation of Root Hair Length and Planar Polarity

Seeds of mutants were surface-sterilized in ethanol and bleach as described above. To

evaluate root hair length and planar polarity, seeds were planted on 48 × 64 mm

coverslips coated with 0.5 x MS, 1% sucrose and 0.4 % (w/v) Gelzan ^{TM} CM (Sigma

Aldrich, USA) according to Dyachok et al. (2016). Coverslips were placed in 9 cm round

Petri dishes and stratified in 4ºC for 48 h. After stratification, the coverslip system with

planted seed were kept in a 24°C growth chamber with 14 hours light (120 µmol m^{-2}s^{-1})/

10 hours dark cycle for 5 to 6 days.

To quantify root hair length, fully-grown root hairs from a region of the primary

root located between 5 to 15 mm from the root tip were photographed with a Nikon

SMZ1500 stereomicroscope. For planar polarity measurements, images of trichoblasts

in which the apical and basal end walls were clearly visible were acquired with a Nikon

Eclipse TE300 inverted microscope using a 20x objective. Root hair lengths were

extracted from digital images using ImageJ (v1.51) software (https://imagej.nih.gov/ij/).

For planar polarity, the distance between the basal trichoblast wall and base of the

emerging root hair (a) and the length of the trichoblast (b) were obtained using Image J.

The ratio of a/b was used as an indicator of relative root hair position with a value of 0

indicative of root hairs emerging at the basal-most end of the trichoblast and 1 indicative

of root hairs emerging at the apical-most end of the trichoblast (Fischer et al., 2006).

Box and violin plots were drawn on R software (R Core Team, 2019) using ggplot

function in ggplot2 package (Wickham, 2016). Two-way Analysis of Variance (ANOVA)

statistical analysis and Tukey's post-hoc tests were performed on R software (R Core

Team, 2019) using car (Fox and Weisberg, 2019) and agricolae (de Mendiburu, 2020)

packages. Pairwise non-parametric Kolmogorov-Smirnov tests were done using ks test

function available in the basic package (R Core Team, 2019).

Microscopy and Image Analysis

Live cell imaging of root hairs using confocal microscopy was performed on 4 or 5-day-old seedlings grown on the 48 mm × 64 mm coverslip system described above. Coverslips with the seedlings were placed horizontally on the stage of an inverted Leica SPX-8 point scanning confocal microscope (Leica Microsystems, Buffalo Grove, Illinois) or an UltraView ERS spinning-disc confocal microscope (Perkin Elmer Life and Analytical Sciences, Waltham, Massachusetts) equipped with 40 × water (numerical aperture=1.10) or 100 × oil (numerical aperture=1.40) immersion objectives. SPI-YPet and YFP-BRK1 were imaged by illuminating roots growing along the coverslip surface with the 514 nm line of the SPX-8 Argon laser and emission detected at 527 nm. Images of root hairs expressing SCAR2-mCherry, BRK1-mRuby3, Lifeact-mRuby and SEC7-RFP were acquired through illumination with the tunable SP8-X white light laser (560-580 nm) and detecting emission at 610 nm. Excitation and emission parameters for GFP (GFP-ARPC5 and mGFP-Lifeact) were 488 nm and 510 nm, respectively. Time-lapse movies or single time point images were collected using Volocity acquisition version 6.3.5 (Improvision) or SPX-8 LAS software, for the UltraView and Leica SP8-X, respectively.

Quantification of fluorescence from root hair images was conducted on 8-bit confocal images acquired at a single fixed focal plane that spanned the median of the cell (SPI-YPet, BRK1-YFP and SEC-RFP) or from maximum projected images (mGFP-Lifeact). For SPI-YPet and SEC-RFP, an oval region of interest (ROI) at the root hair tip was drawn and mean fluorescence within this area was acquired using Image J. Fluorescence was expressed as the ratio of mean fluorescence within the root tip ROI to background fluorescence. For SPI-YPet, the background used was the region adjacent, but outside the root hair tip (Figure 1C) while for SEC-RFP, the background used was an area on the sub-apical region of the root hair tip (Figure 2F). For BRK1-YFP, a ROI was drawn along the apical-most root hair tip that was about 20 pixels-wide using the selection brush tool of image J. The ratio of fluorescence within this area to background fluorescence was obtained (Figure 2C). Root hair growth rate data was

derived from the same root hair images in which fluorescence images were acquired.

For the former, an image of a root hair was taken at time 0 and every 5 minutes

thereafter. The growth rate refers to the displacement of the root hair tip in μm divided

by time elapsed (min).

To quantify tip-focused F- actin, 25 optical sections were taken at 0.5 μm

intervals using the UltraView spinning-disc confocal microscope. Raw Ultraview Z-

stacks were exported to Imaris image analysis software version 9.2.0 (Bitplane).

Transverse sections of the root hair tip were obtained from Z-stacks using the surpass

view interface of the Imaris software and exported as 8-bit TIFF files. From these

images, an ROI spanning the circular area of the root hair tip was drawn using image J

and mean fluorescence was extracted (Figure 3O). The ratio of the tip fluorescence to

background was obtained from 18- 21 root hairs.

For SPI-YPet and BRK1-YFP, a scatter plot to determine the relationship

between growth rate and tip-focused fluorescence was done in R (R Core Team, 2019)

using ggplot function in ggplot2 package (Wickham, 2016). Linear regression analysis

were performed using lme4 package on R software (Bates et al., 2015, R Core Team,

2019).

Trichomes were imaged using a Hitachi Table top TM3000 scanning electron

microscope or with a Nikon SMZ 1500 stereo microscope. Images of trichomes from

newly emerging leaves of 3-week-old plants were collected and branch length was

measured using Image J.

Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome

Initiative or GenBank/ EMBL databases under the following accession numbers: *SPI*

(AT5G61350), *BRK1* (AT2G22640), *ARP2* (AT3G27000), *ARP3* (AT1G13180), *ARPC5*

(AT4G01710), *SCAR1* (AT2G34150), *SCAR2* (AT2G38440), *SCAR3* (AT1G29170),

SCAR4 (AT5G01730)

Supplemental Data

Supplemental Figure 1. Primary Root Growth of *hlb2* Seedlings is Hypersensitive to LatB.

Supplemental Figure 2. *HLB2* Encodes the Beach Domain Containing Protein, SPIRRIG.

Supplemental Figure 3. *SPI-YPet* construct complements *spi*

Supplemental Figure 4. ARPC5-GFP is localized uniformly in the cytoplasm of developing root hairs.

Supplemental Figure 5. Root hair phenotypes of *spi arp2/3* mutants

Supplemental Figure 6. *SPIpro:SPI-YPet* complements the primary root hypersensitivity to LatB, short root hair and distorted trichome phenotypes of *spi*.

Supplemental Figure 7. Distorted trichome defects of *spi arp2*, *spi arp3*, *spi arpc5* resemble *spi*.

Supplemental Movie 1. Time-Lapse Confocal Microscopy of SPI-YPet in a Rapidly Elongating Root Hair.

Supplemental Movie 2. Time-Lapse Confocal Microscopy of SPI-YPet in a Root Hair Bulge Transitioning to Tip Growth.

Supplemental Movie 3. Time-Lapse Confocal Microscopy of SPI-YPet in a Root Hair during Termination of Tip Growth.

Supplemental Movie 4. Spinning-Disc Confocal Microscopy of Lifeact-mGFP in Elongating Root Hairs of Wild type and *spi*.

Supplemental Movie 5. Spinning-Disc Confocal Microscopy of a Rapidly Elongating

Root Hair Co-Expressing SPI-YPet and mRuby3-Lifeact.

Supplemental Movie 6. Time-Lapse Confocal Microscopy of BRK1-YFP Root Hair

Bulge Transitioning to Tip Growth.

Supplemental Movie 7. Time-Lapse Confocal Microscopy of a Root Hair Bulge

Transitioning to Tip Growth and Co-Expressing SPI-YPet and mRuby3-Lifeact.

Supplemental Movie 8. Time-Lapse Confocal Microscopy of a *spi* Root Hair

Expressing BRK1-YFP.

Supplemental Table 1. Statistical results for non-parametric Kolmogorov- Smirnov test

for data in Figure 7C on root hair length in wild-type, *spi,arp and spi arp*

Supplemental Table 2. Statistical results for non-parametric Kolmogorov- Smirnov test

for data in Figure 7D on root hair planar polarity in wild-type, *spi, arp and spi arp.*

ACKNOWLEDGMENTS

We thank Drs. Jose Alonso and Anna Stepanova (North Carolina State University) for

assistance with recombineering and Dr. Magdalena Bezanilla (Dartmouth College) for

the Lifeact-mRuby construct. This work was supported by the National Aeronautics and

Space Administration (NASA grants 80NSSC19K0129 and 80NSSC18K1462) and the

Noble Research Institute to E.B.B.) and by the National Science Foundation (NSF MCB

Grant No.1121893) to D.B.S. We also thank Dr. Larry M. York (Noble Research

Institute) and Dr. Wayne Versaw (Texas A.M. University) for critical comments on the

manuscript.

AUTHOR CONTRIBUTIONS

S.C., T.K, D.B.S and E.B.B. conceptualized and designed the research, and analyzed

and interpreted the data. S.C., J.A.S. and E.B.B. conducted root hair, primary root

growth and trichome assays, microscopy, image and statistical analysis of data. T.K.,

J.A.S., B.R.K., E.L.M. and D.B.S. generated various research reagents. All authors

contributed to writing the manuscript.

FIGURE LEGENDS

Figure 1. A Functional SPI-YPet Fusion Localizes to the Tips of Growing Root Hairs.

(A) SPI-YPet rescues the short root hair phenotype of *spi.* Bar = 100 μm.

(B) Low magnification image shows that SPI-YPet signal is most prominent at the tips of

rapidly expanding root hairs (arrows). SPI-YPet signal is weak during initiation/early

bulge formation and mature root hairs that have terminated tip growth (**). Bar = 50 μm

(C) High magnification images of single root hairs during bulge formation until tip growth

termination. SPI-YPet is enriched at the tip of root hairs that are rapidly growing or

transitioning to tip growth. Faint SPI-YPet signal is found in bulging root hairs or those

that have stopped elongating (**). Bar = 10 μm.

(D) Scatter plot showing correlation analysis of root hair tip SPI-YPet fluorescence and

root hair tip growth. The mean fluorescence in the oval in region 1 divided by the oval in

region 2 as shown in panel **B** represents the fluorescence ratio in the Y axis. Line

shows linear regression fit with R^2 value = 0.308 and p= 1.72 x 10^{-5}. (n=5-7 root hairs

per time point).

Figure 2. SPI-YPet is Localized to Brefeldin A (BFA) Sensitive Post-Golgi

Compartments.

(A) Brightfield and corresponding fluorescence image of an elongating root hair showing

the accumulation of SPI-YPet at the apical dome in solvent control-treated seedlings.

(B) Brightfield and corresponding fluorescence image of an elongating root hair 10 min

after treatment with 50 μM BFA. SPI-YPet at the root hair tip dissipates and forms

fluorescent agglomerates along the subapical regions (arrows).

(C) SPI-YPet signal is undetectable in the elongation zone of solvent control-treated

roots. Within 10 min of BFA application, distinct SPI-YPet fluorescent agglomerates

(arrows) form. Representative low (middle panel) and high (right panel) magnification

images of the elongation zone of BFA-treated roots are shown.

(D) and **(E)** The bulk secretory marker, SEC-RFP, accumulates at the tips of growing

wild-type root hairs (arrowhead), but is absent in *spi* root hairs (**).

(F) Box plot of SEC-RFP root hair tip accumulation expressed as fluorescence ratio.

Ratio values were obtained by dividing mean fluorescence in oval region in 1 over

region 2 (inset). Box limits indicate 25^{th} percentile and 75^{th} percentile, horizontal line is

the mean and whiskers display min and max values. Asterisk (***) indicates statistical

significance ($p<0.001$) as determined by Student's T-test. Bars = 10 μm (**A**, **B** and **D**);

bars in **(C)** = 50 μm. Means (n = 8-10 root hairs) ± SE

Figure 3. The Tip-focused F-actin Meshwork is Disrupted in Root Hairs of *spi*.

(A) to **(D)** Time course of F- actin organization in a wild-type root hair from bulge

formation to rapid tip growth. Weakly fluorescing F-actin networks (**) in the root hair

bulge **(A)** reorganize into prominent tip-focused meshworks as the root hair transitions

to rapid tip growth (arrows in **B** to **D**).

(E) Tip-focused F-actin meshworks (arrow) remain prominent in a long, rapidly

elongating wild-type root hair.

(F) Tip-focused F-actin meshworks in wild-type root hairs are replaced with F-actin

bundles (arrowheads) that protrude to the tip when growth stops.

(G) to **(J)** Time course of F- actin organization in a *spi* root hair bulge that is unable to

transition to tip growth. Distinct F-actin meshworks are unable to form in root hair bulges

that terminate tip growth (**, **I**). Thick F-actin bundles eventually form in these short,

non-growing root hair bulges (arrowhead, **J**).

(K) to **(N)** F-actin organization in slow-growing *spi* root hairs. Some *spi* root hairs show

the tip-focused F-actin meshworks typically observed in wild type (arrow, **K, L**).

However, tip-focused F-actin meshworks in slow-growing *spi* root hairs either dissipate

(**, **M**) or prematurely form thick F-actin bundles that protrude to the tip (arrow, **N**).

(O) Representative maximum projection images and corresponding computer-

generated cross-sections of wild-type and *spi* root hair tips. The fluorescence ratio of

the root hair tip (oval in 1) to background (oval in 2) was used to quantify tip-focused F-

actin meshworks.

(P) Box plot showing tip-focused F-actin fluorescence ratio. Box limits indicate 25[th]

percentile and 75[th] percentile, horizontal line is the mean and whiskers display min and

max values. Asterisk (***) indicates statistical significance ($p<0.001$) as determined by

Student's T-test. Means (n = 18-21 root hairs) ± S.E.

(Q) Elongating root hair simultaneously expressing SPI-YPet and mRuby-Lifeact. Note

that SPI-positive post-Golgi compartments and F-actin meshworks colocalize at the root

hair apex (arrows). Bars = 10 μm.

Figure 4. BRK1 and SCAR2 Mark the Root Hair Initiation Domain and Contribute to

Planar Polarity.

(A) Maximum projection confocal micrograph of the elongation and maturation zone of

an *Arabidopsis* primary root expressing a functional BRK1-YFP fusion. The image was

generated by merging 50 Z sections taken at 0.5 μm intervals. BRK1-YFP accumulates

in the basal end walls and root hair initiation domains (arrows). Bar = 50 μm.

(B) Time course of BRK1-YFP depletion in a developing root hair. BRK1-YFP signal

(arrows) is strongest prior to the formation of a root hair bulge (0 min) and gradually

dissipates as the root hair undergoes rapid tip growth (35-50 min **). Bar = 20 μm.

(C) A scatter plot showing an inverse relationship between BRK1-YFP signal and root

hair growth rate. Ratio of fluorescence of BRK1-YFP (a) to background (b) (inset) was

plotted against root hair growth rate. Line shows linear regression fit with R^2 value =

0.2667 and p= 1.131 x 10^{-7}. (n=5-7 root hairs per time point).

(D) Time course of SCAR2-mCherry in a developing root hair. Like BRK1-YFP, SCAR2-

mCherry signal is strongest at the RHID and early stages of root hair bulge formation (0

min). SCAR2-mCherry signal dissipates when the root hair undergoes rapid tip growth

(40 min **). Bar = 20 μm.

(E) Brightfield microscopy images of representative trichoblasts from 6-d-old seedlings

showing apical shift in root hair position of *brk1* compared to wild type, and

complementation of *brk1* planar polarity phenotypes by BRK1-YFP. Arrowheads mark

the end walls of the trichoblast and asterisks (***) mark the basal wall of the emerged

root hair.

(F) Violin plot of root hair planar polarity in wild type, *brk1, scar1234, BRK1-YFP in brk1*

and *SCAR2-mCherry* in *scar1234* genotypes. Relative root hair position was obtained

by taking the ratio of the distance from the basal trichoblast wall (bottom arrows in **E**) to

the basal root hair wall (***) over the total length of the trichoblasts (i.e. length between

the two arrows). The plot illustrates kernel probability density in which the width

represents distribution of data points. The black dot is the mean and whiskers display

min and max values. Statistical significance was determined using non-parametric, two

sample Kolmogorov-Smirnov (KS) test. Wild type versus *brk1* (***$p < 2.2 \times 10^{-16}$); *brk1*

versus BRK1-YFP in *brk1* (***$p=6.812 \times 10^{-18}$); Wild type versus BRK1-YFP ($p=0.202$

not significant, *ns*); Wild type versus *scar1234* (***$p < 2.2 \times 10^{-16}$); *brk1* versus *scar1234*

($p= 0.699$, *ns*); *scar1234* versus SCAR2-mCherry in *scar1234* (***$p < 2.2 \times 10^{-16}$); Wild

type versus *SCAR2-mCherry* in *scar1234* (*$p=0.012$). n = 90-117 root hairs ±S.E.

1004

Figure 5. Depletion of BRK1 as Root Hairs Transition to Tip Growth is Delayed in *spi*

(A) Dual imaging of BRK1-mRuby3 (arrow) and SPI-YPet shows that dissipation of

BRK1 (**) coincides with accumulation of SPI (arrow) as root hairs transition to rapid tip

growth.

(B) and **(C)** BRK1-YFP signal persists in *spi* root hair bulges (arrows in **B**) that fail to

transition to tip growth and in slow-growing *spi* root hairs (arrows in **C**). Bars = 20 μm.

(D) Accumulation of SPI-YPet at the root hair tip is not altered in *brk1*. Bar = 20 μm

1012

Figure 6. *SPI* Genetically Interacts with *BRK1* during Root Hair Development.

(A) Representative root hair images of wild type, *brk1*, *spi* and *spi brk1*. Note that *brk1*

exhibits wild type-like root hair morphology while *spi brk1* is more similar to *spi.* Bar =

200 μm

(B) and **(C)** Violin plots of root hair length **(B)** and root hair planar polarity **(C)** in wild

type, *brk1, spi* and *spi brk1* genotypes. The plots illustrate kernel probability density in

which the width represents proportion of data points. The black dot is the mean and

whiskers display min and max values. Statistical significance was determined using

non-parametric, two sample Kolmogorov-Smirnov (KS) test. For data in **(B)** wild type

versus *brk1* (*p= 0.00894); wild type versus *spi* (***p<2.2 x 10^{-16}); wild type versus *spi*

brk1 (***p<2.2 x 10^{-16}); *brk1* versus *spi* (***p<2.2 x 10^{-16}); *brk1* versus *spi brk1*

(***p<2.2 x 10^{-16}); *spi* versus *spi brk1* (p=0.2145, not significant, *ns*). For data in **(C)** wild

type versus *brk1* (***p= 9.703 x 10^{-11}); wild type versus *spi* (***p<2.2 x 10^{-16}); wild type

versus *spi brk1* (***p=6.661 x 10^{-16}); *brk1* versus *spi* (***p=1.564 x 10^{-7}); *brk1* versus

spi brk1 (***p=1.32 x 10^{-6}); *spi* versus *spi brk1* (p=0.9774, not significant, *ns*). Means (n

= 83-110 root hairs) ± S.E.

1029

Figure 7. *SPI* and *ARP2/3* Genetically Interact during Root Hair Development

(A) Representative root hair images of wild type, *arp3, spi* and *spi arp3*. Note that *arp3*

exhibits wild type-like root hair morphology. Unlike *spi, spi arp3* appears morphologically

hairless.

(B) Representative high magnification root hair images of *spi* and *spi arp3*. Note that *spi*

arp3 appear morphologically hairless because of a high number of aborted root hair

bulges (arrows). The position in which *spi arp3* root hair bulges emerge is closer to the

basal end walls (***) when compared with *spi*. Bar = 20 μm

(C) and **(D)** Violin plots of root hair planar polarity in wild-type and *arp* mutants. The plot

illustrates kernel probability density in which the width represents distribution of data

points. The black dot is the mean and whiskers display min and max values. Statistical

significance was determined using non-parametric Kolmogorov-Smirnov (KS) test.

Different letters are significantly different based. P values for pair wise comparisons

among genotypes are presented in Supplemental Table 1.

1044

Figure 8. Hypothetical Model Illustrating SPI- and W/SCR-ARP2/3- Modulated Root

Hair Development. Based on their common location at the root hair initiation domain

(RHID), ROP/RAC-GTPase and ROP-GEF3 activate the W/SCR complex. This in turn

activates a pool of ARP2/3 in the RHID to ensure the correct position of root hair

emergence. Genetic interaction studies point to SPI and W/SCR-ARP2/3 functioning in

parallel pathways during root hair initiation and early root hair bulge formation. SPI,

through a yet to be determined mechanism, is recruited to the tips of root hair

outgrowths via post-Golgi vesicles. SPI enrichment at the root hair tip then negatively

regulates the W/SCR-ARP2/3- actin nucleation pathway as the root bulge enlarges by

displacing or degrading BRK1. The antagonistic effect of SPI on BRK1 leads to the

depletion of the active pool ARP2/3. Actin-mediated rapid root hair tip growth then

commences predominantly through a SPI-dependent mechanism.

1057

FIGURES –Chin et al

Figure 1. A Functional SPI-YPet Fusion Localizes to the Tips of Growing Root Hairs.
(A) SPI-YPet rescues the short root hair phenotype of *spi*. Bar = 100 μm.
(B) Low magnification image shows that SPI-YPet signal is most prominent at the tips of rapidly expanding root hairs (arrows). SPI-YPet signal is weak during initiation/early bulge formation and mature root hairs that have terminated tip growth (**). Bar = 50 μm
(C) High magnification images of single root hairs during bulge formation until tip growth termination. SPI-YPet is enriched at the tip of root hairs that are rapidly growing or transitioning to tip growth. Faint SPI-YPet signal is found in bulging root hairs or those that have stopped elongating (**). Bar = 10 μm.
(D) Scatter plot showing correlation analysis of root hair tip SPI-YPet fluorescence and root hair tip growth. The mean fluorescence in the oval in region 1 divided by the oval in region 2 as shown in panel **B** represents the fluorescence ratio in the Y axis. Line shows linear regression fit with R^2 value = 0.308 and p= 1.72 x 10^{-5}. (n=5-7 root hairs per time point).

Figure 2. SPI-YPet is Localized to Brefeldin A (BFA) Sensitive Post-Golgi Compartments.

(A) Brightfield and corresponding fluorescence image of an elongating root hair showing the accumulation of SPI-YPet at the apical dome in solvent control-treated seedlings.

(B) Brightfield and corresponding fluorescence image of an elongating root hair 10 min after treatment with 50 μM BFA. SPI-YPet at the root hair tip dissipates and forms fluorescent agglomerates along the subapical regions (arrows).

(C) SPI-YPet signal is undetectable in the elongation zone of solvent control-treated roots. Within 10 min of BFA application, distinct SPI-YPet fluorescent agglomerates (arrows) form. Representative low (middle panel) and high (right panel) magnification images of the elongation zone of BFA-treated roots are shown.

(D) and **(E)** The bulk secretory marker, SEC-RFP, accumulates at the tips of growing wild-type root hairs (arrowhead), but is absent in *spi* root hairs (**).

(F) Box plot of SEC-RFP root hair tip accumulation expressed as fluorescence ratio. Ratio values were obtained by dividing mean fluorescence in oval region in 1 over region 2 (inset). Box limits indicate 25^{th} percentile and 75^{th} percentile, horizontal line is the mean and whiskers display min and max values. Asterisk (***) indicates statistical significance ($p<0.001$) as determined by Student's T-test. Bars = 10 μm (**A**, **B** and **D**); bars in **(C)** = 50 μm. Means (n = 8-10 root hairs) ± SE

Figure 3. The Tip-focused F-actin Meshwork is Disrupted in Root Hairs of *spi*.
(A) to **(D)** Time course of F- actin organization in a wild-type root hair from bulge formation to rapid tip growth. Weakly fluorescing F-actin networks (**) in the root hair bulge **(A)** reorganize into prominent tip-focused meshworks as the root hair transitions to rapid tip growth (arrows in **B** to **D**).
(E) Tip-focused F-actin meshworks (arrow) remain prominent in a long, rapidly elongating wild-type root hair.
(F) Tip-focused F-actin meshworks in wild-type root hairs are replaced with F-actin bundles (arrowheads) that protrude to the tip when growth stops.
(G) to **(J)** Time course of F- actin organization in a *spi* root hair bulge that is unable to transition to tip growth. Distinct F-actin meshworks are unable to form in root hair bulges that terminate tip growth (**, **I**). Thick F-actin bundles eventually form in these short, non-growing root hair bulges (arrowhead, **J**).

(K) to **(N)** F-actin organization in slow-growing *spi* root hairs. Some *spi*root hairs show the tip-focused F-actin meshworks typically observed in wild type (arrow, **K, L**). However, tip-focused F-actin meshworks in slow-growing *spi* root hairs either dissipate (******, **M**) or prematurely form thick F-actin bundles that protrude to the tip (arrow, **N**).

(O) Representative maximum projection images and corresponding computer-generated cross-sections of wild-type and *spi* root hair tips. The fluorescence ratio of the root hair tip (oval in 1) to background (oval in 2) was used to quantify tip-focused F-actin meshworks.

(P) Box plot showing tip-focused F-actin fluorescence ratio. Box limits indicate 25^{th} percentile and 75^{th} percentile, horizontal line is the mean and whiskers display min and max values. Asterisk (*******) indicates statistical significance ($p<0.001$) as determined by Student's T-test. Means (n = 18-21 root hairs) ± S.E.

(Q) Elongating root hair simultaneously expressing SPI-YPet and mRuby-Lifeact. Note that SPI-positive post-Golgi compartments and F-actin meshworks colocalize at the root hair apex (arrows). Bars = 10 μm.

Figure 4. BRK1 and SCAR2 Mark the Root Hair Initiation Domain and Contribute to Planar Polarity.

(A) Maximum projection confocal micrograph of the elongation and maturation zone of an *Arabidopsis* primary root expressing a functional BRK1-YFP fusion. The image was generated by merging 50 Z sections taken at 0.5 μm intervals. BRK1-YFP accumulates in the basal end walls and root hair initiation domains (arrows). Bar = 50 μm.

(B) Time course of BRK1-YFP depletion in a developing root hair. BRK1-YFP signal (arrows) is strongest prior to the formation of a root hair bulge (0 min) and gradually dissipates as the root hair undergoes rapid tip growth (35-50 min **). Bar = 20 μm.
(C) A scatter plot showing an inverse relationship between BRK1-YFP signal and root hair growth rate. Ratio of fluorescence of BRK1-YFP (a) to background (b) (inset) was plotted against root hair growth rate. Line shows linear regression fit with R^2 value = 0.2667 and p= 1.131 x 10^{-7}. (n=5-7 root hairs per time point).
(D) Time course of SCAR2-mCherry in a developing root hair. Like BRK1-YFP, SCAR2-mCherry signal is strongest at the RHID and early stages of root hair bulge formation (0 min). SCAR2-mCherry signal dissipates when the root hair undergoes rapid tip growth (40 min **). Bar = 20 μm.
(E) Brightfield microscopy images of representative trichoblasts from 6-d-old seedlings showing apical shift in root hair position of *brk1* compared to wild type, and complementation of *brk1* planar polarity phenotypes by BRK1-YFP. Arrowheads mark the end walls of the trichoblast and asterisks (***) mark the basal wall of the emerged root hair.
(F) Violin plot of root hair planar polarity in wild type, *brk1, scar1234, BRK1-YFP in brk1* and *SCAR2-mCherry* in *scar1234* genotypes. Relative root hair position was obtained by taking the ratio of the distance from the basal trichoblast wall (bottom arrows in **E**) to the basal root hair wall (***) over the total length of the trichoblasts (i.e. length between the two arrows). The plot illustrates kernel probability density in which the width represents distribution of data points. The black dot is the mean and whiskers display min and max values. Statistical significance was determined using non-parametric, two sample Kolmogorov-Smirnov (KS) test. Wild type versus *brk1* (***p < 2.2 x 10^{-16}); *brk1* versus BRK1-YFP in *brk1* (***p=6.812 x 10^{-18}); Wild type versus BRK1-YFP (p=0.202 not significant, *ns*); Wild type versus *scar1234* (***p < 2.2 x 10^{-16}); *brk1* versus *scar1234* (p= 0.699, *ns*); *scar1234* versus SCAR2-mCherry in *scar1234* (***p < 2.2 x 10^{-16}); Wild type versus *SCAR2-mCherry* in *scar1234* (*p=0.012). n = 90-117 root hairs ±S.E.

Figure 5. Depletion of BRK1 as Root Hairs Transition to Tip Growth is Delayed in *spi*
(A) Dual imaging of BRK1-mRuby3 (arrow) and SPI-YPet shows that dissipation of BRK1 (**) coincides with accumulation of SPI (arrow) as root hairs transition to rapid tip growth.
(B) and **(C)** BRK1-YFP signal persists in *spi* root hair bulges (arrows in **B**) that fail to transition to tip growth and in slow-growing *spi* root hairs (arrows in **C**). Bars = 20 μm.
(D) Accumulation of SPI-YPet at the root hair tip is not altered in *brk1*. Bar = 20 μm

Figure 6. *SPI* Genetically Interacts with *BRK1* during Root Hair Development.
(A) Representative root hair images of wild type, *brk1*, *spi* and *spi brk1*. Note that *brk1* exhibits wild type-like root hair morphology while *spi brk1* is more similar to *spi.* Bar = 200 μm
(B) and **(C)** Violin plots of root hair length **(B)** and root hair planar polarity **(C)** in wild type, *brk1, spi* and *spi brk1* genotypes. The plots illustrate kernel probability density in which the width represents proportion of data points. The black dot is the mean and whiskers display min and max values. Statistical significance was determined using non-parametric, two sample Kolmogorov-Smirnov (KS) test. For data in **(B)** wild type versus *brk1* (*p= 0.00894); wild type versus *spi* (***p<2.2 x 10^{-16}); wild type versus *spi brk1* (***p<2.2 x 10^{-16}); *brk1* versus *spi* (***p<2.2 x 10^{-16}); *brk1* versus *spi brk1*

(***$p<2.2 \times 10^{-16}$); *spi* versus *spi brk1* (p=0.2145, not significant, *ns*). For data in **(C)** wild type versus *brk1* (***$p= 9.703 \times 10^{-11}$); wild type versus *spi* (***$p<2.2 \times 10^{-16}$); wild type versus *spi brk1* (***$p=6.661 \times 10^{-16}$); *brk1* versus *spi* (***$p=1.564 \times 10^{-7}$); *brk1* versus *spi brk1* (***$p=1.32 \times 10^{-6}$); *spi* versus *spi brk1* (p=0.9774, not significant, *ns*). Means (n = 83-110 root hairs) ± S.E.

Figure 7. *SPI* and *ARP2/3* Genetically Interact during Root Hair Development
(A) Representative root hair images of wild type, *arp3, spi* and *spi arp3*. Note that *arp3* exhibits wild type-like root hair morphology. Unlike *spi*, *spi arp3* appears morphologically hairless.
(B) Representative high magnification root hair images of *spi* and *spi arp3*. Note that *spi arp3* appear morphologically hairless because of a high number of aborted root hair bulges (arrows). The position in which *spi arp3* root hair bulges emerge is closer to the basal end walls (***) when compared with *spi*. Bar = 20 μm
(C) and **(D)** Violin plots of root hair planar polarity in wild-type and *arp* mutants. The plot illustrates kernel probability density in which the width represents distribution of data

points. The black dot is the mean and whiskers display min and max values. Statistical significance was determined using non-parametric Kolmogorov-Smirnov (KS) test. Different letters are significantly different based. P values for pair wise comparisons among genotypes are presented in Supplemental Table 1.

Figure 8. Hypothetical Model Illustrating SPI- and W/SCR-ARP2/3- Modulated Root Hair Development. Based on their common location at the root hair initiation domain (RHID), ROP/RAC-GTPase and ROP-GEF3 activate the W/SCR complex. This in turn activates a pool of ARP2/3 in the RHID to ensure the correct position of root hair emergence. Genetic interaction studies point to SPI and W/SCR-ARP2/3 functioning in parallel pathways during root hair initiation and early root hair bulge formation. SPI, through a yet to be determined mechanism, is recruited to the tips of root hair outgrowths via post-Golgi vesicles. SPI enrichment at the root hair tip then negatively regulates the W/SCR-ARP2/3- actin nucleation pathway as the root bulge enlarges by displacing or degrading BRK1. The antagonistic effect of SPI on BRK1 leads to the depletion of the active pool ARP2/3. Actin-mediated rapid root hair tip growth then commences predominantly through a SPI-dependent mechanism.

Parsed Citations

ALONSO, J. M., STEPANOVA, A. N., LEISSE, T. J., KIM, C. J., CHEN, H., SHINN, P., STEVENSON, D. K., ZIMMERMAN, J., BARAJAS, P., CHEUK, R., GADRINAB, C., HELLER, C., JESKE, A, KOESEMA, E., MEYERS, C. C., PARKER, H., PREDNIS, L., ANSARI, Y., CHOY, N., DEEN, H., GERALT, M., HAZARI, N., HOM, E., KARNES, M., MULHOLLAND, C., NDUBAKU, R., SCHMIDT, I., GUZMAN, P., AGUILAR-HENONIN, L., SCHMID, M., WEIGEL, D., CARTER, D. E., MARCHAND, T., RISSEEUW, E., BROGDEN, D., ZEKO, A, CROSBY, W. L., BERRY, C. C. & ECKER, J. R. 2003. Genome-Wide Insertional Mutagenesis of Arabidopsis thaliana. Science, 301, 653.

Google Scholar: Author Only Title Only Author and Title

BADET, T., LEGER, O., BARASCUD, M., VOISIN, D., SADON, P., VINCENT, R., LE RU, A, BALAGUE, C., ROBY, D. & RAFFAELE, S. 2019. Expression polymorphism at the ARPC4 locus links the actin cytoskeleton with quantitative disease resistance to Sclerotinia sclerotiorum in Arabidopsis thaliana. New Phytol, 222, 480-496.

Google Scholar: Author Only Title Only Author and Title

BALUŠKA, F. 2002. F-Actin-Dependent Endocytosis of Cell Wall Pectins in Meristematic Root Cells. Insights from Brefeldin A-Induced Compartments. Plant Physiology, 130, 422-431.

Google Scholar: Author Only Title Only Author and Title

BALUŠKA, F. 2010. Recent surprising similarities between plant cells and neurons. Plant signaling & behavior, 5, 87-89.

Google Scholar: Author Only Title Only Author and Title

BASCOM, C. S., HEPLER, P. K. & BEZANILLA, M. 2018a. Interplay between Ions, the Cytoskeleton, and Cell Wall Properties during Tip Growth. Plant Physiology, 176, 28-40.

Google Scholar: Author Only Title Only Author and Title

BASCOM, C. S., WINSHIP, L. J. & BEZANILLA, M. 2018b. Simultaneous imaging and functional studies reveal a tight correlation between calcium and actin networks. Proceedings of the National Academy of Sciences, 115, E2869-E2878.

Google Scholar: Author Only Title Only Author and Title

BASU, D., EL-ASSAL SEL, D., LE, J., MALLERY, E. L. & SZYMANSKI, D. B. 2004. Interchangeable functions of Arabidopsis PIROGI and the human WAVE complex subunit SRA1 during leaf epidermal development. Development, 131, 4345-55.

Google Scholar: Author Only Title Only Author and Title

BASU, D., LE, J., EL-ESSAL SEL, D., HUANG, S., ZHANG, C., MALLERY, E. L., KOLIANTZ, G., STAIGER, C. J. & SZYMANSKI, D. B. 2005. DISTORTED3/SCAR2 is a putative arabidopsis WAVE complex subunit that activates the Arp2/3 complex and is required for epidermal morphogenesis. Plant Cell, 17, 502-24.

Google Scholar: Author Only Title Only Author and Title

BASU, D., LE, J., ZAKHAROVA, T., MALLERY, E. L. & SZYMANSKI, D. B. 2008. A SPIKE1 signaling complex controls actin-dependent cell morphogenesis through the heteromeric WAVE and ARP2/3 complexes. Proceedings of the National Academy of Sciences, 105, 4044-4049.

Google Scholar: Author Only Title Only Author and Title

BATES, D., MAECHLER, M., BOLKER, B. & WALKER, S. 2015. Fitting Linear Mixed-Effects Models using lme4. Journal of Statistical Software, 67, 1-48.

Google Scholar: Author Only Title Only Author and Title

BIBIKOVA, T. N., BLANCAFLOR, E. B. & GILROY, S. 1999. Microtubules regulate tip growth and orientation in root hairs of Arabidopsis thaliana. The Plant Journal, 17, 657-665.

Google Scholar: Author Only Title Only Author and Title

BLANCHOIN, L., AMANN, K. J., HIGGS, H. N., MARCHAND, J.-B., KAISER, D. A & POLLARD, T. D. 2000. Direct observation of dendritic actin filament networks nucleated by Arp2/3 complex and WASP/Scar proteins. Nature, 404, 1007-1011.

Google Scholar: Author Only Title Only Author and Title

BRUMOS, J., ZHAO, C., GONG, Y., SORIANO, D., PATEL, A. P., PEREZ-AMADOR, M. A, STEPANOVA, A. N. & ALONSO, J. M. 2020. An Improved Recombineering Toolset for Plants. The Plant Cell, 32, 100-122.

Google Scholar: Author Only Title Only Author and Title

CARMINATI, A, PASSIOURA, J. B., ZAREBANADKOUKI, M., AHMED, M. A, RYAN, P. R., WATT, M. & DELHAIZE, E. 2017. Root hairs enable high transpiration rates in drying soils. New Phytol, 216, 771-781.

Google Scholar: Author Only Title Only Author and Title

CINGOLANI, L. A & GODA, Y. 2008. Actin in action: the interplay between the actin cytoskeleton and synaptic efficacy. Nature Reviews Neuroscience, 9, 344-356.

Google Scholar: Author Only Title Only Author and Title

CLOUGH, S. J. & BENT, A. F. 1998. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. The Plant Journal, 16, 735-743.

Google Scholar: Author Only Title Only Author and Title

CULLINANE, A. R., SCHAFFER, A. A. & HUIZING, M. 2013. The BEACH is hot: a LYST of emerging roles for BEACH-domain containing

proteins in human disease. Traffic, 14, 749-66.
Google Scholar: Author Only Title Only Author and Title

DE MENDIBURU, F. 2020. agricolae: Statistical Procedures for Agricultural Rsearch.

DE SOUZA, N., VALLIER, L. G., FARES, H. & GREENWALD, I. 2007. SEL-2, the C. elegans neurobeachin/LRBA homolog, is a negative regulator of lin-12/Notch activity and affects endosomal traffic in polarized epithelial cells. Development, 134, 691-702.
Google Scholar: Author Only Title Only Author and Title

DEEKS, M. J. & HUSSEY, P. J. 2005. Arp2/3 and SCAR: plants move to the fore. Nat Rev Mol Cell Biol, 6, 954-64.
Google Scholar: Author Only Title Only Author and Title

DEEKS, M. J., KALORITI, D., DAVIES, B., MALHO, R. & HUSSEY, P. J. 2004. Arabidopsis NAP1 is essential for Arp2/3-dependent trichome morphogenesis. Curr Biol, 14, 1410-4.
Google Scholar: Author Only Title Only Author and Title

DENNINGER, P., REICHELT, A, SCHMIDT, V. A. F., MEHLHORN, D. G., ASSECK, L. Y., STANLEY, C. E., KEINATH, N. F., EVERS, J.-F., GREFEN, C. & GROSSMANN, G. 2019. Distinct RopGEFs Successively Drive Polarization and Outgrowth of Root Hairs. Current Biology, 29, 1854-1865.
Google Scholar: Author Only Title Only Author and Title

DJAKOVIC, S. 2006. BRICK1/HSPC300 functions with SCAR and the ARP2/3 complex to regulate epidermal cell shape in Arabidopsis. Development, 133, 1091-1100.
Google Scholar: Author Only Title Only Author and Title

DOYLE, S. M., VAIN, T. & ROBERT, S. 2015. Small molecules unravel complex interplay between auxin biology and endomembrane trafficking. Journal of Experimental Biology, 66, 4971-4982.
Google Scholar: Author Only Title Only Author and Title

DYACHOK, J., PAEZ-GARCIA, A, YOO, C. M., PALANICHELVAM, K. & BLANCAFLOR, E. B. 2016. Fluorescence Imaging of the Cytoskeleton in Plant Roots. In: GAVIN, R. H. (ed.) Cytoskeleton Methods and Protocols: Methods in Molecular Biology series. 3rd ed. Totowa, New Jersey: Humana Press.
Google Scholar: Author Only Title Only Author and Title

DYACHOK, J., SHAO, M.-R., VAUGHN, K., BOWLING, A, FACETTE, M., DJAKOVIC, S., CLARK, L. & SMITH, L. 2008. Plasma Membrane-Associated SCAR Complex Subunits Promote Cortical F-Actin Accumulation and Normal Growth Characteristics in Arabidopsis Roots. Molecular Plant, 1, 990-1006.
Google Scholar: Author Only Title Only Author and Title

DYACHOK, J., ZHU, L., LIAO, F., HE, J., HUQ, E. & BLANCAFLOR, E. B. 2011. SCAR Mediates Light-Induced Root Elongation in Arabidopsis through Photoreceptors and Proteasomes. The Plant Cell, 23, 3610-3626.
Google Scholar: Author Only Title Only Author and Title

EDWARDS, K., JOHNSTONE, C. & THOMPSON, C. 1991. A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. Nucleic acids research, 19, 1349-1349.
Google Scholar: Author Only Title Only Author and Title

EL-ASSAL SEL, D., LE, J., BASU, D., MALLERY, E. L. & SZYMANSKI, D. B. 2004. Arabidopsis GNARLED encodes a NAP125 homolog that positively regulates ARP2/3. Curr Biol, 14, 1405-9.
Google Scholar: Author Only Title Only Author and Title

FASO, C., CHEN, Y. N., TAMURA, K., HELD, M., ZEMELIS, S., MARTI, L., SARAVANAN, R., HUMMEL, E., KUNG, L., MILLER, E., HAWES, C. & BRANDIZZI, F. 2009. A Missense Mutation in the Arabidopsis COPII Coat Protein Sec24A Induces the Formation of Clusters of the Endoplasmic Reticulum and Golgi Apparatus. The Plant Cell, 21, 3655-3671.
Google Scholar: Author Only Title Only Author and Title

FINKA, A, SAIDI, Y., GOLOUBINOFF, P., NEUHAUS, J. M., ZRYD, J. P. & SCHAEFER, D. G. 2008. The knock-out of ARP3a gene affects F-actin cytoskeleton organization altering cellular tip growth, morphology and development in moss Physcomitrella patens. Cell Motil Cytoskeleton, 65, 769-84.
Google Scholar: Author Only Title Only Author and Title

FISCHER, U., IKEDA, Y., LJUNG, K., SERRALBO, O., SINGH, M., HEIDSTRA, R., PALME, K., SCHERES, B. & GREBE, M. 2006. Vectorial Information for Arabidopsis Planar Polarity Is Mediated by Combined AUX1, EIN2, and GNOM Activity. Current Biology, 16, 2143-2149.
Google Scholar: Author Only Title Only Author and Title

FOX, J. & WEISBERG, S. 2019. An {R} Companion to Applied Regression, Third Edition. Thousand Oaks, CA: Sage.
Google Scholar: Author Only Title Only Author and Title

FRANK, M. J. & SMITH, L. G. 2002. A Small, Novel Protein Highly Conserved in Plants and Animals Promotes the Polarized Growth and Division of Maize Leaf Epidermal Cells. Current Biology, 12, 849-853.
Google Scholar: Author Only Title Only Author and Title

GIBBON, B. C., KOVAR, D. R. & STAIGER, C. J. 1999. Latrunculin B Has Different Effects on Pollen Germination and Tube Growth. The Plant Cell, 11, 2349-2364.

Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

GRIERSON, C., NIELSEN, E., KETELAAR, T. & SCHIEFELBEIN, J. 2014. Root Hairs. The Arabidopsis Book, 12.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

HARRIES, P. A, PAN, A. & QUATRANO, R. S. 2005. Actin-related protein2/3 complex component ARPC1 is required for proper cell morphogenesis and polarized cell growth in Physcomitrella patens. Plant Cell, 17, 2327-39.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

HAUBERT, D., GHARIB, N., RIVERO, F., WIEGMANN, K., HÖSEL, M., KRÖNKE, M. & KASHKAR, H. 2007. PtdIns(4,5)P-restricted plasma membrane localization of FAN is involved in TNF-induced actin reorganization. The EMBO journal, 26, 3308-3321.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

HONKANEN, S., JONES, V. A. S., MORIERI, G., CHAMPION, C., HETHERINGTON, A. J., KELLY, S., PROUST, H., SAINT-MARCOUX, D., PRESCOTT, H. & DOLAN, L. 2016. The Mechanism Forming the Cell Surface of Tip-Growing Rooting Cells Is Conserved among Land Plants. Current Biology, 26, 3238-3244.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

HÜLSKAMP, M., MISÉRA, S. & JÜRGENS, G. 1994. Genetic dissection of trichome cell development in Arabidopsis. Cell, 76, 555-566.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

ISNER, J. C., XU, Z, COSTA, J. M., MONNET, F., BATSTONE, T., OU, X., DEEKS, M. J., GENTY, B., JIANG, K. & HETHERINGTON, A. M. 2017. Actin filament reorganisation controlled by the SCAR/WAVE complex mediates stomatal response to darkness. New Phytol, 215, 1059-1067.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

JONES, M. A, SHEN, J.-J., FU, Y., LI, H., YANG, Z & GRIERSON, C. S. 2002. The Arabidopsis Rop2 GTPase Is a Positive Regulator of Both Root Hair Initiation and Tip Growth. The Plant Cell, 14, 763-776.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

JÖRGENS, C. I., GRÜNEWALD, N., HÜLSKAMP, M. & UHRIG, J. F. 2010. A role for ABIL3 in plant cell morphogenesis. The Plant Journal, 62, 925-935.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

KIEFER, C. S., CLAES, A R., NZAYISENGA, J. C., PIETRA, S., STANISLAS, T., HUSER, A, IKEDA, Y. & GREBE, M. 2015. Arabidopsis AIP1-2 restricted by WER-mediated patterning modulates planar polarity. Development, 142, 151-161.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

KOTCHONI, S. O., ZAKHAROVA, T., MALLERY, E. L., LE, J., EL-ASSAL, S. E.-D. & SZYMANSKI, D. B. 2009. The Association of the Arabidopsis Actin-Related Protein2/3 Complex with Cell Membranes Is Linked to Its Assembly Status But Not Its Activation. Plant Physiology, 151, 2095-2109.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

LE, J., EL-ASSAL SEL, D., BASU, D., SAAD, M. E. & SZYMANSKI, D. B. 2003. Requirements for Arabidopsis ATARP2 and ATARP3 during epidermal development. Curr Biol, 13, 1341-7.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

LE, J., MALLERY, E. L., ZHANG, C., BRANKLE, S. & SZYMANSKI, D. B. 2006. Arabidopsis BRICK1/HSPC300 Is an Essential WAVE-Complex Subunit that Selectively Stabilizes the Arp2/3 Activator SCAR2. Current Biology, 16, 895-901.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

LI, J., BLANCHOIN, L. & STAIGER, C. J. 2015. Signaling to actin stochastic dynamics. Annu Rev Plant Biol, 66, 415-40.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

LI, S., BLANCHOIN, L., YANG, Z & LORD, E. M. 2003. The putative Arabidopsis arp2/3 complex controls leaf cell morphogenesis. Plant Physiol, 132, 2034-44.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

LI, X., LI, J. H., WANG, W., CHEN, N. Z, MA, T. S., XI, Y. N., ZHANG, X. L., LIN, H. F., BAI, Y., HUANG, S. J. & CHEN, Y. L. 2014. ARP2/3 complex-mediated actin dynamics is required for hydrogen peroxide-induced stomatal closure in Arabidopsis. Plant Cell Environ, 37, 1548-60.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

LIU, X., LI, Y., WANG, X., XING, R., LIU, K., GAN, Q., TANG, C., GAO, Z, JIAN, Y., LUO, S., GUO, W. & YANG, C. 2017. The BEACH-containing protein WDR81 coordinates p62 and LC3C to promote aggrephagy. Journal of Cell Biology, 216, 1301-1320.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

LUKOWITZ, W., GILLMOR, C. S. & SCHEIBLE, W.-R. 2000. Positional Cloning in Arabidopsis. Why It Feels Good to Have a Genome Initiative Working for You. Plant Physiology, 123, 795-806.
Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

MACHESKY, L. M., ATKINSON, S. J., AMPE, C., VANDEKERCKHOVE, J. & POLLARD, T. D. 1994. Purification of a cortical complex containing two unconventional actins from Acanthamoeba by affinity chromatography on profilin-agarose. The Journal of Cell Biology,

127, 107-115.

Google Scholar: Author Only Title Only Author and Title

MATHUR, J., MATHUR, N., KERNEBECK, B. & HÜLSKAMP, M. 2003. Mutations in actin-related proteins 2 and 3 affect cell shape development in Arabidopsis. The Plant Cell, 15, 1632-1645.

Google Scholar: Author Only Title Only Author and Title

MATTILA, P. K. & LAPPALAINEN, P. 2008. Filopodia: molecular architecture and cellular functions. Nature Reviews Molecular Cell Biology, 9, 446-454.

Google Scholar: Author Only Title Only Author and Title

MOLENDIJK, A. J., BISCHOFF, F., RAJENDRAKUMAR, C. S., FRIML, J., BRAUN, M., GILROY, S. & PALME, K. 2001. Arabidopsis thaliana Rop GTPases are localized to tips of root hairs and control polar growth. The EMBO journal, 20, 2779-2788.

Google Scholar: Author Only Title Only Author and Title

NAKAMURA, M. & GREBE, M. 2018. Outer, inner and planar polarity in the Arabidopsis root. Current Opinion in Plant Biology, 41, 46-53.

Google Scholar: Author Only Title Only Author and Title

NGUYEN, A. W. & DAUGHERTY, P. S. 2005. Evolutionary optimization of fluorescent proteins for intracellular FRET. Nature Biotechnology, 23, 355-360.

Google Scholar: Author Only Title Only Author and Title

NIESMANN, K., BREUER, D., BROCKHAUS, J., BORN, G., WOLFF, I., REISSNER, C., KILIMANN, M. W., ROHLMANN, A. & MISSLER, M. 2011. Dendritic spine formation and synaptic function require neurobeachin. Nature Communications, 2, 557.

Google Scholar: Author Only Title Only Author and Title

PAEZ-GARCIA, A., SPARKS, J.A., DE BANG, L., BLANCAFLOR, E.B. 2018. Plant actin cytoskeleton: New functions from old scaffold. In: SAHI, V., BALUŠKA, F. (ed.) Concepts in Cell Biology - History and Evolution. Springer.

Google Scholar: Author Only Title Only Author and Title

PÉREZ-PÉREZ, J. M., CANDELA, H. & MICOL, J. L. 2009. Understanding synergy in genetic interactions. Trends in Genetics, 25, 368-376.

Google Scholar: Author Only Title Only Author and Title

PERROUD, P. F. & QUATRANO, R. S. 2006. The role of ARPC4 in tip growth and alignment of the polar axis in filaments of Physcomitrella patens. Cell Motil Cytoskeleton, 63, 162-71.

Google Scholar: Author Only Title Only Author and Title

PERROUD, P. F. & QUATRANO, R. S. 2008. BRICK1 is required for apical cell growth in filaments of the moss Physcomitrella patens but not for gametophore morphology. Plant Cell, 20, 411-22.

Google Scholar: Author Only Title Only Author and Title

PRATAP SAHI, V., CIFROVA, P., GARCIA-GONZALEZ, J., KOTANNAL BABY, I., MOUILLE, G., GINEAU, E., MULLER, K., BALUSKA, F., SOUKUP, A, PETRASEK, J. & SCHWARZEROVA, K. 2018. Arabidopsis thaliana plants lacking the ARP2/3 complex show defects in cell wall assembly and auxin distribution. Ann Bot, 122, 777-789.

Google Scholar: Author Only Title Only Author and Title

PREUSS, M. L., SERNA, J., FALBEL, T. G., BEDNAREK, S. Y. & NIELSEN, E. 2004. The Arabidopsis Rab GTPase RabA4b Localizes to the Tips of Growing Root Hair Cells. The Plant Cell, 16, 1589-1603.

Google Scholar: Author Only Title Only Author and Title

RINGLI, C., BAUMBERGER, N., DIET, A, FREY, B. & KELLER, B. 2002. ACTIN2 is essential for bulge site selection and tip growth during root hair development of Arabidopsis. Plant Physiol, 129, 1464-72.

Google Scholar: Author Only Title Only Author and Title

ROBINSON, R. C., TURBEDSKY, K., KAISER, D. A, MARCHAND, J.-B., HIGGS, H. N., CHOE, S. & POLLARD, T. D. 2001. Crystal Structure of Arp2/3 Complex. Science, 294, 1679-1684.

Google Scholar: Author Only Title Only Author and Title

RODAL, A. A, SOKOLOVA, O., ROBINS, D. B., DAUGHERTY, K. M., HIPPENMEYER, S., RIEZMAN, H., GRIGORIEFF, N. & GOODE, B. L. 2005. Conformational changes in the Arp2/3 complex leading to actin nucleation. Nature Structural and Molecular Biology, 12, 26-31.

Google Scholar: Author Only Title Only Author and Title

RUIZ, S., KOEBERNICK, N., DUNCAN, S., FLETCHER, D. M., SCOTSON, C., BOGHI, A, MARIN, M., BENGOUGH, A. G., GEORGE, T. S., BROWN, L. K., HALLETT, P. D. & ROOSE, T. 2020. Significance of root hairs at the field scale - modelling root water and phosphorus uptake under different field conditions. Plant Soil, 447, 281-304.

Google Scholar: Author Only Title Only Author and Title

SAEDLER, R., JAKOBY, M., MARIN, B., GALIANA-JAIME, E. & HÜLSKAMP, M. 2009. The cell morphogenesis gene SPIRRIG in Arabidopsis encodes a WD/BEACH domain protein. The Plant Journal, 59, 612-621.

Google Scholar: Author Only Title Only Author and Title

SAEDLER, R., MATHUR, N., SRINIVAS, B. P., KERNEBECK, B., HÜLSKAMP, M. & MATHUR, J. 2004. Actin Control Over Microtubules Suggested by DISTORTED2 Encoding the Arabidopsis ARPC2 Subunit Homolog. Plant and Cell Physiology, 45, 813-822.

Google Scholar: Author Only Title Only Author and Title

SCHIEFELBEIN, J., HUANG, L. & ZHENG, X. 2014. Regulation of epidermal cell fate in Arabidopsis roots: the importance of multiple feedback loops. Front Plant Sci, 5, 47.
Google Scholar: Author Only Title Only Author and Title

SCHWAB, B., MATHUR, J., SAEDLER, R. R., SCHWARZ, H., FREY, B., SCHEIDEGGER, C. & HÜLSKAMP, M. 2003. Regulation of cell expansion by the DISTORTED genes in Arabidopsis thaliana: actin controls the spatial organization of microtubules. Molecular Genetics and Genomics, 269, 350-360.
Google Scholar: Author Only Title Only Author and Title

SHARAN, S. K., THOMASON, L. C., KUZNETSOV, S. G. & COURT, D. L. 2009. Recombineering: a homologous recombination-based method of genetic engineering. Nature Protocols, 4, 206-223.
Google Scholar: Author Only Title Only Author and Title

SHIBATA, M., BREUER, C., KAWAMURA, A, CLARK, N. M., RYMEN, B., BRAIDWOOD, L., MOROHASHI, K., BUSCH, W., BENFEY, P. N., SOZZANI, R. & SUGIMOTO, K. 2018. GTL1 and DF1 regulate root hair growth through transcriptional repression of ROOT HAIR DEFECTIVE 6-LIKE 4 in Arabidopsis. Development, 145, dev159707.
Google Scholar: Author Only Title Only Author and Title

SPARKS, J. A, KWON, T., RENNA, L., LIAO, F., BRANDIZZI, F. & BLANCAFLOR, E. B. 2016. HLB1 is a tetratricopeptide repeat domain-containing protein that operates at the intersection of the exocytic and endocytic pathways at the TGN/EE in Arabidopsis. The Plant Cell, 28, 746-769.
Google Scholar: Author Only Title Only Author and Title

STEFFENS, A, BRÄUTIGAM, A, JAKOBY, M. & HÜLSKAMP, M. 2015. The BEACH domain protein SPIRRIG is essential for Arabidopsis salt stress tolerance and functions as a regulator of transcript stabilization and localization. PLOS Biology, 13, e1002188.
Google Scholar: Author Only Title Only Author and Title

STEFFENS, A, JAKOBY, M. & HÜLSKAMP, M. 2017. Physical, functional and genetic interactions between the BEACH domain protein SPIRRIG and LIP5 and SKD1 and its role in endosomal trafficking to the vacuole in Arabidopsis. Frontiers in Plant Science, 8.
Google Scholar: Author Only Title Only Author and Title

STEPHAN, O. O. H. 2017. Actin fringes of polar cell growth. J Exp Bot, 68, 3303-3320.
Google Scholar: Author Only Title Only Author and Title

SUN, L., GE, Y., SPARKS, J. A, ROBINSON, Z. T., CHENG, X., WEN, J. & BLANCAFLOR, E. B. 2019. TDNAscan: A Software to Identify Complete and Truncated T-DNA Insertions. Frontiers in Genetics, 10, 685.
Google Scholar: Author Only Title Only Author and Title

SZYMANSKI, D. & STAIGER, C. J. 2018. The Actin Cytoskeleton: Functional Arrays for Cytoplasmic Organization and Cell Shape Control. Plant Physiology, 176, 106-118.
Google Scholar: Author Only Title Only Author and Title

SZYMANSKI, D. B. 2005. Breaking the WAVE complex: the point of Arabidopsis trichomes. Current Opinion in Plant Biology, 8, 103-112.
Google Scholar: Author Only Title Only Author and Title

TEAM, R. C. 2019. R: A language and environment for statistical computing. R Foundation For Statistical Computing. Vienna, Austria.
Google Scholar: Author Only Title Only Author and Title

UHRIG, J. F., MUTONDO, M., ZIMMERMANN, I., DEEKS, M. J., MACHESKY, L. M., THOMAS, P., UHRIG, S., RAMBKE, C., HUSSEY, P. J. & HULSKAMP, M. 2007. The role of Arabidopsis SCAR genes in ARP2-ARP3-dependent cell morphogenesis. Development, 134, 967-977.
Google Scholar: Author Only Title Only Author and Title

VIDALI, L., BURKART, G. M., AUGUSTINE, R. C., KERDAVID, E., TÜZEL, E. & BEZANILLA, M. 2010. Myosin XI Is Essential for Tip Growth in Physcomitrella patens. The Plant Cell, 22, 1868-1882.
Google Scholar: Author Only Title Only Author and Title

WANG, P., RICHARDSON, C., HAWES, C. & PATRICK 2016. Arabidopsis NAP1 Regulates the Formation of Autophagosomes. Current Biology, 26, 2060-2069.
Google Scholar: Author Only Title Only Author and Title

WANG, X., HERBERG, F. W., LAUE, M. M., WÜLLNER, C., HU, B., PETRASCH-PARWEZ, E. & KILIMANN, M. W. 2000. Neurobeachin: A Protein Kinase A-Anchoring,beige/Chediak-Higashi Protein Homolog Implicated in Neuronal Membrane Traffic. The Journal of Neuroscience, 20, 8551-8565.
Google Scholar: Author Only Title Only Author and Title

WICKHAM, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
Google Scholar: Author Only Title Only Author and Title

YANAGISAWA, M., ALONSO, J. M. & SZYMANSKI, D. B. 2018. Microtubule-Dependent Confinement of a Cell Signaling and Actin Polymerization Control Module Regulates Polarized Cell Growth. Current Biology, 28, 2459-2466.e4.

Google Scholar: Author Only Title Only Author and Title

YANAGISAWA, M., DESYATOVA, A. S., BELTETON, S. A., MALLERY, E. L., TURNER, J. A. & SZYMANSKI, D. B. 2015. Patterning mechanisms of cytoskeletal and cell wall systems during leaf trichome morphogenesis. Nature Plants, 1, 15014.
Google Scholar: Author Only Title Only Author and Title

YANAGISAWA, M., ZHANG, C. & SZYMANSKI, D. B. 2013. ARP2/3-dependent growth in the plant kingdom: SCARs for life. Frontiers in Plant Science, 4.
Google Scholar: Author Only Title Only Author and Title

YOO, C. M., QUAN, L., CANNON, A. E., WEN, J. & BLANCAFLOR, E. B. 2012. AGD1, a class 1 ARF-GAP, acts in common signaling pathways with phosphoinositide metabolism and the actin cytoskeleton in controlling Arabidopsis root hair polarity. Plant J, 69, 1064-76.
Google Scholar: Author Only Title Only Author and Title

ZHANG, C., MALLERY, E. L., SCHLUETER, J., HUANG, S., FAN, Y., BRANKLE, S., STAIGER, C. J. & SZYMANSKI, D. B. 2008. Arabidopsis SCARs Function Interchangeably to Meet Actin-Related Protein 2/3 Activation Thresholds during Morphogenesis. The Plant Cell, 20, 995-1011.
Google Scholar: Author Only Title Only Author and Title

ZHANG, X., DYACHOK, J., KRISHNAKUMAR, S., SMITH, L. G. & OPPENHEIMER, D. G. 2005. IRREGULAR TRICHOME BRANCH1 in Arabidopsis encodes a plant homolog of the actin-related protein2/3 complex activator Scar/WAVE that regulates actin and microtubule organization. Plant Cell, 17, 2314-26.
Google Scholar: Author Only Title Only Author and Title

ZHAO, Y., PAN, Z., ZHANG, Y., QU, X., ZHANG, Y., YANG, Y., JIANG, X., HUANG, S., YUAN, M., SCHUMAKER, K. S. & GUO, Y. 2013. The actin-related Protein2/3 complex regulates mitochondrial-associated calcium signaling during salt stress in Arabidopsis. Plant Cell, 25, 4544-59.
Google Scholar: Author Only Title Only Author and Title

ZHOU, R., BENAVENTE, L. M., STEPANOVA, A. N. & ALONSO, J. M. 2011. A recombineering-based gene tagging system for Arabidopsis. The Plant Journal, 66, 712-723.
Google Scholar: Author Only Title Only Author and Title

ZOU, J.-J., ZHENG, Z.-Y., XUE, S., LI, H.-H., WANG, Y.-R. & LE, J. 2016. The role of Arabidopsis Actin-Related Protein 3 in amyloplast sedimentation and polar auxin transport in root gravitropism. Journal of Experimental Botany, 67, 5325-5337.
Google Scholar: Author Only Title Only Author and Title