Thalamic spontaneous activity coordinates the timing of corticothalamic innervation in the visual system

Verónica Moreno-Juan, Francisco J. Martini, Leticia Pérez-Saiz, Álvaro Herrero-Navarro, Miguel Valdeolmillos and Guillermina López-Bendito*

Instituto de Neurociencias de Alicante, Universidad Miguel Hernández-Consejo Superior de Investigaciones Científicas (UMH-CSIC), Sant Joan d'Alacant, Spain.

*Correspondence: g.lbendito@umh.es

1   **Summary**

2   **Sensory processing relies on the correct development of circuits connecting**

3   **thalamus and cortex. Visual corticothalamic axons (CTAs) invade the thalamic**

4   **dorsolateral geniculate nucleus (dLGN) of the thalamus following an early**

5   **postnatal time-regulated programme in mice. Retinal spontaneous activity**

6   **influences the thalamic incursion of CTAs, however, the perinatal thalamus also**

7   **generates intrinsic patterns of spontaneous activity whose role in modulating**

8   **afferent connectivity remains unknown. Here, we found that patterned**

9   **spontaneous activity in the dLGN is critical for the proper spatial and temporal**

10 **innervation of CTAs. When the spontaneous dLGN activity is disrupted *in vivo*,**

11 **CTA innervation is severely delayed until eye-opening. Indeed, visual input**

12 **influenced the temporal development of CTAs by modulating thalamic activity, as**

13 **embryonic enucleation enhanced thalamic calcium waves and accelerated the**

14 **entrance of CTAs into the dLGN. Our results show that patterned spontaneous**

15 **activity in the perinatal thalamus is a key element driving the wiring of visual**

16 **circuits.**

17

18 **Keywords**

19 Thalamus, development, corticogeniculate, corticothalamic, thalamocortical,

20 spontaneous activity, retinal input.

21

Introduction

The formation of sensory circuits relies on the highly regulated spatial and temporal organization of axon connectivity. The visual system has been used extensively as a model to decipher the processes that control the development of neuronal networks, such as axon guidance and/or targeting. Indeed, retinofugal projections have been employed to study in some detail how topographic organization is established in the visual pathway (Fox & Guido, 2011; Huberman et al., 2008; Shatz & Rakic, 1981a; Zipursky & Sanes, 2010). The development of the visual system in mice, depends on feed-forward input from the retina through the retinothalamic axons (RTAs) that reach the dorsolateral geniculate nucleus (dLGN) of the thalamus on embryonic day 15 (E15), ensuring that ipsilateral and contralateral axon segregation is completed by postnatal day 8 (P8: Godement, Salaün, & Imbert, 1984a). By contrast, corticothalamic axons (CTAs) from layer 6 (L6) of the primary visual cortex (V1) reach the dorsal border of the dLGN at E18 and they begin to invade this nucleus around P4. Indeed, after the first postnatal week, CTAs have colonized the entire dLGN (Brooks et al., 2013; Grant, Hoerder-Suabedissen, & Molnár, 2016; Seabrook, El-Danaf, Krahe, Fox, & Guido, 2013).

Numerically, CTAs represent the largest input to the dLGN and it has been demonstrated that they are key regulators of dLGN cell firing (Erişir, Bickford, Van Horn, & Sherman, 1997; Godwin et al., 1996; McCormick & Krosigk, 1992). However, the feedback from the V1 to dLGN develops relatively late with respect to the retinal inputs (Drager & Hubel, 1975; Godement, Salaün, & Imbert, 1984b; Grant et al., 2016; Jacobs et al., 2007; Seabrook et al., 2013; Shatz & Rakic, 1981b). It has been suggested that RTAs and CTAs might interact to ensure the correct development of both projections during the second postnatal week (Bickford et al., 2009; Seabrook et al.,

2013). More recent insights into this process demonstrate that the invasion of the dLGN

by cortical and retinal inputs relies on their mutual interactions at the thalamus (Brooks

et al., 2013; Grant et al., 2016; Jurgens, Bell, McQuiston, & Guido, 2012; Seabrook et

al., 2013; Thompson, Gribizis, Chen, & Crair, 2017). These interactions seem to be

activity-dependent, as disrupting postnatal retinal waves by injecting the cholinergic

agonist epibatidine accelerates CTA entry into the dLGN (Grant et al., 2016). Indeed,

postnatal enucleation of mice leads to earlier innervation of the dLGN by CTAs (Brooks

et al., 2013; Grant et al., 2016; Seabrook et al., 2013). These results suggest that early

peripheral input and the electric messages conveyed are key elements for the timely

innervation and correct development of CTAs in the visual pathway. However, retinal

waves are not the only activity that trains the developing thalamus. It was recently

shown that the perinatal thalamus generates an intrinsic pattern of spontaneous activity

that is known to regulate the organization of sensory systems (Antón-Bolaños et al.,

2019; Castillo-Paterna et al., 2015; Mire et al., 2012; Moreno-Juan et al., 2017),

suggesting this activity may possibly regulate the innervation of the dLGN. Here, we

tested this hypothesis using an *in vivo* approach in which thalamic spontaneous activity

is modified in the dLGN. We found that alterations to the patterns of spontaneous

thalamic activity in the dLGN during prenatal and postnatal stages impair the timing of

CTA innervation of this nucleus. Remarkably, the absence of CTAs in the dLGN is

rescued by eye opening, strongly suggesting that both intrinsic spontaneous activity and

experience-dependent programmes play a role in the development of the

corticothalamic circuits associated with the visual system.

Results

Disrupting the pattern of spontaneous activity in the developing visual thalamus

To determine whether innervation of the dLGN by visual CTAs requires thalamic

spontaneous activity, we disrupted the activity of the thalamus by overexpressing the

inward rectifier potassium channel 2.1 (Kir2.1) in all the principal thalamic sensory

nuclei. As such, we administered tamoxifen to pregnant $Gbx2^{CreER};R26^{Kir2.1-floxed}$ mice

(referred herein as Th^{Kir}) when embryos were at E10.5, inducing the overexpression of

Kir2.1 (fused to mCherry) in embryonic thalamic neurons (Antón-Bolaños et al., 2019).

As expected, the increase in Kir2.1 suppressed the patterned spontaneous activity in the

dLGN of new-born Th^{Kir} mice, this activity normally evident as calcium waves (**Figure**

1A-D: Moreno-Juan et al., 2017). Calcium imaging of *ex vivo* brain slices from Th^{Kir}

mice at P0 demonstrated that neurons from the dLGN only activated asynchronously,

with no large and propagating co-activation witnessed, such as that recorded in control

slices. Therefore, and as shown previously for the somatosensory and auditory nuclei of

the thalamus (Moreno-Juan et al., 2017; Antón-Bolaños et al., 2019), overexpression of

Kir2.1 transforms the pattern of dLGN activity from synchronous to asynchronous at

perinatal stages. Towards the end of the first postnatal week, an additional change in the

pattern of spontaneous activity was recorded in the Th^{Kir} mice *in vivo*. Extracellular

recordings with multichannel electrodes in the thalamus revealed that the firing rate of

dLGN neurons decreased significantly in the Th^{Kir} mice at P6 (**Figure 1E-H**).

Therefore, there is *ex vivo* and *in vivo* evidence that the overexpression of Kir2.1 alters

the properties of spontaneous activity in the visual thalamus during the initial days post-

partum, the time when CTAs reach and invade the dLGN.

Thalamic spontaneous activity mediates the entry of CTAs into the dLGN

In light of the above, we determined whether an abnormal intrinsic pattern of

spontaneous activity in the thalamus could affect the progression of CTAs into the

dLGN. As such, we visualized CTAs in the dLGN through their expression of the

vGlut1 transporter expressed at the terminals of corticothalamic neurons (Grant et al.,

2016; Piñon, Jethwa, Jacobs, Campagnoni, & Molnár, 2009; Varoqui, Schäfer, Zhu,

Weihe, & Erickson, 2002), as well as in the transgenic mouse line *Golli-tau-eGfp* that

labels cortical L6 neurons and their axons (Grant et al., 2016; Jacobs et al., 2007). At

P3, control axons from L6 appeared adjacent to the ventromedial border of the dLGN,

progressively invading the nucleus over successive days (**Figure S1**: Brooks et al.,

2013). While control CTAs had already entered the dLGN by P4, they remained at the

border in the Th^{Kir} mice (**Figure 2A**). In addition, we also measured the invasion of

CTAs in the dLGN in terms of the vGlut1 fluorescence in the dLGN area of control and

Th^{Kir} mice at P4, P6 and P9. The invasion of the dLGN by CTAs was severely perturbed

in the Th^{Kir} mice and while CTAs covered more than half of the nucleus in control

animals at P9, the invasion of this territory by CTAs was reduced by more than 90%

when thalamic spontaneous activity was disturbed (**Figure 2B**). This result was

confirmed by labelling L6 neurons from the V1 with a lentivirus expressing *Gfp*,

showing that virtually no CTAs had invaded the dLGN in Th^{Kir} mice at P9 (**Figure 2C**).

This effect was not due to a developmental delay in the arrival of CTAs in these Th^{Kir}

mice since L6 axons appear to be waiting at the border of the dLGN in both the control

and Th^{Kir} mice at P0 (**Figure 2D**). Together these results indicate that spontaneous

activity in the dLGN regulates the invasion of this structure by CTAs.

Disrupting the patterned thalamic activity leads dLGN neuron death

When analysing CTA invasion in the Th^{Kir} mice, we noticed that perturbing

spontaneous thalamic activity provoked a decrease in the size of the dLGN. While the

dLGN size was normal in Th^{Kir} mice from P0 to P3, it remained stationary from P4

onwards, a period in which the size of the dLGN normally increased in control mice and

when the CTAs invaded the thalamus (**Figure 3A**). To determine whether this lack of

growth of the dLGN is due to an increase in cell-death and/or the enhanced compaction

of dLGN cells, we analysed caspase-3a activity (Ahern et al., 2013; Lossi, Castagna, &

Merighi, 2015) and the cell density in the dLGN of control and Th^{Kir} mice. There was a

significant increase in caspase-3a immunostaining from P3 to P6 in the dLGN of the

Th^{Kir} mice (**Figure 3B and 3C**) but no increase in cell density in this nucleus (**Figure**

S2A and S2B). Hence, these data suggest that the failure to increase the size of the

dLGN in the Th^{Kir} mice might be the result of enhanced cell-death in this nucleus, in

conjunction with an absence of CTAs in the visual thalamus.

The lack of innervation of the dLGN by CTAs is independent of RTAs

It was previously suggested that a hierarchy exists in the temporal invasion of the dLGN

by afferent axons, with RTAs preceding CTAs (Brooks et al., 2013; Guido, 2018;

Jacobs et al., 2007; Seabrook et al., 2013). Furthermore, it was suggested that cortical

input from L6 neurons is required for the correct targeting of RTAs to the dLGN and for

their eye-specific segregation (Shanks et al., 2016; Thompson, Picard, Min, Fagiolini, &

Chen, 2016). Therefore, the mechanisms defining axon interactions might regulate the

invasion of the dLGN by these two sets of projections, whereby RTAs are reorganized

and segregated once CTAs start to innervate the dLGN. Indeed, Cholera-Toxin subunit

B (CTB) injections into the eyes of *Golli-tau-eGfp* mice revealed that RTAs preceded

CTA innervation of the dLGN (**Fig. 4A** and **4B**), supporting previous findings

(Seabrook et al., 2013). As perturbed thalamic spontaneous activity strongly affects the

innervation of the dLGN by CTAs, we tested whether the absence of CTAs in the Th^{Kir}

mouse is a consequence of an early defect in RTA innervation. As such, we performed

anterograde labelling of RTAs by injecting Alexa-647 conjugated CTB into the retina.

Remarkably, we found that the timing of contralateral and ipsilateral RTA invasion of

the dLGN is normal at P2 in Th^{Kir} mice (**Figure 4C and 4D**). Hence, retinal axons

target and invade the dLGN independently of thalamic activity and the absence of CTAs

in the dLGN is not due to a defect of the RTAs.

Retinal inputs modify the spontaneous thalamic activity in the dLGN

Our previous results demonstrated that normal spontaneous activity in the dLGN is

required for CTAs to invade this nucleus. Interestingly, the pattern of the spontaneous

activity in the somatosensory nucleus of the thalamus is thought to be modified after the

removal of retinal input due to embryonic bilateral enucleation (embBE: Moreno-Juan

et al., 2017). Hence, we cannot rule out that the presence of RTAs might somehow also

influence spontaneous activity in the visual thalamus and thereby affect the

development of CTAs. In fact, postnatal deprivation of the visual periphery input

provokes an aberrant development of the CTAs in the dLGN (Grant et al., 2016; Jacobs

et al., 2007; Seabrook et al., 2013). To test this possibility, we performed embryonic

bilateral enucleations at E14.5 to ensure that visual input never reached the thalamus, as

retinal axons reach the dLGN at E15.5 in mice (Moreno-Juan et al., 2017). To detect the

spontaneous activity in the dLGN, we performed calcium imaging *ex vivo* in control and

embBE slices at P1, prior to CTA invasion of the dLGN (**Figure 5A and Figure 5E**).

At this stage we found a higher frequency of spontaneous synchronous calcium events

in the dLGN of embBE mice, which included the appearance of more thalamic calcium

waves than in control mice (**Figure 5B and 5C**). As such, these experiments show that

despite the emergence of thalamic waves that are independent of retinal input, RTAs

exert a modulatory role on the patterned activity of the dLGN that is revealed in

enucleated animals.

We next determined whether innervation of the dLGN by CTAs is affected in

embBE mice by performing enucleations at E14.5 in wild-type and *Golli-tau-eGfp* mice

(**Figure 5D**). In control animals, CTAs began to invade the dLGN from P4 and they

have covered the entire nucleus by around P12, close to the time of eye-opening

(Brooks et al., 2013; Seabrook et al., 2013). By contrast, in embBE mice CTAs

expressing GFP or labelled for vGlut1 immunostaining already covered a significantly

higher proportion of the dLGN (8.85%) at P1 than in their control littermates (3.25%).

This acceleration of dLGN invasion was maintained at all the developmental stages

studied (P2, P3 and P7: **Figure 5E** and **5F, Figure S3**), and it was in accordance with

data found previously in mice enucleated at P0 (Brooks et al., 2013; Grant et al., 2016;

Seabrook et al., 2013).

These results show that the arrival of retinal axons has strong impact on the

developing dLGN. First, they regulate the pattern of spontaneous activity and secondly,

they influence the postnatal innervation of the dLGN by CTAs. However, it remains

unclear whether there is a causal link between these events. In order to address this

question, we performed bilateral embryonic enucleations in the Th^{Kir} mice (**Figure 5G**).

Remarkably, we found that CTA innervation of the dLGN is rescued when RTAs are

eliminated in Th^{Kir} mice. The relative invasion of the dLGN by CTAs in the embBE

Th^{Kir} mice was similar to that of embBE mice (**Figure 5H and 5I**), demonstrating the

existence of a hierarchical organization in the invasion of the dLGN whereby the arrival

of RTAs switches the pattern of spontaneous activity in the nucleus, in turn, delaying

the entry of CTAs.

Visual experience rescues CTA innervation of the dLGN in the *Th^{Kir}* mice

Our results demonstrated that altering the patterns of early thalamic spontaneous

activity affects the development of the corticothalamic connections to the dLGN. The

reduction in spontaneous dLGN activity due to the absence of thalamic calcium waves

induced a severe defect in CTA innervation, which were virtually absent from the

dLGN during the first postnatal week, and at least until P9. Nevertheless, as *Th^{Kir}* mice

survive until adulthood, we wondered whether this effect might persist. Notably, at P30

CTAs had innervated the full extent of the dLGN in *Th^{Kir}* mice as in control littermates

(**Figure S4A**), indicating that the aberrant CTA innervation detected early in

development is reversed at later stages.

Eye-opening occurs in mice at around P13, an event that is crucial for the final

maturation and refinement of the visual system (Hensch, 2005; Hofer, Mrsic-Flogel,

Bonhoeffer, & Hübener, 2009; Hooks & Chen, 2006; Pecka, Han, Sader, & Mrsic-

Flogel, 2014). As CTA invasion of the dLGN is rescued after P9, we assessed whether

eye-opening might play a role in this recovery. We analysed the innervation of the

dLGN by CTAs in both control and *Th^{Kir}* mice at P10, P13 and P14, just before and

after eye-opening. We found that while at P10 when the eyes are still closed, CTAs

poorly innervated the dLGN in *Th^{Kir}* mice relative to the controls (**Figure 6A**).

However, after eye-opening, the CTA invasion in *Th^{Kir}* mice at P13-P14 was

indistinguishable from that of the control mice (**Figure 6B and S4B**).

Visual stimulation through close eyelids evokes glutamatergic retinal waves, and

light deprivation in mice from P8 to P14 provokes deficient retinogeniculate

segregation, demonstrating that activity derived by light stimulus just prior to eye

opening influences the topographic development of the visual system (Tiriac, Smith, &

Feller, 2018). Thus, we hypothesised that visual experience may alter the network

properties of dLGN neurons, allowing CTAs to enter the dLGN. To test this possibility,

we maintained control and Th^{Kir} mice in a dark rearing (DR) environment from P5 until

P15 and we then analysed the CTA innervation of the dLGN (**Figure 6C**). We did not

find major modifications in CTA innervation at P15 in control animals subjected to DR

(**Figure 6D and 6E**). However, in contrast to the Th^{Kir} mice, there was a significant

reduction in CTA invasion of the dLGN in DR Th^{Kir} mice after eye opening (**Figure 6D**

and 6E). Together these results demonstrate that early visual experience is necessary

for the plastic recovery of CTA invasion of the dLGN that occurs in juvenile Th^{Kir} mice.

Discussion

Here we describe a novel aspect in the coordination of afferent connectivity to the

dLGN by intrinsic thalamic spontaneous activity. Using a model in which spontaneous

dLGN activity is perturbed, we show that the timing of CTA entry into this structure

depends on this thalamic activity. In a scenario of impaired thalamic spontaneous

activity, CTA invasion of the dLGN is severely delayed. In addition, the lack of calcium

waves in the dLGN leads to a decrease in its size and a concomitant increase in cell

death, as occurs after embBE or in Math5 mutant mice where retinal input is absent

from embryonic stages (El-Danaf et al., 2015; Moreno-Juan et al., 2017). Hence, in

addition to the trophic factors produced by retinal innervation, spontaneous thalamic

activity may directly regulate dLGN neuronal survival, as in the developing cortex

(Blanquie et al., 2017; Luhmann & Khazipov, 2018). However, it is also possible that

the effect on cell death is indirect and due to the lack of a proper corticogeniculate

innervation, although this will require further study. While CTAs invade the dLGN after

RTAs, it was suggested that CTAs regulate the initial targeting of the visual thalamus

by the retina. In transgenic mice in which the V1 was eliminated and where CTAs

therefore fail to develop, some RGCs did not project to the dLGN (Diao et al., 2018;

Shanks et al., 2016). Moreover, when the activity of L6 neurons was silenced in the V1,

functional modifications were apparent in visually-evoked activity in the thalamus

(Denman & Contreras, 2015). Despite these findings, our work on the Th^{Kir} mice in

which CTAs are absent in the dLGN and RTAs normally innervate this nucleus suggest

the retinothalamic afferents from the CTAs initial develop following an independent

programme.

256   Retinal waves are transmitted to the dLGN and the superior colliculus, and this

transmission is crucial for the correct development of visual topographic maps

(Ackman, Burbridge, & Crair, 2012; Colonnese & Khazipov, 2010; Mooney, Penn,

Gallego, & Shatz, 1996; Siegel, Heimel, Peters, & Lohmann, 2012). During prenatal

development, type I retinal waves coexist with the other patterns of spontaneous activity

in the thalamus. Although the role of type I retinal waves in visual system development

remains poorly understood, here we demonstrate that embryonic ablation of the retinal

input, before the retinal axons reach the dLGN, modulates thalamic wave activity. We

found that the frequency of thalamic calcium waves augments in the dLGN of embBE

mice, which leads to a premature invasion of the dLGN by CTAs. Although the

functional synapses between CTAs and dLGN neurons are established during the

second postnatal week (Seabrook et al., 2013; Bickford et al., 2010), vGlut1 is already

present in the terminals of the CTAs that invade the dLGN of embBE mice at P1. This

transporter has previously been associated with functional excitatory transmission

(Koch et al., 2011; Marcucci, Zou, & Firestein, 2009; Nakamura, Hioki, Fujiyama, &

Kaneko, 2005; Sherry, Wang, Bates, & Frishman, 2003; Wilson et al., 2005). Thus, it is

possible that eliminating retinal input accelerates corticogeniculate maturation in the

dLGN.

Our data from Th^{Kir} mice indicate that spontaneous thalamic activity is crucial

for the timely invasion of the dLGN by CTAs but also, they helped us unravel the

hierarchical organization of RTAs, CTAs and thalamic spontaneous activity. When

retinal input is eliminated in Th^{Kir} mice the lack of CTAs in the dLGN was completely

rescued, as not only the absence of thalamic waves but also, the presence of RTAs in

Th^{Kir} mice prevented the entry of CTAs into the dLGN. As such, it would appear that

RTAs and CTAs compete for dLGN innervation, with RTAs arriving first, and that

spontaneous thalamic activity represents a permissive phenomenon in this context.

The analysis of CTAs in postnatal Th^{Kir} mice shows that the delay in CTA

invasion of the dLGN is reversed at eye-opening, when visually evoked activity

commences and spontaneous retinal waves subside (Demas, Eglen, & Wong, 2003;

Huberman et al., 2008). At this stage, CTAs innervate the dLGN despite the disturbed

thalamic activity, suggesting that visual evoked inputs are sufficient to drive CTA

innervation of the dLGN. Indeed, visual evoked activity is necessary for this recovery as

CTAs did not fully recover in DR Th^{Kir} mice.

In summary, our data suggest patterned spontaneous activity in the perinatal

thalamus provides a fundamental signal that regulates the timing of CTA invasion of the

dLGN, indicating that the thalamus is as a key structure in the development of visual

circuits. Furthermore, these results also suggest the existence of an experience-

dependent mechanism that helps to recover from early defects in circuits within the

visual system.

Acknowledgments

The authors are grateful to L.M. Rodríguez, B. Andrés and R. Susín for their excellent

technical support; to T. Theil and A. Campagnoni for providing the *Golli-tau-eGfp*

mouse line; and to members of G. López-Bendito's laboratory for stimulating

discussions. Funding: this work was supported by grants from the European Research

Council (ERC-2014-CoG-647012) and the Spanish Ministry of Science, Innovation and

Universities (PGC2018/096631-B-I00, BFU2015-64432-R and Severo Ochoa Grant

SEV-2017-0723).

Author Contributions

V.M-J. and G.L-B. conceived the idea and designed the experiments. V.M-J. conducted

the experiments and analyzed the calcium imaging data, corticothalamic innervation,

cell death, CTB tracing, embryonic enucleations and dark-reared experiments. L.P-S

performed the *in vivo* recordings in the Th^{Kir} mice, F.J.M generated the Matlab codes

and F.J.M and L.P-S analyzed this data. A.H-N performed the lentiviral injections in

Th^{Kir} mice. G.L-B. acquired the funding; V.M-J, F.J.M, M.V and G.L-B. wrote the

paper.

Declaration of Interests

The authors have no competing interests to declare

References

Ackman, J. B., Burbridge, T. J., & Crair, M. C. (2012). Retinal waves coordinate

patterned activity throughout the developing visual system. *Nature*, *490*(7419),

219–225. http://doi.org/10.1038/nature11529

Ahern, T. H., Krug, S., Carr, A. V., Murray, E. K., Fitzpatrick, E., Bengston, L., et al.

(2013). Cell death atlas of the postnatal mouse ventral forebrain and hypothalamus:

effects of age and sex. *The Journal of Comparative Neurology*, *521*(11), 2551–

2569. http://doi.org/10.1002/cne.23298

Antón-Bolaños, N., Sempere-Ferràndez, A., Guillamón-Vivancos, T., Martini, F. J.,

Pérez-Saiz, L., Gezelius, H., et al. (2019). Prenatal activity from thalamic neurons

governs the emergence of functional cortical maps in mice. *Science (New York,*

N.Y.), *364*(6444), 987–990. http://doi.org/10.1126/science.aav7617

Bickford, M. E., Slusarczyk, A., Dilger, E. K., Krahe, T. E., Kucuk, C., & Guido, W.

(2009). Synaptic development of the mouse dorsal lateral geniculate nucleus. *The*

Journal of Comparative Neurology, *518*(5), 622–635.

http://doi.org/10.1002/cne.22223

Blanquie, O., Yang, J.-W., Kilb, W., Sharopov, S., Sinning, A., & Luhmann, H. J.

(2017). Electrical activity controls area-specific expression of neuronal apoptosis in

the mouse developing cerebral cortex. *eLife*, *6*. http://doi.org/10.7554/eLife.27696

Brooks, J. M., Su, J., Levy, C., Wang, J. S., Seabrook, T. A., Guido, W., & Fox, M. A.

(2013). A molecular mechanism regulating the timing of corticogeniculate

innervation. *Cell Reports*, *5*(3), 573–581.

http://doi.org/10.1016/j.celrep.2013.09.041

Castillo-Paterna, M., Moreno-Juan, V., Filipchuk, A., Rodríguez-Malmierca, L., Susín,

R., & López-Bendito, G. (2015). DCC functions as an accelerator of

343 thalamocortical axonal growth downstream of spontaneous thalamic activity.

344 *EMBO Reports*, *16*(7), 851–862. http://doi.org/10.15252/embr.201439882

345 Chen, L., Guo, Q., & Li, J. Y. H. (2009). Transcription factor Gbx2 acts cell-

346 nonautonomously to regulate the formation of lineage-restriction boundaries of the

347 thalamus. *Development*, *136*(8), 1317–1326. http://doi.org/10.1242/dev.030510

348 Colonnese, M. T., & Khazipov, R. (2010). "Slow activity transients" in infant rat visual

349 cortex: a spreading synchronous oscillation patterned by retinal waves. *Journal of*

350 *Neuroscience*, *30*(12), 4325–4337. http://doi.org/10.1523/JNEUROSCI.4995-

351 09.2010

352 Demas, J., Eglen, S. J., & Wong, R. O. L. (2003). Developmental loss of synchronous

353 spontaneous activity in the mouse retina is independent of visual experience. *The*

354 *Journal of Neuroscience : the Official Journal of the Society for Neuroscience*,

355 *23*(7), 2851–2860. http://doi.org/10.1523/JNEUROSCI.23-07-02851.2003

356 Denman, D. J., & Contreras, D. (2015). Complex Effects on In Vivo Visual Responses

357 by Specific Projections from Mouse Cortical Layer 6 to Dorsal Lateral Geniculate

358 Nucleus. *Journal of Neuroscience*, *35*(25), 9265–9280.

359 http://doi.org/10.1523/JNEUROSCI.0027-15.2015

360 Diao, Y., Cui, L., Chen, Y., Burbridge, T. J., Han, W., Wirth, B., et al. (2018).

361 Reciprocal Connections Between Cortex and Thalamus Contribute to Retinal Axon

362 Targeting to Dorsal Lateral Geniculate Nucleus. *Cerebral Cortex*, *28*(4), 1168–

363 1182. http://doi.org/10.1093/cercor/bhx028

364 Drager, U. C., & Hubel, D. H. (1975). Physiology of visual cells in mouse superior

365 colliculus and correlation with somatosensory and auditory input. *Nature*,

366 *253*(5488), 203–204. http://doi.org/10.1038/253203a0

El-Danaf, R. N., Krahe, T. E., Dilger, E. K., Bickford, M. E., Fox, M. A., & Guido, W.

(2015). Developmental remodeling of relay cells in the dorsal lateral geniculate

nucleus in the absence of retinal input. *Neural Development*, *10*(1), 19–17.

http://doi.org/10.1186/s13064-015-0046-6

Erişir, A., Bickford, M. E., Van Horn, S. C., & Sherman, S. M. (1997). Relative

numbers of cortical and brainstem inputs to the lateral geniculate nucleus.

Proceedings of the National Academy of Sciences of the United States of America,

94(4), 1517–1520. http://doi.org/10.1073/pnas.94.4.1517

Fox, M. A., & Guido, W. (2011). Shedding light on class-specific wiring: development

of intrinsically photosensitive retinal ganglion cell circuitry. *Molecular*

Neurobiology, *44*(3), 321–329. http://doi.org/10.1007/s12035-011-8199-8

Godement, P., Salaün, J., & Imbert, M. (1984a). Prenatal and postnatal development of

retinogeniculate and retinocollicular projections in the mouse. *The Journal of*

Comparative Neurology, *230*(4), 552–575. http://doi.org/10.1002/cne.902300406

Godement, P., Salaün, J., & Imbert, M. (1984b). Prenatal and postnatal development of

retinogeniculate and retinocollicular projections in the mouse. *The Journal of*

Comparative Neurology, *230*(4), 552–575. http://doi.org/10.1002/cne.902300406

Godwin, D. W., Van Horn, S. C., Eriir, A., Sesma, M., Romano, C., & Sherman, S. M.

(1996). Ultrastructural localization suggests that retinal and cortical inputs access

different metabotropic glutamate receptors in the lateral geniculate nucleus. *Journal*

of Neuroscience, *16*(24), 8181–8192. http://doi.org/10.1523/JNEUROSCI.16-24-

08181.1996

Grant, E., Hoerder-Suabedissen, A., & Molnár, Z. (2016). The Regulation of

Corticofugal Fiber Targeting by Retinal Inputs. *Cerebral Cortex*, *26*(3), 1336–1348.

http://doi.org/10.1093/cercor/bhv315

Guido, W. (2018). Development, form, and function of the mouse visual thalamus.

Journal of Neurophysiology, *120*(1), 211–225. http://doi.org/10.1152/jn.00651.2017

Hensch, T. K. (2005). Critical period plasticity in local cortical circuits. *Nature Reviews*

Neuroscience, *6*(11), 877–888. http://doi.org/10.1038/nrn1787

Hofer, S. B., Mrsic-Flogel, T. D., Bonhoeffer, T., & Hübener, M. (2009). Experience

leaves a lasting structural trace in cortical circuits. *Nature*, *457*(7227), 313–317.

http://doi.org/10.1038/nature07487

Hooks, B. M., & Chen, C. (2006). Distinct roles for spontaneous and visual activity in

remodeling of the retinogeniculate synapse. *Neuron*, *52*(2), 281–291.

http://doi.org/10.1016/j.neuron.2006.07.007

Huberman, A. D., Manu, M., Koch, S. M., Susman, M. W., Lutz, A. B., Ullian, E. M.,

et al. (2008). Architecture and activity-mediated refinement of axonal projections

from a mosaic of genetically identified retinal ganglion cells. *Neuron*, *59*(3), 425–

438. http://doi.org/10.1016/j.neuron.2008.07.018

Jacobs, E. C., Campagnoni, C., Kampf, K., Reyes, S. D., Kalra, V., Handley, V., et al.

(2007). Visualization of corticofugal projections during early cortical development

in a tau-GFP-transgenic mouse. *The European Journal of Neuroscience*, *25*(1), 17–

30. http://doi.org/10.1111/j.1460-9568.2006.05258.x

Jurgens, C. W. D., Bell, K. A., McQuiston, A. R., & Guido, W. (2012). Optogenetic

stimulation of the corticothalamic pathway affects relay cells and GABAergic

neurons differently in the mouse visual thalamus. *PloS One*, *7*(9), e45717.

http://doi.org/10.1371/journal.pone.0045717

Koch, S. M., Cruz, Dela, C. G., Hnasko, T. S., Edwards, R. H., Huberman, A. D., &

Ullian, E. M. (2011). Pathway-specific genetic attenuation of glutamate release

alters select features of competition-based visual circuit refinement. *Neuron*, *71*(2),

235–242. http://doi.org/10.1016/j.neuron.2011.05.045

Lossi, L., Castagna, C., & Merighi, A. (2015). Neuronal cell death: an overview of its

different forms in central and peripheral neurons. *Methods in Molecular Biology*

(Clifton, N.J.), *1254*, 1–18. http://doi.org/10.1007/978-1-4939-2152-2_1

Luhmann, H. J., & Khazipov, R. (2018). Neuronal activity patterns in the developing

barrel cortex. *Neuroscience*, *368*, 256–267.

http://doi.org/10.1016/j.neuroscience.2017.05.025

Marcucci, F., Zou, D.-J., & Firestein, S. (2009). Sequential onset of presynaptic

molecules during olfactory sensory neuron maturation. *The Journal of Comparative*

Neurology, *516*(3), 187–198. http://doi.org/10.1002/cne.22094

McCormick, D. A., & Krosigk, von, M. (1992). Corticothalamic activation modulates

thalamic firing through glutamate "metabotropic" receptors. *Proceedings of the*

National Academy of Sciences of the United States of America, *89*(7), 2774–2778.

http://doi.org/10.1073/pnas.89.7.2774

Mire, E., Mezzera, C., Leyva-Díaz, E., Paternain, A. V., Squarzoni, P., Bluy, L., et al.

(2012). Spontaneous activity regulates Robo1 transcription to mediate a switch in

thalamocortical axon growth. *Nature Neuroscience*, *15*(8), 1134–1143.

http://doi.org/10.1038/nn.3160

Mooney, R., Penn, A. A., Gallego, R., & Shatz, C. J. (1996). Thalamic relay of

spontaneous retinal activity prior to vision. *Neuron*, *17*(5), 863–874.

http://doi.org/10.1016/s0896-6273(00)80218-4

Moreno-Juan, V., Filipchuk, A., Antón-Bolaños, N., Mezzera, C., Gezelius, H., Andrés,

B., et al. (2017). Prenatal thalamic waves regulate cortical area size prior to sensory

processing. *Nature Communications*, *8*(1), 14172–14.

http://doi.org/10.1038/ncomms14172

Nakamura, K., Hioki, H., Fujiyama, F., & Kaneko, T. (2005). Postnatal changes of

vesicular glutamate transporter (VGluT)1 and VGluT2 immunoreactivities and their

colocalization in the mouse forebrain. *The Journal of Comparative Neurology*,

492(3), 263–288. http://doi.org/10.1002/cne.20705

Pecka, M., Han, Y., Sader, E., & Mrsic-Flogel, T. D. (2014). Experience-dependent

specialization of receptive field surround for selective coding of natural scenes.

Neuron, *84*(2), 457–469. http://doi.org/10.1016/j.neuron.2014.09.010

Piñon, M. C., Jethwa, A., Jacobs, E., Campagnoni, A., & Molnár, Z. (2009). Dynamic

integration of subplate neurons into the cortical barrel field circuitry during

postnatal development in the Golli-tau-eGFP (GTE) mouse. *The Journal of

Physiology*, *587*(Pt 9), 1903–1915. http://doi.org/10.1113/jphysiol.2008.167767

Seabrook, T. A., El-Danaf, R. N., Krahe, T. E., Fox, M. A., & Guido, W. (2013).

Retinal input regulates the timing of corticogeniculate innervation. *The Journal of

Neuroscience : the Official Journal of the Society for Neuroscience*, *33*(24), 10085–

10097. http://doi.org/10.1523/JNEUROSCI.5271-12.2013

Shanks, J. A., Ito, S., Schaevitz, L., Yamada, J., Chen, B., Litke, A., & Feldheim, D. A.

(2016). Corticothalamic Axons Are Essential for Retinal Ganglion Cell Axon

Targeting to the Mouse Dorsal Lateral Geniculate Nucleus. *The Journal of

Neuroscience : the Official Journal of the Society for Neuroscience*, *36*(19), 5252–

5263. http://doi.org/10.1523/JNEUROSCI.4599-15.2016

Shatz, C. J., & Rakic, P. (1981a). The genesis of efferent connections from the visual

cortex of the fetal rhesus monkey. *The Journal of Comparative Neurology*, *196*(2),

287–307. http://doi.org/10.1002/cne.901960208

Shatz, C. J., & Rakic, P. (1981b). The genesis of efferent connections from the visual

cortex of the fetal rhesus monkey. *The Journal of Comparative Neurology*, *196*(2),

287–307. http://doi.org/10.1002/cne.901960208

Sherry, D. M., Wang, M. M., Bates, J., & Frishman, L. J. (2003). Expression of

vesicular glutamate transporter 1 in the mouse retina reveals temporal ordering in

development of rod vs. cone and ON vs. OFF circuits. *The Journal of Comparative*

Neurology, *465*(4), 480–498. http://doi.org/10.1002/cne.10838

Siegel, F., Heimel, J. A., Peters, J., & Lohmann, C. (2012). Peripheral and central inputs

shape network dynamics in the developing visual cortex in vivo. *Current Biology :*

CB, *22*(3), 253–258. http://doi.org/10.1016/j.cub.2011.12.026

Thompson, A. D., Picard, N., Min, L., Fagiolini, M., & Chen, C. (2016). Cortical

Feedback Regulates Feedforward Retinogeniculate Refinement. *Neuron*, *91*(5),

1021–1033. http://doi.org/10.1016/j.neuron.2016.07.040

Thompson, A., Gribizis, A., Chen, C., & Crair, M. C. (2017). Activity-dependent

development of visual receptive fields. *Current Opinion in Neurobiology*, *42*, 136–

143. http://doi.org/10.1016/j.conb.2016.12.007

Tiriac, A., Smith, B. E., & Feller, M. B. (2018). Light Prior to Eye Opening Promotes

Retinal Waves and Eye-Specific Segregation. *Neuron*, *100*(5), 1059–1065.e4.

http://doi.org/10.1016/j.neuron.2018.10.011

Varoqui, H., Schäfer, M. K. H., Zhu, H., Weihe, E., & Erickson, J. D. (2002).

Identification of the differentiation-associated Na+/PI transporter as a novel

vesicular glutamate transporter expressed in a distinct set of glutamatergic synapses.

The Journal of Neuroscience : the Official Journal of the Society for Neuroscience,

22(1), 142–155. http://doi.org/10.1523/JNEUROSCI.22-01-00142.2002

Wilson, N. R., Kang, J., Hueske, E. V., Leung, T., Varoqui, H., Murnick, J. G., et al.

(2005). Presynaptic regulation of quantal size by the vesicular glutamate transporter

VGLUT1. *Journal of Neuroscience*, *25*(26), 6221–6234.

http://doi.org/10.1523/JNEUROSCI.3003-04.2005

Zipursky, S. L., & Sanes, J. R. (2010). Chemoaffinity revisited: dscams, protocadherins,

and neural circuit assembly. *Cell*, *143*(3), 343–353.

http://doi.org/10.1016/j.cell.2010.10.009

     Main Figures

Figure 1. Overexpression of Kir2.1 gene in the dLGN disrupts the patterns of

spontaneous activity. (A) Scheme of the experimental design used for calcium imaging

of *ex vivo* slices from control and Th^{Kir} mice at P0. (B) *Left*, maximal projection of an *ex*

vivo thalamic spontaneous calcium wave in the dLGN of a control mouse at P0. *Right*,

quantification of the relative active Region Of Interest (ROIs) in the dLGN over a 15

minute recording (top), and the corresponding raster plot of the active ROIs per frame

(bottom). (C) *Left*, maximal projection of *ex vivo* thalamic spontaneous calcium activity

in the dLGN of Th^{Kir} mouse at P0. *Right*, quantification of the proportion of active ROIs

in the dLGN over 15 minutes recording (top) and the corresponding raster plot of active

ROIs per frame (bottom). (D) *Left,* graph showing the mean event frequency per ROI

per minute in the dLGN of control and Th^{Kir} mice. *Right,* graph showing the mean

frequency of the waves in the dLGN of control and Th^{Kir} mice at P0 (control $n = 7$, Th^{Kir}

$n = 5$). (E) Scheme of the experimental design used for the *in vivo* recording of

spontaneous activity in control and Th^{Kir} mice at P6. (F) Coronal sections of control and

Th^{Kir} mice revealing the position of the recording electrodes in the dLGN. Red

fluorescence corresponds to the mCherry fused to the Kir2.1 gene (labelled by RFP

immunostaining). (G) Sample traces that illustrate the spiking activity in control and

Th^{Kir} mice at P6. (H) Quantification of the firing rate in the dLGN at P6 (control $n = 6$,

Th^{Kir} $n = 8$). Graphs represent the mean $\pm$ SEM, each dot corresponding to a single

value: *P<0.05, **P<0.01, ***P<0.001. Scale bars, 100 μm.

Figure 2. Correct spontaneous activity in the dLGN is necessary for the invasion of

CTAs. (A) Coronal sections of *Golli-tau-eGfp* in control and *Th^{Kir}* mice at P4 showing

the expression of vGlut1 in the dLGN. Insets show GFP immunostaining. (B)

Quantification of the relative dLGN area occupied by vGlut1 in control and *Th^{Kir}* mice

at different postnatal stages (P4 control *n* = 5, *Th^{Kir}* *n* = 5; P6 control *n* = 5, *Th^{Kir}* *n* = 5;

P9 control *n* = 5, *Th^{Kir}* *n* = 3). (C) Coronal sections of control and *Th^{Kir}* mice at P9

showing vGlut1 and Gfp expression in CTAs. A lentivirus expressing *Gfp* was injected

into V1. (D) Coronal sections of *Golli-tau-eGfp* control and *Th^{Kir}* mice at P0 showing

the expression of Gfp in CTAs (see arrowheads) in the dLGN (control *n* = 3, *Th^{Kir}* *n* =

6). Graphs represent the mean ± SEM and each dot correspond to a single value:

*P<0.05, **P<0.01, ***P<0.001. Scale bars, 100 μm.

Figure 3. Significant increase in cell death in the dLGN of Th^{Kir} mice. (A)

Quantification of the dLGN area in controls and Th^{Kir} mice (P0 control $n = 4$, Th^{Kir} $n =$

6; P3 control $n = 3$, Th^{Kir} $n = 5$; P4 control $n = 4$, Th^{Kir} $n = 4$; P6 control $n = 5$, Th^{Kir} $n =$

5). (B) Coronal sections of control and Th^{Kir} mice showing caspase3a positive cells and

mCherry fused to Kir2.1 (labelled by RFP immunostaining) at P0, P3, P4 and P6. Note

that no caspase3a positive cells were found in P0 Th^{Kir} mice. (C) Quantification of

caspase3a positive cells in the dLGN of control and Th^{Kir} mice at P3, P4 and P6 (P3

control $n = 3$, Th^{Kir} $n = 5$; P4 control $n = 4$, Th^{Kir} $n = 4$; P6 control $n = 5$, Th^{Kir} $n = 5$).

The graphs represent the mean ± SEM and each dot corresponds to a single value:

*P<0.05, **P<0.01, ***P<0.001. Scale bars, 100 μm.

Figure 4. RTAs correctly invade the dLGN of Th^{Kir} mice. (A) Scheme of the

experimental design used to label CTAs and RTAs in control P3 and P7 mice.

Injections of CTB-555 and CTB-647 were performed in each eye of *Golli-tau-eGFP*

mice at P0, and they were analysed at P3 and P7 by immunostaining for GFP. (B)

Coronal sections of *Golli-tau-eGfp* mice at P3 and P7 showing GFP expression in CTAs

(green), and the contralateral (CTB-647) and ipsilateral RTAs (CTB-555). (C) Scheme

of the experimental design used to label RTAs in control and Th^{Kir} mice at P2. Injection

of CTB-647 was performed in one eye in control and Th^{Kir} mice at P0 and analysed at

P2. (D) *Left*: Coronal sections showing CTB-647 labelling in the contralateral and

ipsilateral retinal axons in the dLGN of control and Th^{Kir} mice at P2. *Right*:

Quantification of the percentage of contralateral and ipsilateral RTAs in the dLGN at P2

(control $n = 3$, Th^{Kir} $n = 4$). The graphs represent the mean ± SEM and each dot

corresponds to a single value. Scale bars, 100 μm.

Figure 5. Innervation of the dLGN by CTAs is dictated by RTAs and thalamic

spontaneous activity plays a permissive role. (A) Scheme of the experimental design

used for *ex vivo* calcium imaging of the dLGN in control and embBE mice at P1. (B)

Left, quantification of the event frequency of spontaneous calcium activity in the dLGN

in controls and embBE mice at P1. *Right*, quantification of the frequency of thalamic

calcium waves in the dLGN of control and embBE mice at P1 (control n = 5, embBE n

= 5). (C) *Upper panel*, quantification of the percentage of active ROIs over 15 minutes

recordings in control and embBE mice. *Lower panel*, raster plot of the number of active

ROIs per frame. (D) Scheme of the experimental design used to reveal L6 cortical axons

in the *Golli-tau-Gfp* mice that were embryonically enucleated at E14.5. (E) Coronal

sections of control and embBE mice showing GFP immunostaining in CTAs at P1, P3

and P7. (F) Quantification of the percentage of the dLGN area occupied by GFP-

positive axons (P1: control $n = 3$, embBE $n = 3$; P3: control $n = 4$, embBE $n = 6$; P7

control $n = 5$, embBE $n = 5$). (G) Scheme of the experimental design used to reveal

CTAs in embBE Th^{Kir} mice. Tamoxifen was administered at E10.5 and embBE was

performed at E14.5 in control and Th^{Kir} mice. (H) Coronal sections showing vGlut1

immunostaining in the dLGN in control ($n= 5$), Th^{Kir} ($n= 3$), embBE ($n= 5$), and embBE

Th^{Kir} ($n= 4$) mice at P7. (I) Quantification of the relative vGlut1 immunostaining in the

dLGN of control, Th^{Kir}, embBE and embBE Th^{Kir} mice at P7. Note that CTAs invade

the dLGN of the Th^{Kir} embBE mice. Graphs represent the mean $\pm$ SEM and each dot

correspond to a single value: *$P<0.05$, **$P<0.01$, ***$P<0.001$. Scale bars, 100 μm.

Figure 6. Eye-opening recues the entry of CTAs in *Th^{Kir}* mice. (A) *Left:* Coronal

sections from P10 control and *Th^{Kir}* mice showing vGlut1 immunostaining in the dLGN

at P10, before eye opening. *Right:* Quantification of the relative dLGN area occupied by

vGlut1 labelling at P10 (control *n* = 7, *Th^{Kir}* *n* = 6). (B) *Left:* Coronal sections from P13

control and *Th^{Kir}* mice showing vGlut1 immunostaining at P13, after eye opening.

Right: Quantification of the relative dLGN area occupied by vGlut1 labelling at P13

(control *n* = 5, *Th^{Kir}* *n* = 5). (C) Scheme of the experimental design used for the dark

rearing (DR) experiments. Control and *Th^{Kir}* mice were place in a dark room from P5 to

P15. (D) Coronal sections showing the vGlut1 immunostaining in the dLGN of control,

control DR, *Th^{Kir}* and *Th^{Kir}* DR mice. (E) Quantification of the relative dLGN area

occupied by vGlut1 labelling in these four conditions: Control (*n* = 3), *Th^{Kir}* (*n* = 3),

control DR (*n* = 5), and *Th^{Kir}* DR (*n* = 6). The graphs represent the mean ± SEM and

each dot corresponds to a single value: *P<0.05, **P<0.01, ***P<0.001. Scale bars,

100 μm.

Supplementary Figures

Figure S1. Distribution of CTAs and vGlut1 expression in the dLGN. Coronal

sections showing CTAs labelled by *Golli-tau-eGfp* and immunostaining against vGlut1

in the dLGN of P3 (*n* = 6), P6 (*n* = 3) and P7 (*n* = 3) mice. CTAs start to invade the

dLGN around P3. Scale bars, 100 μm.

Figure S2. The density of dLGN cells remains normal in *Th^{Kir}* mice. (A) Coronal

sections showing Nissl staining of control and *Th^{Kir}* mice at P4. (B) Quantification of

the data shown in A. No significant changes in cell density were observed between the

control ($n = 4$) and *Th^{Kir}* mice ($n = 4$). The graph represents the mean ± SEM and each

dot correspond to a single value: *P<0.05, **P<0.01, ***P<0.001. Scale bars, 100 μm.

Figure S3. Early CTA invasion of the dLGN in embBE mice. (A) *Left*, Coronal

sections of the dLGN showing CTAs labelled with GFP in control and embBE *Golli-*

tau-eGfp mice at P2. *Right*, Quantification of the data shown on the left (control *n* = 4,

embBE *n* = 4). (B, C and D) *Left*, Coronal sections of the dLGN showing vGlut1

immunostaining in control and embBE mice at P2 (control *n* = 3, embBE *n* = 3), P3

(control *n* = 3, embBE *n* = 3) and P7 (control *n* = 3, embBE *n* = 3). *Right*,

Quantification of the data shown in the images on the left. Note that in all embBE mice

a higher percentage of CTAs invaded the dLGN. The graphs represent the mean ± SEM

and each dot corresponds to a single value: *P<0.05, **P<0.01, ***P<0.001. Scale bars,

100 μm.

Figure S4. Normal pattern of dLGN invasion by CTAs after eye opening. (A) *Left*,

Coronal sections showing vGlut1 immunostaining in control and Th^{Kir} mice at P30.

Right, Quantification of the dLGN area occupied by vGlut1 expression at P30 (control

$n=$ 3, Th^{Kir} $n=$ 3). No significant differences were found between both conditions. (B)

Left, Coronal sections showing vGlut1 immunostaining in control and Th^{Kir} mice at P14.

Right, Quantification of the dLGN area occupied by vGlut1 expression at P14 (control

$n=$ 3, Th^{Kir} $n=$ 3). No significant differences were found between both conditions. The

graphs represent the mean $\pm$ SEM and each dot corresponds to a single value. Scale

bars, 100 μm.

Materials and Methods

Mouse strains

Wild type mice maintained on an ICR/CD-1 background, were used for the *in vivo* experiments, tracing studies and Ca^{2+} imaging. The day on which the vaginal plug was detected was designated as E0.5. The $R26^{Kir2.1-mCherry}$ mouse line has been described previously (Moreno-Juan et al, 2017) and it was crossed with an inducible thalamic-specific $Gbx2^{CreER}$ line (L. Chen, Guo, & Li, 2009) to generate $Gbx2^{CreER/+}$; $R26^{Kir2.1-mCherry}$ double mutant embryos, referred to as Th^{Kir}. Tamoxifen induction of Cre recombination in the double mutant embryos was performed by gavage administration of tamoxifen (5 mg dissolved in corn oil: Sigma) at E10.5 to specifically target the principal sensory thalamic nucleus. The *Golli-tau-Gfp* mouse line was described previously (Jacobs, 2007) and was crossed with the Th^{Kir} animals. Mutants of either sex were analyzed randomly at stages from P0 to P30. is

Histology

For immunohistochemistry, mice were perfused with 4% paraformaldehyde (PFA) in PBS (0.01M), and their brains were removed and post-fixed overnight in the same fixative. Immunohistochemistry was performed on 80 μm vibratome brain sections that were first incubated for 1h at room temperature in a blocking solution containing 1% BSA (Sigma), 2% Donkey Serum (Biowest) and 0.25% Triton X-100 (Sigma) in PBS 0.01M. Subsequently, the sections were incubated overnight at 4 °C with the primary antibodies: guinea-pig anti-vGlut1 (1:5000, Synaptic Systems, #135304), chicken anti-GFP (1:3000; Aves Labs, #Gfp-1020) and rat anti-RFP (1:1000 Chromotek, #5F8), rabbit anti-Capasa3a (1:150; Abcam Ab2302). The sections were then rinsed three times in PBS (0.01M) and incubated for 2h at room temperature with secondary antibodies:

Alexa-546 donkey anti-rabbit (1:500, ThermoFisher, #A10040), Alexa-488 goat anti-

chicken (1:500, ThermoFisher, #A11039), Alexa-594 donkey anti-rat (1:500,

ThermoFisher, #A21209), Alexa-488 donkey anti-rabbit (1:500, ThermoFisher,

#A21206), Alexa-647 donkey anti-Guinea Pig (1:500, ThermoFisher). Finally, the

sections were counterstained with the fluorescent nuclear dye DAPI (Sigma-Aldrich).

Tracing of retinal axons

For axonal tracing at postnatal stages, P0 pups were anesthetized under hypothermia.

An incision was made over the eyelid of one or both eyes and they were injected with

CTB conjugated to Alexa-647 or Alexa-594 (Mol. Probes C34778 or C22842) using a

nanoliter injector (WPI). Two, three or seven days later, the animals were perfused with

4% PFA in PBS (0.01M), and their brain was dissected out and post-fixed overnight in

the same fixative. Vibratome sections (80 μm) were then counterstained with the

fluorescent nuclear dye DAPI (Sigma-Aldrich).

In utero **enucleation**

For *in utero* enucleation, pregnant females (E14.5) from wildtype colonies or Th^{Kir} mice

were administered tamoxifen at E10 and deeply anesthetized with isoflurane to perform

laparotomies. The embryos were exposed and both eyes were cauterized in half of the

embryos of each litter. The surgical incision was closed and the embryos were allowed

to develop until postnatal stages.

Calcium imaging in slices

The brains of postnatal mice (P0 and P1) were used for calcium imaging experiments,

obtained after they were sacrificed by decapitation. The brains were immediately

dissected out and 300 μm coronal vibratome sections (VT1200 Leica Microsystems

Germany) were obtained in slicing solution (95% O_2 and 5% CO_2) containing (mM):

2.5 KCl, 7 $MgSO_4$, 0.5 $CaCl_2$, 1 NaH_2PO_4, 26 $NaHCO_3$, 11 glucose and 228 sucrose.

The slices were placed at room temperature in standard aCSF (119 NaCl, 5 KCl, 1.3

$MgSO_4$, 2.4 $CaCl_2$, 1 NaH_2PO_4, 11 glucose) in an atmosphere of 95% O_2 and 5% CO_2.

To load the calcium indicator, the slices were incubated at 37 ºC for 30 min in

artificial CSF (aCSF) containing 5 μl of Cal520 AM (AAT Bioquest), 1 mM DMSO

and 20% pluronic acid. The slices were then placed in a recording chamber with 90%

O_2 and 5% CO_2 at 34 ºC and recorded under a Leica DMi8 microscope. We acquired

images with Hamamatsu ORCA-Flash 4.OLT camera every 300 ms for 15 minutes. All

the calcium imaging data was analyzed as described extensively elsewhere (Moreno-

Juan et al., 2017; Antón-Bolaños et al., 2019).

***In vivo* extracellular recordings**

P6 mice were anesthetized with isoflurane (2.5 %), their scalp was removed and the

skull was cleaned. A 3D-printed holder was glued to the skull with cyanoacrylate

adhesive and dental cement, and attached to a stereotaxic apparatus. A 2 x 2 mm

craniotomy was made over the left hemisphere leaving the dura mater intact. The

craniotomy revealed a window above the area of the dLGN (2.0-2.2 mm anterior to

lambda and 2.0 mm from the midline) allowing the perpendicular insertion of the multi-

channel electrodes. During the recordings, the mice were kept warm (36-37 °C) with a

heated blanket and lightly anesthetized with isoflurane (0.5%). Multi-unit activity

(MUA, >300Hz) was recorded with a linear 16 channel electrode separated by 50 μm

(E16+R-50-S1-L6NT, ATLAS). Shank trajectories were stained with DiI (1,1′-

dioctadecyl 3,3,3′,3′-tetramethylindocarbocyanine perchlorate: Invitrogen) diluted in

70% alcohol. Probes were inserted perpendicularly, 2800-3000 µm deep for dLGN

recordings. The electrical signal activity was sampled at 20 kHz with a filter amplifier

and visualized using MC_RACK software (Multi Channel Systems). After the

recording sessions, the animal's brain was dissected out and fixed overnight in 4% PFA.

In order to determine the position of every channel, 100 µm thick coronal sections were

counterstained with the fluorescent nuclear dye DAPI (Sigma-Aldrich) and the data was

analyzed using Matlab^{TM} routines.

Dark reared experiments

729   For the dark reared experiments, Th^{Kir} animals and their control littermates were put

730   into a completely dark environment with their mother from P5 to P15. After this time

731   the animals were rapidly perfused with 4% PFA in PBS (0.01M). The animal's brain

732   was dissected out and post-fixed overnight in the same fixative. Vibratome sections (80

733   µm) were then subjected to immunohistochemistry.

Data analysis and statistics

The analysis of the calcium imaging experiments was described in detail elsewhere

(Moreno-Juan et al., 2017; Antón-Bolaños et al., 2019). ImageJ software was used to

quantify CTA and RTA invasion of the dLGN. Coronal slices (80 µm) were obtained

from the animals subjected to the different conditions and immunostained for vGlut1 or

GFP. At least three dLGN sections were analyzed from each animal, correcting for the

fluorescence background. Threshold images of the fluorescence were analyzed by

ImageJ software, analyzing the particles of the pixels that cover the dLGN and

measuring the proportion of pixels occupied by CTAs in the dLGN.

Caspase3a positive cells were quantified in coronal sections (80 μm) and normalized

to the area of each dLGN. The whole dLGN was measured from P3, P4 and P6 control

and Th^{Kir} mice, and the results were normalized to the mean of the controls,

representing the fold-change in Caspase3a positive cells in the Th^{Kir} mice.

Statistical analysis was carried out in GraphPad Prism6™ and Matlab™, and the

data are presented as the mean and SEM. Statistical comparison between the groups was

performed using an unpaired two-tailed Student's *t* test and when the data failed a

Kolmogorov-Smirnov or a Shapiro Wilk test of normality, a two-tailed, non-parametric

Mann-Whitney U-test was used. A Two-way ANOVA test was performed on the

embryonic enucleation in Th^{Kir} mice and for the DR experiments. No statistical methods

were used to predetermine the sample size, but our sample sizes are considered

adequate for the experiments and consistent with the literature (Brooks et al., 2013;

Grant et al., 2016; Seabrook et al., 2013). The mice were not randomized.

Figure 1

Figure 1D Event frequency: control 0.27 ± 0.03, $n = 7$; Th^{Kir} 0.02 ± 0.005, $n = 5$

(***P<0.001 Two-tailed student's *t*-test). Wave frequency in dLGN: control $0.34 \pm$

0.04, $n = 7$; Th^{Kir} 0 ± 0, $n = 5$ (**P=0.0025, Mann-Whitney *U*-test).

Figure 1H Firing rate dLGN: control 0.16 ± 0.05, $n = 6$; Th^{Kir} 0.07 ± 0.12, $n = 8$

(*P=0.04 Mann-Whitney *U*-test).

Figure 2

Figure 2B vGlut1 at P4: control $10 \pm 1\%$, $n = 5$; Th^{Kir} $1.67 \pm 0.34\%$, $n = 5$ (***P<0.001

Two-tailed student's *t*-test). At P6: control $35.96 \pm 4.47\%$, $n = 5$; Th^{Kir} $1.24 \pm 0.51\%$, n

$= 5$ (***P<0.001, Two-tailed student's *t*-test). At P9: control $62.4 \pm 4.49\%$, $n = 5$; Th^{Kir}

$12.15 \pm 0.74\%$, $n = 3$ (***P<0.001, Two-tailed student's *t*-test).

Figure 3

Figure3A dLGN size at P0: 84326 ± 2872, $n = 4$; Th^{Kir} 83720 ± 2498, $n = 6$ (P=0.88,

Two-tailed student's t-test). At P3: 89800 ± 4558, $n = 3$; Th^{Kir} 80403 ± 2300, $n = 5$

P=0.14, Mann-Whitney U-test. P4: 118410 ± 6262, $n = 4$; Th^{Kir}: 79748 ± 4351, $n = 4$

(*P=0.02, Two-tailed student's t-test). At P6: 193136 ± 5603, $n = 5$; Th^{Kir} 92022 ±

3380, $n = 5$ (***P<0.001, Two-tailed student's t-test).

Figure 3C Caspase3A$^+$ cells normalized to the control data at P3: control 100 ± 14.97%,

$n = 3$; Th^{Kir} 816.3 ± 122.3%, $n = 5$ (*P=0.03, Mann-Whitney U-test). At P4 control 100

± 30.98%, $n = 4$; Th^{Kir} 684.4 ± 139.2%, $n = 4$ (*P=0.028, Mann-Whitney U-test). At P6

control 100 ± 7.71%, $n = 6$; Th^{Kir} 762.6 ± 87.69%, $n = 6$ (***P<0.001, Two-tailed

student's t-test).

Figure 4

Figure 4D Percentage of RTAs in dLGN at P2. Contralateral RTAs in dLGN, control:

69.94 ± 1.56%, $n = 3$; Th^{Kir}: 66.5 ± 1.74%, $n = 4$. P=0.22 Two-tailed student's t-test.

Ipsilateral RTAs in dLGN control: 20.45 ± 2.27%, $n = 3$; Th^{Kir}: 24.34 ± 2.3%, $n = 4$.

P=0.29 Two-tailed student's t-test.

Figure 5

Figure 5B Event frequency: control 0.08 ± 0.02, $n = 5$; embBE 0.23 ± 0.02, $n = 5$

(***P<0.0005, Two-tailed student's t-test). Wave frequency in dLGN: control 0.08 ±

0.03, $n = 5$; embBE 0.31 ± 0.03, $n = 5$ (*P=0.01, Mann-Whitney U-test).

Figure 5F % of CTAs in the dLGN at P1: control 3.26 ± 0.2%, $n = 3$; embBE 8.85 ±

1.44%, $n = 3$ *P=0.018, Two-tailed student's t-test). At P3: control 12.86 ± 1.59%, $n =$

4; embBE 53.92 ± 5.05%, $n = 6$ (***P<0.001, Two-tailed student's t-test). At P7:

control 43.89 ± 2.46%, $n = 4$; embBE 74.82 ± 1.68%, $n = 5$ (*P=0.01, Mann-Whitney

U-test).

Figure 5I % of vGlut1 in the dLGN at P7: control n = 5, embBE n= 5, Th^{Kir} n=3, Th^{Kir}

embBE n= 4. A two-way ANOVA test with Tukey's post hoc analysis indicates that the

interaction between all the conditions are significant, comparisons: control vs Th^{Kir} ***

P<0.001; control vs embBE *** P<0.001; control vs Th^{Kir} embBE *P=0.01; Th^{Kir} vs

embBE *** P<0.001; Th^{Kir} vs Th^{Kir} embBE *** P<0.001; embBE vs Th^{Kir}embBE **

P=0.0025.

Figure 6

Figure 6A %vGlut1 in the dLGN of P10: control 79.28 ± 2.31%, n = 7; Th^{Kir} 18.85. ±

1.85%, n = 6 (***P<0.001, Two-tailed student's t-test).

Figure 6B %vGlut1 in the dLGN of P13: control 76.62 ± 1.23%, n = 5; Th^{Kir} 76.69. ±

0.8%, n = 5 (P=0.96, Two-tailed student's t-test).

Figure 6E % of vGlut1 in the dLGN at P15: control n = 3, control DR n= 5, Th^{Kir} n=3,

Th^{Kir} DR n= 6. A two-way ANOVA test with Tukey's post hoc analysis, comparisons:

control vs Th^{Kir} P= 0.97; control vs control DR P= 0.99; control vs Th^{Kir} DR

***P<0.005; Th^{Kir} vs control DR P= 0.99; Th^{Kir} vs Th^{Kir} DR *** P<0.001; control DR vs

Th^{Kir} DR *** P<0.001.

Figure S2

Figure S2B Nissl$^+$ cells/100 μm^2: control 110.5 ± 4.29, n = 4; Th^{Kir} 124.8 ± 6.34%, n =

4 (P=0.11. Mann-Whitney U-test).

Figure S3

Figure S3A %CTAs in the dLGN of P2 mice: control 7.28 ± 2.24%, n = 4; embBE 39.3

± 3.1%, n = 4 (*P=0.02, Mann-Whitney U-test).

Figure S3B % of vGlut1 in the dLGN at P2: control 7.1 ± 9.92%, n = 3; embBE 44.32±

2.35%, n = 3 (***P<0.001, two-tailed student's t-test).

Figure S3C % of vGlut1 in the dLGN at P3: control $10.68 \pm 1.92\%$, $n = 3$; embBE

$53.06 \pm 4.41\%$, $n = 3$ (***P<0.001, two-tailed student's t-test).

Figure S3D % of vGlut1 in the dLGN at P7: control $39.01 \pm 3.03\%$, $n = 3$; embBE

$71.45 \pm 1.11\%$, $n = 3$ (***P<0.001, two-tailed student's t-test).

Figure S4

Figure S4A %vGlut1 at P30: control $87.26 \pm 3.36\%$, $n = 3$; Th^{Kir} $87.16. \pm 3.28\%$, $n = 3$

(P=0.98, two-tailed student's t-test).

Figure S4B %vGlut1 at P14: control $93.17 \pm 1\%$, $n = 3$; Th^{Kir} $89.31. \pm 1.16\%$, $n = 3$

(P=0.1, two-tailed student's t-test).