Isotropic 3D electron microscopy reference library of whole cells and tissues

C. Shan Xu[1]†, Song Pang[1]*, Gleb Shtengel[1]*, Andreas Müller[2,3,4], Alex T. Ritter[5], Huxley K. Hoffman[6#], Shin-ya Takemura[1], Zhiyuan Lu[1], H. Amalia Pasolli[1§], Nirmala Iyer[1], Jeeyun Chung[7,8], Davis Bennett[1], Aubrey V. Weigel[1], Tobias C. Walther[7,8,9], Robert V. Farese, Jr.[7,8], Schuyler B. van Engelenburg[6], Ira Mellman[5], Michele Solimena[2,3,4,10], Harald F. Hess[1]†

Affiliations

[1]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147, USA

[2]Molecular Diabetology, University Hospital and Faculty of Medicine Carl Gustav Carus, TU Dresden, Dresden, Germany

[3]Paul Langerhans Institute Dresden (PLID) of the Helmholtz Center Munich at the University Hospital Carl Gustav Carus and Faculty of Medicine of the TU Dresden, Dresden, Germany

[4]German Center for Diabetes Research (DZD e.V.), Neuherberg, Germany

[5]Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA

[6]Molecular and Cellular Biophysics Program, Department of Biological Sciences, University of Denver, Denver, CO 80210, USA

[7]Department of Molecular Metabolism, Harvard T.H. Chan School of Public Health, Boston, MA

02115, USA

[8]Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA

[9]Howard Hughes Medical Institute, Boston, MA 02492, USA; Broad Institute of Harvard and

MIT, Cambridge, MA 02124, USA

[10]Max-Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany

†Correspondence to: xuc@janelia.hhmi.org (C.S.X), hessh@janelia.hhmi.org (H.F.H.)

*These authors contributed equally to this work

#Present Address: Department of Cell and Developmental Biology, University of Colorado

Anschutz Medical Campus, Aurora, CO 80045, USA

§Present Address: Electron Microscopy Resource Center, The Rockefeller University, New York,

NY 10065, USA

Summary

Understanding cellular architecture is essential for understanding biology. Electron microscopy (EM) uniquely visualizes cellular structure with nanometer resolution. However, traditional methods, such as thin-section EM or EM tomography, have limitations inasmuch as they only visualize a single slice or a relatively small volume of the cell, respectively. Here, we overcome these limitations by imaging whole cells and tissues with enhanced Focus Ion Beam Scanning Electron Microscopy (FIB-SEM) in high resolution with month-long acquisition duration. We use this approach to generate reference 3D image datasets at 4-nm isotropic voxels for ten different examples, including cultured cells (cancer, macrophages, and T-cells) as well as tissues (mouse pancreatic islets and the *Drosophila* fan-shaped body). We open access to all datasets in [OpenOrganelle](), an interactive web platform that allows accessing both the original 3D EM data, and subsequent organelle segmentation. Together, these data will serve as a reference library to explore comprehensive quantification of whole cells and their constituents, thus addressing questions related to cell identities, cell morphologies, cell-cell interactions, as well as intracellular organelle organization and structure.

Main

Individual cells and organized systems of cells within a tissue can be described with a hierarchy of ultrastructural details, from complex organelle networks down to single protein molecules[1]. While transmission electron microscopy (TEM) and scanning electron microscopy (SEM) images from diamond-knife cut thin sections of metal stained samples have profoundly shaped our understanding of organelles[2], such approaches generate 2D images that can reveal 3D

structures, but may be limited due to constraints from the section thickness (of ~30–300 nm),

allowing only a small fraction of the whole cell to be captured. Furthermore, that sampled

fraction is often viewed at a specific sectioning angle through the cell or tissue, limiting visibility

of many structural aspects unless many sections are examined. Series of sections can be

digitally stitched together to form a 3D volume, but the 30 nm or larger sampling interval from

section to section sacrifices much structural detail. Tilt tomography can overcome this problem

but is confined in volume size to ~ 100 μm^3. Stitching together data of multiple such sections

can extend the effective volume size but is arduous and currently not practical for more than a

dozen sections. Alternatively, diamond-knife cut surfaces of a block face sequentially imaged

with SEM[3] was a pioneering step to access larger volumes, but its z resolution is limited to 25

nm due to electron radiation damage to the block face from beam scanning that prevents

consistent cutting of thinner slices[4]. Thus, high-resolution isotropic 3D whole cell data remain

mostly out of reach.

The constraints of diamond-knife cutting thickness can be removed by using a focused

ion-beam, typically of gallium ions, to abrade arbitrarily fine increments of the block surface as

thin as 1 nm. This technique, combined with SEM, is called FIB-SEM. Here, we advance FIB-SEM

with a 90° angle between the imaging electron beam and the milling ion beam (Fig. 1a), along

with other modifications to transform conventional FIB-SEM that is lacking of long-term

reliability to an enhanced platform that is capable of months to years of continuous imaging

without defects in the final image stack[5,6,7]. We further systematically optimized both ion

milling and electron imaging components to achieve 4-nm sampling intervals in all three

dimensions, thus enabling high-resolution imaging by rendering greater details with volumes

approaching $(100 \ \mu m)^3$. The ability of enhanced FIB-SEM to image whole cells and tissues at 4-

nm isotropic voxels over large volumes makes it an ideal tool to map *in toto* the 3D

ultrastructural relationship of organelles and protein complexes in living systems. The resulting

data are unique in their size and level of 3D detail, and are presented for a variety of cells and

tissues to establish a reference library of cellular architecture.

Finer Resolution

With the dual biological motivations to image with fine resolution *and* encompass the largest

possible volumes, we confront limitations and tradeoffs between these two goals. The isotropic

resolution limits are convoluted by 1) the waist size of the incoming primary electron beam,

and 2) the size of the scattering volume explored by the penetrating primary electrons. To

optimize for point 1, the best beam focus, spherical and chromatic aberrations of the

electromagnetic lenses can be mitigated by operating at modest numerical aperture through

smaller beam currents hence achieving the best x-y resolution. For point 2, lower primary beam

landing energy generates fewer scattering events, resulting in smaller scattering volume which

improves resolutions along all three axes. Any further reduction below the optimal energy will

result in a reduced scattering ratio of the heavy metal stain atoms over lighter background

atoms of the sample resin, thus fading contrast.

Through investigating a multi-dimensional parameter space, we found that a beam

current of 200 to 300 pA and a landing energy of 700 to 900 eV are optimal for isotropic 4 nm

imaging. To characterize the resolution in x-y and z, we analyzed the step transitions at the

edges of gold nano particles on a carbon substrate. The vertical lines in Fig. 1b indicate an x-y

transition distance of 2.0 nm for SEM signal change from 37% to 63%, or of 5.5 nm for

transition from 20% to 80%. One can use ribosomes in a similar way as an *in situ* resolution

standard to get the full 3D edge transition. The ribosome x-y transitions are about 30% larger

than the gold nano particle x-y edges, probably from it being a non-ideal step edge. As an

alternative conservative measure of resolution, this corresponds 12 nm Gaussian full width half

maximum (FWHM) in x-y plane. In addition, Fig. 1b shows mostly isotropic resolution on a

number of different samples where the z transitions are 30% larger than the x-y edge

transitions. The procedure for estimating the resolution and some examples are presented in

Methods Section.

Larger Volumes

Large volumes are crucial to study biology, driven by demands of completeness and sufficient

statistics. In acquiring the largest possible volumes with the finest resolution, we allocate 6,000

to 10,000 electrons per voxel (depending on sample contrast) to achieve a target contrast

(between membrane and resin) to shot noise ratio of three or higher. This translates to a

sampling rate of < 200 kHz (dwell time > 5 µs) needed with a 200-pA primary beam. We set the

sampling voxel size to 4 nm to be consistent with the resolution determined from edge

transitions. These values define a volumetric imaging rate of < $(10 \ \mu m)^3$/day. The continuous

operating duration of conventional FIB-SEM systems is constrained to ~3 days, which inevitably

caps the maximum possible 3D volume to $(\sim15 \ \mu m)^3$.

To access volumes of typical whole cells either in culture or in tissue, reliable long-term

imaging approaching one month with nanometer stability is required. This is particularly

challenging when the highest resolution and the largest possible volumes are desired. Fig. 1c,

places this capability in context of other existing 3D EM methods where bottom axis is sample

size and vertical axis is isotropic voxel size. Note that to access very large volumes such as the

whole *Drosophila* brain of 40×10^6 μm^3, a higher current with lower resolution of 8 nm voxels

was used (Fig. 1c). The enhanced FIB-SEM technology, initially motived by the *Drosophila*

connectome project[8] and demonstrated preliminary value to cell biology[9], delivers continuous

stable and reliable operation for months to years, which is also optimized for fine resolution

whole cell imaging.

Open Data

Access to this new region of large volumes with high resolution imaging requires a considerable

investment in technology development, equipment resources, and imaging duration, but the

resulting wealth of data is unique and comprehensive, covering whole cells with all their

organelles and proteins that can be highlighted with heavy metal stains. Here, we provide ten

such datasets containing seven common wild-type cultured cell samples and three different

tissue samples. To enable the wider scientific community to browse, download, and explore, we

open access the data via OpenOrganelle (https://openorganelle.janelia.org)[10]. For each sample,

metadata and pre-selected views of interest are provided, while cross-sections can be explored

with *neuroglancer*, a built-in visualization tool. This repository also includes further processing,

such as segmentation to categories of interest[10], which forms a basis for quantifying numbers,

sizes, shapes, contacts, locations, proximity or any other parameters and distributions of EM

stained organelles and proteins in the whole cell. While these datasets will serve as the starting

point of a reference library to explore comprehensive cellular architecture, we hope to further

extend such reference datasets to yet other cells and tissues.

Discussion of Exemplary Samples

The initial datasets of seven common wild-type cultured cells and three tissue samples are

tabulated in Table 1. The cells have been cryo-fixed by high pressure freezing (HPF), which

vitrifies a sample on a millisecond time scale to best preserve any dynamic structural details.

This also minimizes any possibility of artifacts that might be of concern with chemical fixation.

The subsequent freeze substitution labels biomolecules with heavy metals to provide contrast

for EM. While thin tissue samples (< 200-µm-thick) can be similarly cryo-fixed, in many cases

thick tissue specimens cannot be cryo-fixed (e.g., large brain tissue) thus requiring various

chemical fixation and staining protocols. We also offer a selection of these tissue samples as

initial reference datasets. The 3D images all used 4 nm voxels sampled at rates of 100 kHz – 2

MHz with ~ 250 pA of primary beam current. The samples with stronger staining were imaged

at much faster rates while maintaining similar image quality. The time to acquire these volumes

ranged from 2 weeks to 5 weeks of uninterrupted imaging.

Individual HeLa Cell

HeLa cells, human cervical cancer cells, are easily cultured and widely used in cell biology

laboratories as a basic model to test diverse hypotheses. Having such cells imaged in their

entirety can provide a reference to which perturbations in growth, genetic, environment, etc.

can be compared. Fig. 2 presents a typical 200 GB 3D dataset on the example of an entire HeLa

cell. The mitochondrial network is shown in green. The bottom row panels show 2D cross-

sections of standard cellular organelles, such as the centrosome (Fig. 2b), the Golgi apparatus

(Fig. 2c), and nuclear envelope (Fig. 2d). Notably, every example illustrates the advantages of

isotropic 3D imaging for cell biology. No single 2D cross-section allows visualizing all centriole

sub-distal appendages, however quick segmentation of the 3D dataset shows them clearly (red,

Fig. 2b). Stereotypical 2D images of the Golgi stacks do not reveal the fenestration details and

long thin tubular extensions, that can only be seen in 3D (magenta – single segmented Golgi

cisterna – Fig. 2c). Polyribosomal chains on the nuclear envelope are mostly hidden in 2D cross-

sections but easily resolved and detailed in 3D (Fig. 2d).

Interacting Cell Pairs: T Cell Attacking Cancer

Cytotoxic T lymphocytes (CTLs) are immune cells that have the capacity to identify and destroy

virus-infected or cancerous cells. Upon encountering a target, CTLs undergo a dramatic

cytoskeletal rearrangement to form an organized intercellular interface called the

immunological synapse (IS). Signaling at the IS directs the polarized release of toxic proteins

housed in specialized secretory lysosomes called lytic granules (LGs), which results in target cell

death. Due to the important role of CTLs in anti-viral and anti-cancer immunity, the IS has been

the subject of intense scrutiny. Many studies of the IS have been performed on artificial planar

lipid bilayers created on glass coverslips. These synthetic two-dimensional surfaces lack the

complex receptor content and membrane topology of a real cancer cell. Studies of IS formation

between actual cells, although more challenging, present more biologically relevant content.

Limited single slice TEM images of CTL:target conjugates have revealed interesting features of

the IS, but do not represent the full scope of this dynamic, three-dimensional structure[11]. Fig.

3a presents a 1.4-TB 3D dataset (16bit) of a mature murine CTL (green) engaging an ID8 ovarian

cancer cell (cyan). The isotropic high-resolution information of FIB-SEM imaging provides a

unique and complete map of the complex membrane topology at the interface between T cell

and target. Fig. 3b shows one cross section demonstrating the stereotypical CTL "cupping" of

the target cell at the IS. Zoomed in images show the variety of features across this interface

including membrane interdigitation (Extended Data Fig. 1a), flat membrane apposition

(Extended Data Fig. 1b), and filopodia of the target cell trapped between the two cells

(Extended Data Fig. 1c). Without the ability to scan in 3D, these filopodia may be mistaken for

vesicles trapped between T cell and target.

Another hallmark of CTL killing is the polarization of the centrosome toward the target

cell[12,13], which can be observed in Fig. 3c. FIB-SEM allows for the label-free localization of all

organelles in the T cell including those responsible for target killing and cytokine secretion.

Individual 2D EM slices may miss the Golgi stacks and lytic granules (Fig. 3d) which can be found

by scanning the FIB-SEM volume around the centrosome.

Tissues: Mouse Pancreatic Islets and *Drosophila* Brain Fan-shaped Body

Pancreatic islets are micro-organs consisting mainly of beta, alpha, delta, polypeptide cells and

endothelial cells. Among them beta cells are in the majority and secrete insulin stored in

secretory granules (SGs) to maintain blood glucose homeostasis. Beta cells have a polyhedral

shape with a longer axis that can easily reach 20 μm[14], posing significant challenges for high

resolution whole cell imaging. Few beta cells have been reconstructed by serial section electron

tomography at medium resolution[15] to allow assessment of subcellular features such as insulin

SG number and shape of mitochondria, but lacking the resolution for imaging of e.g., ER and

microtubules. Large-scale high-resolution FIB-SEM enables the analysis of ultrastructural

differences between beta cells within an islet as well as features that require higher resolution

such as ribosomes and the cytoskeleton. In our studies, especially the preservation of

cytoskeletal elements benefits from fixation by HPF. Fig. 4a displays a 200-GB volume with

several complete beta cells that are entirely included within the 30 x 20 x 25 μm^3 volume. The

large field of view allows for imaging the interaction of beta cells, e.g., full primary cilia and

their connections to neighboring cells and the intermingling of microvilli (Fig. 4b and 4c). At 4-

nm voxel sampling, we observed ultrastructural differences between insulin SGs (Fig. 4d). For

the first time, we report close contacts of ER and insulin SGs in 3D (Fig. 4e). Ultimately, such

high resolution allows the sharp definition of microtubules, single ribosomes and Golgi stacks

for quantitative measurements (Fig. 4f). The seamless imaging results in precisely aligned stacks

furthermore facilitating automated 3D segmentation. Specifically, from these FIB-SEM datasets

has emerged a comprehensive 3D representation of microtubule networks and how they

interact with other organelles[16]. We could show that beta cell microtubules are arranged into

non-radial networks which for the most part do not originate either from centrioles or

endomembranes. Furthermore, glucose stimulation changes the number and length of

microtubule polymers without changing their overall density. Finally, we found insulin SGs to be

enriched near the plasma in association with microtubules, independent of the extracellular

glucose levels.

The **central complex in insect brains** is a highly conserved brain region that controls

various behaviors. Fig. 4g shows higher-resolution images acquired from the fan-shaped body

of the central complex of the fruit fly, *Drosophila*. For a detailed understanding of brain

functions, it is imperative to identify the synaptic levels of neuron connectivity through key

synaptic elements such as presynaptic T-bars and postsynaptic densities (PSDs) (Fig. 4i, 4j). To

investigate if the central complex contains characteristic synaptic motifs, such as polyadic

synapse, rosette synapse, or any new synaptic motif, validating the synaptic structures with

high-resolution images is crucial. However, often a trade-off between imaging a large volume

with low-resolution and acquiring high-resolution with a small volume limits answering

questions in connectomics, where fine synaptic structure identification and large-scale

neuronal tracing are equally important. Higher resolution FIB-SEM images with large volume

therefore provide significant values to the study. Such datasets allow reconstruction of not only

the large neuronal objects and glial cells, but also smaller intracellular components such as

synaptic vesicles, dense-core vesicles (Fig. 4h), and microtubules (Fig. 4k, 4l). Knowing the types

of synaptic vesicles could add important insights to the understanding of the properties of

synaptic connections. As an example, the terminals of dopaminergic neurons in the fly's

olfactory center were shown to contain both clear and dense-core vesicles[17], in congruence

with having more than one neurotransmitter[18].

Summary

While three-dimensional imaging offers tremendous value to elucidate biological structures and

decipher their function, the enhanced FIB-SEM platform has advanced to image not only partial

but entire mammalian cells or multicellular parts of tissues at 4-nm isotropic voxels, hence

revealing a more holistic 3D ultrastructural picture in living systems. We have imaged a variety

of biological samples over volumes of $(100 \ \mu m)^3$ to study the complex morphologies of cells, the

complete interfaces involved in cell-cell interactions, and the shapes, volumes, distributions,

and relationships of intracellular organelles. This has allowed for new biological insights. For

example, the datasets of mouse pancreatic islets (Fig. 4a–4f) unveil a comprehensive 3D

representation of the microtubule network and how it interacts with other organelles, as well

as microtubule remodeling as a result of glucose stimulation[16]. Likewise, the data of HeLa and

U2-OS nuclei were mined to identify and characterize chromatin domains[19]. In a similar spirit,

we feel that guided by other research questions, many insights can be gleaned from this raw

data through customized analysis.

As we envision many fruitful discoveries to be enabled via access to the 3D data in the

larger context of cells in tissues, we aim to expand our portfolio on OpenOrganelle[10] to include

other tissues. The initial datasets of various healthy wild-type cells, can be explored to define

reference features of interest, such as microtubule proximity to certain vesicles, or ER

morphology near a particular organelle or ribosome chain length on curved ER near the Golgi

apparatus. The possibilities seem limitless and are actually daunting. Open access preview on a

wild-type healthy cell can encourage further experiments to compare such features against

perturbations of disease, mutations, development, or environment. Whole cell structural

differences can be quantified and measured against these perturbations. In additional to target

feature analysis, unanticipated new features could also be discovered. Segmentation is a crucial

next step to aid biologists in such analysis. However, the semi-manual attempt on a HeLa cell

(Fig. 2) is time-consuming. A more rapid approach needed on the whole cell scale is presented

in the associated paper[10]. We believe that open access to rich 3D FIB-SEM datasets will provide

important insights, guidance, and inspiration for further exploration to the research community

to address a new broad range of biological questions.

 Reference

[1] Lodish H. et al. *Molecular Cell Biology* (W. H. Freeman, New York, 2016).

[2] Porter, K. R., Claude, A., & Fullam, E. F. A study of tissue culture cells by electron microscopy: methods and preliminary observations. *The Journal of experimental medicine*, **81**(3), 233–246 (1945). https://doi.org/10.1084/jem.81.3.233

[3] Denk W., & Horstmann H. Serial block-face scanning electron microscopy to reconstruct three-dimensional tissue nanostructure. *PLoS Biol* 2004, **2**:e329 (2004). https://doi.org/10.1371/journal.pbio.0020329

[4] Titze B. Techniques to prevent sample surface charging and reduce beam damage effects for SBEM imaging. Dissertation. Heidelberg University. 1–112 (2013). https://doi.org/10.11588/heidok.00015372

[5] Xu C. S. et al. Enhanced FIB-SEM systems for large-volume 3D imaging. *eLife* **6**:e25916 (2017). https://doi.org/10.7554/eLife.25916

[6] Xu, C. S., Hayworth K. J., & Hess H. F. Enhanced FIB-SEM systems for large-volume 3D imaging. US Patent 10,600,615, 24 Mar (2020).

[7] Xu C. S., Pang S., Hayworth K. J., & Hess H. F. Transforming FIB-SEM Systems for Large-Volume Connectomics and Cell Biology. In: Wacker I., Hummel E., Burgold S., Schröder R. (eds) Volume Microscopy. Neuromethods, vol 155 (2020). Humana, New York, NY. https://doi.org/10.1007/978-1-0716-0691-9_12

[8] Xu C. S. et al. A connectome of the adult *Drosophila* central brain. *bioRxiv* 2020.01.21.911859. (2020). https://doi.org/10.1101/2020.01.21.911859

[9] Wu, Y. et al. Contacts between the endoplasmic reticulum and other membranes in neurons. *PNAS* **114**(24) E4859-E4867 (2017). https://doi.org/10.1073/pnas.1701078114

[10] Heinrich, L. et al. Automatic whole cell organelle segmentation in volumetric electron microscopy. (2020).

[11] Stinchcombe J. C., Bossi G., Booth S., & Griffiths G. M. The immunological synapse of CTL contains a secretory domain and membrane bridges. *Immunity*. 2001;15(5):751–761. https://doi.org/10.1016/s1074-7613(01)00234-5

[12] Kupfer A. & Dennert G. Reorientation of the microtubule-organizing center and the Golgi apparatus in cloned cytotoxic lymphocytes triggered by binding to lysable target cells. *J Immunol*. 1984;133(5):2762-2766.

[13] Stinchcombe J. C., Majorovits E., Bossi G., Fuller S. & Griffiths G.M. Centrosome polarization delivers secretory granules to the immunological synapse [published correction appears in Nature. 2006 Nov 9;444(7116):236]. *Nature*. 2006;443(7110):462–465. https://doi.org/10.1038/nature05071

[14] Fava, E. et al. Novel standards in the measurement of rat insulin granules combining electron microscopy, high-content image analysis and in silico modelling. *Diabetologia* **55**, 1013–1023 (2012). https://doi.org/10.1007/s00125-011-2438-4

[15] Noske, A. B., Costin, A. J., Morgan, G. P. & Marsh, B. J. Expedited approaches to whole cell electron tomography and organelle mark-up in situ in high-pressure frozen pancreatic islets. *J. Struct. Biol.* **161**, 298–313 (2008). https://doi.org/10.1016/j.jsb.2007.09.015

[16] Müller, A. et al. Three-dimensional FIB-SEM reconstruction of microtubule-organelle interaction in whole primary mouse beta cells. *bioRxiv* 2020.10.07.329268. (2020). https://doi.org/10.1101/2020.10.07.329268

[17] Takemura, S. Y. et al. A connectome of a learning and memory center in the adult *Drosophila* brain. *eLife* **6**, e26975 (2017). https://doi.org/10.7554/eLife.26975

[18] Aso, Y. et al. Nitric oxide acts as a cotransmitter in a subset of dopaminergic neurons to diversify memory dynamics. *eLife* **8**, e49257 (2019). https://doi.org/10.7554/eLife.49257

[19] Miron, E. et al. Chromatin arranges in chains of mesoscale domains with nanoscale functional topography independent of cohesion. *Sci. Adv.*, **6**, 39 eaba8811 (2020). https://doi.org/10.1126/sciadv.aba8811

Methods

Cultured cell sample preparation

1. HeLa cells

HeLa cells (CCL-2) were purchased from American Type Culture Collection (ATCC). HeLa cells were maintained in EMEM medium (ATCC, 30-2003) supplemented with 10% FBS (Corning, 35-011-CV) and 1X penicillin-streptomycin solution (Corning, 30-002-CI). Trypsinized HeLa cells were seeded on pre-cleaned and edge-coated sapphire coverslips. Immediately prior to freezing, live cells are inspected to ensure cell morphology and viability. After quality assurance the cells are transferred to a water jacketed CO_2 incubator (ThermoFisher, Midi 40) kept at 37° C, 5% CO_2, and 100% humidity while awaiting freezing. Each coverslip is removed from the incubator immediately prior to the freezing procedure. Cell samples were cryo-fixed using a Wohlwend Compact 2 high-pressure freezer (Engineering Office M. Wohlwend GmbH, Bifig 14, 9466 Sennwald, Switzerland). The BSA mixture is used as a cell cryo-protectant to prevent ice crystal formation in the extra cellular space[20]. Once the sample is frozen it can be stored indefinitely in liquid nitrogen for subsequent freeze-substitution and resin embedding. Freeze-substitution (FS) was performed using automated FS machine (AFS2, Leica Microsystems): coverslips were transferred to cryotubes containing FS media (2% OsO_4, 0.1% Uranyl Acetate (UA), and 3% water in acetone) under liquid nitrogen, followed by a programmed FS schedule. Resin embedding was performed immediately after FS. Samples were removed from the AFS2 machine, washed 3 times in anhydrous acetone for a total of 10 min and embedded in Eponate 12 which was polymerized for 48 hours at 60°C. Following EPON embedding, the coverslip is separated from the resin block containing the cells by sequential immersion in liquid nitrogen

and hot water. Once the coverslip is removed, the exposed surface is immediately re-

embedded in Durcupan resin which helps minimize streaks during FIB-SEM imaging[5]. The detail

procedures of coverslip preparation, high pressure freezing, free substitution, and resin

embedding are presented in the Supplementary Information.

2. Cytotoxic T cell and T cell/Cancer cell conjugate (CTLs)

To generate CTLs from OT-I mice (Jackson Labs), splenocytes were isolated and stimulated with

10nM OVA257-264 peptide (AnaSpec, Fremont, CA, USA) in 10% RPMI (RPMI 1640 plus 10%

fetal bovine serum (Fisher), 2mM L-glutamine, 50U/mL penicillin/streptomycin, and 50µM β-

mercaptoethanol. Following 3 days of stimulation, cells were resuspended in complete media

plus 10 IU/mL recombinant human IL-2 (rHIL-2, Roche), and seeded in fresh media at 0.5×10^6

cells/mL every 48 hours. ID8 cancer cell line (ATCC) was grown in RPMI media (Gibco) with 10%

FBS (Gibco).

In-vitro activated CTLs from Ova-transgenic (OT-I) mice were combined with adherent ID8

murine ovarian cancer cells. To facilitate recognition of cancer cells by the OTI-I CTLs, ID8 cells

suspended in complete medium (RPMI with 10% FBS, Gibco) were incubated with $OVA_{257-264}$

SIINFEKL peptide for 1 hour at 37°C. The cells were washed three times with complete medium,

and 10^5 cells were added to each well of a 24-well plate. Each well contained a single sapphire

coverslip coated with human Fibronectin (Corning). The ID8 cells were allowed to settle and

adhere for 2 hours in an incubator at 37°C/5% CO_2. At this time, the media in the well was

replaced with 250uL phenol-red free RPMI (Gibco). Care was taken to ensure that media and

cells were stored in 37°C/5% CO_2 for all incubation periods. Immediately after changing the

media, 10^5 CTLs in 50uL complete medium were added to the well. The CTLs were allowed 7

minutes to find their targets and secrete lytic granules before fixation through high pressure

freezing. The relatively short incubation time favors capture of CTL:target conjugates in an

early-stage of interaction.

The remaining cryofixation, FS, and resin embedding procedures were identical to those

described for HeLa cells. The detail procedures are presented in the Supplementary

Information.

Tissue sample preparation

1.   Mouse pancreatic islets

Pancreatic islets of 9-week-old C57BL/6 mice were isolated as previously described[21]. They

were cultured overnight in standard culture media (RPMI 1640 (Gibco) with 10% FBS, 20 mM

HEPES, 100 U/ml each penicillin and streptomycin) containing 5.5 mM glucose. Prior to high

pressure freezing the islets were subjected to 1 hr incubation in Krebs-Ringer buffer containing

either 3.3 mM or 16.7 mM glucose.

Islets were frozen with a Leica EM ICE high pressure freezer (Leica Microsystems, Germany)

and kept in liquid nitrogen until freeze substitution. They were substituted as previously

published[22] or according to a novel protocol: first the samples were substituted in a cocktail

containing 2% OsO_4, 1% UA, 0.5% glutaraldehyde, 5% H2O in Acetone with 1% Methanol at -

90°C for 24 hours. The temperature was raised to 0°C over 15 hours followed by 4 washes with

100% acetone for 15 min each and an increase in temperature to +22°C. Afterwards the

samples were incubated in 0.2% thiocarbohydrazide in 80% methanol at RT for 60 min followed

by 6 x 10 min washes with 100% acetone. The specimens were stained with 2% OsO_4in acetone

at RT for 60 min followed by incubation in 1% UA in acetone + 10% methanol in the dark at RT

for 60 min. After 4 washes in acetone for 15 min each they were infiltrated with increasing

concentrations of Durcupan resin in acetone followed by incubation in pure Durcupan and

polymerization at 60°C. For quality control the blocks were sectioned with a Leica LC6

ultramicrotome (Leica microsystems) and 300 nm sections were put on slot grids containing a

Formvar film. Tilt series ranging from -63° to +63° were acquired with a F30 electron

microscope (Thermo Fisher) and reconstructed with the IMOD software package[23].

2.   Drosophila brain samples (fan-shaped body and accessory calyx)

Drosophila brain tissues from 5-day-old adult (Genome type: iso Canton S G1 x w1118 iso 5905)

were prepared according to Progressive Lowering of Temperature and Low Temperature

Staining (PLT-LTS) heavy metal enhancement protocol described previously[24].

In details, after tissue dissection and pre-fixation, we osmicated tissue in 1% OsO_4. This was

followed without washing by 1.5% K ferrocyanide, then a complete wash and finally, a transfer

for 15 min to 1% thiocarbohydrazide at 22°C, followed in turn by a complete wash then 2%

osmium for 30 min at 22°C. After osmication we stained in lead aspartate for 30 min at 55°C,

followed by 1 h at 22°C. The tissue was then dehydrated using the PLT method as in Method 3,

followed by low temperature en bloc staining in either 0.2% uranyl acetate in acetone, or 1%

EPTA in 97% ethanol. Specimens were infiltrated and embedded in Durcupan.

Preparation for FIB-SEM

After Durcupan re-embedding, a 3D X-Ray tomogram of the entire block is taken using an

XRadia-510 Versa micro X-Ray system (Carl Zeiss X-ray Microscopy, Inc.). The 3D X-Ray

tomograms allow for selecting Region of Interest (ROI) efficiently. Specifically, it identifies cells

with good morphology (for example a properly shaped cell, a cell in a desired stage of mitosis, a

357 T-cell attacking a cancer cell, etc.), and also determines the proper orientation for the tissue

358 samples. Once a potential ROI is identified, the sample block is re-mounted to the top of a 1

359 mm copper post (using Durcupan) which was in contact with the metal-stained sample for

360 better charge dissipation, as previously described[5]. Each sample is oriented on the copper post

361 to allow the shortest distance along FIB milling direction. A small vertical sample post is

362 trimmed to the ROI with a width of ~100 µm and a depth of 60 ~ 80 µm in the direction of the

363 ion beam for each sample. Sample trimming uses an ultramicrotome (EM UC7, Leica

364 Microsystems), is done in few iterations with X-Ray tomogram collected after every step.

365  Once the desired ROI is trimmed, a thin layer of conductive material of 10-nm gold

366 followed by 100-nm carbon was coated on the trimmed samples using a Gatan PECS 682 High-

367 Resolution Ion Beam Coater. The coating parameters were 6 keV, 200 nA on both argon gas

368 plasma sources, 10 rpm sample rotation with 45-degree tilt.

369 FIB-SEM imaging

370 A Zeiss Gemini 500-Capella Crossbeam system and Zeiss Merlin crossbeam system[5,6] were

371 further modified for this work in the following ways: the FEI Magnum FIB column was replaced

372 with a Zeiss Capella FIB column; the Zeiss Capella FIB column was repositioned at 90 degrees to

373 the SEM column; and the custom NI LabVIEW control software was re-written to switch from

374 the Zeiss RemCon communication protocol to the Zeiss API which reduced the overhead by

375 several seconds per imaging/milling cycle.

376  The coated samples were imaged by a 0.14–0.3 nA electron beam with 0.7–1.2 keV

377 landing energy at 0.1–2 MHz. Detailed conditions are listed in Table 1. In most cases, both

378 backscattered and secondary electron signals were collected by InLens detector to provide

better signal-to-noise ratio. A faster SEM scanning rate was applied on samples with stronger

staining contrast thus larger imaging volumes were obtained. The x-y pixel size was set at 4 nm.

A subsequently applied focused Ga+ beam of 15 nA at 30 keV strafed across the top surface and

ablated away 4 nm of the surface. The newly exposed surface was then imaged again. The

ablation – imaging cycle continued for two to five weeks to complete FIB-SEM imaging one

sample. The sequence of acquired images formed a raw imaged volume, followed by post

processing of image registration and alignment using a Scale Invariant Feature Transform

(SIFT)[25] based algorithm to form isotropic 4 nm voxels. The final aligned stack consists of an

isotropic volume, containing multiple complete cells, which can be viewed in any arbitrary

orientations.

Evaluation of FIB-SEM resolution

In order to evaluate the actual resolution in the acquired FIB-SEM datasets we analyzed the

transitions of the edges of the ribosomes abundant within the cell volumes. Ribosomes are

macromolecular machines consisting of RNA and associated proteins. Their small size (20–30

nm), abundance in the cytoplasm of living cells, and the fact that they are stained with high EM

contrast makes them good candidates for resolution evaluation.

First, we selected volumes of approximately 1000–2000 x 1000 x 1000 pixels in cultured

datasets. These volumes were selected in the middle of datasets in areas with large number of

ribosomes. We then calculated histograms of grey level values in these volumes and subtracted

~0.8 * histogram max to bring the average cytoplasmic material signal level close to 0. We then

used Laplacian of Gaussian (LoG) algorithm (part of Python scikit-image package). It computes

the Laplacian of Gaussian images with successively increasing standard deviation. We had set

the limit of standard deviation for LoG between 1 and 2.5 pixels (all these datasets have 4nm

voxels). Filters were used to exclude the following blobs: average value below zero, edge value

above 0.4 times of max value, 37%–63% transition longer than 15nm, amplitudes below certain

threshold so that at least approximately 3000 blobs per dataset were remaining for analysis.

The summaries of the 37%–63% transitions and 20%–80% transitions for the 3D

ribosome blobs selected in cultured cell datasets are plotted in Fig. 1b. The histograms of the

37%–63% transitions these datasets are shown in Extended Data Fig. 2. The examples of 3D

blobs for each dataset are presented in Extended Data Fig. 3–9. For each dataset the cross-

sections and transition analysis for 3 brightest and 3 dimmest blobs are shown.

Accurate determination of FIB-SEM milling rate

The position of a sample block face can be measured by the SEM beam and the FIB beam

independently. An in-line auto focus routine[6] keeps the SEM images in focus throughout the

entire image acquisition. While the working distance of the SEM beam detects the block face

position by measuring its distance from the SEM objective lens, the FIB milling position under

closed-loop control indicates the block face position directly from the orthogonal angle. As the

sample block face being imaged then milled away layer-by-layer, the SEM working distance and

the FIB beam position track the block face location accordingly. Therefore, one can estimate the

total thickness milled away by FIB based on the changes of either SEM working distance or FIB

milling position from the beginning to the end of the FIB-SEM acquisition. The average milling

rate of z-step can then be calculated as the total milling depth divided by the number of image

frames. Here we report the average z-scaling factors derived from the two measurement

approaches in Extended Data Table 1. It can be used to scale the dataset to generate real

isotropic voxels. For example, HeLa Cell 2017-06 dataset has a z-scaling factor of 1.31 which

means the actual z-step is average at 5.2 nm instead of 4.0 nm.

Data availability

426   All raw and segmented datasets in this work are available on OpenOrganelle website

427   (https://openorganelle.janelia.org).

Code availability

429   FIB-SEM image acquisition LabVIEW codes used in this work are available from

430   https://doi.org/10.7554/eLife.25916.031.

Acknowledgements

We thank K. Hayworth and W. Qiu at Howard Hughes Medical Institute (HHMI) Janelia Research Campus (JRC) for invaluable discussions and data collection support. We gratefully acknowledge P. Rivlin, S. Plaza, and I. Meinertzhagen for JRC EM shared resource and FlyEM project team support on staining protocols development. We thank the electron microscopy facility of MPI-CBG and of the CMCB Technology Platform at TU Dresden for their services. We also thank Y. Wu from Pietro De Camilli's laboratory at Yale for advice.

Funding

C.S.X., S.P., G.S., D.B., S.T., Z.L., A.V.W., H.A.P., N.I., T.C.W., H.F.H. are funded by Howard Hughes Medical Institute (HHMI). A.M. received support from the Carl Gustav Carus Faculty of Medicine at TU Dresden via a MeDDrive GRANT. A.M. and M.S. were supported with funds from the German Center for Diabetes Research (DZD e.V.) by the German Ministry for Education and Research (BMBF), from the Germain-Israeli Foundation for Scientific Research and Development (GIF) (grant I-1429-201.2/2017), from the German Research Foundation (DFG) jointly with the Agence nationale de la recherche (ANR) (grant SO 818/6-1) to M.S.. H.K.H and S.B.v.E. are funded by NIAID grant R01AI138625. R.V.F. is supported by NIH R01GM124348. T.C.W. is supported by NIH R01GM097194. J.C. is a fellow of the Damon Runyon Cancer Research Foundation.

Author Contributions

C.S.X. and H.F.H. supervised the project; C.S.X., S.P., G.S., and H.F.H. wrote the manuscript with input from all co-authors; C.S.X. developed enhanced FIB-SEM platform for large volume high resolution imaging, and optimized imaging conditions; C.S.X., S.P., and G.S. conducted FIB-SEM experiments; C.S.X. and S.P. performed image post processing; S.P., G.S., A.M., A.T.R., H.K.H., S.B.v.E., Z.L., A.V.W., H.A.P., N.I., and J.C. prepared samples; G.S., A.M., A.T.R., and S.T. analyzed data; D.B. prepared and uploaded data to OpenOrganelle website; T.C.W., R.V.F., S.B.v.E., I.M., and M.S. proposed biological questions and provided samples.

Competing interests

Portions of the technology described herein are covered by U.S. Patent 10,600,615 issued to C.S.X. and H.F.H., and assigned to HHMI.

Fig. 1 **Enhanced FIB-SEM configuration, operation, resolution, and volume**. **a**, Sketch of FIB

ablation and SEM imaging. The two operations iterate every minute and repeat tens of

thousands of times to make 3D image stacks, **b**, Resolution characterization: X position of

vertical line indicates the step edge transition as observed on gold nano particles on a carbon

substrate, the round dots are step edge transitions on ribosomes along x-y axis and the z axis.

Each dot represents the average value from more than 3000 ribosomes in one of the seven

cultured cell samples. **c**, Smallest isotropic voxel size vs volume for comparing different volume

EM methods. The light green space indicates the new Resolution-Volume regime uniquely

accessible with enhanced FIB-SEM technology through long term imaging. The regime of this

work at 4 nm isotropic voxel sampling is highlighted in yellow.

Fig. 2 **Interphase HeLa cell**. **a**, FIB-SEM overview with cutaway, and manually segmented

interior features (mitochondria, green; centrosomes, red; one cistern of a Golgi stack, magenta;

a segment of nuclear membrane, blue, polyribosome chains, yellow). Zoomed in cross-sections

of three select areas containing, **b**, Centrosomes, **c**, Golgi stack, and **d**, Nucleus. Scale bars are 1

μm.

Fig. 3 **Murine CTL engaging an ovarian cancer cell**. **a,** FIB-SEM overview with cutaway, and

manually segmented membranes of CTL (green surface and red contour) and cancer cell (cyan

surface and blue contour). Zoomed in cross-sections of three select areas containing **b,** CTL cell

cupping the target cancer cell, **c,** Centrosome, and **d,** Lytic granule. Scale bars are 1 μm.

Fig. 4 **Tissue sample datasets**. **a**, Pancreatic islets treated with 16.7 mM glucose containing

several complete beta cells with detailed structures shown in **b**, A primary cilium with axoneme

(arrow) and centrioles (asterisk), **c**, Intermingled microvilli (arrow), **d**, Ultrastructural diversity

among insulin SGs containing rod-shaped (arrow) or spherical crystals (asterisk), **e**, Contacts of

ER and insulin SGs (arrow), **f**, Microtubules (arrow), Golgi apparatus (asterisks) and ribosomes.

g, *Drosophila* fan-shaped body with detailed structures shown in **h**, Dense-core vesicles of

different sizes (arrows), **i**, Multiple synaptic sites viewed from the side (arrows), **j**, A presynaptic

T-bar viewed from the top (arrow), **k**, A longitudinal-section view of microtubules (arrows), **l**, A

cross-section view of microtubule arrays (arrows). Scale bar, 1 µm in **a**, 500 nm in **b–f**, 5 µm in

g, 200 nm in **h–l**.

Extended Data Fig. 1 **Murine CTL engaging an ovarian cancer cell.** Zooms on regions showing

different synapse topology features. **a**, Interdigitation, **b**, Flat apposition, and **c**, Filopodia

caught between cells. Scale bars are 0.5 μm.

Extended Data Fig. 2 **Edge transition distributions determined from ribosomes in cultured cells**

datasets. Distributions of 37%–63% (**a**) and 20%–80% (**b**) transition distances in X-, Y- (left), Z_{top}-

(center), and Z_{bot}- (right) directions.

Extended Data Fig. 3 **Cross-sections of the example of ribosomes from the dataset HeLa Cell2**

and the profiles with the transition analysis.

Extended Data Fig. 4 **Cross-sections of the example of ribosomes from the dataset HeLa Cell3**

and the profiles with the transition analysis.

Extended Data Fig. 5 **Cross-sections of the example of ribosomes from the dataset**

Macrophage Cell2 and the profiles with the transition analysis.

Extended Data Fig. 6 **Cross-sections of the example of ribosomes from the dataset Jurkat Cell1**

and the profiles with the transition analysis.

Immortalized breat cancer cell (SUM159) 2017-11-21, z-scale 1.14, selected 3D blobs

Extended Data Fig. 7 **Cross-sections of the example of ribosomes from the dataset SUM159**

Cell2 and the profiles with the transition analysis.

Extended Data Fig. 8 **Cross-sections of the example of ribosomes from the dataset T-Cell on**

Cancer Cell and the profiles with the transition analysis.

Extended Data Fig. 9 **Cross-sections of the example of ribosomes from the dataset HeLa Cell4**

and the profiles with the transition analysis.

Tables

Table 1. Descriptions of exemplary datasets with selected imaging conditions

Sample Description	Protocol	Landing Energy (kV)	Imaging Current (nA)	Scan Speed (MHz)	Acquisition Duration (days)	Data Volume: X * Y * Z (μm^3)	Z-scaling factor
Cell 4 nm							
Interphase HeLa cell 2017-06-21	High pressure freezing, freeze-substitution with 2% OsO4 0.1% UA 3% H2O in Acetone, resin embedding in Eponate 12	0.9	0.24	0.1	25	48 * 6 * 25 (7800)	1.31
Interphase HeLa cell 2017-08-09†		1.0	0.24	0.2	31	50 * 4 * 48 (9600)	0.81
Mitotic HeLa cell 2019-05-30		0.9	0.24	0.2	15	30 * 14 * 33 (13300)	1.07
Wild-type THP-1 macrophage 2018-11-11		1.2	0.24	0.2	20	40 * 8 * 44 (14200)	0.84
Immortalized T-cells (Jurkat) 2018-08-10		1.0	0.24	0.2	19	40 * 12 * 34 (16400)	0.86
Immortalized breast cancer cell (SUM159) 2017-11-21‡		1.2	0.24	0.2	19	64 * 11 * 31 (21900)	1.14
Killer T-Cell attacking cancer cell 2020-02-04†		0.7	0.25	0.2	29	74 * 13 * 48 (45500)	0.87
Tissue 4 nm							
Isolated murine pancreatic islets 2019-03-01†	HPF, FS with 2%OsO4, 1% UA, 0.5% GA, 5% H2O in Acetone with 1% Methanol followed by 0.2% TCH in 80% Methanol at RT, 2% OsO4 in Acetone at RT, 1% UA in Acetone + 10% Methanol	0.9	0.14	0.2	15	30 * 20 * 29 (17400)	0.85
Drosophila brain: Fan-shaped body of a 5-day-old male adult 2019-09-14†	Chemical Fixation, ORTO-Lead-EPTA staining via Progressive Lowering of Temperature and Low Temperature Staining (PLT-LTS) heavy metal enhancement protocol	0.7	0.24	1.5	15	45 * 56 * 45 (113400)	1
Drosophila brain: Accessory calyx of a 5-day-old male adult 2019-12-06		0.7	0.3	2	19	50 * 40 * 87 (175000)	0.93

† Figures in main text.

‡ Treated with 0.5 mM oleic acid for 45 mins prior to high pressure freezing to induce the formation of lipid

droplets.

529 Extended Data Table 1. Z scaling factors estimated using SEM working distance and FIB milling

530 voltage.

DataSet	Z (milling direction) pixel scaling factor		
	SEM Working Distance (WD)	FIB milling voltage (MV)	Mean (WD, MV)
Interphase HeLa cell 2017-06-21	1.22	1.40	1.31
Interphase HeLa cell 2017-08-09	0.80	0.83	0.81
Mitotic HeLa cell 2019-05-30	1.07	1.07	1.07
Wild-type THP-1 macrophage 2018-11-11	0.81	0.87	0.84
Immortalized T-cells (Jurkat) 2018-08-10	0.84	0.88	0.86
Immortalized breast cancer cell (SUM159) 2017-11-21	1.13	1.14	1.14
Killer T-Cell attacking cancer cell 2020-02-04	0.88	0.87	0.87
Isolated murine pancreatic islets 2019-03-01	0.86	0.84	0.85
Drosophila Fan-shaped body 2019-09-14	0.98	1.03	1.00
Drosophila Accessory calyx 2019-12-06	0.95	0.92	0.93

 References

[20] Sartori, N., Richter, K., & Dubochet, J. Vitrification depth can be increased more than 10-fold by high-pressure freezing. *J. Microsc.* **172**(1), 55–61 (1993). https://doi.org/10.1111/j.1365-2818.1993.tb03393.x

[21] Gotoh M., Maki T., Kiyoizumi T., Satomi S., & Monaco A. P. An improved method for isolation of mouse pancreatic islets. *Transplantation* **40**, 437–438 (1985). https://doi.org/10.1097/00007890-198510000-00018

[22] Müller, A. et al. A global approach for quantitative super resolution and electron microscopy on cryo and epoxy sections using self-labeling protein tags. *Sci. Rep.* **7**, 23 (2017). https://doi.org/10.1038/s41598-017-00033-x

[23] Kremer, J. R., Mastronarde, D. N., & McIntosh, J. R., Computer visualization of three-dimensional image data using IMOD. *J. Struct. Biol.* **116**, 71–76 (1996). https://doi.org/10.1006/jsbi.1996.0013

[24] Lu Z. et al. En bloc preparation of *Drosophila* brains enables high-throughput FIB-SEM connectomics. *bioRxiv*: 855130 (2019). https://doi.org/10.1101/855130

[25] Saalfeld, S., Cardona, A., Hartenstein, V. & Tomančák, P. As-rigid-as-possible mosaicking and serial section registration of large ssTEM datasets. *Bioinformatics* **26**, i57–63 (2010). https://doi.org/10.1093/bioinformatics/btq219