An open-access volume electron

microscopy atlas of whole cells and tissues

C. Shan Xu[1]†, Song Pang[1]*, Gleb Shtengel[1]*, Andreas Müller[2,3,4], Alex T. Ritter[5], Huxley K.

Hoffman[6]#, Shin-ya Takemura[1], Zhiyuan Lu[1], H. Amalia Pasolli[1§], Nirmala Iyer[1], Jeeyun Chung[7,8],

Davis Bennett[1], Aubrey V. Weigel[1], Melanie Freeman[1‡], Schuyler B. van Engelenburg[6], Tobias C.

Walther[7,8,9], Robert V. Farese, Jr.[7,8], Jennifer Lippincott-Schwartz[1], Ira Mellman[5], Michele

Solimena[2,3,4,10], Harald F. Hess[1]†

[1]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147, USA

[2]Molecular Diabetology, University Hospital and Faculty of Medicine Carl Gustav Carus, TU

Dresden, Dresden, Germany

[3]Paul Langerhans Institute Dresden (PLID) of the Helmholtz Center Munich at the University

Hospital Carl Gustav Carus and Faculty of Medicine of the TU Dresden, Dresden, Germany

[4]German Center for Diabetes Research (DZD e.V.), Neuherberg, Germany

[5]Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA

[6]Molecular and Cellular Biophysics Program, Department of Biological Sciences, University of

Denver, Denver, CO 80210, USA

[7]Department of Molecular Metabolism, Harvard T.H. Chan School of Public Health, Boston, MA

02115, USA

[8]Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA

[9]Howard Hughes Medical Institute, Boston, MA 02492, USA; Broad Institute of Harvard and

MIT, Cambridge, MA 02124, USA

[10]Max-Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany

†Correspondence to: xuc@janelia.hhmi.org (C.S.X), hessh@janelia.hhmi.org (H.F.H.)

*These authors contributed equally to this work

#Present Address: Department of Cell and Developmental Biology, University of Colorado

Anschutz Medical Campus, Aurora, CO 80045, USA

§Present Address: Electron Microscopy Resource Center, The Rockefeller University, New York,

NY 10065, USA

‡Present Address: Advanced Bio-imaging Center, Department of Molecular & Cell Biology,

University of California, Berkeley, CA 94720, USA

Understanding cellular architecture is essential for understanding biology. Electron microscopy (EM) uniquely visualizes cellular structures with nanometer resolution. However, traditional methods, such as thin-section EM or EM tomography, have limitations inasmuch as they only visualize a single slice or a relatively small volume of the cell, respectively. Focused Ion Beam-Scanning Electron Microscopy (FIB-SEM) demonstrated the ability to image cellular samples at 4-nm isotropic voxels with rather limited imageable volume. Here, we present 3D EM images of whole cells and tissues with two orders of magnitude increases in imageable volume at 4-nm voxels. Such data with a combined fine resolution scale and large sample size do not currently exist, and are enabled by the advances in higher precision and stability of FIB milling, together with enhanced signal detection and faster SEM scanning. More importantly, we have generated a volume EM atlas encompassing ten diverse datasets of whole cells and tissues, from cancer cells to immune cells, and from mouse pancreatic islets to *Drosophila* neural tissues. These open-access data (via OpenOrganelle) represent a foundation to nucleate a new field of high-resolution whole-cell volume EM and subsequent analyses, and invite biologists to explore this new paradigm and pose fundamentally new questions.

Individual cells and tissues can be described by a 3D hierarchy of ultrastructural details, from single protein molecules to organelles within complete cellular architecture[1]. Transmission electron microscopy (TEM) images from ~30–300 nm diamond-knife cut sections of metal stained samples provide convenient high-resolution visualization on a small fraction of the whole cell, which have profoundly shaped our understanding of organelles[2]. Series of sections

can be digitally stitched together to form a 3D volume, but the 30 nm or larger sampling

interval sacrifices structural detail. Tilt tomography achieves higher 3D resolution on ~200-nm-

thick sections. Alternatively, diamond-knife cut surfaces of a block face sequentially imaged

with scanning electron microscopy (SEM)[3] was a pioneering step to access larger volumes, but

its z resolution is limited to 25 nm due to electron radiation damage from beam scanning that

prevents consistent cutting of thinner slices[4] in subsequent rounds. This z-resolution constraint

can be overcome using FIB-SEM that offers high isotropic resolution. However, its deficiencies

in imaging speed and duration (several days) cap the maximum imageable volume. Our prior

work (Fig. 1a) reported the transformation of a conventional FIB-SEM lacking long-term

reliability into an enhanced platform capable of months to years of continuous imaging at 8-nm

isotropic voxels without defects in the final image stack[5,6,7,8], but we struggled to obtain

volumes of ~500 μm^3 at 4-nm voxels due to slow SEM scan using backscattered electron

detection via a sample biasing scheme[5,9], and FIB milling instability induced by higher electron

radiation energy density (~400 keV/nm^3). Thus, higher-resolution isotropic 3D whole cell data

remained mostly out of reach.

Here we report imageable volumes greater than 100,000 μm^3 at 4-nm voxels. This newly

accessible paradigm is illustrated in Extended Data Fig. 1. The two orders of magnitude

improvement in volumes compared to our prior work[5] primarily come from: 1) higher precision

and stability of FIB milling control to extend reliable long-term acquisition to 4-nm voxels, and

2) enhanced SEM signal detection using secondary electrons with higher beam current to

achieve faster imaging, hence reducing electron radiation energy density (from ~400 to ~80

keV/nm^3) that further improves FIB milling control. To overcome milling instability, we

integrated a more stable FIB column (Capella from Zeiss), and tightened its FIB emission current

control band to obtain a consistent milling beam profile. We also re-configured the closed-loop

control[5] to reduce the milling variation between FIB reheat cycles. Such FIB milling optimization

allowed for faster SEM imaging[5] using secondary electrons with 5x lower electron radiation

energy density, which subsequentially further mitigated FIB milling artifacts and instability.

Additionally, we investigated a multi-dimensional operation space to optimize image contrast

and isotropic resolution. We found that a beam current of 200 to 300 pA and a landing energy

of 700 to 900 eV are optimal which further accelerate the imaging rate. Such positive synergy

between faster SEM scanning and robust FIB milling has extended reliable imaging acquisition

at 4-nm isotropic voxels from less than a week to months. The detailed instrumentation

improvements are described in the Methods Section. To validate the choice of sampling voxel

size, we characterized the resolution in x-y and z by analyzing the transitions at the edges of

gold nanoparticles on a carbon substrate and ribosomes in cultured cells. The ribosomes in

biological specimens serve as a convenient *in situ* resolution standard. Fig. 1b shows near-

isotropic resolution matching of the 4-nm voxel size with the average transitions from 37% to

63% being 2.5 nm in x-y and 3.1 nm in z, respectively. The x-y transitions determined from

ribosome edges are about 30% larger than those for the gold nanoparticles (vertical lines in Fig.

1b). The procedure for estimating the resolution with examples is presented in the Methods

Section. Furthermore, an improved sample staining protocol yielding higher contrast leads to

additional 5x–10x scanning rate improvement, thereby 200x larger volume is demonstrated on

Drosophila brain samples (Supplementary Video 1). The maximum volume can be seamlessly

extended with longer imaging acquisition.

Enabled by these advances, we here present a 3D atlas of whole cells and tissues at 4-nm

voxels. The initial datasets (Table 1) consist of ten wild-type biological specimens from cancer

cells to immune cells, and from mouse pancreatic islets to *Drosophila* neural tissues, each

requiring weeks of uninterrupted imaging. The resulting data demonstrate the finest possible

isotropic resolution for whole cell volumes. To highlight some features that are only visible at

the higher resolution, a comparison between 4-nm versus 8-nm voxel sampling is illustrated in

Fig. 1c–k. In particular, the nucleosomes inside its nucleus showing finer details are on the

verge of being individually resolved at 4 nm (Fig. 1c); the detailed Golgi cisternae (Fig. 1f), the

close contacts of ER-mitochondria, and the hollow core of microtubules (Fig. 1i) are only

resolvable at 4 nm. Furthermore, Supplementary Video 1 visualizes the value of finer resolution

with 200x larger volume to better illustrate the newly accessible combined larger size-higher

resolution paradigm. By making these data openly available, we invite biologists to contemplate

analyzing such comprehensive whole cell details, and hope to inspire new questions that

harness complete rather than partial cell data to unveil new insights.

HeLa cells, human cervical cancer cells, are easily cultured and widely used in cell biology

laboratories as a basic model to test diverse hypotheses. Fig. 2 presents a typical 3D dataset of

an entire HeLa cell. The top panel provides an overview with manually segmented

mitochondrial network in green (Fig. 2a). The bottom row panels show 2D cross-sections of

other cellular organelles, such as the centrosome (Fig. 2b), the Golgi apparatus (Fig. 2c), and

nuclear envelope (Fig. 2d). No single 2D cross-section allows visualizing all centriole sub-distal

appendages, however quick segmentation of the 3D dataset depicts them clearly (red, Fig. 2b).

Stereotypical 2D images of the Golgi stacks do not reveal the fenestration details nor long thin

tubular extensions that can readily be seen in 3D segmented single Golgi cisterna (magenta, Fig.

2c). Polyribosome chains on the nuclear envelope are mostly hidden in 2D cross-sections, but

easily resolved and detailed in 3D (yellow, Fig. 2d). Evidently, having isotropic 3D images of such

cells in their entirety can provide an accurate reference to which perturbations in growth,

genetic, environment, etc. can be compared.

Cytotoxic T lymphocytes (CTLs) are immune cells that have the capacity to identify and destroy

virus-infected or cancerous cells. Upon encountering a target, CTLs form an organized

intercellular interface called the immunological synapse (IS). Signaling at the IS directs the

polarized release of toxic proteins housed in specialized secretory lysosomes called lytic

granules (LGs), which results in target cell death. Due to the important role of CTLs in anti-viral

and anti-cancer immunity, the IS has been the subject of intense scrutiny. Limited single slice

TEM images of CTL:target conjugates have revealed interesting features of the IS, but do not

represent the full scope of this dynamic, three-dimensional structure[10]. Fig. 3 presents a first

comprehensive 3D dataset of a mature murine CTL (green) engaging an ID8 ovarian cancer cell

(cyan). Although there is a minor membrane damage in the area of IS, which most likely

occurred during high-pressure freezing or freeze-substitution or resin embedding[11], the

isotropic high-resolution information of FIB-SEM imaging provides a unique and complete map

of the complex membrane topology at the interface between T cell and target. Fig. 3b

demonstrates the stereotypical CTL "cupping" of the target cell at the IS through ortho-slices.

Zoomed in images further illustrate the variety of features across this interface including

membrane interdigitation (Extended Data Fig. 2a), flat membrane apposition (Extended Data

143 Fig. 2b), and filopodia of the target cell trapped between the two cells (Extended Data Fig. 2c).

144 Without the ability of high resolution 3D imaging, these filopodia could be mistaken for vesicles

145 trapped between T cell and target. Additionally, Supplementary Video 2 depicts the

146 morphological features in greater details, which further highlights the hallmarks of CTL killing:

147 T-cell cupping the target cancer cell (Fig. 3b), polarization of the centrosome toward the target

148 cell[12,13] (Fig. 3c), and the lytic granules with diverse ultrastructures aggregating near the

149 synapse (Fig. 3d). Such high resolution whole cell FIB-SEM dataset allows for the label-free

150 localization of all organelles in the T cell including those responsible for target killing and

151 cytokine secretion.

152 **Pancreatic islets** are micro-organs consisting mainly of beta, alpha, delta, polypeptide cells and

153 endothelial cells. Among them beta cells are in the majority and secrete insulin stored in

154 secretory granules (SGs) to maintain blood glucose homeostasis. Few beta cells have been

155 reconstructed by serial section electron tomography at medium resolution[14] to allow

156 assessment of subcellular features such as insulin SG number and shape of mitochondria, but

157 lacking the resolution for imaging of e.g., ER and microtubules. Large-scale high-resolution FIB-

158 SEM enables the analysis of ultrastructural differences between beta cells within an islet as well

159 as features that require higher resolution such as ribosomes and the cytoskeleton. Fig. 4a

160 displays several complete beta cells that are entirely included within the 30 x 20 x 25 μm^3

161 volume. The large volume data allow for studying the interaction of beta cells, e.g., full primary

162 cilia and their connections to neighboring cells and the intermingling of microvilli (Fig. 4b, 4c).

163 At 4-nm voxel sampling, we observed ultrastructural differences between insulin SGs (Fig. 4d).

164 For the first time, we report close contacts of ER and insulin SGs in 3D (Fig. 4e). Ultimately, such

high resolution allows the sharp definition of microtubules, single ribosomes, membrane

vesicles, and Golgi cisternae for quantitative measurements (Fig. 4f). The seamless imaging

results in precisely aligned stacks, further facilitating automated 3D segmentation. Specifically,

emerging from these FIB-SEM datasets is a comprehensive 3D representation of microtubule

networks and how they interact with other organelles[15]. We could show that beta cell

microtubules are arranged into non-radial networks which for the most part do not originate

either from centrioles or endomembranes[15]. Moreover, glucose stimulation changes the

number and length of microtubule polymers without changing their overall density[15]. Finally,

we found insulin SGs to be enriched near the plasma membrane in association with

microtubules, independent of the extracellular glucose levels[15].

The **central complex in insect brains** is a highly conserved brain region that controls various

behaviors. Fig. 4g shows higher-resolution images acquired from the fan-shaped body of the

central complex of *Drosophila*. To understand brain functions, it is imperative to identify the

synaptic levels of neuron connectivity over large volumes through key synaptic elements such

as presynaptic T-bars and postsynaptic densities (PSDs) (Fig. 4i, 4j). To investigate if the central

complex contains characteristic synaptic motifs, such as polyadic synapse, rosette synapse, or

any new synaptic motif, validating the synaptic structures with high-resolution data is crucial.

However, a trade-off between volume and resolution often limits capabilities in connectomics

studies, where large-scale neuronal tracing and fine synaptic structure identification are equally

important. High resolution large volume FIB-SEM imaging, shown here, bridges these

capabilities and therefore provides significant possibilities. Such datasets allow reconstruction

of not only the large neuronal objects and glial cells, but also smaller intracellular components

such as synaptic vesicles, dense-core vesicles (Fig. 4h), and microtubules (Fig. 4k, 4l). Knowing

the types of synaptic vesicles could add important insights to understand the properties of

synaptic connections. As an example, the terminals of dopaminergic neurons in the fly's

olfactory center were shown to contain both clear and dense-core vesicles[16], in congruence

with having more than one neurotransmitter[17].

A considerable investment in technology development, equipment resources, and imaging

duration is required to produce large volume high resolution datasets, but the resulting wealth

of such data is unprecedented and comprehensive, covering whole cells with all their organelles

and proteins that can be highlighted with heavy metal stains. Here, we open access to this

volume EM atlas through OpenOrganelle (https://openorganelle.janelia.org)[18]. With access to

these datasets and future ones, biologists can assess whether whole cell structure statistics and

architecture characterization of their interest are visible and possible to measure for evaluating

the merits of further exploration. Additionally, the data can be used as a reference

ultrastructural atlas, providing insights into both known and previously unappreciated

structural details of intracellular organization. Each dataset can be downloaded to mine, while

pre-selected views of interest and cross-sections can be browsed on-line with *neuroglancer*, a

built-in visualization tool. More importantly, these data enable a holistic way of looking into the

cellular world and stand as a foundation of segmentation and analysis efforts that could be

pursued by the scientific community.

In summary, we achieved large imaging volumes over 100,000 μm^3 at the finest possible

isotropic resolution. Thereby, entire mammalian cells and multicellular parts of tissues were

imaged at 4-nm isotropic voxels to reveal a more holistic 3D cellular architecture. Ten diverse

datasets of reference wild-type examples are now made accessible through OpenOrganelle[18].

This allows the study of complex cell morphology, complete interfaces involved in cell-cell

interactions, and the shapes, volumes, distributions, and relationships of intracellular

organelles. Particularly, the examination of a wild-type cell can encourage further experiments

to compare features against perturbations of disease, mutations, development, or environment

at the whole cell level. These datasets and targeted feature analyses not only could lead to

discoveries of unanticipated new features, but also should spur developments of general tools

to break the bottleneck of 3D data mining. Automation of the time-consuming process of

segmentation on the whole cell scale is a crucial next step. Such an approach is presented in the

associated paper[18]. We believe that open access to rich 3D FIB-SEM datasets will provide

important insights, guidance, and inspiration for the research community to further explore and

address a new broad range of biological questions.

 Reference

[1] Lodish, H. et al. *Molecular Cell Biology* (W. H. Freeman, New York, 2016).

[2] Porter, K. R., Claude, A., & Fullam, E. F. A study of tissue culture cells by electron microscopy: methods and preliminary observations. *The Journal of experimental medicine*, **81**(3), 233–246 (1945). https://doi.org/10.1084/jem.81.3.233

[3] Denk, W., & Horstmann, H. Serial block-face scanning electron microscopy to reconstruct three-dimensional tissue nanostructure. *PLoS Biol* 2004, **2**:e329 (2004). https://doi.org/10.1371/journal.pbio.0020329

[4] Titze, B. Techniques to prevent sample surface charging and reduce beam damage effects for SBEM imaging. Dissertation. Heidelberg University. 1–112 (2013). https://doi.org/10.11588/heidok.00015372

[5] Xu, C. S. et al. Enhanced FIB-SEM systems for large-volume 3D imaging. *eLife* **6**:e25916 (2017). https://doi.org/10.7554/eLife.25916

[6] Xu, C. S., Hayworth, K. J., & Hess, H. F. Enhanced FIB-SEM systems for large-volume 3D imaging. US Patent 10,600,615, 24 Mar (2020).

[7] Xu, C. S., Pang, S., Hayworth, K. J., & Hess, H. F. Transforming FIB-SEM Systems for Large-Volume Connectomics and Cell Biology. In: Wacker, I., Hummel, E., Burgold, S., Schröder, R. (eds) Volume Microscopy. Neuromethods, vol 155 (2020). Humana, New York, NY. https://doi.org/10.1007/978-1-0716-0691-9_12

[8] Xu, C. S. et al. A connectome of the adult *Drosophila* central brain. *bioRxiv* 2020.01.21.911859. (2020). https://doi.org/10.1101/2020.01.21.911859

[9] Wu, Y. et al. Contacts between the endoplasmic reticulum and other membranes in neurons. *PNAS* **114**(24) E4859-E4867 (2017). https://doi.org/10.1073/pnas.1701078114

[10] Stinchcombe, J. C., Bossi, G., Booth, S., & Griffiths, G. M. The immunological synapse of CTL contains a secretory domain and membrane bridges. *Immunity*. 2001;15(5):751–761. https://doi.org/10.1016/s1074-7613(01)00234-5

[11] Bullen, A., Taylor, R. R., Kachar, B., Moores, C., Fleck, R. A. & Forge, A. Inner ear tissue preservation by rapid freezing: Improving fixation by high-pressure freezing and hybrid methods. *Hear. Res.* **315**, 49–60 (2014). https://doi.org/10.1016/j.heares.2014.06.006

[12] Kupfer, A. & Dennert, G. Reorientation of the microtubule-organizing center and the Golgi apparatus in cloned cytotoxic lymphocytes triggered by binding to lysable target cells. *J Immunol*. 1984;133(5):2762-2766.

[13] Stinchcombe, J. C., Majorovits, E., Bossi, G., Fuller, S. & Griffiths, G.M. Centrosome polarization delivers secretory granules to the immunological synapse [published correction appears in Nature. 2006 Nov 9;444(7116):236]. *Nature*. 2006;443(7110):462–465. https://doi.org/10.1038/nature05071

[14] Noske, A. B., Costin, A. J., Morgan, G. P. & Marsh, B. J. Expedited approaches to whole cell electron tomography and organelle mark-up in situ in high-pressure frozen pancreatic islets. *J. Struct. Biol.* **161**, 298–313 (2008). https://doi.org/10.1016/j.jsb.2007.09.015

[15] Müller, A. et al. 3D FIB-SEM reconstruction of microtubule-organelle interaction in whole primary mouse β cells. *J. Cell. Biol.* 220 (2): e202010039 (2021). https://doi.org/10.1083/jcb.202010039

[16] Takemura, S. Y. et al. A connectome of a learning and memory center in the adult *Drosophila* brain. *eLife* **6**, e26975 (2017). https://doi.org/10.7554/eLife.26975

[17] Aso, Y. et al. Nitric oxide acts as a cotransmitter in a subset of dopaminergic neurons to diversify memory dynamics. *eLife* **8**, e49257 (2019). https://doi.org/10.7554/eLife.49257

[18] Heinrich, L. et al. Automatic whole cell organelle segmentation in volumetric electron microscopy. *bioRxiv* 2020.11.14.382143 (2020). https://doi.org/10.1101/2020.11.13.382143

Methods

The initial datasets of seven common wild-type cultured cells (from cancer cells to immune cells) and three tissue samples (from mouse pancreatic islets to *Drosophila* neural tissues) are tabulated in Table 1. Detailed procedures are described below.

Cultured cell sample preparation

Cultured cells were cryo-fixed by high pressure freezing (HPF), which vitrified a sample on a millisecond time scale to best preserve any dynamic structural details. Although HPF sometimes can cause segregation artifacts (i.e. freezing damage), it minimizes any possibility of artifacts that might be of concern with chemical fixation. HPF remains the only viable option for non-chemical fixation of samples more than a few hundred nanometers thick. The subsequent freeze substitution labeled biomolecules with heavy metals to provide contrast for EM.

1. HeLa cells

HeLa cells (CCL-2) were purchased from American Type Culture Collection (ATCC). HeLa cells were maintained in EMEM medium (ATCC, 30-2003) supplemented with 10% FBS (Corning, 35-011-CV) and 1X penicillin-streptomycin solution (Corning, 30-002-CI). Trypsinized HeLa cells were seeded on pre-cleaned and edge-coated sapphire coverslips. Immediately prior to freezing, live cells were inspected to ensure cell morphology and viability. After quality assurance the cells were transferred to a water jacketed CO_2 incubator (Thermo Fisher Scientific, Midi 40) kept at 37° C, 5% CO_2, and 100% humidity while awaiting freezing. Each coverslip was removed from the incubator immediately prior to the freezing procedure, and dipped three times in the 25% w/v mixture of Bovine Serum Albumin (BSA) (B4287, Sigma Aldrich)) in the cell medium, which served as a cryo-protectant to prevent ice crystal

formation[19]. The samples were then high-pressure frozen between two Aluminum planchettes

(Technotrade International, 389 and 479) in Wohlwend HPF Compact 02 High Pressure Freezing

Machine (Technotrade International). Once the samples were frozen, they could be stored

indefinitely in liquid nitrogen for subsequent freeze-substitution (FS) and resin embedding (RE).

FS was performed using automated FS machine (AFS2, Leica Microsystems): coverslips were

transferred to cryotubes containing FS media (2% OsO_4, 0.1% Uranyl Acetate (UA), and 3%

water in acetone) under liquid nitrogen, followed by a programmed FS schedule. Resin

embedding was performed immediately after FS. Samples were removed from the AFS2

machine, washed 3 times in anhydrous acetone for a total of 10 min and embedded in Eponate

12 which was polymerized for 48 hours at 60°C. Following EPON embedding, the coverslip was

separated from the resin block containing the cells by sequential immersion in liquid nitrogen

and hot water. EPON was chosen as the initial embedding resin due to less embedding artifacts.

However, EPON generates streaking artifact during FIB-SEM imaging[5]. A thin layer (5–10 μm) of

Durcupan on the specimen surface facing the FIB beam can effectively mitigate such streaks

during FIB-SEM imaging. Therefore, once the coverslip was removed, the exposed surface was

immediately re-embedded in Durcupan ACM resin (Sigma Aldrich, set 44610). The detail

procedures of coverslip preparation, high pressure freezing, free substitution, and resin

embedding are presented in the Supplementary Information.

2. Cytotoxic T cell and T cell/Cancer cell conjugate (CTLs)

To generate CTLs from OT-I mice (Jackson Labs), splenocytes were isolated and stimulated with

10nM OVA257-264 peptide (AnaSpec, Fremont, CA, USA) in 10% RPMI (RPMI 1640 plus 10%

fetal bovine serum (Fisher), 2mM L-glutamine, 50U/mL penicillin/streptomycin, and 50μM β-

mercaptoethanol. Following 3 days of stimulation, cells were resuspended in complete media

plus 10 IU/mL recombinant human IL-2 (rHIL-2, Roche), and seeded in fresh media at 0.5×10^6

cells/mL every 48 hours. ID8 cancer cell line (ATCC) was grown in RPMI media (Gibco) with 10%

FBS (Gibco).

In-vitro activated CTLs from Ova-transgenic (OT-I) mice were combined with adherent ID8

murine ovarian cancer cells. To facilitate recognition of cancer cells by the OTI-I CTLs, ID8 cells

suspended in complete medium (RPMI with 10% FBS, Gibco) were incubated with $OVA_{257-264}$

SIINFEKL peptide for 1 hour at 37°C. The cells were washed three times with complete medium,

and 10^5 cells were added to each well of a 24-well plate. Each well contained a single sapphire

coverslip coated with human Fibronectin (Corning). The ID8 cells were allowed to settle and

adhere for 2 hours in an incubator at 37°C/5% CO_2. At this time, the media in the well was

replaced with 250uL phenol-red free RPMI (Gibco). Care was taken to ensure that media and

cells were stored in 37°C/5% CO_2 for all incubation periods. Immediately after changing the

media, 10^5 CTLs in 50uL complete medium were added to the well. The CTLs were allowed 7

minutes to find their targets and secrete lytic granules before fixation through high pressure

freezing. The relatively short incubation time favors capture of CTL:target conjugates in an

early-stage of interaction.

The remaining cryofixation, FS, and resin embedding procedures were identical to those

described for HeLa cells. The detail procedures are presented in the Supplementary

Information.

Tissue sample preparation

1. Mouse pancreatic islets

Pancreatic islets of 9-week-old C57BL/6 mice were isolated as previously described[20]. They

were cultured overnight in standard culture media (RPMI 1640 (Gibco) with 10% FBS, 20 mM

HEPES, 100 U/ml each penicillin and streptomycin) containing 5.5 mM glucose. Prior to high

pressure freezing the islets were subjected to 1 hr incubation in Krebs-Ringer buffer containing

either 3.3 mM or 16.7 mM glucose.

Islets were frozen with a Leica EM ICE high pressure freezer (Leica Microsystems, Germany) and

kept in liquid nitrogen until freeze substitution. Although freezing damage was observed in

couple nuclei, the preservation of cytoskeletal elements benefits from this fixation. High

pressure frozen islets were substituted as previously published[21] or according to a novel

protocol: first the samples were substituted in a cocktail containing 2% OsO_4, 1% UA, 0.5%

glutaraldehyde, 5% H2O in Acetone with 1% Methanol at -90°C for 24 hours. The temperature

was raised to 0°C over 15 hours followed by 4 washes with 100% acetone for 15 min each and

an increase in temperature to +22°C. Afterwards the samples were incubated in 0.2%

thiocarbohydrazide in 80% methanol at RT for 60 min followed by 6 x 10 min washes with 100%

acetone. The specimens were stained with 2% OsO_4 in acetone at RT for 60 min followed by

incubation in 1% UA in acetone + 10% methanol in the dark at RT for 60 min. After 4 washes in

acetone for 15 min each they were infiltrated with increasing concentrations of Durcupan resin

in acetone followed by incubation in pure Durcupan and polymerization at 60°C. For quality

control the blocks were sectioned with a Leica LC6 ultramicrotome (Leica microsystems) and

300 nm sections were put on slot grids containing a Formvar film. Tilt series ranging from -63°

to +63° were acquired with a F30 electron microscope (Thermo Fisher Scientific) and

reconstructed with the IMOD software package[22].

2. Drosophila brain samples (fan-shaped body and accessory calyx)

While thin tissue samples (< 200-μm-thick) can be cryo-fixed, in many cases thick tissue

specimens can not (e.g., large brain tissue) thus requiring various chemical fixation and staining

protocols. *Drosophila* brain tissues from 5-day-old adult (Genome type: iso Canton S G1 x

w1118 iso 5905) were prepared according to Progressive Lowering of Temperature and Low

Temperature Staining (PLT-LTS) progressive heavy metal enhancement protocol described

previously[23]. In details, after tissue dissection and pre-fixation, we osmicated tissue in 1% OsO_4,

then 1.5% K ferrocyanide, followed by a complete wash. Afterwards, a transfer to 1%

thiocarbohydrazide for 15 min at 22°C, then a complete wash, followed by 2% osmium for 30

min at 22°C. After osmication, we stained in lead aspartate for 30 min at 55°C first, then 1 h at

22°C. Finally, the tissue was dehydrated in a Leica AFS freeze-substitution chamber: the

temperature was dropped from 4°C to −25°C, and the concentration of acetone or ethanol was

increased for 20 min in each of 10%, 30%, 50%, 70%, 80%, 90%, and 97%. The subsequent low

temperature *en bloc* staining was performed in either 0.2% uranyl acetate in acetone, or 1%

EPTA in 97% ethanol. Specimens were infiltrated and embedded in Durcupan.

Preparation for FIB-SEM

After Durcupan re-embedding, a 3D X-Ray tomogram of the entire block was taken using an

XRadia-510 Versa micro X-Ray system (Carl Zeiss X-ray Microscopy, Inc.). The 3D X-Ray

tomograms allowed for robust Region of Interest (ROI) selection. Specifically, it identified cells

with good morphology (for example a properly shaped cell, a cell in a desired stage of mitosis, a

T-cell attacking a cancer cell, etc.), and also determined the proper orientation for the tissue

samples. Once a potential ROI was identified, the sample block was then re-mounted to the top

of a 1 mm copper post (using Durcupan) which was in contact with the metal-stained sample

for better charge dissipation, as previously described[5]. Each sample was oriented on the copper

post to allow the shortest distance along FIB milling direction. A small vertical sample post was

trimmed to the ROI with a width of ~100 μm and a depth of 60–80 μm in the direction of the

ion beam for each sample. Sample trimming used an ultramicrotome (EM UC7, Leica

Microsystems), was done in few iterations with X-Ray tomogram collected after each step.

Once the desired ROI was trimmed, a thin layer of conductive material of 10-nm gold followed

by 100-nm carbon was sputtered using a Gatan PECS 682 High-Resolution Ion Beam Coater. The

coating parameters were 6 keV, 200 nA on both argon gas plasma sources, 10 rpm sample

rotation with 45-degree tilt.

FIB-SEM system advances

A Zeiss Gemini 500 and multiple Zeiss Merlin SEM systems were customized for this work to

enable the two orders of magnitude improvement in imageable volume (from ~500 μm^3 to

greater than 100,000 μm^3) compared to our prior work[5]. Such improvement primarily comes

from: 1) higher precision and stability of FIB milling control to extend reliable long-term

acquisition to 4-nm voxels, and 2) enhanced SEM signal detection using secondary electrons to

achieve faster imaging, hence reduce electron radiation energy density (from ~400 to ~80

keV/nm^3) that further improves FIB milling control.

Specifically, to ensure precise and stable milling at 4-nm sampling, we integrated a more stable

FIB column (Capella from Zeiss) and repositioned at 90 degrees to the SEM column (Fig. 1a).

The Capella FIB column provided higher practical milling currents up to 30 nA compared to 7 nA

of the Magnum FIB column (Thermo Fisher Scientific, previously FEI) used in our prior work[5].

We further tightened its FIB emission current control band from ± 100 pA to ± 50 pA to obtain a

consistent milling beam profile. We also re-configured the milling closed-loop control (Fig. 10 in

Xu et al., 2017[5]) to further reduce the milling variation between FIB reheat cycles thus enabling

long-term acquisition at high electron radiation energy density. This FIB milling optimization

sufficiently reduced milling artifacts hence allowed for the switching of SEM signal detection

from backscattered electrons to secondary electrons for faster imaging (600 V vs. 0 V specimen

bias in Fig. 12 of Xu et al., 2017[5]). The new SEM signal detection scheme lowered electron

radiation energy density by 5x (from ~400 to ~80 keV/nm^3), which in turn further mitigated FIB

milling artifacts and instability. In addition, customized 100-µm-thick molybdenum apertures

were laser machined to define more accurate and consistent FIB milling currents, and to

achieve more than 2x longer lifetime; and the custom NI LabVIEW control software was re-

designed: to include new configurations of FIB milling closed-loop control which improve milling

variation; and to switch from the Zeiss RemCon serial port communication protocol to the Zeiss

API which reduced the overhead by several seconds per imaging/milling cycle.

The optimization of SEM imaging conditions was then guided by the following principles. The

isotropic resolution limits are convoluted by the waist size of the incoming primary electron

beam and the scattering volume of the penetrating primary electrons. While spherical and

chromatic aberrations of the electromagnetic lenses can be mitigated by smaller beam currents

at modest numerical aperture to achieve the best beam focus and x-y resolution, lower primary

beam landing energy generates fewer scattering events, resulting in smaller scattering volume

which improves resolutions along all three axes. However, any further reduction below the

optimal energy will result in a reduced scattering ratio of the heavy metal stain atoms over

lighter background atoms of the sample resin, thus fading contrast. Through investigating a

multi-dimensional operation space, we found that a beam current of 200 to 300 pA and a

landing energy of 700 to 900 eV are optimal for isotropic 4-nm imaging.

Such positive synergy between faster SEM scanning and robust FIB milling has extended reliable

imaging acquisition at 4-nm isotropic voxels from less than a week to months. Primarily limited

by time, the maximum volume can be seamlessly extended.

FIB-SEM imaging

As summarized in Table 1, all samples were imaged by a 0.14–0.3 nA electron beam with 0.7–

1.2 keV landing energy at 0.1–2.0 MHz. In most cases, both backscattered and secondary

electron signals were collected by InLens detector to provide better signal-to-noise ratio. The

time to acquire these volumes ranged from two to five weeks of uninterrupted imaging. Faster

SEM scanning rates (5x–10x scanning speed improvement) were applied on *Drosophila* samples

with stronger staining contrast while maintaining similar image quality, thus achieving

maximum volume of 175,000 μm^3 within three-week imaging.

For all samples, the x-y pixel size was set at 4 nm. A subsequently applied focused Ga+ beam of

15 nA at 30 keV strafed across the top surface and ablated away 4 nm of the surface. The newly

exposed surface was then imaged again. The ablation – imaging cycle continued for two to five

weeks of uninterrupted imaging to complete FIB-SEM imaging one sample. The sequence of

acquired images formed a raw imaged volume, followed by post processing of image

registration and alignment using a Fiji plugin based on Scale Invariant Feature Transform (SIFT)

algorithm[24] to form isotropic 4 nm voxels. The final aligned stack consisted of an isotropic

volume, containing multiple complete cells, which can be viewed in any arbitrary orientations.

Evaluation of FIB-SEM resolution

In order to evaluate the actual resolution in the acquired FIB-SEM datasets we analyzed the

transitions of the edges of the ribosomes abundant within the cell volumes. Ribosomes are

macromolecular machines consisting of RNA and associated proteins. Their small size (20–30

nm), abundance in the cytoplasm of living cells, and the fact that they are stained with high EM

contrast makes them good candidates for resolution evaluation.

First, we selected volumes of approximately 1000–2000 x 1000 x 1000 pixels in cultured cell

datasets. These volumes were selected in the middle of datasets in areas with large number of

ribosomes. We then calculated histograms of grey level values in these volumes and subtracted

~0.8 * histogram max to bring the average cytoplasmic material signal level close to 0. We then

used the Laplacian of Gaussian (LoG) algorithm (part of Python scikit-image package[25]) to select

blobs. We set the limit of standard deviation for LoG between 1 and 2.5 pixels (all these

datasets have 4-nm voxels) to ensure that selected blobs were predominantly ribosomes.

Filters were used to exclude the following blobs: average value below zero (this meant LoG

algorithm failed), edge value above 0.4 times of max value or 37%–63% transition longer than

15nm (the latter two happen where there is another blob or another feature with high signal

level in close proximity—making subsequent edge transition analysis inaccurate), amplitudes

below certain threshold, so that at least approximately 3000 blobs per dataset were remained

for analysis. The edge transitions in all three directions were analyzed for these remaining

blobs, in particular we determined 37%–63% transitions (the value used by Zeiss in their

resolution estimation) and 20%–80% transitions (close to 1 sigma value).

The summaries of the 37%–63% transitions and 20%–80% transitions for the 3D ribosome blobs

selected in cultured cell datasets are plotted in Fig. 1b. The histograms of the 37%–63%

transitions for these datasets are shown in Extended Data Fig. 3. The examples of 3D blobs for

each dataset are presented in Extended Data Fig. 4–10. For each dataset the cross-sections and

transition analysis for 3 brightest and 3 dimmest ribosomes are shown.

Accurate determination of FIB-SEM milling rate

The position of a sample block face was measured by the SEM beam and the FIB beam

independently. An in-line auto focus routine[6] kept the SEM images in focus throughout the

entire image acquisition. While the working distance of the SEM beam detected the block face

position by measuring its distance from the SEM objective lens, the FIB milling position under

closed-loop control indicated the block face position directly from the orthogonal angle. As the

sample block face being imaged then milled away layer-by-layer, the SEM working distance and

the FIB beam position tracked the block face location accordingly. Therefore, we estimated the

total thickness milled away by FIB based on the changes of either SEM working distance or FIB

milling position from the beginning to the end of the FIB-SEM acquisition. The average milling

rate of z-step was then calculated as the total milling depth divided by the number of image

frames. Here we report the average z-scaling factors derived from both measurement

approaches in Extended Data Table 1, which can be used to scale the dataset to generate real

isotropic voxels. For example, HeLa Cell 2017-06-21 dataset has a z-scaling factor of 1.31 which

means the actual z-step is averaged at 5.2 nm instead of 4.0 nm.

Image emulation to visualize the value of finer resolution

To emulate 8-nm images, we first applied a 6-nm Gaussian blur filter and then added Gaussian noise with a standard deviation of 22 to 4-nm images. The higher shot noise at 8-nm sampling was expected due to the 10x lower electron dose of ~12 e/nm^3, compared to that of ~120 e/nm^3 at 4-nm sampling. The 6-nm Gaussian blur and added noise (Fig. 1d, g, and j) were underestimated compared to that of the real 8-nm images (Fig. 1e, h, and k) collected on the same type of specimens. The 8-nm emulated images with comparisons to 4-nm images and real 8-nm images are shown in Fig. 1c–k and Supplementary Video 1.

Data availability

All raw and segmented datasets in this work are available on OpenOrganelle website (https://openorganelle.janelia.org).

Code availability

FIB-SEM image acquisition LabVIEW code used in this work is available from

https://github.com/cshanxu/Enhanced_FIB-SEM

Python code for resolution characterizations using ribosomes is available from

https://github.com/gleb-shtengel/FIB-SEM_resolution_evaluation

Acknowledgements

We thank K. Hayworth and W. Qiu at Howard Hughes Medical Institute (HHMI) Janelia Research Campus (JRC) for invaluable discussions and data collection support. We gratefully acknowledge P. Rivlin, S. Plaza, and I. Meinertzhagen for JRC EM shared resource and FlyEM project team

462   support on staining protocols development. We thank the electron microscopy facility of MPI-

463   CBG and of the CMCB Technology Platform at TU Dresden for their services. We also thank Y.

464   Wu from Pietro De Camilli's laboratory at Yale for advice.

Funding

C.S.X., S.P., G.S., S.T., Z.L., H.A.P., N.I., D.B., A.V.W., M.F., T.C.W., J.L.-S., H.F.H. are funded by

Howard Hughes Medical Institute (HHMI). A.M. received support from the Carl Gustav Carus

Faculty of Medicine at TU Dresden via a MeDDrive GRANT. A.M. and M.S. were supported with

funds from the German Center for Diabetes Research (DZD e.V.) by the German Ministry for

Education and Research (BMBF), from the Germain-Israeli Foundation for Scientific Research

and Development (GIF) (grant I-1429-201.2/2017), from the German Research Foundation

(DFG) jointly with the Agence nationale de la recherche (ANR) (grant SO 818/6-1) to M.S.. H.K.H

and S.B.v.E. are funded by NIAID grant R01AI138625. R.V.F. is supported by NIH R01GM124348.

T.C.W. is supported by NIH R01GM097194. J.C. is a fellow of the Damon Runyon Cancer

Research Foundation.

Author Contributions

C.S.X. and H.F.H. supervised the project; C.S.X., S.P., G.S., and H.F.H. wrote the manuscript with

input from all co-authors; C.S.X. developed enhanced FIB-SEM platform for large volume high

resolution imaging, and optimized imaging conditions; C.S.X., S.P., and G.S. conducted FIB-SEM

experiments; C.S.X. and S.P. performed image post processing; S.P., G.S., A.M., A.T.R., H.K.H.,

S.B.v.E., Z.L., H.A.P., N.I., J.C., A.V.W., and M.F. prepared samples; G.S., A.M., A.T.R., and S.T.

analyzed data; D.B. prepared and uploaded data to OpenOrganelle website; S.B.v.E., T.C.W.,

R.V.F., J.L.-S., I.M., and M.S. proposed biological questions and provided samples.

484   Competing interests

Figures

Fig. 1 **Enhanced FIB-SEM configuration, operation, and resolution**. **a**, Sketch of FIB milling and

SEM imaging. The two operations iterate alternately to generate 3D whole cell image stacks, **b**,

Resolution characterization using transitions at the edges of gold nanoparticles on a carbon

substrate and ribosomes in cultured cells. The vertical dash lines indicate the edge transitions of

gold nanoparticles, the round dots indicate the edge transitions of ribosomes along x-y and z

axes. Each dot represents the average value from more than 3000 ribosomes in one of the

seven cultured cell samples. **c–k**, Comparison of lower current/higher resolution 4-nm sampling

and 8-nm sampling. Left column shows images acquired at 4-nm sampling that matches the

resolutions shown in **b**. The middle column shows the emulated 8-nm images (details in

Methods Section), while the right column shows the real 8-nm sampling images. **c**, Nucleus of

the interphase HeLa cell from Fig. 2d shows fine details of chromatin (Ch), nuclear membrane

(NM), and nuclear pore (NP). Nucleosomes are less resolved in **d,** the emulated 8-nm sampling

image, and in **e**, the real 8-nm sampling image of an interphase HeLa cell. **f**, Golgi (G) cisternae

of the interphase HeLa cell from Fig. 2c are better resolved at 4-nm sampling compared to the

emulated 8-nm sampling image shown in **g** and the interphase HeLa cell imaged at 8-nm

sampling shown in **h**. **i**, *Drosophila* brain sample from Supplementary Video 1 shows well-

resolved hollow core of a microtubule (MT) and the close contract between endoplasmic

reticulum (ER) and mitochondria (Mito), which are not distinguishable in **j**, the emulated 8-nm

sampling image. **k**, The real 8-nm sampling image of blurry microtubules of a *Drosophila* brain

sample with higher shot noise and less resolved microtubules compared to those shown in **j**.

Scale bar, 500 nm in all images.

Fig. 2 **Interphase HeLa cell**. **a**, FIB-SEM overview with cutaway, and manually segmented

interior features (mitochondria, green; centrosomes, red; one cistern of a Golgi stack, magenta;

a segment of nuclear membrane, blue, polyribosome chains, yellow). Zoomed in cross-sections

of three select areas containing: **b**, Centrosomes, **c**, Golgi stack, and **d**, Nucleus, polyribosomes

indicated by arrows, and nuclear pores indicated by asterisks. Scale bars are 1 μm.

519   Fig. 3 **Murine CTL engaging an ovarian cancer cell**. **a,** FIB-SEM overview with cutaway, and

520   manually segmented membranes of CTL (green surface and red contour) and cancer cell (cyan

521   surface and blue contour). Zoomed in cross-sections highlight the signatures of a productive

522   immunological synapse: **b**, CTL cell "cupping" the target cancer cell, **c,** Polarized centrosome,

523   and **d**, Lytic granule showing unique ultrastructures. Scale bars are 1 μm.

Fig. 4 **Tissue sample datasets**. **a**, Pancreatic islets treated with 16.7 mM glucose containing

several complete beta cells with detailed structures shown in **b**, A primary cilium with axoneme

(arrow) and centrioles (asterisk), **c**, Intermingled microvilli (arrow), **d**, Ultrastructural diversity

among insulin SGs containing rod-shaped (arrow) or spherical crystals (asterisk), **e**, Contacts of

ER and insulin SGs (arrow), **f**, Microtubules (arrow), Golgi apparatus (asterisks) and ribosomes.

g, *Drosophila* fan-shaped body with detailed structures shown in **h**, Dense-core vesicles of

different sizes (arrows), **i**, Multiple synaptic sites viewed from the side (arrows), **j**, A presynaptic

T-bar viewed from the top (arrow), **k**, A longitudinal-section view of microtubules (arrows), **l**, A

cross-section view of microtubule arrays (arrows). Scale bar, 1 μm in **a**, 500 nm in **b**–**f**, 5 μm in

g, 200 nm in **h**–**l**.

Extended Data Fig. 1 **Isotropic voxel (representing the minimal voxel size dictated by the worst-case axial resolution) vs. volume for comparing different volume EM methods**. The light green space represents the Resolution-Volume regime accessible with enhanced FIB-SEM technology through long term imaging. The present work of whole cell volumes at 4-nm isotropic voxels matching the resolutions shown in Fig. 1b is colored in yellow, compared to the prior work of smaller volumes colored in red. Adopted from Xu et al., 2017[5] with modifications.

543 Extended Data Fig. 2 **Murine CTL engaging an ovarian cancer cell.** Zooms on regions showing

544 different immunological synapse topology features. **a**, Interdigitation, **b**, Flat apposition, and **c**,

545 Filopodia caught between cells. Scale bars are 0.5 μm.

Extended Data Fig. 3 **Edge transition distributions determined from ribosomes in cultured cells**

datasets. Distributions of 37%–63% (**a**) and 20%–80% (**b**) transition distances in X-, Y- (left), Z_{top}-

(center), and Z_{bot}- (right) directions.

Extended Data Fig. 4 **Cross-sections of the example of ribosomes from the dataset Interphase**

HeLa Cell 2017-06-21 and the profiles with the transition analysis. The top three rows are the

brightest ribosomes and the bottom three rows are the dimmest ribosomes.

Extended Data Fig. 5 **Cross-sections of the example of ribosomes from the dataset Interphase**

HeLa Cell 2017-08-09 and the profiles with the transition analysis. The top three rows are the

brightest ribosomes and the bottom three rows are the dimmest ribosomes.

559 Extended Data Fig. 6 **Cross-sections of the example of ribosomes from the dataset Wild-type**

560 **THP-1 Macrophage 2018-11-11 and the profiles with the transition analysis.** The top three

561 rows are the brightest ribosomes and the bottom three rows are the dimmest ribosomes.

Extended Data Fig. 7 **Cross-sections of the example of ribosomes from the dataset**

Immortalized T-cells (Jurkat) 2018-08-10 and the profiles with the transition analysis. The top

three rows are the brightest ribosomes and the bottom three rows are the dimmest ribosomes.

567   Extended Data Fig. 8 **Cross-sections of the example of ribosomes from the dataset**

568   **Immortalized breast cancer cell (SUM159) 2017-11-21 and the profiles with the transition**

569   **analysis.** The top three rows are the brightest ribosomes and the bottom three rows are the

570   dimmest ribosomes.

Extended Data Fig. 9 **Cross-sections of the example of ribosomes from the dataset Killer T-cell**

attacking cancer cell 2020-02-04 on Cancer Cell and the profiles with the transition analysis.

The top three rows are the brightest ribosomes and the bottom three rows are the dimmest

ribosomes.

Extended Data Fig. 10 **Cross-sections of the example of ribosomes from the dataset Mitotic**

HeLa cell 2019-05-30 and the profiles with the transition analysis. The top three rows are the

brightest ribosomes and the bottom three rows are the dimmest ribosomes.

Tables

Table 1. Descriptions of exemplary datasets with selected imaging conditions

Sample Description	Protocol	Landing Energy (kV)	Imaging Current (nA)	Scan Speed (MHz)	Acquisition Duration (days)	Data Volume: X * Y * Z (um^3)	Z-scaling factor
Cell 4 nm							
Interphase HeLa cell 2017-06-21	High pressure freezing, freeze-substitution with 2% OsO4 0.1% UA 3% H2O in Acetone, resin embedding in Eponate 12	0.9	0.24	0.1	25	48 * 6 * 25 (7800)	1.31
Interphase HeLa cell 2017-08-09†		1.0	0.24	0.2	31	50 * 4 * 48 (9600)	0.81
Mitotic HeLa cell 2019-05-30		0.9	0.24	0.2	15	30 * 14 * 33 (13300)	1.07
Wild-type THP-1 macrophage 2018-11-11		1.2	0.24	0.2	20	40 * 8 * 44 (14200)	0.84
Immortalized T-cells (Jurkat) 2018-08-10		1.0	0.24	0.2	19	40 * 12 * 34 (16400)	0.86
Immortalized breast cancer cell (SUM159) 2017-11-21‡		1.2	0.24	0.2	19	64 * 11 * 31 (21900)	1.14
Killer T-Cell attacking cancer cell 2020-02-04†		0.7	0.25	0.2	29	74 * 13 * 48 (45500)	0.87
Tissue 4 nm							
Isolated murine pancreatic islets 2019-03-01†	HPF, FS with 2%OsO4, 1% UA, 0.5% GA, 5% H2O in Acetone with 1% Methanol followed by 0.2% TCH in 80% Methanol at RT, 2% OsO4 in Acetone at RT, 1% UA in Acetone + 10% Methanol	0.9	0.14	0.2	15	30 * 20 * 29 (17400)	0.85
Drosophila brain: Fan-shaped body of a 5-day-old male adult 2019-09-14†	Chemical Fixation, ORTO-Lead-EPTA staining via Progressive Lowering of Temperature and Low Temperature Staining (PLT-LTS) heavy metal enhancement protocol	0.7	0.24	1.5	15	45 * 56 * 45 (113400)	1
Drosophila brain: Accessory calyx of a 5-day-old male adult 2019-12-06		0.7	0.3	2	19	50 * 40 * 87 (175000)	0.93

584   † Figures in main text.

585   ‡ Treated with 0.5 mM oleic acid for 45 mins prior to high pressure freezing to induce the formation of lipid

586   droplets.

587   Extended Data Table 1. Z scaling factors estimated using SEM working distance and FIB milling

588   voltage.

DataSet	Z (milling direction) pixel scaling factor		
	SEM Working Distance (WD)	FIB milling voltage (MV)	Mean (WD, MV)
Interphase HeLa cell 2017-06-21	1.22	1.40	1.31
Interphase HeLa cell 2017-08-09	0.80	0.83	0.81
Mitotic HeLa cell 2019-05-30	1.07	1.07	1.07
Wild-type THP-1 macrophage 2018-11-11	0.81	0.87	0.84
Immortalized T-cells (Jurkat) 2018-08-10	0.84	0.88	0.86
Immortalized breast cancer cell (SUM159) 2017-11-21	1.13	1.14	1.14
Killer T-Cell attacking cancer cell 2020-02-04	0.88	0.87	0.87
Isolated murine pancreatic islets 2019-03-01	0.86	0.84	0.85
Drosophila Fan-shaped body 2019-09-14	0.98	1.03	1.00
Drosophila Accessory calyx 2019-12-06	0.95	0.92	0.93

 ## References

[19] Sartori, N., Richter, K., & Dubochet, J. Vitrification depth can be increased more than 10-fold by high-pressure freezing. *J. Microsc.* **172**(1), 55–61 (1993). https://doi.org/10.1111/j.1365-2818.1993.tb03393.x

[20] Gotoh, M., Maki T., Kiyoizumi, T., Satomi, S., & Monaco, A. P. An improved method for isolation of mouse pancreatic islets. *Transplantation* **40**, 437–438 (1985). https://doi.org/10.1097/00007890-198510000-00018

[21] Müller, A. et al. A global approach for quantitative super resolution and electron microscopy on cryo and epoxy sections using self-labeling protein tags. *Sci. Rep.* **7**, 23 (2017). https://doi.org/10.1038/s41598-017-00033-x

[22] Kremer, J. R., Mastronarde, D. N., & McIntosh, J. R., Computer visualization of three-dimensional image data using IMOD. *J. Struct. Biol.* **116**, 71–76 (1996). https://doi.org/10.1006/jsbi.1996.0013

[23] Lu, Z. et al. En bloc preparation of *Drosophila* brains enables high-throughput FIB-SEM connectomics. *bioRxiv*: 855130 (2019). https://doi.org/10.1101/855130

[24] Saalfeld, S., Cardona, A., Hartenstein, V. & Tomančák, P. As-rigid-as-possible mosaicking and serial section registration of large ssTEM datasets. *Bioinformatics* **26**, i57–63 (2010). https://doi.org/10.1093/bioinformatics/btq219

[25] van der Walt, S. et al. scikit-image: image processing in Python. *PeerJ*. **2**:e453 (2014). https://doi.org/10.7717/peerj.453