TCRβ Sequencing Reveals Spatial and Temporal Evolution of Clonal CD4 T cell Responses in a Breach of Tolerance Model of Inflammatory Arthritis

Authors: Shaima Al-Khabouri[#], Robert A. Benson*, Catriona T. Prendergast, Joshua I. Gray[$], Thomas D Otto, James M Brewer, Paul Garside[1]

Corresponding author information:

Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary and Life Sciences, Sir Graeme Davies Building, University of Glasgow, 120 University Place, Glasgow G12 8TA, United Kingdom

Email address: shaima.al.khabouri@ki.se

Affiliations:

[1]Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary and Life Sciences, Sir Graeme Davies Building, University of Glasgow, 120 University Place, Glasgow G12 8TA, United Kingdom

[#]Current address: Department of Medicine Solna (MedS), Division of Rheumatology, Center for Molecular Medicine (CMM), Karolinska Institute,171 76 Stockholm, Sweden

*Current address: Antibody Analytics Ltd, BioCity Scotland, Bo'ness Road, Newhouse, ML1 5UH, United Kingdom

[$]Current address: Columbia Center for Translational Immunology, Columbia University, New York NY, USA 10032

KEY MESSAGES

- Selective accumulation of CD4 T cell clones has been observed in arthritic joints of RA patients, indicating the presence of antigen driven accumulation of CD4 T cells in these patients.

- These antigen specific T cell clones are thought to be pathogenic and thus are targets for tolerogenic therapy.

- This study suggests that CD4 TCR clonality is restricted at the early phases of disease and broadens with disease progression. Moreover, changes in CD4 TCR clonality are first reflected in joint draining lymph nodes before being mirrored in the inflamed joint.

- Thus tolerogenic therapies will be more effective when CD4 TCR clonality is restricted i.e. at early disease stages.

- Changes in CD4 TCR clonality in the joint draining lymph node can also be a biomarker for disease state.

ABSTRACT

Objectives: Development of effective tolerogenic therapies for Rheumatoid Arthritis (RA) relies on understanding the evolution of articular antigen specific CD4 T cell responses. TCR clonality of the endogenous CD4 T cell infiltrate in early inflammatory arthritis was assessed to monitor the evolution of the TCR repertoire in the inflamed joint and its associated popliteal lymph node (pLN).

Methods: Mouse models of antigen-induced breach of self-tolerance and chronic polyarthritis were used to recapitulate the early and late phases of RA reported in patients. The infiltrating endogenous, antigen experienced CD4 T cells in inflamed joints and pLNs were analysed using flow cytometry and TCRβ sequencing.

Results: TCR repertoires from pLNs displayed increased clonality and diversity with disease progression, while inflamed joints maintain similar TCR repertoire clonality and diversity with time. Repertoires from late phase pLNs accumulated clones with a diverse range TRBV genes, while inflamed joints at both phases contain clones expressing similar TRBV genes. Repertoires from pLNs and joints at the late phase displayed reduced CDR3β sequence overlap compared to the early disease phase, however correlation analysis revealed the most abundant clones in pLNs accumulate in the joint at the later phase.

Conclusions: CD4 T cell clonality broadens and evolves with progression of inflammatory arthritis and is reflected in pLNs before potentially being mirrored in the joint. These observations imply that antigen specific tolerogenic therapies for RA will be more easily developed and more effective at earlier phases of the disease, when CD4 T cell clonality is least diverse.

Keywords: TCRβ sequencing, T cell clonality, early rheumatoid arthritis

INTRODUCTION

Rheumatoid arthritis (RA) is a chronic inflammatory autoimmune disease characterised by synovial inflammation and cartilage and bone erosion, causing progressive loss of joint function[1]. Antigen presentation and CD4 T activation mechanisms have been shown to play a role in the pathogenesis of RA. This is evidenced by genetic association studies in RA patients showing strong associations of HLA-DRB alleles, T cell activation associated genes such as *PTPN22* and *CTLA-4,* and loci associated with signal transduction such as *STAT4*[2–7]. Moreover, the successes of therapeutics in modulating T cell activation, such as the use of Abatacept, reinforce the importance of CD4 T cells in propagating the disease[8]. As such, there has been increasing interest in targeting pathogenic autoreactive CD4 T cells to provide a more effective line of therapy. Specifically, there has been increasing interest in utilising antigen specific tolerogenic therapies to target autoreactive CD4 T cells in RA to re-establish tolerance and provide long-lasting remission, while retaining protective immunity against pathogens. However, targeting autoreactive CD4 T cells has proven difficult in RA, due to the broad range of antigens implicated in disease, lack of a clear cellular hierarchy of disease drivers, and also patient to patient variation[9–13]. Moreover, it is unknown at what stages of the disease these autoreactive responses develop nor the location in which these responses are primed, thus hampering the development of effective tolerogenic therapies. Antigen specific responses can be identified by detecting expanded clonal T cell populations. Indeed, oligoclonal CD4 T cell responses and skewed TCR repertoires have been reported in arthritic joints[14–17]. More recently, analysis of CD4 T cell behaviour was shown to reflect antigen recognition in inflamed joints at the early stages of disease[18], suggesting that antigen specific CD4 T cell responses may perpetuate and drive progression of RA. However, the development of antigen specific responses and their evolution with the progression of RA is poorly understood. Investigating the evolution TCR clonality of antigen experienced CD4 T cells in RA will provide insight on how antigen specific responses evolve with disease progression. This will inform the development of more effective tolerogenic therapies by indicating the range of clones that would need to be targeted and the disease stage at which these therapies are most likely to be effective. In this study, we employed breach of self-tolerance models of antigen induced inflammatory arthritis[19,20] and chronic polyarthritis[21] in which autoreactive responses develop. These models also recapitulate the early and later stages of the disease – hereafter named the early and late phases respectively. TCRβ sequencing was used to monitor the evolution of antigen specific endogenous CD4 T cell responses in inflamed joints and their associated draining lymph nodes.

RESULTS

Induction of inflammatory arthritis results in local accumulation of endogenous antigen experienced CD4 T cells at both the early and late phases

Induction of the breach of tolerance model of inflammatory arthritis requires the transfer of Th1 polarised ovalbumin (OVA) specific OT-II cells and subsequent immunisations with OVA and heat aggregated OVA (HAO) as outlined in **figure 1 A**. This results in the influx of the transferred OT-II cells and endogenous CD4 T cells of varying antigen specificities, indicated by the range of expressed TCRβ variable genes (TRBV)[18]**.** Assessing the evolution of TCR clonality of the endogenous CD4 T cell population requires the accurate identification of these cells in joints and popliteal lymph nodes (pLN). which were distinguished from the transferred OT-II population using the CD45.1 congenic marker **(supplementary figures 1 and 2)**. Endogenous CD4 T cells were present at a significantly higher number than the transferred OT-II cells in inflamed pLNs and joints at both the early and late phases **(figure 1 B)**. A significant number of the endogenous CD4 T cells at the early phase also displayed an antigen experienced phenotype, indicated by increased CD44 expression, after induction of inflammatory arthritis with HAO **(figure 1 C)**, and were previously shown to produce TNFα and IFNγ following ex-vivo stimulation with PMA/ionomycin[18]. The number of endogenous CD4, CD44hi cells was also increased at the late phase in inflamed joints **(figure 1 C)**.

Accumulation of expanded clones and observed clonality and diversity of CD4 T cell repertoires is antigen driven

To investigate the evolution of the CD4 TCR clonality in this model of inflammatory arthritis, endogenous antigen experienced CD4 T cells i.e. excluding the transferred OT-II cells, were sorted from pLNs and joints of individual mice for both early and late phases. Total RNA was then isolated from the endogenous CD4 T cells and used to sequence the CDR3β region of the TCR, the most variable region of the TCR that is critical for the recognition of specific peptide sequences[22,23]. Repertoires were then characterised by examining the number of unique CDR3β sequences and evaluating the degree expanded clones contributed to the overall antigen experienced repertoire, and also by assessing clonality and diversity of the repertoire. In this study, clonality is defined by the number of unique CDR3β sequences found in the population, and diversity – represented by the Diversity Index (DI) (see Methods) ⬚ represents the relative contribution of each unique CDR3β sequence, or clone, to the population. Thus, a repertoire with a larger number unique CDR3β sequences and therefore clones would be described to have high clonality. This repertoire would also display high diversity if the number of cells contributing to different clonal populations are similar. Due to the low recovery of antigen experienced endogenous CD4 T cells isolated from control pLNs and joints at the early phase, cells

from control pLNs or joints needed to be pooled prior to sequencing. Pooling samples changes the composition of the repertoire and affects T cell repertoire diversity, so comparisons of control CD4 T cell populations were made against pooled data from inflamed joint and pLN samples at the early phase **(figure 2 and table 1)**.

Table 1 – Clonality, diversity, and number of End. CD4 T cells in early inflamed pLNs and joints and PBS controls

Sample	pLN HAO	pLN PBS	Joint HAO	Joint PBS
No. of End. CD4+ T cells	3,113,218	224,978	58,041	1424
Clonality	4181	4347	4549	885
Diversity index (DI)	6.4	16.3	0.72	9.3

Data represents 1 experiment with pooled samples from 3 mice for the PBS sample and 5 mice for the inflamed (HAO) sample.

Clonality of the endogenous CD4 T cell repertoire from inflamed pLNs and controls were comparable (4181 unique CDR3β sequences in the inflamed pLN repertoire vs 4347 in the PBS pLN repertoire), however, the repertoire from the inflamed pLN showed a reduced diversity compared with the control pLN sample (6.4 vs 16.3 respectively). Endogenous CD4 T cell repertoires from inflamed joints displayed higher clonality than controls (4549 vs 885). Despite this, the diversity of the repertoire in the inflamed joint was relatively low compared to the PBS joint (0.72 vs 9.3 respectively). These data therefore demonstrate that accumulation of clones can be detected at the early phases of the disease in both inflamed pLNs and joints.

It is widely accepted that disease progression in RA results in increased joint damage and as a consequence, new antigens are exposed[1]. However, evidence for the accumulation of antigen experienced T cells in the absence of their cognate antigen has been demonstrated in this model[18] and in RA patients[24–26] which may be a result of a chronic inflammatory environment[27,28]. To address whether the accumulation of clones is antigen driven rather than a result of non-antigen induced inflammation, the antigen experienced repertoires of pLNs and joints at the late phase were compared to repertoires isolated from mice challenged with IFA alone **(figure 3 A)**, where inflammation is induced without the presence of antigen. No significant difference was found between the contribution of the top 10 most expanded clones to the repertoires of inflamed pLNs and IFA controls (data not shown) nor when comparing the contribution of clones occurring at the lowest frequency **(figure 3 B)**. To test for antigen driven expansion, clones with a frequency of 2 or more or 10 or more were compared as these were the median frequencies of "expanded" clones in the PBS pLN and joint samples respectively with the assumption being that there should be no antigen-driven clonal expansion of CD4 T cells

in the PBS samples. No difference was found between inflamed pLNs and IFA controls at either of the expanded frequencies selected (*p value 0.32 and 0.23 respectively Welch two sample t-test*). Moreover, repertoires from inflamed pLNs at the late phase and the IFA controls also displayed no differences in clonality or diversity **(figures 3 C and D)**. The top 10 highly expanded clones from repertoires of inflamed joints at the late phase, however, contribute approximately 41% to the overall repertoire compared to joint IFA controls, which only comprise an average of 29% of the entire repertoire **(figures 3 E and F)**. There was no difference in clonality between repertoires from inflamed joints and IFA controls **(figure 3 G)**, but inflamed joints have reduced clonal diversity compared to IFA controls **(figure 3 H)**. Thus, there is evidence to suggest accumulation of expanded clones in inflamed joints at the late phase in addition to the early phase and this accumulation is antigen driven.

The antigen experienced endogenous CD4 T cell repertoire displays disparity in clonal diversity between inflamed pLNs and joints with time

Antigen experienced endogenous CD4 T cell repertoires from pLNs and joints were compared between the early and late phases to investigate how CD4 TCR clonality evolves with disease progression **(figures 4 A and E)**. When comparing inflamed pLNs at the early and late phases, the top 10 most expanded clones contributed approximately 18% to the overall antigen experienced repertoire of inflamed pLNs at the early phase compared to only 2% at the late phase **(figures 4 A and B)**. When comparing clonality and diversity, repertoires from pLNs isolated at the late phase displayed increased clonality and diversity compared with their early counterparts **(figures 2 C and D)** and comprised mainly of low abundance clones. In contrast to the pLNs, antigen experienced CD4 T cell repertoires isolated from inflamed joints at the early and late phases were similar; the top 10 most expanded clones dominated the repertoires and comprised on average 46% and 41% of the overall populations **(figures 4 E and F)**. Furthermore, no differences were observed between the clonality and diversity of the repertoires at the early and late phases **(figures 4 G and H)** and repertoire diversity overall was low in the joint at both phases. Together, these data highlight the disparity in repertoire composition between pLNs and joints at the early and late phases, namely that pLNs at the late phase displayed high repertoire clonality and diversity compared to their early counterparts and also compared to inflamed joints at the same phase.

Disparity in clonality of the antigen experienced endogenous CD4 T cell repertoires is attributed to accumulation of clones with different TCRβ variable (TRBV) genes

Characterisation of the antigen experienced CD4 T cell repertoires from pLNs and joints at the early and late phases provided useful information on the overall distribution of clones

in these repertoires. However, this does not provide any insight on the TCR sequence of these clones, nor on whether these clones have similar or different TCR sequences between the two sites. Moreover, no insight is gained on whether the clonal composition of the antigen experienced repertoires in these two sites change over time. To address this, a PCA analysis was performed on inflamed pLNs and joints at both the early and late phases on the basis of TRBV gene use **(figure 5 A)**. Repertoires isolated from inflamed pLNs at the late phase displayed the largest range in TRBV gene use, while inflamed pLNs and joints at the early phase as well as inflamed joints at the late phase grouped together in terms of TRBV gene expression. Indeed, when quantifying these differences, inflamed pLNs at the late phase had significantly increased presence of TRBV genes 1, 2, 3, 5, and 19, while TRBV genes 12-1, 12-2, and 13-1 were present in the highest frequencies in inflamed pLNs at the early phase **(figure 5 B)**. These differences are also due to changes in TRBV gene use and not in TRBJ genes **(figure 5 C)**. Interestingly, the PCA analysis also highlighted differences between samples in terms of TRBV gene use in inflamed pLNs at the late phase. Accumulation of clones expressing different TRBV genes in inflamed pLNs at the late phase suggests changes in antigen specific responses with time and between inflamed pLNs and their associated tissues. Moreover, these changes in antigen specific responses also differ between individual animals with time.

Antigen associated changes in the inflamed pLN may predict the changes in antigen specificities and clonality of the CD4 T cell repertoire in inflamed joints

To assess whether changes in CD4 TCR clonality reflected changes antigen specificities, the degree of CDR3β amino acid sequence overlap between the antigen specific CD4 T cell repertoires was investigated. Comparing the repertoire overlap between inflamed pLNs and joints at both the early and late phases allowed us to monitor clonal dynamics between these two sites with the progression of inflammatory arthritis. The number of overlapping CDR3β amino acid sequences is represented by the normalised overlap index (see methods) and this decreases between the inflamed pLN and its joint with the progression of inflammatory arthritis **(figures 6 A and B)**. CDR3β sequence overlap decreases significantly between samples when comparing repertoires from inflamed pLN samples at the early phase to inflamed pLNs at the late phase **(supplementary figure 3 A)**, while the degree of CDR3β amino acid sequence overlap remains unchanged between repertoires from inflamed joints at the early and late phases **(supplementary figure 3 B)**. These observations confirm the observed changes in TRBV gene use and also imply changes in antigen specific responses in pLNs, which progress towards polyclonality with disease progression. Correlation analysis between the top 10 most expanded clones in the inflamed joint at the late phase correlated significantly with the most expanded clones in the inflamed pLNs at the same phase **(figure 6 D)**. Furthermore, the 10 most expanded clones in inflamed joints at the early phase and those from IFA

controls did not correlate with the most highly expanded clones in their respective pLNs **(figures 6 C and E)**. Moreover, the 10 most expanded clones in inflamed joints at the early phase were not all present in their respective pLNs **(supplementary table 1)**, but the top 10 expanded clones in inflamed joints at the late phase were all present in inflamed pLNs at the same phase **(supplementary table 2)**. These results indicate that despite pLNs progressing towards polyclonality with the development of disease, the most highly expanded antigen experienced clones accumulate in the inflamed joint. Thus repertoire clonality in the pLN may eventually be mirrored in the inflamed joint with disease progression.

DISCUSSION

Characterising CD4 TCR clonality has been used to monitor progression of disease states in cases of infection[29] and autoimmunity[30,31], including RA[15,32–35]. Although CD4 T cell clonality has been determined in RA patients at different stages of the disease[15,17], the development of antigen specific responses from very early, pre-clinical stages to a more established disease stage remains unknown. Understanding this is important to develop effective tolerogenic therapies as the range of CD4 T cell clones to target and the timepoint this therapy will have the greatest impact can be determined. Thus, we investigated the evolution of CD4 TCR clonality by sequencing CDR3β regions of antigen experienced endogenous CD4 T cells isolated from pLNs and joints at an early and late phases of breach of self-tolerance models of inflammatory arthritis.

Induction of inflammatory arthritis resulted in the accumulation of endogenous CD4 T cells displaying an antigen experienced phenotype. This population has been previously shown to produce TNFα and IFNγ and also harbour autoreactive clones capable of causing bone and cartilage degradation[18–21,36,37]. CDR3β analysis of this this population revealed that TCR repertoires from inflamed joints at both the early and late phases displayed low clonality and diversity and were dominated by few clones present at high frequencies. In contrast, pLNs at the late phase displayed higher clonality and diversity compared with their counterparts at the early phase and were mainly composed of a large number of clones that were present at low frequencies. This indicates that pLNs become more polyclonal with the progression of inflammatory arthritis and this polyclonality occurs in the pLNs before possibly being reflected in inflamed joints. This is reinforced by the observations that the most frequently occurring clones in the inflamed joint at the late phase are also the most frequently occurring clones in the inflamed pLN at the same phase. In addition, a study by Klarenbeek et. al[15] demonstrated that repertoires from synovial fluid of patients with established RA have fewer highly expanded clones compared with the synovial fluid taken from patients 6 months after showing clinical symptoms, demonstrating the reduced clonality of CD4 T cell repertoires in the joints of established RA patients. The polyclonality observed in pLNs at the late phase may reflect changes in antigen specific responses due to the release of neo-antigens resulting from continued joint damage and/or epitope spreading, a phenomenon observed in RA patients[38]. Indeed, PCA analysis of TRBV genes of repertoires isolated from inflamed pLNs and joints at the early and late phases highlighted the expanded range of TRBV genes present in repertoires of pLNs at the late phase, indicating a change in antigen specific responses with the progression of inflammatory arthritis. However, this can also be explained by the re-establishment of lymphatic recirculation. Given that the observations reported were exclusively on the antigen experienced CD4 T cell population,

re-establishment of lymphatic recirculation is an unlikely explanation for the changes in the repertoire clonality and diversity observed. Moreover, the 10 most abundant antigen experienced clones correlate significantly between inflamed pLNs and joints at the late phase and not in IFA controls. Inflamed joints were also shown to have reduced diversity in the antigen experienced repertoire compared to joint IFA controls. By taking these observations together, one can deduce that changes in pLN repertoire clonality, diversity, and TRBV genes is in fact antigen driven.

CDR3β amino acid sequence overlap analysis between repertoires isolated from pLNs and joints at early and late phases of inflammatory arthritis highlighted that fewer clones are shared between pLNs and joints at the late phase compared to the early phase. This indicates that antigen specific responses differ between these sites with time, driven by changes in clonality, diversity, and clones with different TRBV genes in the pLN. Moreover, the reduced clonal overlap at the late phase highlight the fact that inflamed pLNs and joints display different antigen specific responses at the same phase. This disparity has important implications for antigen specific therapies as specific pathogenic clones could be potentially targeted in the lymph node before migrating, accumulating, and possibly performing their effector functions in the joint. Moreover, analysis of clonal compositions can also serve as a biomarker to monitor disease progression and assess the therapeutic impact on changes in CD4 T cell clonality. Indeed, changes in cellular compositions in inguinal lymph nodes has been detected in at-risk and early RA patients[39] and treatment with RA therapeutics have been shown to reduce clonal expansion of CD4 T cells[33].

Together, these data outline the evolution of clonal dynamics between inflamed pLNs and their joints with the progression of inflammatory arthritis in that CD4 TCR clonality, and thus antigen specific responses, begin by being similar between the two sites then change in the pLN with the progression of the disease before potentially being mirrored in the joint in time. These observations have several implications on improving current antigen specific tolerogenic therapies. Firstly, understanding the clonal landscape gives an indication on the range of CD4 T cells that need to be targeted to improve the chances of modulating disease progression. Secondly, targeting expanded CD4 T cell populations early may be more effective therapeutically as the CD4 T cell repertoire is still restricted during the early phases, thus fewer populations would need to be targeted. Lastly, clonality can be a biomarker of disease progression and current therapeutics can be used to 'reset' the clonal landscape to one reflecting the earlier phases of the disease, which may enhance effectiveness of tolerogenic therapies and help reinstate self-tolerance and provide long-lasting remission in RA patients.

ACKNOWLEDGMENTS

Funding: This work was supported by Versus Arthritis. **RAB**, **JMB** and **PG** were supported by the Arthritis Research UK (ARUK) programme grant number 19788 and the Innovative Medicines Initiative EU-funded project Be The Cure (BTCURE) [grant number 115142-2]. **JMB** and **PG** were also supported by the Arthritis Research UK Rheumatoid Arthritis Pathogenesis Centre for Excellence (RACE) (grant number 20298) and the Research into Inflammatory Arthritis Centre Versus Arthritis (RACE) (grant number 22072). This work has received support from the EU/EFPIA Innovative Medicines Initiative 2 Joint Undertaking RTCure grant no. 777357. **SAK** was supported by the Ministry of Higher Education Oman (MOHE) throughout the study.

We thank the staff within the Institute of Infection, Immunity and Inflammation Flow Cytometry Facility and the Central Research Facility at the University of Glasgow for technical assistance. We also thank Dr. Megan K. MacLeod for her insight and for the critical discussions crucial for the development of this project.

AUTHOR CONTRIBUTIONS

S.A.K and R.A.B designed the research and performed the experiments. S.A.K analysed data, constructed figures, and wrote the manuscript. C.T.P and J.I.G provided technical assistance and helped with analysis. T.D.O provided technical assistance with the sequencing data, and J.M.B and P.G designed the research and contributed to writing the paper.

METHODS

Animals

Male and female C57BL/6 mice aged 7-12 weeks were purchased from Envigo (UK), and used as adoptive transfer recipients. OT-II TCR transgenic mice[40] were bred inhouse. All mice were maintained at the University of Glasgow's central research facility and housed under standard housing conditions specified by the UK Home Office.

Induction of inflammatory arthritis

"Early" antigen induced inflammatory arthritis was conducted as previously described[19,20]. Briefly, 2-3x10^6 Th1 polarised OT-II transgenic T cells were transferred i.v. into C57BL/6 recipients. Recipients were immunised with 100µL of 1µg/mL of OVA protein (Worthington Biochemicals) in Freund's complete adjuvant (CFA) (Sigma-Aldrich) subcutaneously after 24 hours. Ten days later, mice were given a periarticular injection in footpads with 50µL of 100µg of heat aggregated OVA (HAO) or PBS as a control. Mice were sacrificed four days later and cells were isolated from joint tissue and draining popliteal lymph nodes (pLN).

The "late" model antigen induced inflammatory arthritis was modified from a model of chronic polyarthritis[21]. 30 days after HAO administration described above, mice were given a second periarticular injection of 100µg of HAO in incomplete Freund's adjuvant (IFA) (Sigma-Aldrich) or IFA alone as a control. Mice were sacrificed 17 days later and cells were isolated from joint tissues and pLNs.

Isolation of cells from tissues

Cells from mouse joints and pLNs were isolated as previously described[18]. Briefly, ankle joints were teased apart and shaken at 110 rpm at 37°C for 25 minutes with 2.68mg/mL collagenase D (Roche) in RPMI 1640 (Gibco, Thermo Fisher Scientific). Joint tissue was then homogenised using a gentleMACS Dissociator (Miltenyi Biotech) then strained to obtain single cell suspensions. PLNs were forced through Nitex mesh (Cadisch Precision Meshes) to obtain single cell suspensions. Samples were washed and stained for flow cytometry. Joint and pLN samples were analysed separately. Joint samples and pLN samples were pooled per mouse.

Flow cytometry

Cells were stained for flow cytometry as described previously[18]. The following antibodies were used: anti-CD4 (clone GK1.5), anti-CD44 (clone IM7), anti-CD45 (clone

30-F11), anti-CD45.1 (clone A20). Data was analysed using FlowJo v10 software (Treestar, Oregon USA).

TCRβ Sequencing

The data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under the accession number PRJEB40509 (https://www.ebi.ac.uk/ena/browser/view/PRJEB40509). CDR3β sequencing was performed by iRepertoire Inc. (Huntsville, AL, USA) on total RNA collected from sorted antigen experienced endogenous CD4 T cells (CD4+, C45.1-, CD44hi) (details in supplemental methods). Barcoded libraries were prepared per sample using barcoded primers covering the V-J TCR region, then amplified, pooled, and sequenced using the Illumina MiSeq platform covering 150 paired end reads (PER). Basic data analysis was performed by iRepertoire. Sequencing data was also prepared for analysis using MiXCR developed by Bolotin et. al [41] and analysed using tcR developed by Nazarov et. al [42] in the R statistical software frame work (R version 3.4.3).

Calculation of the D50 diversity index (DI)

Diversity was calculated using iRepertoire's data analysis guide. Details in supplemental methods.

Calculation of repertoire overlap

A normalised overlap index was used to determine CDR3β amino acid sequence overlap between repertoires. Details in supplemental methods.

Statistics

Data is shown as mean ± SD. Specific test and significance levels are stated in respective figure legends. Statistical analyses were performed using GraphPad Prism version 7 (GraphPad Inc, CA, USA) or using the R statistical software framework (R version 3.4.3).

REFERENCES

Firestein GS, McInnes IB. Immunopathogenesis of Rheumatoid Arthritis. *Immunity* 2017;**46**:183–96. doi:10.1016/j.immuni.2017.02.006

Diaz-Gallo L-M, Martin J. PTPN22 splice forms: a new role in rheumatoid arthritis. *Genome Med* 2012;**4**:13. doi:10.1186/gm312

Holoshitz J. The rheumatoid arthritis HLA-DRB1 shared epitope. *Curr Opin Rheumatol* 2010;**22**:293–8. doi:10.1097/BOR.0b013e328336ba63

Lee H-S, Lee AT, Criswell LA, *et al.* Several regions in the major histocompatibility complex confer risk for anti-CCP-antibody positive rheumatoid arthritis, independent of the DRB1 locus. *Mol Med Camb Mass* 2008;**14**:293–300. doi:10.2119/2007-00123.Lee

Okada Y, Wu D, Trynka G, *et al.* Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature* 2013;**506**:376.

Plenge RM, Padyukov L, Remmers EF, *et al.* Replication of Putative Candidate-Gene Associations with Rheumatoid Arthritis in >4,000 Samples from North America and Sweden: Association of Susceptibility with PTPN22, CTLA4, and PADI4. *Am J Hum Genet* 2005;**77**:1044–60. doi:10.1086/498651

Remmers EF, Plenge RM, Lee AT, *et al.* STAT4 and the Risk of Rheumatoid Arthritis and Systemic Lupus Erythematosus. *N Engl J Med* 2007;**357**:977–86. doi:10.1056/NEJMoa073003

Blair HA, Deeks ED. Abatacept: A Review in Rheumatoid Arthritis. *Drugs* 2017;**77**:1221–33. doi:10.1007/s40265-017-0775-4

van Beers JJBC, Schwarte CM, Stammen-Vogelzangs J, *et al.* The rheumatoid arthritis synovial fluid citrullinome reveals novel citrullinated epitopes in apolipoprotein E, myeloid nuclear differentiation antigen, and β-actin. *Arthritis Rheum* 2013;**65**:69–80. doi:10.1002/art.37720

Burkhardt H, Sehnert B, Bockermann R, *et al.* Humoral immune response to citrullinated collagen type II determinants in early rheumatoid arthritis. *Eur J Immunol* 2005;**35**:1643–52. doi:10.1002/eji.200526000

Ioan-Facsinay A, el-Bannoudi H, Scherer HU, *et al.* Anti-cyclic citrullinated peptide antibodies are a collection of anti-citrullinated protein antibodies and contain overlapping and non-overlapping reactivities. *Ann Rheum Dis* 2011;**70**:188–93. doi:10.1136/ard.2010.131102

James EA, Rieck M, Pieper J, *et al.* Citrulline-specific Th1 cells are increased in rheumatoid arthritis and their frequency is influenced by disease duration and therapy. *Arthritis Rheumatol Hoboken NJ* 2014;**66**:1712–22. doi:10.1002/art.38637

Law S, Street S, Yu C-H, *et al.* T-cell autoreactivity to citrullinated autoantigenic peptides in rheumatoid arthritis patients carrying HLA-DRB1 shared epitope alleles. *Arthritis Res Ther* 2012;**14**:R118. doi:10.1186/ar3848

Ikeda Y, Masuko K, Nakai Y, *et al.* High frequencies of identical T cell clonotypes in synovial tissues of rheumatoid arthritis patients suggest the occurrence of common antigen-driven immune responses. *Arthritis Rheum* 1996;**39**:446–53. doi:10.1002/art.1780390312

Klarenbeek PL, de Hair MJH, Doorenspleet ME, *et al.* Inflamed target tissue provides a specific niche for highly expanded T-cell clones in early human autoimmune disease. *Ann Rheum Dis* 2012;**71**:1088–93. doi:10.1136/annrheumdis-2011-200612

Stamenkovic I, Stegagno M, Wright KA, *et al.* Clonal dominance among T-lymphocyte infiltrates in arthritis. *Proc Natl Acad Sci U S A* 1988;**85**:1179–83.

VanderBorght A, Geusens P, Vandevyver C, *et al.* Skewed T-cell receptor variable gene usage in the synovium of early and chronic rheumatoid arthritis patients and persistence of clonally expanded T cells in a chronic patient. *Rheumatol Oxf Engl* 2000;**39**:1189–201.

Prendergast CT, Patakas A, Al-Khabouri S, *et al.* Visualising the interaction of CD4 T cells and DCs in the evolution of inflammatory arthritis. *Ann Rheum Dis* 2018;**77**:579–88. doi:10.1136/annrheumdis-2017-212279

Benson RA, Patakas A, Conigliaro P, *et al.* Identifying the Cells Breaching Self-Tolerance in Autoimmunity. *J Immunol* 2010;**184**:6378–85. doi:10.4049/jimmunol.0903951

Maffia P, Brewer JM, Gracie JA, *et al.* Inducing Experimental Arthritis and Breaking Self-Tolerance to Joint-Specific Antigens with Trackable, Ovalbumin-Specific T Cells. *J Immunol* 2004;**173**:151–6. doi:10.4049/jimmunol.173.1.151

Conigliaro P, Benson RA, Patakas A, *et al.* Characterization of the anticollagen antibody response in a new model of chronic polyarthritis. *Arthritis Rheum* 2011;**63**:2299–308. doi:10.1002/art.30413

Katayama CD, Eidelman FJ, Duncan A, *et al.* Predicted complementarity determining regions of the T cell antigen receptor determine antigen specificity. *EMBO J* 1995;**14**:927–38.

Turner SJ, Doherty PC, McCluskey J, *et al.* Structural determinants of T-cell receptor bias in immunity. *Nat Rev Immunol* 2006;**6**:883–94. doi:10.1038/nri1977

Fazou C, Yang H, McMichael AJ, *et al.* Epitope specificity of clonally expanded populations of CD8+ T cells found within the joints of patients with inflammatory arthritis. *Arthritis Rheum* 2001;**44**:2038–45. doi:10.1002/1529-0131(200109)44:9<2038::AID-ART353>3.0.CO;2-1

Iannone F, Corrigall VM, Kingsley GH, *et al.* Evidence for the continuous recruitment and activation of T cells into the joints of patients with rheumatoid arthritis. *Eur J Immunol* 1994;**24**:2706–13. doi:10.1002/eji.1830241120

Shadidi KR, Aarvak T, Jeansson S, *et al.* T-cell responses to viral, bacterial and protozoan antigens in rheumatoid inflammation. Selective migration of T cells to synovial tissue. *Rheumatol Oxf Engl* 2001;**40**:1120–5. doi:10.1093/rheumatology/40.10.1120

Brennan FM, Hayes AL, Ciesielski CJ, *et al.* Evidence that rheumatoid arthritis synovial T cells are similar to cytokine-activated T cells: involvement of phosphatidylinositol 3-kinase and nuclear factor kappaB pathways in tumor necrosis factor alpha production in rheumatoid arthritis. *Arthritis Rheum* 2002;**46**:31–41. doi:10.1002/1529-0131(200201)46:1<31::AID-ART10029>3.0.CO;2-5

Brennan FM, Smith NM, Owen S, *et al.* Resting CD4+ effector memory T cells are precursors of bystander-activated effectors: a surrogate model of rheumatoid arthritis synovial T-cell function. *Arthritis Res Ther* 2008;**10**:R36. doi:10.1186/ar2390

Howson LJ, Napolitani G, Shepherd D, *et al.* MAIT cell clonal expansion and TCR repertoire shaping in human volunteers challenged with Salmonella Paratyphi A. *Nat Commun* 2018;**9**:253. doi:10.1038/s41467-017-02540-x

Gomez-Tourino I, Kamra Y, Baptista R, *et al.* T cell receptor β-chains display abnormal shortening and repertoire sharing in type 1 diabetes. *Nat Commun* 2017;**8**:1792. doi:10.1038/s41467-017-01925-2

Risnes LF, Christophersen A, Dahal-Koirala S, *et al.* Disease-driving CD4+ T cell clonotypes persist for decades in celiac disease. *J Clin Invest* 2018;**128**:2642–50. doi:10.1172/JCI98819

Cantaert T, Brouard S, Thurlings RM, *et al.* Alterations of the synovial T cell repertoire in anti-citrullinated protein antibody-positive rheumatoid arthritis. *Arthritis Rheum* 2009;**60**:1944–56. doi:10.1002/art.24635

Pierer M, Rossol M, Kaltenhäuser S, *et al.* Clonal expansions in selected TCR BV families of rheumatoid arthritis patients are reduced by treatment with the TNFα inhibitors etanercept and infliximab. *Rheumatol Int* 2011;**31**:1023–9. doi:10.1007/s00296-010-1402-9

Wagner U, Pierer M, Kaltenhäuser S, *et al.* Clonally expanded CD4+CD28null T cells in rheumatoid arthritis use distinct combinations of T cell receptor BV and BJ elements. *Eur J Immunol* 2003;**33**:79–84. doi:10.1002/immu.200390010

Wagner UG, Koetz K, Weyand CM, *et al.* Perturbation of the T cell repertoire in rheumatoid arthritis. *Proc Natl Acad Sci U S A* 1998;**95**:14447–52. doi:10.1073/pnas.95.24.14447

Nickdel MB, Conigliaro P, Valesini G, *et al.* Dissecting the contribution of innate and antigen-specific pathways to the breach of self-tolerance observed in a murine model of arthritis. *Ann Rheum Dis* 2009;**68**:1059–66. doi:10.1136/ard.2008.089300

Platt AM, Gibson VB, Patakas A, *et al.* Abatacept limits breach of self-tolerance in a murine model of arthritis via effects on the generation of T follicular helper cells. *J Immunol Baltim Md 1950* 2010;**185**:1558–67. doi:10.4049/jimmunol.1001311

Monach PA, Hueber W, Kessler B, *et al.* A broad screen for targets of immune complexes decorating arthritic joints highlights deposition of nucleosomes in rheumatoid arthritis. *Proc Natl Acad Sci* 2009;**106**:15867. doi:10.1073/pnas.0908032106

Rodríguez-Carrio J, Hähnlein JS, Ramwadhdoebe TH, *et al.* Brief Report: Altered Innate Lymphoid Cell Subsets in Human Lymph Node Biopsy Specimens Obtained During the At-Risk and Earliest Phases of Rheumatoid Arthritis. *Arthritis Rheumatol Hoboken NJ* 2017;**69**:70–6. doi:10.1002/art.39811

Barnden MJ, Allison J, Heath WR, *et al.* Defective TCR expression in transgenic mice constructed using cDNA-based alpha- and beta-chain genes under the control of heterologous regulatory elements. *Immunol Cell Biol* 1998;**76**:34–40. doi:10.1046/j.1440-1711.1998.00709.x

Bolotin DA, Poslavsky S, Mitrophanov I, *et al.* MiXCR: software for comprehensive adaptive immunity profiling. *Nat Methods* 2015;**12**:380.

Nazarov VI, Pogorelyy MV, Komech EA, *et al.* tcR: an R package for T cell receptor repertoire advanced data analysis. *BMC Bioinformatics* 2015;**16**. doi:10.1186/s12859-015-0613-1

FIGURE LEGENDS

Figure 1.

Outline of model phases and number of endogenous, OT-II, and endogenous antigen experienced CD4 T cells at both phases. **A)** Illustration of early and late phases of inflammatory arthritis. **B)** Number of endogenous CD4 T cells (CD4+, CD45+, CD45.1-) and OT-II T cells (CD4+, CD45+, CD45.1+) from pLNs and joints of mice undergoing inflammatory arthritis at the early and late phases and respective controls. **C)** Number of antigen experienced endogenous CD4 T cells (CD4+, CD45+, CD45.1-, CD44hi) from pLNs and joints of mice undergoing inflammatory arthritis at the early and late phases and respective controls. Data is representative of mean ± SD with each point representing individual experimental mice. Data represents three independent experiments for the early phase and two independent experiments for the late phase. Groups were compared using 2-way ANOVA and Student's t-tests. Stars represent the following p-values: *<0.05, ** <0.01; ***<0.001; **** <0.0001; ns: not significant. HAO, heat-aggregated OVA; IFA, incomplete Freund's adjuvant.

Figure 2

Characterisation of antigen experienced endogenous CD4 T cell repertoires from inflamed pLNs and joints and PBS controls at the early phase. Proportion of the top 10 most frequently occurring clones, followed by the next 25, 500, 3000, 10,000, and 100,000 most frequent clones contributing to the endogenous antigen experienced CD4+ T cell repertoire in pooled inflamed pLNs and joint samples at the early phase (HAO) and pooled PBS controls. Data represents 1 experiment with pooled samples from 3 mice for the PBS sample and 5 mice for the inflamed (HAO) sample.

Figure 3

Characterisation of antigen experienced endogenous CD4 T cell repertoires from inflamed pLNs and joints and IFA controls at the late phase. Proportion of the top 10 most frequently occurring clones, followed by the next 25, 500, 3000, 10,000, and 100,000 most frequent clones contributing to the endogenous antigen experienced CD4+ T cell repertoire in **A)** inflamed pLNs, and **E)** inflamed joints at the late phase (HAO+IFA) and IFA only controls. **B)** Proportion of the least frequently occurring clones in inflamed pLNs at the late phase and IFA controls. **F)** Percentage contribution of the top 10 most frequently occurring clones in inflamed joints at the late phase and IFA controls. **C and G** clonality, represented as the number of unique CDR3β DNA sequences, in **C)** inflamed pLNs and IFA controls, and **G)** inflamed joints and IFA controls at the late phase. **D and H**

diversity indices of **D)** inflamed pLNs and IFA controls, and **H)** inflamed joints and IFA controls at the late phase. Data is representative of mean ± SD with each point representing individual experimental mice. Data represents 1 experiment. Groups were compared using unpaired Student's t-test. Stars represent the following p values: * <0.05; ns: not significant.

Figure 4

Characterisation of antigen experienced endogenous CD4 T cell repertoires in inflamed pLNs and joints at the early and late phases. Proportion of the top 10 most frequently occurring clones, followed by the next 25, 500, 3000, 10,000, and 100,000 most frequent clones contributing to the endogenous antigen experienced CD4+ T cell repertoire in **A)** inflamed pLNs at the early and late phases, and **E)** inflamed joints at the early and late phases. **B and F** percentage contribution of the top 10 most frequently occurring clones in **B)** inflamed pLNs and **F)** inflamed joints at the early and late phases. **C and G** clonality, represented as the number of unique CDR3β DNA sequences, in **C)** inflamed pLNs, and **G)** inflamed joints at the early and late phases. **D and H** diversity indices of **D)** inflamed pLNs, and **H)** inflamed joints at the early and late phases. Data is representative of mean ± SD with each point representing individual experimental mice. Data represents 1 experiment. Groups were compared using unpaired Student's t-test. Stars represent the following p values: *** <0.001; **** <0.0001; ns: not significant.

Figure 5

Range of TRBV genes expressed in antigen experienced endogenous CD4 T cell repertoires of inflamed pLNs and joints at the early and late phases. **A)** PCA plot of TRBV gene frequency in repertoires of inflamed pLNs and joints at the early and late phases. Frequency of **B)** TRBV gene expression, and **C)** TRBJ gene expression in inflamed pLNs at the early and late phases. Graphs represent 1 experiment with n=5 for both the early and late inflamed groups. Groups were compared using ordinary 2-way ANOVA with Sidak's multiple comparisons test. Data presented as mean ±SD. Stars represent the following p values: * <0.05; ** <0.01; *** <0.001; **** <0.0001; ns: not significant.

Figure 6

Degree of CDR3β amino acid sequence overlap and correlation of most abundant clones between inflamed pLNs and joints at the early and late phases. **A)** Heatmaps of normalised number of CDR3β amino acid sequences (see methods) between inflamed pLNs and joints at the early (left) and late (right) phases. **B)** Graph representing the intersects (diagonal) of A) representing CDR3β sequence overlap between inflamed pLNs

and joint within the same animal. **C-E** Correlation of the top 10 ranked clones in joints with their rank in respective pLN samples The rank of each of the top 10 clones in **C)** inflamed joint at the early phase, **D)** inflamed joint at the late phase, and **E)** IFA control joint samples were plotted against the rank of that clone in respective pLN samples. Data is representative of one experiment. Graph in B) is representative of mean ± SD with each point representing individual experimental mice. * represents p<0.05. In C-E correlations (r) were calculated using Spearman's correlation after testing for normality of the data using the D'Agostino-Pearson test. p represents the p value.

Supplementary figure 1

Gating strategy to identify antigen experienced endogenous CD4 T cells from pLNs and joints. Representative flow cytometry plots identifying antigen experienced endogenous CD4 T cells (CD4+, CD45+, C45.1-, CD44hi) from **A)** inflamed pLNs, and **B)** inflamed joints at the early phase of mice undergoing inflammatory arthritis.

Supplementary figure 2

Representative flow cytometry plots identifying antigen experienced endogenous CD4 T from **A)** pLN, and **B)** joint PBS controls at the early phase, **C)** pLN, and **D)** joints IFA controls at the late phase, and **E)** inflamed pLN, and **F)** inflamed joints at the late phase.

Supplementary figure 3

Degree of CDR3β amino acid sequence overlap between inflamed pLN and inflamed joint samples at the early and late phases. Heatmaps of normalised number of CDR3β amino acid sequences between **A)** inflamed pLNs, and **C)** inflamed joints at the early and late phases. **B and D** Normalised overlap values between **B)** inflamed pLN samples, and **D)** inflamed joint samples at the early and late phases. Data is representative of mean ± SD with each point representing individual experimental mice. Data represents 1 experiment. Groups were compared using unpaired Student's t-test. Stars represent the following p values: ** <0.01; ns: not significant.

Supplementary table 1

The top 10 frequently occurring clones inflamed joint samples at the early phase ranked in terms of frequency in respective pLNs. The top 10 frequently occurring clones in the antigen experienced CD4+ T cell repertoire were found in each of the inflamed joint samples at the early phase and searched for in respective pLN samples. The CDR3β amino acid sequence, associated TRBV gene, read count, and rank of the clone in the

pLN sample was obtained. The rank of each of the top 10 clones in the joint are also included in the table. Data is representative of 1 experiment with n=5.

Supplementary table 2

The top 10 frequently occurring clones inflamed joint samples at the late phase ranked in terms of frequency in respective pLNs. The top 10 frequently occurring clones in the antigen experienced CD4+ T cell repertoire were found in each of the inflamed joint samples at the late phase and searched for in respective pLN samples. The CDR3β amino acid sequence, associated TRBV gene, read count, and rank of the clone in the pLN sample was obtained. The rank of each of the top 10 clones in the joint are also included in the table. Data is representative of 1 experiment with n=5.

Fig 1.

A)

B)

C)

Fig 2.

Fig 3.

Fig 4.

A)

B)

C)

Fig 6

A)

Early inflamed pLNs and joints

Late inflamed pLNs and joints

B)

C) Early inflamed
r = 0.1321
p = 0.5479

D) Late inflamed
r = 0.5298
p < 0.0001

E) IFA control
r = 0.0977
p = 0.5709