TCRβ Sequencing Reveals Spatial and Temporal Evolution of

Clonal CD4 T cell Responses in a Breach of Tolerance Model of

Inflammatory Arthritis

Shaima Al-Khabouri[#], Robert A. Benson*, Catriona T. Prendergast[§], Joshua I.

Gray[$], Thomas D Otto, James M Brewer, Paul Garside[1]

Corresponding author information:

Paul Garside

paul.garside@glasgow.ac.uk

Affiliations:

[1]Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary

and Life Sciences, Sir Graeme Davies Building, University of Glasgow, 120

University Place, Glasgow G12 8TA, United Kingdom

[#]Current address: Department of Medicine Solna (MedS), Division of

Rheumatology, Center for Molecular Medicine (CMM), Karolinska Institute,171 76

Stockholm, Sweden

*Current address: Antibody Analytics Ltd, BioCity Scotland, Bo'ness Road,

Newhouse, ML1 5UH, United Kingdom

§Current address: Novartis Institutes for Biomedical Research (NIBR),

Autoimmunity, Transplantation and Inflammation (ATI), Novartis Campus, 4056,

Basel, Switzerland

[$]Current address: Columbia Center for Translational Immunology, Columbia

University, New York NY, USA 10032

Keywords: TCRβ sequencing, T cell clonality, CD4 T cell, rheumatoid arthritis,

tolerogenic therapy

Word count: 3910

ABSTRACT

Effective tolerogenic intervention in Rheumatoid Arthritis (RA) will rely upon

understanding the evolution of articular antigen specific CD4 T cell responses.

TCR clonality of endogenous CD4 T cell infiltrates in early inflammatory arthritis

was assessed to monitor evolution of the TCR repertoire in the inflamed joint and

associated lymph node (LN). Mouse models of antigen-induced breach of self-

tolerance and chronic polyarthritis were used to recapitulate early and late phases

of RA. The infiltrating endogenous, antigen experienced CD4 T cells in inflamed

joints and LNs were analysed using flow cytometry and TCRβ sequencing. TCR

repertoires from inflamed late phase LNs displayed increased clonality and

diversity compared to early phase LNs, while inflamed joints remained similar with

time. Repertoires from late phase LNs accumulated clones with a diverse range of

TRBV genes, while inflamed joints at both phases contained clones expressing

similar TRBV genes. Repertoires from LNs and joints at the late phase displayed

reduced CDR3β sequence overlap compared to the early disease phase, however

the most abundant clones in LNs accumulate in the joint at the later phase. The

results indicate CD4 T cell repertoire clonality and diversity broadens with

progression of inflammatory arthritis and is first reflected in LNs before mirroring in

the joint. These observations imply that antigen specific tolerogenic therapies

could be more effective if targeted at earlier phases of disease when CD4 T cell

clonality is least diverse.

INTRODUCTION

Rheumatoid arthritis (RA) is a chronic inflammatory autoimmune disease

characterised by synovial inflammation and cartilage and bone erosion, causing

progressive loss of joint function (1). Antigen presentation and CD4 T activation

mechanisms have been shown to play a role in the pathogenesis of RA. This is

evidenced by genetic association studies in RA patients showing strong

associations of HLA-DRB alleles, T cell activation associated genes such as

PTPN22 and *CTLA-4,* and loci associated with signal transduction such as *STAT4*

(2–7). Moreover, the successes of therapeutics in modulating T cell activation,

such as the use of Abatacept, reinforce the importance of CD4 T cells in

propagating the disease (8). As such, there has been increasing interest in

targeting pathogenic autoreactive CD4 T cells using antigen specific tolerogenic

therapies as this line of therapy aims to re-establish tolerance and provide long-

lasting remission, while retaining protective immunity against pathogens. However,

targeting autoreactive CD4 T cells has proven difficult in RA, due to the broad

range of antigens implicated in disease, lack of a clear cellular hierarchy of

disease drivers, and also patient to patient variation (9–13). Moreover, it is

unknown at what stages of the disease these autoreactive responses develop nor

the location in which these responses are primed, thus hampering the

development of effective tolerogenic therapies. Antigen specific responses can be

identified by detecting expanded clonal T cell populations. Indeed, oligoclonal CD4

T cell responses and skewed TCR repertoires have been reported in arthritic joints

(14–17). More recently, analysis of CD4 T cell behaviour was shown to reflect

antigen recognition in inflamed joints at the early stages of disease (18),

suggesting that antigen specific CD4 T cell responses may perpetuate and drive

progression of RA. However, the development of antigen specific responses and

their evolution with the progression of RA is poorly understood. Investigating the

evolution TCR clonality of antigen experienced CD4 T cells in RA will provide

insight on how antigen specific responses evolve with disease progression. This

will inform the development of more effective tolerogenic therapies by indicating

the range of clones that would need to be targeted and the disease stage at which

these therapies are most likely to be effective. In this study, we employed breach

of self-tolerance models of antigen induced inflammatory arthritis (19,20) and

chronic polyarthritis (21) in which autoreactive responses develop. These models

also recapitulate the early and later stages of the disease – hereafter named the

early and late phases respectively. TCRβ sequencing was used to monitor the

evolution of antigen specific endogenous CD4 T cell responses in inflamed joints

and their associated draining lymph nodes.

METHODS

Animals

Male and female C57BL/6 mice aged 7-12 weeks were purchased from Envigo

(UK), and used as adoptive transfer recipients. OT-II TCR transgenic mice (22)

were bred inhouse. All mice were maintained at the University of Glasgow's

central research facility and housed under standard housing conditions specified

by the UK Home Office.

Induction of inflammatory arthritis

"Early" antigen induced inflammatory arthritis was conducted as previously

described (19,20)**.** Briefly, 2-3x10^6 Th1 polarised OT-II transgenic T cells were

transferred i.v. into C57BL/6 recipients. Recipients were immunised with 100µL of

1µg/mL of OVA protein (Worthington Biochemicals) in Freund's complete adjuvant

(CFA) (Sigma-Aldrich) subcutaneously after 24 hours. Ten days later, mice were

given a periarticular injection in footpads with 50µL of 100µg of heat aggregated

OVA (HAO) or PBS as a control. Mice were sacrificed four days later and cells

were isolated from joint tissue and draining popliteal lymph nodes (pLN).

The "late" model antigen induced inflammatory arthritis was modified from a model

of chronic polyarthritis(21). 30 days after HAO administration described above,

mice were given a second periarticular injection of 100µg of HAO in incomplete

Freund's adjuvant (IFA) (Sigma-Aldrich) or IFA alone as a control. Mice were

sacrificed 17 days later and cells were isolated from joint tissues and pLNs.

Isolation of cells from tissues

Cells from mouse joints and pLNs were isolated as previously described (18).

Briefly, ankle joints were teased apart and shaken at 110 rpm at 37°C for 25

minutes with 2.68mg/mL collagenase D (Roche) in RPMI 1640 (Gibco, Thermo

Fisher Scientific). Joint tissue was then homogenised using a gentleMACS

Dissociator (Miltenyi Biotech) then strained to obtain single cell suspensions.

PLNs were forced through Nitex mesh (Cadisch Precision Meshes) to obtain

single cell suspensions. Samples were washed and stained for flow cytometry.

Joint and pLN samples were analysed separately. Joint samples and pLN samples

were pooled per mouse.

Flow cytometry

Cells were stained for flow cytometry as described previously (18). The following antibodies were used: anti-CD4 (clone GK1.5), anti-CD44 (clone IM7), anti-CD45 (clone 30-F11), anti-CD45.1 (clone A20). Data was analysed using FlowJo v10 software (Treestar, Oregon USA).

TCRβ Sequencing

The data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under the accession number PRJEB40509 (https://www.ebi.ac.uk/ena/browser/view/PRJEB40509). CDR3β sequencing was performed by iRepertoire Inc. (Huntsville, AL, USA) on total RNA collected from sorted antigen experienced endogenous CD4 T cells (CD4+, C45.1-, CD44hi). CD4+, CD45.1-, CD44hi were sorted from joint and pLN samples using the BD Aria III FACS sorter (BD Biosciences). Samples were collected separately from individual mice in lysis buffer (Purelink RNA micro kit, Thermofisher) + 1% β-mercaptoethanol (Sigma-Aldrich) and lysed using 29G insulin syringes (VWR). Lysed cells were then stored at -80°C prior to RNA purification. Total RNA was purified using the Purelink RNA micro kit (Thermo Fisher Scientific) as per manufacturer's instructions and quality assessed by measuring A260/280 using a Nanodrop 1000 spectrophotometer (Thermo Fisher Scientific). Barcoded libraries were prepared per sample using barcoded primers covering the V-J TCR region, then amplified, pooled, and sequenced using the Illumina MiSeq platform covering 150 paired end reads (PER). Basic data analysis was performed by iRepertoire. Sequencing data was also prepared for analysis using MiXCR developed by Bolotin et. al (23) and analysed using tcR developed by Nazarov et. al (24) in the R statistical software frame work (R version 3.4.3).

Calculation of the D50 diversity index (DI)

Diversity was calculated using iRepertoire's data analysis guide. For samples where the number of unique CDR3β sequences ≥10,000:

DI = rank of unique CDR3β sequence where 50% of the top 10,000 total reads falls x 100 / 10,000

For samples where the number of unique CDR3β sequences <10,000:

DI = rank of unique CDR3β sequence where 50% of the total reads falls x 100 / no. of unique CDR3β sequences

Calculation of repertoire overlap

The number of shared CDR3β amino acid sequences is represented as a

normalised overlap index taking into consideration the size of the repertoires being

compared. The normalised overlap index was calculated using the tcR R package

(24) using the following calculation:

Normalised overlap index = No. of exact overlapping CDR3β amino acid

sequences/ total reads repertoire 1 x total number of reads repertoire 2

Statistics

Data is shown as mean ± SD. Specific test and significance levels are stated in

respective figure legends. Statistical analyses were performed using GraphPad

Prism version 7 (GraphPad Inc, CA, USA) or using the R statistical software

framework (R version 3.4.3).

RESULTS

Induction of inflammatory arthritis results in local accumulation of

endogenous antigen experienced CD4 T cells at both the early and late

phases

Induction of the breach of tolerance model of inflammatory arthritis requires the

transfer of Th1 polarised ovalbumin (OVA) specific OT-II cells and subsequent

immunisations with OVA and heat aggregated OVA (HAO) as outlined in **figure 1**

A. This results in the influx of the transferred OT-II cells and endogenous CD4 T

cells of varying antigen specificities, indicated by the range of expressed TCRβ

variable genes (TRBV) (18)**.** Assessing the evolution of TCR clonality of the

endogenous CD4 T cell population requires the accurate identification of these

cells in joints and popliteal lymph nodes (pLN), being distinguishable from the

transferred OT-IIs by expression of the congenic marker CD45.1 **(supplementary**

figures 1 and 2). Endogenous CD4 T cells were present at a significantly higher

number than the transferred OT-II cells in inflamed pLNs and joints at both the

early and late phases **(figure 1 B)**. A significant number of endogenous CD4 T

cells at the early phase also displayed an antigen experienced phenotype,

indicated by increased CD44 expression, after induction of inflammatory arthritis

with HAO **(figure 1 C)**, and were previously shown to produce TNFα and IFNγ

following ex-vivo stimulation with PMA/ionomycin (18). The number of endogenous

CD4, CD44hi cells was also increased at the late phase in inflamed joints **(figure 1**

C).

Accumulation of expanded clones and observed clonality and diversity of

CD4 T cell repertoires is antigen driven

To investigate the evolution of CD4 TCR clonality in this model of inflammatory

arthritis, endogenous antigen experienced CD4 T cells i.e. excluding the

transferred OT-II cells, were sorted from pLNs and joints of individual mice at both

early and late phases. Total RNA was then isolated from the endogenous CD4 T

cells and used to sequence the CDR3β region of the TCR, the most variable

region of the TCR critical in recognition of specific peptide sequences (25,26).

Repertoires were then characterised by examining the number of unique CDR3β

sequences and evaluating the degree expanded clones contributed to the overall

antigen experienced repertoire. From this, the clonality and diversity of the CD4 T

cell repertoire can be determined. In this study, clonality is defined by the number

of unique CDR3β sequences found in the population, and diversity – represented

by the Diversity Index (DI) (see Methods) represents the relative contribution of

each unique CDR3β sequence, or clone, to the population. Thus, a repertoire with

a larger number unique CDR3β sequences, and therefore clones, would be

described to have high clonality. This repertoire would also display high diversity

and have a high DI value if the number of cells contributing to different clonal

populations are more evenly distributed. Conversely, the repertoire would be

described to have low diversity and have a low DI value if the repertoire is mainly

comprised of a few clones present in high frequencies relative to the total number

of clones. Due to the low recovery of antigen experienced endogenous CD4 T

cells isolated from control pLNs and joints at the early phase, cells from control

pLNs or joints needed to be pooled prior to sequencing. Pooling samples changes

the composition of the repertoire and affects T cell repertoire diversity, so

comparisons of control CD4 T cell populations were made against pooled data

from inflamed joint and pLN samples at the early phase **(figures 2 A and B)**.

Clonality of the endogenous CD4 T cell repertoire from inflamed pLNs at the early

phase and controls were comparable (4181 unique CDR3β sequences in the

inflamed pLN repertoire vs 4347 in the PBS pLN repertoire). However, the

repertoire from the inflamed pLN showed a reduced diversity compared with the

control pLN sample (6.4 vs 16.3 respectively). Endogenous CD4 T cell repertoires

from inflamed joints displayed higher clonality than controls (4549 vs 885). Despite

this, the diversity of the repertoire in the inflamed joint was relatively low compared

to the PBS joint (0.72 vs 9.3 respectively). These data therefore demonstrate that

accumulation of clones can be detected at the early phases of the disease in both

inflamed pLNs and joints.

Evidence for the accumulation of antigen experienced T cells in the absence of

their cognate antigen has been demonstrated in this model (18) and in RA patients

(27–29) which may be a result of a chronic inflammatory environment (30,31). To

address whether the accumulation of clones is antigen driven rather than a result

of non-antigen induced inflammation, the antigen experienced repertoires of pLNs

and joints at the late phase were compared to repertoires isolated from mice

challenged with IFA alone **(figure 3 A)**, where inflammation is induced without the

presence of antigen. No significant difference was found between the contribution

of the top 10 most expanded clones to the repertoires of inflamed pLNs at the late

phase and IFA controls (data not shown) nor when comparing the contribution of

clones occurring at the lowest frequency **(figure 3 B)**. The number of clones

occurring at a frequency of 2 or more were also compared between inflamed and

IFA pLN samples as this was the median frequency of clones found in the PBS

pLN sample (data not shown). Thus, clonal populations found at higher than this

frequency are assumed to have accumulated due to the presence of antigen. No

clonal accumulation was detected between inflamed pLNs at the late phase and

IFA controls (*p value 0.32 Welch two sample t-test*). Moreover, repertoires from

inflamed pLNs at the late phase and IFA displayed comparable levels of clonality

and diversity **(figures 3 C and D)**.

In late phase arthritic joints, the top 10 highly expanded clones constituted

approximately 41% of the overall repertoire compared to joint IFA controls, where

the top clones only comprised an average of 29% of the entire repertoire **(figures**

3 E and F). No difference in clonality was detected between repertoires from

inflamed joints and IFA controls **(figure 3 G)**, but inflamed joints have reduced

clonal diversity compared to IFA controls **(figure 3 H)**. This would suggest that

antigen driven accumulation of expanded clones in inflamed joints can also be

observed at the late phase.

The antigen experienced endogenous CD4 T cell repertoire displays

disparity in clonal diversity between inflamed pLNs and joints with time

Antigen experienced endogenous CD4 T cell repertoires from pLNs and joints

were compared between the early and late phases to investigate how CD4 TCR

clonality evolves with disease progression **(figures 4 A and E)**. When comparing

inflamed pLNs at the early and late phases, the top 10 most expanded clones

contributed approximately 18% to the overall antigen experienced repertoire of

inflamed pLNs at the early phase compared to only 2% at the late phase **(figures**

4 A and B). When comparing clonality and diversity, repertoires from pLNs

isolated at the late phase displayed increased clonality and diversity compared

with their early counterparts **(figures 2 C and D)** and comprised mainly of low

abundance clones. In contrast to the pLNs, antigen experienced CD4 T cell

repertoires isolated from inflamed joints at the early and late phases were similar;

the top 10 most expanded clones dominated the repertoires and comprised on

average 46% and 41% of the overall populations **(figures 4 E and F)**.

Furthermore, no differences were observed between the clonality and diversity of

the repertoires at the early and late phases **(figures 4 G and H)** and repertoire

diversity overall was low in the joint at both phases. Together, these data highlight

the disparity in repertoire composition between pLNs and joints at the early and

late phases, namely that pLNs at the late phase displayed high repertoire clonality

and diversity compared to their early counterparts and also compared to inflamed

joints at the same phase.

Disparity in clonality of the antigen experienced endogenous CD4 T cell

repertoires is attributed to accumulation of clones with different TCRβ

variable (TRBV) genes

Characterisation of the antigen experienced CD4 T cell repertoires from pLNs and

joints at the early and late phases provided useful information on the overall

distribution of clones in these repertoires. However, this does not provide any

insight on the TCR sequence of these clones, nor on whether these clones have

similar or different TCR sequences between the two sites. Moreover, no insight is

gained on whether the clonal composition of the antigen experienced repertoires in

these two sites change over time. To address this, a PCA analysis was performed

on inflamed pLNs and joints at both the early and late phases on the basis of

TRBV gene use **(figure 5 A)**. Repertoires isolated from inflamed pLNs at the late

phase displayed the largest range in TRBV gene use, while inflamed pLNs and

joints at the early phase as well as inflamed joints at the late phase grouped

together in terms of TRBV gene expression. Indeed, when quantifying these

differences, inflamed pLNs at the late phase had significantly increased presence

of TRBV genes 1, 2, 3, 5, and 19, while TRBV genes 12-1, 12-2, and 13-1 were

present in the highest frequencies in inflamed pLNs at the early phase **(figure 5**

B). These differences are also due to changes in TRBV gene use and not in TRBJ

genes **(figure 5 C)**. Interestingly, the PCA analysis also highlighted inter-sample

differences in TRBV gene use in inflamed pLNs at the late phase. Accumulation of

clones expressing different TRBV genes in inflamed pLNs at the late phase

suggests changes in antigen specific responses with time and also between

inflamed pLNs and their associated tissues. Moreover, these changes in antigen

specific responses also differ between individual animals with time.

Antigen associated changes in the inflamed pLN may predict the changes in

antigen specificities and clonality of the CD4 T cell repertoire in inflamed

joints

To assess whether changes in CD4 TCR clonality reflected changes antigen

specificities, the degree of CDR3β amino acid sequence overlap between the

antigen specific CD4 T cell repertoires was investigated. Comparing the repertoire

overlap between inflamed pLNs and joints at both the early and late phases

allowed us to monitor clonal dynamics between these two sites with the

progression of inflammatory arthritis. The number of overlapping CDR3β amino

acid sequences is represented by the normalised overlap index (see methods) and

this decreased between the inflamed pLN and its joint with the progression of

inflammatory arthritis **(figures 6 A and B)**. CDR3β sequence overlap decreased

significantly between samples when comparing repertoires from inflamed pLN

samples at the early phase to inflamed pLNs at the late phase **(supplementary**

figure 3 A), while the degree of CDR3β amino acid sequence overlap remained

unchanged between repertoires from inflamed joints at the early and late phases

(supplementary figure 3 B). These observations confirm the observed changes

in TRBV gene use and also imply changes in antigen specific responses in pLNs,

which progress towards polyclonality with disease progression. Correlation

analysis revealed that the top 10 most expanded clones in the inflamed joint at the

late phase correlated significantly with the most expanded clones in the inflamed

pLNs at the same phase **(figure 6 D)**. Furthermore, the 10 most expanded clones

in inflamed joints at the early phase and those from IFA controls did not correlate

with the most highly expanded clones in their respective pLNs **(figures 6 C and**

E). Moreover, the 10 most expanded clones in inflamed joints at the early phase

were not all present in their respective pLNs **(supplementary table 1)**, but the top

10 expanded clones in inflamed joints at the late phase were all present in

inflamed pLNs at the same phase **(supplementary table 2)**. These results

indicate that despite pLNs progressing towards polyclonality with the development

of disease, the most highly expanded antigen experienced clones accumulate in

the inflamed joint. Thus repertoire clonality in the pLN may eventually be mirrored

in the inflamed joint with disease progression.

DISCUSSION

Characterising CD4 TCR clonality has been used to monitor progression of

disease states in cases of infection (32) and autoimmunity (33,34), including RA

(15,35–38). Although CD4 T cell clonality has been determined in RA patients at

different stages of the disease (15,17), the development of antigen specific

responses from very early, pre-clinical stages to a more established disease stage

remains unknown. Understanding this is important to develop effective tolerogenic

therapies as the range of CD4 T cell clones to target and the timepoint this therapy

will have the greatest impact can be determined. Thus, we investigated the

evolution of CD4 TCR clonality by sequencing CDR3β regions of antigen

experienced endogenous CD4 T cells isolated from pLNs and joints at an early

and late phases of breach of self-tolerance models of inflammatory arthritis.

Induction of inflammatory arthritis resulted in the accumulation of endogenous

CD4 T cells displaying an antigen experienced phenotype. This population has

been previously shown to produce TNFα and IFNγ and also harbour autoreactive

clones capable of causing bone and cartilage degradation (18–21,39,40). CDR3β

analysis of this this population revealed that TCR repertoires from inflamed joints

at both the early and late phases displayed low clonality and diversity and were

dominated by few clones present at high frequencies. In contrast, pLNs at the late

phase displayed higher clonality and diversity compared with their counterparts at

the early phase and were mainly composed of a large number of clones that were

present at low frequencies. This indicates that pLNs become more polyclonal with

the progression of inflammatory arthritis and this polyclonality occurs in the pLNs

before possibly being reflected in inflamed joints. This is reinforced by the

observations that the most frequently occurring clones in the inflamed joint at the

late phase are also the most frequently occurring clones in the inflamed pLN at the

same phase. In addition, a study by Klarenbeek et. al (15) demonstrated that

repertoires from synovial fluid of patients with established RA have fewer highly

expanded clones compared with the synovial fluid taken from patients 6 months

after showing clinical symptoms, demonstrating the reduced clonality of CD4 T cell

repertoires in the joints of established RA patients. The polyclonality observed in

pLNs at the late phase may reflect changes in antigen specific responses due to

the release of neo-antigens resulting from continued joint damage and/or epitope

spreading, a phenomenon observed in RA patients (41). Indeed, PCA analysis of

TRBV genes of repertoires isolated from inflamed pLNs and joints at the early and

late phases highlighted the expanded range of TRBV genes present in repertoires

of pLNs at the late phase, indicating a possible change in antigen specific

responses with the progression of inflammatory arthritis. However, this can also be

explained by the re-establishment of lymphatic recirculation. Given that the

observations reported were exclusively on the antigen experienced CD4 T cell

population, re-establishment of lymphatic recirculation is an unlikely explanation

for the changes in the repertoire clonality and diversity observed. Moreover, the 10

most abundant antigen experienced clones correlate significantly between

inflamed pLNs and joints at the late phase and not in IFA controls. Inflamed joints

were also shown to have reduced diversity in the antigen experienced repertoire

compared to joint IFA controls. By taking these observations together, one can

deduce that changes in pLN repertoire clonality, diversity, and TRBV genes is in

fact antigen driven.

CDR3β amino acid sequence overlap analysis between repertoires isolated from

pLNs and joints at early and late phases of inflammatory arthritis highlighted that

fewer clones are shared between pLNs and joints at the late phase compared to

the early phase. This indicates that antigen specific responses differ between

these sites with time, driven by changes in clonality, diversity, and clones with

different TRBV genes in the pLN. Moreover, the reduced clonal overlap at the late

phase highlight the fact that inflamed pLNs and joints display different antigen

specific responses at the same phase. This disparity has important implications for

antigen specific therapies as specific pathogenic clones could be potentially

targeted in the lymph node before migrating, accumulating, and possibly

performing their effector functions in the joint. Moreover, analysis of clonal

compositions can also serve as a biomarker to monitor disease progression and

assess the therapeutic impact on changes in CD4 T cell clonality. Indeed, changes

in cellular compositions in inguinal lymph nodes has been detected in at-risk and

early RA patients (42) and treatment with RA therapeutics have been shown to

reduce clonal expansion of CD4 T cells (36).

Together, these data outline the evolution of clonal dynamics between inflamed

pLNs and their joints with the progression of inflammatory arthritis in that CD4

TCR clonality, begin by being similar between the two sites then change in the

pLN with the progression of the disease before potentially being mirrored in the

joint in time. These observations have several implications on improving current

antigen specific tolerogenic therapies. Firstly, understanding the clonal landscape

gives an indication on the range of CD4 T cells that need to be targeted to improve

the chances of modulating disease progression. Secondly, targeting expanded

CD4 T cell populations early may be more effective therapeutically as the CD4 T

cell repertoire is still restricted during the early phases, thus fewer populations

would need to be targeted. Lastly, clonality can be a biomarker of disease

progression and current therapeutics can be used to 'reset' the clonal landscape

to one reflecting the earlier phases of the disease, which may enhance

effectiveness of tolerogenic therapies and help reinstate self-tolerance and provide

long-lasting remission in RA patients.

ETHICS STATEMENT

This study was carried out at the University of Glasgow and performed in

accordance with the UK Home Office regulations and approved by the University

of Glasgow Ethics Committee.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

ACKNOWLEDGMENTS

We thank the staff within the Institute of Infection, Immunity and Inflammation Flow

Cytometry Facility and the Central Research Facility at the University of Glasgow

for technical assistance. We also thank Dr. Megan K. MacLeod for her insight and

for the critical discussions crucial for the development of this project.

FUNDING

This work was supported by Versus Arthritis. **RAB**, **JMB** and **PG** were supported

by the Arthritis Research UK (ARUK) programme grant number 19788 and the

Innovative Medicines Initiative EU-funded project Be The Cure (BTCURE) [grant

number 115142-2]. **JMB** and **PG** were also supported by the Arthritis Research

UK Rheumatoid Arthritis Pathogenesis Centre for Excellence (RACE) (grant

number 20298) and the Research into Inflammatory Arthritis Centre Versus

Arthritis (RACE) (grant number 22072). This work has received support from the

EU/EFPIA Innovative Medicines Initiative 2 Joint Undertaking RTCure grant no.

777357. **SAK** was supported by the Ministry of Higher Education Oman (MOHE)

throughout the study.

AUTHOR CONTRIBUTIONS

S.A.K and R.A.B designed the research and performed the experiments. S.A.K

analysed data, constructed figures, and wrote the manuscript. C.T.P and J.I.G

provided technical assistance and helped with analysis. T.D.O provided technical

assistance with the sequencing data, and J.M.B and P.G designed the research

and contributed to writing the paper.

REFERENCES

1.   Firestein GS, McInnes IB. Immunopathogenesis of Rheumatoid Arthritis. *Immunity*
(2017) **46**:183–196. doi:10.1016/j.immuni.2017.02.006

2.   Diaz-Gallo L-M, Martin J. PTPN22 splice forms: a new role in rheumatoid arthritis.
Genome Med (2012) **4**:13. doi:10.1186/gm312

3.   Holoshitz J. The rheumatoid arthritis HLA-DRB1 shared epitope. *Curr Opin*
Rheumatol (2010) **22**:293–298. doi:10.1097/BOR.0b013e328336ba63

4.   Lee H-S, Lee AT, Criswell LA, Seldin MF, Amos CI, Carulli JP, Navarrete C,
Remmers EF, Kastner DL, Plenge RM, et al. Several regions in the major
histocompatibility complex confer risk for anti-CCP-antibody positive rheumatoid
arthritis, independent of the DRB1 locus. *Mol Med Camb Mass* (2008) **14**:293–300.
doi:10.2119/2007-00123.Lee

5.   Okada Y, Wu D, Trynka G, Raj T, Terao C, Ikari K, Kochi Y, Ohmura K, Suzuki A,
Yoshida S, et al. Genetics of rheumatoid arthritis contributes to biology and drug
discovery. *Nature* (2013) **506**:376.

6.   Plenge RM, Padyukov L, Remmers EF, Purcell S, Lee AT, Karlson EW, Wolfe F,
Kastner DL, Alfredsson L, Altshuler D, et al. Replication of Putative Candidate-Gene
Associations with Rheumatoid Arthritis in >4,000 Samples from North America and
Sweden: Association of Susceptibility with PTPN22, CTLA4, and PADI4. *Am J Hum*
Genet (2005) **77**:1044–1060. doi:10.1086/498651

7.   Remmers EF, Plenge RM, Lee AT, Graham RR, Hom G, Behrens TW, de Bakker
PIW, Le JM, Lee H-S, Batliwalla F, et al. STAT4 and the Risk of Rheumatoid Arthritis
and Systemic Lupus Erythematosus. *N Engl J Med* (2007) **357**:977–986.
doi:10.1056/NEJMoa073003

8.   Blair HA, Deeks ED. Abatacept: A Review in Rheumatoid Arthritis. *Drugs* (2017)
77:1221–1233. doi:10.1007/s40265-017-0775-4

9.   van Beers JJBC, Schwarte CM, Stammen-Vogelzangs J, Oosterink E, Božič B,
Pruijn GJM. The rheumatoid arthritis synovial fluid citrullinome reveals novel
citrullinated epitopes in apolipoprotein E, myeloid nuclear differentiation antigen, and
β-actin. *Arthritis Rheum* (2013) **65**:69–80. doi:10.1002/art.37720

10. Burkhardt H, Sehnert B, Bockermann R, Engström A, Kalden JR, Holmdahl R.
Humoral immune response to citrullinated collagen type II determinants in early

rheumatoid arthritis. *Eur J Immunol* (2005) **35**:1643–1652.
doi:10.1002/eji.200526000

11. Ioan-Facsinay A, el-Bannoudi H, Scherer HU, van der Woude D, Ménard HA, Lora M, Trouw LA, Huizinga TWJ, Toes REM. Anti-cyclic citrullinated peptide antibodies are a collection of anti-citrullinated protein antibodies and contain overlapping and non-overlapping reactivities. *Ann Rheum Dis* (2011) **70**:188–193. doi:10.1136/ard.2010.131102

12. James EA, Rieck M, Pieper J, Gebe JA, Yue BB, Tatum M, Peda M, Sandin C, Klareskog L, Malmström V, et al. Citrulline-specific Th1 cells are increased in rheumatoid arthritis and their frequency is influenced by disease duration and therapy. *Arthritis Rheumatol Hoboken NJ* (2014) **66**:1712–1722. doi:10.1002/art.38637

13. Law S, Street S, Yu C-H, Capini C, Ramnoruth S, Nel HJ, van Gorp E, Hyde C, Lau K, Pahau H, et al. T-cell autoreactivity to citrullinated autoantigenic peptides in rheumatoid arthritis patients carrying HLA-DRB1 shared epitope alleles. *Arthritis Res Ther* (2012) **14**:R118. doi:10.1186/ar3848

14. Ikeda Y, Masuko K, Nakai Y, Kato T, Hasanuma T, Yoshino SI, Mizushima Y, Nishioka K, Yamamoto K. High frequencies of identical T cell clonotypes in synovial tissues of rheumatoid arthritis patients suggest the occurrence of common antigen-driven immune responses. *Arthritis Rheum* (1996) **39**:446–453. doi:10.1002/art.1780390312

15. Klarenbeek PL, de Hair MJH, Doorenspleet ME, van Schaik BDC, Esveldt REE, van de Sande MGH, Cantaert T, Gerlag DM, Baeten D, van Kampen AHC, et al. Inflamed target tissue provides a specific niche for highly expanded T-cell clones in early human autoimmune disease. *Ann Rheum Dis* (2012) **71**:1088–1093. doi:10.1136/annrheumdis-2011-200612

16. Stamenkovic I, Stegagno M, Wright KA, Krane SM, Amento EP, Colvin RB, Duquesnoy RJ, Kurnick JT. Clonal dominance among T-lymphocyte infiltrates in arthritis. *Proc Natl Acad Sci U S A* (1988) **85**:1179–1183.

17. VanderBorght A, Geusens P, Vandevyver C, Raus J, Stinissen P. Skewed T-cell receptor variable gene usage in the synovium of early and chronic rheumatoid arthritis patients and persistence of clonally expanded T cells in a chronic patient. *Rheumatol Oxf Engl* (2000) **39**:1189–1201.

18. Prendergast CT, Patakas A, Al-Khabouri S, McIntyre CL, McInnes IB, Brewer JM, Garside P, Benson RA. Visualising the interaction of CD4 T cells and DCs in the evolution of inflammatory arthritis. *Ann Rheum Dis* (2018) **77**:579–588. doi:10.1136/annrheumdis-2017-212279

19. Benson RA, Patakas A, Conigliaro P, Rush CM, Garside P, McInnes IB, Brewer JM. Identifying the Cells Breaching Self-Tolerance in Autoimmunity. *J Immunol* (2010) **184**:6378–6385. doi:10.4049/jimmunol.0903951

20. Maffia P, Brewer JM, Gracie JA, Ianaro A, Leung BP, Mitchell PJ, Smith KM, McInnes IB, Garside P. Inducing Experimental Arthritis and Breaking Self-Tolerance to Joint-Specific Antigens with Trackable, Ovalbumin-Specific T Cells. *J Immunol* (2004) **173**:151–156. doi:10.4049/jimmunol.173.1.151

21. Conigliaro P, Benson RA, Patakas A, Kelly SM, Valesini G, Holmdahl R, Brewer JM, McInnes IB, Paul Garside. Characterization of the anticollagen antibody response in a new model of chronic polyarthritis. *Arthritis Rheum* (2011) **63**:2299–2308. doi:10.1002/art.30413

22. Barnden MJ, Allison J, Heath WR, Carbone FR. Defective TCR expression in transgenic mice constructed using cDNA-based alpha- and beta-chain genes under the control of heterologous regulatory elements. *Immunol Cell Biol* (1998) **76**:34–40. doi:10.1046/j.1440-1711.1998.00709.x

23. Bolotin DA, Poslavsky S, Mitrophanov I, Shugay M, Mamedov IZ, Putintseva EV, Chudakov DM. MiXCR: software for comprehensive adaptive immunity profiling. *Nat Methods* (2015) **12**:380.

24. Nazarov VI, Pogorelyy MV, Komech EA, Zvyagin IV, Bolotin DA, Shugay M, Chudakov DM, Lebedev YB, Mamedov IZ. tcR: an R package for T cell receptor repertoire advanced data analysis. *BMC Bioinformatics* (2015) **16**: doi:10.1186/s12859-015-0613-1

25. Katayama CD, Eidelman FJ, Duncan A, Hooshmand F, Hedrick SM. Predicted complementarity determining regions of the T cell antigen receptor determine antigen specificity. *EMBO J* (1995) **14**:927–938.

26. Turner SJ, Doherty PC, McCluskey J, Rossjohn J. Structural determinants of T-cell receptor bias in immunity. *Nat Rev Immunol* (2006) **6**:883–894. doi:10.1038/nri1977

27. Fazou C, Yang H, McMichael AJ, Callan MFC. Epitope specificity of clonally expanded populations of CD8+ T cells found within the joints of patients with

inflammatory arthritis. *Arthritis Rheum* (2001) **44**:2038–2045. doi:10.1002/1529-0131(200109)44:9<2038::AID-ART353>3.0.CO;2-1

28. Iannone F, Corrigall VM, Kingsley GH, Panayi GS. Evidence for the continuous recruitment and activation of T cells into the joints of patients with rheumatoid arthritis. *Eur J Immunol* (1994) **24**:2706–2713. doi:10.1002/eji.1830241120

29. Shadidi KR, Aarvak T, Jeansson S, Henriksen JE, Natvig JB, Thompson KM. T-cell responses to viral, bacterial and protozoan antigens in rheumatoid inflammation. Selective migration of T cells to synovial tissue. *Rheumatol Oxf Engl* (2001) **40**:1120–1125. doi:10.1093/rheumatology/40.10.1120

30. Brennan FM, Hayes AL, Ciesielski CJ, Green P, Foxwell BMJ, Feldmann M. Evidence that rheumatoid arthritis synovial T cells are similar to cytokine-activated T cells: involvement of phosphatidylinositol 3-kinase and nuclear factor kappaB pathways in tumor necrosis factor alpha production in rheumatoid arthritis. *Arthritis Rheum* (2002) **46**:31–41. doi:10.1002/1529-0131(200201)46:1<31::AID-ART10029>3.0.CO;2-5

31. Brennan FM, Smith NM, Owen S, Li C, Amjadi P, Green P, Andersson A, Palfreeman AC, Hillyer P, Foey A, et al. Resting CD4+ effector memory T cells are precursors of bystander-activated effectors: a surrogate model of rheumatoid arthritis synovial T-cell function. *Arthritis Res Ther* (2008) **10**:R36. doi:10.1186/ar2390

32. Howson LJ, Napolitani G, Shepherd D, Ghadbane H, Kurupati P, Preciado-Llanes L, Rei M, Dobinson HC, Gibani MM, Teng KWW, et al. MAIT cell clonal expansion and TCR repertoire shaping in human volunteers challenged with Salmonella Paratyphi A. *Nat Commun* (2018) **9**:253. doi:10.1038/s41467-017-02540-x

33. Gomez-Tourino I, Kamra Y, Baptista R, Lorenc A, Peakman M. T cell receptor β-chains display abnormal shortening and repertoire sharing in type 1 diabetes. *Nat Commun* (2017) **8**:1792. doi:10.1038/s41467-017-01925-2

34. Risnes LF, Christophersen A, Dahal-Koirala S, Neumann RS, Sandve GK, Sarna VK, Lundin KEA, Qiao S-W, Sollid LM. Disease-driving CD4+ T cell clonotypes persist for decades in celiac disease. *J Clin Invest* (2018) **128**:2642–2650. doi:10.1172/JCI98819

35. Cantaert T, Brouard S, Thurlings RM, Pallier A, Salinas GF, Braud C, Klarenbeek PL, de Vries N, Zhang Y, Soulillou J-P, et al. Alterations of the synovial T cell repertoire in anti-citrullinated protein antibody-positive rheumatoid arthritis. *Arthritis Rheum* (2009) **60**:1944–1956. doi:10.1002/art.24635

36. Pierer M, Rossol M, Kaltenhäuser S, Arnold S, Häntzschel H, Baerwald C, Wagner U. Clonal expansions in selected TCR BV families of rheumatoid arthritis patients are reduced by treatment with the TNFα inhibitors etanercept and infliximab. *Rheumatol Int* (2011) **31**:1023–1029. doi:10.1007/s00296-010-1402-9

37. Wagner U, Pierer M, Kaltenhäuser S, Wilke B, Seidel W, Arnold S, Häntzschel H. Clonally expanded CD4+CD28null T cells in rheumatoid arthritis use distinct combinations of T cell receptor BV and BJ elements. *Eur J Immunol* (2003) **33**:79–84. doi:10.1002/immu.200390010

38. Wagner UG, Koetz K, Weyand CM, Goronzy JJ. Perturbation of the T cell repertoire in rheumatoid arthritis. *Proc Natl Acad Sci U S A* (1998) **95**:14447–14452. doi:10.1073/pnas.95.24.14447

39. Nickdel MB, Conigliaro P, Valesini G, Hutchison S, Benson R, Bundick RV, Leishman AJ, McInnes IB, Brewer JM, Garside P. Dissecting the contribution of innate and antigen-specific pathways to the breach of self-tolerance observed in a murine model of arthritis. *Ann Rheum Dis* (2009) **68**:1059–1066. doi:10.1136/ard.2008.089300

40. Platt AM, Gibson VB, Patakas A, Benson RA, Nadler SG, Brewer JM, McInnes IB, Garside P. Abatacept limits breach of self-tolerance in a murine model of arthritis via effects on the generation of T follicular helper cells. *J Immunol Baltim Md 1950* (2010) **185**:1558–1567. doi:10.4049/jimmunol.1001311

41. Monach PA, Hueber W, Kessler B, Tomooka BH, BenBarak M, Simmons BP, Wright J, Thornhill TS, Monestier M, Ploegh H, et al. A broad screen for targets of immune complexes decorating arthritic joints highlights deposition of nucleosomes in rheumatoid arthritis. *Proc Natl Acad Sci* (2009) **106**:15867. doi:10.1073/pnas.0908032106

42. Rodríguez-Carrio J, Hähnlein JS, Ramwadhdoebe TH, Semmelink JF, Choi IY, van Lienden KP, Maas M, Gerlag DM, Tak PP, Geijtenbeek TBH, et al. Brief Report: Altered Innate Lymphoid Cell Subsets in Human Lymph Node Biopsy Specimens Obtained During the At-Risk and Earliest Phases of Rheumatoid Arthritis. *Arthritis Rheumatol Hoboken NJ* (2017) **69**:70–76. doi:10.1002/art.39811

FIGURE LEGENDS

Figure 1.

Outline of model phases and number of endogenous, OT-II, and endogenous

antigen experienced CD4 T cells at both phases. **A)** Illustration of early and late

phases of inflammatory arthritis. **B)** Number of endogenous CD4 T cells (CD4+,

CD45+, CD45.1-) and OT-II T cells (CD4+, CD45+, CD45.1+) from pLNs and

joints of mice undergoing inflammatory arthritis at the early and late phases and

respective controls. **C)** Number of antigen experienced endogenous CD4 T cells

(CD4+, CD45+, CD45.1-, CD44hi) from pLNs and joints of mice undergoing

inflammatory arthritis at the early and late phases and respective controls. Data is

representative of mean ± SD with each point representing individual experimental

mice. Data represents three independent experiments for the early phase and two

independent experiments for the late phase with n=5 for each experiment. Groups

were compared using 2-way ANOVA and Student's t-tests. Stars represent the

following p-values: *<0.05, ** <0.01; ***<0.001; **** <0.0001; ns: not significant.

HAO, heat-aggregated OVA; IFA, incomplete Freund's adjuvant.

Figure 2

Characterisation of antigen experienced endogenous CD4 T cell repertoires from

inflamed pLNs and joints and PBS controls at the early phase. Proportion of the

top 10 most frequently occurring clones, followed by the next 25, 500, 3000,

10,000, and 100,000 most frequent clones contributing to the endogenous antigen

experienced CD4+ T cell repertoire in pooled inflamed pLNs and joint samples at

the early phase (HAO) and pooled PBS controls. Data represents 1 experiment

with pooled samples from 3 mice for the PBS sample and 5 mice for the inflamed

(HAO) sample.

Figure 3

Characterisation of antigen experienced endogenous CD4 T cell repertoires from

inflamed pLNs and joints and IFA controls at the late phase. Proportion of the top

10 most frequently occurring clones, followed by the next 25, 500, 3000, 10,000,

and 100,000 most frequent clones contributing to the endogenous antigen

experienced CD4+ T cell repertoire in **A)** inflamed pLNs, and **E)** inflamed joints at

the late phase (HAO+IFA) and IFA only controls. **B)** Proportion of the least

frequently occurring clones in inflamed pLNs at the late phase and IFA controls. **F)**

Percentage contribution of the top 10 most frequently occurring clones in inflamed

joints at the late phase and IFA controls. **C and G** clonality, represented as the

number of unique CDR3β DNA sequences, in **C)** inflamed pLNs and IFA controls,

and **G)** inflamed joints and IFA controls at the late phase. **D and H** diversity

indices of **D)** inflamed pLNs and IFA controls, and **H)** inflamed joints and IFA

controls at the late phase. Data is representative of mean ± SD with each point

representing individual experimental mice. Data represents 1 experiment with n=5

for the inflamed group and n=3 for the control group. Groups were compared using

unpaired Student's t-test. Stars represent the following p values: * <0.05; ns: not

significant.

Figure 4

Characterisation of antigen experienced endogenous CD4 T cell repertoires in

inflamed pLNs and joints at the early and late phases. Proportion of the top 10

most frequently occurring clones, followed by the next 25, 500, 3000, 10,000, and

100,000 most frequent clones contributing to the endogenous antigen experienced

CD4+ T cell repertoire in **A)** inflamed pLNs at the early and late phases, and **E)**

inflamed joints at the early and late phases. **B and F** percentage contribution of

the top 10 most frequently occurring clones in **B)** inflamed pLNs and **F)** inflamed

joints at the early and late phases. **C and G** clonality, represented as the number

of unique CDR3β DNA sequences, in **C)** inflamed pLNs, and **G)** inflamed joints at

the early and late phases. **D and H** diversity indices of **D)** inflamed pLNs, and **H)**

inflamed joints at the early and late phases. Data is representative of mean ± SD

with each point representing individual experimental mice. Data represents 1

experiment with n=5 for both the early and late phase groups. Groups were

compared using unpaired Student's t-test. Stars represent the following p values:

*** <0.001; **** <0.0001; ns: not significant.

Figure 5

Range of TRBV genes expressed in antigen experienced endogenous CD4 T cell

repertoires of inflamed pLNs and joints at the early and late phases. **A)** PCA plot

of TRBV gene frequency in repertoires of inflamed pLNs and joints at the early and

late phases. Frequency of **B)** TRBV gene expression, and **C)** TRBJ gene

expression in inflamed pLNs at the early and late phases. Graphs represent 1

experiment with n=5 for both the early and late inflamed groups. Groups were

compared using ordinary 2-way ANOVA with Sidak's multiple comparisons test.

Data presented as mean ±SD. Stars represent the following p values: * <0.05; **

<0.01; *** <0.001; **** <0.0001; ns: not significant.

Figure 6

Degree of CDR3β amino acid sequence overlap and correlation of most abundant

clones between inflamed pLNs and joints at the early and late phases. **A)**

Heatmaps of normalised number of CDR3β amino acid sequences (see methods)

between inflamed pLNs and joints at the early (left) and late (right) phases. **B)**

Graph representing the intersects (diagonal) of A) representing CDR3β sequence

overlap between inflamed pLNs and joint within the same animal. **C-E** Correlation

of the top 10 ranked clones in joints with their rank in respective pLN samples. The

rank of each of the top 10 clones in **C)** inflamed joint at the early phase, **D)**

inflamed joint at the late phase, and **E)** IFA control joint samples were plotted

against the rank of that clone in respective pLN samples. Data is representative of

one experiment with n=5 for the early and late phase inflamed groups and n=3 for

controls. Graph in B) is representative of mean ± SD with each point representing

individual experimental mice. * represents p<0.05. In C-E correlations (r) were

calculated using Spearman's correlation after testing for normality of the data

using the D'Agostino-Pearson test. p represents the p value.

Supplementary figure 1

Gating strategy to identify antigen experienced endogenous CD4 T cells from

pLNs and joints. Representative flow cytometry plots identifying antigen

experienced endogenous CD4 T cells (CD4+, CD45+, C45.1-, CD44hi) from **A)**

inflamed pLNs, and **B)** inflamed joints at the early phase of mice undergoing

inflammatory arthritis.

Supplementary figure 2

Representative flow cytometry plots identifying antigen experienced endogenous

CD4 T from **A)** pLN, and **B)** joint PBS controls at the early phase, **C)** pLN, and **D)**

joints IFA controls at the late phase, and **E)** inflamed pLN, and **F)** inflamed joints at

the late phase.

Supplementary figure 3

Degree of CDR3β amino acid sequence overlap between inflamed pLN and

inflamed joint samples at the early and late phases. Heatmaps of normalised

number of CDR3β amino acid sequences between **A)** inflamed pLNs, and **C)**

inflamed joints at the early and late phases. **B and D** Normalised overlap values

between **B)** inflamed pLN samples, and **D)** inflamed joint samples at the early and

late phases. Data is representative of mean ± SD with each point representing

individual experimental mice. Data represents 1 experiment with n=5 for both the

early and late inflamed groups. Groups were compared using unpaired Student's t-

test. Stars represent the following p values: ** <0.01; ns: not significant.

Supplementary table 1

The top 10 frequently occurring clones inflamed joint samples at the early phase

ranked in terms of frequency in respective pLNs. The top 10 frequently occurring

clones in the antigen experienced CD4+ T cell repertoire were found in each of the

inflamed joint samples at the early phase and searched for in respective pLN

samples. The CDR3β amino acid sequence, associated TRBV gene, read count,

and rank of the clone in the pLN sample was obtained. The rank of each of the top

10 clones in the joint are also included in the table. Data is representative of 1

experiment with n=5.

Supplementary table 2

The top 10 frequently occurring clones inflamed joint samples at the late phase

ranked in terms of frequency in respective pLNs. The top 10 frequently occurring

clones in the antigen experienced CD4+ T cell repertoire were found in each of the

inflamed joint samples at the late phase and searched for in respective pLN

samples. The CDR3β amino acid sequence, associated TRBV gene, read count,

and rank of the clone in the pLN sample was obtained. The rank of each of the top

10 clones in the joint are also included in the table. Data is representative of 1

experiment with n=5.

Fig 1.

Fig 2.

A

B

Clonality, diversity, and number of End. CD4 T cells in early inflamed pLNs and joints and PBS controls

Sample	pLN HAO	pLN PBS	Joint HAO	Joint PBS
No. of End. CD4+ T cells	3,113,218	224,978	58,041	1424
Clonality	4181	4347	4549	885
Diversity index (DI)	6.4	16.3	0.72	9.3

Data represents 1 experiment with pooled samples from 3 mice for the PBS sample and 5 mice for the inflamed (HAO) sample.

Fig 3.

Fig 4.

Fig 5

A

B

C

Fig 6