Perphenazine-macrocycle conjugates rapidly sequester the Aβ42 monomer and inhibit amyloid formation

Sarah R. Ball[1], Julius S. P Adamson[2], Michael A. Sullivan[1], Manuela R. Zimmermann[3], Victor Lo[1], Maximo Sanz-Hernandez[4], Frank Jiang[2], Ann H. Kwan[5], Eryn L. Werry[2, 6], Tuomas P. J. Knowles[3, 7], Michael Kassiou[2], Georg Meisl[3], Matthew H. Todd[8*], Peter J. Rutledge[2*] and Margaret Sunde[1*]

[1]School of Medical Sciences, The University of Sydney, Sydney, Australia. [2]School of Chemistry, The University of Sydney, Sydney, Australia. [3]Centre for Misfolding Diseases, Department of Chemistry, University of Cambridge, Cambridge UK. [4]Department of Life Sciences, Imperial College London, South Kensington, SW7 2AZ, UK. [5]School of Life and Environmental Sciences, The University of Sydney, Sydney, Australia. [6]Brain and Mind Centre, The University of Sydney. [7]Cavendish Laboratory, University of Cambridge, JJ Thomson Avenue, CB4 0HE, UK. [8]School of Pharmacy, University College London, London UK.

*Co-corresponding authors: Margaret Sunde, Peter Rutledge, and Matthew Todd

Email: margaret.sunde@sydney.edu.au; peter.rutledge@sydney.edu.au; matthew.todd@ucl.ac.uk

Keywords

amyloid, inhibition, kinetic analysis, monomer sequestration, oligomers, Alzheimer's disease

Author Contributions

Conceptualization: Sarah R. Ball, Julius S. Adamson, Manuela R. Zimmermann, Frank Jiang, Georg Meisl, Matthew H. Todd, Peter J. Rutledge and Margaret Sunde.
Data Curation: Sarah R. Ball, Julius S. Adamson and Maximo Sanz-Hernandez.
Formal Analysis: Sarah R. Ball, Michael A. Sullivan, Manuela R. Zimmermann, Maximo Sanz-Hernandez and Georg Meisl.
Funding Acquisition: Peter J. Rutledge and Margaret Sunde.
Investigation: Sarah R. Ball, Julius S. Adamson, Michael A. Sullivan, Victor Lo, Maximo Sanz-Hernandez, Frank Jiang, Peter J. Rutledge and Margaret Sunde.
Methodology: Sarah R. Ball, Julius S. Adamson, Michael A. Sullivan, Manuela R. Zimmermann, Victor Lo, Maximo Sanz-Hernandez, Frank Jiang, Ann H. Kwan, Tuomas P. Knowles, Georg Meisl, Eryn L. Werry, Matthew H. Todd, Peter J. Rutledge and Margaret Sunde.
Project Administration: Peter J. Rutledge and Margaret Sunde.
Resources: Michael Kassiou, Peter J. Rutledge and Margaret Sunde.
Supervision: Ann H. Kwan, Tuomas P. Knowles, Eryn L. Werry, Michael Kassiou, Matthew H. Todd, Peter J. Rutledge and Margaret Sunde.
Validation: Sarah R. Ball, Julius S. Adamson, Frank Jiang and Margaret Sunde.
Visualization: Sarah R. Ball, Julius S. Adamson, Michael A. Sullivan, Manuela R. Zimmermann, Maximo Sanz-Hernandez, Georg Meisl, Peter J. Rutledge and Margaret Sunde.
Writing - Original Draft Preparation: Sarah R. Ball, Julius S. Adamson, Michael A. Sullivan, Peter J. Rutledge and Margaret Sunde.
Writing - Review & Editing: Sarah R. Ball, Julius S. Adamson, Michael A. Sullivan, Manuela R. Zimmermann, Victor Lo, Maximo Sanz-Hernandez, Ann H. Kwan, Tuomas P. Knowles, Georg Meisl, Eryn L. Werry, Michael Kassiou, Matthew H. Todd, Peter J. Rutledge and Margaret Sunde.

Abstract

Alzheimer's disease is imposing a growing social and economic burden worldwide and effective therapies are required. Strategies aimed at the removal of fibrillar plaques formed by the amyloid-β peptide have not proved therapeutic and the focus has shifted to approaches that target the cytotoxic oligomeric amyloid-β species that are populated before fibrils are deposited. We have designed and synthesized perphenazine-cyclam conjugates that specifically and rapidly bind to the monomeric form of Aβ42, reducing the production of both cytotoxic oligomers and amyloid fibrils. We have applied detailed kinetic analysis and NMR spectroscopy to show that the perphenazine-cyclam conjugates divert the Aβ42 monomer into amorphous aggregates that are not toxic to differentiated SH-SY5Y cells *in vitro.* Unlike most other amyloid inhibitors studied to date, these conjugates inhibit oligomer and fibril assembly even in the presence of pre-formed fibrillar seeds, demonstrating that they act through a monomer sequestration mechanism. These modular, three-dimensional conjugates therefore effectively prevent monomer-dependent secondary nucleation, the autocatalytic process that generates the majority of toxic oligomers.

Introduction

Alzheimer's disease (AD) is the most common form of dementia and one of the most devastating diseases of the current age. There are currently no approved disease-modifying therapies for this neurodegenerative disease. Although the full sequence of critical molecular events that lead to neuronal death is yet to be defined, self-assembly of the amyloid-β peptide (Aβ) into soluble oligomers and amyloid fibrils has been shown to generate neurotoxic oligomeric species[1-3]. These oligomers impact neuronal activity through effects on cell membranes, calcium homeostasis, and glutamate reuptake, amongst other deleterious mechanisms[4-10]. The failure of clinical trials aimed at reducing the accumulation of mature Aβ fibrils in extracellular plaques has seen efforts shift towards intervention at earlier stages in the pathway, with the aim of inhibiting formation of these toxic, pre-fibrillar oligomers[11].

Aβ amyloid formation is a complex, multiple-step process, initiated by primary nucleation from Aβ monomers. This is followed by recruitment of additional monomers, leading to the formation of multiple different soluble oligomers with a range of sizes and toxicities[12]. As amyloid fibrils form, they offer sites for elongation at their ends, and for secondary nucleation on their lateral surfaces[13, 14], generating further oligomeric species. Monomer-dependent secondary nucleation is a significant source of cytotoxic oligomeric Aβ42 species and is the dominant nucleation process even in cerebrospinal fluid and the presence of phospholipid membranes[15].

Effective inhibition of Aβ-associated toxicity is likely to require a mechanism that is effective against all steps of the pathway[16]. Several small-molecule compounds have been shown to suppress primary nucleation or to accelerate fibril formation[17-19] and have shown promise in animal models but are yet to show efficacy in clinical trials[20]. Some small molecules increase the population of neurotoxic Aβ oligomers[21] and other proposed inhibitors are not effective in the presence of pre-formed fibrils[17, 22]. The molecular chaperone domain pro-SPC and certain antibody fragments are inhibitors of secondary nucleation[23, 24], as are some small molecules[18, 25]. The efforts to identify effective modulators of the Aβ assembly process have been hampered by the use of poorly characterised preparations of the peptide and by testing against Aβ40, the less amyloidogenic and less clinically relevant isoform, rather than Aβ42 [26-28]. The growing impact of AD necessitates continued efforts directed at the development of therapeutic modulators.

We have shown previously that cyclam derivatives with triazole-linked pendant groups can adopt a range of different, defined pendant geometries depending on the connectivity of the triazole linker and the metal ion coordinated by the macrocycle[29-31]. Combining these insights with the documented capacity of polymeric perphenazine conjugates to modulate Aβ aggregation[28], we designed a series of molecules in which cyclam is linked via click-derived triazoles to pendant groups that are known to interact with Aβ42.

Quantitative kinetic analysis demonstrates that this series of molecules effectively and rapidly sequesters the monomeric form of Aβ42. We show that this monomer sequestration activity inhibits all downstream events that are dependent on monomer concentration, including suppression of the formation of toxic oligomers and amyloid fibril assembly, even in the presence of preformed fibril seeds. These novel molecules redirect the Aβ42 formation pathway towards the generation of amorphous aggregates that are not toxic to differentiated, neuron-like SH-SY5Y cells. This study provides the first demonstration of a reduction in neurotoxicity resulting from the designed diversion of the Aβ42 monomer into non-toxic oligomers.

Results

Perphenazine-cyclam conjugates inhibit Aβ42 amyloid assembly

A series of cyclam conjugates was designed to enable a systematic investigation of the Aβ aggregation and amyloid assembly pathways. Six parent cyclam derivatives were prepared: the mono-substituted naphthyl (**C1**), dopamine (**C4**) and perphenazine (**C7**) analogues, and corresponding bis-substituted compounds with naphthyl (**C10**), dopa (**C13**) and perphenazine (**C16**) pendants, along with their zinc(II)

and copper(II) complexes (mono-naphthyl **C2**, **C3**; mono-dopa **C5**, **C6**; mono-perphenazine **C8**, **C9**;
bis-naphthyl **C11**, **C12**; bis-dopa **C14**, **C15**; and bis-perphenazine **C17**, **C18**) (Fig. 1A and SI Fig. S1A).
Initial screening of these compounds showed the bis-perphenazine derivative (ligand **C16**) to have
promising capacity to suppress amyloid formation by both Aβ40 and Aβ42 (SI, Fig. S1B and C).

We prepared the bis-perphenazine ligand (**C16**) and its zinc (**C17**) and copper (**C18**) analogues and
determined the ability of these three cyclam conjugates to inhibit amyloid formation by Aβ42 (SI, Fig.
S1C,D). We have characterised the inhibitory activity of the molecules against Aβ42 amyloid assembly.
This isoform is clinically associated with an increase in amyloid deposition[32] and we have utilised highly
purified, monomeric recombinant Aβ42, prepared using the methods of Linse and colleagues[33, 34]. Aβ42
prepared according to these protocols displays highly reproducible kinetics of amyloid assembly. The
three cyclam conjugates **C16**, **C17** and **C18** were found to be potent inhibitors of Aβ42 amyloid
formation, as monitored by ThT fluorescence (Fig. 1; SI Fig. S2 and Fig. S3).

Fig. 1. Time courses of fibril formation by Aβ42 show concentration-dependent inhibition of aggregation
by **C16**, **C17** and **C18**. *(A)* The lead compounds used in this study combine a cyclam macrocycle
(purple) with pendant perphenazine moieties (cyan), conjugated via a click-derived triazole (black), and
deployed either as the unmetallated HCl salt **C16**, the zinc perchlorate complex **C17**, or the copper(II)
perchlorate **C18**. *(B)* 10 µM monomeric Aβ42 in the absence or presence of either 1% DMSO (dark
grey) or 10 – 50 µM of **C16** (pink), **C17** (blue) and **C18** (green). Lower section displays an enlarged 10-

128 hour window. N = 3 (SI Fig S2). *(C)* 2.5 µM monomeric Aβ42 in the presence of either 1% DMSO (dark
129 grey) or 0.1, 1, 2, 3 or 5 µM of compounds. N = 4 (SI Fig S3). All samples were measured in triplicate
130 under quiescent conditions at 37 °C. Traces have been corrected for time zero and data normalised as
131 a function of the maximum fluorescence intensity, using GraphPad Prism.
132
The inhibitors demonstrated a concentration-dependent effect, and at a 1:5 molar ratio of
Aβ42:compound, using 10 µM Aβ42, amyloid assembly was almost completely suppressed, as judged
by the reduction in the level of Thioflavin T (ThT) fluorescence observed (Fig. 1B). Assays at lower
Aβ42 concentration, using a range of inhibitor concentrations that allowed the plateau phase of amyloid
assembly to be reached, revealed that all three compounds also introduce an increase in lag phase.
The length of the lag phase increases with increasing inhibitor concentration (Fig. 1C). The zinc(II)
complex **C17** displayed the strongest inhibition, followed by copper(II) complex **C18**, and the
uncomplexed ligand **C16**, highlighting the capacity of the central metal ion to influence the
three-dimensional shape of the inhibitor – in particular the relative orientation of the two pendant groups
[29, 30, 35] – and resulting impact on the mechanism of Aβ42 amyloid inhibition. The exact nature of this
inferred variation in shape between **C16**, **C17** and **C18** remains to be elucidated.

Perphenazine alone does not display strong inhibition of Aβ42 amyloid formation under these
conditions, highlighting the fact that it is not the perphenazine moiety *per se* that is responsible for the
inhibitory activity (SI Fig. S4). The observed inhibition is also not due to an effect of unchelated metal
ions directly on Aβ42. Previous studies have demonstrated that Zn(II) and Cu(II) cations can themselves
inhibit Aβ42 amyloid formation[36-38], reflected by an increase in the lag phase or a decrease in the
observed ThT intensity (SI Fig. 5A). However, we observed that **C17** and **C18** remain effective inhibitors
of Aβ amyloid formation even in the presence of the chelating agent EDTA, which would sequester any
free metal ions and block their inhibitory capacity (SI Fig. 5B).

The perphenazine conjugates divert Aβ42 from amyloid assembly towards amorphous
aggregates

A sensitive protein concentration assay confirmed that when the plateau of assembly was reached, in
samples containing Aβ42 only, or Aβ42 in the presence of a 5-fold molar excess of inhibitor, all Aβ42
was converted to an insoluble form that was readily pelleted by centrifugation (SI Fig. S6). Transmission
electron microscopy was used to examine the final morphology of the Aβ42 material present after ~40 h
incubation with compounds or 1% DMSO as control. An inhibitor concentration dependent shift from
the characteristic fibrillar morphology of Aβ42 amyloid (Fig. 2A) to amorphous aggregates (Fig. 2B, C
& D) was observed. The amorphous aggregates had a beads-on-a-string appearance, with the "beads"
approximately 20 nm wide and the "strings" >1 µm in length. Only amorphous material was observed
when Aβ42 was incubated in the presence of **C16**, even at 1:1 molar ratio of **C16**:Aβ42, while a
combination of fibrillar and amorphous material was observed in samples prepared at 1:1 molar ratios
of **C17**:Aβ42 and **C18**:Aβ42 (SI Fig. S7).

Fig. 2. Transmission electron microscopy images showing representative examples of the morphology of fibrils formed by 10 µM Aβ42 alone (A) or aggregates formed by 10 µM Aβ42 in the presence of 50 µM **C16** (B), **C17** (C) and **C18** (D). Scale bar represents 200 nm.

The perphenazine conjugates reduce neurotoxicity associated with Aβ oligomers

Many groups have demonstrated that soluble intermediate species formed by Aβ42 *in vitro* are neurotoxic[1-3, 39]. Accordingly, the ability of compounds **C16**–**C18** to prevent the formation of toxic Aβ42 species was indexed using SH-SY5Y cells which had been differentiated to an AD-appropriate phenotype by treatment with retinoic acid and brain-derived neurotrophic factor[40]. Lyophilised, monomeric Aβ42 was treated with hexafluoro-2-propanol (HFIP), lyophilised, and then resuspended in PBS, before being diluted in cell media and added immediately to cells. Aβ42 fibrils were prepared by incubating monomeric Aβ42 solution in cell media at 37 °C for 24 hours, a time point significantly past the plateau level in unshaken ThT experiments using cell media (SI Fig. S8). The compounds were added to differentiated SH-SY5Y cells alone or added together with Aβ42 in a monomeric or fibrillar form. After 24 hours, the number of metabolically-active cells was measured using a CTB fluorescence assay. The CellTiter-Blue® (CTB) assay of cell viability was chosen, given published concerns regarding the use of MTT with Aβ42[1].

By themselves, compounds **C16**, **C17** and **C18** displayed no significant toxicity towards differentiated SH-SY5Y cells, relative to vehicle only (Fig. 3A). Exposure of these cells to freshly prepared monomeric Aβ42 resulted in significant ($P < 0.001$) toxicity compared to vehicle, as expected due to the generation of oligomeric species, while the addition of **C16**, **C17** or **C18** led to significant protection against this neurotoxicity (Fig. 3B). Perphenazine was not cytotoxic by itself and imparted a similar level of protection as **C16**, **C17** and **C18**. These results contrast with the observed neurotoxicity of epigallocatechin gallate (EGCG) at the same concentrations and the lack of protection by this polyphenol against cytotoxicity induced by addition of monomeric Aβ42 (Fig. 3A,B). When Aβ42 was prepared in fibrillar form and then added to cells, no cytotoxicity was observed with fibrils alone or in the presence of compounds (Fig. 3C). Following from these experiments, independent concentration response assays were performed with **C16**. The IC_{50} for **C16** was determined to be 2.33 µM (n = 3; SD = 0.89) (Fig. 3D). Here we have used an established assay of neurotoxicity, with differentiated SH-SY5Y cells and well-characterised preparations of monomeric and fibrillar Aβ42. Although the origins of the neurotoxicity resulting from the conversion of Aβ42 from monomeric to fibrillar form *in vitro* are

198 not defined, these results provide clear evidence that the generation of amorphous Aβ42 aggregates in
199 the presence of **C16**–**C18** prevents the formation of neurotoxic species.

200
Fig. 3. Cell viability was measured using the CellTiter Blue® assay after 24 h incubation with vehicle or
10 µM compound, with or without 20 µM Aβ42. (*A*) Compounds alone. (*B*) Compounds incubated with
monomeric Aβ42. (*C*) Compounds incubated with fibrillar Aβ42. The figure displays the mean ± SD of
N ≥ 3 independent experiments. One-way ANOVA followed by a Bonferroni's multiple comparisons test
at the 0.05 level was used to determine the difference between each condition and vehicle control (****
$p < 0.0001$, *** $p < 0.001$, * $p < 0.05$, NS = non-significant). (*D*) Representative concentration-response
curve for **C16** incubated with Aβ42 showing mean ± SD of duplicates. $IC_{50} = 2.33$ µM (SD = 0.89). N =
3.
209

C16, C17 and C18 inhibit Aβ amyloid formation through sequestration of Aβ42 monomers

The microscopic processes underlying amyloid formation of Aβ42 are primary nucleation, elongation
and secondary nucleation[41, 42]. Their relative importance can be quantified through their rate constants,
obtained by fitting integrated rate laws within the framework of chemical kinetics[43]. Data measured at
2.5 µM and 10 µM Aβ42, in the presence of a range of concentrations of inhibitor, were analysed using
the online fitting software AmyloFit. The rate constants of the unperturbed systems agreed well with
similar experiments[13]. To probe the mechanism of action of the inhibitory compounds **C16**, **C17** and
C18, we varied systematically one rate constant in the rate law, or the level of available monomer,
respectively[43]. By comparing how well these modified rate laws can describe the data, a mechanism
that primarily involves inhibition of primary nucleation could be ruled out for **C16**, **C17** and **C18** (Fig. 4A,
B and SI Fig. S9-13) as the origin of the changes in the kinetic behaviour in the presence of the inhibitor.

Fig. 4. Chemical kinetics analysis reveals inhibition through monomer sequestration. To elucidate the major mechanism of inhibition, normalised aggregation curves measured in thioflavin T fluorescence assays were fitted under the constraint that deviation from the unperturbed system was allowed in only one microscopic step or the free monomer concentration, respectively. Representative curves shown in A and B, all fits shown in the SI. *(A)* The observed inhibition cannot be described through inhibition of primary nucleation. *(B)* Inhibition through monomer sequestration is a plausible mechanism based on the kinetic profiles. *(C)* Microscopic steps underlying amyloid formation, with the suggested mode of inhibition through monomer sequestration.

The introduction of small numbers of preformed fibril seeds at the beginning of the aggregation reaction can be used to bypass primary nucleation. These compounds remain effective as inhibitors of Aβ42 amyloid fibril formation even when exogenous fibril seeds are present (Fig. 5). In the presence of 2% or 5% seeds, and the presence of **C16–C18** at 1:2.5 or 1:5 Aβ42:compound molar ratio resulted in inhibition of Aβ42 assembly, similar to the level of inhibition observed in the absence of pre-formed seeds (Fig. 5 and SI Fig. S14). In addition to ruling out the blocking of primary nucleation as the source of the inhibitory effect, the results observed in the presence of seeds, when the aggregation reaction reaches completion much sooner, suggest that binding of the compounds to the monomeric form of Aβ42 occurs rapidly.

Fig. 5 Characterisation of the effects of compounds on seeded Aβ42 aggregation. 10 μM monomeric
Aβ42, in the absence of seeds (black) or presence of 2% (top panel) or 5% (bottom panel) seeds and
1% DMSO (dark grey), 25 or 50 μM of compounds. N = 3 (SI Figs S14). All samples were measured
in triplicate under quiescent conditions at 37 °C. Traces have been corrected for time zero and data
normalised as a function of the maximum fluorescence intensity, using GraphPad Prism.

Based on the fits to the normalised kinetic data alone, mechanisms based on inhibition of secondary
nucleation, inhibition of elongation, and monomer sequestration were similarly likely as the main modes
of action of compounds **C16**–**C18** (SI Fig. S9-13). However, we found a clear correlation between the
ThT intensity at completion of the aggregation reaction and the effective monomer concentration
determined from the fits (Fig. 6A, B, C insets). As the fitting was performed on normalised kinetic data,
these two quantities constitute independent measures of the same physical property. The correlation
between an equilibrium measure, the plateau level, and a parameter estimated on the basis of
normalised kinetics, the effective monomer concentration, provides further independent evidence for
monomer sequestration by these inhibitors. In addition to accounting for the delay in aggregation and
decrease in ThT signal, monomer sequestration by the inhibitors also provides a mechanistic
explanation for the appearance of amorphous material as discussed above. These fits were made
assuming that this monomer sequestration proceeds much faster than aggregation and can thus be
modelled by a reduced effective monomer concentration.

Further support for monomer sequestration as the dominant mode of inhibition of Aβ42 aggregation by
C16, **C17** and **C18** was gained by obtaining consistent values for the apparent dissociation constant of
monomer and inhibitor, K_D, by orthogonal methods. First, the effective monomer concentration obtained
through the chemical kinetics analysis at different monomer and inhibitor concentrations was used to
construct monomer-inhibitor binding curves (Fig. 6A, B, C). To fit these data, we assumed a simple one-
to-one binding, yielding K_D values of 7.6 μM (95% CI, 6.6–9.1) for **C16**, 5.7 μM (95% CI, 4.6–7.1) for
C17 and 7.4 μM (95% CI, 6.2–9.1) for **C18**.

Secondly, we utilised one-dimensional [1]H NMR spectroscopy to determine an apparent binding
constant for the interaction between the inhibitors and Aβ42. For each of **C16**, **C17** and **C18**, inhibitor
was titrated into a solution containing 10 μM unlabelled Aβ42, over the inhibitor concentration range of
1–50 μM. A decay profile in the protein signal intensity across the spectrum was observed with
increasing concentration of all three inhibitors (Fig. 6 and SI Fig. S15). A 5:1 ratio of inhibitor:Aβ42
resulted in a reduction of proton signals to baseline, consistent with the sequestration of monomer into

very large and/or insoluble assemblies, the precipitation of material during the course of the NMR titrations, and the amorphous aggregates observed by TEM (Fig. 3). The signal decay curves were fitted to a quadratic equation for the calculation of the apparent K_D of each compound, yielding 4.6 μM (95% CI, 3.1–6.5) for **C16**, 3.4 μM (95% CI, 2.2–4.9) for **C17** and 3.7 μM (95% CI, 2.4–5.3) for **C18**. The observed effect of these perphenazine cyclam conjugates on Aβ42 assembly is different to that of the polyphenol EGCG, where the presence of EGCG results in only a small decrease in the intensity of the observed 1H spectrum from Aβ42, up to 31% signal loss at a 5-fold excess of small molecule (Fig. 6D and SI Fig. S15). This is evidence that upon addition of EGCG, the monomeric Aβ42 species is still the most prevalent in the sample. In contrast, when the less aggregation-prone Aβ40 interacts with EGCG, specific interactions are observed by solution NMR and it forms insoluble oligomers that are in equilibrium with soluble Aβ40[44].

Fig. 6. Dissociation constants determined from ThT assays and ¹H NMR. Monomer-inhibitor binding curves for *(A)* **C16**, *(B)* **C17** and *(C)* **C18**, with correlation between the effective free monomer concentration and the absolute increase in ThT signal shown as insets. One-dimensional ¹H NMR spectroscopy to determine apparent K_D for compounds. Unlabelled 10 μM Aβ42 (dark grey) was incubated with 1–50 μM *(D)* **C16**, *(E)* **C17**, *(F)* **C18** and *(G)* **EGCG**, and spectra recorded at 4 °C. Methyl proton region is displayed, all regions analysed are shown in SI Fig. 15. Increasing concentration of

292 **C16–C18** resulted in a decay in protein signal intensity. Binding curves for *(H)* **C16,** *(I)* **C17** and *(J)* **C18**
293 constructed from signal decay curves and used to calculate the apparent K_D.
294
A monomeric sample of recombinant [15]N-labeled Aβ42 was prepared and analysed in the absence and
presence of **C16** to probe the regions of the polypeptide involved in the interaction. The [15]N HSQC
spectrum collected in the absence of any added compound reflected the high quality of the recombinant,
monomeric Aβ42 sample and the chemical shifts were superimposable upon those reported by others
previously for Aβ42[45] (SI Fig. S16). Addition of **C16** resulted in an immediate reduction in signal intensity
that was uniform across all peaks in the spectrum. Protein that is not sequestered into an amorphous
aggregate remains in the solution at the 1:1 ratio of protein:inhibitor. No peak-specific changes in signal
intensity or chemical shift were detected that could be attributed to binding of **C16** to a distinct region
of monomeric Aβ42. Instead, the loss of signal is consistent with the binding of monomer to **C16**
resulting in the generation of a conformation that is highly aggregation-prone and rapidly forms large
structures, undetectable by solution NMR.

Discussion

We have designed a class of compounds that specifically sequester the monomeric form of Aβ42 into
non-fibrillar assemblies that are different to the cytotoxic oligomeric forms associated with Aβ42 amyloid
formation. The process by which monomeric Aβ42 converts to the mature fibrillar form detected in
amyloid plaques involves multiple oligomeric and prefibrillar species that may act as reactants,
products, intermediates, and catalysts of the process and contribute to toxicity to different extents. The
ability to interrogate a change in aggregation kinetics caused by an inhibitor, and thereby identify the
mechanism of inhibition, offers new opportunities to understand the consequences of intervention at
defined points within the Aβ42 assembly pathway. Many studies of peptide-based inhibitors that
incorporate "disruption elements" report success in preventing the formation of the β-structure that
stabilises amyloid fibrils and can influence elongation[46] and inhibitors have been described that
specifically inhibit secondary nucleation[18], but the impact of these interventions on residual Aβ monomer
concentration has not always been described. The kinetic theory of protein aggregation inhibition
reveals that inhibitors such as the cyclam-perphenazine conjugates described here, which sequester
monomeric Aβ42, will reduce the rates of primary nucleation, elongation and secondary nucleation, and
are therefore likely to have a large impact on amyloid assembly overall[16].

In addressing AD the sequestration of monomer into non-toxic assemblies and consequent reduction
of oligomer formation, could be beneficial. Studies of the effect of EGCG on Aβ42 indicate that it
converts the peptide to off-pathway oligomers and remodels pre-formed fibrils, although these effects
are not rapid[26, 47, 48]. The intrinsically disordered and dynamic nature of monomeric Aβ42 has impeded
the use of traditional approaches that focus on identifying chemical moieties that bind to defined sites
on the target. Recently a small molecule (10074-G5) that binds to the monomeric form of Aβ42 has
been reported[16, 22].The binding of 10074-G5 to Aβ42, with K_D of ~40 μM, sequesters Aβ42 in its soluble
form and reduces its hydrophobicity. In contrast, we observed an immediate uniform reduction in signal
intensity across the spectrum when **C16** was added to Aβ42 at a 1:1 ratio, indicating that **C16**
sequesters Aβ42 in a polymeric, insoluble assembly. The small molecule 10074-G5 reduces functional
deficits in a *C. elegans* model of Aβ42-associated toxicity, however its impact on neuronal toxicity is yet
to be reported and it does not display a concentration-dependent inhibition of seeded assembly *in vitro*.
Thus, further work is required to define the relative benefit of Aβ42 sequestration in monomeric versus
non-toxic oligomeric form.

The inventive modular design of the compounds described here offers a toolkit for future elucidation of
the key moieties and properties that influence Aβ42 monomer binding and subsequent formation of
amorphous aggregates. Compounds that were effective at Aβ42 monomer sequestration protected
against Aβ42-induced cytotoxicity. This scaffold provides a base for future rational design of more
potent compounds that act through the same mechanism, yet which are accessible via a divergent
synthetic strategy. This class of compounds can provide wide mass-to-charge and hydrophobic-to-

hydrophilic surface area ratios, while the attachment of both rigid and flexible pendants provides a great diversity of possible interactions with client protein targets. The switchable geometry allows maintenance of a constant periphery while varying the gross molecular shape. Given that self-assembly and aggregation of proteins are three-dimensional processes, this level of structural control has significant potential to contribute to effective inhibition. AD is one of over 50 human disorders considered to be proteinopathies. The critical pathogenic species associated with protein aggregation and amyloid formation in each of these disorders likely have unique features. However, a similar dissection of assembly pathways with modular scaffolds decorated with protein-specific binding moieties may reveal points for therapeutic intervention in other protein aggregation-related diseases.

Materials and Methods

Synthesis of lead compounds C16–C18.
Full synthetic procedures and characterisation data are detailed in the Supporting Information.

Reaction scheme 1: Cyclam (1,4,8,11-tetraazacyclotetradecane) **1** was converted to the di-Boc-di-propargyl derivative **2** in 68% yield over four steps using previously reported methods[49, 50]. Perphenazine **3** was converted to the azide **4** (79%) by activating its primary alcohol with diphenylphosphoryl azide (DPPA) and displacing with sodium azide in dimethylformamide (DMF). The azide **4** (2 equivalents) and bis-alkyne **2** were coupled using a copper-catalysed azide/alkyne cycloaddition reaction (CuAAC) to yield the protected ligand **5** (46%), before the Boc groups were removed using 4 M HCl in dioxane to afford **C16** (74%). The HCl salt **C16** was converted to its zinc(II) (**C17**) and copper(II) (**C18**) complexes by first treating with Ambersep 900 hydroxide form resin to generate the free amine, then stirring overnight at room temperature with either zinc(II) perchlorate hexahydrate (for **C17**, 63%) or copper(II) perchlorate hexahydrate (for **C17**, 64%). All compounds were dissolved in 100% DMSO and stored at 4 °C until use. Reagents and conditions: i. $CuSO_4.5H_2O$ (0.2 equiv.), sodium ascorbate (0.5 equiv.), THF/H_2O (2:3), rt, 16 h, 46%; ii. 4 M HCl in dioxane (excess), 0 °C, 1 h, 74%; iii. Ambersep 900 hydroxide form resin, MeOH, 15 min, then $Zn(ClO_4)_2$, MeOH, reflux, 16 h, 63%; iv. Ambersep 900 hydroxide form resin, MeOH, 15 min, then $Cu(ClO_4)_2$, MeOH, reflux, 16 h, 64%.

Aβ42 Overexpression and purification. Overexpression and purification of unlabelled Aβ42 was performed according to Walsh and colleagues[34]. Briefly, large cultures of *Escherichia coli* BL21 Gold (DE3) were incubated at 37 °C with shaking, induced with 0.5 mM IPTG and harvested by centrifugation. Purification involved a series of sonication and centrifugation steps followed by resuspension of inclusion bodies in 8M urea. Anion exchange chromatography using diethylaminoethyl cellulose beads was performed and protein eluted with increasing concentrations of NaCl (up to 150 mM). Aβ42 elution was assessed by electrophoresis, using by Novex™ 10–20% Tricine SDS-PAGE gels. Fractions containing elevated levels of Aβ42 were combined and filtered through a 30 kDa molecular mass cut-off centrifugal filtration device (Amicon® Ultra-15). The combined sample was concentrated using a 3 kDa molecular mass cut-off centrifugal filtration device (Amicon® Ultra-15), snap frozen, lyophilised and stored at –20 °C. [15]N Aβ42 expression was performed according to Habchi and colleagues[18], with purification as above.

Size exclusion chromatography. An aliquot of lyophilised Aβ42 protein was solubilised in 6 M GuHCl until no large aggregates were visible. The sample was centrifuged at ~3300 x *g* for 3 min, bath

sonicated for 5 min in ice-water and centrifuged again at 16000 x g for 10 min at 4 °C. 500 µL of sample
was drawn up carefully with a syringe and loaded onto a Superdex™ 75 Increase 10/300 GL size
exclusion column running on an ÄKTA FPLC system. Protein was eluted at 0.5 mL/min in 50 mM
ammonium acetate pH 8.5. Fractions (0.5 mL) were collected in Protein LoBind Eppendorf tubes and
immediately placed on ice. Fractions were analysed for Aβ42 content using SDS-PAGE. Concentration
was measured via spectrophotometer (ε275 = 1450). Fractions containing Aβ42 were combined and
either used immediately or snap frozen and lyophilised.

Thioflavin T assays. Protein samples (Aβ42) were prepared in triplicate in non-binding 96-well half
area plates (Corning 3881) in 25 mM or 50 mM NaH_2PO_4 pH 7.4, 40 µM ThT to a final volume of 80 µL.
Plate preparation was performed at 4 °C to minimise aggregation. Samples were incubated at 37 °C in
a POLARstar Omega microplate reader (BMG Labtech), with excitation at 440 nm and the fluorescence
emission recorded at 480 nm under quiescent conditions.

Seeding. Seeds were prepared by incubating 10 µM Aβ42 in 50 mM NaH_2PO_4, pH 7.4 at 37 °C for 5 h.
Protein solution was subject to 6 x 10 s sonication with 20 s rest on ice between cycles. The seeds were
maintained on ice and used immediately.

Enhanced BCA assay details. Kinetic assays were run to completion as judged by the plateau in ThT
fluorescence (~5 h). Contents of wells were removed and centrifuged at 16000 x g for 10 min to pellet
fibrils and aggregates. Protein concentration was determined using a Pierce® Bicinchoninic Acid Protein
Assay Kit according to manufacturer's enhanced protocol. Monomeric Aβ42 and compounds alone
were used as controls.

TEM. Samples were prepared on carbon-coated copper grids (200 mesh) with formvar support film
(ProSciTech Pty Ltd.) that had been exposed to UV light (230 nm) for 10 min before sample application
and stained with 2% aqueous uranyl acetate solution. Samples were examined with a FEI Tecnai T12
electron microscope operating at 120 kV and images captured with a Veleta CCD camera and
processed using RADIUS 2.0 imaging software (EMSIS GmbH).

Preparation of Aβ42 for cell assay and NMR. Lyophilised, purified protein was resuspended in HFIP
at 1 mg/mL and incubated for 30 min at room temperature. The protein was snap frozen, lyophilised,
and stored at –80 °C until use.

Cell Culture and Differentiation. SH-SY5Y cells were cultured in in Dulbecco's Modified Eagle
Medium: Nutrient Mixture F-12 (DMEM/F12; Invitrogen, Carlsbad, CA, USA) supplemented with 10%
fetal bovine serum (FBS; Gibco, Uxbridge, UK) at 37 °C and 5% CO_2. Cells were fed every 2-3 days
and passaged using Trypsin-EDTA (0.25%). Cells were detached using Trypsin-EDTA, centrifuged (5
min, 125 x g), counted and then plated in a Cell-bind 96-well plate at a density of 0.5 x 10^4 cells/well in
10 µM all-trans retinoic acid (RA; Sigma-Aldrich) in DMEM/F12 with 10% FBS. Media was replaced
daily for 4 days (RA is light sensitive, so all steps involving RA were completed in the dark). After a 4-
day incubation with RA, media was switched to DMEM/F12 (no serum) with 50 ng/mL BDNF (Sigma-
Aldrich) for 3 days. After a total of 7 days of differentiation the cells were used for the experiments.

Aβ42 preparation, compound treatment and CellTiter-Blue® viability assay. 24 h prior to the cells
finishing differentiation, an aliquot of Aβ42 was centrifuged at 16,000 x g for 10 min and dissolved in
PBS at 200 µM, then diluted to 20 µM in DMEM/F12 with 10% FBS. The solution was stored in the
incubator (37 °C, 5% CO_2) in a low-bind Eppendorf to promote Aβ42 fibrillization. After cell
differentiation was complete, for the Aβ42 monomeric preparation, a second aliquot of Aβ42 was
centrifuged at 16,000 x g for 10 min, dissolved in PBS at 200 µM and immediately diluted to a final
concentration of 20 µM in DMEM/F12 (with 10% FBS). Experimental compounds (10 µM, 0.1%) or
DMSO control were added to the fibrillar or monomeric Aβ42. Differentiation media was aspirated,
100 µL of each condition added to the cells and incubated for 24 h (37 °C, 5% CO_2). In the monomeric
preparation, the time between dissolving the Aβ42 and applying onto the cells was minimised (less than
2 min total). After 24 h, CellTiter-Blue® (CTB; Promega G8080) (20 µL/well) was added and incubated

for 4 h. The fluorescent emission at 590 nm was measured using a POLARstar Omega microplate-
reader (BMG Labtech); viable cells metabolise this agent and reduce it to a fluorescent compound.

Statistical Analysis. One-way ANOVA (analysis of variance) followed by a Bonferroni's multiple
comparisons test was used to determine the differences between mean CTB values of differentiated
SH-SY5Y cells treated with Aβ42 and/or test compounds relative to vehicle control. Significance
threshold was defined as $p < 0.05$.

Kinetic analysis. The kinetic data obtained in ThT assays were analysed using the online fitting
software AmyloFit, following the cited protocol[43]. The model 'secondary nucleation dominated,
unseeded' was used, with the reaction orders set to 2 as typical for Aβ42 and under the constraint of
allowing deviations from the unperturbed kinetics in only one parameter at a time to investigate which
microscopic step or which species of the aggregation mixture was most likely affected by the inhibitors.
Repeats of the experiment were analysed separately and combined at the end to estimate uncertainty.

K_d determination. To determine the value of the dissociation constant, K_d, between free monomer and
inhibitor, a one-to-one binding was assumed and the free monomer concentration, m, expressed as a
function of the total inhibitor concentration, I_{tot}, the total monomer concentration, m_{tot}, and K_d,

$$m = \frac{-I_{tot} + m_{tot} - K_d + \sqrt{(I_{tot} - m_{tot} + K_d)^2 + 4\,K_d\,m_{tot}}}{2}.$$

The free monomer concentration was estimated as the effective monomer concentration in the kinetic
assays or the relative one-dimensional 1H NMR signal, respectively. Likelihoods were calculated using
grid approximations, under the assumption of Gaussian noise.

NMR / K_d Analysis. All NMR data were collected at 4 °C using unlabelled 10 μM Aβ42 or ^{15}N-labelled
Aβ42 in 20 mM NaH_2PO_4 pH 7.0, 10% D_2O, 1 mM DSS. Individual titrations with **C16**, **C17**, **C18** or
EGCG were carried out by incubating unlabelled 10 μM Aβ42 with increasing concentrations of the
compounds, ranging from 0 to 50 μM. 9 titration points were recorded for each compound. The NMR
signals from one-dimensional ^{1}H spectra were used to monitor the binding. Each spectrum was
recorded using 512 scans. A ^{1}H–^{15}N heteronuclear single quantum coherence (HSQC) spectrum was
also recorded for ^{15}N-labelled Aβ42 in the presence of 10 μM **C16**. The HSQC spectrum was recorded
overnight, for 240 scans, with a spectral width of 25 ppm (^{15}N offset at 117.5 ppm). All spectra were
obtained using a 18.8 T (800 MHz ^{1}H frequency) Advance III HD Bruker spectrometer equipped with a
cryoprobe.

Acknowledgements

The work was funded by Australian Research Council Discovery Project grant (DP200102463) to MS,
AK and PJR. SRB, JSPA and MAS were supported by Research Training Program Scholarships from
the Australian Government (Department of Education, Skills and Employment). MK, MAS and ELW's
research is supported by the National Health and Medical Research Council of Australia (NHMRC;
APP1132524). MK is an NHMRC Principal Research Fellow (APP1154692). MRZ acknowledges
support from the Herchel Smith Fund. The authors thank Elva (Meng) Shi and Dr Malcolm Spain for
assistance with synthesis and characterisation of compounds and Dr Bill Bubb for comments on the
manuscript. The authors acknowledge the facilities and the scientific and technical assistance of staff
within the Sydney Analytical and Sydney Microscopy & Microanalysis Core Research Facilities at the
University of Sydney. We acknowledge and pay respect to the Gadigal people of the Eora Nation, the
traditional owners of the land on which we research, teach, and collaborate at the University of Sydney.

References
477

1.   I. Benilova, E. Karran and B. De Strooper, *Nature neuroscience*, 2012, **15**, 349-357.

2.   S. J. C. Lee, E. Nam, H. J. Lee, M. G. Savelieff and M. H. Lim, *Chemical Society Reviews*, 2017,
46, 310-323.

3.   D. M. Walsh, I. Klyubin, J. V. Fadeeva, W. K. Cullen, R. Anwyl, M. S. Wolfe, M. J. Rowan and D.
J. Selkoe, *Nature*, 2002, **416**, 535-539.

4.   F. Chiti and C. M. Dobson, *Nature chemical biology*, 2009, **5**, 15-22.

5.   J. Fantini, H. Chahinian and N. Yahi, *Protein Science*, 2020, **29**, 1748-1759.

6.   P. Jadiya, D. W. Kolmetzky, D. Tomar, A. Di Meco, A. A. Lombardi, J. P. Lambert, T. S. Luongo,
M. H. Ludtmann, D. Praticò and J. W. Elrod, *Nature communications*, 2019, **10**, 3885.

7.   I. F. Tsigelny, Y. Sharikov, V. L. Kouznetsova, J. P. Greenberg, W. Wrasidlo, T. Gonzalez, P.
Desplats, S. E. Michael, M. Trejo-Morales, C. R. Overk and E. Masliah, *Journal of Alzheimer's*
disease : JAD, 2014, **39**, 583-600.

8.   I. F. Tsigelny, Y. Sharikov, W. Wrasidlo, T. Gonzalez, P. A. Desplats, L. Crews, B. Spencer and
E. Masliah, *The FEBS journal*, 2012, **279**, 1000-1013.

9.   B. Winner, R. Jappelli, S. K. Maji, P. A. Desplats, L. Boyer, S. Aigner, C. Hetzer, T. Loher, M.
Vilar, S. Campioni, C. Tzitzilonis, A. Soragni, S. Jessberger, H. Mira, A. Consiglio, E. Pham, E.
Masliah, F. H. Gage and R. Riek, *Proceedings of the National Academy of Sciences*, 2011, **108**,
4194-4199.

10.   B. Zott, M. M. Simon, W. Hong, F. Unger, H.-J. Chen-Engerer, M. P. Frosch, B. Sakmann, D. M.
Walsh and A. Konnerth, *Science*, 2019, **365**, 559-565.

11.   M. Serra-Batiste, M. Ninot-Pedrosa, M. Bayoumi, M. Gairí, G. Maglia and N. Carulla,
Proceedings of the National Academy of Sciences, 2016, **113**, 10866-10871.

12.   T. C. T. Michaels, A. Šarić, S. Curk, K. Bernfur, P. Arosio, G. Meisl, A. J. Dear, S. I. A. Cohen, C.
M. Dobson, M. Vendruscolo, S. Linse and T. P. J. Knowles, *Nature Chemistry*, 2020, **12**, 445-
451.

13.   S. I. A. Cohen, S. Linse, L. M. Luheshi, E. Hellstrand, D. A. White, L. Rajah, D. E. Otzen, M.
Vendruscolo, C. M. Dobson and T. P. J. Knowles, *Proceedings of the National Academy of*
Sciences, 2013, **110**, 9758-9763.

14.   J. E. Gillam and C. E. MacPhee, *Journal of physics. Condensed matter : an Institute of Physics*
journal, 2013, **25**, 373101.

15.   S. Linse, *Biophys Rev*, 2017, **9**, 329-338.

16.   T. C. T. Michaels, A. Šarić, G. Meisl, G. T. Heller, S. Curk, P. Arosio, S. Linse, C. M. Dobson, M.
Vendruscolo and T. P. J. Knowles, *Proceedings of the National Academy of Sciences*, 2020,
DOI: 10.1073/pnas.2006684117, 202006684.

17.   J. Habchi, P. Arosio, M. Perni, A. R. Costa, M. Yagi-Utsumi, P. Joshi, S. Chia, S. I. A. Cohen, M.
B. D. Müller, S. Linse, E. A. A. Nollen, C. M. Dobson, T. P. J. Knowles and M. Vendruscolo,
Science Advances, 2016, **2**, e1501244.

18.   J. Habchi, S. Chia, R. Limbocker, B. Mannini, M. Ahn, M. Perni, O. Hansson, P. Arosio, J. R.
Kumita, P. K. Challa, S. I. A. Cohen, S. Linse, C. M. Dobson, T. P. J. Knowles and M.
Vendruscolo, *Proceedings of the National Academy of Sciences*, 2017, **114**, E200-E208.

19.   R. Limbocker, S. Chia, F. S. Ruggeri, M. Perni, R. Cascella, G. T. Heller, G. Meisl, B. Mannini, J.
Habchi, T. C. T. Michaels, P. K. Challa, M. Ahn, S. T. Casford, N. Fernando, C. K. Xu, N. D. Kloss,
S. I. A. Cohen, J. R. Kumita, C. Cecchi, M. Zasloff, S. Linse, T. P. J. Knowles, F. Chiti, M.
Vendruscolo and C. M. Dobson, *Nature communications*, 2019, **10**, 225.

20.   M. Ettcheto, A. Cano, P. R. Manzine, O. Busquets, E. Verdaguer, R. D. Castro-Torres, M. L.
García, C. Beas-Zarate, J. Olloquequi, C. Auladell, J. Folch and A. Camins, *Molecular*
neurobiology, 2020, **57**, 1814-1827.

21.   R. Cataldi, S. Chia, K. Pisani, F. S. Ruggeri, C. K. Xu, T. Šneideris, M. Perni, S. Sarwat, P. Joshi, J.
R. Kumita, S. Linse, J. Habchi, T. P. J. Knowles, B. Mannini, C. M. Dobson and M. Vendruscolo,
Commun Biol, 2021, **4**, 19-19.

22. G. T. Heller, F. A. Aprile, T. C. T. Michaels, R. Limbocker, M. Perni, F. S. Ruggeri, B. Mannini, T.
Löhr, M. Bonomi, C. Camilloni, A. De Simone, I. C. Felli, R. Pierattelli, T. P. J. Knowles, C. M.
Dobson and M. Vendruscolo, *Science Advances*, 2020, **6**, eabb5924.
23. S. I. A. Cohen, P. Arosio, J. Presto, F. R. Kurudenkandy, H. Biverstål, L. Dolfe, C. Dunning, X.
Yang, B. Frohm, M. Vendruscolo, J. Johansson, C. M. Dobson, A. Fisahn, T. P. J. Knowles and
S. Linse, *Nature Structural & Molecular Biology*, 2015, **22**, 207-213.
24. A. Munke, J. Persson, T. Weiffert, E. De Genst, G. Meisl, P. Arosio, A. Carnerup, C. M.
Dobson, M. Vendruscolo, T. P. J. Knowles and S. Linse, *Proceedings of the National Academy*
of Sciences, 2017, **114**, 6444.
25. H. Liu, L. Yu, X. Dong and Y. Sun, *Journal of Colloid and Interface Science*, 2017, **491**, 305-312.
26. F. L. Palhano, J. Lee, N. P. Grimster and J. W. Kelly, *Journal of the American Chemical Society*,
2013, **135**, 7503-7510.
27. L. M. Young, J. C. Saunders, R. A. Mahood, C. H. Revill, R. J. Foster, L.-H. Tu, D. P. Raleigh, S. E.
Radford and A. E. Ashcroft, *Nature Chemistry*, 2015, **7**, 73-81.
28. L. Zhu, Y. Song, P.-N. Cheng and J. S. Moore, *Journal of the American Chemical Society*, 2015,
137, 8062-8068.
29. E. Tamanini, K. Flavin, M. Motevalli, S. Piperno, L. A. Gheber, M. H. Todd and M. Watkinson,
Inorganic Chemistry, 2010, **49**, 3789-3800.
30. J. K.-H. Wong, S. Ast, M. Yu, R. Flehr, A. J. Counsell, P. Turner, P. Crisologo, M. H. Todd and P.
J. Rutledge, *ChemistryOpen*, 2016, **5**, 375-385.
31. M. Yu, T. M. Ryan, S. Ellis, A. I. Bush, J. A. Triccas, P. J. Rutledge and M. H. Todd, *Metallomics*,
2014, **6**, 1931-1940.
32. S. L. Risacher, N. Fandos, J. Romero, I. Sherriff, P. Pesini, A. J. Saykin and L. G. Apostolova,
Alzheimer's & dementia (Amsterdam, Netherlands), 2019, **11**, 510-519.
33. S. Linse, *Methods in molecular biology (Clifton, N.J.)*, 2020, **2141**, 731-754.
34. D. M. Walsh, E. Thulin, A. M. Minogue, N. Gustavsson, E. Pang, D. B. Teplow and S. Linse, *The*
FEBS journal, 2009, **276**, 1266-1281.
35. M. Yu, G. Nagalingam, S. Ellis, E. Martinez, V. Sintchenko, M. Spain, P. J. Rutledge, M. H. Todd
and J. A. Triccas, *Journal of Medicinal Chemistry*, 2016, **59**, 5917-5921.
36. E. House, J. Collingwood, A. Khan, O. Korchazkina, G. Berthon and C. Exley, *Journal of*
Alzheimer's disease : JAD, 2004, **6**, 291-301.
37. M.-C. Lee, W.-C. Yu, Y.-H. Shih, C.-Y. Chen, Z.-H. Guo, S.-J. Huang, J. C. C. Chan and Y.-R. Chen,
Scientific Reports, 2018, **8**, 4772.
38. M. Mold, L. Ouro-Gnao, B. M. Wieckowski and C. Exley, *Scientific Reports*, 2013, **3**, 1256.
39. C. Haass and D. J. Selkoe, *Nature Reviews Molecular Cell Biology*, 2007, **8**, 101-112.
40. J. Krishtal, O. Bragina, K. Metsla, P. Palumaa and V. Tõugu, *PLOS ONE*, 2017, **12**, e0186636.
41. S. I. Cohen, M. Vendruscolo, M. E. Welland, C. M. Dobson, E. M. Terentjev and T. P. Knowles,
The Journal of chemical physics, 2011, **135**, 065105.
42. T. P. J. Knowles, C. A. Waudby, G. L. Devlin, S. I. A. Cohen, A. Aguzzi, M. Vendruscolo, E. M.
Terentjev, M. E. Welland and C. M. Dobson, *Science*, 2009, **326**, 1533-1537.
43. G. Meisl, J. B. Kirkegaard, P. Arosio, T. C. Michaels, M. Vendruscolo, C. M. Dobson, S. Linse
and T. P. Knowles, *Nature protocols*, 2016, **11**, 252-272.
44. J. M. Lopez del Amo, U. Fink, M. Dasari, G. Grelle, E. E. Wanker, J. Bieschke and B. Reif,
Journal of molecular biology, 2012, **421**, 517-524.
45. J. Roche, Y. Shen, J. H. Lee, J. Ying and A. Bax, *Biochemistry*, 2016, **55**, 762-775.
46. J. R. Horsley, B. Jovcevski, K. L. Wegener, J. Yu, T. L. Pukala and A. D. Abell, *The Biochemical*
journal, 2020, **477**, 2039-2054.
47. J. Bieschke, J. Russ, R. P. Friedrich, D. E. Ehrnhoefer, H. Wobst, K. Neugebauer and E. E.
Wanker, *Proceedings of the National Academy of Sciences*, 2010, **107**, 7710-7715.
48. D. E. Ehrnhoefer, J. Bieschke, A. Boeddrich, M. Herbst, L. Masino, R. Lurz, S. Engemann, A.
Pastore and E. E. Wanker, *Nature Structural & Molecular Biology*, 2008, **15**, 558-566.

49.   M. Spain, J. K. H. Wong, G. Nagalingam, J. M. Batten, E. Hortle, S. H. Oehlers, X. F. Jiang, H. E.
Murage, J. T. Orford, P. Crisologo, J. A. Triccas, P. J. Rutledge and M. H. Todd, *Journal of*
Medicinal Chemistry, 2018, **61**, 3595-3608.
50.   M. Yu, Q. Yu, P. J. Rutledge and M. H. Todd, *ChemBioChem*, 2013, **14**, 224-229.