Revisiting the notion of deleterious sweeps

Parul Johri[1,‡], Brian Charlesworth[2,‡], Emma K. Howell[1], Michael Lynch[1,3,*], and Jeffrey D. Jensen[1,*]

[1]School of Life Sciences, Arizona State University, Tempe, AZ 85287, United States
[2]Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, EH9 3FL, United Kingdom
[3]Center for Mechanisms of Evolution, The Biodesign Institute, Arizona State University, Tempe, AZ, United States

[‡] authors contributed equally
[*] co-corresponding authors

keywords: population genetics, selective sweeps, genetic hitchhiking

ABSTRACT

It has previously been shown that, conditional on its fixation, the time to fixation of a semi-dominant deleterious autosomal mutation in a randomly mating population is the same as that of an advantageous mutation. This result implies that deleterious mutations may generate selective sweep effects. Although their fixation probabilities greatly differ, the much larger input of deleterious relative to beneficial mutations suggests that this phenomenon could be important. We here examine how the fixation of mildly deleterious mutations affects levels and patterns of polymorphism at linked sites, and how this class of sites may contribute to divergence between-populations and species. We find that, while deleterious sweeps are unlikely to represent a significant proportion of outliers in polymorphism-based genomic scans within populations, minor shifts in the frequencies of deleterious mutations can influence the proportions of private variants and the value of F_{ST} after a recent population split. As sites subject to deleterious mutations are necessarily found in functional genomic regions, interpretations in terms of recurrent positive selection may require reconsideration.

INTRODUCTION

Among the most important results in theoretical population genetics, now nearly a century old, are the fixation probabilities of new beneficial and deleterious mutations, which were obtained by Fisher (1922, 1930), Haldane (1927), and Wright (1931), using different approaches. Their results were later generalized by Kimura (1957, 1962, 1964), using the backward diffusion equation. A somewhat lesser known result concerns the trajectories of these selected mutations. Specifically, Maruyama & Kimura (1974) found that, conditional on fixation, the time that a beneficial autosomal mutation with selection coefficient $+s$ and dominance coefficient h spends in a given interval of allele frequency in a randomly mating population is the same as that for a deleterious mutation with selection coefficient $-s$ and dominance coefficient $1 - h$, provided that the conditions for the validity of the diffusion equation approximation hold. Thus, given that the effects of selective sweeps on variability at linked neutral sites are related to their speed of transit through a population (Maynard Smith & Haigh 1974; Stephan 2019), the fixation of a deleterious mutation by genetic drift can generate a similar selective sweep effect to that caused by the fixation of a beneficial mutation, for mutations with the same magnitude of selection coefficient. Moreover, Tajima (1990) demonstrated that, on average, there is a ~42% mean reduction in diversity at a site where a neutral mutation has recently become fixed by genetic drift. While the mean time to fixation for this class of mutation is well known to be $4N_e$ generations (Kimura & Ohta 1969); this is of course associated with a wide variance of approximately $4.64N_e^2$ (Kimura 1970), such that neutral mutations may also fix relatively rapidly (in less than N_e generations) and generate an appreciable but highly localized sweep effect (see Tables 1 and 2 of Tajima 1990).

Naturally, the probabilities of deleterious and beneficial fixations differ strongly. However, given that the input of deleterious mutations is much higher than the input of beneficial mutations

each generation (see reviews by Eyre-Walker & Keightley 2007; Bank *et al.* 2014a), the potential

contribution of such deleterious sweeps to levels and patterns of nucleotide variation, as well as

divergence between populations and species, remains an important open question (Charlesworth

2020a). An alternative way of viewing this issue, as discussed by Gillespie (1994) and

Charlesworth & Eyre-Walker (2007), is that under a model of constant selection and reversible

mutation between two alternative nucleotide variants, statistical equilibrium with respect to the

frequencies of sites fixed for the alternatives implies equal rates of beneficial and deleterious

substitutions per unit time. It is important to note, however, that only deleterious mutations with

selection coefficients on the order of the reciprocal of the population size have significant

probabilities of fixation (Fisher 1930; Kimura 1964), implying that the substitutions concerned

involve only very weakly selected mutations. The effects on diversity statistics of sweeps of very

weakly selected mutations, including those of deleterious mutations, appears to have been

investigated previously only by Mafessoni & Lachmann (2015).

A starting point for investigating this problem is the distribution of fitness effects of new

mutations (the DFE). There is substantial evidence from both empirical and experimental studies

that the DFE of new mutations is bimodal - consisting of a strongly deleterious mode, and a weakly

deleterious / neutral mode that may contain a beneficial tail under certain conditions (*e.g.,* Crow

1993; Lynch *et al.* 1999; Sanjuán 2010; Jacquier *et al.* 2013; Bank *et al.* 2014b). While the

calculations and simulations presented below represent a general approach to addressing this topic,

we have necessarily chosen a specific DFE realization and species for illustration. Specifically,

Johri *et al.* (2020) recently presented an approximate Bayesian (ABC) approach, which represents

the first joint-estimator of the DFE shape together with population history that corrects for the

effects of background selection (BGS; Charlesworth *et al.* 1993). They estimated that a substantial

proportion of new mutations in coding regions have mildly deleterious effects on fitness,

emphasizing the importance of further understanding the consequences of such mutations in

dictating observed polymorphism and divergence. Furthermore, they found it unnecessary to

invoke a beneficial mutational class in order to fit the data from the African population of

Drosophila melanogaster that they considered.

This analysis provides a basis for exploring the possible implications of sweeps of

deleterious mutations, a topic that has been rather neglected. Here, we re-examine this question,

considering both single and recurrent substitution models, which we use to examine the possibility

that genomic scans for positive selection may in fact also be identifying deleterious fixations, when

based on: a) levels and patterns of variation; b) population differentiation; and c) species-level

divergence. Our results suggest that, while this phenomenon is unlikely to be a major factor in

polymorphism-based scans, it may be a serious confounder in among-population-based analyses.

METHODS

<u>Analytical calculations</u>

For convenience, we assume a Wright-Fisher population of N randomly mating individuals

throughout the analytical section. The selection coefficient for homozygotes is denoted by s, with

$s = s_a$ or $s = -s_d$ (s_a and $s_d > 0$), representing selection for and against homozygotes, respectively.

In particular, the distribution of fitness effects (DFE) for semi-dominant deleterious mutations is

assumed to be discrete, with four fixed classes of mutations given by $0 < 2Ns_d \leq 1$, $1 < Ns_d \leq 10$,

$10 < 2Ns_d \leq 100$ and $100 < 2Ns_d \leq 2N$, where mutations are assumed to follow a uniform

distribution within each class. These assumptions concerning the distribution of s_d were made in

order to simplify integration over the DFE (Johri *et al*. 2020).

Probability of fixation: The fixation probability (P_{fix}) of a new semi-dominant mutation with an

initial frequency of $1/2N$ in a Wright-Fisher population of size N was calculated using Equation

10 of Kimura (1962):

$$P_{fix} = \frac{1 - e^{s}}{1 - e^{2Ns}} \approx \frac{s}{e^{2Ns} - 1} \qquad (1)$$

125   Note that this equation assumes demographic equilibrium and independence between the selected

126   sites, as well as $|s| \ll 1$.

127   *Contributions to divergence*: Under the DFE model described above, the number of fixations N_{fix}

128   (i) expected per generation per site for a given DFE class i of deleterious mutations is given by

129   the following expression:

$$N_{fix}(i) = 2N\mu \times f_i \times \frac{1}{s_{d2} - s_{d1}} \int_{s_{d1}}^{s_{d2}} P_{fix}(s)ds \qquad (2)$$

where μ is the total mutation rate per site per generation, f_i represents the proportion of new

mutations belonging to the i^{th} DFE class and s_{d2} and s_{d1} represent the upper and lower bounds to

the DFE class, respectively. The integral in equation (2) was evaluated analytically by means of

an infinite series representation (derivation provided in Appendix 1), which was validated using

the "integrate" function in R (R Core Team 2018).

Waiting and fixation times: The waiting time (t_w) between fixations under a Poisson process was

calculated as follows:

$$t_w = \frac{1}{L \times N_{fix}} \qquad (3)$$

where L represents the number of functional sites under consideration.

The expected time to fixation of a semi-dominant mutation was calculated by numerically

integrating equation (17) of Kimura & Ohta (1969), using Simpson's rule (Atkinson 1989).

Reduction in diversity due to a single sweep: The expected reduction in pairwise nucleotide site

diversity at the end of a sweep ($-\Delta\pi$), relative to the diversity in the absence of selection, was

calculated using equation (14a) of Charlesworth (2020b) for a non-zero rate of recombination

(omitting a factor that describes the effect of background selection):

$$-\Delta\pi = (1 - P_r) - T_s - P_{rs}T_d - (P_r - P_{rs})T_r \qquad (4)$$

where T_s is the mean coalescent time (in units of N) for a pair of alleles sampled at the end of a

sweep, T_d is the duration of the deterministic phase of the sweep (*i.e.*, excluding the initial and

final stochastic phases) in units of N, T_r is the mean time to a recombination event during the sweep,

conditioned on the occurrence of a recombination event; P_r is the probability of at least one

recombination event transferring a sampled allele onto the wild-type background during the sweep,

and P_{rs} is the probability that there is only a single such recombination event. For large values of

the ratio of the rate of recombination between the neutral and selected site (r) to the magnitude of

the selection coefficient, this expression can become negative, in which case it is reset to zero.

We also used simulations based on equations (27) of Tajima (1990), which provide

recursion equations for the expectation of the pairwise diversity at a neutral locus linked to a

selected locus, conditional on a given trajectory of allele frequency change at the selected locus,

as described by Charlesworth (2020b). These equations require only the validity of the diffusion

equation approximation. Binomial sampling of allele frequencies post-selection in each generation

was used to generate the trajectories of change at the selected locus, using the standard selection

equation for a single locus to calculate the deterministic change in allele frequency each

generation. Application of the recursion equations to a trajectory of allele-frequency change

simulated in this way gives one realization of $\Delta\pi$, and the overall expected value of $\Delta\pi$ can be

obtained from the mean of the simulated values over a large number of replicates. It was found

that 1000 replicates gave very accurate estimates of $\Delta\pi$, with ratios of the standard errors to the

means of < 5% for the parameter sets used here.

Population genetic parameters used for analytical calculations: The parameters for the

calculations were chosen to match those estimated from *D. melanogaster* populations, estimated

from exonic sites. Mutations occurred at a rate μ per basepair, and were assumed to be a mixture

of neutral, nearly neutral and weakly deleterious mutations. For this analysis; $\mu = 3 \times 10^{-9}$ per

basepair (bp) per generation (Keightley *et al.* 2009, 2014). The sex-averaged rate of crossing over

per bp (r_c) was assumed to be equal to 10^{-8} per generation (Fiston-Lavier *et al.* 2010), the effective

population size was 10^6 (Arguello *et al*. 2019; Johri *et al*. 2020), and 10 generations per year were

assumed. Given estimates of neutral divergence between *D. melanogaster* and *D. simulans*, this

means that $t \sim 21.3 \times 10^6$ generations elapsed since their common ancestor (Li *et al*. 1999; Halligan

& Keightley 2006), corresponding to 2.13 M years.

Because non-crossover associated gene conversion is an important source of recombination

between closely linked sites in *Drosophila*, it was assumed to occur uniformly across the genome,

independently of local differences in the rate of crossing over, as indicated by the data on *D,*

melanogaster (Comeron *et al*. 2012; Miller *et al*. 2016). The sex-averaged rate of initiation per bp

of conversion events was $r_g = 10^{-8}$ per bp per generation, and there was an exponential distribution

of tract length with a mean of $d_g = 440$ bp (Comeron *et al*. 2012; Miller *et al*. 2016). The net rate

of recombination between sites separated by z bp, $r(z)$, is the sum of the contributions from

crossing over and non-crossover gene conversion, given by the formula of Frisse $et\ al.$ (2001):

$$r(z) = r_c z + 2 r_g d_g [1 - \exp\left(-\frac{z}{d_g}\right)] \qquad (5)$$

For some of the results presented below, only the net rate of recombination r, or its value

scaled by the effective population size, $\rho = 2Nr$, was used. In the presence of gene conversion

with the parameters described above, the corresponding value of z can then be obtained by

Newton-Raphson iteration of the equation $r(z) - r = 0$, assuming that the rate of gene conversion

does not vary with the rate of crossing over,

<u>Simulations</u>

Simulating individual fixations in a single population: SLiM 3.3.1 (Haller & Messer 2019) was

used to simulate a genomic element of length 10 kb. For the single sweep model, in order to

quantify the hitchhiking effect of a single fixation, selection only acted on a single site in the

middle of the region, with all other sites evolving neutrally. Simulations were performed for five

different values of the scaled selection coefficient: $2Ns = 0, -1, +1, -5,$ and $+5$. Population genetic

parameters resembling those of $D.\ melanogaster$ were utilized (as defined above) for illustrative

purposes. In order to perform simulations efficiently, population size was scaled down by a factor

of 100 while the mutation and recombination rates were correspondingly scaled up by the same

factor. Simulations were run for a burn-in period of 10^5 generations ($10N_{sim}$) after which a

mutation with a scaled selection coefficient of $2Ns$ was introduced at the selected site. Simulations

in which the introduced mutation reached fixation were retained for analysis. Fifty diploid

individuals were sampled at the completion of the simulations, and population genetic summary

statistics were calculated using *Pylibseq* (Thornton 2003).

Simulating multiple populations and F_{ST}-based analyses: Based on a recently inferred

demographic history of African and European populations of *D. melanogaster* using populations

sampled from Beijing, the Netherlands, and Zimbabwe (Arguello *et al*. 2019), an ancestral

population of size of 1.95×10^6 was simulated, which split into two populations (6.62×10^4

generations ago) of constant size: 3.91×10^6 and 4.73×10^5, representing African and European

populations, respectively. Note that, although Arguello *et al*. inferred recent growth in both

populations, we have assumed constant sizes in order to avoid confounding effects of such growth

on the fixation probabilities of mutations, and we also assumed no migration between the two

populations. For the purpose of F_{ST} analyses, a functional region of size 10 kb was simulated, in

which mutations had selection coefficients given by the DFE inferred in Johri *et al.* (2020).

Specifically, the DFE was given by a discrete distribution of four fixed bins with 24.7% of

mutations belonging to the effectively neutral class (f_0: $0 \leq 2Ns_d <1$), 49.4% to the weakly

deleterious class (f_1: $1 \leq 2Ns_d < 10$), 3.9% to the moderately deleterious class (f_2: $10 \leq 2Ns_d <$

100), and 21.9% to the strongly deleterious class of mutations (f_3: $100 < 2Ns_d \leq 2N$).

In this two-population framework, four separate scenarios were tested: 1) the DFE

remained scaled to the ancestral population size (making selection effectively weaker in the

smaller derived (European) population); 2) the DFE was rescaled after the population split with

respect to subpopulation-specific sizes (*i.e.*, both populations experienced equal proportions of

mutations belonging to each DFE class as defined in terms of $2Ns_d$), such that selection was

equally strong in both populations - this is an arbitrary biological model that is simply chosen for

comparison, as one would naturally expect selection to be effectively weaker in the derived

population as in scenario 1; 3) in addition to this neutral and deleterious DFE, 1% of all mutations

were mildly beneficial with selective effects drawn uniformly from the interval $1 \leq 2Ns_a \leq 10$,

where s_a is the increase in fitness of the mutant homozygote; and 4) in addition to this neutral and

deleterious DFE, 1% of all mutations were strongly beneficial with $2Ns_a = 1000$. In scenarios 3

and 4, the deleterious and beneficial DFEs were scaled to the respective sizes of each population

post-split, so that the proportions of beneficial mutations were the same in the two populations. In

order to assess the role of population bottlenecks in generating neutral outliers, a fifth scenario was

simulated in which there was no selection and the demographic history was that inferred by Li &

Stephan (2006) - a model that involves a much larger size reduction than inferred in the Arguello

et al. (2019) model utilized above. The parameters of both demographic models are provided in

Supp Table 1.

Fifty diploid individuals were sampled from both populations in order to calculate F_{ST}. All

sites that would be considered polymorphic in the metapopulation were used to calculate F_{ST} (*i.e.*,

sites fixed either in one population or both populations (for different alleles) were also included in

F_{ST} calculations). F_{ST} was calculated in sliding windows across the genomic region for a) windows

containing a constant number of SNPs using the package PopGenome (Pfeifer *et al*. 2014) in R,

and b) for windows representing the same total number of bases using *Pylibseq* 0.2.3 (Thornton

2003). F_{ST} was calculated for both cases by the method of Hudson *et al*. (1992). F_{ST} was also

calculated individually for different mutation types (*i.e.*, for neutral, weakly deleterious and

beneficial mutations, by simply restricting the calculations to segregating sites of the specific

mutation type). Although there will be an upper bound to the F_{ST} values obtained in this way,

which is determined by the frequency of the most frequent allele in the metapopulation (Jakobsson

et al. 2013), the detection of outliers should not be affected by this procedure and should mimic

the scenario for natural populations of *Drosophila*.

Simulating multiple species, and divergence-based analyses: McDonald-Kreitman (MK) tests

(McDonald & Kreitman 1991) were performed to investigate the degree to which substitutions of

mildly deleterious mutations might affect the inference of divergence due to positive selection. A

population resembling a *D. melanogaster* African population was simulated under demographic

equilibrium. Ten independent replicates of a 10-kb protein-coding region were considered such

that every third position was neutral (representing synonymous sites) while all other sites

represented nonsynonymous positions. Nonsynonymous sites experienced purifying selection

given by a DFE that comprised only the non-neutral bins (*i.e.,* f_1, f_2, and f_3, in the same proportions

described above), while the neutral sites experienced mutations that belonged to class f_0. For

comparison, simulations were performed in which 10% or 20% of nonsynonymous mutations were

also neutral. In all cases, the simulation was run for $20N$ generations (where $N = 10^4$). The number

of segregating nonsynonymous (P_N) and synonymous (P_S) sites were estimated by sampling 50

diploid individuals from the population from the functional and neutral regions respectively. The

number of fixed substitutions occurring at the functional sites (D_N) and neutral sites (D_S) were

calculated post burn-in (*i.e.*, after $10N$ generations), and then rescaled to the number of generations

since the *D. melanogaster* ancestor.

In order to correct for mildly deleterious mutations segregating in populations, the

proportion of adaptive substitutions (α) was also inferred by implementing a variant of the test

referred to as the asymptotic MK test. Messer & Petrov (2013) suggested plotting the derived

allele frequency (x) of variants against α inferred using the number of segregating sites ($P(x)$) at

that derived allele frequency (*i.e.*, $\alpha(x) = \frac{P(x)_N}{P(x)_S} \times \frac{D_S}{D_N}$), and showed that the asymptote of this

curve would tend towards the true value of α. The asymptotic MK test was performed using a

web-based tool available at: http://benhaller.com/messerlab/asymptoticMK.html (Haller &

Messer 2017). For the purpose of the asymptotic MK test, values of P_N/P_S were binned with a

bin size of 0.05, and the curve-fitting (of $\alpha(x)$ with respect to x) was restricted to derived allele

frequencies between 0.1-1.0

RESULTS & DISCUSSION

<u>Theoretical expectations</u>

As has been long appreciated (Fisher 1930; Wright 1931), equation (1) implies that the probability

of fixation of a mutation in the strongly deleterious DFE classes is vanishingly small. However,

the weakly deleterious class may contribute substantially to divergence (Figure 1a). We derived

an analytical approximation for the probability of fixation for mutations with a DFE represented

as a combination of four non-overlapping uniform distributions (see Methods and Appendix 1). In

a Wright-Fisher population, the ratio of the mean probability of fixation of mutations with fitness

effects between $1 < 2Ns_d \leq 5$, relative to the probability of fixation of effectively neutral mutations

($0 < 2Ns_d \leq 1$), is 0.27 (note that the value of N is irrelevant if the diffusion approximation holds).

This ratio rapidly declines to 0.01 for mutations with fitness effects $5 < 2Ns_d \leq 10$ (Figure 1a).

Taking the recently estimated DFE for the Zambian *D. melanogaster* population (Johri *et*

al., 2020) as an example, a question of interest concerns the probabilities of fixation of different

classes of mutations and their contributions to population- and species-level divergence (Supp

Figure 1). In the absence of positive selection, the weakly deleterious class of mutations would be

expected to contribute 7.2% of the total divergence in exonic regions, while effectively neutral

mutations would be expected to contribute 92.8%. If we were to assume that approximately 50%

of all substitutions in *Drosophila* have been caused by positive selection (Eyre-Walker and

Keightley 2009; Campos *et al*. 2017), weakly deleterious mutations are still likely to have

contributed 3.5% of the total divergence in functional regions and possibly much more in regions

experiencing reduced selective constraints.

<u>Hitchhiking effects of deleterious sweeps: levels and patterns of variation</u>

We next considered the fixation times of these mutations contributing to divergence, as well as the

expected waiting time between fixations. Using population parameters relevant for *D.*

melanogaster (see Methods), the neutral and deleterious DFE of Johri *et al.* (2020), and assuming

that 60% of the *D. melanogaster* genome (of size 140 Mb) is functional, equation (3) shows that

the genome-wide waiting time between successive weakly deleterious fixations is ~ 83

generations. Hence, the ~2.13 million years estimated for the time since the *D. melanogaster* and

D. simulans split is expected to equate to many such fixations.

As expected, there is no significant difference (p = 0.16; Student's *t*-test) in fixation times

between the mildly beneficial (3.2N generations; SD: 1.5N) *vs* mildly deleterious mutations (2.9N

generations; SD: 1.1N) with the same fitness effects ($2Ns_a$=$2Ns_d$=5) (Supp Table 2). Importantly

however, the variance in fixation times of weakly selected mutations is extremely large, such that

the faster tails of the fixation time distributions for both $2Ns_d$ = 5 and $2Ns$ = 0 occupy ~N

generations, which corresponds to a sweep effect of the same size as the mean for $2Ns_a$ = 30

(Figure 1b). In other words, weakly deleterious and neutral fixations can match the sweep effects

of comparatively strongly selected beneficial fixations.

We evaluated the impact of these fixations on observed genomic variation in two ways.

The first corresponds to a model of a single sweep event, which is relevant to the literature on

detecting signatures of individual fixations, as done in genomic scans for positively selected loci

(*e.g.*, Harr *et al.* 2002; Glinka *et al.* 2003; Haddrill *et al.* 2005; Nielsen *et al.* 2005; see also Jensen

2009; Stephan 2019). Such scans operate under the assumption that the selective sweeps in

question have achieved fixation immediately prior to sampling, due to the rapid loss of signal as

the time since fixation increases (Kim & Stephan 2002; Przeworski 2002). The second corresponds

to a recurrent substitution model, which is relevant to the recurrent sweep literature that attempts

to quantify the effect of selective sweeps on genome-wide patterns of variability and their relation

to rates of recombination (*e.g.*, Wiehe & Stephan 1993; Kim 2006; Andolfatto 2007; Macpherson

et al. 2007; Jensen *et al.* 2008; Campos & Charlesworth 2019; Charlesworth 2020b; see review by

Sella *et al.* 2009).

The reduction of diversity resulting from the fixation of a mildly deleterious or beneficial

semi-dominant mutation with a very small selective effect is not expected to be substantially

different from that caused by the fixation of a selectively neutral mutation (Tajima 1990). The

widely-used approximation of Barton (2000) for the reduction in diversity relative to neutrality

caused by the sweep of a semi-dominant mutation, $\Delta\pi = (2Ns_a)^{(-4r/s)}$, suggests that, when $2Ns_a$

= 5, diversity will only be reduced by more than 20% in a region for which $r/s \leq 0.2$,

corresponding to approximately 20 bp for a typical *D. melanogaster* recombination rate of 3 x 10^{-}

^{8} per bp, including the contribution from gene conversion as given by equation (6), and assuming

an effective population size of $\geq 10^6$.

However, this formula assumes that fixations are so fast that swept alleles that have failed

to recombine onto a wild-type background experience no coalescent events during the duration of

the sweep (Hartfield & Bataillon 2020; Charlesworth 2020b), which is unlikely to be true for

weakly selected mutations. We therefore used equation (4) for analytical predictions (Figure 2,

Table 1, Supp Table 3), which is based on the results of Charlesworth (2020b). But this equation

is also likely to be inaccurate with weak selection, because it assumes that the trajectory of allele-

frequency change is close to that for the deterministic case, except for the initial and final stochastic

phases at the two extremes of allele frequencies. We therefore also used simulations based on

equations (27) of Tajima (1990) to predict sweep effects, as described in the Methods.

With the full simulations of a 10-kb region, in which we assumed a population size of 10^6

and mutation rate of 3×10^{-9} /site/generation, with $2Ns_a = 5$ the nucleotide diversity 10 bp ($\sim\rho =$

0.2) around the selected site (*i.e.*, 5 bp in both directions) was 0.0058 (SE: 0.0012), corresponding

to a reduction of 48% below the neutral value. For $2Ns = 10$, the 10-bp nucleotide diversity was

0.0083 (SE: 0.0017), corresponding to a reduction of 69%. The observed reduction in both cases

almost fully recovers to the expected level under neutrality within 500 bp ($\rho \approx 10$; Figure 2). A

similar pattern is seen in Table 1 and Supp Table 3, which shows both the analytical predictions

and those based on Tajima's equations, which agree surprisingly well except at the two highest

rates of recombination displayed. For comparison, the results of simulating neutral fixations are

also shown in Table 1 and Figure 2. The case with $Ns = 10$ is at the higher limit of what is likely

to be produced by sweeps of deleterious mutations, given that the ratio of the fixation probability

given by equation (1) to the neutral value of $1/(2N)$ is then approximately 0.0045, compared to

0.034 with $Ns = 5$.

Mafessoni & Lachmann (2015) showed that that fixations of weakly selected, highly

dominant favorable mutations - or highly recessive deleterious mutations - could reduce diversity

at linked sites by a *smaller* amount than fixations of neutral mutations, with a maximum effect

when $2Ns$ is approximately 2 (and $h = 1$ for favorable mutations, and 0 for deleterious ones). We

have confirmed this unexpected observation using Tajima algorithm simulations (Supp Table 4),

finding that it exists even for $2Ns = 5$ (Supp Figure 2; Supp Table 4). However, our use of $h =$

0.5 and $2Ns \geq 2.5$ means that this phenomenon is absent from the results presented here.

Because gene conversion is an important contributor to recombination between closely

linked sites (Miller *et al.* 2016), the effects of fixations in the presence of gene conversion are

restricted to a region that is about one-third of the distance in the absence of gene conversion

(Table 1). Thus, a greater than 20% reduction in nucleotide diversity for $2Ns = 5$ [$2Ns = 10$] was

observed up to $\rho = 0.8$ [$\rho = 1.6$], corresponding to 14 bp [27 bp] with gene conversion, and 40 bp

[80 bp] without gene conversion. This quite localized effect is similar for weakly beneficial,

weakly deleterious, and neutral mutations.

Given the relatively faster mean speed (3.1N generations) of fixation of weakly selected

semi-dominant mutations compared to the neutral expectation (4N generations), they should also

result in small distortions of the SFS at closely linked neutral sites. We observed a slight skew

towards rare variants (as measured by Tajima's D, Supp Table 5) restricted to ~50 base pairs from

the selected site immediately after fixation (for $2Ns_d = 2Ns_a = 5$). This highly localized

distortion of the SFS is probably too weak to play any important role in generating false positives

in genomic scans. Indeed, owing to the inherent stochasticity involved in the underlying processes,

such scans generally only have power to detect very strongly selected fixations (often requiring

values of $2Ns_a > 1000$ in order to observe appreciable true positive rates: Crisci *et al.* 2013).

Viewed in another way, given that the false-positive rates associated with genomic scans may often

be inflated well-above true-positive rates owing to the underlying demographic history of the

population (*e.g.,* Teshima *et al.* 2006; Thornton & Jensen 2007; Crisci *et al.* 2013; Harris *et al.*

2018), demography is probably a much stronger confounder than deleterious sweeps in

polymorphism-based scans.

Finally, using a recurrent fixation model, we considered the steady-state impact of weakly

deleterious sweeps (Figure 3). Here we used a model of a gene with five exons of 100 codons each,

with 70% of exonic mutations subject to selection, which were separated from each other by 100

bp introns, as described by Campos & Charlesworth (2019) and Charlesworth (2020b). Five

equally large classes of sites subject to deleterious mutations were modeled, with the lowest scaled

selection coefficient being Ns_d = 2.5 and the largest Ns_d = 10; as described above, a uniform

distribution of Ns_d values was assumed within each class. The theoretical predictions used the

method for predicting the effects of recurrent sweeps of Charlesworth (2020b), based on equation

(4) above. For consistency with the simulations of population divergence described below, the

population size was assumed to be 1.95 x 10^6, giving a neutral diversity of 0.0234 with the

assumed mutation rate of 3 x 10^{-9}. Substitution rates of deleterious nonsynonymous mutations

were calculated from equations (1) and (2). The expected number of nonsynonymous substitutions

per gene over 2*N* generations was 0.506, and the ratio of nonsynonymous to synonymous

substitutions was 0.0412, which is somewhat less than half the mean value for comparisons of *D.*

melanogaster and its close relatives (*e.g.*, Campos *et al.* 2017). The major source of these

substitutions is the class with Ns_d between 2.5 and 4.375, which accounts for 76% of all

substitutions, reflecting the fact that mutations in this class have the highest fixation probabilities.

The results for single sweeps shown in Table 1 suggested that the theoretical predictions

will tend to overestimate sweep effects, so that the results in Figure 3 must be treated with some

caution. With the standard *D. melanogaster* rate of crossing over per bp of 1 x 10^{-8}, the mean

reduction in nucleotide diversity at synonymous sites caused by deleterious sweeps is

approximately 5% with gene conversion and 8% in its absence (Figure 3). For the lowest rates of

crossing over and no gene conversion, average reductions can reach ~19%, suggesting that the

fixation of mildly deleterious mutations could play a significant role in organisms or genomic

regions with highly reduced rates of recombination. In this low recombination environment

however, selective interference would also become a factor - which would be expected to

accelerate the rate of deleterious fixations, while also making deleterious variants behave more

like neutral mutations. Regardless of these details, however, it seems unlikely that substitutions of

deleterious mutations will have more than a minor effect on average diversity in the *Drosophila*

genome overall, particularly compared with the effects of population history, background

selection, and sweeps of positively selected mutations. Furthermore, the findings of Mafessoni &

Lachmann (2015) suggest that strongly recessive deleterious mutations will have even smaller

effects than those studied here (see Supp. Figure 2).

<u>The contribution of deleterious mutations to population- and species-level divergence-based scans</u>

Given the potentially substantial contribution of the weakly deleterious class to observed fixations,

it is also of interest to consider their impact on divergence-based analyses (*e.g.,* methodology

related to d_N/d_S) and population differentiation-based scans (*e.g.,* methodology related to F_{ST}). We

examined properties related to inferring the proportion of substitutions fixed by positive selection

(α) by performing MK tests in the presence of mildly deleterious mutations. Specifically, we

simulated two scenarios: 1) 50% of mutations at nonsynonymous sites were weakly deleterious,

and 50% were neutral, and 2) nonsynonymous sites experienced deleterious mutations that

followed the DFE inferred by Johri *et al*. (2020).

Although, as expected, the presence of mildly deleterious mutations substantially increases

values of d_N/d_S (Supp Table 6) relative to stronger purifying selection (*i.e.*, larger proportions of

moderately and strongly deleterious mutations), it also leads to strongly negative values of α when

performing MK tests. This is due to the fact that mildly deleterious mutations often segregate in

the population at low frequency, inflating the total number of segregating nonsynonymous

polymorphisms (P_N) significantly (Supp Table 6). As such, the presence of mildly deleterious

mutations can result in negative values of α in the absence of positive selection, as expected. This

is consistent with previous studies that have proposed a derived allele frequency cutoff (Fay *et al*.

2001, 2002; Andolfatto 2005; but see Charlesworth & Eyre-Walker 2008) to correct for

segregating mildly deleterious alleles, as well as proposed modification of the traditional MK test

(*e.g.,* the asymptotic MK test; Messer & Petrov 2013). Nevertheless, under the asymptotic MK

test, α remains underestimated (Supp Table 6) when the proportion of mildly deleterious mutations

is sufficiently high.

In order to study inter-population effects, we simulated a model similar to that recently

inferred by Arguello *et al.* (2019), which represents the European and African split of *D.*

melanogaster (see Methods), and overlaid it with the estimated DFE of Johri *et al.* (2020). Under

this model (*i.e.,* in the absence of positive selection), roughly 50% of SNPs identified as F_{ST} outliers

(defined as representing the upper 1% or 2.5% tails) are mildly deleterious (Figure 4a; Supp Figure

3). Two models of purifying selection are given for comparison (Table 2): 1) a more biologically

realistic model in which selection coefficients are scaled to the ancestral population size in defining

the DFE classes, such that selection is effectively weaker in the smaller derived population; and 2)

an arbitrary model in which the DFE is rescaled such that selective effects are equally strong in

the larger ancestral and the smaller derived populations (which, under the chosen demographic

model, differ from one another by roughly an order of magnitude). It should be noted that in the

model based on Arguello *et al.*, the time post-split between the African and European population

is extremely brief, such that there are few or no substitutions post-split (Supp Table 7). Thus, F_{ST}

values are almost entirely dictated by allele frequency differences with respect to co-segregating

mutations (Supp Table 7).

Mean F_{ST} values at both neutral and weakly deleterious sites are larger with the rescaled

purifying selection model, whereas the unscaled model yields similar values to the purely neutral

model (Table 1 and Supp Tables 8, 10). In addition, the frequency of private SNPs at all sites is

higher for the rescaled model than the unscaled model, with the neutral model having the lowest

value of the three (Supp Table 7). In order to determine whether these weakly deleterious mutations

were associated with a decrease in diversity at linked sites, potentially leading to an increase in F_{ST}

values (Cruickshank & Hahn 2014), we evaluated the relationship between F_{ST} and nucleotide

diversity using neutral variants alone (Supp Figure 4; see also the comparison with directly

selected sites in Supp Figures 5-6). The lack of a strong negative correlation suggests that sweep-

like effects of deleterious mutations on diversity at linked sites are not primarily responsible for

the observed effects on F_{ST} and the frequency of private alleles. In contrast, there is a negative

relationship between nucleotide diversity and F_{ST} after the strong population bottleneck

represented by the neutral model of Li & Stephan (2006).

The lack of sweep-like effects of deleterious mutations is not surprising in view of the time-

scale to fixation required for weakly selected mutations, which is of the order of the coalescent

time, $2N$ generations; the split times for both the African and European populations are both only

a small fraction of their respective coalescent times. Thus, neither new neutral nor weakly

deleterious mutations are likely to have reached fixation in either population, consistent with the

results shown in the last column of Supp Table 7. In addition, there is no time for large changes in

π as a result of the altered N values.

Instead, it is more likely that BGS effects on the frequencies of segregating mutations

explain these patterns ($B = 0.18$ and 0.15 in the African and European populations, respectively,

obtained from the ratios of neutral π values in the absence of purifying selection to those without:

Supp Table 9). In the absence of rescaling of the DFEs for deleterious mutations (the more

biologically plausible case), the enhanced N for the African population means that fewer

deleterious mutations behave as effectively neutral, so that there is less effect of drift on their

frequencies compared with the ancestral population; the reverse is true for the European

population. As far as linked neutral variants are concerned, there is likely to be a greater BGS

effect in the African than in the ancestral population, and vice-versa for the European population.

F_{ST} relative to the purely neutral case is thus subject to two opposing factors, which presumably

explains the lack of any strong effect of this model of purifying selection on F_{ST} in Table 2 and

Supp Table 8. However, the lower effective population size induced by BGS means that rare

variants are more likely to be lost after the population split than under neutrality, explaining the

increased proportion of private variants compared with neutrality with purifying selection (Supp

Table 7). Because of the effects of BGS, the relative N_e values of the African and European

populations are less disparate than the relative N values, so that the African/European ratio of the

proportions of private SNPs is smaller than under neutrality (Supp Table 7).

The effect of rescaling is to keep the proportion of deleterious mutations that are effectively

neutral the same in the two descendant populations, with the absolute strength of selection being

higher in the African population, so that overall there is a stronger BGS effect in this population.

This results in a higher overall fraction of private SNPs, and a larger enrichment of private SNPs

in the African population, compared with the neutral case (Supp Table 7). There is a corresponding

increase in mean F_{ST} for both neutral and weakly deleterious variants compared with the neutral

case (Table 2 and Supp Table 8).

It is also of interest to consider the contribution of deleterious mutations to outliers in the

presence of positive selection. When a class of weakly positively selected sites was added to the

DFE (*i.e.* beneficial mutations have fitness effects $1 < 2Ns_a \leq 10$, and comprise 1% of new

mutations), these adaptive mutations contributed little to the observed outliers ($< 5\%$) - with mildly

deleterious and neutral mutations strongly represented amongst outliers (Figure 4b). Conversely,

when positive selection is very strong (*i.e.*, mutations have selective effects with $2Ns_a=1000$, and

comprise 1% of new mutations), the majority (~75%) of outliers are drawn from this beneficial

class (Figure 4c). Yet, even in the presence of this exceptionally strong and frequent positive

selection, ~10% of outliers remain in the mildly deleterious class. It is also noteworthy that under

this model of strong, recurrent positive selection, little variation within populations is observed,

owing to the severity of the selective sweeps (Supp Table 8), and inter-population allele

frequencies are only weakly correlated with one another (Supp Table 7).

Because any model, including equilibrium neutrality, will have outliers based on empirical

p-values, we further quantified the properties of outliers (2.5% and 1%) relative to the genome-

wide mean. Under models of purifying selection, the values of F_{ST} outliers were ~2.3-4.5 fold

larger than mean F_{ST} values (Table 2; Supp Table 10), but differed only slightly from the neutral

case. Under the strongly bottlenecked neutral model of Li & Stephan (2006), in which the

European population experiences a substantial decrease in population size during the bottleneck,

the genome-wide mean F_{ST} values obtained are much higher, as would be expected, though outlier

F_{ST} values were smaller in relative magnitude (~1.5-2.7 fold higher than the means). Under models

including weak positive selection, only slight increases in genome-wide F_{ST} values were observed;

whereas the strong positive selection model greatly increased genome-wide values. However,

outlier values under both positive selection models were ~2.4-4.6 fold higher than the respective

means. This suggests that recurrent positive selection does not generate substantially larger effect

sizes for outlier F_{ST} values than neutrality or purifying selection, although the proportion of outliers

is larger with strong positive selection.

CONCLUSIONS

In this paper, we have examined the expected impact of deleterious fixations on polymorphism-

and divergence-based scans for selection. Amongst the class of weakly deleterious mutations that

have some chance of reaching fixation ($1 < 2Ns_d \leq 10$), the resulting sweep effects are highly

localized, as expected: on the order of a few dozen base pairs for the parameters considered here.

This suggests that deleterious sweeps of this kind are unlikely to be detected in genomic scans

based on localized deficits of variation or strongly skewed site frequency spectra. Given the

theoretically expected symmetry between beneficial and deleterious sweeps (Maruyama &

Kimura 1974; Mafessoni & Lachmann 2015; Charlesworth 2020a), this is expected, as common

polymorphism-based methods generally have little power unless selection is exceptionally strong

(*e.g.,* Kim & Stephan 2002; Jensen *et al.* 2005; Crisci *et al*. 2013). However, our results suggest

that studies which estimate the frequency and strength of classic selective sweeps using patterns

of diversity around substitutions (Hernandez *et al*. 2011; Sattath *et al*. 2011; Elyashiv *et al*.

2016), which assume that reductions are entirely caused by the fixation of positively selected

mutations, ought to take into account reductions caused by neutral and weakly deleterious

substitutions.

Furthermore, for among-population comparisons based on F_{ST}, mildly deleterious

mutations contribute significantly to observed outliers, even in the presence of positive selection,

particularly in the case of a recent population size reduction. This appears to be the true regardless

of whether selective effects are equally strong in both populations (Figure 4), or if selection is

relaxed in the smaller derived population (Supp Figure 3). Because such effects are localized to

functional sites (*i.e.,* those genomic regions experiencing purifying selection), this effect may be

particularly pernicious in the sense that this class of outlier will not fall in non-functional regions,

where they are often attributed to demographic effects. Rather, owing to the common tendency of

constructing biological narratives (true or otherwise) around functional outliers (Pavlidis *et al*.

2012), these results suggest that adaptive story-telling may arise from weakly deleterious outliers.

However, we have only considered one specific scenario, and the generality of this conclusion

remains to be established.

557
558

559   **ACKNOWLEDGMENTS**

560
561   This work was funded by National Institutes of Health grant R01GM135899 to JDJ.

APPENDIX 1

Calculating the probability of fixation with uniformly distributed fitness effects

For a selection coefficient $s \ll 1$, the fixation probability of a deleterious mutation when $N_e \neq N$ is approximated by:

$$P_{fix} \approx \frac{s\left(\frac{N_e}{N}\right)}{\exp(2N_e s) - 1} \qquad (A1)$$

For large N_e, this should be a good approximation, since when $2N_e s \geq 10$, the fixation probability is negligible, so that the contribution from the bin with $2N_e s \geq 10$ can be ignored. For $2N_e s < 10$, s is sufficiently small that this equation is an accurate approximation.

The integral of this expression over a given interval of $2N_e s$ values can be found as follows; for convenience, x is substituted for s and a for $2N_e$. For $a > 0$, we need to evaluate the following indefinite integral:

$$I = \int x(e^{ax} - 1)^{-1} \, dx$$

Integration by parts give:

$$I = \frac{1}{a}\left[x\ln(1 - e^{-ax}) - \int \ln(1 - e^{-ax}) \, dx\right] \qquad (A2)$$

For $a > 0$, the logarithm can be expanded as a power series in $\exp(-ax)$, which can be integrated term by term:

$$\int \ln(1 - e^{-ax}) \, dx = \int\left(-e^{-ax} - \frac{1}{2}e^{-2ax} - \frac{1}{3}e^{-3ax} - \cdots\right)dx$$

$$= a^{-1} \sum_{i=1}^{\infty} i^{-2} e^{-iax} \qquad (A3)$$

The final expression for the integral of equation (1) is thus:

$$\frac{1}{2N}[s \ln(1 - e^{-2N_e s}) - (2N_e)^{-1} \sum_{i=1}^{\infty} i^{-2} e^{-2iN_e s}] \qquad (A4)$$

The contribution to the mean fixation probability from the interval s_i to s_{i+1} is obtained

by dividing this expression by $(s_{i+1} - s_i)$.

For $s_0 = 0$, equation (A4) is invalid. However, the integral between 0 and a small positive

value of s, s_ε, can be found as follows. For $2N_e s \ll 1$, the initial integrand can be approximated

by $a^{-1}(1 - ax/2)$, so that equation 2 can be replaced with:

$$a^{-1}x - \frac{1}{4}x^2$$

and the indefinite integral of the fixation probability becomes:

$$[(2N_e)^{-1}s - \frac{1}{4}s^2](N_e/N) = \frac{s}{2N}[1 - \frac{1}{2}N_e s)] \qquad (A5a)$$

The contribution to the integral of P_{fix} between $s_0 = 0$ and s_1 from the interval $(0, s_\varepsilon)$ is thus:

$$[(2N_e)^{-1}s_\varepsilon - \frac{1}{2}s_\varepsilon^2](N_e/N) = \frac{s_\varepsilon}{2N}[1 - \frac{1}{2}(N_e s_\varepsilon)] \qquad (A5b)$$

The corresponding mean fixation probability over this interval is:

$$\frac{1}{2N}[1 - \frac{1}{2}s(N_e s_\varepsilon)]$$

This is slightly smaller than the neutral value, $1/(2N)$, as would be expected when $2N_e s \ll 1$.

Thus, for the interval (s_0, s_1) with $s_0 = 0$, equation (A5b) should be used for the interval

$(0 \le s \le s_\varepsilon)$, and equation (A4) with integration limits s_ε and s_1 for the remainder of the interval.

FIGURES AND TABLES

Table 1: Reduction in nucleotide diversity at linked neutral sites relative to the neutral value after the fixation of a weakly deleterious or advantageous semi-dominant mutation. The distance between the selected and neutral site is presented in units of the scaled recombination rate (ρ), as well as in the number of bases corresponding to *Drosophila*-like populations.

Distance from selected site in scaled recombination rate (ρ)	Corresponding number of bases in *Drosophila*		Expected diversity reduction	Expected diversity reduction		Expected diversity reduction	
	Without gene conversion	With gene conversion	$2Ns = 0$	$2Ns = 5$		$2Ns = 10$	
				A	B	A	B
0	0	0	0.426	0.459	0.496	0.523	0.534
0.05	2.5	0.8	0.393	0.429	0.471	0.502	0.514
0.10	5	1.7	0.372	0.409	0.446	0.479	0.495
0.20	10	3.3	0.325	0.371	0.402	0.452	0.459
0.40	20	6.7	0.261	0.301	0.328	0.393	0.395
0.80	40	13.5	0.184	0.221	0.228	0.304	0.298
1.60	80	27.2	0.100	0.127	0.138	0.200	0.182
3.20	160	55.6	0.047	0.063	0.100	0.107	0.101
6.40	320	116	0.018	0.025	0.077	0.044	0.069
12.8	640	254	0.006	0.009	0.043	0.014	0.040

The double entries in the two right-hand columns are the results of A) 1000 replicate simulations using equation (27) of Tajima (1990), and B) the analytical approximation of Charlesworth (2020b, equation 14). For the simulations, a population size of 100 and a scaled neutral mutation rate per bp of 2.5 x 10^{-6} were assumed. 1000 replicate runs were performed. Note that if the diffusion approximation is valid, only the scaled parameter values (ρ, $2Ns$ and $\theta = 4Nu = 0.001$) are relevant.

658 **Table 2**: Mean genome-wide and outlier F_{ST} values (calculated in windows of 500 bp (labeled
659 'Constant total sites'), or 10 SNPs (labeled 'Constant SNPs')).
660

Evolutionary model	Mean F_{ST}		Outlier F_{ST} (1%)		Outlier F_{ST} (1%)/ Mean F_{ST}	
	Constant total sites	Constant SNPs	Constant total sites	Constant SNPs	Constant total sites	Constant SNPs
Neutrality	0.0374	0.0363	0.0874	0.1628	2.3369	4.4838
Purifying selection	0.0365	0.0352	0.0880	0.1454	2.4110	4.1296
Purifying selection (rescaled DFE)	0.0666	0.0613	0.1585	0.2637	2.3799	4.2999
With 1% weak positive selection	0.0578	0.0523	0.1403	0.2412	2.4273	4.6150
With 1% strong positive selection	0.2261	0.1976	0.7673	0.7455	3.3936	3.7731
Neutral values, with the Li & Stephan (2006) bottleneck	0.2828	0.2557	0.4154	0.6956	1.4689	2.7204

662 These results, with the exception of the Li & Stephan row, use the Arguello *et al.* (2019)
663 parameters for European and African *D. melanogaster* population history, under different DFE
664 configurations. Additional details may be found in the text and Supp Table 10.
665
666

Figure 1: (a) Probability of fixation of weakly deleterious mutations relative to that of neutral mutations. (b) The distribution of fixation times (conditional on fixation) of mutations with varying selective effects, obtained from 100 simulated replicates. Fixation times are measured as the time taken for the mutant allele to spread from frequency $1/2N$ to frequency 1. Black solid circles are the means of the distributions obtained from simulations, and red solid circles are the mean expectations obtained from numerically integrating the expression of Kimura & Ohta (1969). The dominance coefficient is 0.5 for all mutations except in the cases "dom" and "rec" where $h = 1$ and 0, respectively.

Figure 2: Recovery of nucleotide diversity per site (π) relative to that expected under neutrality (π_0), around a recent fixation (shown at position 0 on the x-axis). The target site has experienced (a) a neutral fixation ($2Ns = 0$; black lines), (b) a weakly deleterious fixation ($2Ns_d = 5$; red line), and (c) a weakly beneficial fixation ($2Ns_a = 5$; blue line). Solid lines represent mean values of 100 replicates, shaded regions correspond to 1 SE above and 1 SE below the mean. Solid circles show the theoretical predictions using equation (14) of Charlesworth 2020b; and crosses correspond to simulations based on equation (27) of Tajima (1990).

Figure 3: Predicted mean reductions of nucleotide diversity at linked neutral sites compared to neutrality ($-\Delta\pi$), due to the recurrent fixation of weakly deleterious semi-dominant mutations with $2.5 \leq 2Ns_d \leq 10$. Nucleotide diversity at neutral sites was averaged across a gene comprised of five 300-bp exons and 100-bp introns, in which all intronic sites and 30% of exonic sites were neutral.

Figure 4: Allele frequencies of SNPs in simulated population 1 (European) *vs* population 2
(African), using parameters of the Arguello *et al.* (2019) model, where the selective effects of all
mutations were rescaled with respect to their population sizes after the split (*i.e.,* keeping the
strength of selection constant in both populations). Genomic elements experienced (a) purifying
selection following the DFE inferred by Johri *et al.* (2020); (b) the same DFE, but with the
addition of 1% beneficial mutations with selective effects between $1 < 2Ns_a \le 10$; or (c) the
same DFE, but with the addition of 1% beneficial mutations with selective effects of $2Ns_a =$
1000. *Left panel*: Allele frequency plots for 10 (out of 100) replicates simulated. Colored open
circles represent F_{ST} outliers. Green depicts effectively neutral mutations (belonging to class 0),
blue depicts beneficial mutations, and warm colors depict deleterious mutations (belonging to
classes 1, 2, and 3), with red representing weakly deleterious mutations. *Right panel*: The
distribution of fitness effects of outlier mutations for the corresponding scenarios, showing the
mean and standard deviation for all 100 replicates. Sites that were fixed in both populations for
the same allele were not included in this analysis.

CITATIONS

Andolfatto P., 2005 Adaptive evolution of non-coding DNA in *Drosophila*. Nature 437: 1149-
1152.

Andolfatto P., 2007 Hitchhiking effects of recurrent beneficial amino acid substitutions in the
Drosophila melanogaster genome. Genome Res. 17: 1755–1762.

Arguello J.R., S. Laurent, and A.G. Clark, 2019 Demographic history of the human commensal
Drosophila melanogaster. Genome Biol. Evol. 11: 844–854.
717
718   Atkinson, K.E., 1989 Introduction to numerical analysis. John Wiley, New York, NY.
719
720   Bank, C., M. Foll, A. Ferrer-Admetlla, G. Ewing, and J.D. Jensen, 2014a Thinking too positive?
721   Revisiting current methods in population genetic selection inference. Trends Genet. 30: 540-6.
722
723   Bank C., R.T. Hietpas, A. Wong, D.N. Bolon, and J.D. Jensen, 2014b A Bayesian MCMC
724   approach to assess the complete distribution of fitness effects of new mutations: uncovering the
725   potential for adaptive walks in challenging environments. Genetics 196: 841–852.
726
727   Barton N.H., 2000 Genetic hitchhiking. Phil. Trans. R. Soc. Lond. B 355: 1553-62.
728
729   Campos J.L., L. Zhao, and B. Charlesworth, 2017 Estimating the parameters of background
730   selection and selective sweeps in *Drosophila* in the presence of gene conversion. Proc. Natl.
731   Acad. Sci. U.S.A. 114: E4762–E4771.
732
733   Campos J.L., and B. Charlesworth, 2019 The effects on neutral variability of recurrent selective
734   sweeps and background selection. Genetics 212: 287–303.
735
736   Charlesworth B., M.T. Morgan, and D. Charlesworth, 1993 The effect of deleterious mutations
737   on neutral molecular variation. Genetics 134: 1289–1303.
738
739   Charlesworth B., 2020a How long does it take to fix a favorable mutation, and why should we
740   care? Am. Nat. 195: 753–771.
741
742   Charlesworth B., 2020b How good are predictions of the effects of selective sweeps on levels of
743   neutral diversity? bioRxiv 2020.05.27.119883.
744
745   Charlesworth J., and A. Eyre-Walker, 2007 The other side of the nearly neutral theory, evidence
746   of slightly advantageous back-mutations. Proc. Natl. Acad. Sci. USA 104: 16992-997.
747
748   Charlesworth J., and A. Eyre-Walker, 2008 The McDonald–Kreitman test and slightly
749   deleterious mutations. Mol. Biol. Evol. 25: 1007–1015.
750
Comeron J.M., R. Ratnappan, and S. Bailin, 2012 The many landscapes of recombination in
Drosophila melanogaster. PLoS Genet. 8:e1002905.

Crisci J., Y.-P. Poh, S. Mahajan, and J.D. Jensen, 2013 The impact of equilibrium assumptions
on tests of selection. Front. Genet. 4: 235.

Crow J.F., 1993 Mutation, mean fitness, and genetic load. Oxf. Surv. Evol. Biol. 9: 3-42.

Cruickshank T.E., and M.W. Hahn, 2014 Reanalysis suggests that genomic islands of speciation
are due to reduced diversity, not reduced gene flow. Mol. Ecol. 23: 3133–3157.

Elyashiv E., S. Sattath, T.T. Hu, A. Strutsovsky, G. McVicker, et al., 2016 A genomic map of the
effects of linked selection in *Drosophila*. PLoS Genet. 12: e1006130.

Eyre-Walker A., and P.D. Keightley, 2007 The distribution of fitness effects of new mutations.
Nat. Rev. Genet. 8: 610-8.

Eyre-Walker A., and P.D. Keightley, 2009 Estimating the rate of adaptive molecular evolution in
the presence of slightly deleterious mutations and population size change. Mol. Biol. Evol. 26:
2097–2108.

Fay J., G.J. Wycoff, and C.-I Wu, 2001 Positive and negative selection on the human genome.
Genetics 158: 1227-1234

Fay J., G.J. Wycoff, and C.-I Wu, 2002 Testing the neutral theory of molecular evolution with
genomic data from *Drosophila*. Nature 415: 1024-1026.

Fisher R.A. 1922. On the dominance ratio. Proceedings of the Royal Society of Edinburgh 42:
321-341.

Fisher R.A., 1930 *The genetical theory of natural selection*. Clarendon Press, Oxford.

Fiston-Lavier A.-S., N.D. Singh, M. Lipatov, and D.A. Petrov, 2010 *Drosophila melanogaster*
recombination rate calculator. Gene 463: 18–20.

Frisse L., R.R. Hudson, A. Bartoszewicz, J.D. Wall, J. Donfack, and A. Di Rienzo, 2001 Gene
conversion and population histories may explain the contrast between polymorphism and linkage
disequilibrium levels. Am. J. Hum. Genet. 69: 831-43.

Gillespie J.H., 1994 Substitution processes in molecular evolution. III. Deleterious alleles.
Genetics 138: 943-952.

Glinka S.L., L. Ometto, S. Mousset, W. Stephan, and D. De Lorenzo, 2003 Demography and
natural selection have shaped genetic variation in *Drosophila melanogaster*: a multi-locus
approach. Genetics 165:1269–1278.

Haddrill P.R., K.R. Thornton, B. Charlesworth, and P. Andolfatto, 2005 Multilocus patterns of
nucleotide variability and the demographic and selection history of *Drosophila melanogaster*
populations. Genome Res. 15:790–799.

Haldane J.B.S. 1927. A mathematical theory of natural and artificial selection. Part V. Selection
and mutation. Math. Proc. Camb. Philos. Soc 23:838-844.

Haller B.C., and P.W. Messer, 2017 asymptoticMK: A web-based tool for the asymptotic
McDonald–Kreitman test. G3 7: 1569–1575.
798
799 Haller B.C., and P.W. Messer, 2019 SLiM 3: Forward genetic simulations beyond the Wright-
800 Fisher model. Mol. Biol. Evol. 36: 632–637.
801
Halligan D.L., and P.D. Keightley, 2006 Ubiquitous selective constraints in the *Drosophila*
genome revealed by a genome-wide interspecies comparison. Genome Res. 16: 875–884.

Harr B., M. Kauer, and C. Schlotterer, 2002 Hitchhiking mapping: a population-based fine-
mapping strategy for adaptive mutations in *Drosophila melanogaster*. Proc. Natl. Acad. Sci.
USA 99: 12949–12954.

Harris R.B., A. Sackman, and J.D. Jensen, 2018 On the unfounded enthusiasm for soft selective
sweeps II: examining recent evidence from humans, flies, and viruses. PLoS Genetics 14(12):
e1007859.
810
811 Hartfield M., and T. Bataillon, 2020 Selective sweeps under dominance and inbreeding. G3 10:
812 1063-75.
813
814 Hernandez R.D., J.L. Kelley, E. Elyashiv, S.C. Melton, A. Auton, et al., 2011 Classic selective
815 sweeps were rare in recent human evolution. Science 331: 920–924.
816
817 Hudson R.R., M. Slatkin, and W.P. Maddison, 1992 Estimating levels of gene flow from DNA
818 sequence data. Genetics 13: 583-589.
819
820 Jacquier H., A. Birgy, H.L. Nagard, Y. Mechulam, E. Schmitt, et al., 2013 Capturing the
821 mutational landscape of the beta-lactamase TEM-1. Proc. Natl. Acad. Sci. USA 110: 13067–
822 13072.
823
824 Jakobsson M., M.D. Edge, and N.A. Rosenberg, 2013 The relationship between F_{ST} and the
825 frequency of the most frequent allele. Genetics 193: 515–528.
826
Jensen J.D., Y. Kim, V.B. DuMont, C.F. Aquadro, and C.D. Bustamante, 2005 Distinguishing
between selective sweeps and demography using DNA polymorphism data. Genetics 170:1401–
1410.

Jensen J.D., K.R. Thornton, and P. Andolfatto, 2008 An approximate Bayesian estimator
suggests strong recurrent selective sweeps in *Drosophila*. PLoS Genet. 4:e1000198.

Jensen J.D., 2009 On reconciling single and recurrent hitchhiking models. Genome Biol. Evol. 1:
320-4.
834
835   Johri P., B. Charlesworth, and J.D. Jensen, 2020 Toward an evolutionarily appropriate null
836   model: jointly inferring demography and purifying selection. Genetics 215: 173–192.
837
838   Keightley P.D., U. Trivedi, M. Thomson, F. Oliver, S. Kumar, et al., 2009 Analysis of the
839   genome sequences of three *Drosophila melanogaster* spontaneous mutation accumulation lines.
840   Genome Res. 19: 1195–1201.
841
Keightley P.D., R.W. Ness, D.L. Halligan, and P.R. Haddrill, 2014 Estimation of the
spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family.
Genetics 196: 313–320.

Kim Y., 2006 Allele frequency distribution under recurrent selective sweeps. Genetics 172:
1967–1978.

Kim Y., and W. Stephan, 2002 Detecting a local signature of genetic hitchhiking along a
recombining chromosome. Genetics 160: 765–777.

Kimura M., 1957 Some problems of stochastic processes in genetics. Ann. Math. Stat. 28: 882-
901.
851
852   Kimura M., 1962 On the probability of fixation of mutant genes in a population. Genetics 47:
853   713–719.
854
855   Kimura M., 1964 Diffusion models in population genetics. J. Appl. Probab. 1: 177-232.
856
857   Kimura, M., 1970 The length of time required for a selectively neutral mutation to reach fixation
858   through random frequency drift in a finite population. Genet. Res. 15: 131-133.
859
860   Kimura M., and T. Ohta, 1969 The average number of generations until fixation of a mutant gene
861   in a finite population. Genetics 61: 763–771.
862
863   Li H., and W. Stephan, 2006 Inferring the demographic history and rate of adaptive substitution
864   in *Drosophila*. PLoS Genet. 2: e166.
865
866   Li Y.J., Y. Satta, and N. Takahata, 1999 Paleo-demography of the *Drosophila melanogaster*
867   subgroup: application of the maximum likelihood method. Genes Genet. Syst. 74: 117–127.
868
Lynch M., D. Blanchard, D. Houle, T. Kibota, S. Schultz, L. Vassilieva, and J. Willis, 1999
Spontaneous deleterious mutations. Evolution 53: 645-63.

Macpherson J.M., G. Sella, J.C. Davis, and D.A. Petrov, 2007 Genome-wide spatial
correspondence between nonsynonymous divergence and neutral polymorphism reveals
extensive adaptation in *Drosophila*. Genetics 177: 2083–2099.

Mafessoni F., and M. Lachmann, 2015 Selective strolls: fixation and extinction in diploids
are slower for weakly selected mutations than for neutral ones. Genetics 201: 1581-1589.

Maruyama T., and M. Kimura, 1974 A note on the speed of gene frequency changes in reverse
directions in a finite population. Evolution 28: 161-163.

Maynard Smith J., and J. Haigh, 1974 The hitch-hiking effect of a favourable gene. Genet. Res.
23: 23-35.

McDonald J.H., and M. Kreitman, 1991 Adaptive protein evolution at the Adh locus in
Drosophila. Nature 351:652–654.
882
883   Messer P.W., and D.A. Petrov, 2013 Frequent adaptation and the McDonald–Kreitman test.
884   Proc. Natl. Acad. Sci. U.S.A. 110: 8615–8620.
885
Miller D.E., C.B. Smith, N.Y. Kazemi, A.J. Cockrell, A.V. Arvanitakis, J.P. Blumenstiel, S.L.
Jaspersen, and R.S. Hawley, 2016 Whole-genome analysis of individual meiotic events in
Drosophila melanogaster reveals that noncrossover gene conversions are insensitive to
interference and the centromere effect. Genetics 203:159–171.

Nielsen R., S. Williamson, Y. Kim, M.J. Hubisz, A.G. Clark, and C.D. Bustamante, 2005
Genomics scans for selective sweeps using SNP data. Genome Res. 15: 1566–1575.

Pavlidis P., J.D. Jensen, W. Stephan, and A. Stamatakis, 2012 A critical assessment of
storytelling: Gene Ontology categories and the importance of validating genomic scans. Mol.
Biol. Evol. 29: 3237–3248.
895
Pfeifer B., U. Wittelsbürger, S.E. Ramos-Onsins, and M.J. Lercher, 2014 PopGenome: an
efficient Swiss army knife for population genomic analyses in R. Mol. Biol. Evol. 31: 1929–
1936.

Przeworski M., 2002 The signature of positive selection at randomly chosen loci. Genetics 160:
1179–1189.

R Core Team, 2018 R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Sanjuán R., 2010 Mutational fitness effects in RNA and single-stranded DNA viruses: common
patterns revealed by site-directed mutagenesis studies. Philos. Trans. R. Soc. Lond. B Biol. Sci.
365: 1975-82.
906
Sattath S., E. Elyashiv, O. Kolodny, Y. Rinott, and G. Sella, 2011 Pervasive adaptive protein
evolution apparent in diversity patterns around amino acid substitutions in *Drosophila simulans*.
PLoS Genet. 7:e1001302.

Sella G., D.A. Petrov, M. Przeworski, and P. Andolfatto, 2009 Pervasive natural selection in the
Drosophila genome. PLoS Genet. 5:e1000495.

Stephan W., 2019 Selective sweeps. Genetics 211: 5-13.

Tajima F., 1990 Relationship between DNA polymorphism and fixation time. Genetics 125:
447–454.

Teshima K.M., G. Coop, and M. Przeworski, 2006 How reliable are empirical genome scans for
selective sweeps? Genome Res. 16: 702–712.

Thornton K., 2003 Libsequence: a C++ class library for evolutionary genetic analysis.
Bioinformatics 19: 2325–2327.
919
920   Thornton K.R., and J.D. Jensen, 2007 Controlling the false positive rate in multi-locus genome
921   scans for selection. Genetics 175: 737-750.

922   Wiehe T.H., and W. Stephan, 1993 Analysis of a genetic hitchhiking model and its application to
923   DNA polymorphism data from *Drosophila melanogaster*. Mol. Biol. Evol. 10: 842–854.

924   Wright S., 1931 Evolution in Mendelian populations. Genetics 16:97-159.

SUPPLEMENTARY MATERIAL

Supp Table 1: Parameters of the demographic models of *D. melanogaster* simulated for the F_{ST} analyses.

	Arguello *et al.* 2019	Li & Stephan 2006
Ancestral population size	1.95×10^6	8.603×10^6
Time of split of European population	6.62×10^4 generations ago	1.58×10^5 generations ago
European bottleneck size	-	2.2×10^3
European bottleneck duration	-	3.4×10^3 generations
European current size	4.73×10^5	1.075×10^6
African current size	3.91×10^6	8.603×10^6

Supp Table 2: The means and standard deviations of fixation times (conditional on fixation) of mutations with varying selective effects, obtained from 100 simulated replicates. Fixation times are measured as the time taken for the mutant allele to spread from frequency $1/2N$ to frequency 1. Unless otherwise indicated, h=0.5.

			Time to fixation (in $2N$ generations)				
		First quartile	**Median**				**Analytical**
$2Ns$	**min**	**(25%)**	**(50%)**	**Mean**	**Max**	**SD**	**Mean**
0	0.57	1.30	1.82	2.08	5.77	1.04	2.00
-1	0.45	1.20	1.67	1.96	6.19	1.05	1.97
1	0.58	1.14	1.52	1.73	4.09	0.99	1.97
-5	0.53	1.14	1.36	1.55	3.37	0.74	1.56
5	0.58	1.15	1.49	1.66	5.42	0.57	1.56
5 (*h*=0)	0.43	0.99	1.19	1.31	2.82	0.47	-
5 (*h*=1)	0.49	1.21	1.57	1.83	6.91	0.99	-
10	0.50	0.77	0.97	1.04	2.18	0.36	1.11
20	0.33	0.56	0.70	0.73	1.40	0.23	0.70
30	0.25	0.44	0.51	0.53	1.05	0.14	0.53
40	0.23	0.38	0.43	0.44	0.80	0.10	0.42

Supp Table 3: Reduction in diversity (relative to a value of 1 for neutrality) as (A) obtained by
Tajima's (1990) simulations and (B) as approximated by numerical results using equation (14) in
Charlesworth 2020b.

ρ	$2Ns_a = 4$		$2Ns_a = 8$		$2Ns_a = 20$		$2Ns_a = 40$		$2Ns_a = 80$	
	A	**B**	**A**	**B**	**A**	**B**	**A**	**B**	**A**	**B**
0	0.448	0.513	0.494	0.512	0.620	0.633	0.734	0.741	0.830	0.830
0.005	0.445	0.511	0.492	0.510	0.619	0.632	0.732	0.740	0.830	0.830
0.05	0.422	0.486	0.473	0.490	0.606	0.618	0.723	0.730	0.823	0.824
0.10	0.398	0.461	0.453	0.469	0.592	0.604	0.713	0.720	0.816	0.817
0.25	0.398	0.417	0.401	0.412	0.552	0.563	0.684	0.690	0.796	0.796
0.50	0.266	0.306	0.333	0.335	0.493	0.501	0.638	0.643	0.763	0.764
1.00	0.179	0.201	0.240	0.229	0.399	0.399	0.558	0.559	0.702	0.702
2.50	0.077	0.114	0.114	0.111	0.231	0.215	0.384	0.372	0.550	0.548
5.00	0.031	0.093	0.049	0.079	0.113	0.102	0.222	0.201	0.375	0.368
10.0	0.010	0.057	0.016	0.051	0.041	0.053	0.113	0.080	0.190	0.177

986 **Supp Table 4:** Reduction of diversity ($\Delta\pi$) at ρ ($= 4Nr$) distance from the selected site for
987 varying dominance (h) and selection coefficients ($2Ns$) calculated using the algorithm presented
988 in Tajima (1990). For the purpose of these calculations N was assumed to be 100 and $\theta = 0.001$.

$\Delta\pi$		0	0.05	0.1	0.2	0.4	0.8	1.6	3.2	6.4	12.8
$2Ns = 0$; 100000 replicates											
	Mean	0.4215	0.3907	0.3635	0.3178	0.2508	0.1706	0.0966	0.0451	0.0178	0.0062
	SE	3.3E-04	3.5E-04	3.6E-04	3.8E-04	3.9E-04	3.6E-04	2.7E-04	1.6E-04	7.1E-05	2.4E-05
$2Ns_a = 2\ [2Ns_d = 2]$; 100000 replicates											
h=0	**Mean**	0.4430	0.4157	0.3909	0.3482	0.2825	0.1990	0.1164	0.0556	0.0220	0.0076
[h=1]	**SE**	3.3E-04	3.4E-04	3.5E-04	3.7E-04	3.8E-04	3.5E-04	2.7E-04	1.6E-04	7.6E-05	3.1E-05
h=0.1	**Mean**	0.4400	0.4121	0.3871	0.3440	0.2781	0.1949	0.1149	0.0547	0.0217	0.0075
[h=0.9]	**SE**	3.3E-04	3.4E-04	3.6E-04	3.7E-04	3.8E-04	3.5E-04	2.7E-04	1.6E-04	7.4E-05	3.0E-05
h=0.2	**Mean**	0.4371	0.4089	0.3836	0.3401	0.2741	0.1913	0.1109	0.0527	0.0208	0.0072
[h=0.8]	**SE**	3.3E-04	3.5E-04	3.6E-04	3.7E-04	3.8E-04	3.5E-04	2.7E-04	1.6E-04	7.4E-05	2.7E-05
h=0.3	**Mean**	0.4344	0.4057	0.3801	0.3364	0.2704	0.1880	0.1085	0.0514	0.0203	0.0070
[h=0.7]	**SE**	3.3E-04	3.5E-04	3.6E-04	3.8E-04	3.8E-04	3.5E-04	2.7E-04	1.6E-04	7.4E-05	2.9E-05
h=0.4	**Mean**	0.4317	0.4029	0.3769	0.3328	0.2664	0.1843	0.1061	0.0501	0.0198	0.0069
[h=0.6]	**SE**	3.3E-04	3.5E-04	3.6E-04	3.8E-04	3.9E-04	3.5E-04	2.7E-04	1.6E-04	7.3E-05	2.9E-05
h=0.5	**Mean**	0.4289	0.3996	0.3734	0.3290	0.2623	0.1807	0.1035	0.0488	0.0192	0.0067
[h=0.5]	**SE**	3.3E-04	3.5E-04	3.6E-04	3.8E-04	3.9E-04	3.5E-04	2.7E-04	1.6E-04	7.2E-05	2.6E-05
h=0.6	**Mean**	0.4264	0.3967	0.3699	0.3251	0.2585	0.1773	0.1011	0.0474	0.0187	0.0065
[h=0.4]	**SE**	3.3E-04	3.5E-04	3.6E-04	3.8E-04	3.9E-04	3.5E-04	2.7E-04	1.6E-04	7.2E-05	2.8E+00
h=0.7	**Mean**	0.4236	0.3935	0.3665	0.3210	0.2545	0.1737	0.0987	0.0461	0.0182	0.0064
[h=0.3]	**SE**	3.3E-04	3.5E-04	3.6E-04	3.8E-04	3.9E-04	3.5E-04	2.6E-04	1.5E-04	7.1E-05	2.8E-05
h=0.8	**Mean**	0.4211	0.4032	0.3769	0.3326	0.2673	0.1896	0.0963	0.0449	0.0177	0.0062
[h=0.2]	**SE**	3.3E-04	3.3E-04	3.5E-04	3.6E-04	3.7E-04	3.3E-04	2.6E-04	1.5E-04	7.1E-05	3.1E-05
h=0.9	**Mean**	0.4218	0.3868	0.3595	0.3132	0.2463	0.1666	0.0938	0.0436	0.0171	0.0060
[h=0.1]	**SE**	3.4E-04	3.5E-04	3.6E-04	3.8E-04	3.9E-04	3.5E-04	2.6E-04	1.5E-04	7.1E-05	3.0E-05
h=1	**Mean**	0.4158	0.3842	0.3562	0.3097	0.2429	0.1633	0.0915	0.0425	0.0167	0.0059
[h=0]	**SE**	3.3E-04	3.5E-04	3.7E-04	3.8E-04	3.9E-04	3.5E-04	2.6E-04	1.5E-04	7.1E-05	2.7E-05
$2Ns_a = 5\ [2Ns_d = 5]$; 1000 replicates											
h=0	**Mean**	0.4916	0.4724	0.4422	0.4006	0.3516	0.2554	0.1544	0.0812	0.0321	0.0109
[h=1]	**SE**	3.1E-03	3.2E-03	3.3E-03	3.3E-03	3.6E-03	3.4E-03	2.8E-03	1.8E-03	8.2E-04	3.1E-04
h=0.9	**Mean**	0.4249	0.4022	0.3755	0.3357	0.2712	0.1899	0.1119	0.0523	0.0213	0.0074
[h=0.1]	**SE**	3.5E-03	3.7E-03	3.7E-03	4.0E-03	4.1E-03	3.7E-03	2.7E-03	1.7E-03	7.9E-04	3.0E-04
h=1	**Mean**	0.4201	0.3932	0.3717	0.3268	0.2677	0.1774	0.1046	0.0530	0.0185	0.0068
[h=0]	**SE**	3.5E-04	3.5E-03	3.9E-03	3.9E-03	4.2E-03	3.7E-03	2.8E-03	1.7E-03	7.7E-04	3.0E-04
$2Ns_a = 10\ [2Ns_d = 10]$; 1000 replicates											
h=0	**Mean**	0.5701	0.4724	0.4422	0.4006	0.3516	0.2554	0.1544	0.0812	0.0321	0.0109
[h=1]	**SE**	2.5E-03	3.2E-03	3.3E-03	3.3E-03	3.6E-03	3.4E-03	2.8E-03	1.8E-03	8.2E-04	3.1E-04
h=1	**Mean**	0.4507	0.4315	0.4124	0.3796	0.3231	0.2452	0.1548	0.0794	0.0319	0.0109
[h=0]	**SE**	1.2E-03	1.2E-03	1.2E-03	1.3E-03	1.3E-03	1.2E-03	9.8E-04	6.1E-04	2.9E-04	1.1E-04

Supp Table 5: Distortion of the SFS at neutral sites linked to the selected site at a distance measured by ρ, and calculated immediately after the fixation of a weakly selected mutation. 100 replicates were used to obtain these values. Tajima's D was calculated in non-overlapping sliding windows of 50 base pairs, with the distance from the selected site corresponding to the mid-point of the window.

ρ	Number of bases from the selected site in *Drosophila*	$2Ns = 0$ Tajima's D mean	$2Ns = 0$ Tajima's D SE	$2Ns = +5$ Tajima's D mean	$2Ns = +5$ Tajima's D SE	$2Ns = -5$ Tajima's D mean	$2Ns = -5$ Tajima's D SE
1	25	-0.1493	0.0712	-0.2097	0.0671	-0.1852	0.0867
3	75	-0.0979	0.0696	-0.0827	0.0717	-0.0476	0.0919
5	125	0.00165	0.0731	-0.1042	0.0712	-0.0176	0.0871
7	175	-0.0355	0.0657	-0.0731	0.0699	-0.0193	0.0869
9	225	-0.0864	0.0712	-0.0214	0.0729	-0.1359	0.0816

Supp Table 6: Number of polymorphic (P) and fixed (D) sites at nonsynonymous (N) and synonymous (S) sites, and the estimated proportion of substitutions fixed by positive selection (α), for an increasing proportion of mildly deleterious mutations at nonsynonymous sites. The ratio of the rate of divergence at nonsynonymous relative to that at synonymous sites is denoted by d_N/d_S.

	% weakly deleterious NS sites	P_N	D_N	P_S	D_S	d_N/d_S	α	α_{asym} [95% CI]
Neutral	0%	522	439	255	221	0.99	-0.05	-0.06 [-0.21, 0.09]
Weakly deleterious	50%	447	261	240	237	0.55	-0.58	-0.35 [-0.72, 0.03]
Johri *et al*. DFE	66%	281	60	173	254	0.12	-5.22	-3.55 [-5.10, -2.01]

Supp Table 7: Composition of shared and private SNPs between populations (calculated in sliding windows of 500 bp).

	Fraction of shared SNPs	Fraction of private SNPs in African population	Fraction of private SNPs in European population	Fraction of fixed differences
Neutral	0.538	0.355	0.108	0.000
Purifying selection (rescaled)	0.363	0.521	0.116	0.000
Purifying selection (not rescaled)	0.417	0.440	0.143	0.000
Weak positive selection	0.379	0.502	0.119	0.000
Strong positive selection	0.040	0.640	0.319	0.001
Neutral bottleneck	0.242	0.706	0.052	0.000

1049 **Supp Table 8**: Mean F_{ST} and nucleotide diversity in the African and European population, for
1050 neutral, weakly deleterious, and beneficial mutations separately, calculated for windows of 500
1051 bp.

Mutation type	Model	F_{ST}		Nucleotide diversity			
		Mean	SD	AFR (Mean)	AFR (SD)	EUR (Mean)	EUR (SD)
Neutral	**Neutral**	0.0374	0.0226	0.0317	0.0070	0.0298	0.0071
	Purifying selection (rescaled)	0.0689	0.0556	0.0056	0.0022	0.0046	0.0021
	Purifying selection (not rescaled)	0.0363	0.0276	0.0057	0.0021	0.0054	0.0021
	Weak positive selection	0.0594	0.0479	0.0055	0.0022	0.0047	0.0021
	Strong positive selection	0.1065	0.1815	0.0002	0.0002	0.0002	0.0002
	Neutral bottleneck	0.2828	0.0762	0.1392	0.0145	0.0633	0.0219
Weakly deleterious	**Purifying selection (rescaled)**	0.0621	0.0457	0.0062	0.0021	0.0045	0.0019
	Purifying selection (not rescaled)	0.0361	0.0245	0.0063	0.0021	0.0059	0.0021
	Weak positive selection	0.0540	0.0369	0.0062	0.0021	0.0045	0.0020
	Strong positive selection	0.1489	0.2059	0.0004	0.0003	0.0003	0.0003
Beneficial	**Weak positive selection**	0.0385	0.0616	0.0004	0.0005	0.0004	0.0005
	Strong positive selection	0.3139	0.2952	0.0004	0.0003	0.0002	0.0003

1052

1053

1054

1055

1056

1057

1058

1059 **Supp Table 9**: Average nucleotide diversity (π) per site of African and European populations
1060 simulated for F_{ST} analyses.

Model	Population	π	
		Mean	Standard deviation
Neutral	African	0.1618	0.0086
	European	0.1538	0.0094
Weak positive selection	African	0.1416	0.0062
	European	0.1175	0.0161
Strong positive selection	African	0.0378	0.0128
	European	0.0281	0.0104
Purifying selection (DFE rescaled)	African	0.1131	0.0060
	European	0.0869	0.0095
Purifying selection (DFE not rescaled)	African	0.1079	0.0050
	European	0.0997	0.0054
Neutral bottleneck	African	0.1851	0.0026
	European	0.0841	0.0159

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

Supp Table 10: Mean genome-wide and outlier F_{ST} values for European and African *D. melanogaster* populations (expanded from Table 2) - for window sizes of both 10 and 20 SNPs, and the 1% and 2.5% tails of the distribution.

Evolutionary model	Window size	Mean F_{ST}	Outlier F_{ST} (2.5%)	Outlier F_{ST} (1%)	Outlier F_{ST} (2.5%) / Mean F_{ST}	Outlier F_{ST} (1%) / Mean F_{ST}
Neutral	10 SNPs	0.0363	0.1585	0.1628	4.3662	4.4838
	20 SNPs	0.0369	0.1411	0.1471	3.8203	3.9844
Purifying selection (rescaled DFE)	10 SNPs	0.0613	0.2497	0.2637	4.0707	4.2999
	20 SNPs	0.0646	0.2234	0.2234	3.4564	3.4564
Purifying selection	10 SNPs	0.0352	0.1467	0.1454	4.1673	4.1296
	20 SNPs	0.0361	0.1311	0.1311	3.6288	3.6288
Neutral bottleneck	10 SNPs	0.2557	0.6459	0.6956	2.5260	2.7204
	20 SNPs	0.2681	0.6071	0.6560	2.2650	2.4472
With weak positive selection	10 SNPs	0.0523	0.2256	0.2412	4.3154	4.6150
	20 SNPs	0.0556	0.1977	0.1977	3.5525	3.5525
With strong positive selection	10 SNPs	0.1976	0.7455	0.7455	3.7731	3.7731
	20 SNPs	0.2617	0.6498	0.6498	2.4832	2.4832

1084
1085 **Supp Figure 1**: Shown above is the specific DFE of deleterious mutations assumed in this study
1086 (from Johri *et al.* 2020), color-coded to display the probability of fixation of deleterious
1087 mutations relative to the probability of fixation of neutral mutations. Mutations with selective
1088 effects shown in blue regions have an appreciable probability of fixation, while those shown in
1089 red are highly unlikely to fix (scale is displayed on right).
1090

1091
1092 **Supp Figure 2**: Reduction in neutral diversity from the selected site, post-fixation of a mutation,
1093 with $2Ns_a = 5$ and (a) $h = 0$ and (b) $h = 1$. Solid lines represent mean values of 100 replicates,
1094 shaded regions correspond to 1 SE above and 1 SE below the mean. Crosses correspond to
1095 simulations based on equations (27) of Tajima (1990).
1096
1097
1098

1099
Supp Figure 3: Allele frequencies of SNPs in European (population 1) *vs* African (population 2)
populations - under the parameters inferred by Arguello *et al.* (2019), and where the selective
effects of all mutations are not re-scaled after the split (thus selection is stronger in the African
population). Simulated genomic elements experience purifying selection given by the DFE
inferred in Johri *et al.* (2020). Colored open circles represent outliers. Green depicts effectively
neutral mutations while warm colors depict deleterious mutations, with red representing weakly
deleterious mutations. *Left panel*: Allele frequency plots are shown for 10 (out of 100) replicates.
Right panel: The distribution of fitness effects of outlier mutations displaying the mean and
standard deviation for all 100 replicates.
1109
1110

1111

Supp Figure 4: The relationship between F_{ST} and nucleotide diversity calculated for neutral mutations alone, for the African and European populations simulated under six scenarios: (a) neutrality, (b) purifying selection (rescaled DFE), (c) purifying selection (not rescaled DFE), (d) purifying selection and weak positive selection, (e) purifying selection and strong positive selection, and (f) the neutral bottleneck of Li & Stephan 2006. On each, the upper left corner gives mean F_{ST} values with the SD in parentheses. The upper right corners give the corresponding mean value of π with the SD in parentheses. The red line represents the best fitting linear regression. Note the difference in scale between scenarios.

1120

Supp Figure 5: The relationship between F_{ST} and nucleotide diversity calculated for weakly
deleterious mutations alone, in the African and European populations simulated under four
scenarios: (a) purifying selection (rescaled DFE), (b) purifying selection (not rescaled DFE), (c)
purifying selection and weak positive selection, and (d) purifying selection and strong positive
selection. On each, the upper left corner gives mean F_{ST} values with the SD in parentheses. The
upper right corner gives the corresponding mean value of π with the SD in parentheses. The red
line represents the best fit linear regression. Note the difference in scale between scenarios.

1128

Supp Figure 6: The relationship between F_{ST} and nucleotide diversity calculated for only beneficial mutations in the African and European populations simulated under two scenarios: (a) purifying selection and weak positive selection, and (b) purifying selection and strong positive selection. On each, the upper left corner gives mean F_{ST} values with the SD in parentheses. The upper right corner gives the corresponding mean value of π with the SD in parentheses. The red line represents the best fit linear regression. Note the difference in scale between scenarios.

1135