1 **Microbial function is related to behavior of an invasive anuran**
2

Jia Zhou[1], Tiffanie Maree Nelson[2], Carlos Rodriguez Lopez[3], Shao Jia Zhou[1,4], Georgia
Ward-Fear[5], Katarina C. Stuart[6], Lee Ann Rollins[6]

[1]School of Agriculture, Food and Wine, University of Adelaide, Waite Campus, PMB 1 Glen
Osmond, South Australia, Australia, 5064.

[2]Queensland Facility for Advanced Bioinformatics, School of Medicine, Menzies Health
Institute Queensland, Building G40, Level 9, Gold Coast Campus, Griffith University,
Southport, QLD, Australia, 4215.

[3]Environmental Epigenetics and Genetics Group, Department of Horticulture, College of
Agriculture, Food and Environment, University of Kentucky, Lexington, KY, USA, 40546.

[4]Robinson Research Institute, University of Adelaide, South Australia, Australia, 5005

[5] Department of Biological Sciences, Macquarie University, North Ryde, NSW, Australia,
2109.

[6]Ecology & Evolution Research Centre, School of Biological, Earth and Environmental
Sciences, University of New South Wales, Biological Sciences South (E26) UNSW,
Kensington, Australia, 2052

Running title: Microbial function relates to host behavior

Corresponding author: Jia Zhou (jia.zhou@adelaide.edu.au)

Abstract

Invasive species cause negative environmental and economic impacts worldwide. Their

management may be improved by clarifying the role of behavior in advancing invasions. Gut

microbial communities are known to affect behavior of wild populations, but their impact on

behavior underlying invasiveness remains unexplored. Invasive populations of the cane toad

(*Rhinella marina*) in Australia have expanded across the continent and exhibit variation in

behavioral traits along their expansion trajectory, making this an ideal system to investigate

the relationship between gut microbes and behaviors. We collected wild female toads from

six locations in Queensland (n = 30) and Western Australia (n = 30), and conducted simple

tests on behavioral traits previously associated with invasion ability. We investigated the

relationships between toad gut microbiota, behavioral traits and the presence and intensity of

co-introduced lungworms (*Rhabdias pseudosphaerocephala*) in toad samples from both ends

of their Australian range. Based on 16S rRNA sequencing data, we found that microbiota in

cane toad colons were dominated by the phyla Bacteroidetes, Proteobacteria, Firmicutes,

Verrucomicrobia, and Fusobacteria. We found significant differences in microbiota

composition (p-value < 0.001) between regions and in predicted microbial functional groups

(p-value = 0.002). The occurrence of lungworms was strongly associated with variation in

both microbial composition and microbial functions. However, the behavioral traits were

associated with microbial functional variation, but not microbial compositional variation.

These results support the "holobiont concept" (investigating the assemblage associated with a

host) to fully understand drivers of invasion and highlight the need for experimental

manipulations to detect causal relationships between microbiota, parasites and host behavior.

Keywords: gut microbiota, invasive behavior, Cane toad, 16S rRNA, predicted

microbial functions, parasite

Introduction

An animal's behavior can be influenced by the bacterial microbiome within its intestines. To

understand this, scientists have experimentally manipulated the microbiome in host species.

For example, gut microbiota transfer has been found to modify exploratory behavior of germ-

free mice to resemble that of the donors (Bercik et al., 2011), and to influence emotional

reactivity in Japanese quail (*Coturnix japonica*; Kraimi et al., 2019). Further, altering gut

microbiota by treatment with a sterile diet or with antibiotics changed chemical social cues

and triggered aggressive behavior in leaf-cutting ants (*Acromyrmex echinatior*; Teseo et al.,

2019). Remarkably, host behavioral changes can be due to only an individual bacterial

species. Mono-colonization with one dominant species, *Lactobacillus brevis*, can decrease

walking speed and daily activity of germ-free or antibiotic-treated fruit flies (*Drosophila*

melanogaster; Schretter et al., 2018). Dietary administration with the probiotic *Lactobacillus*

rhamnosus IMC 501 to zebrafish (*Danio rerio*) caused shoaling behavior to be more uniform

and exploratory than that of the control groups (Borrelli et al., 2016) and administration of

Bifidobacteria reduced stress-related behavior of BALB/c mice (Savignac, Kiely, Dinan, &

Cryan, 2014). The gut microbiota can contribute to host adaptability (Hauffe & Barelli, 2019)

through mating choices (Sharon et al., 2010) and food resource usage (Vogel et al., 2017).

For example, gut microbiota influenced foraging decisions in fruit flies (*Drosophila*

melanogaster; Wong et al., 2017) and were associated with different behavioral tasks, such as

foraging, food processing and nursing in honey bees (Jones et al., 2018).

Empirical evidence for the role of intestinal microbiota on host behavior is increasing, yet to

date have focused primarily on human or domesticated models. Studying wild models in-situ

can elucidate the microbial functional value of this biological phenomenon, particularly how

it relates to host fitness and subsequent ecological or evolutionary processes (Davidson,

Raulo, & Knowles, 2020). This also allows us to understand the impact of microbes on host

invasive behavior (Stuart, Shine, & Brown, 2019), particularly when encountering novel

environmental stressors (Sampson & Mazmanian, 2015). While this idea is not well-tested,

there is evidence that native and invasive populations may differ in their microbiota in

unexpected ways. For example, despite sharing the same environment, microbial

communities in the hindguts of invasive Asian silver carp (*Hypophthalmichthys molitrix*)

differ significantly from those in indigenous planktivorous gizzard shad (*Dorosoma*

cepedianum). The invasive species has much higher abundance of the order *Bacteroidales*,

with the genera *Bacillariophyta* and *Clostridium* being significantly more abundant in native

than invasive fish (Ye, Amberg, Chapman, Gaikowski, & Liu, 2014). It is not clear how the

observed differences in gut bacteria contribute to invasiveness, especially the comparison of

gut microbiota regarding host invasiveness has only been conducted in two different species

(Fontaine & Kohl, 2020a; Qu et al., 2020). Research on gut microbiota variation between

populations with different invasion characters in the same species would help to elucidate

such patterns.

Host behavior appears to be more closely linked to the microbiota of the colon rather than

that of the small intestine. From a mechanistic perspective, it is known that the microbial

community in the colon can ferment complex carbohydrates, such as dietary fibre, into short

chain fatty acids (SCFAs) that are known to have neuroactive properties (Dinan, Stilling,

Stanton, & Cryan, 2015). Certain microbial species in the colon can cause changes to brain

metabolites: in pigs, *Ruminococcus* spp. can influence brain N-acetylaspartate through the

mediation of serum cortisol (Mudd, Berding, Wang, Donovan, & Dilger, 2017). In the cane

toad, reactive nerve cell bodies (which may be caused by the different microbial

communities) were observed to be common in the submucosa of colon and less common in

small intestine (Z. S. Li, Furness, Young, & Campbell, 1992). In addition to these explicit

links to behavior, there is evidence that colon microbiota may have a greater effect on hosts

than microbiota in other parts of the digestive system. The 'stationary phase' is the point at

which the maintenance of the bacterial population size reaches equilibrium. Following

regular nutrient provisioning, microbial communities replicate quickly and reach the

stationary phase after only 20 minutes in the colon as compared to more than 2.2 hours in

other organs of the digestive system. Microbiota also have longer transit times in the colon

(10-60 hours versus less than 3 hours in all other digestive organs; Fetissov, 2017), increasing

their opportunity to impact their host.

Intrinsic factors such as genetics, age, sex and parasite load can interact to maintain and

shape the host's gut microbiota community (Kozik, Nakatsu, Chun, & Jones-Hall, 2017; Org

et al., 2016; Wang et al., 2016). In mice, significantly different microbiota are observed in

narrow age ranges, such as between 4-5 weeks and 6-7 weeks (Kozik et al., 2017). Males and

females can also possess different microbiota (in mice: Kozik et al., 2017, and cane toads:

Zhou et al., 2020). Further, host health is associated with gut microbiota and the presence of

some intestinal parasites can be predicted by studying gut microbial composition. For

example, *Blastocystis* colonization is negatively associated with *Bacteroides* (Stensvold &

van der Giezen, 2018). Because parasites are also known to impact host behaviors (Herbison,

2017; Klein, 2003), it is important to consider interactions of parasites and gut bacteria host

behavioral variation.

Extrinsic environmental factors such as habitat and diet also influence the composition and

variation of gut microbiota (Bletz et al., 2016; Carmody et al., 2015; Marques et al., 2010).

Amphibians are good models to study these changes because they utilize both aquatic and

terrestrial habitats and their skin is permeable, making them highly sensitive to environmental variation (Hopkins, 2007). For example, the gut microbiota of fire salamander (*S. salamandra*) larvae differs depending on whether they live in ponds or streams (Bletz et al., 2016). Translocating animals to the opposite habitat type shifts their gut microbiota to resemble that of residents. Further, gut microbiomes can be similar across phylogenetically distinct, but sympatric amphibian species (such as *Fejervarya limnocharis* and *Babina adenopleura* inhabiting both farmlands and forests) and mirror environmental conditions (in both species bacterial diversity was higher in farmland populations; Huang, Chang, Huang, Gao, & Liao, 2017). Because diet is a key environmental factor affecting gut microbial community assemblages, these changes are likely to be influenced by differences in diet across environments (Scott, Gratz, Sheridan, Flint, & Duncan, 2013; Zmora, Suez, & Elinav, 2019). In summary, both environmental factors and host factors should be considered when studying intestinal microbiota variation.

Previously, drivers of variation in invasion-related behaviors have been studied, including genetics, morphology, habitat, diet, prior experience and parasites (Brown, Kelehear, Pizzatto, & Shine, 2016; Child, Phillips, & Shine, 2008; Hudson, Brown, & Shine, 2017; Selechnik et al., 2019; Stuart et al., 2019). However, there have been no investigations of gut microbiota as a potential driver of behavioral shifts across an invasive range in any species. Cane toads are an ideal system for these investigations because they display significant differences across Australia in behaviors likely to be related to increased capacity for dispersal ability; invasion front toads are bolder and more exploratory (Gruber, Brown, Whiting, & Shine, 2017; Stuart et al., 2019). A range of differences in dispersal-related morphology and performance metrics also have been shown: for example, wild-caught invasion-front toads have longer legs and toads with longer legs move greater distances (Benjamin L Phillips, Brown, Webb, & Shine, 2006) and toad morphology and performance

are linked (Hudson, Vidal-García, Murray, & Shine, 2020), endurance is greater in invasion-

front adult toads held in captivity (Llewelyn, Phillips, Alford, Schwarzkopf, & Shine, 2010),

invasion-front toads move in a more consistent direction than those from the range-core

(Brown, Phillips, & Shine, 2014). However, the relationships between these traits are

complex and may be affected by factors such as sex (Hudson et al., 2020) or rearing

environment (Stuart et al., 2019). Understanding how microbiota may contribute to invasion

success may be key to successful management of these species.

Here, we investigated the intestinal microbiome of invasive cane toads in Australia to

determine whether there were significant differences in the identity or relative composition of

gut microbiota in cane toads from range-core (i.e. the introduction site) versus invasion-front

populations. We predicted that extrinsic and intrinsic factors would impact microbial

communities, including environmental conditions and host parasite prevalence and intensity.

We also investigated how microbial variation correlated with cane toad behavior. Given our

knowledge of differences in cane toad behavior across the Australian range, we predicted that

the microbiota of animals captured from the invasion-front populations would differ to that of

the range-core populations, and that this would be correlated with their behavior. We discuss

the potential implications of these impacts on invasion ability in this iconic invader.

Methodology

Study species, sample collection and behavioral assays

Cane toads (*Rhinella marina*) are large anurans (measuring up to 23 cm, 1.25 kg) introduced

into Australia in 1935 as a biocontrol for pests of sugar cane crops (Shine, 2010). They are

actively invading northern Australia and critically threaten many native predator species who

are fatally poisoned when they attempt to eat them (Shine, 2010). We hand captured 60 wild

adult females from three sites at the invasion front edge in Western Australia ('invasion-

front': Kununurra 15.776566° S, 128.744293° E, Old Theda 14.790795° S, 126.497624° E,

Mary pool 18.72528° S, 126.870096° E. N=30, collected November 2018) and three sites in

Queensland (QLD) ('range-core', Rossville 15.697069° S, 145.254385° E, Croydon

18.207536° S, 142.245702° E, Lucinda 18.530149° S, 146.331264° E. N=30, collected

December 2018) for this study. The University of Adelaide Animal Ethics Committee

approved all animals that had been used in this research (approval number: S-2018-056).

We conducted brief behavioral assays while collecting toads based on the methodologies

described in a common-garden experiment on this species (Stuart et al., 2019) including: (1)

struggle score (number of arm or leg kicks after being picked up until toad remains still for 5

seconds and struggle likelihood) and (2) righting behavior (within two minutes after the toad

is placed on its dorsal side: time to right itself, number of limb kicks ('righting effort'), and

righting effort likelihood). In that common-garden experiment, righting time was correlated

with speed (r^2=0.09, p=0.004) and stamina (r^2=0.07, p=0.01), and righting effort was

correlated with speed (r^2=0.05, p=0.02) after body size correction (Stuart, pers. comm.); these

data demonstrate the relevance of the simple assays used here to traits more clearly linked to

invasion potential in this species. After behavioral assays were conducted, we placed the

animals into individual, moist, calico bags and weighed, measured (snout urostyle length;

SUL) and euthanised them by injecting tricaine methanesulfonate (MS222) buffered with

bicarbonate of soda to balance pH. After euthanasia, we used a clean surface and clean tools

to dissect each animal. From each animal, we preserved 0.3cm of colon near the cloaca

(including gut content) in 95% ethanol. We also collected data on two toad parasites: an

encysted gut parasites (*Physalopterinae*: physalopterine larvae; Kelehear & Jones, 2010) and

the co-introduced toad lungworm, which passes into the toad gut as eggs and then hatches

into larvae in the gut (Baker, 1979). We examined the gut for evidence of the former. To

investigate the latter, we dissected and inverted the lungs (right lobe when facing the ventral

side) and counted lungworms. Body size and parasite data are hereafter collectively referred

to as 'host characteristics'.

Analyzing morphological and ecological data

We compared host characteristics, host behavioral traits and environmental factors between

range-core and invasion-front toads. Within each of these data sets, we conducted correlation

tests to exclude highly corrected variables for downstream analyses. Because body length

(SUL) and weight were significantly correlated, we only included SUL in further analyses.

For host characteristics and behavioral traits, we used SUL as a covariate in generalized

linear models (GLM). We compared count and occurrence data (i.e. absence=0, presence=1)

for host parasite infection and behavioral traits. To analyze occurrence data, we used GLM

with logistic regression (R version 4.0.2, R Core Team, 2020) and with negative binomial

regression *glm.nb* in the "MASS" package (Venables & Ripley, 2002) to analyze over-

dispersed count data, when the conditional variance exceeded the conditional mean. We

obtained environmental data for 19 bioclimatic variables for each sampling site from

WorldClim (Table S1, Fick & Hijmans, 2017). We analyzed these data using the Shapiro-

Wilk test to assess multivariate normality using "RVAideMemoire" package (Herve, 2020)

and then we used a nonparametric Wilcoxon signed rank test to compare the median of the

difference between groups because the data were not normally distributed.

DNA extraction, library preparation and sequencing

We extracted microbial DNA from colon contents following the manufacturer's protocol for

the DNeasy PowerSoil kit (Qiagen) and prepared 16S rRNA amplicon libraries by following

guidelines for the Illumina MiSeq System. We used Zymo isolated DNA (D6305) as a

community positive control and MilliQ water as a PCR negative control to identify

environmental contamination during the library preparation process (Eisenhofer et al., 2019).

Libraries were prepared based on the hypervariable (V3-V4) region of the 16S rRNA gene

from each sample using primers 341F (5' –

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG-

3') and 785R (5'-

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAA

TCC-3') (Herlemann et al., 2011). The library preparation details have been reported

previously (Zhou et al., 2020). The libraries were sequenced at the Ramaciotti Centre for

Genomics (University of New South Wales, Kensington, Sydney) on the Illumina MiSeq

platform targeting 2x300bp paired-end sequence reads.

Identifying and classifying bacteria community

We downloaded demultiplexed FASTQ data from Illumina's BaseSpace cloud storage, then

created an amplicon sequence variants (ASVs) abundance table using the open-source

QIIME2 pipeline (Bolyen et al., 2018). Demultiplexed sequence counts from samples and the

positive control ranged between 138,989 and 305,662; the count of the PCR negative control

was 67,506. The DADA2 pipeline (Callahan et al., 2016), implemented in QIIME2, was used

to filter and trim the first 20 bases from each read and truncate sequences to 200 bases. The

remaining sequences were dereplicated, then forward/reverse reads were merged, chimeras

were removed, and finally ASVs were generated for downstream analysis (Callahan et al.,

2016). After quality filtering, reads from colon samples and the positive control ranged

between 103,245 and 245,059 counts; the PCR negative control yielded 6,727 reads. The

taxonomic assignment of ASVs was performed using Greengenes version 13_8 (DeSantis et

al., 2006). Data were pruned to remove representatives classified to Archaea (N = 28),

chloroplast (N = 17), mitochondria (N = 186), and 151 unassigned ("kingdom") ASVs

implemented in the package 'phyloseq' (McMurdie & Holmes, 2013). We also removed the

ASVs with prevalence of less than four, which makes the logged counts per sample more

evenly distributed. The remaining 9,878 taxa were classified to the Kingdom Bacteria with

62.62% assigned to phylum level and 39.65% assigned to family level.

Characterizing diversity and microbial function of bacteria communities

We calculated observed ASVs (DeSantis et al., 2006), evenness (Pielou, 1966), and Shannon

(Shannon, 1948) indices through QIIME2 for alpha diversity (a measure of diversity within

individuals). We used a Wilcoxon signed-rank test to compare Shannon Index of toad

microbiomes from different locations/sites and used a GLM model to check the correlation

between Shannon Index and locations/sites with consideration of SUL as a covariate.

In order to explore the bacterial taxa that may contribute to the observed differences in

community, the Core50 gut community (Bletz et al., 2016) was calculated as follows: the

ASV table was filtered to include only the ASVs present in a minimum of 50% of individual

toads from each site. This calculation was performed separately for three sites from invasion-

front toads: Kununurra (gut Core50: n = 111 ASVs), Old Theda (gut Core50: n = 118 ASVs),

and Mary Pool (gut Core50: n = 129 ASVs); three sites from range-core toads: Rossville (gut

Core50: n = 148 ASVs), Croydon (gut Core50: n = 86 ASVs), and Lucinda (gut Core50: n =

117 ASVs). Then we subsequently compiled filtered ASVs of six sites to avoid excluding

ASVs that may be specific to only one site. In combination, the gut Core50 contained 325

unique ASVs. We visualized relative abundance of bacteria (abundance >2%) in different

sample types, classified to the phylum and family level.

To compare microbial communities between samples, we used a Hellinger transformation

implemented in the package "microbiome" (Valverde, Makhalanyane, & Cowan, 2014) in R

and then calculated and generated a weighted UniFrac dissimilarity matrix, which was

visualized using Principal Coordinate Analysis (PCoA) plots using commands from the

package "Phyloseq" (McMurdie & Holmes, 2013). We used the *adonis* command from the

package "Vegan" to perform permutational multivariate analysis of variance (perMANOVA)

to check whether the microbial communities of toads from each location were significantly

different. We used the command *betadisper* in the package "Vegan" (Oksanen et al., 2019) to

check the homogeneity of group variances, an assumption of perMANOVA. After finding

significant differences between invasion-front and range-core toads, we performed pairwise

comparisons between six sites from both locations using the command

pairwise.perm.manova function in "RVAideMemoire" package with the Wilks Lambda

(Nath & Pavur, 1985) and corrections for multiple testing (Herve, 2020) were conducted

using the Hochberg procedure (Hochberg, 1988).

We used PICRUST2 (Douglas et al., 2019) to predict microbial functions for Core50 ASVs

and generated pathway abundances considering the taxonomic contributions of ASVs. We

removed all pathways with less than 0 prevalence and 474 out of 484 pathways remained for

analysis. Then we used a Hellinger transformation and calculated and plotted Bray-Curtis

dissimilarity matrix through package "Phyloseq". We performed permutational multivariate

analysis of variance (perMANOVA) to check whether the pathway abundance of toads from

each location and site were significantly different and we used the command *betadisper* to

check the homogeneity of group variances.

Correlating intrinsic and extrinsic factors with host microbiota

To identify the association of individual host characteristics, host behavioral traits and

environmental factors with bacterial community and predicted microbial functions, statistical

analysis was conducted using the function envfit in the package "Vegan" on Bray Curtis

dissimilarities. Bray Curtis dissimilarities were visualized with non-metric multidimensional

scaling (nMDS) plots. Host characteristics and behavioral traits were fitted to the ordination

plots using the function envfit. Environmental variables that were significantly associated

with microbiota taxonomy and microbial function were also fitted to the ordination plots

using the function envfit and smooth surfaces of the same environmental factors were added

on the nMDS plot with the ordisurf function in package "Vegan".

After analysing all variables individually, we also conducted distance-based redundancy

analysis (dbRDA) combining all host characteristics and behavioral traits to identify

relationships between variables with respect to their impact on gut microbiome. dbRDA

performs constrained ordination directly on a distance or dissimilarity matrix with the

function *capscale* in the "Vegan" package in R. We checked the correlation between each

pair of traits. Bray-Curtis dissimilarity matrixes of ASVs and predicted microbial functions

were ordinated and the results were analysed using redundancy analysis with constraining

variables that were not highly correlated (correlation coefficient < 0.85 and > -0.85) to

estimate their explanatory proportion. We did not include environmental factors in the

capscale analysis because these data are not independent (toads from the same site have the

same environmental factors).

Identifying differences in microbial taxa and their predicted microbial functions in range-

core versus range front toads

To identify the significant differences in ASVs and predicted microbial functions between

range-core and invasion-front toads, we performed differential abundance testing using the

function *DESeq* in the package "DESeq2" (Love, Huber, & Anders, 2014). We report

significant differences as the log fold change of a taxa in range-core toads versus those from

the invasion-front. We set ASVs and microbial functions abundance in invasion-front toads

as the control group. Results for all statistical tests were considered significant where p-

values <0.05.

Results

Morphological characteristics and parasite load in range-core versus invasion-front toads

The SUL (body length) of wild-caught invasion-front toads was longer than that of range-

core toads (edge: mean = 103.51mm, SD = 9.42; core: mean = 96.24mm, SD = 12.5; t = -

2.543, df = 53.897, p-value = 0.014). There was no significant differences in lungworm and

gut parasite prevalence or intensity between the range-core and invasion-front toads we

collected (Tables 1, S2): in all sampled toads, we report the number of lungworms (mean =

2.56, SD = 5.37), the occurrence of lungworms (mean = 0.52, SD = 0.50), the number of gut

parasites (mean = 3.26, SD = 5.35) and occurrence of gut parasites (mean = 0.5, SD = 0.51).

Behavioral traits of range-core versus invasion-front toads

Range-core toads were found to have higher struggle scores when compared with invasion-

front toads (Table 1, S2). We found range-core toads were more likely to struggle than

invasion-front toads (core: mean = 0.83, SD = 0.37; edge: mean = 0.27, SD = 0.43; p-value =

0.008) and, in those who did struggle, the number of struggle movements was significantly

higher for range-core toads than for invasion-front toads (core: mean = 2.63, SD = 2.52;

edge: mean = 2.1, SD = 4.54; p-value = 0.002). The righting effort likelihood was higher for

range-core toads than for invasion-front toads (core: mean = 0.73, SD = 0.45; edge: mean =

0.5, SD = 0.51; p-value = 0.036) (Tables 1, S2). The righting effort and righting time did not

differ between range-core and invasion-front toads (Table 1, S2): across all sampled toads,

we report the righting effort (mean = 2.2, SD = 3.56) and righting time (mean = 18.88sec, SD

= 41.32).

Environmental conditions experienced by range-core and invasion-front toads

Cane toads at the invasion-front experienced significantly hotter (six temperature

measurements, see Table 1, S2 for statistics) and dryer (seven precipitation measurements,

see Table 1, S2 for statistics) conditions year round than those in the range-core. Moreover,

cane toads at the invasion-front experienced more changeable weather than those in range-

core (three temperature variation measurement and one precipitation variation measurement,

see Table 1, S2 for statistics). However, one temperature measurement (min temperature of

coldest month: mean = 14.65, SD = 1.77) and one temperature variation measurement

(isothermality, higher isothermality refering to more variable temperature: mean = 55.17, SD

= 2.29) were not different in both environments included here (Table 1).

DNA sequence data quality

The number of observed species in the PCR negative control (N = 36) was much lower

compared to that of toad colon samples (mean = 336, SD = 137.8), which indicated minimal

contamination throughout the library preparation processes (Table S3). The number of

observed species (N = 19) in the community positive control was also low compared to

experimental samples. The eight species used as microbial community standards constituted

more than 99.8% of ASVs from our sequenced community positive control, as shown in

Figure S1, indicating minimal PCR bias in our library preparation process.

Comparison of microbial composition and diversity within individuals

The alpha diversity (diversity within individual toads) was not significantly different between

range-core and invasion-front toads: observed species (*wilcox.test*, p-value = 0.631), Pielou's

evenness (*wilcox.test*, p-value = 0.307), Shannon (*wilcox.test*, p-value = 0.230). Lucinda had

the highest Shannon diversity, followed by Old Theda, Mary Pool, Rossville, Croydon and

Kununurra (Figure S2B). Shannon diversity in individual samples was not correlated with

measured behavioral traits (*glm*, covariate = SUL, p-values > 0.050).

Relative abundance of core50 microbiota in large intestine

A total of 89.23% ASVs were assigned to the level of phylum and dominant phyla (average

abundance > 2%) were Firmicutes, Bacteroidetes, Proteobacteria, Fusobacteria, and

Verrucomicrobia (Figure S3, Table S4). We discuss the microbial functional significance of

these groups in the discussion. Across all individuals, we found that: (1) the most dominant

phyla were Firmicutes (average abundance = 47.89%, SD = 19.25%) and Bacteroidetes

(average abundance = 39.05%, SD = 19.26%) (Table S4); (2) the phylum Bacteroidetes were

consistently present in most individuals, except for two toads (Toad 14, 21) from the

invasion-front and two toads (Toad 43, 48) from the range-core (Figure 1); and (3) there was

one invasion-front toad (Toad 21) that showed a higher abundance of Proteobacteria

(77.35%) when compared with other individuals (average abundance = 5.13%, SD = 7.99%).

A total of 70.77% ASVs were assigned to the family level and dominant families (average

abundance > 2%) included Bacteroidaceae, Lachnospiraceae, Porphyromonadaceae,

Clostridiaceae, Enterobacteriaceae, Erysipelotrichaceae, Fusobacteriaceae, Bacillaceae,

Ruminococcaceae, Verrucomicrobiaceae, Turicibacteraceae, and Veillonellaceae (Figure S4,

Table S5). Toad 21, who had a high abundance of Proteobacteria, also had a higher

abundance of Enterobacteriaceae (77.34%) when compared with other individuals (average

abundance = 4.55%, SD = 7.86%) (Figure S5).

A total of 40.62% ASVs were assigned to the genus level and dominant genera (average

abundance > 2%) included *Bacteroides*, *Parabacteroides*, *Clostridium*, *Epulopiscium*,

Akkermansia, *Turicibacter*, *Bacillus* (Figure S6, Table S6).

Comparison of microbial community taxonomy and microbial function between

individuals with Core50 microbiota

Microbial taxonomic communities were significantly different between locations (Figure 2A,

$R^2 = 0.095$, F = 6.092, p-value < 0.001). Microbial communities were marginally different

between most sites (p-values < 0.050), except that Old Theda was not significantly different

from Kununurra, Mary Pool, Croydon and Lucinda (Figure 2A). Among 230 ASVs that

assigned to family level, there were 124 ASVs that significantly differed between range-core

and invasion-front toads' colon (Table S7). The number of significantly different ASVs in

each phylum were: Bacteroidetes (60 ASVs), Firmicutes (55 ASVs), Proteobacteria (7

ASVs), Actinobacteria (1 ASVs), Verrucomicrobia (1 ASV) (Table S7, Figure 4A).

Among the identified 474 predicted microbial functions, we found significant differences

between invasion-front and range-core toads (Figure 2B; $R^2 = 0.064$, F = 4.110, p-value=

0.002). Pairwise tests between sampling sites indicated that toads from Kununurra had

significantly different microbial functions to toads from Rossville (p-value= 0.009) and

Lucinda (p-value = 0.046), and the other sites were not significantly different (p-values >

0.050) (Figure 2B). The top six most abundant pathways were pentose phosphate pathway

(non-oxidative branch), adenosine deoxyribonucleotides de novo biosynthesis II, guanosine

deoxyribonucleotides de novo biosynthesis II, pyruvate fermentation to isobutanol

(engineered), Calvin-Benson-Bassham cycle, and adenine and adenosine salvage III (Figure

3). There were 84 out of 474 microbialfunctions that were significant different between

invasion-front and range-core toads (Table S8, Figure 4B). Range-core toads have

significantly higher abundance of microbial function in superpathway of pyrimidine

ribonucleosides degradation (log2FoldChange = 5.98) and lower abundance of microbial

functions in phosphopantothenate biosynthesis III (log2FoldChange = -4.98), superpathway

of sialic acids and CMP-sialic acids biosynthesis (log2FoldChange = -4.89) and factor 420

biosynthesis (log2FoldChange = -4.72) than invasion-front toads (Table S8, Figure 4B). Out

of the top 30 abundant microbial functions, range-core have significantly lower abundance of

microbial function in urate biosynthesis/inosine 5'-phosphate degradation (log2FoldChange =

-0.10) than invasion-front toads (Figure 3, Table S8).

Associations between host, environment and intestinal microbiota

We assessed which host or environmental factors are strongly associated with microbiota

taxonomy and microbial function. When comparing the association between individual host

characteristics and behaviors with microbiota variation, only the occurrence of lungworms

was significantly associated with the microbial taxonomic community ($R^2 = 0.128$, p-value =

0.023) (Tables S9, S10; Figure S7A, B). Among the 21 environmental factors investigated

here, longitude ($R^2 = 0.194$, p-value = 0.001), isothermality ($R^2 = 0.193$, p-value = 0.002),

and mean temperature of driest quarter ($R^2 = 0.146$, p-value = 0.015) were strongly

associated with microbial taxonomic community. The mean temperature of driest quarter (R^2

$= 0.187$, p-value = 0.004) and isothermality ($R^2 = 0.175$, p-value = 0.007) were also strongly

associated with microbial functions (Table S10). Because longitude is a proxy for sampling

region, we only present mean temperature of driest quarter and isothermality in figure S7C,

D. Host characteristics and behaviors are independent measurements, yet environmental

factors are the same for all 10 toads from each site. The variable isothermality was identified

as having a possible correlation with microbial taxa and microbial functions at the range of

54.5 – 57.5 (Figures S7C, D). The contour lines of mean temperature of driest quarter show

less linear relationship than it does for isothermality with microbial taxa, which is further

supported by envfit results (Figure S7C). The contour lines of mean temperature of driest

quarter show the strongest linear relationship with microbial functions (Figure S7D).

However, the deviance explained by this ordisurf model is the lowest (18.3%) compared to

other three ordisurf models (isothermality x microbial taxa, isothermality x microbial

functions, and mean temperature of driest quarter x microbial taxa).

We calculated correlations between the factors that were significantly associated with

microbial taxa and microbial functions as: (1) the occurrence of lungworms is correlated with

isothermality (t = 2.3234, df = 58, p-value = 0.02369, coefficient = 0.292, method =

"pearson"), (2) the occurrence of lungworms is not correlated with mean temperature of the

driest quarter (t = 1.0534, df = 58, p-value = 0.2965, coefficient = 0.137), (3) isothermality is

highly correlated with mean temperature (t = 8.3496, df = 58, p-value = 1.594e-11,

coefficient = 0.739).

We used a redundancy analysis to assess the association of all host variables. The best model

which was associated with the most variation in the microbiota taxonomic community

included only the occurrence of lungworms (AIC = 178.58), whereas the best model which

was associated with the most variation in predicted microbial functions included righting

effort likelihood (AIC = 53.613), the occurrence of lungworms (AIC = 54.297) and righting

time (AIC = 56.912). The combination of these three factors was associated with 17.8% of

variation in the predicted microbial functions (Figure 5).

Discussion

Our research is the first of its kind to investigate how the gut microbiome of an invasive

species changes 1) as a function of intrinsic (host) and extrinsic (environmental) factors, 2)

with invasion context (established resident versus invading populations), and 3) importantly,

with relation to aspects of animal behavior that influence dispersal, and thus wider invasion

success. There is growing acknowledgement that microbiota influences animal behavior in

diverse ways (Davidson, Cooke, Johnson, & Quinn, 2018; Davidson et al., 2020) yet until

now, the links between microbiota and behaviors that influence invasiveness remain

unexplored. We found that whilst the alpha diversity in microbiota was similar in the

invasion-front and range-core individuals, there were significant differences in both the

composition and microbial function of gut microbiota in toad populations across the

Australian invasive range. Microbial functions better explain cane toad righting behavior than

does microbial composition. Interestingly, the presence of the co-introduced parasitic

lungworm and environmental factors related to temperature could contribute to the variation

of both gut microbial community composition and microbial functions, highlighting the

importance of studying how these factors interact.

Microbial taxa

Four phyla (Firmicutes, Bacteroidetes, Proteobacteria, and Fusobacteria) were found to be

dominant (Mean > 2%) in toad samples from both ends of the range, similar to results of a

previous study of cane toads from both of these regions held in captivity in a common facility

(Zhou et al., 2020). The previous three phyla are commonly observed as dominant in the

intestine of frogs and toads (Chang, Huang, Lin, Huang, & Liao, 2016; Kohl, Amaya,

Passement, Dearing, & McCue, 2014; Wiebler, Kohl, Lee, & Costanzo, 2018), which

suggests these three phyla may play important roles for maintaining the health of toads, and

that these taxa are not dependent on particular host or environmental conditions. Even though

not described as dominant phyla, Fusobacteria were found in the intestine of other frogs and

toads (Chai, Dong, Chen, & Wang, 2018; Kohl et al., 2014). Fusobacteria are dominant phyla

in the fish intestine (Colombo, Scalvenzi, Benlamara, & Pollet, 2015; Kohl et al., 2014) and

Fusobacteria (*Cetobacterium*) were previously reported to be associated with changes in

immunity of Cuban tree frogs (Knutie, Wilkinson, Kohl, & Rohr, 2017). Interestingly, we

found a higher abundance of the phylum Verrucomicrobia in range-core toads than in toads

from the invasion-front (core: mean = 3.98%, SD = 7.39%; edge: mean = 1.29%, SD =

2.62%) (Table S4, Figure S3). Species within the phylum Verrucomicrobia are commonly

found in the intestinal microbiota of frogs or toads (Jiménez & Sommer, 2017; J. Li et al.,

2019) and one dominant genus within this phylum in anurans is *Akkermansia* (Chai et al.,

2018; Kohl, Cary, Karasov, & Dearing, 2013), which is also identified in range-core wild-

caught toads (Figure S6, Table S6). One particular species (*A. muciniphila*) is common in the

human intestinal tract, specializing in the degradation of intestinal mucins (Derrien, Collado,

Ben-Amor, Salminen, & de Vos, 2008) and therefore often considered a to be associated with

a 'healthy' metabolic status in humans (Dao et al., 2016). The reduction in members of the

phylum Verrucomicrobia in invasion-front toads could be a result of an altered habitat or

different available dietary options. Previously, members of the phylum Verrucomicrobia

were not found to be dominant in the gut of captive toads from the invasion-front and range-

core (Zhou et al., 2020).

Range-edge cane toads, despite their ability to spread rapidly, are known to experience health

problems, such as severe spinal arthritis (Brown, Shilton, Phillips, & Shine, 2007). Emerging

research in humans has highlighted the association of the gut microbiome on inflammation

and the development of musculoskeletal disorders such as arthritis which has a significant

impact on mobility (Szychlinska, Di Rosa, Castorina, Mobasheri, & Musumeci, 2019). In the

study of cane toad skin microbiota, a higher proportions of bacteria with antifungal properties

(fungal pathogen: *Batrachochytrium dendrobatidis*-inhibitory) were detected in toads from

range-core than those from invasion-front, which known to be shaped by site-specific

pathogen pressures (Weitzman et al., 2019). Interestingly, even though cane toads are known

to eat their skins, the 14 core *Batrachochytrium dendrobatidis*-inhibitory bacteria found on

cane toad skin were not observed in its core gut bacteria, except family Enterobacteriaceae

(*Bufotyphonius*-inhibitory_9).

We found significant differences in the diversity and composition of microbial taxonomy

between range-core and invasion-front toads even though their dominant phyla and alpha

diversity were similar. ASVs in the family Veillonellaceae were found to be significantly

higher in invasion-front toads, compared to range-core toads (Figure 4A). ASVs in this

family have been found to influence host metabolic regulation. For example, in an

experiment that altered air temperature for Brandt's voles (*Lasiopodomys brandtii*); in colder

temperatures, voles which huddled, had a higher abundance of Veillonellaceae as well as

higher concentrations of short-chain fatty acids (SCFAs) in their intestines when compared

with non-huddling voles (Zhang et al., 2018). This family is known to produce SCFAs, such

as propionic acid (Gamage et al., 2017; Louis & Flint, 2017), and then contribute to increased

locomotor activity (Thomas et al., 2012). The relationship between the bacterial members of

this family and host metabolic regulation, suggests that invasion-front toads may use the

SCFAs to fuel their invasion by producing SCFAs, such as propionic acid and increasing

locomotor activity (Llewelyn et al., 2010). ASVs from another family Clostridiaceae

(Hugenholtz et al., 2018) that belongs to SCFA-producing bacteria were also higher in

invasion-front toads than those from the range-core. Furthermore, the family Veillonellaceae

may be associated with host sociality. A significant reduction of Veillonellaceae has been

observed in children with Autism Spectrum Disorder, often known for desiring social

isolation, when compared to a neurotypical group of children (Kang et al., 2013). Higher

abundance of Veillonellaceae in invasion-front toads could foster their "bolder" personality,

retaining a higher propensity for exploration and risk-taking (González-Bernal, Brown, &

Shine, 2014; Gruber et al., 2017).

Several other ASVs that differed across the cane toad's range have previously been related to

behavior. ASVs from family Peptococcaceae were found to be significantly higher in

invasion-front toads than in range-core ones (Figure 4A). Peptococcaceae have been found to

be related with host neurotransmitter (noradrenaline linking visual awareness to external

world events, (Gelbard-Sagiv, Magidov, Sharon, Hendler, & Nir, 2018): for example,

Peptococcaceae in the caecum showed a significantly positive correlation with noradrenaline

levels in mice (Houlden et al., 2016). ASVs from family Bacillaceae were found to be

significantly lower in invasion-front toads than range-core (Figure 4A). Bacillaceae have

been found to be related to host anxiety: for example, this species was found in higher

abundance in methamphetamine-treated rats versus the control (Ning, Gong, Xie, & Ma,

2017) and in higher abundance in excercised mice versus sedentary group (Choi et al., 2013).

Higher abundance of Bacillaceae could contribute to higher anxiety-like behaviors thus

heightening the stress response (Ning et al., 2017) and may be correlated with a decreased

exploratory behavior in new environments (Golla, Østby, & Kermen, 2020). In summary,

invasion-front toads possessed gut microbial communities that in other studies have been

associated with SCFAs production and neurotransmitters, suggesting the potential to increase

locomotor ability, visual awareness to external world events and propensity for exploration

and risk-taking. In comparison, range-core toads possessed taxa are associated with higher

anxiety-like behavior, suggesting decreased propensity to explore.

Predicted microbial functions

Variation in the predicted microbial functional groups observed between locations and cites

was less obvious than in community composition (Figure 2). Other studies have found similar

results, suggesting that microbial function is more likely to be preserved than composition

(e.g. in fire salamanders, Bletz et al., 2016). This could result from different gut microbiota

harboring similar microbial functions and may contribute to increasing resilience and

persistence of microbial functional stability of gut microbiota (Bletz et al., 2016; Huang et

al., 2017; Lozupone, Stombaugh, Gordon, Jansson, & Knight, 2012). For invasive species,

this can also contribute to its adaptive potential in novel environments. We found significant

differences in microbial functions between range-core and invasion-front toads even though

their most abundant microbial functions were similar. There are significant differences in

microbial functional pathways that contribute to the microbe food sources and metabolism.

As for microbe food sources, invasion-front toads have significantly lower abundance of

microbial function in the superpathway of pyrimidine ribonucleosides degradation. This

pyrimidine ribonucleosides degradation provides nitrogen source for microbes (West, 1996)

and plays an important role in perturbations in the uridine monophosphate (UMP)

biosynthetic pathways. This allows the bacterial cell to sense signals such as starvation,

nucleic acids degradation, and availability of exogenous pyrimidines, and to adapt the

production of the extracellular matrix to the changing environmental conditions (Garavaglia,

Rossi, & Landini, 2012). This may be related to the disappear of Verrucomicrobia as a

dominant taxon. As for microbe metabolism, invasion-front toads have a higher abundance of

microbial functions in factor 420 biosynthesis; cofactor 420 is critical to bacterial metabolism

and mediates a variety of important redox transformations involved in bacterial persistence,

antibiotic biosynthesis, pro-drug activation and methanogenesis (Bashiri et al., 2019).

Moreover, there are significant differences in microbial functional pathways that contribute

to host health. Invasion-front toads have higher abundance of microbial functions that are

beneficial to general host health and immunity: (1) phosphopantothenate biosynthesis, which

is the first step for the bacteria production of coenzyme A (CoA), an indispensable enzyme

cofactor for all living organisms (Sibon & Strauss, 2016); and (2) superpathway of sialic

acids biosynthesis, which generates sialic acids that play multifarious roles in immunity

including acting as host receptors and pathogen decoys for viruses and bacteria (Varki &

Gagneux, 2012) and are especially critical for preventing neural tissue damage (Liao, Klaus,

& Neumann, 2020). Despite this higher abundance of microbial functions that could enable

invasion-front toads to expand their range, there were some indications that these toads may

also face some health challenges. Invasion-front toad microbiota have significantly higher

abundance of urate biosynthesis function (urate biosynthesis/inosine 5'-phosphate

degradation, the only significantly different one out of the top 30 abundant functions), which

plays a significant, but modest, role in determining variation in serum urate levels (Sinnott-

Armstrong, Naqvi, Rivas, & Pritchard, 2020). Excessively high levels of urate can result in

the formation of needle-like crystals of urate in the joints (gout), which may be related to the

finding that invasion-front cane toads have a higher incidence of severe spinal arthritis

thought to be caused by high levels of movement in combination with morphological

differences found in that population (Brown et al., 2007).

Associations between gut microbiota, environmental isothermaliy and host lungworm

Higher environmental isothermality increased the prevalence of cane toad lungworms, and

both factors could contribute to changes in the cane toads' colon microbiota. It is interesting

that even though environmental isothermality and lungworm infection intensity are not

significantly different between range-core and invasion-front toads, environmental

isothermality and occurrence of lungworms are significantly associated with the microbial

taxonomic community and predicted microbial function. Environmental temperature can alter

microbial taxonomic community and predicted microbial function, resulting in increased

phenotypic plasticity and persistence in harsh conditions (Fontaine & Kohl, 2020b). For

example, gut microbiota of invasion bullfrog tadpoles responds more rapidly to

environmental temperature than noninvasion green frogs (Fontaine & Kohl, 2020b).

Environmental temperature alters the digestive performance and gut microbiota of red-

backed salamanders (*Plethodon cinereus*: Fontaine, Novarro, & Kohl, 2018) and in a study

of western clawed frogs (*Xenopus tropicalis*) the authors observed a higher prevalence of

Verrucomicrobia in the gut of frogs inhabiting a warmer environment, and an opposite

pattern in Proteobacteria (J. Li et al., 2019). The effects of temperature on Verrucomicrobia,

Proteobacteria, and Akkermansia observed in that study was consistent with previous studies

in mammals (J. Li et al., 2019). Pathogens and parasites impact the composition of the host

microbiota and have also been shown to modify host behavior in a manner that improves

their probability of transmission and survival (Gegear, Otterstatter, & Thomson, 2006;

House, Vyas, & Sapolsky, 2011; Poulin, 2010). Lungworms can affect cane toad locomotor

performance and reduced host endurance, presumably because of the reduced oxygen supply

from infected lungs (Pizzatto & Shine, 2012). Lungworms are also known to alter a cane

toad's thermal preference and manipulate the timing and location of defecation, thereby

enhancing lungworm egg production and larvae survival (Finnerty, Shine, & Brown, 2018).

Lungworms are reported to lag behind their host on the invasion front by 2-3 years (Ben L

Phillips et al., 2010) and to affect righting behavior (prolongs righting time; Finnerty, 2017).

In the current study, although we collected invasion-front toads in areas where toads had been

present for less than one year, we found no difference in lungworm presence or intensity

between the invasion-front and range-core toads, nor did we find differences in righting times

between these populations. In combination, these data suggests a need for further studies to

determine how isothermality and occurrence of lungworms may affect cane toad gut

microbiota variation, and whether this influences toad behavior.

Associations between gut microbiota and host righting reflex behavior

Interestingly, behaviors including righting effort likelihood and righting time were not

associated with the gut microbial taxonomy composition but were associated with predicted

microbial functions. It is possible that multiple identified taxa share the same microbial

function or that one taxa contributes to multiple microbial functions, which might obscure the

relationship between behavior and taxonomic composition. Nevertheless, these relationships

we found between gut microbial function and righting behaviors may be related to toads'

health and/or rearing conditions. A dampened stress response (lower level of corticosterone)

in invasion-front toads exposed to stressors (Brown, Kelehear, Shilton, Phillips, & Shine,

2015) could be a result of higher abundance of microbial functions that are beneficial to

general host health and immunity, especially the superpathway of sialic acids biosynthesis,

which generates sialic acids that are especially critical for preventing neural tissue damage

(Liao et al., 2020). Further, invasion-front toads respond less to stressful stimuli and to be

more reluctant to flee than those from range-core toads in simulated predation trials,

exhibiting "bold" anti-predator responses when exposed to novel predators (Hudson, et al.,

2017). Studies showed that dampened stress responses are related to higher propensity for

exploratory behavior (Golla et al., 2020), which has been linked to dispersal ability (Gruber

et al., 2017). Rearing conditions also have been shown to affect righting behavior: juvenile

toads whose parents were collected on the invasion-front that were raised in high exercise

regimes in a common-garden setting righted themselves more quickly than those whose

parents were collected from the range core, whereas the opposite trend was seen in controls

(Stuart et al., 2019). It is also possible that the lower likelihood of righting effort we found in

invasion-front toads could be caused by the higher incidence of severe spinal arthritis (Brown

et al., 2007) described above, which could be related to the higher abundance of microbial

functions (urate biosynthesis function) that contribute to the formation of needle-like crystals

of urate in the joints (Sinnott-Armstrong et al., 2020). Although manipulative studies are

needed to clarify causal relationships between stress responses, proactive behaviors, and gut

microbial functions, our results indicate that behavior and toad colon microbial functions are

related, suggesting that gut microbiome should be considered as a potentially important

driver of invasion.

Conclusion

In this study, we observed significant differences in the microbial community and predicted

microbial functions between range-core and invasion-front Australian cane toad gut

microbiota. Both environmental factors (isothermality and mean temperature of driest

quarter) and host factors (occurrence of lungworms) were significantly associated with these

differences. Also, we found that behavioral traits were not associated with the gut microbiota

taxonomy, but were associated with their predicted microbial functions. Owing to the

predictive nature of PICRUSt and its use here with a non-model host, metagenomic or

transcriptomic sequencing will be an important future research direction for verifying our

results related to gut microbial function (Bletz et al., 2016). In model organisms, the gut

microbiome has been shown to impact host metabolic activity, brain function and the

pathogenesis of disease. Our research demonstrates that the cane toad gut microbiota are

linked to behaviors that may be important to invasion, providing a new perspective of species

invasions. Manipulative experiments (e.g. fecal transplants) coupled with longer term

behavioral measurements would be useful to clarify relationships between microbiota and

host behaviors related to invasion ability in this species.

Data Accessibility Statement
All the data and supporting information will be made available online.
1. Supplementary Figures S7.
2. Supplementary Tables S11.

Author Contributions

1. Designed research: JZ, TMN, CRL, SJZ, LAR
2. Performed Research: JZ, CRL, GWF, KS, LAR
3. Analyzed data: JZ, TMN, CRL, SJZ, KS, LAR
4. Wrote the paper: JZ, TMN, CRL, SJZ, GWF, KS, LAR

Acknowledgements

This project was supported by the Holsworth Wildlife Research Endowment (Ecological
Society of Australia) and Adelaide Graduate Research Scholarship (University of Adelaide)
to JZ, and a Scientia Fellowship to LAR (UNSW). Ms. Alice Russo, and Ms. Rita
Kurpiewski contributed to field sample collections. We would like to thank our colleagues
from University of Adelaide, University of New South Wales, Macquarie University,
University of Kentucky and others who provided insight and expertise that assisted this
research.

Bibliography

Baker, M. R. (1979). The free living and parasitic development of *Rhabdias spp*. (Nematoda:
        Rhabdiasidae) in amphibians. *Canadian Journal of Zoology*, 57, 161-178.

Bashiri, G., Antoney, J., Jirgis, E. N. M., Shah, M. V., Ney, B., Copp, J., … Jackson, C. J.
        (2019). A revised biosynthetic pathway for the cofactor F420 in prokaryotes. *Nature
        Communications*, *10*(1), 1558. doi:10.1038/s41467-019-09534-x

Bercik, P., Denou, E., Collins, J., Jackson, W., Lu, J., Jury, J., … Collins, S. M. (2011). The
        intestinal microbiota affect central levels of brain-derived neurotropic factor and
        behavior in mice. *Gastroenterology*, *141*(2), 599–609, 609.e1.
        doi:10.1053/j.gastro.2011.04.052

Bletz, M. C., Goedbloed, D. J., Sanchez, E., Reinhardt, T., Tebbe, C. C., Bhuju, S., …
        Steinfartz, S. (2016). Amphibian gut microbiota shifts differentially in community
        structure but converges on habitat-specific predicted functions. *Nature
        Communications*, *7*, 13699. doi:10.1038/ncomms13699

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C., Al-Ghalith, G. A., …
        Caporaso, J. G. (2018). QIIME 2: Reproducible, interactive, scalable, and extensible
        microbiome data science. doi:10.7287/peerj.preprints.27295v1

Borrelli, L., Aceto, S., Agnisola, C., De Paolo, S., Dipineto, L., Stilling, R. M., … Fioretti, A.
        (2016). Probiotic modulation of the microbiota-gut-brain axis and behaviour in
        zebrafish. *Scientific Reports*, *6*, 30046. doi:10.1038/srep30046

Brown, G. P., Kelehear, C., Pizzatto, L., & Shine, R. (2016). The impact of lungworm
        parasites on rates of dispersal of their anuran host, the invasive cane toad. *Biological
        Invasions*, *18*(1), 103–114. doi:10.1007/s10530-015-0993-1

Brown, G. P., Kelehear, C., Shilton, C. M., Phillips, B. L., & Shine, R. (2015). Stress and
        immunity at the invasion front: a comparison across cane toad (*Rhinella marina*)
        populations. *Biological Journal of the Linnean Society*, *116*(4), 748–760.
        doi:10.1111/bij.12623

Brown, G. P., Phillips, B. L., & Shine, R. (2014). The straight and narrow path: the evolution

of straight-line dispersal at a cane toad invasion front. *Proceedings of The Royal*
Society B, *281*(1795). doi:10.1098/rspb.2014.1385

Brown, G. P., Shilton, C., Phillips, B. L., & Shine, R. (2007). Invasion, stress, and spinal
arthritis in cane toads. *Proceedings of the National Academy of Sciences of the United*
States of America, *104*(45), 17698–17700. doi:10.1073/pnas.0705057104

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S.
P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data.
Nature Methods, *13*(7), 581–583. doi:10.1038/nmeth.3869

Carmody, R. N., Gerber, G. K., Luevano, J. M., Gatti, D. M., Somes, L., Svenson, K. L., &
Turnbaugh, P. J. (2015). Diet dominates host genotype in shaping the murine gut
microbiota. *Cell Host & Microbe*, *17*(1), 72–84. doi:10.1016/j.chom.2014.11.010

Chai, L., Dong, Z., Chen, A., & Wang, H. (2018). Changes in intestinal microbiota of Bufo
gargarizans and its association with body weight during metamorphosis. *Archives of*
Microbiology, *200*(7), 1087–1099. doi:10.1007/s00203-018-1523-1

Chang, C.-W., Huang, B.-H., Lin, S.-M., Huang, C.-L., & Liao, P.-C. (2016). Changes of diet
and dominant intestinal microbes in farmland frogs. *BMC Microbiology*, *16*, 33.
doi:10.1186/s12866-016-0660-4

Child, T., Phillips, B. L., & Shine, R. (2008). Abiotic and biotic influences on the dispersal
behavior of metamorph cane toads (Bufo marinus) in tropical Australia. *Journal of*
Experimental Zoology. Part A, Ecological Genetics and Physiology, *309*(4), 215–224.
doi:10.1002/jez.450

Choi, J. J., Eum, S. Y., Rampersaud, E., Daunert, S., Abreu, M. T., & Toborek, M. (2013).
Exercise attenuates PCB-induced changes in the mouse gut microbiome.
Environmental Health Perspectives, *121*(6), 725–730. doi:10.1289/ehp.1306534

Colombo, B. M., Scalvenzi, T., Benlamara, S., & Pollet, N. (2015). Microbiota and mucosal
immunity in amphibians. *Frontiers in Immunology*, *6*, 111.
doi:10.3389/fimmu.2015.00111

Dao, M. C., Everard, A., Aron-Wisnewsky, J., Sokolovska, N., Prifti, E., Verger, E. O., …
Clément, K. (2016). Akkermansia muciniphila and improved metabolic health during
a dietary intervention in obesity: relationship with gut microbiome richness and
ecology. *Gut*, *65*(3), 426–436. doi:10.1136/gutjnl-2014-308778

Davidson, G. L., Cooke, A. C., Johnson, C. N., & Quinn, J. L. (2018). The gut microbiome as
a driver of individual variation in cognition and functional behaviour. *Philosophical*
Transactions of the Royal Society of London. Series B, Biological Sciences,
373(1756). doi:10.1098/rstb.2017.0286

Davidson, G. L., Raulo, A., & Knowles, S. C. L. (2020). Identifying Microbiome-Mediated
Behaviour in Wild Vertebrates. *Trends in Ecology & Evolution*.
doi:10.1016/j.tree.2020.06.014

Derrien, M., Collado, M. C., Ben-Amor, K., Salminen, S., & de Vos, W. M. (2008). The
Mucin degrader Akkermansia muciniphila is an abundant resident of the human
intestinal tract. *Applied and Environmental Microbiology*, *74*(5), 1646–1648.
doi:10.1128/AEM.01226-07

DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., …
Andersen, G. L. (2006). Greengenes, a chimera-checked 16S rRNA gene database
and workbench compatible with ARB. *Applied and Environmental Microbiology*,
72(7), 5069–5072. doi:10.1128/AEM.03006-05

Dinan, T. G., Stilling, R. M., Stanton, C., & Cryan, J. F. (2015). Collective unconscious: how
gut microbes shape human behavior. *Journal of Psychiatric Research*, *63*, 1–9.
doi:10.1016/j.jpsychires.2015.02.021

Douglas, G. M., Maffei, V. J., Zaneveld, J., Yurgel, S. N., Brown, J. R., Taylor, C. M., …

787   Langille, M. G. I. (2019). PICRUSt2: An improved and extensible approach for
788   metagenome inference. *BioRxiv*. doi:10.1101/672295
789   Eisenhofer, R., Minich, J. J., Marotz, C., Cooper, A., Knight, R., & Weyrich, L. S. (2019).
790   Contamination in low microbial biomass microbiome studies: issues and
791   recommendations. *Trends in Microbiology*, *27*(2), 105–117.
792   doi:10.1016/j.tim.2018.11.003
793   Fetissov, S. O. (2017). Role of the gut microbiota in host appetite control: bacterial growth to
794   animal feeding behaviour. *Nature Reviews. Endocrinology*, *13*(1), 11–25.
795   doi:10.1038/nrendo.2016.150
796   Fick, S. E. and Hijmans, R. J. (2017) WorldClim 2: new 1km spatial resolution climate
797   surfaces for global land areas. *International Journal of Climatology, 37* (12): 4302-
798   4315. doi.org/10.1002/joc.5086
799   Finnerty, P. B. (2017). *The physiological and ecological impacts of lungworm infection on*
800   *invasive cane toads* (Undergraduate thesis). University of Sydney.
801   Finnerty, P. B., Shine, R., & Brown, G. P. (2018). Survival of the feces: Does a nematode
802   lungworm adaptively manipulate the behavior of its cane toad host? *Ecology and*
803   *Evolution*, *8*(9), 4606–4618. doi:10.1002/ece3.3870
804   Fontaine, S. S., & Kohl, K. D. (2020a). Gut microbiota of invasive bullfrog tadpoles responds
805   more rapidly to temperature than a noninvasive congener. *Molecular Ecology*, *29*(13),
806   2449–2462. doi:10.1111/mec.15487
807   Fontaine, S. S., & Kohl, K. D. (2020b). Gut microbiota of invasive bullfrog tadpoles
808   responds more rapidly to temperature than a noninvasive congener. *Molecular*
809   *Ecology*, *29*(13), 2449–2462. doi:10.1111/mec.15487
810   Fontaine, S. S., Novarro, A. J., & Kohl, K. D. (2018). Environmental temperature alters the
811   digestive performance and gut microbiota of a terrestrial amphibian. *The Journal of*
812   *Experimental Biology*, *221*(Pt 20). doi:10.1242/jeb.187559
813   Gamage, H. K. A. H., Tetu, S. G., Chong, R. W. W., Ashton, J., Packer, N. H., & Paulsen, I.
814   T. (2017). Cereal products derived from wheat, sorghum, rice and oats alter the infant
815   gut microbiota in vitro. *Scientific Reports*, *7*(1), 14312. doi:10.1038/s41598-017-
816   14707-z
817   Garavaglia, M., Rossi, E., & Landini, P. (2012). The pyrimidine nucleotide biosynthetic
818   pathway modulates production of biofilm determinants in Escherichia coli. *Plos One*,
819   *7*(2), e31252. doi:10.1371/journal.pone.0031252
820   Gegear, R. J., Otterstatter, M. C., & Thomson, J. D. (2006). Bumble-bee foragers infected by
821   a gut parasite have an impaired ability to utilize floral information. *Proceedings of the*
822   *Royal Society B*, *273*(1590), 1073–1078. doi:10.1098/rspb.2005.3423
823   Gelbard-Sagiv, H., Magidov, E., Sharon, H., Hendler, T., & Nir, Y. (2018). Noradrenaline
824   modulates visual perception and late visually evoked activity. *Current Biology*,
825   *28*(14), 2239–2249.e6. doi:10.1016/j.cub.2018.05.051
826   Golla, A., Østby, H., & Kermen, F. (2020). Chronic unpredictable stress induces anxiety-like
827   behaviors in young zebrafish. *Scientific Reports*, *10*(1), 10339. doi:10.1038/s41598-
828   020-67182-4
829   González-Bernal, E., Brown, G. P., & Shine, R. (2014). Invasive cane toads: social
830   facilitation depends upon an individual's personality. *Plos One*, *9*(7), e102880.
831   doi:10.1371/journal.pone.0102880
832   Gruber, J., Brown, G., Whiting, M. J., & Shine, R. (2017). Geographic divergence in
833   dispersal-related behaviour in cane toads from range-front versus range-core
834   populations in Australia. *Behavioral Ecology and Sociobiology, 71* (38).
835   doi.org/10.1007/s00265-017-2266-8
Hauffe, H. C., & Barelli, C. (2019). Conserve the germs: the gut microbiota and adaptive

potential. *Conservation Genetics (Print)*, *20*(1), 19–27. doi:10.1007/s10592-019-
01150-y
Herbison, R. E. H. (2017). Lessons in Mind Control: Trends in Research on the Molecular
Mechanisms behind Parasite-Host Behavioral Manipulation. *Frontiers in Ecology and*
Evolution, *5*. doi:10.3389/fevo.2017.00102
Herlemann, D. P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J., & Andersson, A. F.
(2011). Transitions in bacterial communities along the 2000 km salinity gradient of
the Baltic Sea. *The ISME Journal*, *5*(10), 1571–1579. doi:10.1038/ismej.2011.41
Herve, M. (2020). Testing and plotting procedures for biostatistics. *Package*
"RVAideMemoire.". https://CRAN.R-project.org/package=RVAideMemoire
Hochberg, Y. (1988). A sharper Bonferroni procedure for multiple tests of significance.
Biometrika, *75*(4), 800–802. doi:10.1093/biomet/75.4.800
Hopkins, W. A. (2007). Amphibians as models for studying environmental change. *ILAR*
Journal / National Research Council, Institute of Laboratory Animal Resources,
48(3), 270–277. doi:10.1093/ilar.48.3.270
Houlden, A., Goldrick, M., Brough, D., Vizi, E. S., Lénárt, N., Martinecz, B., … Denes, A.
(2016). Brain injury induces specific changes in the caecal microbiota of mice via
altered autonomic activity and mucoprotein production. *Brain, Behavior, and*
Immunity, *57*, 10–20. doi:10.1016/j.bbi.2016.04.003
House, P. K., Vyas, A., & Sapolsky, R. (2011). Predator cat odors activate sexual arousal
pathways in brains of Toxoplasma gondii infected rats. *Plos One*, *6*(8), e23277.
doi:10.1371/journal.pone.0023277
Huang, B.-H., Chang, C.-W., Huang, C.-W., Gao, J., & Liao, P.-C. (2017). Composition and
functional specialists of the gut microbiota of frogs reflect habitat differences and
agricultural activity. *Frontiers in Microbiology*, *8*, 2670.
doi:10.3389/fmicb.2017.02670
Hudson, C. M., Brown, G. P., & Shine, R. (2017). Evolutionary shifts in anti-predator
responses of invasive cane toads (Rhinella marina). *Behavioral Ecology and*
Sociobiology, *71*(9), 134. doi:10.1007/s00265-017-2367-4
Hudson, C. M., Vidal-García, M., Murray, T. G., & Shine, R. (2020). The accelerating
anuran: evolution of locomotor performance in cane toads (Rhinella marina,
Bufonidae) at an invasion front. *Proceedings of the Royal Society B*, *287*(1938),
20201964. doi:10.1098/rspb.2020.1964
Hugenholtz, F., Davids, M., Schwarz, J., Müller, M., Tomé, D., Schaap, P., … Kleerebezem,
M. (2018). Metatranscriptome analysis of the microbial fermentation of dietary milk
proteins in the murine gut. *Plos One*, *13*(4), e0194066.
doi:10.1371/journal.pone.0194066
Jiménez, R. R., & Sommer, S. (2017). The amphibian microbiome: natural range of variation,
pathogenic dysbiosis, and role in conservation. *Biodiversity and Conservation*, *26*(4),
763–786. doi:10.1007/s10531-016-1272-x
Jones, J. C., Fruciano, C., Marchant, J., Hildebrand, F., Forslund, S., Bork, P., … Hughes, W.
O. H. (2018). The gut microbiome is associated with behavioural task in honey bees.
Insectes Sociaux, *65*(3), 419–429. doi:10.1007/s00040-018-0624-9
Kang, D.-W., Park, J. G., Ilhan, Z. E., Wallstrom, G., Labaer, J., Adams, J. B., & Krajmalnik-
Brown, R. (2013). Reduced incidence of Prevotella and other fermenters in intestinal
microflora of autistic children. *Plos One*, *8*(7), e68322.
doi:10.1371/journal.pone.0068322
Kelehear, C., & Jones, H. I. (2010). Nematode larvae (order Spirurida) in gastric tissues of
Australian anurans: a comparison between the introduced cane toad and sympatric
native frogs. *Journal of Wildlife Diseases*, *46*(4), 1126–1140. doi:10.7589/0090-3558-

46.4.1126
Klein, S. L. (2003). Parasite manipulation of the proximate mechanisms that mediate social
behavior in vertebrates. *Physiology & Behavior*, *79*(3), 441–449. doi:10.1016/s0031-
9384(03)00163-x
Knutie, S. A., Wilkinson, C. L., Kohl, K. D., & Rohr, J. R. (2017). Early-life disruption of
amphibian microbiota decreases later-life resistance to parasites. *Nature*
Communications, *8*(1), 86. doi:10.1038/s41467-017-00119-0
Kohl, K. D., Amaya, J., Passement, C. A., Dearing, M. D., & McCue, M. D. (2014). Unique
and shared responses of the gut microbiota to prolonged fasting: a comparative study
across five classes of vertebrate hosts. *FEMS Microbiology Ecology*, *90*(3), 883–894.
doi:10.1111/1574-6941.12442
Kohl, K. D., Cary, T. L., Karasov, W. H., & Dearing, M. D. (2013). Restructuring of the
amphibian gut microbiota through metamorphosis. *Environmental Microbiology*
Reports, *5*(6), 899–903. doi:10.1111/1758-2229.12092
Kozik, A. J., Nakatsu, C. H., Chun, H., & Jones-Hall, Y. L. (2017). Age, sex, and TNF
associated differences in the gut microbiota of mice and their impact on acute TNBS
colitis. *Experimental and Molecular Pathology*, *103*(3), 311–319.
doi:10.1016/j.yexmp.2017.11.014
Kraimi, N., Calandreau, L., Zemb, O., Germain, K., Dupont, C., Velge, P., … Leterrier, C.
(2019). Effects of gut microbiota transfer on emotional reactivity in Japanese quails
(Coturnix japonica). *The Journal of Experimental Biology*, *222*(Pt 10).
doi:10.1242/jeb.202879
Li, J., Rui, J., Li, Y., Tang, N., Zhan, S., Jiang, J., & Li, X. (2019). Ambient temperature
alters body size and gut microbiota of Xenopus tropicalis. *Science China. Life*
Sciences. doi:10.1007/s11427-019-9540-y
Li, Z. S., Furness, J. B., Young, H. M., & Campbell, G. (1992). Nitric Oxide Synthase
Immunoactivity and NADPH Diaphorase Enzyme Activity in Neurons of the
Gastrointestinal Tract of the Toad, Bufo marinus. *Archives of Histology and Cytology*.
doi: 10.1679/aohc.55.333
Liao, H., Klaus, C., & Neumann, H. (2020). Control of innate immunity by sialic acids in the
nervous tissue. *International Journal of Molecular Sciences*, *21*(15).
doi:10.3390/ijms21155494
Llewelyn, J., Phillips, B. L., Alford, R. A., Schwarzkopf, L., & Shine, R. (2010). Locomotor
performance in an invasive species: cane toads from the invasion front have greater
endurance, but not speed, compared to conspecifics from a long-colonised area.
Oecologia, *162*(2), 343–348. doi:10.1007/s00442-009-1471-1
Louis, P., & Flint, H. J. (2017). Formation of propionate and butyrate by the human colonic
microbiota. *Environmental Microbiology*, *19*(1), 29–41. doi:10.1111/1462-
2920.13589
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
doi:10.1186/s13059-014-0550-8
Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K., & Knight, R. (2012).
Diversity, stability and resilience of the human gut microbiota. *Nature*, *489*(7415),
220–230. doi:10.1038/nature11550
Marques, T. M., Wall, R., Ross, R. P., Fitzgerald, G. F., Ryan, C. A., & Stanton, C. (2010).
Programming infant gut microbiota: influence of dietary and environmental factors.
Current Opinion in Biotechnology, *21*(2), 149–156. doi:10.1016/j.copbio.2010.03.020
McMurdie, P. J., & Holmes, S. (2013). phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. *Plos One*, *8*(4), e61217.

937   doi:10.1371/journal.pone.0061217

938 Mudd, A. T., Berding, K., Wang, M., Donovan, S. M., & Dilger, R. N. (2017). Serum cortisol

939   mediates the relationship between fecal Ruminococcus and brain N-acetylaspartate in

940   the young pig. *Gut Microbes*, *8*(6), 589–600. doi:10.1080/19490976.2017.1353849

941 Nath, R., & Pavur, R. (1985). A new statistic in the one-way multivariate analysis of

942   variance. *Computational Statistics & Data Analysis*, *2*(4), 297–315.

943   doi:10.1016/0167-9473(85)90003-9

944 Ning, T., Gong, X., Xie, L., & Ma, B. (2017). Gut Microbiota Analysis in Rats with

945   Methamphetamine-Induced Conditioned Place Preference. *Frontiers in Microbiology*,

946   *8*, 1620. doi:10.3389/fmicb.2017.01620

947 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Wagner,

948   H. (2019). vegan: Community Ecology Package (R package version 2.5-6). Computer

949   software, vegan: Community Ecology Package. https://CRAN.R-

950   project.org/package=vegan

951 Org, E., Mehrabian, M., Parks, B. W., Shipkova, P., Liu, X., Drake, T. A., & Lusis, A. J.

952   (2016). Sex differences and hormonal effects on gut microbiota composition in mice.

953   *Gut Microbes*, *7*(4), 313–322. doi:10.1080/19490976.2016.1203502

954 Phillips, Ben L, Kelehear, C., Pizzatto, L., Brown, G. P., Barton, D., & Shine, R. (2010).

955   Parasites and pathogens lag behind their host during periods of host range advance.

956   *Ecology*, *91*(3), 872–881. doi:10.1890/09-0530.1

957 Phillips, Benjamin L, Brown, G. P., Webb, J. K., & Shine, R. (2006). Invasion and the

958   evolution of speed in toads. *Nature*, *439*(7078), 803. doi:10.1038/439803a

959 Pielou, E. C. (1966). The measurement of diversity in different types of biological

960   collections. *Journal of Theoretical Biology*, *13*, 131–144. doi:10.1016/0022-

961   5193(66)90013-0

962 Pizzatto, L., & Shine, R. (2012). Lungworm infection modifies cardiac response to exercise

963   in cane toads. *Journal of Zoology*, *287*(2), 150–155. doi:10.1111/j.1469-

964   7998.2012.00898.x

965 Poulin, R. (2010). Parasite manipulation of host behavior: an update and frequently asked

966   questions. *Advances in the Study of Behavior*, *41*, 151-186. doi.org/10.1016/S0065-

967   3454(10)41005-0

968 Qu, Y.-F., Wu, Y.-Q., Zhao, Y.-T., Lin, L.-H., Du, Y., Li, P., … Ji, X. (2020). The invasive

969   red-eared slider turtle is more successful than the native Chinese three-keeled pond

970   turtle: evidence from the gut microbiota. *PeerJ*, *8*, e10271. doi:10.7717/peerj.10271

971 R Core Team. (2020). R: A language and environment for statistical computing (R version

972   4.0.2). Computer software, R Foundation for Statistical Computing, Vienna, Austria.

973   https://www.R-project.org/

974 Sampson, T. R., & Mazmanian, S. K. (2015). Control of brain development, function, and

975   behavior by the microbiome. *Cell Host & Microbe*, *17*(5), 565–576.

976   doi:10.1016/j.chom.2015.04.011

977 Savignac, H. M., Kiely, B., Dinan, T. G., & Cryan, J. F. (2014). Bifidobacteria exert strain-

978   specific effects on stress-related behavior and physiology in BALB/c mice.

979   *Neurogastroenterology and Motility*, *26*(11), 1615–1627. doi:10.1111/nmo.12427

980 Schretter, C. E., Vielmetter, J., Bartos, I., Marka, Z., Marka, S., Argade, S., & Mazmanian, S.

981   K. (2018). A gut microbial factor modulates locomotor behaviour in Drosophila.

982   *Nature*, *563*(7731), 402–406. doi:10.1038/s41586-018-0634-9

983 Scott, K. P., Gratz, S. W., Sheridan, P. O., Flint, H. J., & Duncan, S. H. (2013). The influence

984   of diet on the gut microbiota. *Pharmacological Research*, *69*(1), 52–60.

985   doi:10.1016/j.phrs.2012.10.020

Selechnik, D., Richardson, M. F., Shine, R., DeVore, J. L., Ducatez, S., & Rollins, L. A.

(2019). Increased adaptive variation despite reduced overall genetic diversity in a

rapidly adapting invader. *Frontiers in Genetics*, *10*, 1221.

doi:10.3389/fgene.2019.01221

Shannon, C. E. (1948). A mathematical theory of communication. *Bell System Technical

Journal*, *27*(3), 379–423. doi:10.1002/j.1538-7305.1948.tb01338.x

Sharon, G., Segal, D., Ringo, J. M., Hefetz, A., Zilber-Rosenberg, I., & Rosenberg, E. (2010).

Commensal bacteria play a role in mating preference of Drosophila melanogaster.

Proceedings of the National Academy of Sciences of the United States of America,

107(46), 20051–20056. doi:10.1073/pnas.1009906107

Shine, R. (2010). The ecological impact of invasive cane toads (Bufo marinus) in Australia.

The Quarterly Review of Biology, *85*(3), 253–291. doi:10.1086/655116

Sibon, O. C. M., & Strauss, E. (2016). Coenzyme A: to make it or uptake it? *Nature Reviews.

Molecular Cell Biology*, *17*(10), 605–606. doi:10.1038/nrm.2016.110

Sinnott-Armstrong, N., Naqvi, S., Rivas, M. A., & Pritchard, J. K. (2020). GWAS of three

molecular traits highlights core genes and pathways alongside a highly polygenic

background. *BioRxiv*. doi:10.1101/2020.04.20.051631

Stensvold, C. R., & van der Giezen, M. (2018). Associations between Gut Microbiota and

Common Luminal Intestinal Parasites. *Trends in Parasitology*, *34*(5), 369–377.

doi:10.1016/j.pt.2018.02.004

Stuart, K. C., Shine, R., & Brown, G. P. (2019). Proximate mechanisms underlying the rapid

modification of phenotypic traits in cane toads (*Rhinella marina*) across their

invasive range within Australia. *Biological Journal of the Linnean Society*, *126*(1),

68–79. doi:10.1093/biolinnean/bly150

Szychlinska, M. A., Di Rosa, M., Castorina, A., Mobasheri, A., & Musumeci, G. (2019). A

correlation between intestinal microbiota dysbiosis and osteoarthritis. *Heliyon*, *5*(1),

e01134. doi:10.1016/j.heliyon.2019.e01134

Teseo, S., van Zweden, J. S., Pontieri, L., Kooij, P. W., Sørensen, S. J., Wenseleers, T., …

Sapountzis, P. (2019). The scent of symbiosis: gut bacteria may affect social

interactions in leaf-cutting ants. *Animal Behaviour*, *150*, 239–254.

doi:10.1016/j.anbehav.2018.12.017

Thomas, R. H., Meeking, M. M., Mepham, J. R., Tichenoff, L., Possmayer, F., Liu, S., &

MacFabe, D. F. (2012). The enteric bacterial metabolite propionic acid alters brain

and plasma phospholipid molecular species: further development of a rodent model of

autism spectrum disorders. *Journal of Neuroinflammation*, *9*, 153. doi:10.1186/1742-

2094-9-153

Valverde, A., Makhalanyane, T. P., & Cowan, D. A. (2014). Contrasting assembly processes

in a bacterial metacommunity along a desiccation gradient. *Frontiers in

Microbiology*, *5*, 668. doi:10.3389/fmicb.2014.00668

Varki, A., & Gagneux, P. (2012). Multifarious roles of sialic acids in immunity. *Annals of the

New York Academy of Sciences*, *1253*, 16–36. doi:10.1111/j.1749-6632.2012.06517.x

Venables, W. N., & Ripley, B. D. (2002). Modern Applied Statistics with S. Springer-Verlag.

New York. http://www.stats.ox.ac.uk/pub/MASS4/

Vogel, H., Shukla, S. P., Engl, T., Weiss, B., Fischer, R., Steiger, S., … Vilcinskas, A.

(2017). The digestive and defensive basis of carcass utilization by the burying beetle

and its microbiota. *Nature Communications*, *8*, 15186. doi:10.1038/ncomms15186

Wang, J., Thingholm, L. B., Skiecevičienė, J., Rausch, P., Kummen, M., Hov, J. R., …

Franke, A. (2016). Genome-wide association analysis identifies variation in vitamin D

receptor and other host factors influencing the gut microbiota. *Nature Genetics*,

48(11), 1396–1406. doi:10.1038/ng.3695

Weitzman, C. L., Kaestli, M., Gibb, K., Brown, G. P., Shine, R., & Christian, K. (2019).

Disease exposure and antifungal bacteria on skin of invasive cane toads, australia.
Emerging Infectious Diseases, *25*(9), 1770–1771. doi:10.3201/eid2509.190386
West, T. P. (1996). Degradation of pyrimidine ribonucleosides by Pseudomonas aeruginosa.
Antonie Van Leeuwenhoek, *69*(4), 331–335. doi:10.1007/bf00399622
Wiebler, J. M., Kohl, K. D., Lee, R. E., & Costanzo, J. P. (2018). Urea hydrolysis by gut
bacteria in a hibernating frog: evidence for urea-nitrogen recycling in Amphibia.
Proceedings of the Royal Society B, *285*(1878). doi:10.1098/rspb.2018.0241
Wong, A. C.-N., Wang, Q.-P., Morimoto, J., Senior, A. M., Lihoreau, M., Neely, G. G., …
Ponton, F. (2017). Gut Microbiota Modifies Olfactory-Guided Microbial Preferences
and Foraging Decisions in Drosophila. *Current Biology*, *27*(15), 2397–2404.e4.
doi:10.1016/j.cub.2017.07.022
Ye, L., Amberg, J., Chapman, D., Gaikowski, M., & Liu, W.-T. (2014). Fish gut microbiota
analysis differentiates physiology and behavior of invasive Asian carp and indigenous
American fish. *The ISME Journal*, *8*(3), 541–551. doi:10.1038/ismej.2013.181
Zhang, X.-Y., Sukhchuluun, G., Bo, T.-B., Chi, Q.-S., Yang, J.-J., Chen, B., … Wang, D.-H.
(2018). Huddling remodels gut microbiota to reduce energy requirements in a small
mammal species during cold exposure. *Microbiome*, *6*(1), 103. doi:10.1186/s40168-
018-0473-9
Zhou, J., Nelson, T. M., Rodriguez Lopez, C., Sarma, R. R., Zhou, S. J., & Rollins, L. A.
(2020). A comparison of nonlethal sampling methods for amphibian gut microbiome
analyses. *Molecular Ecology Resources*. doi:10.1111/1755-0998.13139
Zmora, N., Suez, J., & Elinav, E. (2019). You are what you eat: diet, health and the gut
microbiota. *Nature Reviews. Gastroenterology & Hepatology*, *16*(1), 35–56.
doi:10.1038/s41575-018-0061-2

Tables and Figures

Table 1. Comparison of host characteristics, behavioral traits and environmental

factors between range-core and invasion-front toads.

Variables		wilcox.test	GLM (SUL as covariate)		
			glm.nb	glm	binomial
Host characteristics	Body length (SUL)	_	_	0.01099 *	_
	Lungworms	_	0.8600	_	_
	Occurrences of lungworms	_ _	_	_	0.07783 .
	Gut parasites	_	0.817578	_	_
	Occurrences of gut parasites	_	_	_	0.604488
Behavioural traits	Struggle score	_ _	0.002196 **	_	_
	Struggle likelihood	_	_	_	0.008477 **
	Righting effort	_	0.3203	_	_
	Righting effort likelihood	_	_	_	0.03569 *
	Righting time	_	0.5067	_	_
Environmental factors	Longitude	1.596e-11***	_	_	_
	Latitude	0.458	_	_	_
	Annual Mean Temperature	1.608e-07 ***	_	_	_
	Mean Diurnal Range	0.0001836 ***	_	_	_
	Isothermality	0.4515	_	_	_
	Temperature Seasonality	0.02501 *	_	_	_
	Max Temperature of Warmest Month	1.608e-07 ***	_	_	_
	Min Temperature of Coldest Month	0.458	_	_	_
	Temperature Annual Range	0.0001836 ***	_	_	_
	Mean Temperature of Wettest Quarter	1.608e-07 ***	_	_	_
	Mean Temperature of Driest Quarter	0.02099 *	_	_	_
	Mean Temperature of Warmest Quarter	1.608e-07 ***	_	_	_
	Mean Temperature of Coldest Quarter	0.0001836 ***	_	_	_
	Annual Precipitation	1.608e-07 ***	_	_	_
	Precipitation of Wettest Month	1.608e-07 ***	_	_	_

Precipitation of Driest Month	8.069e-12 ***	_	_	_
Precipitation Seasonality	0.02501 *			
Precipitation of Wettest Quarter	1.608e-07 ***	_	_	_
Precipitation of Driest Quarter	5.222e-06 ***	_	_	_
Precipitation of Warmest Quarter	1.596e-11 ***	_	_	_
Precipitation of Coldest Quarter	5.222e-06 ***	_	_	_

Note

Negative binomial regression (glm.nb) was used for over-dispersed count data, that is when the conditional variance exceeds the conditional mean. Shapiro-wilk test to assess multivariate normality using RVAideMemoire package in R (Herve, 2018). If it is below 0.05, the data significantly deviate from a normal distribution. Wilcox.test used for the data that's not normally distributed and ttest for the data that's normally distributed. Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1. SUL: snout urostyle length (body length).

1104
1105 **Figure 1. Core bacterial community composition of different individual toads.** Relative
1106 abundance plot shows phyla (>2%). WA: invasion-front; QLD: range-core.
1107
1108

1109
1110
1111
1112 **Pairwise comparisons using permutation MANOVAs on a weighted UniFrac distance**
1113 **matrix of Core50 microbiota**

	Kununurra	**Old Theda**	**Mary Pool**	**Rossville**	**Croydon**
Old Theda	0.0608	-	-	-	-
Mary Pool	0.0045	0.0584	-	-	-
Rossville	0.0011	0.0340	0.0175	-	-
Croydon	0.0020	0.0608	0.0011	0.0011	-
Lucinda	0.0011	0.1352	0.0011	0.0324	0.0175

1114 P value adjustment method: Hochberg (1988)
1115
1116
1117

1118

1119

1120 **Pairwise comparisons using permutation MANOVAs on a Bray Curtis distance matrix**
1121 **of Core50 microbial functions**

	Kununurra	**Old Theda**	**Mary Pool**	**Rossville**	**Croydon**
Old Theda	0.298	-	-	-	-
Mary Pool	0.298	0.298	-	-	-
Rossville	0.009	0.298	0.298	-	-
Croydon	0.105	0.298	0.298	0.298	-
Lucinda	0.046	0.298	0.298	0.298	0.298

1122 P value adjustment method: Hochberg (1988)

1123

Figure 2. Beta diversity by locations and sites. Principle coordinate analysis plot of
weighted UniFrac distance of microbial community (R^2 = 0.095, F = 6.092, p-value < 0.001)
(A) and Bray Curtis distance of predicted functional groups (R^2 = 0.064, F = 4.110 , p-value=
0.002) (B) from 60 cane toad individuals of the invasion-front (Kununurra, Old Theda, and
Mary Pool) and the range-core (Rossville, Croydon, and Lucinda). Pairwise comparisons
were conducted on both distance matrixes using permutation MANOVAs. P values were
adjusted with Hochberg method.

1131

1132

1133
Figure 3. Heatmap for top 30 functional group abundance.
Heatmap indicates the top 30 functional groups in the intestinal samples from range-core and
invasion-front toads. Abundance indicates the raw count of functional groups inferred from
taxonomic 16S sequences using PICRUSt where light blue is high abundance and dark blue
is lower abundance. Functional pathways that differ significantly between range-core and
invasion-front toads are highlighted in bold.
1140

1141

1142
1143
Figure 4. Significant different bacterial taxa and predicted functions between range-
core (QLD) and invasion-front (WA) toads' colon. Significant differences were identified
between locations via differential abundance testing based on a negative binomial
distribution. The dots represent the average log-2 fold change (x axis) abundance and positive
log_2 fold changes signify increased abundance in range-core, and negative log_2 fold changes
display increased abundance in invasion-front. Bacterial taxa (A) were classified to the
taxonomic level of family (y axis) and coloured by taxonomic level of phylum. Family name
in bracket is proposed taxonomy by Greengenes. Only ASVs that could be matched to a
known bacterial family and with a log2FoldChange value higher than 20 or lower than -20
are presented. Predicted functions (B) with a log2FoldChange value higher than 3 or lower
than -3 are presented.
1155

1156
1157
Figure 5. Main variables that affect predicted function differentiation among
individuals. CAP (*capscale*) plot displays the combination of variables that explained the
greatest variation in the predicted functions through model selection, using 60 cane toad
individuals from the invasion-front (Kununurra, Old Theda, and Mary Pool) and the range-
core (Rossville, Croydon, and Lucinda). The final model explained 17.8% of variation in the
microbial predicted functions, which includes righting effort likelihood (AIC = 53.613),

occurrence of lung worms (AIC = 54.297) and righting time (AIC = 56.912) explained the
greatest variation.
1166
1167
1168
1169