The gut bacteria of an invasive amphibian respond to the dual challenges of range-expansion and parasite attack

Jia Zhou[1,6*,] Tiffanie Maree Nelson[2], Carlos Rodriguez Lopez[3], Shao Jia Zhou[1,4], Georgia

Ward-Fear[5], Katarina C. Stuart[6], Richard Shine[5], Lee Ann Rollins[6]

[1]School of Agriculture, Food and Wine, University of Adelaide, Waite Campus, PMB 1 Glen

Osmond, South Australia, Australia, 5064.

[2]Queensland Facility for Advanced Bioinformatics, School of Medicine, Menzies Health Institute

Queensland, Building G40, Level 9, Gold Coast Campus, Griffith University, Southport, QLD,

Australia, 4215.

[3]Environmental Epigenetics and Genetics Group, Department of Horticulture, College of Agriculture,

Food and Environment, University of Kentucky, Lexington, KY, USA, 40546.

[4]Robinson Research Institute, University of Adelaide, South Australia, Australia, 5005

[5]Department of Biological Sciences, Macquarie University, North Ryde, NSW, Australia, 2109.

[6]Ecology & Evolution Research Centre, School of Biological, Earth and Environmental Sciences,

University of New South Wales, Biological Sciences South (E26) UNSW, Kensington, Australia,

2052

* Correspondence:

Jia Zhou

E-mail: jiazhou0116@gmail.com

Keywords: microbiome, Cane toad, 16S rRNA, host-parasite

Word count: 6334.

Abstract

Gut bacterial communities influence, and are influenced by, the behaviour and ecology of their hosts.

Those interactions have been studied primarily in humans and model organisms, but we need field

research to understand the relationship between an organism's gut bacteria and its ecological

challenges, such as those imposed by rapid range expansion (as in invasive species) and the presence

of host-manipulating parasites. Cane toads (*Rhinella marina*) provide an excellent model system in

this respect, because the species' ongoing colonization of Australia has enforced major changes in

phenotypic traits (including behaviour), and lungworm parasites (*Rhabdias pseudosphaerocephala*)

modify host gut function in ways that enhance the viability of lungworm larvae. We collected female

toads from across the species' invasive range and studied their morphology, behaviour, parasite

infection status and gut bacterial community. Range-core *versus* range-edge toads differed in

morphology, behaviour, gut bacterial composition and predicted gut bacterial function but did not

differ in the occurrence of parasite infection nor in the intensity of infection. Toads infected with

lungworms differed from uninfected conspecifics in gut bacterial composition and diversity. Our

study demonstrates strong associations between gut bacterial community and host ecology and

behaviour.

Introduction

The bacterial community within an organism's intestines can be strongly influenced by host

behaviour and ecology, such as habitat selection and diet [1–5]. But that interaction runs both ways

because gut bacteria can influence behaviour and ecology of the host. For example, transferring gut

contents can modify the recipient host's behaviour (exploratory behaviour, *Mus musculus* [6];

emotional reactivity, *Coturnix japonica* [7]). Similarly, altering gut microbial communities by

administering antibiotics or altering dietary composition triggered aggressive behaviour in leaf-

cutting ants (*Acromyrmex echinatior* [8]). Remarkably, changes in only a single bacterial species

within the gut can affect behaviour of the host (e.g., *Drosophila melanogaster* [9]; *Danio rerio* [10];

M. musculus [11]). Gut microbiota can also affect mating choices [12] and foraging [13–15].

To date, most evidence for effects of intestinal bacteria on host behaviour comes from studies on

humans and "model organisms". To elucidate the functional significance of this phenomenon, we

need to extend such studies to free-ranging animals, incorporating a wider range of taxa [16].

Invasive species offer good models for such research, because novel challenges in the invaded range

create an opportunity to compare closely-related organisms exposed to profoundly different

environments [17,18]. Host-parasite relationships also may be disrupted during biological invasions,

due to processes such as "enemy release" (loss of co-evolved native pathogens from the native range

[19]). Parasites can manipulate host behaviour and physiology in ways that enhance parasite fitness

but reduce host fitness [20]. Thus, if the gut bacterial community provides a mechanism for such

effects, parasite-infected individuals should exhibit different gut bacteria than uninfected

conspecifics.

These ideas suggest two predictions: (i) that the gut bacterial community should differ between

populations of an invasive species (e.g., between the range-core and the invasion-front); and (ii) that

the gut bacterial community should differ between parasitized hosts and non-parasitized conspecifics.

The colonization of Australia by cane toads (*Rhinella marina*) provides a robust opportunity to test

these predictions. Since their release in north-eastern Australia in 1935, toads have dispersed into

areas that are much hotter and more seasonally arid than in the native range or the initial release sites

[21]; and toads have brought with them a native-range nematode lungworm (*Rhabdias*

pseudosphaerocephala) that can have devastating impacts on host viability, and induces behavioural

and physiological changes in the host [22]. Notably, infected toads produce copious watery faeces

[22]; hence, we expect lungworm-infected toads to exhibit different gut bacterial communities than

uninfected individuals.

Methodology

Study species, sample collection and behavioural assays

Cane toads are native to South America, and were introduced into Australia in 1935 as a biocontrol

for pests of sugar cane crops [23]. As the toads spread through tropical Australia, they fatally

poisoned many native predators [23]. Toads from range-core populations (eastern Australia) differ

from invasion-front conspecifics (in north-western Australia) in phenotypic traits that confer

increased dispersal ability, such as endurance [24], limb morphology [25], boldness and exploratory

behaviour [17,26]. Toads from the invasion-front also have lower rates of infection of the co-

introduced parasitic lungworm [27]. Drivers of variation in invasion-related behaviours in this

species include genetics, morphology, habitat, diet, prior experience and parasites [17,28–31].

However, the possible role of gut bacteria as a potential driver of behavioural shifts across the

invasive range has not been studied.

We hand-captured 60 adult females from three sites on the invasion-front and three sites in the range-

core (Table S1). We conducted brief behavioural assays upon collection including: (i) struggle score

(number of kicks after being captured until toad remains still for 5 seconds) and struggle likelihood;

and (ii) righting effort (time to right itself, number of kicks within two minutes after toad is placed on

its dorsal side, and righting effort likelihood [17]. These measures are predictive of traits including

speed and stamina (K. Stuart, pers. comm.), suggesting that these simple assays may reveal a toad's

dispersal potential. We then placed animals into individual, moist, calico bags and weighed,

measured (snout urostyle length; SUL) and euthanised them by injecting tricaine methanesulfonate

(MS222) buffered with bicarbonate of soda.

We dissected the toads and scored the presence of two types of toad parasite: the gut-encysted

physalopterine larvae [32] and adult lungworms [22,33]. Lungworm larvae pass through the gut, but

are less easily detected and reliably counted than are adult lungworms. From each toad, we removed

0.3cm of colon near the cloaca (including gut contents) and preserved it in 95% ethanol (see

Supplementary Material for justification of sampling protocols).

Analyses

We compared host morphology and behaviour between regions (range-core *versus* invasion-front),

and as a function of infection status (lungworm infected *versus* non-infected). Because body length

(SUL) and mass were correlated, we only included SUL as our measure of host morphology in

further analyses. We used a t-test to compare mean SUL between regions and infected/non-infected

toads. For associations between region or infection status with host characteristics or behavioural

traits, we used SUL as a covariate in generalized linear models (GLM). See Supplementary Material

for details of statistical analyses.

Laboratory methods and data pre-processing for characterizing gut bacterial community composition
and predicted functions are described in Supplementary Material. Briefly, we calculated within-
individual (alpha) bacterial diversity and between-site (beta) bacterial diversity. For the latter
variable, we subset our data to include the Core50 gut community (Amplified Sequence Variants
(ASVs) present in a minimum of 50% of toads from each site [2]). We predicted bacterial functions
and generated pathway abundance based on Core50 ASVs. We compared bacterial composition and
predicted function between regions, and between lungworm-infected and non-infected toads. We
identified differences in individual ASVs and predicted bacterial functions between range-core and
invasion-front toads and identified associations between host characteristics (including infection with
parasites) with bacterial communities and predicted bacterial functions. Analyses were conducted in
QIIME2 [34], PICRUST2 [35], and R packages in R version 4.0.2 [36].

Results

Ecological traits

Wild-caught invasion-front toads were larger than range-core toads (Tables S2, S3; mean SUL t =
2.54, df = 53.90, p = 0.014). Neither counts nor presence of parasites (lungworm and gut) differed
significantly across the range (Table S2). Range-core toads were more likely to struggle (Tables S2,
S3; p = 0.008, 95% CI: core [-0.174, 0.005], edge [-0.023, 0.199]) and, in those that did struggle, the
number of struggle movements was higher for range-core toads (Tables S2, S3; p = 0.002, 95% CI:
core [-0.057, 0.026], edge [0.046, 0.2]). Range-core toads also were more likely to attempt to right
themselves (p = 0.036, 95% CI: core [-0.092, 0.04], edge [-0.006, 0.19]), but righting effort and
righting time did not differ significantly between geographic regions (Tables S2, S3).
132
133 Because there were no significant differences in prevalence or intensity of lungworm between the
134 range-core and invasion-front toads (Table S2), we combined samples to analyse correlates of
135 lungworm infection. Infected toads were similar in SUL to non-infected toads (Tables S3, 4; t = 0.86,
136 df = 57.19, p = 0.393), with no significant behavioural differences between the two groups (Table S3,
137 4).
138
Gut bacterial community composition and predicted bacterial function
Alpha diversity did not differ significantly between regions (Supplementary Material), but beta
diversity of bacterial taxonomic communities differed between regions (Figures 1A, S1; R^2 = 0.050,

F = 3.050, p < 0.001) and sampling sites (Table S5; all p-values < 0.001). Among 230 ASVs that

were assigned to family level, the abundance of 124 ASVs differed between the colons of range-core

versus invasion-front toads (Table S6). The number of significantly different ASVs in each phylum

were: Bacteroidetes (60 ASVs), Firmicutes (55 ASVs), Proteobacteria (7 ASVs), Actinobacteria (1

ASVs), Verrucomicrobia (1 ASV) (Table S6, Figure 2A).

Among the identified 474 predicted bacterial functions, we found significant differences between

invasion-front and range-core toads (Figure 1B; R^2 = 0.064, F = 4.110, p-value= 0.002). Pairwise

tests between sampling sites indicated that Kununurra toads had different bacterial functions to

Rossville (p-value= 0.009) and Lucinda toads (p-value = 0.046), but no other sites differed

functionally from each other (Table S7; all other p-values > 0.05). In total, 84 predicted bacterial

functions differed between invasion-front and range-core toads (Table S8, Figure 2B). Range-core

toads had more abundant bacterial function in the superpathway of pyrimidine ribonucleosides

degradation (log2FoldChange = 5.98) and less abundant bacterial function in phosphopantothenate

biosynthesis III (log2FoldChange = -4.98), superpathway of sialic acids and CMP-sialic acids

biosynthesis (log2FoldChange = -4.89) and factor 420 biosynthesis (log2FoldChange = -4.72) than

did invasion-front toads (Table S8, Figure 2B). Among the 30 most abundant functions, range-core

toads had lower bacterial function in urate biosynthesis/inosine 5'-phosphate degradation

(log2FoldChange = -0.10) than did invasion-front toads (Figure 3, Table S8).

Associations between ecological traits and intestinal bacteria

To assess correlates of gut bacterial composition and function, we compared characteristics of

individual hosts to bacterial variation. Only the occurrence of lungworms was significantly

associated with the bacterial composition (R^2 = 0.128, p = 0.02) (Table 1; Figure S2A, B).

In a redundancy analysis combining ecological traits measured here, the model that explained the

most variation in the bacterial community assemblage included only the occurrence of lungworms

(AIC = 178.58). The best model to explain variation in predicted bacterial functions included the

likelihood of righting (AIC = 53.613), the occurrence of lungworms (AIC = 54.297) and righting

time (AIC = 56.912). The combination of these three factors explained 17.8% of total variation in

predicted bacterial functions (Figure 4).

In explicit tests of whether bacterial community assemblages differed in infected *versus* non-infected

toads, we found a significant association with lungworm occurrence (Table 2, p=0.005). Intensity of

lungworm infection was not significantly associated with gut bacterial community but did have a

significant interaction with region in this analysis (Table 2, p=0.04).

Discussion

Bacteria influence animal behaviour in diverse ways [16,37], but the ecological drivers of variation in

gut bacterial composition remain largely unstudied. Our analyses of cane toads from two regions

within their invasive range documents substantial variation in community assemblage and function of

gut bacteria. Importantly, that variation was associated with two traits that we predicted to influence

gut bacterial assemblages: invasion history and parasite infection. Interestingly, toad behaviour

differed across the invasive range, and toad righting behaviour was associated with bacterial function

but not with parasite infection.

Geographic divergence in gut bacteria

First, we consider the differences in gut bacteria between toads from the invasion-front and the

range-core. Although these populations have been separated by less than a century, the toads have

diverged remarkably in morphology, physiology and behaviour and much of that divergence is

heritable [38,39]. Some of those shifts likely reflect evolutionary pressures for increased rates of

dispersal, due to adaptive (natural selection) and non-adaptive (spatial sorting) mechanisms [40,41].

Other geographically variable aspects of toad phenotypes likely are responses to different climatic

conditions in the newly-invaded regions (hot, seasonally arid) compared to the range-core (cooler,

more equable climate) [42]. Similar geographic divergence has been reported for the microbiome on

the toad's skin [43]. Our data illustrate that the invasion of Australia by cane toads has been

accompanied by substantial divergence in gut bacterial communities. Alpha diversity in gut bacteria

was similar in invasion-front and range-core individuals, but there were differences in both the gut

bacteria composition and predicted bacterial function between toad populations across the species'

Australian invasive range. Predicted bacterial functions better explained cane toad righting behaviour

than did gut bacterial community composition. Intriguingly, similarity between gut bacterial

communities between individuals within regions in Australia is related to the similarity of their host's

epigenome, and this relationship is strengthened in populations where genetic diversity is lowest,

such as on the invasion front [44]. Relationships between gut bacterial communities and their hosts

are complex, and that a clear understanding of these relationships requires careful consideration of

numerous environmental, host and gut bacterial factors.

The diversity and composition of bacterial communities differed between range-core and invasion-

front toads, despite an overall similarity in their dominant phyla and alpha diversity. ASVs in the

family Veillonellaceae were higher at the invasion-front (Figure 2A). The abundance of this bacterial

family may influence host metabolic regulation. For example, in Brandt's voles (*Lasiopodomys*

brandtii) exposed to colder temperatures, voles which huddled had more Veillonellaceae and more

short-chain fatty acids (SCFAs) in their intestines than did non-huddling voles [45]. This family

produces SCFAs such as propionic acid [46,47], which can increase locomotor activity [48]. The link

to host metabolic regulation suggests that invasion-front toads might fuel their invasion in this way

[24]. ASVs from another family of SCFA-producing bacteria, Clostridiaceae [49], were also higher

in invasion-front toads than those from the range-core. Furthermore, the family Veillonellaceae may

be associated with host sociality. A reduction of Veillonellaceae has been observed in children with

Autism Spectrum Disorder, often known for desiring social isolation [50]. Higher abundance of

Veillonellaceae in invasion-front toads could foster their "bolder" personality, retaining a higher

propensity for exploration and risk-taking [26,51].

Several other ASVs that differed across the toad's range also may affect behaviour. ASVs from the

family Peptococcaceae, more common in invasion-front toads (Figure 2A), are related to host

neurotransmitter levels (noradrenaline linking visual awareness to external world events [52]). For

example, Peptococcaceae levels in the caecum are positively correlated with noradrenaline levels in

mice [53]. ASVs from family Bacillaceae, lower in invasion-front toads (Figure 2A), might be

related to host anxiety (e.g., abundant in methamphetamine-treated rats [54], and in exercised *versus*

sedentary mice [55]). Abundant Bacillaceae might induce anxiety-like behaviours, thus intensifying

the stress response [54] and decreasing exploratory behaviour in new environments [56]. In

summary, invasion-front toads possessed gut bacterial communities that in other studies have been

associated with SCFAs production and neurotransmitters. That pattern supports the idea that gut

microbes in invasion-front toads may increase locomotor ability, alertness and propensity for

exploration and risk-taking. In comparison, range-core toads possessed bacterial taxa that have been

associated with anxiety, and a decreased propensity to explore.

Geographic variation was less obvious in the predicted bacterial functional groups than in community

composition (Figure 1), consistent with the hypothesis that bacterial function is more conservative

than taxonomic composition (e.g. in fire salamanders [2]). Different gut microbiota can have similar

bacterial functions, increasing resilience and functional stability [2,3,57]. Despite this broad

similarity, bacterial functions differed between range-core and invasion-front toads. These

differences included those involved in functional pathways related to food sources and metabolism.

Invasion-front toads had less bacterial function in the superpathway of pyrimidine ribonucleosides

degradation, which provides a nitrogen source for microbes [58] and plays an important role in

perturbations in the uridine monophosphate (UMP) biosynthetic pathways. This pathway allows the

bacterial cell to sense signals such as starvation, nucleic acid degradation, and availability of

exogenous pyrimidines, and to adapt the production of the extracellular matrix to changing

environmental conditions [59]. This function might help to explain the disappearance of

Verrucomicrobia as a dominant taxon. As for microbe metabolism, invasion-front toads have higher

abundance of bacterial functions in factor 420 biosynthesis, critical to bacterial metabolism and

mediating important redox transformations involved in bacterial persistence, antibiotic biosynthesis,

pro-drug activation, and methanogenesis [60].

We also detected geographic variation in bacterial functional pathways that contribute to host health.

Invasion-front toads exhibited bacterial functions beneficial to host health and immunity: (i)

phosphopantothenate biosynthesis (involved in bacterial production of coenzyme A [61]); and (ii)

superpathway of sialic acids biosynthesis (involved in immunity including acting as host receptors

and pathogen decoys for viruses and bacteria [62] and especially critical for preventing neural tissue

damage [63]). Despite this abundance of health-promoting bacterial functions, these toads may also

face health challenges. Invasion-front toad bacteria had a higher abundance of urate biosynthesis

function (urate biosynthesis/inosine 5'-phosphate degradation, the only significantly different one out

of the top 30 abundant functions), which affects serum urate levels [64]. High levels of urate can

result in the formation of needle-like crystals of urate in the joints (gout), perhaps related to severe

spinal arthritis in invasion-front cane toads [65].

Associations between lungworms and host gut bacteria

Pathogens and parasites impact the composition of the host microbiota and can modify host

behaviour in a manner that improves parasite transmission and survival [66–68]. Lungworms can

affect cane toad locomotor performance and reduce host endurance, curtailing oxygen supply from

infected lungs [69]. Lungworms also can alter a cane toad's thermal preference and manipulate the

timing and location of defecation, thereby enhancing lungworm egg production and larval survival

[22]. Lungworms are reported to lag behind their host on the invasion-front by 2-3 years [27] and to

affect righting behaviour (prolongs righting time [70]). In the current study, although we collected

invasion-front toads in recently invaded areas, we found no difference in lungworm presence or

intensity between the invasion-front *versus* range-core toads, nor did we find behavioural differences

in lungworm-infected *versus* uninfected toads.

Infection by parasitic lungworms was associated with differences in gut bacteria. Here, the direction

of causation is less ambiguous than is the case for geographic variation in the gut bacteria. It seems

unlikely that a toad's bacteria affect its probability of carrying adult lungworms, although bacterial-

driven differences in habitat selection might create such a link. Instead, we suggest that the presence

of lungworms induces a shift in gut bacteria. Consistent with that hypothesis, experimental trials have

shown that lungworms modify gastric function in their hosts, changing the volume and consistency

of faeces produced in ways that enhance survival of larval lungworms [22]. Shifts in the microbiome

inside the gut might be either causes or consequences of that shift in gastric function. Moreover, *C.*

elegans are known to prefer specific bacterial foods [71], suggesting that lungworm larvae may also

feed selectively on bacteria in the gut, generating differences in bacterial communities between

lungworm-infected toads *versus* non-infected conspecifics. Additionally, gut bacteria may affect

lungworms via microbiome-induced shifts in host immunity [72].

Associations between host behaviours and gut bacteria

Interestingly, behaviours including righting effort likelihood and righting time were associated more

closely with predicted gut bacterial functions than with bacterial taxonomic composition. Multiple

identified taxa may share the same bacterial function, or one taxon may contribute to multiple

bacterial functions, obscuring the relationship between host behaviour and bacterial taxonomic

composition. Nonetheless, these relationships we found between gut bacterial function and righting

behaviours may be related to toad health and/or rearing conditions. A dampened stress response

(lower corticosterone levels) in invasion-front toads [73] could result from higher abundance of

bacterial functions beneficial to host health and immunity, especially the superpathway of sialic acids

biosynthesis [63]. Further, invasion-front toads are more reluctant to flee in simulated predation trials

[74]. Dampened stress responses can be related to more exploratory behaviour [56], and to greater

dispersal ability [26]. Rearing conditions also affect righting behaviour [17]. Although manipulative

studies are needed to clarify causal relationships between stress responses, proactive behaviours, and

gut bacterial functions, our results indicate that host behaviour and gut bacterial functions are related,

suggesting that gut bacteria may be an important driver of invasion.

Our study has identified patterns rather than testing alternative hypotheses about underlying causal

processes. To clarify causal mechanisms underlying the geographic divergence in gut bacteria across

the toads' Australian range, future studies could use reciprocal transplantation to examine if (and

how) their gut bacteria respond to novel environmental conditions. Breeding these animals, and

raising their offspring under common-garden conditions, could reveal the degree to which a toad's

gut bacteria is driven by host genetics *versus* their rearing conditions [75,76]. To clarify the

hypothesis that changes in gut bacteria mediate the ability of lungworm parasites to modify host gut

function, we could implant colon contents from infected into uninfected toads. In short, our discovery

of strong associations between gut bacteria and important facets of toad ecology provides the

opportunity to move to hypothesis-testing experimental studies.

Our research illustrates that during invasion, as a species expands across a novel and variable

landscape, a complex relationship between host behaviour, its parasite community, and its

microbiome may unfold. A clearer understanding of these relationships and how they influence the

rate of expansion are key to understanding the role of the holobiont during invasion [77]. Such

studies also will advance our understanding of co-evolution and may facilitate innovative approaches

to invasive species management.

Ethics

Approved by University of Adelaide Animal Ethics Committee (S-2018-056).

Author Contributions

Designed research: JZ, TMN, CRL, SJZ, LAR; performed research: JZ, CRL, GWF, KS, LAR;

analyzed data: JZ, TMN, CRL, SJZ, KS, LAR; drafted manuscript: JZ, RS, TMN, CRL, SJZ, GWF,

KS, LAR.

Funding

This project was supported by the Holsworth Wildlife Research Endowment and University of

Adelaide Graduate Research Scholarship to JZ, and a UNSW Scientia Fellowship to LAR. CMRL is

partially supported by the National Institute of Food and Agriculture (AFRI Competitive Grant

1018617 and Hatch Program 1020852).

Acknowledgments

We thank Dr. Alice Russo and Ms. Rita Kurpiewski for field assistance, and Dr. Eve Slavich (UNSW

Stats Central) for statistical advice.

Data Availability Statement

Supplementary methods and results available online. Code available at:

https://github.com/jiazhou0116/gut-microbiome-analyses-2. Sequence data are available in NCBI

Sequence Read Archive (PRJNA670039). Raw ecological data available on Dryad

(doi:10.5061/dryad.v15dv41tw).

Bibliography

[1]   Hopkins WA. Amphibians as models for studying environmental change. ILAR J 2007;48:270–7. doi:10.1093/ilar.48.3.270.

[2]   Bletz MC, Goedbloed DJ, Sanchez E, Reinhardt T, Tebbe CC, Bhuju S, et al. Amphibian gut microbiota shifts differentially in community structure but converges on habitat-specific predicted functions. Nat Commun 2016;7:13699. doi:10.1038/ncomms13699.

[3]   Huang B-H, Chang C-W, Huang C-W, Gao J, Liao P-C. Composition and functional specialists of the gut microbiota of frogs reflect habitat differences and agricultural activity. Front Microbiol 2017;8:2670. doi:10.3389/fmicb.2017.02670.

[4]   Scott KP, Gratz SW, Sheridan PO, Flint HJ, Duncan SH. The influence of diet on the gut microbiota. Pharmacol Res 2013;69:52–60. doi:10.1016/j.phrs.2012.10.020.

[5]   Zmora N, Suez J, Elinav E. You are what you eat: diet, health and the gut microbiota. Nat Rev Gastroenterol Hepatol 2019;16:35–56. doi:10.1038/s41575-018-0061-2.

[6]   Bercik P, Denou E, Collins J, Jackson W, Lu J, Jury J, et al. The intestinal microbiota affect central levels of brain-derived neurotropic factor and behavior in mice. Gastroenterology 2011;141:599–609, 609.e1. doi:10.1053/j.gastro.2011.04.052.

[7]   Kraimi N, Calandreau L, Zemb O, Germain K, Dupont C, Velge P, et al. Effects of gut microbiota transfer on emotional reactivity in Japanese quails (Coturnix japonica). J Exp Biol 2019;222. doi:10.1242/jeb.202879.

[8]   Teseo S, van Zweden JS, Pontieri L, Kooij PW, Sørensen SJ, Wenseleers T, et al. The scent of symbiosis: gut bacteria may affect social interactions in leaf-cutting ants. Anim Behav 2019;150:239–54. doi:10.1016/j.anbehav.2018.12.017.

[9]   Schretter CE, Vielmetter J, Bartos I, Marka Z, Marka S, Argade S, et al. A gut microbial factor modulates locomotor behaviour in Drosophila. Nature 2018;563:402–6. doi:10.1038/s41586-018-0634-9.

[10]   Borrelli L, Aceto S, Agnisola C, De Paolo S, Dipineto L, Stilling RM, et al. Probiotic modulation of the microbiota-gut-brain axis and behaviour in zebrafish. Sci Rep 2016;6:30046. doi:10.1038/srep30046.

[11]   Savignac HM, Kiely B, Dinan TG, Cryan JF. Bifidobacteria exert strain-specific effects on stress-related behavior and physiology in BALB/c mice. Neurogastroenterol Motil 2014;26:1615–27. doi:10.1111/nmo.12427.

[12]   Sharon G, Segal D, Ringo JM, Hefetz A, Zilber-Rosenberg I, Rosenberg E. Commensal bacteria play a role in mating preference of Drosophila melanogaster. Proc Natl Acad Sci USA 2010;107:20051–6. doi:10.1073/pnas.1009906107.

[13]   Vogel H, Shukla SP, Engl T, Weiss B, Fischer R, Steiger S, et al. The digestive and defensive

basis of carcass utilization by the burying beetle and its microbiota. Nat Commun
2017;8:15186. doi:10.1038/ncomms15186.

[14]   Wong AC-N, Wang Q-P, Morimoto J, Senior AM, Lihoreau M, Neely GG, et al. Gut
Microbiota Modifies Olfactory-Guided Microbial Preferences and Foraging Decisions in
Drosophila. Curr Biol 2017;27:2397–2404.e4. doi:10.1016/j.cub.2017.07.022.

[15]   Jones JC, Fruciano C, Marchant J, Hildebrand F, Forslund S, Bork P, et al. The gut
microbiome is associated with behavioural task in honey bees. Insectes Soc 2018;65:419–29.
doi:10.1007/s00040-018-0624-9.

[16]   Davidson GL, Raulo A, Knowles SCL. Identifying Microbiome-Mediated Behaviour in Wild
Vertebrates. Trends Ecol Evol (Amst) 2020. doi:10.1016/j.tree.2020.06.014.

[17]   Stuart KC, Shine R, Brown GP. Proximate mechanisms underlying the rapid modification of
phenotypic traits in cane toads (*Rhinella marina*) across their invasive range within Australia.
Biological Journal of the Linnean Society 2019;126:68–79. doi:10.1093/biolinnean/bly150.

[18]   Wagener C, Mohanty NP, Measey J. The gut microbiome facilitates ecological adaptation in
an invasive vertebrate. BioRxiv 2020. doi:10.1101/2020.12.10.418954.

[19]   Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ. Is invasion success explained by the
enemy release hypothesis? Ecol Lett 2004;7:721–33. doi:10.1111/j.1461-0248.2004.00616.x.

[20]   Poulin R. Parasite manipulation of host behavior: an updateand frequently asked questions.
Advances In The Study Of Behavior, vol. 41, Elsevier; 2010, p. 151–86. doi:10.1016/S0065-
3454(10)41005-0.

[21]   Tingley R, Ward-Fear G, Schwarzkopf L, Greenlees MJ, Phillips BL, Brown G, et al. New
weapons in the toad toolkit: A review of methods to control and mitigate the biodiversity
impacts of invasive cane toads (*rhinella marina*). Q Rev Biol 2017;92:123–49.
doi:10.1086/692167.

[22]   Finnerty PB, Shine R, Brown GP. Survival of the feces: Does a nematode lungworm
adaptively manipulate the behavior of its cane toad host? Ecol Evol 2018;8:4606–18.
doi:10.1002/ece3.3870.

[23]   Shine R. The ecological impact of invasive cane toads (Bufo marinus) in Australia. Q Rev Biol
2010;85:253–91. doi:10.1086/655116.

[24]   Llewelyn J, Phillips BL, Alford RA, Schwarzkopf L, Shine R. Locomotor performance in an
invasive species: cane toads from the invasion front have greater endurance, but not speed,
compared to conspecifics from a long-colonised area. Oecologia 2010;162:343–8.
doi:10.1007/s00442-009-1471-1.

[25]   Hudson CM, Vidal-García M, Murray TG, Shine R. The accelerating anuran: evolution of
locomotor performance in cane toads (Rhinella marina, Bufonidae) at an invasion front. Proc R
Soc Lond, B, Biol Sci 2020;287:20201964. doi:10.1098/rspb.2020.1964.

[26]   Gruber J, Brown G, Whiting MJ, Shine R. Geographic divergence in dispersal-related

behaviour in cane toads from range-front versus range-core populations in Australia.
Behavioral Ecology And 2017.

[27] Phillips BL, Kelehear C, Pizzatto L, Brown GP, Barton D, Shine R. Parasites and pathogens
lag behind their host during periods of host range advance. Ecology 2010;91:872–81.
doi:10.1890/09-0530.1.

[28] Child T, Phillips BL, Shine R. Abiotic and biotic influences on the dispersal behavior of
metamorph cane toads (Bufo marinus) in tropical Australia. J Exp Zool A Ecol Genet Physiol
2008;309:215–24. doi:10.1002/jez.450.

[29] Selechnik D, Richardson MF, Shine R, DeVore JL, Ducatez S, Rollins LA. Increased adaptive
variation despite reduced overall genetic diversity in a rapidly adapting invader. Front Genet
2019;10:1221. doi:10.3389/fgene.2019.01221.

[30] Brown GP, Kelehear C, Pizzatto L, Shine R. The impact of lungworm parasites on rates of
dispersal of their anuran host, the invasive cane toad. Biol Invasions 2016;18:103–14.
doi:10.1007/s10530-015-0993-1.

[31] Hudson CM, Brown GP, Shine R. Evolutionary shifts in anti-predator responses of invasive
cane toads (Rhinella marina). Behav Ecol Sociobiol (Print) 2017;71:134. doi:10.1007/s00265-
017-2367-4.

[32] Kelehear C, Jones HI. Nematode larvae (order Spirurida) in gastric tissues of Australian
anurans: a comparison between the introduced cane toad and sympatric native frogs. J Wildl
Dis 2010;46:1126–40. doi:10.7589/0090-3558-46.4.1126.

[33] Baker MR. The free-living and parasitic development of *Rhabdias* spp. (Nematoda:
Rhabdiasidae) in amphibians. Can J Zool 1979;57:161–78. doi:10.1139/z79-014.

[34] Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet C, Al-Ghalith GA, et al. QIIME 2:
Reproducible, interactive, scalable, and extensible microbiome data science 2018.
doi:10.7287/peerj.preprints.27295v1.

[35] Douglas GM, Maffei VJ, Zaneveld J, Yurgel SN, Brown JR, Taylor CM, et al. PICRUSt2: An
improved and extensible approach for metagenome inference. BioRxiv 2019.
doi:10.1101/672295.

[36] R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria; 2020.

[37] Davidson GL, Cooke AC, Johnson CN, Quinn JL. The gut microbiome as a driver of
individual variation in cognition and functional behaviour. Philos Trans R Soc Lond B, Biol Sci
2018;373. doi:10.1098/rstb.2017.0286.

[38] Rollins LA, Richardson MF, Shine R. A genetic perspective on rapid evolution in cane toads
(Rhinella marina). Mol Ecol 2015;24:2264–76. doi:10.1111/mec.13184.

[39] Shine R. The impact of an invasive amphibian: the cane toad Rhinella marina. In: Heatwole H,
Rowley J, editors. Status of Conservation and Decline of Amphibians: Australia, New Zealand,

and Pacific Islands, Victoria, Australia: CSIRO Publishing; 2018, p. 107–24.

[40]   Shine R, Brown GP, Phillips BL. An evolutionary process that assembles phenotypes through
space rather than through time. Proc Natl Acad Sci USA 2011;108:5708–11.
doi:10.1073/pnas.1018989108.

[41]   Brown GP, Kelehear C, Shine R. The early toad gets the worm: cane toads at an invasion front
benefit from higher prey availability. J Anim Ecol 2013;82:854–62. doi:10.1111/1365-
2656.12048.

[42]   Kosmala GK, Brown GP, Shine R. Thin-skinned invaders: geographic variation in the
structure of the skin among populations of cane toads (Rhinella marina). Biological Journal of
the Linnean Society 2020;131:611–21. doi:10.1093/biolinnean/blaa128.

[43]   Weitzman CL, Kaestli M, Gibb K, Brown GP, Shine R, Christian K. Disease exposure and
antifungal bacteria on skin of invasive cane toads, australia. Emerging Infect Dis
2019;25:1770–1. doi:10.3201/eid2509.190386.

[44]   Zhou J, Tesfamicael K, Zhou SJ, Rollins LA, Rodriguez Lopez CM. Genetic similarity
enhances the strength of the relationship between gut bacteria and host DNA methylation.
BioRxiv 2021. doi:10.1101/2021.07.10.451923.

[45]   Zhang X-Y, Sukhchuluun G, Bo T-B, Chi Q-S, Yang J-J, Chen B, et al. Huddling remodels gut
microbiota to reduce energy requirements in a small mammal species during cold exposure.
Microbiome 2018;6:103. doi:10.1186/s40168-018-0473-9.

[46]   Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota.
Environ Microbiol 2017;19:29–41. doi:10.1111/1462-2920.13589.

[47]   Gamage HKAH, Tetu SG, Chong RWW, Ashton J, Packer NH, Paulsen IT. Cereal products
derived from wheat, sorghum, rice and oats alter the infant gut microbiota in vitro. Sci Rep
2017;7:14312. doi:10.1038/s41598-017-14707-z.

[48]   Thomas RH, Meeking MM, Mepham JR, Tichenoff L, Possmayer F, Liu S, et al. The enteric
bacterial metabolite propionic acid alters brain and plasma phospholipid molecular species:
further development of a rodent model of autism spectrum disorders. J Neuroinflammation
2012;9:153. doi:10.1186/1742-2094-9-153.

[49]   Hugenholtz F, Davids M, Schwarz J, Müller M, Tomé D, Schaap P, et al. Metatranscriptome
analysis of the microbial fermentation of dietary milk proteins in the murine gut. PLoS One
2018;13:e0194066. doi:10.1371/journal.pone.0194066.

[50]   Kang D-W, Park JG, Ilhan ZE, Wallstrom G, Labaer J, Adams JB, et al. Reduced incidence of
Prevotella and other fermenters in intestinal microflora of autistic children. PLoS One
2013;8:e68322. doi:10.1371/journal.pone.0068322.

[51]   González-Bernal E, Brown GP, Shine R. Invasive cane toads: social facilitation depends upon
an individual's personality. PLoS One 2014;9:e102880. doi:10.1371/journal.pone.0102880.

[52]   Gelbard-Sagiv H, Magidov E, Sharon H, Hendler T, Nir Y. Noradrenaline modulates visual

perception and late visually evoked activity. Curr Biol 2018;28:2239–2249.e6.
doi:10.1016/j.cub.2018.05.051.

[53]   Houlden A, Goldrick M, Brough D, Vizi ES, Lénárt N, Martinecz B, et al. Brain injury induces
specific changes in the caecal microbiota of mice via altered autonomic activity and
mucoprotein production. Brain Behav Immun 2016;57:10–20. doi:10.1016/j.bbi.2016.04.003.

[54]   Ning T, Gong X, Xie L, Ma B. Gut Microbiota Analysis in Rats with Methamphetamine-
Induced Conditioned Place Preference. Front Microbiol 2017;8:1620.
doi:10.3389/fmicb.2017.01620.

[55]   Choi JJ, Eum SY, Rampersaud E, Daunert S, Abreu MT, Toborek M. Exercise attenuates
PCB-induced changes in the mouse gut microbiome. Environ Health Perspect 2013;121:725–
30. doi:10.1289/ehp.1306534.

[56]   Golla A, Østby H, Kermen F. Chronic unpredictable stress induces anxiety-like behaviors in
young zebrafish. Sci Rep 2020;10:10339. doi:10.1038/s41598-020-67182-4.

[57]   Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and
resilience of the human gut microbiota. Nature 2012;489:220–30. doi:10.1038/nature11550.

[58]   West TP. Degradation of pyrimidine ribonucleosides by Pseudomonas aeruginosa. Antonie
Van Leeuwenhoek 1996;69:331–5. doi:10.1007/bf00399622.

[59]   Garavaglia M, Rossi E, Landini P. The pyrimidine nucleotide biosynthetic pathway modulates
production of biofilm determinants in Escherichia coli. PLoS One 2012;7:e31252.
doi:10.1371/journal.pone.0031252.

[60]   Bashiri G, Antoney J, Jirgis ENM, Shah MV, Ney B, Copp J, et al. A revised biosynthetic
pathway for the cofactor F420 in prokaryotes. Nat Commun 2019;10:1558.
doi:10.1038/s41467-019-09534-x.

[61]   Sibon OCM, Strauss E. Coenzyme A: to make it or uptake it? Nat Rev Mol Cell Biol
2016;17:605–6. doi:10.1038/nrm.2016.110.

[62]   Varki A, Gagneux P. Multifarious roles of sialic acids in immunity. Ann N Y Acad Sci
2012;1253:16–36. doi:10.1111/j.1749-6632.2012.06517.x.

[63]   Liao H, Klaus C, Neumann H. Control of innate immunity by sialic acids in the nervous tissue.
Int J Mol Sci 2020;21. doi:10.3390/ijms21155494.

[64]   Sinnott-Armstrong N, Naqvi S, Rivas MA, Pritchard JK. GWAS of three molecular traits
highlights core genes and pathways alongside a highly polygenic background. BioRxiv 2020.
doi:10.1101/2020.04.20.051631.

[65]   Brown GP, Shilton C, Phillips BL, Shine R. Invasion, stress, and spinal arthritis in cane toads.
Proc Natl Acad Sci USA 2007;104:17698–700. doi:10.1073/pnas.0705057104.

[66]   House PK, Vyas A, Sapolsky R. Predator cat odors activate sexual arousal pathways in brains
of Toxoplasma gondii infected rats. PLoS One 2011;6:e23277.

doi:10.1371/journal.pone.0023277.

[67]   Poulin R. Parasite manipulation of host behavior: an updateand frequently asked questions.
528   Advances In The Study Of Behavior, vol. 41, Elsevier; 2010, p. 151–86. doi:10.1016/S0065-
529   3454(10)41005-0.

530   [68]   Gegear RJ, Otterstatter MC, Thomson JD. Bumble-bee foragers infected by a gut parasite have
531   an impaired ability to utilize floral information. Proc Biol Sci 2006;273:1073–8.
532   doi:10.1098/rspb.2005.3423.

533   [69]   Pizzatto L, Shine R. Lungworm infection modifies cardiac response to exercise in cane toads. J
534   Zool 2012;287:150–5. doi:10.1111/j.1469-7998.2012.00898.x.

535   [70]   Finnerty PB. The physiological and ecological impacts of lungworm infection on invasive cane
536   toads. Undergraduate thesis. University of Sydney, 2017.

537   [71]   Abada EA, Sung H, Dwivedi M, Park B-J, Lee S-K, Ahnn J. C. elegans behavior of preference
538   choice on bacterial food. Mol Cells 2009;28:209–13. doi:10.1007/s10059-009-0124-x.

539   [72]   Zheng D, Liwinski T, Elinav E. Interaction between microbiota and immunity in health and
540   disease. Cell Res 2020;30:492–506. doi:10.1038/s41422-020-0332-7.

541   [73]   Brown GP, Kelehear C, Shilton CM, Phillips BL, Shine R. Stress and immunity at the invasion
542   front: a comparison across cane toad (*Rhinella marina*) populations. Biological Journal of the
543   Linnean Society 2015;116:748–60. doi:10.1111/bij.12623.

544   [74]   Hudson CM, Brown GP, Shine R. Evolutionary shifts in anti-predator responses of invasive
545   cane toads (Rhinella marina). Behav Ecol Sociobiol (Print) 2017;71:134. doi:10.1007/s00265-
546   017-2367-4.

547   [75]   Zhang C, Zhang M, Wang S, Han R, Cao Y, Hua W, et al. Interactions between gut
548   microbiota, host genetics and diet relevant to development of metabolic syndromes in mice.
549   ISME J 2010;4:232–41. doi:10.1038/ismej.2009.112.

550   [76]   Fan P, Bian B, Teng L, Nelson CD, Driver J, Elzo MA, et al. Host genetic effects upon the
551   early gut microbiota in a bovine model with graduated spectrum of genetic variation. ISME J
552   2020;14:302–17. doi:10.1038/s41396-019-0529-2.

553   [77]   Arnaud-Haond S, Aires T, Candeias R, Teixeira SJL, Duarte CM, Valero M, et al. Entangled
554   fates of holobiont genomes during invasion: nested bacterial and host diversities in Caulerpa
555   taxifolia. Mol Ecol 2017;26:2379–91. doi:10.1111/mec.14030.

556   [Preprint]   Zhou J, Nelson TM, Rodriguez Lopez C, Zhou SJ, Ward-Fear G, Stuart KC, et al. The gut
557   bacteria of an invasive amphibian respond to the dual challenges of range-expansion and
558   parasite attack. BioRxiv 2020. doi:10.1101/2020.11.16.385690.

562 **Tables and Figures**

563 **Table 1a-d.** The association between a) single host factors and microbial community, b) single host
564 factors and predicted microbial function, c) single behavioural trait and microbial community and d)
565 single behavioural trait and predicted microbial function. Significant p-values denoted by an asterisk.

	MDS1	MDS2	r2	Pr(>r)
a. Host factor/microbial community				
SUL	0.170	0.985	0.071	0.119
BodyWeight	-0.009	1.000	0.051	0.231
Lungworms	0.254	-0.967	0.020	0.585
Occurrence of lungworms	0.453	-0.892	0.128	0.023*
b. Host factor/microbial function				
SUL	0.680	0.733	0.021	0.556
BodyWeight	0.610	0.793	0.041	0.311
Lungworms	-0.791	0.612	0.007	0.839
Occurrence of lungworms	-0.827	-0.562	0.059	0.187
c. Behavioural trait/microbial community				
Struggle score	-0.814	0.580	0.044	0.268
Struggle likelihood	-0.880	-0.474	0.021	0.555
Righting effort	-0.124	0.992	0.075	0.121
Righting effort likelihood	0.499	0.867	0.0255	0.474
Righting time	0.933	0.360	0.0615	0.174
d. Behavioural trait/microbial function				

Struggle score	0.376	0.927	0.051	0.221
Struggle likelihood	0.183	0.983	0.017	0.606
Righting effort	0.571	0.821	0.070	0.141
Righting effort likelihood	0.399	0.917	0.035	0.362
Righting time	-0.956	-0.294	0.004	0.894

569 **Table 2.** Association between gut microbiota variation and lungworm occurrence and intensity, based
570 on Bray Curtis dissimilarity values for microbial community assemblages.

	Df	SumOfSqs	R2	F	Pr(>F)
Location	1	1.314	0.091	6.026	<0.001***
Lungworm_occurrence	1	0.482	0.033	2.213	0.005**
Lungworms_intensity	1	0.150	0.010	0.688	0.861
Location:Lungworm_occurrence	1	0.331	0.023	1.520	0.075
Location:Lungworm_intensity	1	0.357	0.025	1.638	0.043*
Residual	54	11.774	0.817		
Total	59	14.409	1.000		

Significance codes: 0 '***' ≤0.001 '**' ≤0.01 '*' ≤0.05.

Figure 1. Beta diversity by location. Principle coordinate analysis plot of Bray Curtis distance of microbial community (A) and Bray Curtis distance of predicted functional groups (B) from 60 cane toad individuals of the invasion-front (Kununurra, Old Theda, and Mary Pool) and the range-core (Rossville, Croydon, and Lucinda).

Figure 2. Significantly different bacterial taxa and predicted functions between range-core
(QLD) and invasion-front (WA) toads' colon. Significant differences were identified between
locations via differential abundance testing based on a negative binomial distribution. The dots
represent the average log-2 fold change (x axis) abundance and positive $\log_2$ fold changes signify
increased abundance in range-core, and negative $\log_2$ fold changes display increased abundance in
invasion-front. Bacterial taxa (A) were classified to the taxonomic level of family (y axis) and
coloured by taxonomic level of phylum. Family name in bracket is proposed taxonomy by
Greengenes. Only ASVs that could be matched to a known bacterial family and with a
log2FoldChange value higher than 20 or lower than -20 are presented. Predicted functions (B) with a
log2FoldChange value higher than 3 or lower than -3 are presented.

Figure 3. Heatmap for top 30 functional group abundance.

Heatmap indicates the top 30 functional groups in the intestinal samples from range-core and
invasion-front toads. Abundance indicates the raw count of functional groups inferred from
taxonomic 16S sequences using PICRUSt where light blue is high abundance and dark blue is lower
abundance. Functional pathways that differ significantly between range-core and invasion-front toads
are highlighted in bold. Range-core includes Rossville, Croyden, and Lucinda; invasion-front
includes Kununurra, Old Theda, and Mary Pool.

Figure 4. Main variables that affect predicted function differentiation among individuals. CAP (*capscale*) plot displays the combination of variables that explained the greatest variation in the predicted functions through model selection, using 60 cane toad individuals from the invasion-front (Kununurra, Old Theda, and Mary Pool) and the range-core (Rossville, Croydon, and Lucinda). The final model explained 17.8% of variation in the microbial predicted functions, which includes righting effort likelihood (AIC = 53.613), occurrence of lung worms (AIC = 54.297) and righting time (AIC = 56.912) explained the greatest variation.

A

Family

Verrucomicrobiaceae

Rur

Rh

Porph

P

La

Erysi

Enter

Bacteroidaceae

Bacillaceae

[Odoribacteraceae]

[Mogibacteriaceae]

-20   0   20

Log$_2$ fold change, p.value=0.05

Phylum

● Firmicutes

● Bacteroidetes

● Proteobacteria

● Verrucomicrobia

B

Predicted Functions: range-core vs invasion-front

Log_2 fold change, p.value=0.05

righting_effort_likelihood

Lungworms_occurrence

righting_time

Site

Kununurra

Old Theda

Mary Pool

Rossville

Croyden

Lucinda