SEX-SPECIFIC PLASTICITY AND THE NUTRITIONAL GEOMETRY OF INSULIN-

SIGNALING GENE EXPRESSION IN *DROSOPHILA MELANOGASTER*

Jeanne M.C. McDonald[1,2]*, Pegah Nabili[2]*, Lily Thorsen[2]*, Alexander Shingleton[2,3]

[1] Department of Ecology and Evolutionary Biology, Cornell University, Corson Hall Ithaca, NY 14853

[2] Department of Biology, Lake Forest College, 555 North Sheridan Road, Lake Forest, IL 60045

[3] Department of Biological Sciences, University of Illinois at Chicago, 840 W Taylor St, Chicago, IL 60607

* These authors contributed equally to this study

Keywords: Sexual-size dimorphism, sex-specific plasticity, nutrition, body size, insulin/IGF-signaling,

TOR-signaling

1. Abstract

Background: Sexual-size dimorphism (SSD) is replete among animals, but while the selective pressures that drive the evolution of SSD have been well studied, the developmental mechanisms upon which these pressures act are poorly understood. Ours and others' research has shown that SSD in *Drosophila* reflects elevated levels of nutritional plasticity in females versus males, such that SSD increases with dietary intake and body size, a phenomenon called sex-specific plasticity (SSP). Additional data indicate that while body size in both sexes responds to variation in protein level, only female body size is sensitive to variation in carbohydrate level. Here we explore whether these difference in sensitivity at the morphological level are reflected by differences in how the insulin/IGF-signaling (IIS) and TOR-signaling pathways respond to changes in carbohydrates and proteins in females versus males, using a nutritional geometry approach.

Results: The IIS-regulated transcripts of *4E-BP* and *InR* most strongly correlated with body size in females and males respectively, but neither responded to carbohydrate level and so could not explain the sex-specific response to body size to dietary carbohydrate. Transcripts regulated by TOR-signaling did, however, respond to dietary carbohydrate in a sex-specific manner. In females, expression of *dILP5* positively correlated with body size, while expression of *dILP2,3* and *8,* was elevated on diets with a low concentration of both carbohydrate and protein. In contrast, we detected lower levels of dILP2 and 5 protein in the brains of females fed on low concentration diets. We could not detect any effect of diet on *dILP* expression in males.

Conclusion: Although females and males show sex-specific transcriptional responses to changes in protein and carbohydrate, the patterns of expression do not support a simple model of the regulation of body-size SSP by either insulin- or TOR-signaling. The data also indicate a complex relationship between carbohydrate and protein level, *dILP* expression and dILP peptide levels in the brain. In general, diet

quality and sex both affect the transcriptional response to changes in diet quantity, and so should be

considered in future studies that explore the effect of nutrition on body size.

2.   Background

Sexual Size Dimorphism (SSD), the difference in body size between males and females, is perhaps the

most familiar and widespread form of sexual dimorphism. This condition is extremely variable among

species. For example, a female blanket octopi can weigh 10,000- 20,000 times more than male [1], while a

male southern elephant seals can weigh seven times more than a female [2]. Further, the degree of SSD is

highly evolutionarily labile, and can vary between closely related species or among populations within

species, sometimes dramatically. For example, among populations of the Australian carpet python,

females range from being less than 1.5x to more than 10x the size of males [3]. While intraspecific

variation in SSD is likely due, in part, to genetic differences between populations, it may also be a

consequence of sex-specific differences in the phenotypic plasticity of body size. Environmental variation

can account for the vast majority of variation in body size within a population [4, 5], and if males and

females differ in the extent to which the environment affects body size, this will generate changes in SSD

across environments [6]. While there is considerable evidence that males and females differ in the extent

of their body size response to a variety of environmental variables – a phenomenon called *Sex-Specific*

Plasticity (SSP)– the role that SSP plays in the developmental generation and evolution of sexual size

dimorphism is largely unknown. This is compounded by a general lack of knowledge regarding the

proximate developmental mechanisms that generate differences in body size between males and females.

Perhaps the most important, and certainly the best understood, environmental factor that regulates body

size is developmental nutrition. In all animals where it has been studied, the nutritional regulation of

growth is mediated via the insulin/IGF-signaling (IIS) pathway, a highly conserved receptor tyrosine

kinase pathway, the components of which pre-date the Metazoa [7]. Insulin-like peptides are released in

response to nutrition and bind to receptors of dividing cells to initiate a signal-transduction cascade that

regulates the expression of positive and negative growth regulator. The IIS pathway positively regulates

the activity of TOR-signaling, which also responds directly to cellular levels of amino acid. Activation of

IIS/TOR-signaling results in an increase in both growth rate and (typically) adult body size at high levels

of nutrition, while deactivation of IIS/TOR-signaling does the opposite at low levels of nutrition.

Although both males and female typically display nutritional plasticity, in many species one sex is more

nutritionally plastic than the other. In birds and mammals, male growth tends to be affected more by

nutritional stress than female growth [8], while in arthropods, the reverse appears to be true [9]. This is

accompanied by the general trend of male-biased SSD in birds and mammals versus female-biased SSD in

arthropods [10, 11]. Generally, therefore, the larger sex tends to be more nutritionally plastic than the

smaller sex, and SSD increases with body size within a population [11]. This suggests that the

mechanisms that generate SSD at least partially overlap with those that generate sex-specific nutritional

plasticity.

Several adaptive hypotheses that explain why selection acts differently on males versus females may

explain a relationship between SSD and SSP [9]. Under the adaptive canalization hypothesis [12], traits

that are under strong stabilizing selection should be more canalized and less plastic than traits that are

not. Thus, if there were strong stabilizing selection on the body size of one sex but not the other, then this

could lead to differences in size plasticity. In contrast, the condition dependence hypothesis predicts that

traits under strong directional sexual selection will exhibit greater sensitivity to environmental condition

[13, 14] Consistent with this latter hypothesis is the observation that exaggerated traits used by males to

compete for and attract mates are often typically highly plastic relative to the same trait in females (e.g.

[14]). The condition-dependence hypothesis was formulated to explain the condition dependence of

individual traits within the body and their response to directional sexual selection. Whether it can explain

the correlation between SSP and SSD for the body as a whole, and where body size is not obviously

subject to sexual selection (for example for female-biased SSD) is unclear.

While these adaptive hypotheses provide an ultimate evolutionary mechanism for a correlation between

SSD and SSP, several studies have hinted at the proximate developmental mechanism linking the two

phenomena. A study by Emlen et al [15] found that exaggerated male traits – specifically the horns of the

beetle *Trypoxylus dichotomus* – are more insulin sensitive than other traits in the body. This accounted for

both the trait's increase in size in males relative to females, and their elevated nutritional plasticity.

Consequently, the trait becomes more sexually dimorphic as trait size increases. In *Drosophila*

melanogaster, where females are larger and more nutritionally plastic than males, SSD also appears to be

regulated by the IIS pathway. This is based on three pieces of evidence: (1) SSD is eliminated in flies with

suppressed IIS through hypomorphic mutation of the insulin receptor (*InR*) [16]; (2) well-fed females

have higher IIS activity than males [17], and; (3) SSD requires sex-specific difference in the neurosecretory

cells that produce insulin-like peptides (dILPs), the hormone that activates the IIS pathway [18]. Because

SSD is eliminated in flies with a loss of InR activity, it follows that female body size is more sensitive to

changes in insulin-signaling than male body size. Consequently, as for the beetle horn, SSP of body size

in *Drosophila* may result from sex-specific differences in insulin-sensitivity. What is unknown, however, is

whether the differences in nutritional plasticity between males and females is evident at the level of

IIS/TOR-signaling activity itself.

To address this question, we looked at the expression of genes that regulate and are transcriptionally

regulated by the IIS and TOR-signaling pathways across a nutritional landscape, where nutritional

quality and quantity varies. In a previous study [19], we looked at the effect of changes in protein-to-

carbohydrate ratio (diet quality) and total food concentration (diet quantity) on the size of the wing,

maxillary palp, femur of the first leg and thorax. For all traits, female trait size was more sensitive to

changes in either diet quality or quantity than male trait size. Intriguingly, however, these sex-specific

differences could be attributed to the each trait's size response to changes in carbohydrate versus protein

concentration: males and female trait sizes were equally sensitive to changes in protein concentration, but

only female trait size was detectibly sensitive to changes in carbohydrate concentration. Here we test the

hypothesis that these observed sex-specific differences in body-size plasticity are reflected by

corresponding differences in response to diet at the level of the IIS/TOR-signaling pathways.

118   3.   Results

3.1.   Female and male body size responds differently to changes in carbohydrates but not

proteins.

We found that, consistent with prior results [19], males and females differed in their response to dietary

carbohydrate versus protein (Figure 1, Supplementary Figure 1, Table 1). Specifically, both male and

female body size responded to changes in dietary protein as a negative quadratic, with body size

increasing as protein concentration increased, but at a decreasing rate (Table 1). In contrast, only female

body size responded to dietary carbohydrate concentration, this time as a positive quadratic, such that

body size declined with increasing carbohydrate, but at a decreasing rate (Table 1). Correspondingly,

including an interaction between sex and carbohydrate when modeling the relationship between diet and

body size significantly improved model fit, while including an interaction between sex and protein did

not further improve fit (Table 2). Thus male and female body size appears to respond differently to

carbohydrate but not to protein. As a result, sexual size dimorphism (SSD) varied across the nutritional

geometry landscape (Figure 1, Supplementary Figure 1).

Figure 1: The effect of protein and carbohydrate concentration on female and male body size and

sexual size dimorphism (SSD). (A, B) Surfaces show the fitted relationship between body size,

carbohydrate level and protein level in female and male flies, based on the statistical model specified by

the equation above each chart. T = body size, P = protein, C = carbohydrate, ε = error, subscripts refer to

levels within each factor. (C) Surface shows the difference in female and male body size (SSD) across the

same nutritional landscape, using fitted values from A and B. Points indicate diets tested and dotted lines

connect diets with equal protein-to-carbohydrate ratios (1:14.6, 1:7.2, 1:3.5, 1:1.7, 1.3:1, 1.4 :1). * indicates

approximate composition of standard cornmeal-molasses medium. Corresponding thin-plate spline plots

are shown in Supplementary Figure 1.

3.2.   Female and male IIS/TOR gene expression responds differently to changes in diet

We collected expression data for eight genes that either regulate or are regulated by the IIS/TOR signaling

pathway: insulin receptor (*InR*); eukaryotic translation initiation factor 4E binding protein (*4E-BP/Thor*);

Drosophila insulin like peptides 2, 3, 5 and 8 (*dILP 2, 3, 5,8),* Absent, small, or homeotic discs-like protein

(*Ash2L*), and *CG3071* (the Drosophila homolog of UTP15/SAW). Expression of both *InR* and *4E-BP* is

negatively regulated by the IIS pathway [26], *dILPs 2,3,5* encode peptides that ostensibly activate the IIS

pathway, while *dILP8* retards development by inhibiting the synthesis and release of ecdysone [27].

Finally expression of both *Ash2L* and *CG3071* has been shown to be negatively and positive regulated by

TOR signaling, respectively [28]. Expression of only a few of these genes correlated directly with body

size: *4EBP* and *dILP5* negatively and positively correlated with body size in females, respectively, while

expression of *InR* negatively correlated with body size in males (Table 3). Nevertheless, a multivariate

analysis of gene expression across all diets supported a significant interaction between the effects of sex

and carbohydrate, as a linear factor, and sex and protein, as a quadratic factor, on gene expression (Table

4). We therefore explored how sex and diet affected the expression of individual genes. Specifically, we

looked to see whether sex-specific responses in the expression of genes that either regulate or are

regulated by the IIS/TOR-signalling pathways reflected the sex-specific response of body size to changes

in diet.

Genes Regulated by IIS*.* Both *InR* and *4E-BP* expression responded to changes only in dietary protein but

not carbohydrate (Figure 2). *InR* expression in males and females had a positive quadratic response to

protein level, decreasing as protein level increased, but at a declining rate (Table 5). The expression of *InR*

was lower across all diets in males than in females, but there was no significant interaction between the

effects of sex and protein level on gene expression (Table 5). In contrast, *4E-BP* expression was sensitive

to protein level only in females, decreasing linearly as protein increased, but was unaffected by diet in

males (Table 4). Consequently, there was a significant interaction between the effect of sex and protein

level on *4E-BP* expression (Table 5).

Genes Regulated by TOR Signaling. Expression of both *Ash2L* and *CG3071* had a complex and sex-

specific response to diet. Carbohydrate significantly affected *Ash2L* expression as a linear relationship,

but in opposite directions in either sex: Increased carbohydrate increased *Ash2L* expression in males but

decreased *Ash2L* expression in females (Figure 2, Table 4). *Ash2L* expression was not affected by protein

in males, but had a positive quadratic response to protein in females, decreasing as protein increased but

at a declining rate. There was therefore a significant interaction between sex and diet on the expression of

Ash2L, although only between sex and carbohydrate and not sex and protein (Table 5). *CG3071* also

showed a sex-specific expression response to diet. In females, *CG3071* expression had a negative

quadratic response to protein level (increasing at a declining rate as protein increased), a positive

quadratic response to carbohydrate level (decreasing at a declining rate as carbohydrate increased), with

a significant carbohydrate: protein interaction (Table 4). In contrast, *CG3071* expression was not affected

by diet in males, which lead to a significant diet-by-sex interaction effect on *CG3071* expression (Table 5).

Figure 2: The effect of protein and carbohydrate concentration on the expression of IIS and TOR
transcriptionally-regulated genes in females and males. Surfaces show the fitted relationship between
gene expression, carbohydrate level and protein level in female and male flies, based on the statistical
model specified by the equation above each chart (Table 4). *P* = protein, *C* = carbohydrate, ε = error,
subscripts refer to levels within each factor. Expression of (A, A') *InR*, and (B, B') *4E-BP*, both negatively
regulated by the activity of the IIS via the Forkhead transcription factor FOXO. (C, C') Expression of
Ash2L, ostensibly negatively regulated by the activity of TOR signaling. (D, D') Expression of *CG3071*,
ostensibly positively regulated by the activity of TOR signaling. Points indicate diets tested and dotted
lines connect diets with equal protein-to-carbohydrate ratios (1:16, 1:8, 1:4, 1:2, 1:1, 2:1). Corresponding
thin-plate spline plots are shown in Supplementary Figure 2.

Genes That Regulate IIS/TOR Signaling. The expression of *dILP2* showed a significant response to diet in

females, declining linearly with increasing protein, and at a decreasing rate with increasing carbohydrate,

as a positive quadratic. There was also a significant carbohydrate-by-protein interaction effect on *dILP2*

expression in females. There was no significant effect of diet on *dILP2* expression in males, although this

appears to be a consequence of low statistical power, due to large amount of within-diet variation in

dILP2 expression among male samples compared to females sample. Fitting a one-way ANOVA of *dILP2*

expression against diet, where diet is each protein-to-carbohydrate ratio at each concentration, the

residual variance (a measure of average variance within diets) was significantly higher in males than in

females, by a ratio of 8.68 (Supplementary Table 1). Correspondingly, we had very low power (0.22) to

detect a significant sex-by-diet interaction for *dILP2* expression, based on the observed effect size

(Cohen's f^2 = 0.023, see Supplementary Material for details). Further, there was no evidence that

expression of *dILP2* was different between males and females, independent of diet (Table 5).

♀ ♂

A $dILP2_{ijk} = -C_i + C_i^2 - P_j + C_i{:}P_j + \varepsilon_{ijk}$ A′   dILP2

B $dILP3_{jk} = -P_j + \varepsilon_{jk}$ B′   dILP3

C $dILP5_{ijk} = -C_i + P_j - P_j^2 + \varepsilon_{ijk}$ C′   dILP5

D $dILP8_{jk} = -P_j + P_j^2 + \varepsilon_{jk}$ D′   dILP8

Figure 3: The effect of protein and carbohydrate concentration on the expression of dILPs in females and males. Surfaces show the fitted relationship between gene expression, carbohydrate level and protein level in female and male flies, based on the statistical model specified by the equation above each chart (Table 4). P = protein, C = carbohydrate, ε = error, subscripts refer to levels within each factor. (A) *dILP2.* (B) *dILP3.* (C) *dILP 5.* (D) *dILP8.* Points indicate diets tested and dotted lines connect diets with equal protein-to-carbohydrate ratios (1:16, 1:8, 1:4, 1:2, 1:1, 2:1). Corresponding thin-plate spline plots are shown in Supplementary Figure 3.

The expression of *dILP3* decreased linearly with protein in females, but there was no detectable effect of

diet on *dILP3* expression in males (Figure 2, Table 4). There was also no detectable difference in

expression between the sexes (Table 5). As for *dILP2*, variation in expression of *dILP3* among samples

within diets was significantly higher in males than in females (Supplementary Table 1), and a power

analysis gave a power of only 0.05 to detect a sex-by-protein interaction on *dILP3* expression, based on the

observed effect size (Cohen's f^2 <0.001).

The expression of *dILP5* was only marginally affected by diet in females, with a significant negative

quadratic effect of protein. We could detect no significant effect of either carbohydrate or protein on

dILP5 expression in males. As for *dILP2* and *3*, *dILP5* expression was more variable among male samples

within diets than among female samples (Supplementary Table 1), and a power analysis gave a power of

only 0.23 to detect a sex-by-protein interaction on *dILP5* expression, based on the observed effect size

(Cohen's f^2 = 0.022). Nevertheless, there was sex-specific expression independent of diet, such that *dILP5*

expression was higher in males than in females (Table 5).

The expression of *dILP8* responded as a positive quadratic in response to protein in females, declining as

protein increased at a decreasing rate, but did not show any response to any aspect of diet in males

(Figure 2, Table 4). The lack of a corresponding response to protein in males was not due to a lack of

statistical power: a power analysis suggested that the power to detect the female effect size of protein on

dILP8 expression (Cohen's f^2 = 0.46) in males was 0.99. Further, *dILP8* expression was not more variable

among male samples within diets than among female samples (Supplementary Table 1). Consequently,

we detected a significant sex-by-protein interaction on *dILP8* expression (Table 5).

3.3.   dILP protein levels in the brain increase with increasing food concentration

Our finding that *dILP* expression in females was generally higher in larvae fed on lower food

concentrations was surprising, given that dILPs (apart from dILP8) are canonically positive regulators of

growth in response to nutrition. We therefore explored the relationship between diet, *dILP* expression

and the levels of dILP peptide in the insulin-producing cells (IPCs) of the brain. Staining for dILP2 and

dILP5 peptide was higher in larvae fed a higher food concentration (360g/l) versus a lower food

concentration (45g/l), with a 1:2 protein: carbohydrate ratio (Figure 4).

Figure 4: The effect of diet on dILP2/5 expression and peptide levels in the brain in females and males.
(A,B) The relationship between diet and *dILP2* and *dILP5* expression. Points indicate 45 and 360 g/l food
concentration at 1:2 protein: carbohydrate ratio. (A', B') Representative images of dILP2 and dILP5 levels
in the insulin-producing cells (IPCs) of female larvae reared on different diets. (A″, B″) Mean normalized
level of dILP2 and dILP5 in the IPCs of larvae reared on different diets. *P*-values are for a pooled t-test.

4.   Discussion

Previous studies indicate that in *Drosophila,* body size is more nutritionally plastic in females than in

males, a phenomenon referred to as sex-specific plasticity (SSP) [19]. Nutritional plasticity is canonically

regulated by the insulin/IGF-signaling and TOR signaling pathways in *Drosophila* and most other

animals. We hypothesized, therefore, that differences in nutritional plasticity between the sexes reflects

sex-specific differences in the activation and activity of these pathways. In this study, we applied a

nutritional geometry framework to test this hypothesis, and explored the sex-specific effects of protein

and carbohydrates on body size and the expression of genes that activate and are transcriptionally

regulated by IIS/TOR. Our results indicate that, broadly, gene expression is more nutritionally sensitive in

females than in males, although the effect of nutrition on the expression of IIS and TOR-signaling genes is

both quantitively and qualitatively different between males and females.

The simplest hypothesis to account for sex-specific differences in the nutritional plasticity of body size in

Drosophila is that females have the same response to nutrition as males, but at a higher sensitivity. Under

this hypothesis, the response surfaces of body size and gene expression across a nutritional landscape of

varying protein and carbohydrate concentrations should be the same shape in both sexes, but with lower

gradients in males. This would be evident as lower parameter values in males versus females. This is not

supported by the data, however. In particular, body size in males shows more or less the same sensitivity

to changes in protein concentration as females, but has no detectable sensitivity to changes in

carbohydrate concentration. This in turn suggests that an important aspect of sex-specific nutritional

plasticity in *Drosophila* is differences in the response to variation in dietary carbohydrates.

The best understood regulator of body size with respect to nutrition are the insulin/IGF-signaling and

TOR-signaling pathways [29, 30]. The IIS pathway is activated by circulating dILPs, some of which are

released in a nutrition-dependent manner [27]. The TOR-signaling pathway is in part regulated by IIS,

but also responds directly to circulating amino acids [31–33]. Because the SSP of body size in flies appears

to primarily driven by a differential response to carbohydrates but not to protein, we might expect that

the effect of diet on body size is mediated primarily by IIS rather than TOR-signaling. Our data do not

support this hypothesis. We used the expression of *InR* and *4E-BP* as a measure of IIS activity: Both are

transcriptionally regulated by the forkhead transcription factor FOXO, which is activated when nutrition

and IIS activity is low [26, 34]. If sex-specific plasticity is mediated by IIS then we would expect to see sex-

specific differences in *Inr* and *4EBP* expression in response to changes in carbohydrate but not to protein.

In both males and females, however, *InR* expression responded only to protein level – and to the same

extant – but did not respond to carbohydrate level. Expression of *4E-BP* only responded to protein level

in females, but did not respond to carbohydrate level in either sex. Collectively, variation in the

expression of *4E-BP* and *InR* were the strongest correlates with body size in females and males

respectively (Table 3), consistent with the hypothesis that the IIS is the major regulator of body size with

respect to nutrition in *Drosophila*. However, the expression of these genes do not reflect sex-specific

differences in the effect of carbohydrate on body size, and so do not explain sex-specific plasticity.

In contrast to *InR* and *4E-BP*, the expression of *Ash2L* and *CG3071* both responded to carbohydrates in a

sex-specific manner. However, the response is not consistent with a previous report on the regulation of

the these genes' expression by TOR-signaling [28]. This study reported that *Ash2L* is negatively regulated

by TOR-signaling while *CG3071* is positively regulated. If TOR-signaling increases with protein, then we

would expect *Ash2L* expression to decrease correspondingly, which is true in females. However, *Ash2L*

expression also decreases with increasing carbohydrate, which is difficult to reconcile with the negative

effects that carbohydrate has on female growth. Even more challenging to interpret is the observation that

Ash2L expression increases with carbohydrate but not protein in males, even though male do not appear

to have a growth response to carbohydrate. Thus the relationship between *Ash2L* expression and body

size is not a simple one. The expression pattern of *CG3071* is similarly unclear. While *CG3071* also has a

carbohydrate and protein response in females, this response is qualitatively similar to the response of

Ash2L, which is not expected if TOR-signaling positively regulates *CG3071* expression but negatively

regulates *Ash2L* expression. Thus the role that TOR signaling plays in regulating SSD and SSP is

equivocal based on our data. Problematically, the mechanism by which TOR-signaling regulates *Ash2L*

and *CG3071* expression has not been fully explored, which makes interpreting changes in *Ash2L* and

CG3071 expression with diet even more challenging.

Our data suggests a complex relationship between sex, diet, *dILP* expression and dILP retention levels in

the IPCs. In females, there were stark differences in the pattern of expression among the dILPS. Of all the

dILPs, *dILP5* expression was most strongly correlated with body size, increasing with increasing protein

level but decreasing with increasing carbohydrate level. We also found that dILP5 levels in the IPCs were

lower in larvae fed on lower food concentrations. Thus, chronic reductions in nutrition reduce *dILP5*

expression and dILP5 IPCs level. This is consistent with dILP5 being a positive regulator of growth in

response to nutrition [35]. The negative effect of low nutrition on *dILP5* expression has been observed in

previous studies [30, 35, 36]. However, at least one of these studies reported an increase in dILP5

retention in the IPCs of larvae that were either starved for 24 hours or reared on a low protein diet since

birth [36], which is inconsistent with our data. The low protein diet in this study was generated by

reducing the amount of protein from ~16g/l to ~0.8g/l, while keeping the concentration of carbohydrate at

~60g/l, thereby decreasing the protein-to-carbohydrate ratio. In our study we reduced overall food

concentration from 360g/l to 45g/l while retaining a 1:2 protein-to-carbohydrate ratio. It is possible,

therefore, that the retention of dILP5 in the IPCs in the earlier study is due to a low protein-to-

carbohydrate ratio. Indeed, a second earlier study showed that a chronic high sugar diet with low

protein-to-carbohydrate ratio also increased dILP5 levels in the IPCs [37], although this time

accompanied by an increase in *dILP5* expression. Thus the relationship between *dILP5* expression and

dILP5 levels in the IPCs may depend not only on the overall food concentration, but also on food

composition, in particular the protein-to-carbohydrate level.

Previous studies suggest that *dILP2* expression is unaffected by reduced nutrition [30, 36] (although see

supplementary data in [38]), but that dILP2 peptides are retained in the IPCs in larvae that have been

starved 24h or are reared on a low protein diet [36, 39, 40]. A high sugar diet is also associated with

increased retention of dILP2 peptide in the IPCs, but with an increase in *dILP2* expression [37]. In our

study, however, we found that *dILP2* expression declined with an increase in both protein and

carbohydrate, and that dILP2 peptide was lower in the IPCs of larvae fed on lower food concentrations.

Our observed expression of *dILP3* was also inconsistent with previous studies that found that acute

starvation reduced expression [30], while rearing on a high protein diet increased expression [41]. In

contrast, we found that, as for *dILP2*, *dILP3* expression declined with increasing protein, although was

unaffected by carbohydrate level, which is surprising given that circulating sugars promote the release of

dILP3 from the IPCs [40]. Our finding that *dILP2* and *dILP3* expression is lowest on low protein diets that

restrict growth, and that dILP2 peptides levels are higher when *dILP2* expression is lower, suggests that

dILP2 and dILP3 may negatively regulate their own expression, which is a common characteristic of

hormone regulation.

Finally, the expression of *dILP8* showed a similar pattern to *dILP3,* increasing with a decrease in protein

concentration. dILP8 is involved in regulating growth and developmental timing through its inhibitory

effects on ecdysone synthesis [42, 43]. It is released from imaginal discs in response to damage or growth

perturbation, as well as showing periodic changes in expression throughout development . Ecdysone

synthesis is also inhibited by low nutrition early in the third larval instar, leading to a delay in

metamorphosis [44]. The observation that females on low protein diets also show elevated levels of *dILP8*

expression, suggests that dILP8 may play a role in regulating growth and/or development in response to

nutrition, at least in females.

We did not detect an effect of diet on the expression of any of the *dILP*s in male larvae. This did not,

however, translate into a significant interaction between diet and sex on *dILP* expression, and therefore

does not help explain the sex-specific differences in nutritional plasticity of body size. This was because

variation in *dILP 2,3,5* and *8* expression among samples within diets was significantly higher for male

samples than for females samples (Supplementary Table 1), reducing the statistical power to detect diet-

by-sex interactions. The only exception was for *dILP8*, such that *dILP8* expression was significantly more

plastic in females than in males. Nevertheless, *dILP8* expression in females did not correlate with

carbohydrate level, and so is unlikely to explain the differential effect of carbohydrate on body size in

females versus males. The elevated variation in *dILP* expression among male samples was not seen for

expression of *4eBP*, *InR*, *Ash2L*, or *CG3072*, and is therefore not likely due to sample degradation. It is also

not due to lower levels of expression in males relative to females. It is possible, therefore, that *dILP*

expression levels are more developmentally dynamic in males then females, and that we are capturing

aspects of that instability by measuring gene expression at only a single point in development.

Collectively, there appears to be no simple explanatory relationship between the sex-specific nutritional

geometry of body size with the sex-specific nutritional geometry of IIS/TOR-signaling gene expression.

Further, many of our findings do not replicate what has been reported in previous studies. An earlier

study looking at *dILP* expression in adults across a nutritional landscape saw similarly complex and non-

intuitive relationships between diet and gene expression [23]. What is very clear from both studies is that

the protein-to-carbohydrate ratio affects the transcriptional response of IIS/TOR-signaling genes to

changes in total diet, and our study indicates that it does so in a sex-specific way. This has important

implications for interpreting those studies that have looked at changes in the expression of IIS/TOR-

signaling genes in response to changes in nutrition. First, many studies simply dilute diet to reduce

overall nutrition. Problematically, while most *Drosophila* labs use 'standard' food recipes, the use of

different types of yeast, cornmeal, mollasses *etc* means that the composition of these diets may be unique

to most research groups [45]. Our data indicate that the effect of diet dilution on both body size and

IIS/TOR-signaling gene expression – and potentially the activity of the IIS/TOR-signaling pathways – will

depend on the P:C ratio of that specific diet. Second, these responses are sex-specific, and studies that do

not consider sex may come to different conclusions than studies that do. Third, the relationship between

dILP expression, dILP retention and diet is similarly complex, and may also depend on the composition

of the diet being manipulated.

One important caveat with our, and almost all other studies of IIS/TOR-signaling gene expression during

development, is that we measured expression at a single developmental time point, at the very beginning

of larval wandering. Previous studies have shown that the activity of the IIS and TOR-signaling pathways

change dynamically during development, and that different dILPs are expressed at different life stages.

This may also account for differences between our and other studies of the effects of nutrition on

IIS/TOR-signaling gene expression. Ideally, one would like to conduct a multidimensional study of gene

expression across both time and a nutritional landscape, and to tie this with a corresponding study of

dILP levels both in the IPCs and circulating in the hemolymph. Such a study would not only help

elucidate the relationship between IIS/TOR-signaling and the sex-specific effects of nutrition on body

size, but also help us understand better how nutrition regulates the transcription, translation, storage and

release of dILPs.

5.   Conclusions

Our study provides a foundation for future work that looks at the sex-specific effects of diet quantity and quality on body size and on the developmental mechanisms that regulate body size. Our data show significant differences in the effect of diet on gene expression in males and females, although they do not suggest a simple explanatory relationship between gene expression and body size. Future studies on sex-specific plasticity in *Drosophila*, and other animals, should therefore consider not only the nutritional geometry of gene expression, but also of protein levels, pathway activity and other developmental parameters such as growth rate and growth duration, ideally at multiple time points throughout development.

6.   Materials and Methods

The goal of the study was to test the hypothesis that sex-specific differences in body-size plasticity across a nutritional landscape in *Drosophila* correspond to sex-specific differences in the response to diet at the level of the IIS/TOR-signaling pathways, using the expression of genes that regulate or are regulated by IIS/TOR-signaling to assay pathway activity.

6.1.   Fly Stocks and Maintenance

Flies used for the nutritional geometry of body size were as described in [19]. Flies used for the nutritional geometry of gene expression were an isogenic line of white-eyed but otherwise wild-type flies (VDRC 60000). Flies were maintained as a stock at 17˚C on standard on standard media of 45 g of molasses, 75 g of sucrose, 70 g of cornmeal, 20 g of yeast extract, 10 g of agar, 1100 ml of water, and 25 ml of a 10% Nipagin solution per liter of fly food, under constant light.

6.2. *Diet Manipulation for Nutritional Geometry of Body Size*

Flies were reared on 24 diets: six protein-to-carbohydrate ratios (1:14.6, 1:7.2, 1:3.5, 1:1.7, 1.3:1, 1.4 :1), at four food concentrations: (45, 90, 180 and 360 g l^{-1}), with six replicate vials per diet, as described in [19]. We measured four non-genital traits in each fly: the length of the first femur, the area of the wing, the area of the maxillary palp and the length of the thorax (see [19] for details). Data were collected from, on average, 17 flies from each sex at each diet, with no more than 30 flies coming from any one vial. To assay the effect of diet on overall body size, we conducted a principle component analysis on all log-transformed non-genital morphological data from all flies, and used each fly's value for the first principle component (PC1) as a proxy for overall body size [20, 21].

6.3. *Diet Manipulation for Nutritional Geometry of Gene Expression*

Flies were reared on 28 different diets: seven protein-to-carbohydrate ratios (1:16, 1:8, 1:4, 1:2, 1:1, 1.5:1 and 2:1) at four food concentrations (45, 90, 180, and 360 g/l). To generate the diets we made an instant-yeast solution (Lesaffre SAF-Instant Red: 44% protein, 33% carbohydrate) and a sucrose solution (Mallinokrodt, Paris, Kentucky) at each food concentration. We also added 0.5% agar, 10% nipagen (10% p-hydroxy benzoic acid methyl ester in 95% ethanol) and 1% propionic acid (Sigma Life Science, St. Louis, Missouri) to each solution, to control consistency and prevent fungal growth. The solutions were autoclaved and mixed at different ratios to achieve the desired protein-to-carbohydrate ratios for each diet (supplemental Table 1). 10 ml of each of the 28 different diets were aliquoted into separate 25 x 95 mm vials.

Populations of 50 flies were placed in mating cages with 60mm petri dishes of standard lab food. The flies were allowed to lay eggs for 24 hrs. The adult flies were removed and the eggs were left on the petri dish for another 24hrs until the larvae had hatched. The first instar larvae were then transferred in groups of 50 into each food vial. There were at least five replicate vials for each diet.

All cultures were maintained at 25°C at constant light under 60-70% humidity. Larvae were collected

from the food vials when they had just initiated wandering, to ensure that all larvae were collected at the

same developmental stage regardless of diet. All larvae were sexed and flash-frozen to -80°C within five

minutes of being removed from the vial. The larvae were stored at -80°C until they were used for RNA

extractions.

_6.4._   **_Gene Expression Quantification_**

We used qPCR to assay gene expression in male and female larvae collected at the beginning of larval

wandering. Larvae were pooled into cohorts of 15 individuals, and we collected 3 cohorts of each sex for

each of the 28 diets, with the exception of the 1:16 360g/L diet, which only generated enough larvae for 2

cohorts of each sex. For each sample, RNA was extracted using Trizol (Invitrogen, Grand Island, NY,

USA) and treated with DNase 1 (Invitrogen) before being reverse-transcribed with High Capacity cDNA

Reverse Transcription Kit synthesis according to manufactures instructions (Thermo Fischer Scientific,

Carlsbad, CA, USA). We used SYBR Green PCR master mix (Thermo Fischer Scientific, Carlsbad, CA,

USA) to conduct qPCR using a CFX Connect Real-time System (Bio-Rad Laboratories, Inc., Hercules, CA).

mRNA abundance was calculated using the standard curve method [22]. Relative gene expression was

calculated by comparing the measured mRNA abundance of the gene of interest to that of ribosomal

protein 49 (RP49). Prior studies have indicated that RP49 is insensitive to changes in nutritional

conditions [23]. In two samples (1:16 180g and 1:4 45g male), gene expression levels were an order of

magnitude above all other samples across all amplicons. Data from these samples was therefore excluded

from subsequent analysis. The complete protocol is provided in the supplementary material, along with

primer sequences.

6.5. *dILP2 and dILP5 Levels in the Brain*

The following antibodies were used: dILP2 rat (1:800); dILP5 rabbit (1:800) (both gifts of P. Leopold); AlexaFluor 594 Donkey anti-Rat IgG (1:500); AlexaFluor 594 Goat anti-Rabbit IgG (1:500) (both from Life Technologies).

Larvae were reared on either 45g/l or 360g/l at a protein:carbohydrate ratio of 1:2, as described above. Brains were dissected in 1X PBS on ice from wandering female third instar larvae. The brains were fixed for 30 minutes in 4% paraformaldehyde and washed in 0.3% Triton-X in PBS (PBT) overnight at 4°C. They were then washed for 1 hr in 2% NGS in 0.2% I-Block in PBT (BBT/NGS) and incubated with primary antibody diluted in BBT/NGS overnight at 4°C. Subsequently, brains were washed four time in 0.2% BSA/I-Block in PBT (BBT), 15 minutes per wash. Brain were then incubated in secondary antibody at room temperature for 2 hours. Finally, brains were washed in four times in PBT for 15 minutes per wash and mounted with vector shield + DAPI.

We acquired images using Olympus Fluoview FV101 confocal microscope. For both 45g/L and 360g/L dILP2 images, 35% DAPI and 35% Alexa 594 intensity settings were used. For both 45g/L and 360g/L dILP5 images, 30% DAPI and 30% Alexa 594 intensity settings were used. We used a 1.5 micron step size to capture a Z-stack of images for each IPC (one per brain), which was flattened to a single image using a maximum-intensity projection in FIJI software. The intensity of the IPC in both the Alexa 594 and DAPI channel was then used to generate a normalized measure of the level of dILP2 or dILP5 in each brain. We repeated this for five brains for dILP2 and dILP5 at each diet level (nine for dILP5 at 45g/l). dILP levels were then compared between diets using a standard pooled *t*-test.

479   *6.6.   Statistical Analysis*

Nonlinear response surfaces are routinely modelled using the second-order polynomial regression [19,

23–25]. To determine the effect of protein and carbohydrate level on body size or gene expression we

therefore fit the model:

$$T_{ijk} = C_i + P_j + C_i^2 + P_j^2 + C_iP_j + \varepsilon_{ijk}$$

where T is trait (body size or expression level), C is carbohydrate concentration, P is protein concentration

and ε is error (subscripts are levels within variables). Models for body size included replicate vial as a

random factor and were fit using *lmer* in the *lme4* package in *R.* Models for expression level were fit using

lm in the *base* package of *R.* For each trait, the significance of each parameter was tested with a type III

ANOVA using the *Anova* function in the *car* package for standard models and the *summary* function in

the *lmerTest* package for mixed models. Any non-significant parameters were subsequently removed

from the model before the data were reanalyzed using the simplified model. Models that did not contain

interactions were analyzed with a type II ANOVA. We subsequently plotted the fitted values of the

simplest model against carbohydrate and protein levels. We also plotted thin-plate splines of the raw

measurements against carbohydrate and protein levels. Finally, we plotted sexual size dimorphism (SSD)

as the difference between the female and male fitted values against carbohydrate and protein level, as

well a thin-plate splines of the difference between mean female and male values at each diet.

To test whether the response to carbohydrates or protein differed between the sexes we fit two models:

$$\text{Model 1: } T_{ijk} = S_k + (C_i + P_j + C_i^2 + P_j^2 + C_iP_j) + \varepsilon_{ijk}$$

$$\text{Model 2: } T_{ijk} = S_k*(C_i + P_j + C_i^2 + P_j^2 + C_iP_j) + \varepsilon_{ijk}$$

where S is sex. Models were compared using ANOVA (for standard models of expression level) or

parametric boostrapping (for mixed models of body size). If inclusion of sex as an interactive factor rather

than an additive factor significantly improved the fit of the model, we concluded that the relationship

between trait and diet varied between males and females. If the relationship between trait and diet could

be explained by a simpler model, we tested whether inclusion of sex as an interactive rather than an

additive factor in the simplified model improved the fit of the model. We also conducted a similar

multivariate analysis on the effect of diet and sex on the expression of multiple genes, using MANOVA to

compare models. Finally, we used the fitted values from the relationship between diet and body size to

test the relationship between body size and the expression of each gene, using the linear model:

$$T_{ij...p} = A_i + B_j + ... + H_p + \varepsilon_{ij...p}$$

where T is body size (fitted values from the linear mixed model of the relationship between body size and

diet) and A through H is the expression of the genes measured in our analysis.

All analyses were conducted in R and the data and scripts for the analyses are provided on Dryad. Gene

expression levels were log-transformed prior to analysis. For all analyses, we plotted residual against

fitted values to confirm homogeneity of variance and generated a QQ plot to confirm that the residuals

were approximately normally distributed.

7.   Bibliography

1. Norman MD, Paul D, Finn J, Tregenza T. First encounter with a live male blanket octopus: The world's most sexually size-dimorphic large animal. New Zeal J Mar Fresh. 2010;36:733–6.

2. Bininda-Emonds ORP, Gittleman JL. Are pinnipeds functionally different from fissiped carnivores? The importance of phylogenetic comparative analyses. Evolution. 2000;54:1011–23.

3. Pearson D, Shine R, Williams A. Geographic variation in sexual size dimorphism within a single snake species (Morelia spilota, Pythonidae). Oecologia. 2002;131:418 426. doi:10.1007/s00442-002-0917-5.

4. Gebhardt-Henrich SG, Noordwijk AJ van. Nestling growth in the Great Tit I. Heritability estimates under different environmental conditions. J Evolution Biol. 1991;4:341 362.

5. Noordwijk AJV, Balen JHV, Scharloo W. Heritability of body size in a natural population of the Great Tit (Parus major) and its relation to age and environmental conditions during growth. Genet Res. 1988;51:149–62.

6. Stillwell RC, Davidowitz G. Sex differences in phenotypic plasticity of a mechanism that controls body size: implications for sexual size dimorphism. Proc Royal Soc B Biological Sci. 2010;277:3819 3826.

7. Pertseva MN, Shpakov AO. Conservatism of the Insulin Signaling System in Evolution of Invertebrate and Vertebrate Animals. J Evol Biochem Phys+. 2002;38:547 561. doi:10.1023/a:1022008932029.

8. Badyaev AV. Growing apart: An ontogenetic perspective on the evolution of sexual size dimorphism. Trends Ecol Evol. 2002;17:369 378.

9. Stillwell RC, Blanckenhorn WU, Teder T, Davidowitz G, Fox CW. Sex differences in phenotypic plasticity affect variation in sexual size dimorphism in insects: From physiology to evolution. Annu Rev Entomol. 2010;55:227–45.

10. Abouheif E, Fairbairn DJ. A comparative analysis of allometry for sexual size dimorphism: Assessing Rensch's rule. Am Nat. 1997;149.

11. Horne CR, Hirst AG, Atkinson D. A synthesis of major environmental-body size clines of the sexes within arthropod species. Oecologia. 2019;190:343 353. doi:10.1007/s00442-019-04428-7.

12. Fairbairn D. Allometry for sexual size dimorphism: Testing two hypotheses for Rensch's rule in the water strider Aquarius remigis. The American Naturalist. 2005;166:S69 S84.

13. Rowe L, Houle D. The lek paradox and the capture of genetic variance by condition dependent traits. Proc Royal Soc Lond Ser B Biological Sci. 1996;263:1415–21. doi:10.1098/rspb.1996.0207.

14. Bonduriansky R. The evolution of condition-dependent sexual dimorphism. Am Nat. 2007;169:9 19.

15. Emlen DJ, Warren IA, Johns A, Dworkin I, Lavine LC. A mechanism of extreme growth and reliable signaling in sexually selected ornaments and weapons. Science. 2012;337:860 864.

16. Testa ND, Ghosh SM, Shingleton AW. Sex-specific weight loss mediates sexual size dimorphism in Drosophila melanogaster. Plos One. 2013;8:e58936.

17. Rideout EJ, Narsaiya MS, Grewal SS. The Sex Determination Gene transformer Regulates Male-Female
Differences in Drosophila Body Size. Plos Genet. 2015;11:e1005683. doi:10.1371/journal.pgen.1005683.

18. Sawala A, Gould AP. The sex of specific neurons controls female body growth in Drosophila. Plos Biol.
2017;15:e2002252. doi:10.1371/journal.pbio.2002252.

19. Shingleton AW, Masandika JR, Thorsen LS, Zhu Y, Mirth CK. The sex-specific effects of diet quality versus
quantity on morphology in Drosophila melanogaster. Roy Soc Open Sci. 2017;4:170375. doi:10.1098/rsos.170375.

20. Bitner-Mathe BC, Peixoto AA, Klaczko LB. Morphological variation in a natural population of Drosophila
mediopunctata: altitudinal cline, temporal changes and influence of chromosome inversions. Heredity. 1995;75:54–
61.

21. Somers K. Multivariate allometry and removal of size with principal components analysis. Systematic Zoology.
1986;35:359–68. http://links.jstor.org/sici?sici=0039-
7989(198609)35%3A3%3C359%3AMAAROS%3E2.0.CO%3B2-4.

22. Larionov A, Krause A, Miller W. A standard curve based method for relative real time PCR. Bmc
Bioinformatics. 2005;6:62.

23. Post S, Tatar M. Nutritional Geometric Profiles of Insulin/IGF Expression in Drosophila melanogaster. Plos
One. 2016;11:e0155628. doi:10.1371/journal.pone.0155628.

24. Lee KP, Simpson SJ, Clissold FJ, Brooks R, Ballard JWO, Taylor PW, et al. Lifespan and reproduction in
Drosophila: New insights from nutritional geometry. Proc National Acad Sci. 2008;105:2498 2503.

25. Rodrigues MA, Martins NE, Balancé LF, Broom LN, Dias AJS, Fernandes ASD, et al. Drosophila melanogaster
larvae make nutritional choices that minimize developmental time. J Insect Physiol. 2015;81 J. Exp. Biol. 216
2013:69 80.

26. Puig O, Marr MT, Ruhf ML, Tjian R. Control of cell number by Drosophila FOXO: downstream and feedback
regulation of the insulin receptor pathway. Gene Dev. 2003;17:2006 2020. doi:10.1101/gad.1098703.

27. Semaniuk U, Piskovatska V, Strilbytska O, Strutynska T, Burdyliuk N, Vaiserman A, et al. Drosophila insulin-
like peptides: from expression to functions – a review. Entomol Exp Appl. 2020.

28. Guertin DA, Guntur KVP, Bell GW, Thoreen CC, Sabatini DM. Functional genomics identifies TOR-regulated
genes that control growth and division. Curr Biol. 2006;16:958 970.

29. Brogiolo W, Stocker H, Ikeya T, Rintelen F, Fernandez R, Hafen E. An evolutionarily conserved function of the
Drosophila insulin receptor and insulin-like peptides in growth control. Curr Biol. 2001;11:213 221.
doi:10.1016/s0960-9822(01)00068-9.

30. Ikeya T, Galic M, Belawat P, Nairz K, Hafen E. Nutrient-dependent expression of insulin-like peptides from
neuroendocrine cells in the CNS contributes to growth regulation in Drosophila. Curr Biol. 2002;12:1293 1300.

31. Zhang H, Stallock JP, Ng JC, Reinhard C, Neufeld TP. Regulation of cellular growth by the Drosophila target of
rapamycin dTOR. Gene Dev. 2000;14:2712–24.

32. Oldham S, Hafen E. Insulin/IGF and target of rapamycin signaling: a TOR de force in growth control. Trends
Cell Biol. 2003;13:79 85. doi:10.1016/s0962-8924(02)00042-9.

33. Hennig KM, Colombani J, Neufeld TP. TOR coordinates bulk and targeted endocytosis in the Drosophila
melanogaster fat body to regulate cell growth. J Cell Biology. 2006;173:963 974. doi:10.1083/jcb.200511140.

34. Puig O, Tjian R. Transcriptional feedback control of insulin receptor by dFOXO/FOXO1. Gene Dev.
2005;19:2435 2446. doi:10.1101/gad.1340505.

35. Okamoto N, Nishimura T. Signaling from Glia and Cholinergic Neurons Controls Nutrient-Dependent
Production of an Insulin-like Peptide for Drosophila Body Growth. Dev Cell. 2015;35:295 310.

36. Geminard C, Rulifson EJ, Léopold P. Remote control of insulin secretion by fat cells in Drosophila. Cell Metab.
2009;10:199 207.

37. Pasco MY, Léopold P. High Sugar-Induced Insulin Resistance in Drosophila Relies on the Lipocalin Neural
Lazarillo. Plos One. 2012;7:e36583. doi:10.1371/journal.pone.0036583.

38. Buhler K, Clements J, Winant M, Bolckmans L, Vulsteke V, Callaerts P. Growth control through regulation of
insulin-signaling by nutrition-activated steroid hormone in Drosophila. Development. 2018;145:dev.165654.
doi:10.1242/dev.165654.

39. Sano H, Nakamura A, Texada MJ, Truman JW, Ishimoto H, Kamikouchi A, et al. The Nutrient-Responsive
Hormone CCHamide-2 Controls Growth by Regulating Insulin-like Peptides in the Brain of Drosophila
melanogaster. Plos Genet. 2015;11:e1005209.

40. Kim J, Neufeld TP. Dietary sugar promotes systemic TOR activation in Drosophila through AKH-dependent
selective secretion of Dilp3. Nat Commun. 2015;6:6846. doi:10.1038/ncomms7846.

41. Millington JW, Chao C, Sun Z, Basner-Collins PJ, Brownrigg GP, Wat LW, et al. Female-specific upregulation
of insulin pathway activity mediates the sex difference in Drosophila body size plasticity. Biorxiv.
2020;:2020.04.22.054239.

42. Garelli A, Gontijo AM, Miguela V, Caparros E, Dominguez M. Imaginal Discs Secrete Insulin-Like Peptide 8 to
Mediate Plasticity of Growth and Maturation. Science. 2012;336:579 585.

43. Colombani J, Andersen DS, Léopold P. Secreted peptide Dilp8 coordinates Drosophila tissue growth with
developmental timing. Science. 2012;336:582 585.

44. Lee GJ, Han G, Yun HM, Lim JJ, Noh S, Lee J, et al. Steroid signaling mediates nutritional regulation of
juvenile body growth via IGF-binding protein in Drosophila. Proc National Acad Sci. 2018;115:201718834.

45. Lesperance DNA, Broderick NA. Meta-analysis of Diets Used in Drosophila Microbiome Research and
Introduction of the Drosophila Dietary Composition Calculator (DDCC). G3 Genes Genomes Genetics.
2020;10:g3.401235.2020.

8. Supplementary Figures

Supplementary Figure 1: Thin plate spline of the effect of protein and carbohydrate concentration on female and male body size and sexual size dimorphism (SSD). (A, B) Surfaces shows the relationship between body size, carbohydrate level and protein level in female and male flies (C) Surface shows a thin plate spline of the difference in female and male body size (SSD) across the same nutritional landscape, using fitted values from A and B. Points indicate diets tested and dotted lines connect diets with equal protein-to-carbohydrate ratios (1:14.6, 1:7.2, 1:3.5, 1:1.7, 1.3:1, 1.4 :1).

Supplementary Figure 2: Thin plate spline of the effect of protein and carbohydrate concentration on

the expression of IIS and TOR transcriptionally-regulated genes in females and males. Surfaces shows

the relationship between gene expression, carbohydrate level and protein level in female and male flies.

Expression of (A, A′) *InR*, and (B, B′) *4E-BP*, both negatively regulated by the activity of the IIS via the

Forkhead transcription factor FOXO. (C, C′) Expression of *Ash2L*, ostensibly negatively regulated by the

activity of TOR signaling. (D, D′) Expression of *CG3071*, ostensibly positively regulated by the activity of

TOR signaling. Points indicate diets tested and dotted lines connect diets with equal protein-to-

carbohydrate ratios (1:16, 1:8, 1:4, 1:2, 1:1, 2:1).

Supplementary Figure 3: Thin plate spline of the effect of protein and carbohydrate concentration on

the expression of dILPs in females and males. Surfaces shows the relationship between gene expression,

carbohydrate level and protein level in female and male flies. (A) *dILP2*. (B) *dILP3.* (C) *dILP 5.* (D) *dILP8*.

Points indicate diets tested and dotted lines connect diets with equal protein-to-carbohydrate ratios (1:16,

1:8, 1:4, 1:2, 1:1, 2:1). Corresponding thin-plate spline plots are shown in Supplementary Figure 1.

9.   Tables

Table 1: Effects of protein (P), carbohydrate (C), and their squares and product on body size in females and males.

SEX[A]		C	P	C²	P²	C*P
FEMALE	β[B]	-0.0044	0.0068	0.00001	-0.00003	0.00001
	X²	17.04***	0.02[NS]	21.45***	36.25***	4.80*
MALE	β[B]		0.0081		-0.00004	
	X²		11.408***		11.513**	

[A] For each sex, we fitted a linear mixed effect orthogonal polynomial regression with replicate vial as a random factor and removed non-significant parameters.

[B] Parameter values for the response surface are for the non-orthogonal polynomial regression to facilitate interpretation

[C] Type III Wald chi-square test with associated P-values: NS P > 0.05, * P < 0.05, ** P < 0.01, *** P < 0.001.

Table 2: Effect of including sex as an interactive versus additive term when modeling the influence of diet on body size

MODEL[A]	DF[B]	AIC[C]	BIC[C]	LL	MODEL COMPARSONS	LRT [D]	P [E]
A: $T_{ijkl} = S_k + C_i + C_{i^2} + P_j + P_{j^2} + C_iP_j + R_l$	9	-1059.0	-1017.3	538.48			
B: $T_{ijkl} = C_i + C_{i^2} + S_k (P_j + P_{j^2}) + C_iP_j + R_l$	11	-1057.7	-1006.8	539.85	A. v. B	2.73	0.256
C: $T_{ijkl} = S_k (C_i + C_{i^2}) + P_j + P_{j^2} + C_iP_j + R_l$	11	-1066.7	-1015.8	544.34	A. v. C.	11.71	**0.006**
D: $T_{ijkl} = S_k (C_i + C_{i^2} + P_j + P_{j^2} + C_iP_j) + R_l$	14	-1067.7	-1002.9	547.85	C. v. D.	7.03	0.071

[A] *T* is the body size, *S* is sex, *C* is carbohydrate, *P* is protein, *R* is replicate vial (random factor). The models differ by having sex as an interactive versus an additive factor.
[B] Estimated degrees of freedom for each model.
[C] AIC, BIC, log likelihood (LL) calculated using ML fit.
[D] Likelihood ratio test (LRT) statistic is for model comparisons.
[E] *P*-value for LRT is calculated by parametric bootstrapping using ML fit. Significant *P*-values are shown in bold.

662 **Table 3: Coefficients and partial-eta-squared for multiple regression of body size against gene**
663 **expression in males and females.**

GENE	FEMALE			MALE		
	Coefficient [A]	η_p^2	P [B]	Coefficient	η_p^2	P
InR	-0.014	0.004	0.638	-0.038	0.168	**0.047**
4E-BP	-0.123	0.147	**0.003**	0.018	0.013	0.593
Ash2L	0.014	0.003	0.660	0.019	0.034	0.391
CG3071	-0.053	0.038	0.137	-0.020	0.017	0.545
dILP2	0.039	0.018	0.308	-0.019	0.072	0.206
dILP3	-0.010	0.001	0.866	-0.018	0.080	0.181
dILP5	0.054	0.071	**0.042**	0.047	0.146	0.066
dILP8	-0.025	0.076	0.035	<0.001	<0.001	0.970

665 [A] Coefficients for the multiple regression $T_{ij...p} = A_i + B_j + … + H_p$ where T is body size (fitted values from the linear
666 mixed model of the relationship between body size and diet) and A through H is the expression of the genes
667 measured in our analysis.
668 [B] Significant P-values are shown in bold.

674 **Table 4: Effect of including sex as an interactive versus additive term when modeling the influence of**
675 **diet on the expression of multiple genes**

MODEL[A]	DF [B]	VAR [C]	*PILLAI*[D]	*F* [E]	*P* [F]
$T_{ijk} = S_k + (C_i + P_j + P_j^2)$	92	0.53979			
$T_{ijk} = S_k\,(C_i + P_j + P_j^2)$	89	0.51856	0.5284	2.1254	**0.002**
$T_{ijk} = S_k\,(C_i + C_i^2 + P_j + P_j^2 + C_iP_j)$	85	0.51583	0.3791	1.0599	0.384

[A] *T* is a matrix of expression levels across eight genes, *S* is sex, *C* is carbohydrate, *P* is protein, *R* is replicate vial
(random factor). The models differ by having sex as an interactive versus an additive factor.
[B] Estimated degrees of freedom for each model.
[C] Generalized variance
[E] Pillai's Trace is against model specified in row above.
[D] Aproximate *F*-statistic is against model specified in row above.
[E] *P*-value for LRT is calculated by parametric bootstrapping using ML fit. Significant *P*-values are shown in bold.

Table 5: Effects of protein (*P*), carbohydrate (*C*), and their squares and product on gene expression in

females and males.

GENE	SEX		C	C²	P	P²	C*P
InR	female	β^A			-0.025	0.0001	
		t-value^B			**-3.883*** **	**2.904** **	
	male	β			-0.033	0.0002	
		t-value			**-2.466* **	**2.347* **	
4E-BP	female	β			-0.005		
		t-value			-3.614***		
	male	β					
		t-value					
Ash2L	female	β	-1.64258		-0.018	0.0001	
		t-value	**-2.180* **		**-2.160* **	**2.076* **	
	male	β	1.7724				
		t-value	**2.651** **				
CG3071	female	β	-0.014	0.00002	-0.049	0.0002	0.00013
		t-value	**-4.252*** **	**2.083* **	**-5.137 *** **	**4.313*** **	**3.175** **
	male	β					
		t-value					
dILP2	female	β	-0.016	0.00004	-5.255		0.00012
		t-value	**-2.692** **	**3.013** **	**-3.095** **		**2.524* **
	male	β					
		t-value					
dILP3	female	β			-0.005		
		t-value			**-2.950** **		
	male	β					
		t-value					
dILP5	female	β	-1.5897		0.019	-0.0001	
		t-value	-1.573		0.156	**-2.087* **	
	male	β					
		t-value					
dILP8	female	β			-0.054	0.0003	
		t-value			**-5.101*** **	**2.917** **	
	male	β					
		t-value					

690   [A] Parameter values for the response surface are for the non-orthogonal polynomial regression to facilitate
691   interpretation.
692   [B] t-tests are for orthogonal polynomials. [NS] $P > 0.05$, * $P < 0.05$, * $P < 0.01$, *** $P < 0.001$. Significant t-values are shown
693   in bold.

Table 6: Effect of including sex as an interactive versus additive term when modeling the influence of diet on the expression of individual genes.

GENE	MODEL[A]	DF[B]	RSS[C]	F[D]	P[E]
InR	$T_{jk} = P_j + P_{j^2}$	148	100.209		
	$T_{jk} = S_k + P_j + P_{j^2}$	147	96.398	5.7500	**0.0178**
	$T_{jk} = S_k (P_j + P_{j^2})$	145	96.120	0.2098	0.811
4E-BP	$T_{jk} = S_k + P_j$	153	52.219		
	$T_{jk} = S_k P_j$	152	48.604	11.305	**0.001**
Ash2l	$T_{ik} = S_k + C_i$	152	80.009		
	$T_{ik} = S_k * C_i$	151	72.610	15.888	**0.0001**
	$T_{ik} = S_k (C_i + P_j + P_{j^2})$	147	68.459	2.229	0.069
CG3071	$T_{ijk} = S_k + C_i + C_{i^2} + P_j + P_{j^2}$	123	52.497		
	$T_{ijk} = S_k (C_i + C_{i^2} + P_j + P_{j^2})$	118	43.322	4.998	**0.0003**
dILP2	$T_{ijk} = C_i + C_{i^2} + P_j$	123	157.57		
	$T_{ijk} = S_k + C_i + C_{i^2} + P_j$	122	156.47	0.8623	0.3550
	$T_{ijk} = S_k (C_i + C_{i^2} + P_j + C_iP_j)$	119	152.96	0.9097	0.4386
dILP3	$T_{jk} = P_j$	152	136.38		
	$T_{jk} = S_k + P_j$	151	134.35	2.2632	0.1346
	$T_{jk} = S_k P_j$	150	134.35	<0.0001	0.9803
dILP5	$T_{ijk} = C_i + P_j + P_{j^2}$	148	147.40		
	$T_{ijk} = S_k + C_i + P_j + P_{j^2}$	147	142.85	4.6873	**0.0320**
	$T_{ijk} = S_k (C_i + P_j + P_{j^2})$	144	139.80	1.0458	0.37431
dILP8	$T_{jk} = S_k + P_j + P_{j^2}$	121	205.98		
	$T_{jk} = S_k (P_j + P_{j^2})$	119	190.40	4.8693	**0.0093**

[A] *T* is a expression level of gene, *S* is sex, *C* is carbohydrate, *P* is protein. The models differ by having sex as an interactive versus an additive factor. The models chosen are the simplest where one or either sex have significant parameter values.

[B] Degrees of freedom for each model.

[C] Residual Sum Square

[D] *F*-statistic is against model specified in row above.

[E] Significant P-values are shown in bold.

Supplementary Table 1: Variation in gene expression among samples within diets for male and female samples.

	4E-BP	*INR*	*ASH2L*	*CG3071*	*DILP2*	*DILP3*	*DILP5*	*DILP8*
FEMALE RMSE[A]	0.4184	0.5834	0.5233	0.5070	0.0505	0.4012	0.7208	1.3167
MALE RMSE[A]	0.5041	0.6971	0.6720	0.4876	1.4900	1.2494	0.9775	1.0351
P-**VALUE[B]**	0.2752	0.3739	0.1016	0.1628	**<0.0001**	**<0.0001**	**0.0389**	**0.0018**

[A] Root mean square error from the model $T = D$ where T is expression level of gene and D is diet (protein and carbohydrate combination).

[B] *P*-value for *F* test comparing residual variance for males and female level of gene expression, after fitting the model $T = D$. Significant P-values are shown in bold