Probing the aggregated effects of purifying selection per individual on 1,250 medical

phenotypes in the UK biobank

Ha My T. Vy,[1,¶] Daniel M. Jordan,[1,2,¶], Daniel J. Balick[3,4,¶], Ron Do[1,2*]

[1] The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine

at Mount Sinai, New York, NY, USA

[2] Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount

Sinai, New York, NY, USA

[3] Division of Genetics, Brigham and Women's Hospital, Harvard Medical School,

Boston, MA, USA

[4] Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA

* Corresponding author

Ron Do, PhD

Annenberg Building, Floor 18 Room 80A

1468 Madison Ave

New York, NY-10029

Phone Number: 212-241-6206

Fax Number: 212-849-2643

Email: ron.do@mssm.edu

[¶] These authors contributed equally to this work

Abstract

Understanding the relationship between natural selection and phenotypic

variation has been a long-standing challenge in human population genetics. With the

emergence of biobank-scale datasets, along with new statistical metrics to approximate

strength of purifying selection at the variant level, it is now possible to correlate a proxy

of individual relative fitness with a range of medical phenotypes. We calculated a per-

individual deleterious load score by summing the total number of derived alleles per

individual after incorporating a weight that approximates strength of purifying selection.

We assessed four methods for the weight, including GERP, phyloP, CADD, and fitcons.

By quantitatively tracking each of these scores with the site frequency spectrum, we

identified phyloP as the most appropriate weight. The phyloP-weighted load score was

then calculated across 15,129,142 variants in 335,161 individuals from the UK Biobank

and tested for association on 1,250 medical phenotypes. After accounting for multiple

test correction, we observed a strong association of the load score amongst coding

sites only on 28 traits including body mass, adiposity and metabolic rate. We further

observed that the association signals were driven by common variants (derived allele

frequency > 5%) with high phyloP score (phyloP > 2). Finally, through permutation

analyses, we showed that the deleterious load score had an excess of nominally

significant associations on many medical phenotypes. These results suggest a broad

impact of deleterious load on medical phenotypes and highlight the deleterious load

score as a new tool to disentangle the complex relationship between natural selection

and medical phenotypes.

Author summary

This study aims to augment our understanding between the complex relation between natural selection and human phenotypic variation. We developed a load score to approximate the relative fitness of an individual and correlate it with a set of medical phenotypes. Association tests between the load score amongst coding sites and 1,250 phenotypes in a sample of 335,161 individuals from the UK Biobank showed a strong association with 28 traits including body mass, adiposity and metabolic rate. Furthermore, an excess of nominal associations at suggestive levels was observed between the load score and medical phenotypes than would be expected under a null model. These results suggest that the aggregate effect of deleterious mutations as measured by the load score has a broad effect on human phenotypes.

Introduction

One of the primary questions of interest in the study of human population genetics is the relation between natural selection and the evolution of human phenotypes, from quantitative traits to complex disease. With the emergence of biobank-scale datasets, along with new statistical metrics to approximate strength of purifying selection at the variant level, it is now possible to both estimate the net impact of deleterious mutations for each individual in a large population sample and correlate it to a range of medical phenotypes exhibited by that individual. This provides an opportunity to simultaneously study the genetics of individuals within a relatively homogenous population and the potential impact of natural selection on annotated phenotypes.

A large body of literature exists on evolution and estimation of the deleterious

mutation load from human population samples, with particular emphasis on cross-

ancestry comparisons [1-6]. Rather than a comparison between human populations, we

aimed to assess the distribution of deleterious loads—the sum of all purifying selective

effects in each individual's genome—within a single human population. While the

mutation load generally represents a population-wide average of this quantity, we

estimated the same object for each individual in the population to produce a "load

score" that counts the net effect of deleterious variation in each individual's genome, a

count of derived alleles weighted by an estimate of the selective disadvantage for each

variant. When compared to the mean of the population, this per-individual load score

can be interpreted as a component of the relative fitness of each individual.

In this study, we aim to augment our understanding of the relation between

natural selection and human phenotypes by focusing on the net impact of purifying

selection on the fitness of each individual, and correlating this quantity to the set of

phenotypes acting on that individual. Previously this has been difficult for two reasons:

first, we do not have a direct measure of the fitness of individual humans that can be

estimated from genetic information, and second, there were no large databases

available to quantify the wide range of phenotypes possessed by each individual.

Biobank-scale datasets that contain both individual genotypes and phenotypes, such as

the UK Biobank [7, 8], finally provides access to both large-scale phenotypic

descriptions of each individual and some part of their genetic sequence.

We ventured to apply computational tools that predict aspects of purifying

selection for individual alleles to published genotypes of 335,161 white British

individuals from the UK Biobank to estimate the fitness impact of derived variation

present in each imputed genome in this sample. Most of the variation in the sample

exists at appreciable frequencies, and is likely under relatively small selective

disadvantage, but in aggregate the fitness impact can be substantial. Using this

representation of each individual's relative fitness, we probed correlations between the

impact of common deleterious variation in an individual's genome and their personal

phenotypic makeup. This provides a different lens into questions about the relation

between fitness vis-à-vis mutation load and human traits by looking at a per-individual

measure correlated to fitness, rather than focusing on the distribution of selective effects

in the population as a whole. This allows us to ask which phenotypes, if any, are highly

correlated to the aggregation of deleterious variation, and probe the relation between

the ensemble of phenotypes and fitness loss due to common variation in individuals.

Results

Comparison of four deleteriousness prediction scoring methods

The additive effects of deleterious variation can be quantified in aggregate by a

genome-wide score representing the net action of purifying selection on an individual

under the assumption that effects of individual variants can be summed additively.

Multiple methods have been developed to characterize purifying selection, including

methods that predict deleterious selection acting on the level of a single allele (fitCons

[9], FATHMM-MKL [10], deltaSVM [11], Funseq2 [12]), methods that measure

evolutionary conservation (phyloP [13], phastCons [14], GERP++ [15], SiPhy [16]) and

methods that predict the effect of an allele on molecular function (CADD [17], DANN

[18], GenoCanyon [19], Eigen and EigenPC [20]). Although these scores are formulated

as tests for strong selection or for molecular function, rather than as estimates of the

strength of selection, they are also correlated to the strength of selection, and are often

used as proxies for strength of selection [4, 21, 22]. In this study, we compared the

predicted deleteriousness of alleles for four widely used scoring methods that

approximate deleteriousness of a variant-- GERP++, phyloP, CADD, and fitCons-- with

their effects on allelic frequency in human population genetic data to select the most

appropriate measure for computation of the additive load.

Under negative or purifying selection, natural selection acts to reduce the

population frequency of deleterious mutations. This effect is more able to overcome

genetic drift as the strength of selection increases. As a result, we expect to observe a

higher number of rare alleles and a lower number of common alleles in regions of the

genome that are under negative selection, relative to putatively neutral regions. This

can be seen as a shift of the allele frequency spectrum (AFS) towards rare alleles, with

a steeper slope of the AFS indicating stronger purifying selection. We evaluated the

extent to which each scoring method captures the deleteriousness of an allele by

grouping alleles by the scores provided by each method and measuring the slope of the

resulting AFS. The more strongly a score is related to the strength of selection, the

more marked the increase in slope will be for high-scoring alleles relative to lower

scoring alleles.

Figure 1. Derived allele frequency spectra of different score categories for each deleteriousness

prediction scoring method.

For each scoring method, polymorphic sites are grouped into score intervals by the value of the score

annotated at the sites. Each solid line represents derived allele frequency spectrum of polymorphic sites

belonging to one score interval and three dashed lines represent derived allele frequency spectra of three

control categories: synonymous (syn), missense (mis), and loss of function (LOF) variants.

We evaluated this correlation using whole genome sequencing data from a non-

Finnish European population in the Genome Aggregation Database (gnomAD) [23]. For

each scoring method, we grouped alleles by score and compared the non-normalized

derived allele frequency spectra for each group (see Methods). Fig 1 plots the log of the

derived allele frequency (DAF), and shows a consistent pattern across all scores: the

higher the score, the steeper the slope of the log DAF (S1 Table). This indicates that,

for all scores shown, higher scores are associated with sites under stronger negative

selection, as expected. While all four scores show this pattern, CADD and phyloP show

clearer separations between DAF spectra than fitCons and GERP++. In the case of

fitCons, this underperformance is likely due to its incorporation of functional genomic

signatures that may increase its performance at identifying functional regions, but

detract from its performance at identifying sites under purifying selection. In the case of

GERP++, the underperformance may be due to the difference in multiple sequence

alignments used: the most recent available phyloP scores are calculated from an

alignment of 100 vertebrate sequences, while the most recent available GERP++

scores are calculated from an alignment of 36 mammalian sequences. For comparison,

we also calculated DAF spectra for synonymous sites, missense sites, and loss of

function sites (LOF). Based on this comparison, phyloP scores between 5 and 7.5

appear to be similarly deleterious to missense sites (nearly the same DAF slope) and

phyloP scores greater than 7.5 appear to be under similar purifying selection to LOF

sites. The equivalent numbers for CADD are 20 to 25 for missense and greater than 30

for LOF.

To further compare between CADD and phyloP, we examined the DAF

distribution for protein coding variants and noncoding variants separately. Both scores

performed similarly for coding variants, but phyloP showed better separation in

noncoding variants (S1 Fig). This is as expected, since CADD uses more features when

scoring coding variants than noncoding variants, while the phyloP method is identical for

coding and noncoding sites. For this reason, we concluded that phyloP has the most

consistent relationship between score and strength of negative selection, and selected

phyloP as our weight for our load score computation.

Per-individual load scores in UK Biobank

We calculated a per-individual deleterious load score by summing the total

number of derived alleles per individual, weighting each derived allele by its phyloP

score to account for the strength of purifying selection. We considered three load

scores: a genome-wide load score, a coding-specific load score, and a non-coding-

specific load score. Each score was computed for 335,161 unrelated, white-British

ancestry individuals in the UK Biobank using 6,774,062 variants from imputed

genotypes (95,850 coding and 6,678,212 non-coding) with positive phyloP scores

(positive scores denote uniform purifying selection, while negative scores denote clade-

specific selection). The observed population distribution across all sampled individuals

appear very close to normal for each of our three scores (Kolmogorov-Smirnov Tests P-

values = 0.32; 0.55; 0.20 for all variants, non-coding, and coding, respectively, Fig 2).

This is the expected result if the phyloP scores of derived alleles are identically

distributed across the entire population, due to the Central Limit Theorem. By contrast, if

the white-British population contained distinct subpopulations with dramatically different

distributions of phyloP scores among derived alleles, we would expect to see a sum of

multiple normal distributions with different means, resulting in a skewed or multi-modal

distribution.

Figure 2. Distribution of load score.

Histogram of three load scores computed from three sets of variants: coding variants (coding load score), non-coding variants (non-coding load score), and both coding and non-coding variants (genome-wide load score). Each load score was computed for 335,161 unrelated, white-British ancestry individuals.

Significant association between load score of coding variants and anthropometric and metabolic traits

To explore the overall effect of deleterious mutations on specific clinically measured phenotypes, we tested the association of each of the three load scores (genome-wide, coding and non-coding) with 681 traits and 569 clinical phenotypes, after adjusting for age, sex, genotyping chip, and assessment center. To account for potential confounders, we further included a set of geographical and socioeconomic variables available in the UK Biobank data as additional covariates (S2 Table). We note that many of these variables are significantly associated with the load score but the effects are small. Nonetheless, careful consideration was taken to add these as covariates in our association tests (S2 Table).

We discovered no phenotype significantly associated with either the genome-wide load score or non-coding load score (Bonferroni P value threshold = 1.3×10^{-5}).

212 However, 28 traits were significantly associated with the load score calculated from

213 coding SNPs; these included body mass, metabolic rate, and several adiposity traits

214 such as body mass index and waist circumference (Table 1). Some of these traits have

215 been found under directional selection in contemporary populations [24-26].

217
218
219 **Table 1. Association between coding load score and 28 traits.**
220

Trait	Sample size	beta	SE	P-value
Arm fat-free mass (right)	329238	-0.0073	0.0012	6.61×10^{-10}
Arm fat-free mass (left)	329182	-0.0074	0.0012	7.93×10^{-10}
Arm predicted mass (left)	329170	-0.0073	0.0012	1.12×10^{-9}
Leg fat mass (left)	329295	-0.0090	0.0015	1.39×10^{-9}
Basal metabolic rate	329326	-0.0074	0.0012	2.83×10^{-9}
Arm predicted mass (right)	329235	-0.0069	0.0012	3.76×10^{-9}
Weight	334221	-0.0098	0.0017	3.98×10^{-9}
Whole body water mass	329333	-0.0068	0.0012	8.68×10^{-9}
Leg fat mass (right)	329311	-0.0086	0.0015	1.02×10^{-8}
Whole body fat-free mass	329306	-0.0067	0.0012	1.24×10^{-8}
Whole body fat mass	328780	-0.0103	0.0018	1.80×10^{-8}
Leg fat percentage (left)	329297	-0.0064	0.0011	2.66×10^{-8}
Trunk fat-free mass	329057	-0.0065	0.0012	3.55×10^{-8}
Trunk predicted mass	329019	-0.0064	0.0012	5.07×10^{-8}
Trunk fat mass	329118	-0.0100	0.0019	1.21×10^{-7}
Leg predicted mass (right)	329303	-0.0063	0.0012	1.54×10^{-7}
Leg fat-free mass (right)	329303	-0.0063	0.0012	1.96×10^{-7}
Leg fat-free mass (left)	329280	-0.0062	0.0012	3.18×10^{-7}
Leg predicted mass (left)	329275	-0.0062	0.0012	3.77×10^{-7}
Leg fat percentage (right)	329316	-0.0058	0.0012	5.74×10^{-7}
Arm fat mass (right)	329242	-0.0092	0.0018	6.38×10^{-7}
Body mass index (BMI)	334097	-0.0092	0.0019	7.25×10^{-7}
Arm fat mass (left)	329188	-0.0090	0.0018	1.15×10^{-6}
Waist circumference	334612	-0.0080	0.0016	1.35×10^{-6}
Body fat percentage	329134	-0.0066	0.0014	2.98×10^{-6}
Hip circumference	334579	-0.0088	0.0019	3.29×10^{-6}
Impedance of arm (left)	329313	0.0061	0.0013	5.25×10^{-6}

Impedance of arm (right)	329297	0.0058	0.0013	1.25×10^{-5}

SE: standard error

Stratification by derived allele frequency showed that these association signals

are more pronounced when limiting to variants that are common (DAF > 5%) but not

close to fixation (DAF < 70%), while stratification by phyloP score shows that they are

more pronounced when limiting to variants with higher phyloP scores (phyloP>2, S3 and

S4 Tables). We therefore performed an additional stratification analysis by both DAF

and phyloP score (S5 Table). We observed that the signals are mostly driven by

common variants (5<=DAF<70%) with higher phyloP score (phyloP>2). This class of

variants notably contributes a large fraction (mean: 0.38 and sd: 0.005 per individual)

towards the per individual coding load score. This analysis necessarily excludes

extremely rare alleles, which are not well captured by the process of genotyping and

imputation. It is not clear how significant the aggregate contribution of these alleles to

the per individual load score would be.

To assess the effect of our weighting procedure, we calculated an unweighted

load score, the per-individual mutation burden, that simply counts derived alleles with no

reference to phyloP or other measures of selection. When using this score, all

significant association signals observed for the coding load score disappeared and no

significant association for genome-wide and non-coding unweighted score was detected

(S2 Fig). We further tested the associations with burden scores while restricting to only

rare variants (DAF < 5%) or only common variants (5% < DAF < 70%, S2 Fig), however

no significant association was observed. This is likely due to the domination of the

mutation burden by alleles under effectively no purifying selection, highlighting the need

for a weighting scheme to identify correlations to the relative per-individual fitness.

To assess whether the observed significant associations are sensitive to

reference bias, we included as a covariate the number of non-reference sites per

individual in our association testing for the top results in Table 2 (S6 Table). Association

results were very consistent, suggesting that reference bias is not likely a confounder.

Similarly, associations between the phenotypes and load score remain significant when

restricted to variants at which reference alleles are the same as predicted ancestral

alleles (S7 Table). We also re-computed load scores using phyloPNH scores, which are

phyloP scores calculated without human reference genome [4], and obtained similar but

slightly less significant results, with all the 28 phenotypes yielded p-value $< 6.13 \times 10^{-4}$

(S8 Table).

Associations with load score are enriched for nominal associations with disease

Phenome wide association test results showed that no single disease is significantly

associated with the load score (all $P > 0.05$ after accounting for multiple tests using

Bonferroni correction). However, rather than the load score having a strong effect on a

single disease, we hypothesized that the load score may have subtle effects on many

diseases, leading to an excess of weak associations that do not individually reach

statistical significance. To test this hypothesis, we compared the number of phenotypes

nominally associated with the load score (p-value < 0.05 without multiple test correction)

to a null distribution generated by random permutation of individual load score values

(Methods). For this analysis, we restricted to associations with clinical phenotypes

defined by ICD (International Statistical Classification of Diseases) codes. Out of 364

ICD-10 codes, 28, 25, and 26 ICD codes (S9 Table) were found to be nominally

associated with coding load score, non-coding load score, and genome-wide load score

respectively. The number of nominally significant associations for each load score is

significantly larger than the expected number under the null model, supporting this

hypothesis (Fig 3). This analysis was statistically significant for both the coding load

score (P=0.02) and the genome-wide load score (P=0.04) but not the non-coding load

score (P>0.05) (S3 Fig) for these weak phenotype associations. We repeated the

permutation analysis for the unweighted burden score as negative controls. As

expected, enrichment of week association between burden scores and diseases are not

statistically significant (S4 Fig).

Figure 3. Enrichment of clinical phenotypes nominally associated with coding load score. Null

distribution of the number of clinical phenotypes weakly associated with coding load score was obtained

from 2,000 permutations in total. For each permutation, coding load score was shuffled randomly among

335,161 samples and the number of association was the count of phenotypes which yielded a p-value <

0.05 in the association tests between permuted load score and 1,250 phenotypes. Red dashed line

indicates the observed number of clinical phenotypes nominally associated with coding load score (n =

28).

Discussion

In this study, we have described a polygenic load score that estimates the deleterious

load carried by an individual, and applied this score to 335,161 white British individuals

from the UK Biobank. Our analysis produced two major results: First, while we found no

significant associations between individual medical phenotypes and the genome-wide

load score, we found that more phenotypes are nominally associated with the load

score than would be expected under a null model (Figure 3). This suggests that the

deleterious load has a broad effect on the human phenome, rather than being

specifically associated with a small number of phenotypes. This is consistent with

Fisher's Geometric Model of fitness, which proposes that the fitness of a population is

determined by overall phenotypic distance from a theoretical optimal point in a

phenotype space that potentially encompasses the organism's entire phenome [27, 28].

Second, by restricting to protein coding variation, we found significant associations

between the coding-only deleterious load score and a variety of adiposity phenotypes,

along with other anthropometric phenotypes and phenotypes related to metabolic rate.

This suggests that adiposity may be under polygenic selection driven by a large number

of coding variants in humans. This is consistent with previous results obtained from the

UK Biobank using an unrelated methodology [24]. We found no similar associations with

the noncoding deleterious load score, which is in contrast to numerous studies finding

significant genetic associations in noncoding regions, including associations with the

same adiposity traits we found associated with our coding load score. Since our derived

allele frequency spectrum analysis (Fig. 1) suggests that sites with higher phyloP scores

are under purifying selection in noncoding regions as well as coding, the lack of

significance in non-coding regions cannot be interpreted as a lack of purifying selection

in these regions or poor sensitivity to selection in these regions. It may instead indicate

that selection acts on phenotype associations in noncoding regions in a different way

from how it acts in coding regions, possibly due to the small effect size of individual

noncoding variants.

There are several limitations to our method. We computed the load score from

imputed genotypes rather than sequenced whole genomes, which gives us little

information about extremely rare variants in the population, masking potentially large

contributions to the load from variants under the strongest selection. As a future topic of

research, the same methodology can be applied to include rare variants, which would

shed light on the relative contribution of common and rare variation to the phenotypic

associations of load. Previous studies have shown that rare variation contributes

substantially to differences in deleterious load between human populations, so we may

expect it to have a significant impact on individual load in this context as well [1, 2].

Furthermore, the phyloP score used to estimate the deleteriousness of alleles measures

only the likelihood that a site is evolving under constraint in vertebrates, and is not a

direct estimate of the selective effect of a variant in humans. It is possible that the use of

vertebrate-level conservation has reduced our ability to identify recent selection on

human phenotypes, particularly those that are human specific. However, the fact that

selection on adiposity traits was also detected by a method [24, 25] that does not rely

on phyloP suggests that this result is not spurious. This feature of the phyloP score also

makes it difficult to measure the effect of dominant or recessive selection, which may

contain additional important insights. Finally, we did not incorporate any measure of

positive selection in the computation of the load score. Scores similar to phyloP that

could be used to detect positive selection do exist, but they rely on measures of

nucleotide diversity and haplotype structure within larger regions of the genome, and

are difficult to apply to single nucleotides as would be required to incorporate them into

this analysis [29]. Methods to detect positive selection in the human lineage on finer

scales are an area of ongoing research, and such methods could be incorporated into

this approach in the future. All these methodological constraints limit the range of

variation we identify as contributing to an individual's burden score, and this limitation

may be biased with respect to trait associations. In particular, we might expect rare

variation or positive selection to reveal a different set of trait associations than the ones

we find here by investigating common variation under purifying selection. This could

potentially expand the scope of associated phenotypes beyond adiposity traits, a

possibility that is also supported by the presence of nominal associations with many

phenotypes unrelated to adiposity (S9 Table). Nevertheless, we do expect common

variation under purifying selection to underlie a large fraction of common disease

phenotypes, and therefore to provide valuable insights about the action of natural

selection in humans.

One potentially exciting application for this approach is applying it to different

populations to discover of population-specific insights into phenotypic associations with

deleterious load. Since PhyloP scores can be calculated without any reference to

specific human populations [4], there is no reason in principle that this method could not

be applied to biobank data from other populations, given a sufficient number of

samples. However, a few cautions are necessary. First, it is well known that

comparisons of genetic associations and polygenic risk are unreliable across different

ancestries [30], that signals of polygenic selection can easily be confounded by

population structure or admixture [31, 32], and that mutation load specifically differs

substantially between populations based on their demographic history [1, 2]. This

makes it difficult to compare load scores directly between individuals of different

ancestry, and also would likely make it difficult to apply this approach to admixed

populations or populations with heterogeneous ancestry. Second, the approach of

genotyping and imputation is entirely dependent on the availability of appropriate

genotyping arrays and imputation panels, neither of which is necessarily available for all

populations. It will be essential to use sequencing data for any population that is not well

represented in these resources. Finally, many traits are strongly influenced by social,

cultural, and environmental factors which may differ dramatically across populations,

resulting in differences between populations that are not necessarily related to natural

selection in a straightforward way. This is certainly true of the adiposity traits we identify

in this study. Results of such studies should therefore be interpreted with caution.

The deleterious load score presented here provides a new approach to

investigate the complex relationship between natural selection acting on individuals,

individual medical phenotypes, and the human phenome at large. We expect that as the

available biobank data continues to grow in size and scope, this method can be applied

to larger and more diverse populations to gain additional insights into how load varies

between different populations, possibly empowering population-specific medical

discoveries with deleterious load.

Material and Methods

The dependence of derived allele frequency on deleteriousness score

We evaluated the dependence of derived allele frequency of single nucleotide

polymorphisms (SNPs) discovered in the whole genome sequences of 7,509 non-

Finnish European individuals in the GnomAD data set [23] on each of the four candidate

annotations for the presence of purifying selection: GERP++ [15], phyloP [13], CADD

[17], and fitcons [9]. 88,060,485 SNPs with less than one percent of missing data were

considered. Functional effects and deleterious scores at each SNP were annotated

using Whole Genome Sequence Annotator (WGSA) v0.7 [33]. We used functional

effects annotated by Variant Effect Predictor (VEP) and determined derived and

ancestral allele status based on the six-way EPO (Enredo, Pecan, Ortheus) multiple

alignments of primate species.

For each deleteriousness score, we divided the SNPs into multiple groups with

arbitrarily defined intervals based on the range of each score. The intervals used were:

(0 ~ 0.2, 0.2 ~ 0.4, 0.4 ~ 0.6, 0.6 ~ 0.8) for fitcons, (-9.5 ~ -7.5, -7.5 ~ -5.5, -5.5 ~ -3.5, -

3.5 ~ -1.5, -1.5 ~ 0.5, 0.5 ~ 2.5, 2.5 ~ 4.5, 4.5 ~ 6.5) for GERP, (0 ~ 5, 5 ~ 10, 10 ~ 15,

15 ~ 20, 20 ~ 25, 25 ~ 30, 30 ~ 35) for CADD, and (-5 ~ -2.5, -2.5 ~ 0, 0 ~ 2.5, 2.5 ~

5.0, 5.0 ~ 7.5, 7.5~ 10) for phyloP.

Load score calculation

The load score of each individual was calculated by adding up the number

(dosage in case of imputed SNPs) of derived alleles at each SNP, weighted by the

phyloP score at that site, across the entire genome. Derived alleles were determined

based on the six-way EPO alignment, as described above. Since we are focusing on

the effect of purifying selection, only SNPs with positive phyloP score (positive scores

denote uniform purifying selection, while negative scores denote clade-specific

selection) were included. In this paper, we computed three load scores using three

different SNP sets: the coding load score summed only over coding variants, the non-

coding load score summed only over non-coding variants, and the genome-wide load

score computed from both coding and non-coding variants. All load scores were

computed using PRSice-2 software [34] under an additive model. Coding and

noncoding variants were defined based on VEP annotation.

Genotypic and phenotypic data

The UK Biobank consists of genotype, phenotype, and demographic data of

more than 500,000 individuals recruited across the United Kingdom. Individual

genotypes were generated from either the Affymetrix Axiom UK Biobank array

(~450,000 individuals) or the UK BiLEVE array (~50,000 individuals), each contains

~0.9 million markers. Additional variants were then imputed using the Haplotype

Reference Consortium (HRC) combined with the UK10K haplotype resource, with a

total of ~96 million variants available in the latest released imputed data (version 3). To

compute per-individual load scores, we restricted to variants with imputation quality

INFO score >= 0.9. We excluded samples that were outliers in heterozygosity or

missing rates, samples with putative sex chromosome aneuploidy, and samples with

self-reported non-white British ancestry. We also excluded one individual from each pair

of samples with relatedness up to the third degree. This produced a subsample of

335,161 individuals. All information used to exclude samples is included in the UK

Biobank resource page.

UK Biobank provides a wide range of medical phenotypes from base line

assessment, biochemical assays, dietary questionnaire, and health records. In the

present study, we focused on 2,419 phenotypes which had been selected for heritability

estimation by the Neale group (http://www.nealelab.is/blog/2017/9/15/heritability-of-

2000-traits-and-disorders-in-the-uk-biobank). This subgroup covers phenotypes in most

of the core categories, including early life and reproductive factors, family history,

cognitive function, physical measures, lifestyle and health outcomes.

Association between load scores and 1,250 phenotypes in UKBB

Association between load scores and phenotypes was conducted using

PHESANT [35], a package designed to process phenotypes and run phenome scans in

UK Biobank. The PHESANT pipeline loads each input phenotype as continuous,

ordered categorical, unordered categorical, or binary based on the information in the UK

Biobank data dictionary; preprocesses and re-categorizes the phenotype data based on

predefined rules; and assigns the appropriate regression test for the data type (linear

regression / lm R function for continuous, ordered logistic regression / polr R function for

ordered categorical, multinomial logistic regression / multinom R function for unordered

categorical, and binomial regression / glm R function with family = binomial for binary).

We used PHESANT to process all 2,419 available traits after raising the number

of valid cases or controls for binary traits from 50 to 500 and the minimum number of

individuals with valid data from 500 to 5000. This resulted in 1,250 phenotypes (75

continuous, 104 ordered categorical, 36 unordered categorical, and 1,035 binary

phenotypes) remaining for our association analysis. Besides the commonly used

covariates of age, sex, genotype chip, assessment center and 40 principal components,

we added five variables as covariates in all association tests that might denote

population structure: birth location, home area population density, Townsend

deprivation index, and UK deprivation index.

Association of 28 adiposity traits with coding load scores stratified by phyloP

score and derived allele frequency

To explore which variants drive the association between the 28 adiposity traits

and the coding load score, we stratified variants by derived allele frequency (rare

variants, 0 to 0.05; intermediate frequency variants, 0.05 to 0.3; common variants, 0.3 to

0.7; and variants near fixation, 0.7 to 1) and phyloP score (0 to 2, 2 to 4, 4 to 6, 6 to 8,

and 8 to 10). Simultaneous stratification was performed with four groups of SNPs:

DAF<0.05 and phyloP$\leq$2, DAF<0.05 and phyloP>2, DAF$\geq$0.05 and phyloP$\leq$2, DAF$\geq$0.05

and phyloP>2.

Permutation of phenome-wide association analysis and creation of null distribution

A null distribution of the number of clinical phenotypes weakly associated with load score was created by repeatedly running the association test between load scores and phenotypes after randomly shuffling the load scores of individuals within the tested sample. The phenotypes included in this permutation analysis were all primary International Statistical Classification of Diseases (ICD-10) codes (field 41202), excluding three environmental-related categories: "Injury, poisoning and certain other consequences of external causes" S00-T88; "External causes of morbidity" V00-Y99; and "Factors influencing health status and contact with health services" Z00-Z99. The same set of covariates used in phenome-wide association study (PHEWAS) tests above was applied. For each permutation, the number of phenotypes nominally associated with the load score (p-value<0.05) was then computed. The permutation p-value was calculated as the fraction of permutations for which the number of nominally associated traits was at least as large as the observed number of nominally associated traits.

Acknowledgments

This research has been conducted using the UK Biobank Resource under Application Number '16218'.

Funding

RD is supported by R35GM124836 from the National Institute of General Medical

Sciences of the National Institutes of Health, and R01HL139865 from the National

Heart, Lung, and Blood Institute of the National Institutes of Health.

Conflict of interest

RD has received research support from AstraZeneca and Goldfinch Bio, being a

scientific co-founder and equity holder for Pensieve Health and being a consultant for

Variant Bio, all not related to this work.

References

493
1.   Henn BM, Botigue LR, Bustamante CD, Clark AG, Gravel S. Estimating the mutation load
in human genomes. Nat Rev Genet. 2015;16(6):333-43. Epub 2015/05/13. doi:
10.1038/nrg3931. PubMed PMID: 25963372; PubMed Central PMCID: PMCPMC4959039.
2.   Henn BM, Botigue LR, Peischl S, Dupanloup I, Lipatov M, Maples BK, et al. Distance from
sub-Saharan Africa predicts mutational load in diverse human genomes. Proc Natl Acad Sci U S
A. 2016;113(4):E440-9. Epub 2015/12/30. doi: 10.1073/pnas.1510805112. PubMed PMID:
26712023; PubMed Central PMCID: PMCPMC4743782.
3.   Lohmueller KE, Indap AR, Schmidt S, Boyko AR, Hernandez RD, Hubisz MJ, et al.
Proportionally more deleterious genetic variation in European than in African populations.
Nature. 2008;451(7181):994-7. Epub 2008/02/22. doi: 10.1038/nature06611. PubMed PMID:
18288194; PubMed Central PMCID: PMCPMC2923434.
4.   Fu W, Gittelman RM, Bamshad MJ, Akey JM. Characteristics of neutral and deleterious
protein-coding variation among individuals and populations. The American Journal of Human
Genetics. 2014;95(4):421-36.
5.   Do R, Balick D, Li H, Adzhubei I, Sunyaev S, Reich D. No evidence that selection has been
less effective at removing deleterious mutations in Europeans than in Africans. Nat Genet.
2015;47(2):126-31. Epub 2015/01/13. doi: 10.1038/ng.3186. PubMed PMID: 25581429;
PubMed Central PMCID: PMCPMC4310772.
6.   Simons YB, Turchin MC, Pritchard JK, Sella G. The deleterious mutation load is
insensitive to recent population history. Nat Genet. 2014;46(3):220-4. Epub 2014/02/11. doi:
10.1038/ng.2896. PubMed PMID: 24509481; PubMed Central PMCID: PMCPMC3953611.
7.   Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK biobank: an open
access resource for identifying the causes of a wide range of complex diseases of middle and
old age. PLoS Med. 2015;12(3):e1001779. Epub 2015/04/01. doi:

10.1371/journal.pmed.1001779. PubMed PMID: 25826379; PubMed Central PMCID:
PMCPMC4380465.
8.   Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK Biobank
resource with deep phenotyping and genomic data. Nature. 2018;562(7726):203-9. Epub
2018/10/12. doi: 10.1038/s41586-018-0579-z. PubMed PMID: 30305743; PubMed Central
PMCID: PMCPMC6786975.
9.   Gulko B, Hubisz MJ, Gronau I, Siepel A. A method for calculating probabilities of fitness
consequences for point mutations across the human genome. Nat Genet. 2015;47(3):276-83.
Epub 2015/01/20. doi: 10.1038/ng.3196. PubMed PMID: 25599402; PubMed Central PMCID:
PMCPMC4342276.
10.   Shihab HA, Rogers MF, Gough J, Mort M, Cooper DN, Day IN, et al. An integrative
approach to predicting the functional effects of non-coding and coding sequence variation.
Bioinformatics. 2015;31(10):1536-43. Epub 2015/01/15. doi: 10.1093/bioinformatics/btv009.
PubMed PMID: 25583119; PubMed Central PMCID: PMCPMC4426838.
11.   Lee D, Gorkin DU, Baker M, Strober BJ, Asoni AL, McCallion AS, et al. A method to
predict the impact of regulatory variants from DNA sequence. Nat Genet. 2015;47(8):955-61.
Epub 2015/06/16. doi: 10.1038/ng.3331. PubMed PMID: 26075791; PubMed Central PMCID:
PMCPMC4520745.
12.   Fu Y, Liu Z, Lou S, Bedford J, Mu XJ, Yip KY, et al. FunSeq2: a framework for prioritizing
noncoding regulatory variants in cancer. Genome Biol. 2014;15(10):480. Epub 2014/10/03. doi:
10.1186/s13059-014-0480-5. PubMed PMID: 25273974; PubMed Central PMCID:
PMCPMC4203974.
13.   Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution
rates on mammalian phylogenies. Genome Res. 2010;20(1):110-21. Epub 2009/10/28. doi:
10.1101/gr.097857.109. PubMed PMID: 19858363; PubMed Central PMCID: PMCPMC2798823.
14.   Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, et al.
Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome
Res. 2005;15(8):1034-50. Epub 2005/07/19. doi: 10.1101/gr.3715005. PubMed PMID:
16024819; PubMed Central PMCID: PMCPMC1182216.
15.   Davydov EV, Goode DL, Sirota M, Cooper GM, Sidow A, Batzoglou S. Identifying a high
fraction of the human genome to be under selective constraint using GERP++. PLOS
Computational Biology. 2010;6(12).
16.   Garber M, Guttman M, Clamp M, Zody MC, Friedman N, Xie X. Identifying novel
constrained elements by exploiting biased substitution patterns. Bioinformatics.
2009;25(12):i54-62. Epub 2009/05/30. doi: 10.1093/bioinformatics/btp190. PubMed PMID:
19478016; PubMed Central PMCID: PMCPMC2687944.
17.   Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework
for estimating the relative pathogenicity of human genetic variants. Nat Genet. 2014;46(3):310-
5. Epub 2014/02/04. doi: 10.1038/ng.2892. PubMed PMID: 24487276; PubMed Central PMCID:
PMCPMC3992975.
18.   Quang D, Chen Y, Xie X. DANN: a deep learning approach for annotating the
pathogenicity of genetic variants. Bioinformatics. 2015;31(5):761-3. Epub 2014/10/24. doi:
10.1093/bioinformatics/btu703. PubMed PMID: 25338716; PubMed Central PMCID:
PMCPMC4341060.

19.   Lu Q, Hu Y, Sun J, Cheng Y, Cheung K-H, Zhao H. A statistical framework to predict
functional non-coding regions in the human genome through integrated analysis of annotation
data. Sci Rep. 2015;5:10576.
20.   Ionita-Laza I, McCallum K, Xu B, Buxbaum JD. A spectral approach integrating functional
genomic annotations for coding and noncoding variants. Nat Genet. 2016;48(2):214-20. Epub
2016/01/05. doi: 10.1038/ng.3477. PubMed PMID: 26727659; PubMed Central PMCID:
PMCPMC4731313.
21.   Racimo F, Schraiber JG. Approximation to the distribution of fitness effects across
functional categories in human segregating polymorphisms. PLoS Genet.
2014;10(11):e1004697. Epub 2014/11/07. doi: 10.1371/journal.pgen.1004697. PubMed PMID:
25375159; PubMed Central PMCID: PMCPMC4222666.
22.   Huang YF, Siepel A. Estimation of allele-specific fitness effects across human protein-
coding sequences and implications for disease. Genome Res. 2019;29(8):1310-21. Epub
2019/06/30. doi: 10.1101/gr.245522.118. PubMed PMID: 31249063; PubMed Central PMCID:
PMCPMC6673719.
23.   Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The
mutational constraint spectrum quantified from variation in 141,456 humans. Nature.
2020;581(7809):434-43.
24.   Sanjak JS, Sidorenko J, Robinson MR, Thornton KR, Visscher PM. Evidence of directional
and stabilizing selection in contemporary humans. Proc Natl Acad Sci U S A. 2018;115(1):151-6.
Epub 2017/12/20. doi: 10.1073/pnas.1707227114. PubMed PMID: 29255044; PubMed Central
PMCID: PMCPMC5776788.
25.   Byars SG, Ewbank D, Govindaraju DR, Stearns SC. Colloquium papers: Natural selection
in a contemporary human population. Proc Natl Acad Sci U S A. 2010;107 Suppl 1(suppl
1):1787-92. Epub 2009/10/28. doi: 10.1073/pnas.0906199106. PubMed PMID: 19858476;
PubMed Central PMCID: PMCPMC2868295.
26.   Beauchamp JP. Genetic evidence for natural selection in humans in the contemporary
United States. Proc Natl Acad Sci U S A. 2016;113(28):7774-9. Epub 2016/07/13. doi:
10.1073/pnas.1600398113. PubMed PMID: 27402742; PubMed Central PMCID:
PMCPMC4948342.
27.   Sella G, Barton NH. Thinking about the evolution of complex traits in the era of genome-
wide association studies. Annu Rev Genom Hum Genet. 2019;20:461-93.
28.   Fisher RA. The genetical theory of natural selection. Oxford, UK: Clarendon Press; 1930.
29.   Booker TR, Jackson BC, Keightley PD. Detecting positive selection in the genome. BMC
Biol. 2017;15(1):98. Epub 2017/11/01. doi: 10.1186/s12915-017-0434-y. PubMed PMID:
29084517; PubMed Central PMCID: PMCPMC5662103.
30.   Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, et al. Human
Demographic History Impacts Genetic Risk Prediction across Diverse Populations. Am J Hum
Genet. 2017;100(4):635-49. Epub 2017/04/04. doi: 10.1016/j.ajhg.2017.03.004. PubMed PMID:
28366442; PubMed Central PMCID: PMCPMC5384097.
31.   Berg JJ, Harpak A, Sinnott-Armstrong N, Joergensen AM, Mostafavi H, Field Y, et al.
Reduced signal for polygenic adaptation of height in UK Biobank. Elife. 2019;8. Epub
2019/03/22. doi: 10.7554/eLife.39725. PubMed PMID: 30895923; PubMed Central PMCID:
PMCPMC6428572.

32.   Sohail M, Maier RM, Ganna A, Bloemendal A, Martin AR, Turchin MC, et al. Polygenic
adaptation on height is overestimated due to uncorrected stratification in genome-wide
association studies. Elife. 2019;8. Epub 2019/03/22. doi: 10.7554/eLife.39702. PubMed PMID:
30895926; PubMed Central PMCID: PMCPMC6428571.
33.   Liu X, White S, Peng B, Johnson AD, Brody JA, Li AH, et al. WGSA: an annotation pipeline
for human genome sequencing studies. J Med Genet. 2016;53(2):111-2. Epub 2015/09/24. doi:
10.1136/jmedgenet-2015-103423. PubMed PMID: 26395054; PubMed Central PMCID:
PMCPMC5124490.
34.   Choi SW, O'Reilly PF. PRSice-2: Polygenic Risk Score software for biobank-scale data.
Gigascience. 2019;8(7). Epub 2019/07/16. doi: 10.1093/gigascience/giz082. PubMed PMID:
31307061; PubMed Central PMCID: PMCPMC6629542.
35.   Millard LAC, Davies NM, Gaunt TR, Davey Smith G, Tilling K. Software Application Profile:
PHESANT: a tool for performing automated phenome scans in UK Biobank. Int J Epidemiol.
2018;47(1):29-35. Epub 2017/10/19. doi: 10.1093/ije/dyx204. PubMed PMID: 29040602;
PubMed Central PMCID: PMCPMC5837456.
621
622

Supporting information

S1 Fig. Derived allele frequency spectra of coding and non-coding variants for different CADD and phyloP score categories. Top row: Derived allele frequency spectrum of coding variants. Bottom row: Derived allele frequency spectrum of non-coding variants. Each solid line represents derived allele frequency spectrum of polymorphic sites belonging to one score category and three dashed lines represent derived allele frequency spectra of three control categories: synonymous (syn), missense (mis), and loss of function (LOF) variants.

S2 Fig. Phenotypic association of load (phyloP-weighted) and burden (unweighted) scores. Quantile-quantile plot of -log10 p-values for the phenotypic association of A) load scores (weighted by phyloP); B) burden scores (unweighted); C)

burden scores restricted to rare variants (DAF<5%); and D) burden scores restricted to

common variants (5%<=DAF<70%).

S3 Fig: Enrichment of clinical phenotypes nominally associated with genome-

wide load score and non-coding load score. Null distribution of the number of

clinical phenotypes weakly associated with genome-wide load score (left) and non-

coding load score (right) was obtained from 2,000 permutations each. For each

permutation, the load score was shuffled randomly among 335,161 samples and the

number of associations on the x-axis was the count of phenotypes which yielded p-

value < 0.05 in the association tests between the permuted load score and 1,250

phenotypes. The red dashed lines indicates the observed number of clinical phenotypes

nominally associated with genome-wide load score (n = 26, left) and non-coding load

score (n = 25, right).

S4 Fig: Enrichment of clinical phenotypes nominally associated with burden

scores. Null distributions of the number of clinical phenotypes weakly associated with

burden scores were obtained using the same procedure to obtain the null distributions

for load scores (Figure 3 and Figure S3). The red dashed lines indicates the observed

number of clinical phenotypes nominally associated with genome-wide burden score (n

= 22, left), coding burden score (n=20, middle), and non-coding load score (n = 20,

right).

S1 Table. Linear regression between slopes and score categories.

S2 Table. Association between load score and the first 10 principal components.

S3 Table. Derived allele frequency stratification analysis.

S4 Table. phyloP score stratification analysis.

S5 Table. phyloP score and DAF stratification analysis.

S6 Table. Association between coding load score and the 28 phenotypes with number

of non-reference variants included as a covariate.

S7 Table. Association between load scores restricted to sites where the human genome

reference allele is the ancestral allele and 28 phenotypes.

S8 Table. Association between coding load scores computed from phyloPNH and 28

phenotypes.

S9 Table. Clinical phenotypes weekly associated with load scores.

671