High Content Analysis uncovers functionally relevant metastable phenotypic states in human Endothelial Cells monolayers

Francois Chesnais[1], Juliette Le Caillec[1], Errin Roy[2], Davide Danovi[2], Lorenzo Veschini[1#]

1 Vascular Cells Dynamics Lab, ACRS, Faculty of Dentistry Oral & Craniofacial Sciences, King's College London.
2 Stem Cell Hotel, Centre for Stem Cells & Regenerative Medicine, King's College London

Correspondence to lorenzo.1.veschini@kcl.ac.uk

Abstract

Endothelial cells (EC) present distinct cell properties in different tissues. Heterogeneity within the same vascular bed has been proposed to underpin emergent behaviours in EC monolayers. Quantification and functional relevance of EC phenotypic variance are challenging to assess. Here we developed a novel EC profiling tool (ECPT) to uniquely enable single EC profiling within a monolayer and providing spatial and relational information regarding cell proliferation, inter-endothelial Junctions and NOTCH activation. We used ECPT to characterise differential phenotypes in arterial, venous and microvascular EC populations. Our analysis highlighted extensive heterogeneity within individual monolayers and revealed VEGF-modulable metastability of NOTCH signalling which in turn regulates inter-endothelial junction's stability. We suggest that accounting adaptive emerging endothelial behaviours is necessary to develop specific tissue engineering approaches and identify more effective and EC-specific therapeutic targets for cardiovascular diseases and cancer.

Introduction

Endothelial cells (EC) form the inner layer of blood and lymphatic vessels and play major roles in tissue development, angiogenesis, inflammation and immune cell trafficking[1]. EC are functionally plastic and rapidly adapt to changes in the environment to preserve homeostasis. Local EC dysregulation is a hallmark of pathological states such as atherosclerosis, ischemia and cancer[2]. EC from different tissues and vascular beds (such as arteries, capillaries, veins) exhibit distinct metabolism, morphology and gene expression[3] and contribute in diverse ways to tissue development and regeneration[4,5]. It is well established that EC are phenotypically heterogenous not only among different tissues, reflecting specialised organ-specific functions[6], but also within the same tissue. Pools of EC substitute senescent cells, maintaining endothelial homeostasis; the role of endothelial progenitor cells with high repopulating potential has been highlighted, in particular in large vessels endothelia[7]. Inter-endothelial adherens junctions (IEJ) are dynamically regulated by VE cadherin (VEC) shuffling between the cell membrane and intracellular compartments. This process presents variations across vascular beds[3] and involves molecular mechanisms including VEGF receptors, cytoskeletal proteins and NOTCH family members. VEC and NOTCH signalling are well studied in angiogenesis and development. Nonetheless, defining the role of these molecular mechanisms in EC monolayer maintenance is key to understand vessel homeostasis in different organs in the human body. Understanding the heterogeneity of EC will also have important implications to realise the full potential of tissue engineering[8]. So far, a comprehensive analysis of EC cultures exploring and quantifying phenotypic variance has proven prohibitively difficult because of lack of adequate tools.

Single cell analysis is important to identify and characterise intermediate cell states[9,10] and to demonstrate correspondence between phenotypes and function[11] despite challenges in discriminating functional phenotypes variance from biological noise[12]. Single cell transcriptomic data is becoming available (mostly in mouse[13]) and could in principle advance the characterisation of human EC [14] and provide an overview of molecular processes in distinct EC populations. Nonetheless, spatial information is essential to map variable phenotypes to function and is also required to understand higher-level functions depending on cell-cell

connectivity. Of note, cell cycle and rapid adaptative phenomena, such as immediate increase
of endothelial permeability upon VEGF stimulation, rely on built-in sensing mechanisms that
depend on cell-cell interaction[15]. The study of human EC models has largely concentrated on
single cells, due to the limited availability of tools to study interactions at the population level.
Importantly, EC monolayers maintain their integrity over years while exerting a variety of
system-level functions which are emerging properties of cells in contact[16]. It has been in fact
proposed that endothelial adaptability and diversity of functions within an EC population
depends on cell heterogeneity[17].
64
Here, we present a novel comprehensive high-content analysis (HCA) platform to characterise
single EC within an endothelial monolayer capturing a wealth of cell features, context features
and subcellular features including extensive analysis of IEJ. This unbiased approach allows
quantification of EC diversity and feature variance comprehensively[12]. We set up an
endothelial cell profiling tool (ECPT) based on image analysis. Using an innovative machine
learning based method (Appendix 1), we addressed automated and accurate classification of
IEJ using junctional VEC immunostaining, and evaluation of NOTCH activation at the single
cell level. In total, we present here data from 4 independent acquisitions in triplicate technical
replicates detailing >7.5 M cell objects from >100,000 cells and >6000 images on a 29-
dimensional parameter space. This approach demonstrates the feasibility of using ECPT as
an exploratory platform to quantify heterogeneity in EC.
We take advantage of ECPT to address whether intermediate EC phenotypes exist within
homogeneous EC populations and whether these are functionally relevant. Single EC profiling
at such scale have not been reported and we here demonstrate its validity to uncover key
relationships across features and to comprehensively characterise cell behaviour across
distinct EC types.
81
82 **Results:**
83
HUVEC, HAoEC and HPMEC present heterogeneous phenotypes and subpopulations
can be clustered by automated analysis
While heterogeneity of EC has been proposed, there are few examples of studies that
achieved successful quantification phenotypic variance of EC within the same vascular bed
by transcriptomic analysis[18]. We previously reported the value of specific VEC and NOTCH
quantification to benchmark EC in intact monolayers[19] and we now extended our observations
including relational information between individual cells which are impossible to retrieve with
current single cell transcriptomic analysis.
Here, we seeded primary HUVEC, HAoEC and HPMEC under low FBS conditions for 48 hours
in the absence or presence of supra-physiologic VEGF. We aimed for sufficient cells to reach
confluency upon adhesion and spreading (~24 h) and we cultured for a further 24 h to enable
formation of stable IEJ. EC demonstrate a uniform cobblestone morphology under low
magnification, confirming monolayer stabilisation. In analysing these cultures, we focussed on
proliferation, junctions and NOTCH activation. We stained for VEC, activated NOTCH (NICD),
actin cytoskeleton and nuclei. Qualitative inspection at high magnification revealed remarkable
morphological differences between the three cell types in culture. HUVEC appear smaller and
display a relatively disorganised cytoskeleton (Supplementary Figure 1); HPMEC in contrast
are largest and often demonstrate cortical distribution of the cytoskeleton; HAoEC show
irregular shapes and sizes and often present as cells with stress fibres (Supplementary Figure
1).
104
Altogether these qualitative observations point to a complex interplay of proliferation, IEJ and
NOTCH that mirrors only in part previous studies referring to EC heterogeneity and distinct
traits[3,19,20]. Measuring and scoring individual EC phenotypes to demonstrate, explain and
quantify EC heterogeneity at single cell resolution is a prohibitively tedious and biased
approach if not automated. Therefore, we set to assemble a dedicated automated platform
that can seamlessly and comprehensively quantify variance of phenotypic parameters in EC

in appropriate multidimensional space to ultimately define specific cell populations. To serve
this purpose, we developed a method to profile EC phenotypes at single cell level based on
extracted HCA-features, Endothelial Cell Profiling Tool (ECPT, Figure 1 and Appendix 1).
114
A list of 29 features extending and refining our previous proof-of-principle efforts[19] was chosen
including proliferation and context feature and machine learning-aided classification of
junctions and of NOTCH activation.
We resourced to tSNE based dimensionality reduction techniques to capture variance within
our multivariate datasets and to identify discrete populations among heterogeneous cells. We
first asked whether the three EC cell types analysed cluster using the chosen features. EC
from different cell types formed distinct clusters (Supplementary Figure 2A) suggesting that
our parameters of choice (Supplementary Table 1) capture key differences in the three EC
populations. Nonetheless, similarities across the populations are also observed (e.g., mean
VEC intensity, Supplementary Figure 2B).
125
Altogether, tSNE enabled us to (1) specify diversities between EC populations and (2)
demonstrate a remarkable heterogeneity within the same populations. To understand the
origin of this heterogeneity we dissected individual sets of features dependent on the same
biological mechanism. To subset our database according to tSNE clustering we also
developed a dedicated open-source Shiny app (https://shiny.rstudio.com) allowing interactive
and iterative selection of single cells or groups of cells on tSNE scatterplots. To understand in
detail phenotypic differences, we proceeded with the exploration of features focussing on
selected traits to quantify variance in proliferation, junction stability and NOTCH activation.
134
Variance in proliferation: HUVEC, HAoEC and HPMEC present different proliferation
potential validated by context features
By culturing HUVEC, HAoEC and HPMEC under the same standardised conditions, we
noticed that the three EC types present different intrinsic proliferation rates. We quantified
proliferating cells within the different EC types using ECPT (Figure 2B, C Supplementary
Figure 3). Proliferation rate was higher in HAoEC than in HUVEC at baseline (average 13.61
vs 11.72 % of dividing cells per field) and lower in HPMEC (average 6.81%). VEGF induced
a small increase in proliferation rate in all EC. We then asked whether EC proliferation was
mostly due to clonal proliferation as suggested[7,21] or whether EC were retaining general
proliferation potential and how this might differ in distinct EC cell types. In previous work form
our group we demonstrated that the tendency to form colonies in hiPSC correlates with
differential gene expression programs[22]. Following a similar strategy and the method
described below and in Supplementary Figure 4 we evaluated the tendency to form densely
packed cell clusters as a proxy of EC clonal proliferation. We developed a novel measure of
nuclei clustering as a proxy of clonal proliferation based on the assumption that fast
proliferating clones will group with reduced cytoplasm and reduced internuclear distance
(Supplementary Figure 4). We extracted subsets of cells by manual annotation of plots of
interest (Figure 1D, Appendix 1) and explored further subsets of the database according to six
clusters identified by tSNE (Figure 3A).
Figure 3B shows the mapping of the nuclei clustering index (NCI, defined as the number of
nuclei within a distance less than ~1/3 the mean cell diameter, Appendix 1). In other words,
within our definitions an NCI equal or above 2 indicates that a cell has two or more very close
neighbours, and this is strongly indicative that the cell resides within a group of packed cells
(Supplementary Figure 4). We then calculated average second cell neighbour distance
measurements for each cluster and found that the cells within clusters 1 and 2 had the lowest
average distance to second neighbours, those in cluster 3 and 5 an intermediate average
distance and clusters 4 and 6 were the most distanced cells (Figure 3F). Therefore, cells in
tSNE clusters 1 and 2 displayed both nuclei clustering and decreased 2^{nd} neighbour distance
indicating that cells within these clusters are more closely packed than the rest of the
population. Since closely packed cells reflect compact colonies, we asked whether these
colonies might represent highly proliferative subclones. Figure 3D shows mapping of

proliferating cells on the tSNE plot and it appears that cells in compact colonies (clusters 1
and 2) have a higher frequency of proliferating cells. Subsequently, we calculated enrichment
of dividing cells in the tSNE clusters. We found that LM cells were enriched in clusters 1 and
2 and G2/M cells in clusters 3 and 4 (Figure 3C, D). This validated our observations. Notably,
cluster 3 also contained cells with a nuclei clustering index >2 (Figure 3B), had decreased 2nd
neighbour distance, albeit higher than cluster 1 and 2, and an increased frequency of LM cells.
Furthermore, cluster 3 and 4 contains >95% of all G2/M cells whereby cells in cluster 4 have
both high 2nd neighbour distance and a nuclei clustering index <2. The same pattern was
observed in all EC lines when analysed individually (Supplementary Figure 5A).
Overall, this data demonstrates that 1) highly proliferating EC clones exist within EC
monolayers (clusters 1, 2 and 3) and 2) that clonal proliferation is not the only modality of EC
proliferation, as isolated G2/M and LM cells were also found scattered throughout the
monolayers (clusters 4, 5 and 6), (Figure 3C-F).
When we looked at differences between HAoEC, HPMEC and HUVEC we found that all cells
examined had the capacity to proliferate both within colonies and in isolation. Furthermore,
proliferation in isolation was more frequent than in colonies in all cell populations examined
(Supplementary Figure 5B). Overall, all the EC lines we cultured continued proliferating up to
8 passages with little signs of clonal proliferation (i.e., sparse EC from all lines proliferated
homogeneously, not shown). Although few seemingly highly proliferative clusters were
identifiable, dividing cells were mostly found interspersed with quiescent cells within stabilised
monolayers in all cell lines (Supplementary Figure 5B) suggesting that proliferative capacity is
largely retained across the EC monolayers, including HPMEC which are more specialised and
can proliferate, albeit at a reduced rate.
189
Distinct EC types display heterogeneous activation states
The evaluation of EC activation state has been previously based on actin cytoskeleton
organisation[23] (Supplementary Figure 1). EC with stress fibres have been considered
motile/migratory while those in which peripheral cytoskeleton overlap with cell junctions are
considered quiescent. The former EC phenotypes are usually associated with cell elongation
and unstable IEJ.
In line with previous observations[19], VEGF induced elongation in all cells examined
(Supplementary Figure 6). At simple microscopic observation, HPMEC homogeneously
demonstrated features of quiescence such as cortical cytoskeleton supporting stable junctions
and surprisingly despite evidence of cell elongation VEGF did not evidently affect junctional
stability in HPMEC. HAoEC and HUVEC instead presented a more heterogeneous phenotype,
appeared more active and chronic VEGF treatment promoted migratory phenotypes in both
cell types. Furthermore, VEGF treatment induced more stress fibres in HAoEC but not in
HUVEC in contrast to previous observations under acute VEGF stimulation[24]. To substantiate
these observations, we measured EC activation by ECPT. To this purpose, we trained a
machine-learning based software (Cell Profiler Analyst, CPA based on fast gentle boosting)
to classify quiescent, activated or stressed EC (Figure 4A, Supplementary Figure 7, Appendix
1). Under this classification, 40.6 % of HPMEC were quiescent cells, substantially more than
HAoEC (20.5%) and HUVEC (16.2%). An opposite trend was observed for stressed cells
(HAoEC, 25.4%; HPMEC, 13.7%; HUVEC 23.2%). VEGF had opposite effects on HAoEC
and HUVEC, quiescent cells were reduced (-8.2%) in HAoEC and increased (+4%) in HUVEC
while stressed cells were increased in HAoEC (+5.3%) and reduced it in HUVEC (-7.1%),
(Figure 4B, Supplementary figure 7). VEGF treatment had little effects on HPMEC overall
quiescence save a small increase in quiescent cells (+2.2%).
Dividing (Late mitotic) cells, as shown in Figure 2A were binned in a specific category which
was used to refine cell cycle analysis based on DNA intensity (Appendix 1) overcoming a
specific limitation of previous approaches[25]. Overall, these data demonstrate that HPMEC
largely do not acquire a migratory phenotype and tend to be more quiescent than HAoEC and
HUVEC. Chronic supra physiologic VEGF treatment increased the proportion of activated
HAoEC yet interestingly it promoted quiescence in the two other EC lines analysed
(Supplementary Figure 7) suggesting that the effects of VEGF are context specific.

Furthermore, all EC populations had a significant proportion (46.1-55.6%) of activated cells
(i.e., an intermediate, neither migratory nor quiescent and non-proliferating, state
characterised by reduced connectivity) further demonstrating that EC monolayers are
composed of heterogeneous EC.
225
Variance in junctions: EC from different vessels display heterogeneous junction
composition
IEJ morphology is an immediate index of EC activation in contrast to quiescence. In general,
quiescent EC (i.e., non-proliferating, non-migrating) are also those with stabilised IEJ while
migrating or proliferating EC demonstrate jagged or discontinuous junctions[23,26]. Average
population measurements have been used by us and others[19,24,27] to assess stability of EC
monolayers which in turn correlate with functions such as trans-endothelial permeability to
large proteins[28].
Assessing IEJ at the single cell level is key to highlight subtle nuances in inter-endothelial cells
connectivity within the same monolayer. Nonetheless, measuring and classifying junctional
signal by image analysis is technically challenging and no currently available software is able
to evaluate multiple images in a standardised way[29,30]. To overcome this obstacle, we have
implemented within ECPT a junction analysis method (Appendix 1). Our method subdivides
the junctional inter-cellular space in several discrete areas scored using an expert-trained
machine learning algorithm (CPA with Fast gentle Boosting, Appendix 1). In line with the
literature in the field, we classified each junction binning in three categories: jagged, linear or
stabilised (Figure 5A). Stabilised junctions are visually reminiscent of the reticular junctions
previously described[26]. To facilitate data representation, we calculated a single continuous
measure accounting for the proportions of different IEJ categories (junctional stability index,
STB Index, Appendix 1).
246
Comparing Figure 1E (tSNE mapping of the three EC lines) with Figure 5B (tSNE mapping of
STB Index) it appears that the different EC lines map on areas with different STB Index a large
proportion of HPMEC mapped to area with high STB Index indicating that their junctions are
mostly composed of stabilised and linear junctions in contrast to the other two lines where
junction's composition was very heterogeneous. To quantify differences, we evaluated the
density distribution of the STB index across the three cell types (Figure 5C). It is evident that
both HAoEC and HUVEC are very heterogeneous in respect to IEJ, and that many cells have
a composition of different junctions. The mean STB Index was -4.7 (62.1 median IQR) for
HAoEC (prevalence of linear junctions) and -23 (64.6 median IQR) for HUVEC (mixture of
jagged and linear junctions). In contrast, HPMEC had prevalence of linear and stabilised
junctions (30 mean) and were more homogeneous (44.2 median IQR).
Chronic exposure to 50ng/ml VEGF increased STB index in all cells to different extents (+2.2,
+5.6 and +16.2 in HAoEC, HPMEC and HUVEC respectively), HPMEC junctions were also
more homogeneous upon VEGF treatment (36.9 median IQR) (Figure 5C).
In vitro conditions might not support full stabilisation of IEJ therefore these results should be
interpreted as differences in intrinsic EC potential to form stable monolayers in contrast to
plastic but unstable (and potentially leaky) ones, rather than a faithful reflection of *in vivo*
reality. Within these limitations, our results demonstrate that HPMECs' junctions are
intrinsically more stable producing homogeneously stable monolayers in contrast to HAoEC
and HUVEC which tend to form heterogeneous and less stable monolayers which is
compatible with their differential functions, i.e., a reduced requirement to tightly regulate
paracellular transit in EC from large vessels.
269
Variance in NOTCH activation: Quantification of NOTCH activation in EC monolayers
highlights metastable cell states
In line with previous results on HEY1/HEY2 expression in HUVEC[31], we hypothesised that EC
within monolayer would have a homogeneous level of NOTCH signalling either low/absent in
venous EC (HUVEC) and high in arterial EC as high NOCTH signalling is associated with the

arterial phenotype[32]. One previous study has reported appreciable baseline NICD levels in
HPMEC by western blotting[33] but not in comparison with other EC lines.
Computational models of NOTCH signalling pathway dynamics suggest that the NOTCH
system can generate either mono-stable, bi-stable or unstable/metastable phenotypes.
Unstable/metastable dynamics have never been described in EC[34,35,36]. A bi-stable scenario
would entail that most non-dividing cells in a monolayer will be either NOTCH active or non-
active which should promote formation of regular patterns of ON and OFF cells (Figure 6A).
Conversely, if the system cannot reach a stable equilibrium (non-stability, metastability), we
would expect to observe distribution between different activation states including cells without
NICD (NOTCH OFF), actively signalling cells with varying NICD levels and cells with high
nuclear NICD (NOTCH ON) (Figure 6A).
In order to distinguish between these possibilities, we measured the amount of NICD puncta
in the cytoplasm and nuclei of single EC using ECPT. Figure 6B shows the distribution of
NOTCH among four classes derived from cytoplasmic and nuclear NOTCH puncta
measurements. The thresholds were selected empirically, <50 total NICD intensity reflects
presence of 1-5 NICD puncta, in comparison, most highly active cells had ~50-80 NICD puncta
and ~1600 total NICD intensity (Figure 6C, Appendix 1). Interestingly, we found a continuous
nuance of differential NICD intensities across this range (0-~1600, Fig 6C). Our results
indicate that irrespectively of origin, EC within monolayers are heterogeneous in NOTCH
activation (Classification in Figure 6B). In line with current literature, HUVEC demonstrated
higher degree of inactive cells (OFF) and lower percentage of highly active and ON cells
(Figure 6B, Supplementary Figure 8) in comparison to HAoEC and HPMEC. NOTCH activity
in pulmonary microvascular EC has not been compared to other EC previously, we found that
HPMEC had the highest degree of high NOTCH highly active (~31.3%) and ON cells (~17.7%)
among the EC types examined (Supplementary Figure 8). Interestingly, all EC had a
significative presence of all four categories including relevant portions of "ON" and "OFF" cells
demonstrating that the system is not mono-stable (Fig 6A). Analysis of static images cannot
provide information regarding the exact dynamics of the system, nonetheless NICD staining
conveys a temporal information as NICD puncta are necessarily indicating previous signalling
from neighbouring cells and they are inherently fated to translocate to the nucleus or to be
degraded before reaching the nucleus. When we plotted the distribution of all cells across
nuclear to cytoplasm NICD ratio (Figure 6D and Supplementary Figure 9), we observed two
peaks corresponding to OFF and ON cells and a significant proportion of cells with
intermediate states including cells with significant nuclear NICD and further ongoing signalling
in the cytoplasm. Qualitative inspection of the plots in Figure 6D and Supplementary Figure 9
immediately shows that 1) the plots are only compatible with unstable/metastable cell states
at the population level (Figure 6A) and 2) all EC examined follow a similar pattern suggesting
that NOTCH metastability might be a general feature of endothelia.
To assess the functional relevance of this phenomenon we evaluated correlations between
differential NOTCH activation states and the phenotypical features described in previous
results. NOTCH activation has been associated with increased junctional stability in EC[37].
When we examined correlation between NOTCH activity and junction stability at the single
cell level, we did not find any evidence of correlation between the two phenomena
(Supplementary Figure 10,11). Therefore, high level of NICD does not appear to affect
Junctional stability within the same (receiver) cell. We then moved our focus on cell
neighbourhoods by calculating field averaged measures of Junctional stability index and the
frequency of NICD classes (OFF, Low, High, or ON) and we found that the average stability
index increased in fields with high frequency of High cells and decreased in fields with high
frequency of Low cells (Figure 7 A and Supplementary Figure 8). Overall, our data suggest
that sustained NOTCH signalling in one cell favours junctional stabilisation in its
neighbourhood while low levels of NOTCH signalling are associated with junctional
destabilisation in neighbouring cells.
To confirm this observation, we evaluated the effect of VEGF in each cell type since, as shown
before, VEGF increases average STB_Index in all cells (Figure 5C). As shown in
Supplementary Figure 8B, we found that VEGF induced a significant reduction in the

frequency of NOTCH Low cells in HAoEC and HUVEC, this was accompanied by an increase
in NOTCH High cells suggesting lateral inhibition mechanisms. Consistently, junctional
stabilisation was less pronounced in HAoEC which have a high frequency of high cells at
baseline therefore lateral inhibition could be dampened by excess and continuous NOTCH
signalling in neighbouring cells.
These data demonstrate that VEGF treatment can modulate NOTCH signalling in HAoEC and
HUVEC with similar effects of promoting junctional stabilisation but to different extents
depending on initial conditions.
Finally, to verify that junctional stabilisation is associated with acquisition of overall quiescent
EC phenotype we calculated the correlation between field averaged frequency of quiescent
and stressed cells and junction stability index. Our results overall demonstrate a positive
correlation between junction stability index and the frequency of quiescent cells and a negative
correlation between the index and the frequency of stressed (migratory) cells in line with
previous literature (Figure 7B).
Taken together, our data demonstrate that chronic supra-physiologic VEGF induces junction
stabilisation in HAoEC and HUVEC via increase of NOTCH signalling coupled with lateral
inhibition in neighbouring cells and that this effect is dependent from initial conditions (intrinsic
NOCTH signalling). This effect is mediated by selective reduction of Low NOTCH signalling
EC and therefore, by demonstrating that heterogeneity in NOTCH activation produces
physiologically relevant and distinct phenotypes our data also provide a clear experimental
evidence that EC metastability within the same monolayer is functionally relevant. Finally,
chronic VEGF stimulation didn't significantly affect overall pattern of NOTCH active/inactive
cells demonstrating that metastability of NOTCH signalling in EC monolayers is robust to
chronic and supra-physiologic perturbation.
354
Discussion
EC exert a surprising variety of specialised functions[3,6] some of which are limited to specific
organs while others are common, and this variety of functions is reflected in EC phenotypical
heterogeneity. EC heterogeneity across different organs is explained by the activation of
differential genetic programs (e.g., arterial, venous, capillary and lymphatic EC)[20,45]. This EC
diversity arises during development, is preserved throughout homeostasis and is thought to
be led by micro environmental factors. Heterogeneity within the same vascular bed where
microenvironmental factors are stable (or undergoing minor fluctuations) has been observed[17]
and deemed crucial to understand variable functions within the same EC type. Strategies such
as Hedge Betting (where heterogeneity expands the repertoire of possible responses) or
Swarm Sensing (where pioneer cells can redirect the response of neighbouring cells in a
cascade) are thought to be employed by homogeneous cells population to expand the
repertoire of sensory and adaptative responses[11].
368
These mechanisms require similar cells to preserve discrete, different and adaptable
phenotypes and the time scale at which these phenotypes change (minutes) is much smaller
than that of gene expression (hours). Thus, to justify rapid phenotype switches it is necessary
to involve inherently dynamic processes such as junctional plasticity.
Direct evaluation of dynamic changes in cell connectivity, functions and phenotype ideally
requires observing live cells at a timescale compatible with that of the phenomena under
investigation and this poses considerable technical hurdles. Therefore, despite great
advances in live-microscopy, comprehensive cell phenotyping of single cells at population
level is still vastly relying on fixed immuno-stained cells. Although, observation of end-point
cells cannot provide information on the exact dynamics of the system, extensive profiling of
single cells can inform about cell dynamic behaviours and help to develop hypotheses which
can be demonstrated experimentally or with mathematical models and computational
methods[38].
382
Aiming to grasp a better understanding of endothelial cells' heterogeneity, connectivity and
emerging dynamic behaviours[16,17] we developed ECPT and characterised three EC lines

under standard conditions or supra-physiological (like those observed in cancer) levels of
VEGF. In line with previously published data[20] our data demonstrate that Human Aortic,
Umbilical Vein or Pulmonary EC present clearly distinct phenotypes (Figure 2-6), maintained
at steady state, independently from microenvironmental stimuli. The three cell types have
distinct intrinsic proliferative potential and demonstrate diffuse ability to proliferate, in contrast
to the idea that EC undergo terminal differentiation and lose proliferative capacity[7].
Our findings are consistent with a model of monolayer maintenance hosting intermediate EC
populations with different proliferation potential. This effect could be achieved by a stem-
precursor-differentiated hierarchy as proposed by Patel et al.[21], or by an alternative model
positing proliferating EC and quiescent EC as in metastable states (Figure 2-3 and
Supplementary Figure 12). Thus, single EC could undergo a temporary reversion of the
differentiation programme in response to specific conditions, generate an adequate progeny
and return to quiescence (Supplementary Figure 12)[39]. Such state would eliminate the need
for a defined intermediate EC phenotype and would be more efficient at repairing localised
endothelial damage. Our data support this latter concept as we found a significant fraction of
proliferating EC far from closely packed and highly proliferative colonies (Figure 3) and is also
supported by EC monolayer scratch assays where most cells at the edge of the scratch
promptly activate, migrate and start proliferating until confluence is reached (not shown)[40] and
by many evidences that EC proliferation is contact inhibited and chiefly regulated by
decreased EC connectivity and lateral inhibition (during angiogenesis)[41,42], implicating that the
default EC state might be proliferation not quiescence and that in turn quiescence should be
achieved actively[39]. Under our experimental conditions, both HUVEC and HAoEC had a
similar sustained proliferative potential (on average >10% cells in cell cycle for each
microscopic field) and as expected VEGF induced increased proliferation in all EC examined.
The observed induction was small (~1% in all cells) as under our experimental conditions EC
are confluent and thus largely contact inhibited. Nonetheless ongoing sustained proliferation
in a confluent monolayer suggest that EC might require additional stimuli (some of which might
be absent *in vitro*) to further inhibit proliferation. Overall, our data on proliferation support a
model where EC can enter cell cycle at need notwithstanding opposing signalling including
EC-EC contact. We also demonstrate that clusters of fast proliferating EC existed within
confluent monolayers and under our experimental conditions (confluent EC) is highly unlikely
that enough space is created to allow more than one/two cell divisions before insurgence of
contact inhibition. Therefore, these observations are consistent with the existence of highly
clonogenic EC sparsely scattered throughout the monolayer. It remains to be elucidated
whether these colony forming EC are a distinct subpopulation (i.e., a true EC precursor driven
by a specific transcriptional programme) or the result of stochastic fluctuations in expression
of genes related to cell cycle. In this sense, the Rafii laboratory[43] has recently demonstrated
that exogenous overexpression of ETV2 in primary EC reprogrammes cells toward a highly
clonogenic phenotype. Therefore, reactivation of ETV2 might be a potential physiologic
mechanism to rejuvenate EC pools by creating new EC precursors but evidences of this
occurring in physiologic conditions in vivo has not been reported so far and warrants further
investigations.

Inter-endothelial junctions (IEJ) are among the main drivers of EC biology mediating high-level
functions such as endothelial barrier permeability and leukocyte trafficking and low-level
functions such as contact inhibition of proliferation. Analysis of IEJ in three different EC
populations highlighted a high degree of junctional heterogeneity in HAoEC and HUVEC, while
HPMEC junctions where homogeneously stable. Here, we adopted the "stabilised"
nomenclature as these junctions are structurally more complex and wider than linear ones
which we interpreted as areas where EC membranes overlap forming a junctional plate
(Figure 5A). The amount of membrane engaged in the junction is associated with higher
junction strength since a larger area will correspond to more intercellular VEC-VEC

interactions which in turn are responsible for maintenance of adhesion[44]. Inter-endothelial
connectivity is also a regulator of contact-dependent signalling pathways. In line with previous
literature, we show that HAoEC (arterial) had more linear (stabilised) junctions than HUVEC
(venous) which instead tended to form highly dynamic jagged junctions. IEJ chiefly regulate
traffic of molecules and solutes in and out of capillaries and inhibit coagulation by preventing
exposure of underlying sub-intimal layer in arteries and veins. Importantly, HPMEC had
stabilised junctions and HAoEC had mostly linear junctions and presented stress fibres. This
is reminiscent of the unique role of the pulmonary and arterial vessels in tightly regulate solute
exchange or adapt to high flow respectively.

Distinctive features among different EC populations seem to be hard coded within cell gene
expression program as cells from individual donors had vastly overlapping phenotypes and
the features where maintained in all lines upon several passages in culture. Nonetheless,
analysis of single cell data highlighted extensive heterogeneity within the same population. If
decision whether to enter cell cycle or to maintain stable junction were dependent on
differential gene expression, we would expect to find a limited collection of homogeneous
phenotypes whereas we find an apparently continuous nuance of EC phenotypes within the
same EC monolayer. Furthermore, it is established that these decisions need to be taken at
a faster pace than gene expression could justify[17].

As mentioned, rapid phenotype switch between the same cell population is justified by
involving dynamic genetic circuits dominated by oscillatory or stochastic behaviours[10]. We
focussed our attention on NOTCH signalling as its role in EC biology and its crosstalk with the
junctional complex is well recognised[37]. Furthermore, NOTCH has been proposed as a master
switch which dynamics can deploy either mono-stable, bi-stable or metastable cell states[42]. In
the context of angiogenesis, it has been demonstrated that leading tip cells induce NOTCH
signalling in trailing stalk cells via Dll4-NOTCH1 leading to lateral inhibition of the tip cell
phenotype[42] whereas EC within a stabilised monolayer cannot acquire a tip cell phenotype
and all EC within a monolayer receive in principle similar NOTCH stimulation from neighbours.
Three scenarios could thus arise 1) mono-stable, where most non dividing cells accumulate
NICD to a characteristic carrying capacity proportional to basal expression levels of NOTCH
agonists such as Dll4 (e.g., high in arteries and low/absent in vein), 2) bi-stable, where cells
oscillate between ON and OFF states due to lateral inhibition/induction until they ultimately
settle on one of the two and 3) non-stable/metastable, where cells do not settle on specific
states and instead constantly oscillate generating in the process differential levels of
signalling. In a mono-stable scenario cells will favour a specific phenotype until perturbation
occurs. In a bi-stable scenario ON and OFF cells will tend to acquire characteristic distributions
or patterns (e.g., boundaries). Metastability is more flexible as the acquisition of phenotype
can depend on both the overall intensity of the original signal and the frequency at which this
oscillates, and this favour the emergence of biological noise and stochastic behaviours which
in principle can deploy a much higher degree of variability[17].
EC monolayers have been previously described as either NOTCH active or inactive especially
in the context of arteriovenous specification[45] and downstream NOTCH genes are often used
as surrogate markers of the arterial phenotype[46]. Fluctuations in NOTCH signalling have been
hypothesised in EC monolayers and demonstrated in EC during sprouting angiogenesis[42].
NOTCH signalling has been linked to both inhibition of EC proliferation[47] and increased
junction stabilisations[37] but NOTCH regulation in EC monolayer at the single cell resolution
has not been investigated previously.
By measuring NICD in a relatively large collection of single cell measurements, we found that
different EC within their monolayer acquire metastable NOTCH activation states which can be
modulated by chronic VEGF treatment and are used to regulate overall junctional stability of
the monolayer. Since NOTCH signalling is titratable[48] different extent of signalling could
generate different phenotypes with ON state. Therefore, our data provide a first demonstration
of recent hypotheses[16,17] where heterogeneity and metastability were postulated as relevant

mechanism for swarm sensing behaviours in EC monolayers and as underlying mechanisms
for EC phenotypical plasticity and adaptability.
We also show for the first time that microvascular EC can possess high levels of NOTCH
activity as demonstrated by our data on HPMEC, which cannot be explained by the currently
established model positing high NOTCH activity as a mediator of arterial phenotype. In other
words, despite sustained NOTCH signalling is a hallmark of arterial cells it cannot be
accounted as the sole or main pathway involved in the establishment of the arterial phenotype
warranting revision of protocols to generate stem cells derived arterial or venous EC[32].
NOTCH signalling in HPMEC wasn't affected by VEGF (Supplementary Figure 8 B) possibly
because the frequencies of NOTCH Low and NOTCH High cells were already the highest and
lowest respectively (within our experiments) at baseline and these levels might represent set
boundaries of possible VEGF modulation. Overall, our results regarding VEGF-mediated
NOTCH modulation in different EC lines strongly support the idea that the net effects of chronic
VEGF stimulation are sensibly dependent from initial conditions (i.e., the intrinsic baseline
NOTCH signalling levels) and might lead to appreciably different responses in different EC.
504
Altogether, our data provide the first direct evidence that metastable cell states orchestrate
the stability and plasticity of EC monolayers, and that EC within monolayers use NOTCH and
junction's machinery to constantly probe their neighbourhood relaying spatial and relational
information to the gene expression machinery and deploying different phenotypes in a
localised and context dependent manner. We suggest that phenotypical diversity within a
homogeneous EC population can represent a cellular implementation of "Bet hedging",
"Swarm sensing" or "Response distribution" mechanisms[15] to deploy a globally stable but
locally adaptable tissue such as the endothelium.
513
Our in vitro approach allows focussing on intrinsic EC mechanisms of monolayer maintenance
and plasticity excluding variability caused by microenvironmental factors and it has been
highlighted that the heterogeneity of different blood vessels found in vivo is sustained by EC
rather than perivascular cells heterogeneity[18]. Therefore, it seems highly probable that the
robust metastability in EC phenotypes described above has a direct relevance to more
complex in vivo context. It will be important to elucidate in future studies how the crosstalk
with perivascular cells and microenvironment can affect emerging endothelial behaviours.
521
522 These results imply that pharmacological interventions targeting the endothelium need to
523 account for dynamic adaptive mechanisms which might otherwise unpredictably dampen or
524 exacerbate the efficacy of EC targeting drugs (e.g., VEGF or VEGF receptors inhibitors).
525 Otherwise, adaptive mechanisms might be exploited to precisely targeting EC subsets of
526 interest while leaving most of the other EC un-perturbed.
527
In conclusion, we have built on our previous work to create ECPT, a powerful platform to
dissect single cell level phenotypical composition of EC monolayers enabling for the first time
to demonstrate EC heterogeneity as a driving force of endothelial plasticity. We envisage that
coupling ECPT with live imaging setups and single cells transcriptomic analysis will open the
way to a much deeper understanding of emerging dynamic endothelial behaviours and thus
to develop novel more effective therapies for prevention of cardiovascular diseases and the
treatment of cancer.
535
536
537 **Methods**
538
<u>Cell culture</u>
HUVEC, HAoEC and HDMEC (PromoCell) were plated on 10 µg/mL fibronectin (from human
plasma, Promocell)-coated flasks, grown in EGMV2 medium (Promocell) in the absence of
antibiotics, detached with Accutase (Thermo Fisher Scientific, Waltham, MA), and used by
passage 5. For experiments, 4 × 10^4 ECs per well were seeded in fibronectin-coated 96-well

plates (µclear, Greiner) and cultured for 48 h under basal (EGMV2, Promocell) or activated
(EGMV2 + 50 ng/mL VEGFA, Peprotech, London, UK) conditions in triplicate. The ECs formed
confluent monolayers at the time of immunostaining and image acquisition.
547
Immunostaining
Cells were fixed with 2% paraformaldehyde in phosphate buffered saline (PBS) for 10 minutes
at room temperature. Cells were blocked 1h with PBS supplemented with 1% fetal bovine
serum (FBS) and permeabilised with 0.1% Triton X 100. Cells were then incubated for 1h at
room temperature with primary antibodies against CDH5 (Ve-Cadherin Novusbio NB600-
1409, 1 µg/mL final) and activated NOTCH (NICD, Abcam ab8925, 1 µg/mL final). CDH5 is a
lineage marker for EC and cells were all CDH5+, the anti-activated NOTCH1 reveals NOTCH
intracellular domain (NICD) puncta. Plates were washed and incubated 1h with 1 µg/mL
secondary Alexa 488-conjugated antibody and Alexa-555-conjugated antibody (Thermo),
Hoechst 33342 (1 µg/mL, Sigma) and Phalloidin-Atto 647N (Sigma).
558
Image Acquisition
We obtained images from slides with an Operetta CLS system (PerkinElmer, Waltham, MA)
equipped with a 40×water-immersion lens (NA 1.1). In each well, 3 areas were acquired. Each
area is composed of nine microscopic fields at 40× magnification (Supplementary Figure 1).
We standardised acquisition parameters (led power, exposure time) throughout different
experiments and used HUVEC as a standard for calibration in all experiments. We analysed
an image database containing 6028 images (1507 fields in four fluorescence channels)
extracted from four independent experiments conducted on EC lines from two different donor
each. Three intraexperiment replicates were conducted for each experiment.
568
569
Endothelial Cell Profiling Tool
We used a combination of machine learning-aided image segmentation (ImageJ)[49] and an
image-based cell profiling tool (CellProfiler)[50] to extract the phenotype of single EC in
monolayers. Our workflow enables to measure EC morphology (Area, perimeter, shape
descriptors, cell neighbours), NICD, VE-Cadherin and DNA intensities and to characterise
inter-endothelial junctions (IEJ). In the present study, we chose to analyse only selected
features with recognised functions in EC biology. Image texture features were measured and
only used during the training of machine learning algorithm[51,52] which in turn were used to
classify cell activation states and junction morphology. ECPT scripts and methods are detailed
in Appendix 1 and available at https://github.com/exr98/HCA-uncovers-metastable-
phenotypes-in-hEC-monolayers
581
We Imported and cleaned the results into R studio excluding artifacts and mis-segmented cell
objects. We then calculated derived continuous (Junctional stability index) and categorical
(Cell Cycle and NOCTH activation) parameters (Supplementary Table 1). Following guidelines
suggested by Caceido et al.[53] We pre-processed the database to exclude miss-segmented
cells and to normalise the measurements prior to dimensionality reduction or single factor
analysis. Our final dataset was composed of measurements of 29 parameters from 104416
single cells (Supplementary Table 1)
We conducted PCA analysis on cluster of ECPT parameters (Cell morphology, intensity,
neighbourhood and junctional status, Supplementary Table 1) and generated 12 PCA
components which we fed into the tSNE algorithm.
We performed tSNE using different perplexity values (5, 10, 20, 40, 60, 80) and ran the
algorithm with 1000 iterations, all tSNE plots displayed correspond to 10 perplexity.
594
595 We created a shiny application for interactive selection of data of interest from t-SNE
596 scatterplots (Appendix 1). This application[54] allowed specific cluster analysis by subclustering
597 and further analysis in the R environment[55]. All files containing the code required to reproduce

the shiny application using our dataset are available on https://github.com/exr98/HCA-uncovers-metastable-phenotypes-in-hEC-monolayers.

Statistical analysis

To compare multiple groups, we used one-way ANOVA followed by Tuckey's HSD post-hoc test. We considered $p < 0.05$ (*) significative and $p < 0.01$ (**), $p < 0.001$ (***) and $p < 0.0001$ (****) highly significative. To evaluate linear correlation between continuous variables we calculated Pearson's Correlation coefficient (R) and considered $p < 0.001$ as highly statistically significative.

References

1- Potente, M., Gerhardt, H. and Carmeliet, P., 2011. Basic and Therapeutic Aspects of Angiogenesis. *Cell*, 146(6), pp.873-887.

2- Park-Windhol, C. and D'Amore, P., 2016. Disorders of Vascular Permeability. *Annual Review of Pathology: Mechanisms of Disease*, 11(1), pp.251-281.

3- Augustin, H. and Koh, G., 2017. Organotypic vasculature: From descriptive heterogeneity to functional pathophysiology. *Science*, 357(6353), p.eaal2379.

4- Kusumbe, A., Ramasamy, S. and Adams, R., 2014. Coupling of angiogenesis and osteogenesis by a specific vessel subtype in bone. *Nature*, 507(7492), pp.323-328.

5- Itkin, T., Gur-Cohen, S., Spencer, J., Schajnovitz, A., Ramasamy, S., Kusumbe, A., Ledergor, G., Jung, Y., Milo, I., Poulos, M., Kalinkovich, A., Ludin, A., Golan, K., Khatib, E., Kumari, A., Kollet, O., Shakhar, G., Butler, J., Rafii, S., Adams, R., Scadden, D., Lin, C. and Lapidot, T., 2016. Distinct bone marrow blood vessels differentially regulate haematopoiesis. *Nature*, 532(7599), pp.323-328.

6- Rafii, S., Butler, J. and Ding, B., 2016. Angiocrine functions of organ-specific endothelial cells. *Nature*, 529(7586), pp.316-325.

7- Yoder, M., 2017. Endothelial stem and progenitor cells (stem cells): (2017 Grover Conference Series). *Pulmonary Circulation*, 8(1), p.204589321774395.

8- Greco Song, H., Rumma, R., Ozaki, C., Edelman, E. and Chen, C., 2018. Vascular Tissue Engineering: Progress, Challenges, and Clinical Promise. *Cell Stem Cell*, 22(4), p.608.

9- Siu, D., Lee, K., Lo, M., Stassen, S., Wang, M., Zhang, I., So, H., Chan, G., Cheah, K., Wong, K., Hsin, M., Ho, J. and Tsia, K., 2020. Deep-learning-assisted biophysical imaging cytometry at massive throughput delineates cell population heterogeneity. *Lab on a Chip*, 20(20), pp.3696-3708

10- MacLean, A., Hong, T. and Nie, Q., 2018. Exploring intermediate cell states through the lens of single cells. *Current Opinion in Systems Biology*, 9, pp.32-41.

11- Dueck, H., Eberwine, J. and Kim, J., 2015. Variation is function: Are single cell differences functionally important? *BioEssays*, 38(2), pp.172-180.

12- Eling, N., Morgan, M. and Marioni, J., 2019. Challenges in measuring and understanding biological noise. *Nature Reviews Genetics*, 20(9), pp.536-548.

13- Kalucka, J., de Rooij, L., Goveia, J., Rohlenova, K., Dumas, S., Meta, E., Conchinha, N., Taverna, F., Teuwen, L., Veys, K., García-Caballero, M., Khan, S., Geldhof, V., Sokol, L., Chen, R., Treps, L., Borri, M., de Zeeuw, P., Dubois, C., Karakach, T., Falkenberg, K., Parys, M., Yin, X., Vinckier, S., Du, Y., Fenton, R., Schoonjans, L., Dewerchin, M., Eelen, G., Thienpont, B., Lin, L., Bolund, L., Li, X., Luo, Y. and Carmeliet, P., 2020. Single-Cell Transcriptome Atlas of Murine Endothelial Cells. *Cell*, 180(4), pp.764-779.e20.

14- Tikhonova, A., Dolgalev, I., Hu, H., Sivaraj, K., Hoxha, E., Cuesta-Domínguez, Á., Pinho, S., Akhmetzyanova, I., Gao, J., Witkowski, M., Guillamot, M., Gutkin, M., Zhang, Y., Marier, C., Diefenbach, C., Kousteni, S., Heguy, A., Zhong, H., Fooksman, D., Butler, J., Economides, A., Frenette, P., Adams, R., Satija, R., Tsirigos, A. and Aifantis, I., 2019. The bone marrow microenvironment at single-cell resolution. *Nature*, 569(7755), pp.222-228.

15- Acar, M., Mettetal, J. and van Oudenaarden, A., 2008. Stochastic switching as a survival strategy in fluctuating
environments. *Nature Genetics*, 40(4), pp.471-475.

16- McCarron, J., Lee, M. and Wilson, C., 2017. The Endothelium Solves Problems That Endothelial Cells Do Not
Know Exist. *Trends in Pharmacological Sciences*, 38(4), pp.322-338.

17- McCarron, J., Wilson, C., Heathcote, H., Zhang, X., Buckley, C. and Lee, M., 2019. Heterogeneity and
emergent behaviour in the vascular endothelium. *Current Opinion in Pharmacology*, 45, pp.23-32.

18- Chavkin, N. and Hirschi, K., 2020. Single Cell Analysis in Vascular Biology. *Frontiers in Cardiovascular
Medicine*, 7.

19- Wiseman, E., Zamuner, A., Tang, Z., Rogers, J., Munir, S., Di Silvio, L., Danovi, D. and Veschini, L., 2019.
Integrated Multiparametric High-Content Profiling of Endothelial Cells. *SLAS DISCOVERY: Advancing the
Science of Drug Discovery*, 24(3), pp.264-273.

20- Chi, J., Chang, H., Haraldsen, G., Jahnsen, F., Troyanskaya, O., Chang, D., Wang, Z., Rockson, S., van de
Rijn, M., Botstein, D. and Brown, P., 2003. Endothelial cell diversity revealed by global expression profiling.
Proceedings of the National Academy of Sciences, 100(19), pp.10623-10628.

21- Patel, J., Seppanen, E., Rodero, M., Wong, H., Donovan, P., Neufeld, Z., Fisk, N., Francois, M. and
Khosrotehrani, K., 2017. Functional Definition of Progenitors Versus Mature Endothelial Cells Reveals Key SoxF-
Dependent Differentiation Process. *Circulation*, 135(8), pp.786-805.

22- Vigilante, A., Laddach, A., Moens, N., Meleckyte, R., Leha, A., Ghahramani, A., Culley, O., Kathuria, A., Hurling,
C., Vickers, A., Wiseman, E., Tewary, M., Zandstra, P., Durbin, R., Fraternali, F., Stegle, O., Birney, E., Luscombe,
N., Danovi, D. and Watt, F., 2019. Identifying Extrinsic versus Intrinsic Drivers of Variation in Cell Behavior in
Human iPSC Lines from Healthy Donors. *Cell Reports*, 26(8), pp.2078-2087.e3.

23- Millán, J., Cain, R., Reglero-Real, N., Bigarella, C., Marcos-Ramiro, B., Fernández-Martín, L., Correas, I. and
Ridley, A., 2010. Adherens junctions connect stress fibres between adjacent endothelial cells. *BMC Biology*, 8(1).

24- Veschini, L., Crippa, L., Dondossola, E., Doglioni, C., Corti, A. and Ferrero, E., 2011. The vasostatin-1 fragment
of chromogranin A preserves a quiescent phenotype in hypoxia-driven endothelial cells and regulates tumor
neovascularization. *The FASEB Journal*, 25(11), pp.3906-3914.

25- Roukos, V., Pegoraro, G., Voss, T. and Misteli, T., 2015. Cell cycle staging of individual cells by fluorescence
microscopy. *Nature Protocols*, 10(2), pp.334-348

26- Fernández-Martín, L., Marcos-Ramiro, B., Bigarella, C., Graupera, M., Cain, R., Reglero-Real, N., Jiménez, A.,
Cernuda-Morollón, E., Correas, I., Cox, S., Ridley, A. and Millán, J., 2012. Crosstalk Between Reticular Adherens
Junctions and Platelet Endothelial Cell Adhesion Molecule-1 Regulates Endothelial Barrier
Function. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 32(8).

27- Veschini, L., Belloni, D., Foglieni, C., Cangi, M., Ferrarini, M., Caligaris-Cappio, F. and Ferrero, E., 2006.
Hypoxia-inducible transcription factor–1 alpha determines sensitivity of endothelial cells to the proteosome inhibitor
bortezomib. *Blood*, 109(6), pp.2565-2570.

28- Ferrero, E., Scabini, S., Magni, E., Foglieni, C., Belloni, D., Colombo, B., Curnis, F., Villa, A., Ferrero, M. and
Corti, A., 2004. Chromogranin A protects vessels against tumor necrosis factor α-induced vascular leakage. *The
FASEB Journal*, 18(3), pp.554-556.

29- Brezovjakova, H., Tomlinson, C., Mohd Naim, N., Swiatlowska, P., Erasmus, J., Huveneers, S., Gorelik,
J., Bruche, S. and Braga, V., 2019. Junction Mapper is a novel computer vision tool to decipher cell–cell contact
phenotypes. *eLife*, 8.

30- Seebach, J., Taha, A., Lenk, J., Lindemann, N., Jiang, X., Brinkmann, K., Bogdan, S. and Schnittler, H., 2015.
The CellBorderTracker, a novel tool to quantitatively analyze spatiotemporal endothelial junction dynamics at the
subcellular level. *Histochemistry and Cell Biology*, 144(6), pp.517-532.

31- Aranguren, X., Beerens, M., Coppiello, G., Wiese, C., Vandersmissen, I., Lo Nigro, A., Verfaillie, C., Gessler,
M. and Luttun, A., 2013. COUP-TFII orchestrates venous and lymphatic endothelial identity by homo- or hetero-
dimerisation with PROX1. *Journal of Cell Science*, 126(5), pp.1164-1175.

32- Zhang, J., Chu, L., Hou, Z., Schwartz, M., Hacker, T., Vickerman, V., Swanson, S., Leng, N., Nguyen, B.,
Elwell, A., Bolin, J., Brown, M., Stewart, R., Burlingham, W., Murphy, W. and Thomson, J., 2017. Functional

characterization of human pluripotent stem cell-derived arterial endothelial cells. *Proceedings of the National Academy of Sciences*, 114(30), pp.E6072-E6078.

33- Zong, D., Li, J., Cai, S., He, S., Liu, Q., Jiang, J., Chen, S., Long, Y., Chen, Y., Chen, P. and Ouyang, R., 2018. Notch1 regulates endothelial apoptosis via the ERK pathway in chronic obstructive pulmonary disease. *American Journal of Physiology-Cell Physiology*, 315(3), pp.C330-C340.

34- Falo-Sanjuan, J. and Bray, S., 2019. Decoding the NOTCH signal. *Development, Growth & Differentiation*, 62(1), pp.4-14.

35- Agrawal, S., Archer, C. and Schaffer, D., 2009. Computational Models of the NOTCH Network Elucidate Mechanisms of Context-dependent Signaling. *PLoS Computational Biology*, 5(5), p.e1000390.

36- Vujovic, F., Hunter, N. and Farahani, R., 2019. NOTCH pathway: a bistable inducer of biological noise?. *Cell Communication and Signaling*, 17(1).

37- Bentley, K., Franco, C., Philippides, A., Blanco, R., Dierkes, M., Gebala, V., Stanchi, F., Jones, M., Aspalter, I., Cagna, G., Weström, S., Claesson-Welsh, L., Vestweber, D. and Gerhardt, H., 2014. The role of differential VE-cadherin dynamics in cell rearrangement during angiogenesis. *Nature Cell Biology*, 16(4), pp.309-321.

38- Altschuler, S. and Wu, L., 2010. Cellular Heterogeneity: Do Differences Make a Difference?. *Cell*, 141(4), pp.559-563.

39- Marescal, O. and Cheeseman, I., 2020. Cellular Mechanisms and Regulation of Quiescence. *Developmental Cell*, 55(3), pp.259-271.

40- Yang, Y., Jamilpour, N., Yao, B., Dean, Z., Riahi, R. and Wong, P., 2016. Probing Leader Cells in Endothelial Collective Migration by Plasma Lithography Geometric Confinement. *Scientific Reports*, 6(1).

41- Bocci, F., Onuchic, J. and Jolly, M., 2020. Understanding the Principles of Pattern Formation Driven by NOTCH Signaling by Integrating Experiments and Theoretical Models. *Frontiers in Physiology*, 11.

42- Ubezio, B., Blanco, R., Geudens, I., Stanchi, F., Mathivet, T., Jones, M., Ragab, A., Bentley, K. and Gerhardt, H., 2016. Synchronization of endothelial Dll4-NOTCH dynamics switch blood vessels from branching to expansion. *eLife*, 5.

43- Palikuqi, B., Nguyen, D., Li, G., Schreiner, R., Pellegata, A., Liu, Y., Redmond, D., Geng, F., Lin, Y., Gómez-Salinero, J., Yokoyama, M., Zumbo, P., Zhang, T., Kunar, B., Witherspoon, M., Han, T., Tedeschi, A., Scottoni, F., Lipkin, S., Dow, L., Elemento, O., Xiang, J., Shido, K., Spence, J., Zhou, Q., Schwartz, R., De Coppi, P., Rabbany, S. and Rafii, S., 2020. Adaptable haemodynamic endothelial cells for organogenesis and tumorigenesis. *Nature*, 585(7825), pp.426-432.

44- Giannotta, M., Trani, M. and Dejana, E., 2013. VE-Cadherin and Endothelial Adherens Junctions: Active Guardians of Vascular Integrity. *Developmental Cell*, 26(5), pp.441-454.

45- Adams, R. and Alitalo, K., 2007. Molecular regulation of angiogenesis and lymphangiogenesis. *Nature Reviews Molecular Cell Biology*, 8(6), pp.464-478.

46- Mack, J. and Iruela-Arispe, M., 2018. NOTCH regulation of the endothelial cell phenotype. *Current Opinion in Hematology*, 25(3), pp.212-218.

47- Fang, J., Coon, B., Gillis, N., Chen, Z., Qiu, J., Chittenden, T., Burt, J., Schwartz, M. and Hirschi, K., 2017. Shear-induced NOTCH-Cx37-p27 axis arrests endothelial cell cycle to enable arterial specification. *Nature Communications*, 8(1).

48- Henrique, D. and Schweisguth, F., 2019. Mechanisms of NOTCH signaling: a simple logic deployed in time and space. *Development*, 146(3), p.dev172148.

49- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J., White, D., Hartenstein, V., Eliceiri, K., Tomancak, P. and Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. *Nature Methods*, 9(7), pp.676-682.

50- Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang IH, Friman O, Guertin DA, Chang JH, Lindquist RA, Moffat J, Golland P, Sabatini DM (2006) CellProfiler: image analysis software for identifying and quantifying cell phenotypes. Genome Biology 7:R100.

51-Jones, T., Kang, I., Wheeler, D., Lindquist, R., Papallo, A., Sabatini, D., Golland, P. and Carpenter, A., 2008.
CellProfiler Analyst: data exploration and analysis software for complex image-based screens. *BMC*
Bioinformatics, 9(1), p.482.

52-Jones, T., Carpenter, A., Lamprecht, M., Moffat, J., Silver, S., Grenier, J., Castoreno, A., Eggert, U., Root, D.,
Golland, P. and Sabatini, D., 2009. Scoring diverse cellular morphologies in image-based screens with iterative
feedback and machine learning. *Proceedings of the National Academy of Sciences*, 106(6), pp.1826-1831.

53- Caicedo, J., Cooper, S., He0igwer, F., Warchal, S., Qiu, P., Molnar, C., Vasilevich, A., Barry, J., Bansal, H.,
Kraus, O., Wawer, M., Paavolainen, L., Herrmann, M., Rohban, M., Hung, J., Hennig, H., Concannon, J., Smith, I.,
Clemons, P., Singh, S., Rees, P., Horvath, P., Linington, R. and Carpenter, A., 2017. Data-analysis strategies for
image-based cell profiling. *Nature Methods*, 14(9), pp.849-863.
803

804   54- Winston Chang, Joe Cheng, JJ Allaire, Yihui Xie and Jonathan McPherson (2020). shiny: Web Application
805   Framework for R. R package version 1.5.0. https://CRAN.R-project.org/package=shiny
806
807   55- R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical
808   Computing, Vienna, Austria. URL https://www.R-project.org/
809
810
811

Figure 1 : ECPT workflow.

A) Acquisition of high resolution (40X OM) images on each channel (represented in false colours, original images are grey scale).

B) Features extraction and representation (ImageJ, Cell Profiler, Cell Profiler Analyst, R Studio).

C) tSNE plots of features normalised and grouped by biological function. Colouring in panels (Top to bottom) correspondent to Cell cycle, STB junctions, NOTCH Activation classes and Cell activation classes.

D) Shiny app (https://shiny.rstudio.com) enabling exploration of datasets, data extraction and iterative exploration.

Figure 2: A) Late mitotic cells, LM. LM, were counted by AI aided cell classification (CPA, Fast gentle boosting). B) Cell cycle analysis of HUVEC, HAoEC and HPMEC in basal conditions or upon treatment with 50 ng/ml VEGF. C) Average of dividing (G2/M+S+LM) HUVEC, HAoEC and HPMEC per microscopic field in basal conditions or upon treatment with 50 ng/ml VEGF.

Figure 3: A) tSNE clusters mapped on tSNE scatterplot. B) Mapping of Nuclei Clustering Index in tSNE scatterplot. C) Mapping of Cell cycle categories in tSNE scatterplot. D) Average 2nd neighbour distance per well in the six clusters. E) Percent of late mitotic (LM) cells per well in the six clusters. F) Percent of G2/M cells per well in the six clusters. ** p<0.001, $ p<0.0001 against clusters 4 and 6, # p<0.0001 against clusters 4,5,6, & p<0.0001 against clusters 1,2,5,6.

Figure 4: A) Examples of quiescent, activated and stressed EC. B) Cell activation classes in HUVEC, HAoEC and HPMEC in basal conditions or upon treatment with 50 ng/ml VEGF.

Figure 5: A) Examples of stabilised, linear and jagged or discontinuous junctions. B) Mapping of junctional stability index (a) and cell types (b) on tSNE scatterplot. C) Distribution of STB Index in HUVEC, HAoEC and HPMEC either under basal conditions (black) or 50ng/ml VEGF (red).

Figure 6: A) Schematic representing the possible configurations of notch activation at the population level. B) Distribution of NOTCH activation classes in HAoEC, HPMEC and HUVEC in basal conditions or upon treatment with 50 ng/ml VEGF C) Total vs Nuclear NICD intensity scatter plot. Cells without NICD are concentrated on the origin of the axes. Cells aligned along y axis (Nu Fraction =0) have varying degree of cytoplasmic NOTCH puncta and none in the nuclei. Cells on the diagonal (Nu fraction =1) have varying degrees of NOCTH puncta in the nuclei and none in the cytoplasm. Colours indicates the threshold used in the definition of categories in B, the threshold is set to 50 corresponding to 1-2 NOTCH puncta either in cytoplasm or nucleus. D) Density plot of Nuclear fraction of NICD intensity. The two peaks of NOTCH OFF and NOTCH ON (NICD Nu Fraction = 1) cells are evident. The majority of cells had intermediate activation states (NICD Nu Fraction >0 and < 1)

Figure 7: A) Field averaged measurements of Junctional stability index and percentage of NICD "Low" or "ON". Stability junction index positively correlated (R=0.43, P<0.001) with the frequency of ON cells and anticorrelated with the frequency of Low cells (R= -0.3, P<0.001). B) Field averaged measures of Junctional stability index and percentage of "quiescent" or "Stressed" cells.

Supplementary Figure 1: Microphotographs comparing HUVECs, HAoEC and HPMEC untreated immunostained for VE-cadherin (green), NICD (orange), F-actin (red) and DAPI (blue); (scale bar: 25µm)

Original Parameters		PCAs
Cell Metadata		
Cell Activation Class		A
Cell Orientation		A
Cell Area		A
Cell Perimetre	A	A
Cell Max Length		A
Cell Min Length		A
Widt/length Ratio		A
Number of Neighbours		B
Distance from closest Neighbour		B
Distance from second closest Neighbour		B
Nuclei Clustering Index	B	B
Nuclei Distance from closest Neighbour		B
Nuclei Distance from second closest Neighbour		B
Nuclear Area		
DNA Intensity		D
Cell Cycle Class		
Cell CDH5 intensity		D
Junctional CDH5 Intensity		D
CDH5 Junction/Cytoplasm Ratio		
Total Number of Junctions	C	
Percent of Stabilised Junctions		
Percent of Linear Junctions		
Percent of Jagged Junctions		
Junction Stability Index		
Cytoplasmic NICD Puncta Intensity		D
Nuclear NICD Puncta Intensity		D
Total NICD Puncta Intensity		
NOTCH Activation Class		
Nuclear NICD Fraction		

Supplementary Table 1: HCIAEC parameters. Highlighted parameters (groups A-D) were used in individual PCAs (prcomp package). For each group the first 3 PCA components were selected (capturing >90% total variance for each group). 12 PCA components were used in tSNE computation (Rtsne package with pca option set to FALSE).

Supplementary Figure 2: A) HUVEC, HAoEC and HPMEC, density distribution of CDH5 intensities. B) HUVEC, HAoEC and HPMEC, density estimate of VEC intensity by cell type.

Supplementary Figure 3: HAoEC, HPMEC and HUVEC distributions of DNA intensities measurements (Coloured by experiment)

Supplementary Figure 4: Schematics representing the logic behind assessment of clonal proliferation by NCI and validation with cell 2^{nd} NB distance. NCI = Nuclei clustering index.

Supplementary Figure 5: A) LM and G2/M cells percentages in tSNE clusters. G2/M cells were highly enriched in clusters 3 and 4 (Nu Clustering < 2), LM cells were enriched in Cluster 2 (Nu Clustering >=2). B) Average LM number per well either in colonies (IC) or isolated (not in colony, NIC), p<0.001 for all IC-NIC comparisons.

Supplementary Figure 6: HUVEC, HAoEC and HPMEC, density distributions of width to length ratio as index of cell elongation. VEGF induced significant reduction of WLR in all cell lines indicating cell elongation.

Supplementary Figure 7: HAoEC, HPMEC and HUVEC Activation classes

Supplementary Figure 8: HUVEC, HAoEC and HPMEC, NOTCH activation comparisons. A) Comparisons between cell types at baseline. B) Comparisons between Cell Types upon treatment with VEGF. * P<0,05, ** p<0.01, *** P<0.001

Supplementary Figure 9: HUVEC, HAoEC and HPMEC, NOTCH Nuclear fraction distribution

Supplementary Figure 10: HUVEC, HAoEC and HPMEC, Total NCID vs Nuclear NICD scatterplot. Coloured according to STB_Index. No correlation between NICD activity and STB index is evident.

Supplementary Figure 11: HUVEC, HAoEC and HPMEC, STB index in the four NOTCH activation classes. Average stability index is similar in all classes.

Supplementary Figure 12: Possible models of cell proliferation. a) Unidirectional model in which only endothelial cell stem cells (ECSC) and endothelial precursor cells (EPC) can proliferate while fully differentiate EC cannot proliferate. b) Cyclic model in which every EC within monolayers remains quiescent until contact inhibited. Local loss of contact in the presence or absence of supplementary stimulation (i.e. VEGF) relieve contact inhibition of proliferation and the EC can re-enter cell cycle as a proliferating EC (PEC).