Unified neural pathways that gate affective pain and multisensory innate threat signals to the amygdala

Sukjae Joshua Kang[1,5], Shijia Liu[1,2,5], Mao Ye[1,5], Dong-Il Kim[1], Jong-Hyun Kim[1], Tae Gyu Oh[3], Jiahang Peng[1, 2], Ronald M. Evans[3], Kuo-Fen Lee[1], Martyn Goulding[4], Sung Han[1,2] *

[1] Peptide Biology Laboratories, The Salk Institute for Biological Studies, La Jolla, CA 92037, USA
[2] Section of Neurobiology, Division of Biological Sciences, University of California, San Diego, La Jolla, CA 92093, USA
[3] Howard Hughes Medical Institute, Gene Expression Laboratories, The Salk Institute for Biological Studies, La Jolla, CA 92037, USA
[4] Molecular Neurobiology Laboratories, The Salk Institute for Biological Studies, La Jolla, CA 92037, USA
[5] These authors contributed equally

*Correspondence to:
Sung Han, Ph.D.
Peptide Biology Laboratories,
Salk Institute for Biological Studies
10010 N. Torrey Pines Rd.
La Jolla, CA 92037, USA
Email: sunghan@salk.edu
Phone: (858) 453-4100, x1856

Abstract
Perception of aversive sensory stimuli such as pain and innate threat cues is essential for animal survival.
The amygdala is critical for aversive sensory perception, and it has been suggested that multiple parallel
pathways independently relay aversive cues from each sensory modality to the amygdala. However, a
convergent pathway that relays multisensory aversive cues to the amygdala has not been identified. Here,
we report that neurons expressing calcitonin gene-related peptide (CGRP) in the parvocellular
subparafasicular thalamic nucleus (SPFp) are necessary and sufficient for affective-motivational pain
perception by forming a spino-thalamo-amygdaloid pain pathway. In addition, we find that this thalamic
CGRP pain pathway, together with well-known parabrachio-amygdaloid CGRP pain pathway, is critical
for the perception of multisensory innate threat cues. The discovery of unified pathways that collectively
gate aversive sensory stimuli from all sensory modalities may provide critical circuit-based insights for
developing therapeutic interventions for affective pain- and innate fear-related disorders.

Introduction

Pain is a complex sensory and emotional experience caused by tissue-damaging noxious stimuli that produce immediate avoidance behavior, as well as long-lasting aversive memories so that future damage can be avoided (Julius and Basbaum, 2001; Melzack and Casey, 1968). Therefore, the perception of pain results in behavioral outcomes similar to those associated with the perception of threats. Indeed, most research on Pavlovian threat learning has used electric foot shock, a painful noxious stimulus, as a threat cue. It has also been suggested that pain and threat perceptions interact with each other (Elman and Borsook, 2018). Individuals with pain asymbolia, who have deficits in perceiving affective-motivational aspects of pain due to damage to limbic structures, show compromised perception of threats (Berthier et al., 1988; Rubins and Friedman, 1948). Alternatively, people with affective pain disorders, such as migraine and fibromyalgia, are often hypersensitive to sensory inputs and perceive normal sensory signals as aversive (Bar-Shalita et al., 2019; López-Solà et al., 2017). Therefore, it is likely that there are unified neural circuits and brain areas that process both pain-causing noxious stimuli and threat-producing aversive sensory cues (Price, 1999).

The amygdala is a key limbic structure that integrates sensory stimuli with an internal state to generate appropriate emotional responses (Janak and Tye, 2015; LeDoux, 2012, 2000). It is activated by aversive sensory stimuli, including noxious stimuli (Ren and Neugebauer, 2010; Simons et al., 2014; Veinante et al., 2013), and lesioning the amygdala greatly attenuates the perception of multimodal sensory threats (Bach et al., 2015; Blanchard and Blanchard, 1972; Dal Monte et al., 2015) and pain (Gao et al., 2004; Helmstetter, 1992; Manning and Mayer, 1995; Tanimoto et al., 2003). Therefore, the amygdala may serve as a pivotal node in integrating and unifying all threat cues from different sensory modalities, including pain-causing noxious stimuli. Recent studies suggest that aversive sensory stimuli from each sensory modality relay threat cues to the distinct brain areas including the hypothalamus and amygdala through parallel non-overlapping pathways. These include somatosensory (Barsy et al., 2020; Choi et al., 2020; Han et al., 2015; Sato et al., 2015), visual (Salay et al., 2018; Wei et al., 2015; Zhou et al., 2019), auditory (Barsy et al., 2020), gustatory (Carter et al., 2013; Kim et al., 2017; Wang et al., 2018) and olfactory (Rosen et al., 2015; Tong et al., 2020). However, little is known about the convergent neural circuits that relay and integrate multimodal aversive sensory signals, including nociceptive stimuli, to the amygdala.

Noxious stimuli from the periphery are relayed to the brain through two ascending pain pathways, the spino-parabrachial pathway and the spino-thalamic pathway (Bushnell et al., 2013). It is a well-established idea that the spino-thalamic pathway is involved in sensory and discriminative pain perception and that the spino-parabrachial pathway is involved in the perception of affective and motivational pain. This is because the former projects to the somatosensory cortex and the latter projects to the amygdala (Basbaum et al., 2009). Among multiple areas of the amygdala, the capsular subdivision of the central nucleus of the amygdala (CeC) is known as the nociceptive amygdala since it is activated by nociceptive stimuli and receives direct input from the parabrachial nucleus (PBN) through the spino-parabrachial pathway (Gauriau and Bernard, 2002; Neugebauer, 2015). Nevertheless, it remains unclear how the nociceptive information is relayed to other areas of the amygdala involved in pain processing, such as the amygdala-striatum transition area (AStr) (Xiu et al., 2014) and the lateral nucleus of the amygdala (LA) (Bernard et al., 1992; Thompson and Neugebauer, 2017). In particular, the LA is critically involved in pain-induced aversive learning (LeDoux, 2007; Ressler and Maren, 2019), but the detailed pain pathway that relays nociceptive information to the LA has not been fully understood.

Although the thalamus has been implicated in sensory and discriminative pain (Dado et al., 1994; Zhang
and Giesler, 2005), some thalamic nuclei, such as the ventromedial posterior thalamus (VMpo) in primates,
or the triangular subdivision of the posterior thalamus (PoT) in rodents, are thought to be involved in
affective and motivational pain perception by relaying nociceptive signals to the limbic areas (Craig et al.,
1994, 2000; Gauriau and Bernard, 2004a; Price, 2002). However, the involvement of these nuclei in
affective and motivational pain perception is still inconclusive (Willis et al., 2002), mainly because the
thalamus has many small, functionally distinct nuclei without clear anatomical boundaries. Therefore, to
understand the roles of the thalamus in affective pain processing, it is critical to identify genetically defined
populations of thalamic neurons that directly convey spinal nociceptive input to the limbic areas, such as
the amygdala.

Calcitonin gene-related peptide (CGRP) is a 37-amino acid neuropeptide produced by peripheral neurons
and mediates vasodilation and nociception (Russell et al., 2014; Russo, 2015). It is also produced in the
brain and plays an essential role in aversive learning and pain perception (Palmiter, 2018; Shinohara et al.,
2017; Yu et al., 2009). CGRP-expressing neurons are highly clustered in two brain areas: the external lateral
subdivision of the PBN (PBel) and the parvocellular subparafascicular nucleus (SPFp) (D'Hanis et al., 2007;
Dobolyi et al., 2005); (Experiment 79587715, Allen Brain Atlas). Previous studies have shown that CGRP
neurons in the PBel (CGRP^{PBel}) are critically involved in transmitting affective pain signals during aversive
learning (Han et al., 2015) and in transmitting visceral aversive signals to the CeA (Chen et al., 2018). On
the other hand, the latter is a relatively unexplored area. The SPFp is an elongated structure that extends
from the anteromedial to posterolateral thalamus (D'Hanis et al., 2007). It has been speculated that CGRP
neurons in the medial part of the SPFp (CGRP^{SPFp}) may play a role in sexual behaviors (Coolen et al., 2003a,
2003b), whereas those in the posterolateral part of the CGRP^{SPFp} may be involved in emotional behaviors,
based on anatomical projections to the amygdala (D'Hanis et al., 2007; LeDoux et al., 1985; Yasui et al.,
1991). The neighboring thalamic nucleus, such as the posterior intralaminar nucleus (PIL), or medial
subdivision of medial geniculate nucleus (MGm) have been implicated to relay noxious stimuli to the
amygdala during aversive learning (Barsy et al., 2020; Shi and Davis, 1999).   However, it has not been
tested whether the CGRP^{SPFp} neurons are involved in conveying pain signals to the amygdala during
affective-motivational pain perception. Furthermore, it is unknown to what extent CGRP^{PBel} and CGRP^{SPFp}
neurons play similar roles in conveying aversive sensory information to the amygdala.

Here, we report that CGRP^{SPFp} neurons receive direct monosynaptic inputs from projection neurons within
the dorsal horn of the spinal cord and project their axons to multiple regions within the amygdala (namely
(the AStr and the LA) and to the posterior insular cortex, but not to the somatosensory cortex. These neurons
are activated by multimodal nociceptive stimuli. Silencing these neurons substantially attenuates affective
and motivational pain perception, and activating these neurons induces aversion and aversive memory.
Furthermore, CGRP^{SPFp} neurons, together with CGRP^{PBel} neurons, are collectively activated by aversive
sensory stimuli from all sensory modalities (visual, auditory, somatosensory, gustatory, and olfactory), and
silencing these neurons attenuates the perception of all aversive sensory stimuli. Taken together, CGRP
neurons within the SPFp and PBel not only form two affective pain pathways, namely the spino-thalamo-
amygdaloid and the spino-parabracho-amygdaloid pathways, but they also convey aversive sensory signals
from all sensory modalities to the amygdala during threat perception.

Results
CGRP^{SPFp} and CGRP^{PBel} neurons connect multiple sensory relay areas to the amygdala
The amygdala is critically involved in the affective-motivational pain perception. However, it is not fully
understood by which the nociceptive information is conveyed to the amygdala. The spinal dorsal horn
projection neurons relay nociceptive signals to the brain (Basbaum et al., 2009; Todd, 2010). One example
is *Tacr1*-expressing neurons in the spinal dorsal horn (Barik et al., 2020; Chiang et al., 2020; Choi et al.,
2020; Deng et al., 2020). To identify direct spino-recipient areas in the brain, we genetically labeled only
the spinal *Tacr1* neurons with tdTomato fluorescent protein by the triple crossing of *Tacr1^{Cre}*, *Cdx2^{FlpO}*, and
Ai65 (Rosa-CAG-FrtSTOPFrt-LoxSTOPLox-tdTomato; *dsTomato*) mice as described previously (Bourane
et al., 2015) (Figure S1A). The tdTomato-expressing cell bodies were only observed in the spinal dorsal
horn (Figure S1A). Fluorescently labeled axonal terminals were observed in multiple brain areas, including
the PBN, SPFp, the posterior complex of the thalamus (Po), the ventral posterolateral nucleus of the
thalamus (VPL), superior colliculus (SC), periaqueductal gray (PAG), dorsal column nuclei (DCN), and
ventrolateral medulla (VLM) (Figure S1B). We then asked which of these areas project to the amygdala.
By searching through the Allen Mouse Connectivity Atlas (http://connectivity.brain-map.org/), we found
that the SPFp and PBel project to the LA and CeA, respectively. Interestingly, CGRP neurons are found in
both the SPFp and PBel, and CGRP^{PBel} neurons are known to play a role in affective pain perception (Han
et al., 2015). Therefore, we sought to dissect and compare the roles of these CGRP circuits in processing
nociceptive sensory information and relaying this information to the amygdala.
To identify regions that lie downstream of CGRP neurons in the SPFp and PBel, we injected Cre-
dependent AAVs encoding EYFP or mCherry into the SPFp and PBel of the same *Calca^{Cre}* mouse that
expresses Cre-recombinase in the CGRP-expressing neurons (*Calca* gene encodes CGRP), respectively
(Figure 1A). Coronal slices around AP -1.1 showed an intermingled green and red expression pattern in the
CeA and LA (Figure S2A). However, posterior slices (AP -1.5) revealed distinct patterns of EYFP and
mCherry in the amygdala. While the CGRP^{SPFp} synaptic terminals were found in the AStr, LA, and medial
amygdala (MEA), the CGRP^{PBel} terminals were most abundant in the CeA, and basomedial amygdala
(BMA) (Figure 1B). CGRP^{SPFp} neurons did not project to the somatosensory cortex, instead they projected
to the auditory cortex and the dorsal regions of the posterior insular cortex (pIC), whereas CGRP^{PBel} neurons
projected to the bed nuclei of the stria terminalis (BNST), ventral posteromedial nucleus of the thalamus
parvicellular part (VPMpc), parasubthalamic nucleus (PSTN), and the ventral portion of the pIC (Figure
S2B).
To identify upstream brain regions that directly project their axons to the CGRP^{SPFp}, or CGRP^{PBel} neurons,
we performed cell-type-specific monosynaptic retrograde tracing using pseudotyped rabies virus (Kim et
al., 2016). We injected AAV8-hSyn-FLEX-TVA-P2A-GFP-2A-oG into the SPFp or PBel of *Calca^{Cre}* mice
and then waited three weeks before injecting EnvA-ΔG-rabies-mCherry into the same region (Figure 1C).
Five days after the rabies injection, mice were sacrificed to detect starter cells in both regions (Figure 1D)
and mCherry-labeled direct upstream neurons in the brain and spinal cord (Figure 1E). Histological analyses
revealed that CGRP^{SPFp} neurons received inputs from superficial and deep layers of the spinal dorsal horn
in multiple spinal cord segments, but most abundantly from the cervical segment (Figures 1F-1H). By
contrast, fewer neurons in the spinal cord project to the CGRP^{PBel} neurons compared to those projecting to
the CGRP^{SPFp} neurons (Figures 1I-1K). Other than the spinal cord, CGRP^{SPFp} neurons received inputs from
sensory relay areas including the SC, inferior colliculus (IC), vestibular nucleus (VN), and trigeminal spinal
nucleus (SpV), as well as other regions such as the hypothalamus and cortex (Figures S3A, S3C, and S3E).
CGRP^{PBel} neurons also received projections from sensory relay areas, including the SC, IC, VN, SpV, and

the nucleus tractus solitarius (NTS), as well as other areas such as the amygdala (in particular the CeA) and
the hypothalamus including the lateral hypothalamus (LHA), zona inserta (ZI), and PSTN (Figures S3B,
S3D, and S3F). Thus, these two populations of CGRP neurons may receive multimodal sensory inputs.
Our results show that both CGRP^{SPFp} and CGRP^{PBel} neurons receive monosynaptic inputs from the spinal
dorsal horn, other sensory-related regions, hypothalamus, and amygdala (CGRP^{PBel} neurons in particular).
However, in terms of output patterns, CGRP^{SPFp} neurons project to the LA and AStr, while CGRP^{PBel}
neurons project to the CeA, thereby forming complementary parallel sensory pathways to the amygdala.
179
CGRP^{SPFp} and CGRP^{PBel} neurons are activated by multimodal nociceptive stimuli
Next, we investigated the response of CGRP^{SPFp} and CGRP^{PBel} neurons to multimodal nociceptive stimuli,
such as mechanical, thermal, and inflammatory stimuli using the fiber photometry *in vivo* calcium imaging
(Figure 2A). AAV-DIO-GCaMP6m was injected into the SPFp or PBel of *Calca^{Cre}* mice, and an optic fiber
(400 μm, 0.37 NA) was implanted above the injection site (Figures 2B and 2C). Previous *in vivo*
electrophysiology studies have shown that nociceptive signals can be conveyed from the spinal cord to the
brain under anesthesia (Gauriau and Bernard, 2004b; Peschanski et al., 1981). Therefore, we performed the
fiber photometry experiments under light anesthesia to remove other emotional and movement artifacts.
Various intensities of mechanical stimuli (0, 50, 100, 200, and 300 g of pressure delivered by a pressure
meter, not von Frey hairs) were applied for 5 seconds to the ipsi- or contra-lateral paws or tail, resulting in
intensity-dependent increases in calcium signals (Area under curve analysis; A.U.C.) in both CGRP^{SPFp}
(Figures 2D-2F) and CGRP^{PBel} neurons (Figures 2G-2I), but there was a significantly decreased calcium
response in the CGRP^{PBel} neurons by 300 g stimulation which created the inverted U-shaped intensity
response curve. Nevertheless, CGRP^{SPFp} and CGRP^{PBel} neurons display different dynamics to noxious
stimuli. CGRP^{PBel} neurons reached maximum response at lower stimulation intensity compared to the
CGRP^{SPFp} neurons (Figures S4A and S4B), but the latter responded faster to the stimuli, which was observed
by greater initial rise slope of calcium peak in the CGRP^{SPFp} neurons compared to the CGRP^{PBel} neurons
(Figure S4C). Although contralateral stimulation evoked greater responses than the ipsilateral stimulation,
both the CGRP^{SPFp} and CGRP^{PBel} neurons were robustly activated by noxious stimuli from both sides,
contrary to the conventional lateralized ascending sensory pain pathways. (Figures S4D-S4G).
Heat stimuli (25, 35, 45, and 55ºC) also induced intensity-dependent calcium increases in both CGRP^{SPFp}
(Figure 2J-2L) and CGRP^{PBel} neurons (Figure 2M-2O). The maximum calcium peak values were higher
overall in CGRP^{PBel} neurons than in CGRP^{SPFp} neurons, both for contralateral (Figure S4H) and ipsilateral
stimulations (Figure S4I). However, the CGRP^{SPFp} neurons again responded faster to 55 ºC stimuli than
CGRP^{PBel} neurons (Figure S4J). Although the difference between ipsilateral and contralateral stimulation
was observed in CGRP^{SPFp} neurons, these neurons were robustly activated by thermal noxious stimuli from
both sides (Figures S4K–S4N). Interestingly, calcium responses induced by mechanical stimuli were much
stronger than those induced by thermal stimuli in CGRP^{SPFp} neurons; the opposite was observed for
CGRP^{PBel} neurons. These results indicate that the CGRP^{SPFp} and CGRP^{PBel} neurons may play different roles
in conveying mechanical and thermal pain.
To assess the effects of inflammatory pain, we injected 10 μl of 4% formalin into the contralateral
forepaw, and calcium activity was recorded with fiber photometry under light anesthesia (Figures 2P-2U).
We observed increases in activity during both the initial acute pain phase (5–10 min; Figures 2Q and 2T)
and the inflammatory phase (15–45 min; Figures 2R and 2U). Activation of these neurons by formalin was
confirmed by Fos immunostaining, which again showed that bilateral CGRP^{SPFp} and CGRP^{PBel} neurons

were activated by unilateral formalin injection with slightly higher response in the contralateral neurons
(Figures S5A-S5G).
Our data indicate that multimodal nociceptive stimuli bilaterally activate CGRP^SPFp and CGRP^PBel
neurons and that these two populations differentially respond to nociceptive inputs of distinct modalities.
Moreover, there is a potential feedback inhibitory circuit in the CGRP^PBel neurons, as we observed activity
decreases in response to 300 g of mechanical pain and during the acute phase of the formalin test.
221
222 Pain-induced synaptic plasticity change in CGRP^SPFp and CGRP^PBel neurons
As we observed that noxious stimuli activate CGRP^SPFp and CGRP^PBel neurons, we hypothesized that pain
might alter the glutamatergic synaptic strength of these neurons. We performed *ex vivo* electrophysiology
to test pain-induced synaptic plasticity changes. To induce pain, we injected 50 µl of 5% formalin into the
upper lip. Twenty-four hours after the injection, we prepared acute brain slices that contained the SPFp
(Figure 3A) or PBel (Figure 3I). We then measured the AMPA/NMDA ratio, an index of glutamatergic
synaptic strength, for both CGRP^SPFp and CGRP^PBel neurons using the whole-cell patch-clamp recording.
For both CGRP^SPFp (Figures 3B and 3C) and CGRP^PBel neurons (Figures 3J and 3K), the AMPA/NMDA
ratio increased (indicating long-term potentiation) in mice treated with formalin compared to controls. No
differences were observed in the paired-pulse ratio (which indicates a presynaptic mechanism) for
CGRP^SPFp neurons (Figures 3D and 3E), but an increase in this ratio was found for CGRP^PBel neurons
(Figures 3L and 3M). To determine whether the AMPA current was affected, we recorded AMPA-mediated
mini EPSCs (Figures 3F and 3N). mEPSC amplitude (Figure 3G), but not frequency (Figure 3H), increased
in CGRP^SPFp neurons of mice subjected to pain compared with controls. For CGRP^PBel neurons, the pain did
not affect mEPSC amplitude (Figure 3O) but decreased frequency (Figure 3P).
These data together suggest that CGRP^SPFp and CGRP^PBel neuronal synapses differentially increase the
strength of glutamatergic signaling in response to pain.
239
240 The transcriptome profiling of CGRP^SPFp and CGRP^PBel neurons
To further investigate whether CGRP^SPFp and CGRP^PBel neurons exhibit specific transcriptomic profiles
associated with the affective pain perception, we conducted cell-type-specific transcriptomic profiling of
CGRP neurons in the SPFp and PBel regions. The *Calca^CreER* mouse line was crossed with the RiboTag
mouse line (Sanz et al., 2009) that has a floxed allele of hemagglutinin (HA)-tagged *Rpl22* gene. As a
result, the HA-tagged ribosomal protein, RPL22 is Cre-dependently expressed in the CGRP-expressing
neurons. After fresh brain tissues containing the SPFp or PBel region were collected and homogenized, the
ribosome-associated transcriptome was captured by immunoprecipitation with anti-HA antibody-
conjugated magnetic beads. Precipitated / unprecipitated total RNAs were sequenced to profile active
transcriptome enriched / deenriched in CGRP^SPFp and CGRP^PBel neurons. RNA sequencing results revealed
that the *Calca* gene that encodes CGRP is highly enriched in both CGRP^SPFp and CGRP^PBel-specific
transcriptome, which served as a positive control. Moreover, genes encoding neuropeptide were specifically
enriched in CGRP^PBel neurons (Figure 4D), and genes encoding markers for inhibitory neurons or glia were
de-enriched in both regions (Figure 4E), confirming that these neurons are glutamatergic neurons. Notably,
several pain-related genes, in particular those encoding membrane proteins, were enriched in both
populations (Figures 4A-4C). Interestingly, genes associated with affective pain disorders, such as *Scn9a*,
and *Faah* for congenital insensitivity to pain (CIP), and *Cacna1a* for migraine (Nassar et al., 2004; van den
Maagdenberg et al., 2004; Cravatt and Lichtman, 2004) are highly enriched in these neurons (Figures 4A-
4C). Expression of the proteins encoded by these genes (Na$_V$1.7, Ca$_V$2.1, and FAAH) in the SPFp and PBel

was confirmed via immunohistochemistry (Figures 4F-4H). These data indicate that both CGRP^{SPFp} and
CGRP^{PBel} neurons express genes involved in pain perception, further supporting our results that these
neurons form ascending pain pathways.

CGRP^{SPFp} and CGRP^{PBel} neurons are activated by multisensory innate threat stimuli

Our retrograde tracing results indicate that both CGRP^{SPFp} and CGRP^{PBel} neurons receive inputs from areas
conveying sensory information from multiple sensory modalities, such as visual stimulus via the SC,
auditory stimulus via the IC, gustatory stimulus via NTS, and nociceptive stimuli via the SC and SpV. We,
therefore, examined whether they are also activated by multisensory innate threat stimuli. We used the fiber
photometry system to measure neural activity in response to five different aversive sensory stimuli. AAV-
DIO-GCaMP7s was injected into the SPFp or PBel of *Calca^{Cre}* mice, and an optic fiber (400 µm, 0.37 NA)
was implanted above the injection site to measure calcium activity of CGRP^{SPFp} or CGRP^{PBel} neurons
(Figures 5A and 5B). A somatosensory stimulus was first tested by applying a foot shock in a cued fear
conditioning test (Figure 5C). We associated a non-aversive low-volume tone (70 dB) with the shock to
minimize the tone's aversive effect (Figure S6A). For both CGRP^{SPFp} and CGRP^{PBel} neurons, immediate
increases in neural activity were detected in response to the 2-s foot shock, but not during habituation or
during the cue test (when the tone was on; Figures 5D–5E). Freezing was observed during the cue test,
indicating that fear memory was formed (Figures S6B and S6C). To assess the auditory threat, an intense
sound (85 dB) was delivered for 2 s (Figure 5F). Time-locked calcium responses were detected at the onset
of an 85-dB intense sound, but not a 70-dB sound for both CGRP^{SPFp} (Figure 5G) and CGRP^{PBel} neurons
(Figure 5H). For an innate visual stimulus, a 2-s looming stimulus was given three times with 10-s intervals.
Both CGRP^{SPFp} (Figure 5J) and CGRP^{PBel} neurons (Figure 5K) displayed an increase of activity in response
to the looming (large disk) compared to the control (small disk) stimulus. As an innate olfactory stimulus
test, we exposed mice to a cotton swab soaked with trimethylthiazoline (TMT; Figure 5L). CGRP^{SPFp}
neurons did not respond to TMT (Figure 5M), whereas CGRP^{PBel} neurons exhibited a slight increase in
activity (Figure 5N). Finally, a gustatory stimulus was administered by exposing mice to quinine (vs. water;
Figure 5O). When overnight water restricted mice licked quinine solution (0.5 mM), CGRP^{SPFp} neurons did
not respond, compared with water controls (Figure 5P), but CGRP^{PBel} neurons exhibited an increase in
calcium activity (Figure 5Q). The calcium peak amplitude analysis shows that the calcium responses
between the CGRP^{SPFp} and CGRP^{PBel} neurons were not significantly different by somatosensory and
auditory stimuli (Figures S7A and S7B). However, the CGRP^{SPFp} neurons showed a greater response to the
visual stimulus compared to the CGRP^{PBel} neurons (Figure S7C). In contrast, the CGRP^{PBel} neurons showed
greater responses to the olfactory and gustatory stimuli compared to the CGRP^{SPFp} neurons (Figures S7D
and S7E).

Our results indicate that the CGRP^{SPFp} and CGRP^{PBel} neurons are both involved in the perception of
innate multisensory threat but respond differently to inputs from distinct modalities. CGRP^{PBel} neurons were
activated by all five aversive sensory stimuli, whereas the CGRP^{SPFp} neurons were only activated by
somatosensory, auditory, and visual aversive stimuli.

Silencing CGRP neurons attenuates responses to multimodal threat stimuli

Our rabies tracing and fiber photometry results imply that CGRP^{SPFp} and CGRP^{PBel} neurons are critically
involved in innate threat perception. Thus, we next tested whether these neurons are necessary for innate
threat perception. We silenced these neurons by bilateral injection of AAV-DIO-TetTox::GFP into the
SPFp or PBel of *Calca^{Cre}* mice and measured behavioral responses to pain stimuli, and multimodal aversive

threat stimuli (Figure 6A). We first performed the formalin assay to test the affective pain perception (Figure 6B). Following injection of 4% formalin into the forepaw, mice in which CGRP[SPFp] or CGRP[PBel] neurons were silenced spent less time licking the injected paw (Figure 6C). In addition, the CGRP[SPFp] silenced group exhibited decreased thermal sensitivity in the 55 °C hot plate test (Figure S8A and S8B), decreased mechanical sensitivity in the electronic von Frey test (Figures S8C and S8D), and decreased freezing in response to the contextual fear conditioning test (Figures S8E and S8F). Interestingly, previous results with CGRP[PBel]-silenced mice by TetTox exhibited no changes in thermal or mechanical thresholds but decreased freezing in the fear conditioning test (Han et al., 2015), suggesting that both CGRP[SPFp] and CGRP[PBel] neurons are necessary for affective pain perception. However, CGRP[SPFp] neurons, not CGRP[PBel] neurons, are also necessary for sensory pain perception. The elevated plus maze (EPM) test shows that silencing the CGRP[SPFp] or CGRP[PBel] neurons decreased anxiety-like behaviors in mice (Figures S8G and S8H).

To test the role of these neurons in multisensory threat perception, the same group of mice were subjected to multiple aversive sensory threat cues, as described in Figure 5. Levels of immediate freezing in response to the aversive somatosensory stimulus (2-s, 0.6 mA electric foot shock) were significantly reduced in both the CGRP[SPFp] and CGRP[PBel] TetTox groups compared to the EYFP control groups (Figure 6D). In the auditory threat test with 85-dB intense sound, EYFP control mice displayed freezing behavior, but freezing levels were reduced in both the CGRP[SPFp] and CGRP[PBel] TetTox groups (Figure 6E). Defensive behaviors (freezing) were also attenuated in response to a looming visual stimulus in the CGRP[SPFp] TetTox group compared with controls, but no difference in freezing was observed between the CGRP[PBel] TetTox group and controls (Figure 6F). Interestingly, CGRP[PBel] neurons were activated by looming (Figure 5K), but their silencing was not enough to attenuate the animals' response to a visual threat, indicating that they play a less significant role in transmitting aversive visual stimulus to the amygdala compared to the CGRP[SPFp] neurons. The aversive olfactory test was performed using a two-chamber system, with one chamber containing water-soaked cotton and the other containing TMT-soaked cotton. Silencing the CGRP[SPFp] neurons did not affect the perception of the aversive olfactory cue, as these mice and EYFP controls both avoided the TMT chamber, while the CGRP[PBel] TetTox group exhibited no aversion to TMT, spending equal amounts of time in the water and TMT chambers (Figure 6G). The gustatory test was performed as a two-bottle choice test between water and quinine solution. The CGRP[SPFp] TetTox group consumed minimal quinine solution (0.5 mM), like controls, whereas the CGRP[PBel] TetTox group showed much less aversion to quinine (Figure 6H).

These results indicate that both the CGRP[SPFp] and CGRP[PBel] neurons are necessary for the perception of innate sensory threat cues, as well as affective pain.

Activating CGRP[SPFp]/CGRP[PBel] to amygdala pathways induces negative valence

Next, we performed optogenetic gain-of-function experiments to test whether activation of these neurons is sufficient to induce negative affect in mice. We bilaterally injected AAV-DIO-ChR2 into the SPFp of *Calca[Cre]* mice and implanted optic fibers (200 μm, NA 0.22) above the injection site (Figure 7A). 20-Hz photo-stimulation of CGRP[SPFp] neurons did not change responses in the hot plate thermal sensitivity test and the electronic von Frey mechanical threshold test for sensory and discriminative pain perception (Figures S8I-S8L). To test whether these neurons encode negative valence, we performed the real-time place aversion (RTPA) test. Optogenetic stimulation is delivered only when the test mouse stays on one side of a two-chamber apparatus (Stamatakis and Stuber, 2012; Figure 7B). Optogenetic activation of the CGRP[SPFp] neurons induced aversion to the photo-stimulated chamber, suggesting that these neurons play a

role in negative emotion or affective-motivational pain (Figures 7B and 7C). Next, we replaced the foot shock with photo-stimulation (20 Hz) as the unconditioned stimulus (US) in the context and cued fear conditioning test. This was to assess whether activation of CGRP^{SPFp} neurons was sufficient to induce fear behaviors. Context-dependent optogenetic conditioning was achieved by 10 mins of photo-stimulation in an open field arena; freezing behavior was then assessed in the same context 24 h after the conditioning (Figure S8M). The ChR2 group exhibited more freezing than the control group, suggesting that CGRP^{SPFp} activation can act as the US (Figure S8N). For cue-dependent optogenetic conditioning, photo-stimulation was associated with a tone as a non-noxious conditioned stimulus (CS+) in a fear conditioning chamber (Figure S8O). Both context and cue tests were performed after the conditioning. The ChR2 group exhibited more freezing in both context (Figure S8P) and cue tests (Figure S8Q). Together with the optogenetic conditioning results of CGRP^{PBel} neurons in the previous result (Han et al., 2015), these results indicate that activation of both CGRP^{SPFp} and CGRP^{PBel} neurons is sufficient to induce negative valence associated with affective-motivational pain perception.

We then sought to characterize the functional downstream of the CGRP^{SPFp} neurons. To examine the functional connectivity of anatomical downstream regions from the CGRP^{SPFp} neurons, we performed *ex vivo* electrophysiology recording. AAV-DIO-ChR2-EYFP was injected into the SPFp of *Calca*^{Cre} mice (Figure S9A). After four weeks of ChR2 expression, we performed whole-cell recordings of neurons from AStr, LA, and pIC to measure optogenetically-evoked excitatory/inhibitory postsynaptic currents (EPSC/IPSC; Figure S9B). We found that CGRP^{SPFp} neurons form functional glutamatergic synapses with neurons within AStr, LA (Figure S9C), and pIC (data not shown). Moreover, the onset of IPSCs lagged 4–5 ms compared to the onset of EPSCs, indicating a feed-forward inhibition circuit. The number of cells that had both EPSCs and IPSCs, EPSCs only, IPSCs only, and non-responsive were also counted (Figures S9D-S9L). To investigate whether these connections form functional circuits that encode negative valence, we optogenetically stimulated axonal terminals from the CGRP^{SPFp} neurons and performed behavioral tests. AAV-DIO-ChR2-EYFP was injected into the SPFp of the Calca^{Cre} mice, and optical fibers were implanted into the postsynaptic areas of CGRP^{SPFp} neurons, namely the LA, AStr, and pIC (Figure 7D). Optogenetic activation of each of these three projections induced aversion in the RTPA experiment (Figures 7E and S9M), as observed in direct photo-stimulation of CGRP^{SPFp} cell bodies (Figure 7C). Cue-dependent optogenetic conditioning of the downstream circuits was then performed and only the CGRP$^{SPFp \to LA}$ circuit caused significant freezing in the context test (Figure 7F). The other two projections only showed a trend (Figure S9N). Increased freezing was observed in the cue test for all three projections (Figures 7G and S9O), but the most prominent effect was observed with the CGRP$^{SPFp \to LA}$ circuit.

The same behavioral experiments were performed as above with the CGRP^{PBel} neurons to compare their role in encoding negative valence with the CGRP^{SPFp} neurons. First, photo-stimulation of the CGRP^{PBel} neuronal cell body (Figure 7H) induced aversion during the RTPA test (Figures 7I and 7J). Then, we investigated the CGRP$^{PBel \to CeA}$ circuit as in Figure 7D-7H by optogenetic terminal stimulation (Figure 7K). Photo-stimulation of CGRP$^{PBel \to CeA}$ terminals induced aversion in the RTPA test (Figure 7L), and freezing in the optogenetic conditioning context, and cue tests (Figures 7M and 7N).

These results satisfy the idea that CGRP$^{SPFp \to LA}$ and CGRP$^{PBel \to CeA}$ circuits induce negative valence associated with either affective pain or innate sensory threat cues.

Discussion

We report a genetically defined population of neurons that express the neuropeptide CGRP in the SPFp and PBel mediate perception of affective pain and innate sensory threat cues by relaying aversive sensory signals from the spinal cord and other sensory relay areas to the amygdala. These analyses provide the first evidence of convergent innate threat pathways that relay multisensory aversive sensory cues to the amygdala.

CGRP neurons convey affective pain signals to the amygdala

Perception of pain protects us from physical harm by locating the source of a harmful stimulus. Painful experiences also elicit emotional and motivational responses, which help us remember these events and avoid similar stimuli in the future (Yeh et al., 2018). Thus, pain is not just a simple sensory process but also a complex cognitive process that generates sensory and emotional responses. This unique aspect of pain gives rise to the concept of two aspects of pain: sensory-discriminative and affective-motivational (Auvray et al., 2010; Melzack and Casey, 1968). It is thought that the sensory-discriminative aspect of pain is processed within the sensory cortex via the spino-thalamic tract (STT), and the affective-motivational aspect of pain is processed within the amygdala via the spino-parabrachial tract (SPT). Indeed, previous studies have shown that the PBN-to-CeA circuit is critical for affective-motivational pain perception (Han et al., 2015; Sato et al., 2015). However, it has also been suggested that the thalamus is actively involved in affective-motivational pain perception (Craig, 2003; Willis et al., 2002). Importantly, posterior regions of the thalamus (e.g., the VMpo in primates and humans, as well as the SPFp and PoT in rodents) are anatomically connected to limbic areas (Craig, 1998; Gauriau and Bernard, 2004a) and activated by noxious stimuli (Craig et al., 1994; Peschanski et al., 1981). Therefore, the thalamus likely also plays a critical role in affective-motivational pain perception. Our results demonstrate that a genetically defined population of neurons expressing the neuropeptide CGRP within the SPFp receives monosynaptic inputs from projection neurons within the spinal dorsal horn. They then project to specific nuclei within the amygdala, namely the AStr and LA (Figure 1). Multimodal nociceptive stimuli activate these neurons in an intensity-dependent manner in anesthetized mice (Figure 2). Inactivating these neurons attenuates the perception of affective-motivational pain, and pain signals induces the changes in synaptic plasticity of these neurons (Figure 3). These data indicate that CGRP^{SPFp} neurons create the spino-thalamo-amygdaloid affective pain pathway.

Unlike the STT, the SPT has been well-characterized as an affective-motivational pain pathway. Recent studies have shown that the lateral PBN receives direct nociceptive inputs from projection neurons within the spinal dorsal horn (Barik et al., 2020; Chiang et al., 2020; Choi et al., 2020; Deng et al., 2020). The dorsolateral PBN (PBdl) predominantly receives nociceptive inputs from the spinal cord and then projects to multiple limbic structures, such as the PAG, VMH, and ILN, thereby producing emotional and physiological changes in response to pain signals (Chiang et al., 2020; Deng et al., 2020). Although the PBdl does not directly project to the amygdala, it indirectly sends pain signals to the CeA through the PBel (Deng et al., 2020). In particular, CGRP^{PBel} neurons are critical for relaying aversive unconditioned stimuli to the CeA during aversive fear learning (Han et al., 2015). It has been shown that the dynorphin neurons in the PBdl project to the PBel (Chiang et al., 2020), and a recent study has shown that spinal projection neurons that express GPR83 directly innervate CGRP^{PBel} neurons to relay noxious signals (Choi et al., 2020). Therefore, it is clear that CGRP^{PBel} neurons relay nociceptive information from the PBdl and the spinal cord to the CeA. Our retrograde tracing results show that CGRP^{PBel} neurons receive less direct inputs from the spinal cord than the CGRP^{SPFp} neurons (Figure 1), and these neurons are activated slower than the CGRP^{SPFp}

neurons (Figure S4C and S4J), implying that the CGRP^{PBel} neurons may receive nociceptive inputs more
predominantly through the PBdl than through directly from the spinal cord. The CGRP^{PBel} neurons also
exhibit intensity-dependent activation by multimodal nociceptive stimuli in anesthetized mice (Figure 2)
and induce changes in synaptic plasticity by pain signals (Figure 3). Therefore, together with previous
studies, our results reaffirm that CGRP^{PBel} neurons comprise the spino-parabrachio-amygdaloid pain
pathway.
437
Comparing the CGRP^{SPFp} and CGRP^{PBel} neuronal activity in response to multimodal nociceptive stimuli at
various intensities in anesthetized mice provides us with novel insights into understanding central affective
pain pathways. First, unlike sensory pain signals that are conveyed to the contralateral side of the
somatosensory cortex, these neurons are activated by both contralateral and ipsilateral noxious stimuli
indicating that the affective pain pathway may not be strictly lateralized (Figure S4). Second, the CGRP^{SPFp}
neurons more robustly respond to the mechanical stimulus, whereas the CGRP^{PBel} neurons respond more
robustly to the thermal stimulus suggesting that these two parallel pathways may convey different
modalities of nociceptive information (Figure 2). Lastly, the CGRP^{PBel} neurons are activated at lower
stimulus intensity but respond slowly to nociceptive signals compared with the CGRP^{SPFp} neurons (Figures
2 and S4). Therefore, our results demonstrate that CGRP-expressing neurons in two brain areas (the SPFp
and PBel) play complementary roles in relaying multimodal nociceptive signals from the spinal cord to the
amygdala through two parallel ascending pain pathways, which are critical for affective-motivational pain
perception.
451

CGRP neurons convey multisensory innate threat cues to the amygdala
Both pain and innate sensory threats motivate animals to execute immediate avoidance behaviors to escape
the threatening or tissue-damaging situation and produce long-lasting aversive memories. Indeed,
Pavlovian threat conditioning uses noxious electric foot shock, an acute painful stimulus that motivates the
animals to create aversive memory (LeDoux, 2012; Maren, 2001). However, the neural circuit mechanisms
by which noxious information is conveyed to the amygdala during aversive learning is not fully understood.
Recent advances in the neural circuit-based understanding of innate predator threat perception suggest that
innate threat cues from each sensory modality are conveyed to discrete brain areas, such as the amygdala
and hypothalamus through parallel pathways (Canteras, 2002; Gross and Canteras, 2012; Kunwar et al.,
2015; Silva et al., 2013), which do not overlap with the unconditioned stimulus (affective pain) pathway in
Pavlovian threat learning (Silva et al., 2016). However, it is beneficial for animals to use unified neural
circuits that encode multimodal aversive sensory signals including pain because animals search for and
detect imminent threats using multiple sensory modalities simultaneously. Moreover, previous clinical and
animal studies have shown that pain and threat perceptions interact with each other (Berthier et al., 1988;
Crook et al., 2014; Elman and Borsook, 2018; Lister et al., 2020). Therefore, it is plausible that there is a
unified mechanism that conveys multisensory threat cues from multiple sensory modalities to the amygdala.
Our results demonstrate that CGRP-expressing neurons in the PBel and the SPFp not only relay nociceptive
stimuli to the amygdala during aversive learning, but also convey innate sensory threat cues from various
sensory modalities (Figures 5 and 6). The CGRP^{SPFp} neurons relay aversive sensory cues from the
somatosensory, visual, and auditory modalities to the LA, AStr, and pIC. By contrast, CGRP^{PBel} neurons
relay aversive cues from all sensory modalities (somatosensory, visual, auditory, olfactory, and gustatory)
(Figures 5, 6, and S3). In addition, previous studies have shown that CGRP^{PBel} neurons are activated by
hypercapnic conditions (high CO_2 levels) (Kaur et al., 2017; Yokota et al., 2015), and aversive visceral

cues, such as lithium chloride and lipopolysaccharide (Carter et al., 2013; Paues et al., 2001). Therefore, it
is tempting to speculate that CGRP[SPFp] neurons relay exteroceptive threat signals, whereas CGRP[PBel]
neurons relay both exteroceptive and interoceptive threat cues to the amygdala.

Consistent with a previous report (Han et al., 2015), our data provide strong evidence that the CGRP[PBel]
neurons mediate aversive learning via their projection to the CeA (Figure 7K-7N). Our results seem to be
contradictory to a recent study by Bowen et al., (2020), which argued that optogenetic stimulation of
CGRP[PBel] axon terminals in the VPMpc, instead of the CeA, evokes strong aversive memory. This
discrepancy can be explained by the differences in optogenetic stimulation protocols. Whereas we used a
40 Hz light stimulation for 10 s as described (Han et al., 2015), Bowen et al. used a 30 Hz light train for
only 2 s. Moreover, it is worth noting that axonal bundles from all glutamatergic neurons in the PBel that
project to forebrain regions, including the CeA, BNST, and PSTN pass through the VPMpc (Huang et al.,
2020), which suggests that optogenetic stimulation within the VPMpc activates both CGRP[PBel] axonal
terminals and axon bundles passing through this area. Therefore, an alternative explanation of their
observation is that concurrent stimulation of all downstream areas produces stronger aversive memory than
stimulating CeA alone.

Together, our results demonstrate that the US pathways for relaying electric foot shock cues during the
Pavlovian fear learning also collectively relay innate threat cues from multiple sensory modalities.

Pain-threat interactions in affective pain disorders
Our results show that the affective-motivational pain and multisensory threat stimuli arrive in the amygdala
via the same neural pathways. Interestingly, pain-threat interactions have been reported in many human
clinical cases. People with pain asymbolia, caused by damage to limbic brain areas, have the normal sensory
perception of noxious stimuli, but they have impaired affective pain perception (Berthier et al., 1988;
Rubins and Friedman, 1948). Interestingly, pain asymbolia patients often display deficits in perceiving
general threats (Klein, 2015; Price, 2000), indicating that the perception of affective pain and other sensory
threat cues share the same neural substrate. People with congenital insensitivity to pain (CIP) are insensitive
to all sensory and affective components of pain, but they also display profound deficits in general threat
perception, which is the primary cause of their short life expectancy (McMurray, 1955; Nagasako et al.,
2003). CIP is caused by loss-of-function mutations in genes critical for pain transmission, such as *Scn9a*
(Dabby, 2012; Fischer and Waxman, 2010; Lampert et al., 2010) and *Faah* (Drissi et al., 2020). *Scn9a*
encodes voltage-gated sodium channel type 7 (Na$_V$1.7), and *Faah* encodes fatty acid amide hydrolase, both
of which are critical for pain transmission in the spinal cord (Cajanus et al., 2016; Kim et al., 2006;
Nantermet and Henze, 2011; Nassar et al., 2004). However, the functional loss of these genes in the spinal
neurons cannot explain the insensitivity to general threats exhibited by CIP patients. Surprisingly, our cell
type-specific transcriptome analysis revealed that *Scn9a* and *Faah* transcripts are highly enriched in both
CGRP[PBel] and CGRP[SPFp] neurons (Figure 4). Thus, mutations in these genes may prevent these neurons
from relaying sensory threat signals to the amygdala, thereby causing insensitivity to general threats in CIP.
This speculation should be addressed by testing the causal relationship between mutations in these genes
in CGRP neurons and threat perception.

Opposite clinical cases also exist. People with affective pain disorders, such as migraine, and fibromyalgia
experience chronic pain and suffer from hypersensitivity to normal sensory stimuli (Demarquay and

Mauguière, 2016; Harriott and Schwedt, 2014; Harte et al., 2016; López-Solà et al., 2017). Further, normal
sensory stimuli often trigger or aggravate their pain symptoms (Bar-Shalita and Cermak, 2020; Bar-Shalita
et al., 2019). Surprisingly, the gene *Cacna1a*, which has been linked to migraines, is highly enriched in
both CGRP[PBel] and CGRP[SPFp] neurons (Figure 4). Further studies should address the causal relationship
between loss of *Cacna1a* function in CGRP neurons and sensory hypersensitivity in migraine. In addition,
CGRP signaling is a proven therapeutic target for treating migraine (Ashina, 2020). Therefore, we speculate
that the CGRP-expressing neurons characterized in this study may serve as the functional substrate for
sensory hypersensitivity in migraine. Identifying membrane proteins commonly expressed in both the
CGRP[PBel] and CGRP[SPFp] neurons (Figure 4C) may provide potential candidates of therapeutic targets for
treating affective pain-, and innate threat-related disorders, such as migraine, fibromyalgia, phobias, panic
disorder, and post-traumatic stress disorder. Indeed, the CGRP receptor antagonists or neutralizing
monoclonal antibodies are promising drugs for treating migraine, but they may also serve as potential
therapeutic interventions for treating the disorders mentioned above.
532
Conclusion
Our findings demonstrate that the CGRP[SPFp] neurons create a novel spino-thalamo-amygdaloid affective
pain pathway and, together with the previously characterized CGRP[PBel] neurons, serve as complementary
parallel pathways for conveying the unconditioned stimulus during Pavlovian threat learning, as well as
multisensory innate threat cues from all sensory modalities to the amygdala (Figure S10). The discovery of
a unified threat perception system that conveys multimodal interoceptive and exteroceptive aversive
sensory stimuli greatly enhances our understanding of the neural mechanisms of innate threat perception.
These insights also provide opportunities to discover novel targets for developing therapeutic interventions
against affective pain- and innate threat-related disorders.
542

Acknowledgments

We thank Dr. D. O'Keefe, Ms. C. Jia, and Han lab members for critical discussions during manuscript preparation. S.H. is supported by 1R01MH116203 from NIMH and the Bridge to Independence award from the Simons Foundation Autism Research Initiative (SFARI #388708). S.L. is supported by the Salk Women & Science Special Award, the Mary K. Chapman Foundation, and the Jesse & Caryl Philips Foundation.

Author Contributions

S.H. conceived of the idea and secured funding. S.H., S.J.K., S.L., and M.Y. designed and performed the experiments. S.H. and S.J.K wrote the manuscript. S.J.K. performed most of the CGRP[SPFp] experiments. S.L., and S.J.K. performed CGRP[PBel] experiments. M.Y. performed electrophysiology and CGRP[PBel] RTPA. D.I.K., and J.H.K. performed RiboTag experiments, and T.G.O analyzed it. R.M.E. provided resources for Ribotag analysis. J.P. performed spinal projection histology experiments. M.G. provided *Cdx2[FlpO]* mouse line. K.F.L. provided resources for microscopy.

Declaration of Interests

The authors declare no competing interests.

References

Ashina, M. (2020). Migraine. N. Engl. J. Med. *383*, 1866–1876.

Auvray, M., Myin, E., and Spence, C. (2010). The sensory-discriminative and affective-motivational aspects of pain. Neurosci. Biobehav. Rev. *34*, 214–223.

Bach, D.R., Hurlemann, R., and Dolan, R.J. (2015). Impaired threat prioritisation after selective bilateral amygdala lesions. Cortex *63*, 206–213.

Barik, A., Sathyamurthy, A., Thompson, J., Seltzer, M., Levine, A., and Chesler, A. (2020). A spinoparabrachial circuit defined by Tacr1 expression drives pain. BioRxiv 2020.07.15.205484.

Bar-Shalita, T., and Cermak, S.A. (2020). Multi-sensory Responsiveness and Personality Traits Predict Daily Pain Sensitivity. Front. Integr. Neurosci. *13*, 77.

Bar-Shalita, T., Granovsky, Y., Parush, S., and Weissman-Fogel, I. (2019). Sensory Modulation Disorder (SMD) and Pain: A New Perspective. Front. Integr. Neurosci. *13*.

Barsy, B., Kocsis, K., Magyar, A., Babiczky, Á., Szabó, M., Veres, J.M., Hillier, D., Ulbert, I., Yizhar, O., and Mátyás, F. (2020). Associative and plastic thalamic signaling to the lateral amygdala controls fear behavior. Nat. Neurosci. *23*, 625–637.

Basbaum, A.I., Bautista, D.M., Scherrer, G., and Julius, D. (2009). Cellular and Molecular Mechanisms of Pain. Cell *139*, 267–284.

Bernard, J.F., Huang, G.F., and Besson, J.M. (1992). Nucleus centralis of the amygdala and the globus pallidus ventralis: electrophysiological evidence for an involvement in pain processes. J. Neurophysiol. *68*, 551–569.

Berthier, M., Starkstein, S., and Leiguarda, R. (1988). Asymbolia for pain: A sensory-limbic disconnection syndrome. Ann. Neurol. *24*, 41–49.

Blanchard, D.C., and Blanchard, R.J. (1972). Innate and conditioned reactions to threat in rats with amygdaloid lesions. J. Comp. Physiol. Psychol. *81*, 281–290.

Bourane, S., Grossmann, K.S., Britz, O., Dalet, A., Del Barrio, M.G., Stam, F.J., Garcia-Campmany, L., Koch, S., and Goulding, M. (2015). Identification of a Spinal Circuit for Light Touch and Fine Motor Control. Cell *160*, 503–515.

Bowen, A.J., Chen, J.Y., Huang, Y.W., Baertsch, N.A., Park, S., and Palmiter, R.D. (2020). Dissociable control of unconditioned responses and associative fear learning by parabrachial CGRP neurons. ELife *9*, e59799.

Bushnell, M.C., Čeko, M., and Low, L.A. (2013). Cognitive and emotional control of pain and its disruption in chronic pain. Nat. Rev. Neurosci. *14*, 502–511.

Cajanus, K., Holmström, E.J., Wessman, M., Anttila, V., Kaunisto, M.A., and Kalso, E. (2016).
Effect of endocannabinoid degradation on pain: role of: FAAH: polymorphisms in experimental
and postoperative pain in women treated for breast cancer. PAIN *157*, 361–369.

Canteras, N.S. (2002). The medial hypothalamic defensive system: Hodological organization and
functional implications. Pharmacol. Biochem. Behav. *71*, 481–491.

Carter, M.E., Soden, M.E., Zweifel, L.S., and Palmiter, R.D. (2013). Genetic identification of a
neural circuit that suppresses appetite. Nature *advance online publication*.

Chen, J.Y., Campos, C.A., Jarvie, B.C., and Palmiter, R.D. (2018). Parabrachial CGRP Neurons
Establish and Sustain Aversive Taste Memories. Neuron *100*, 891-899.e5.

Chiang, M.C., Nguyen, E.K., Canto-Bustos, M., Papale, A.E., Oswald, A.-M.M., and Ross, S.E.
(2020). Divergent Neural Pathways Emanating from the Lateral Parabrachial Nucleus Mediate
Distinct Components of the Pain Response. Neuron *106*, 927-939.e5.

Choi, S., Hachisuka, J., and Ginty, D. (2020). Parallel ascending spinal pathways for affective
touch and pain. Nature *In press*.

Coolen, L.M., Veening, J.G., Wells, A.B., and Shipley, M.T. (2003a). Afferent connections of the
parvocellular subparafascicular thalamic nucleus in the rat: Evidence for functional subdivisions.
J. Comp. Neurol. *463*, 132–156.

Coolen, L.M., Veening, J.G., Petersen, D.W., and Shipley, M.T. (2003b). Parvocellular
subparafascicular thalamic nucleus in the rat: Anatomical and functional compartmentalization. J.
Comp. Neurol. *463*, 117–131.

Craig, A.D. (1998). A new version of the thalamic disinhibition hypothesis of central pain. Pain
Forum *7*, 1–14.

Craig, A.D. (Bud) (2003). PAIN MECHANISMS: Labeled Lines Versus Convergence in Central
Processing. Annu. Rev. Neurosci. *26*, 1–30.

Craig, A.D., Bushnell, M.C., Zhang, E.-T., and Blomqvist, A. (1994). A thalamic nucleus specific
for pain and temperature sensation. Nature *372*, 770–773.

Craig, A.D., Chen, K., Bandy, D., and Reiman, E.M. (2000). Thermosensory activation of insular
cortex. Nat. Neurosci. *3*, 184–190.

Cravatt, B.F., and Lichtman, A.H. (2004). The endogenous cannabinoid system and its role in
nociceptive behavior. J. Neurobiol. *61*, 149–160.

Crook, R.J., Dickson, K., Hanlon, R.T., and Walters, E.T. (2014). Nociceptive Sensitization
Reduces Predation Risk. Curr. Biol. *24*, 1121–1125.

Dabby, R. (2012). Pain Disorders and Erythromelalgia Caused by Voltage-Gated Sodium Channel
Mutations. Curr. Neurol. Neurosci. Rep. *12*, 76–83.

Dado, R.J., Katter, J.T., and Giesler, G.J. (1994). Spinothalamic and spinohypothalamic tract
neurons in the cervical enlargement of rats. II. Responses to innocuous and noxious mechanical
and thermal stimuli. J. Neurophysiol. *71*, 981–1002.

Dal Monte, O., Costa, V.D., Noble, P.L., Murray, E.A., and Averbeck, B.B. (2015). Amygdala
lesions in rhesus macaques decrease attention to threat. Nat. Commun. *6*, 10161.

Demarquay, G., and Mauguière, F. (2016). Central Nervous System Underpinnings of Sensory
Hypersensitivity in Migraine: Insights from Neuroimaging and Electrophysiological Studies.
Headache J. Head Face Pain *56*, 1418–1438.

Deng, J., Zhou, H., Lin, J.-K., Shen, Z.-X., Chen, W.-Z., Wang, L.-H., Li, Q., Mu, D., Wei, Y.-C.,
Xu, X.-H., et al. (2020). The Parabrachial Nucleus Directly Channels Spinal Nociceptive Signals
to the Intralaminar Thalamic Nuclei, but Not the Amygdala. Neuron *107*, 923.

D'Hanis, W., Linke, R., and Yilmazer-Hanke, D. m. (2007). Topography of thalamic and
parabrachial calcitonin gene-related peptide (CGRP) immunoreactive neurons projecting to
subnuclei of the amygdala and extended amygdala. J. Comp. Neurol. *505*, 268–291.

Dobolyi, A., Irwin, S., Makara, G., Usdin, T.B., and Palkovits, M. (2005). Calcitonin gene-related
peptide-containing pathways in the rat forebrain. J. Comp. Neurol. *489*, 92–119.

Drissi, I., Woods, W.A., and Woods, C.G. (2020). Understanding the genetic basis of congenital
insensitivity to pain. Br. Med. Bull. *133*, 65–78.

Elman, I., and Borsook, D. (2018). Threat Response System: Parallel Brain Processes in Pain vis-
à-vis Fear and Anxiety. Front. Psychiatry *9*.

Fischer, T.Z., and Waxman, S.G. (2010). Familial pain syndromes from mutations of the Nav1.7
sodium channel. Ann. N. Y. Acad. Sci. *1184*, 196–207.

Gao, Y.-J., Ren, W.-H., Zhang, Y.-Q., and Zhao, Z.-Q. (2004). Contributions of the anterior
cingulate cortex and amygdala to pain- and fear-conditioned place avoidance in rats: Pain *110*,
343–353.

Gauriau, C., and Bernard, J.-F. (2002). Pain pathways and parabrachial circuits in the rat. Exp.
Physiol. *87*, 251–258.

Gauriau, C., and Bernard, J.-F. (2004a). Posterior Triangular Thalamic Neurons Convey
Nociceptive Messages to the Secondary Somatosensory and Insular Cortices in the Rat. J. Neurosci.
24, 752–761.

Gauriau, C., and Bernard, J.-F. (2004b). A comparative reappraisal of projections from the
superficial laminae of the dorsal horn in the rat: The forebrain. J. Comp. Neurol. *468*, 24–56.

Gross, C.T., and Canteras, N.S. (2012). The many paths to fear. Nat. Rev. Neurosci. *13*, 651–658.

Han, S., Soleiman, M.T., Soden, M.E., Zweifel, L.S., and Palmiter, R.D. (2015). Elucidating an Affective Pain Circuit that Creates a Threat Memory. Cell *162*, 363–374.

Harriott, A.M., and Schwedt, T.J. (2014). Migraine is Associated With Altered Processing of Sensory Stimuli. Curr. Pain Headache Rep. *18*, 458.

Harte, S.E., Ichesco, E., Hampson, J.P., Peltier, S.J., Schmidt-Wilcke, T., Clauw, D.J., and Harris, R.E. (2016). Pharmacologic attenuation of cross-modal sensory augmentation within the chronic pain insula: PAIN *157*, 1933–1945.

Helmstetter, F.J. (1992). The Amygdala Is Essential for the Expression of Conditional Hypoalgesia. Behav. Neurosci. *106*, 518–528.

Huang, D., Grady, F.S., Peltekian, L., and Geerling, J.C. (2020). Efferent projections of VGLUT2 , FOXP2 , and PDYN parabrachial neurons in mice. J. Comp. Neurol. cne.24975.

Janak, P.H., and Tye, K.M. (2015). From circuits to behaviour in the amygdala. Nature *517*, 284–292.

Julius, D., and Basbaum, A.I. (2001). Molecular mechanisms of nociception. Nature *413*, 203–210.

Kaur, S., Wang, J.L., Ferrari, L., Thankachan, S., Kroeger, D., Venner, A., Lazarus, M., Wellman, A., Arrigoni, E., Fuller, P.M., et al. (2017). A Genetically Defined Circuit for Arousal from Sleep during Hypercapnia. Neuron *96*, 1153-1167.e5.

Kim, E.J., Jacobs, M.W., Ito-Cole, T., and Callaway, E.M. (2016). Improved Monosynaptic Neural Circuit Tracing Using Engineered Rabies Virus Glycoproteins. Cell Rep. *15*, 692–699.

Kim, H., Mittal, D.P., Iadarola, M.J., and Dionne, R.A. (2006). Genetic predictors for acute experimental cold and heat pain sensitivity in humans. J. Med. Genet. *43*, e40–e40.

Kim, J., Zhang, X., Muralidhar, S., LeBlanc, S.A., and Tonegawa, S. (2017). Basolateral to Central Amygdala Neural Circuits for Appetitive Behaviors. Neuron *93*, 1464-1479.e5.

Klein, C. (2015). What Pain Asymbolia Really Shows. Mind *124*, 493–516.

Kunwar, P.S., Zelikowsky, M., Remedios, R., Cai, H., Yilmaz, M., Meister, M., and Anderson, D.J. (2015). Ventromedial hypothalamic neurons control a defensive emotion state. ELife *4*, e06633.

Lampert, A., O'Reilly, A.O., Reeh, P., and Leffler, A. (2010). Sodium channelopathies and pain. Pflüg. Arch. - Eur. J. Physiol. *460*, 249–263.

LeDoux, J. (2007). The amygdala. Curr. Biol. *17*, R868–R874.

LeDoux, J. (2012). Rethinking the Emotional Brain. Neuron *73*, 653–676.

LeDoux, J.E. (2000). Emotion Circuits in the Brain. Annu. Rev. Neurosci. *23*, 155–184.

LeDoux, J.E., Ruggiero, D.A., and Reis, D.J. (1985). Projections to the subcortical forebrain from
anatomically defined regions of the medial geniculate body in the rat. J. Comp. Neurol. *242*, 182–
213.

Lister, K.C., Bouchard, S.M., Markova, T., Aternali, A., Denecli, P., Pimentel, S.D., Majeed, M.,
Austin, J.-S., de C. Williams, A.C., and Mogil, J.S. (2020). Chronic pain produces hypervigilance
to predator odor in mice. Curr. Biol. *30*, R866–R867.

López-Solà, M., Woo, C.-W., Pujol, J., Deus, J., Harrison, B.J., Monfort, J., and Wager, T.D.
(2017). Towards a neurophysiological signature for fibromyalgia. PAIN *158*, 34–47.

Manning, B.H., and Mayer, D.J. (1995). The central nucleus of the amygdala contributes to the
production of morphine antinociception in the formalin test. PAIN® *63*, 141–152.

Maren, S. (2001). Neurobiology of Pavlovian Fear Conditioning. Annu. Rev. Neurosci. *24*, 897–
931.

McMurray, G.A. (1955). Congenital insensitivity to pain and its implications for motivational
theory. Can. J. Psychol. Can. Psychol. *9*, 121–131.

Melzack, R., and Casey, K. (1968). Sensory, Motivational, and Central Control Determinants of
Pain. In Skin Senses, pp. 423–439.

Nagasako, E.M., Oaklander, A.L., and Dworkin, R.H. (2003). Congenital insensitivity to pain: an
update. PAIN *101*, 213–219.

Nantermet, P.G., and Henze, D.A. (2011). Recent Advances Toward Pain Therapeutics. In Annual
Reports in Medicinal Chemistry, (Elsevier), pp. 19–32.

Nassar, M.A., Stirling, L.C., Forlani, G., Baker, M.D., Matthews, E.A., Dickenson, A.H., and
Wood, J.N. (2004). Nociceptor-specific gene deletion reveals a major role for Nav1.7 (PN1) in
acute and inflammatory pain. Proc. Natl. Acad. Sci. *101*, 12706–12711.

Neugebauer, V. (2015). Amygdala Pain Mechanisms. In Pain Control, (Springer, Berlin,
Heidelberg), pp. 261–284.

Palmiter, R.D. (2018). The Parabrachial Nucleus: CGRP Neurons Function as a General Alarm.
Trends Neurosci. *41*, 280–293.

Paues, J., Engblom, D., Mackerlova, L., Ericsson-Dahlstrand, A., and Blomqvist, A. (2001).
Feeding-related immune responsive brain stem neurons: association with CGRP. NeuroReport *12*,
2399–2403.

Peschanski, M., Guilbaud, G., and Gautron, M. (1981). Posterior intralaminar region in rat:
Neuronal responses to noxious and nonnoxious cutaneous stimuli. Exp. Neurol. *72*, 226–238.

Price, D.D. (1999). Multisensory integration in pain and consciousness. Pain Forum *8*, 130–132.

Price, D.D. (2000). Psychological and Neural Mechanisms of the Affective Dimension of Pain.
Science *288*, 1769–1772.

Price, D.D. (2002). Central Neural Mechanisms that Interrelate Sensory and Affective Dimensions
of Pain. Mol. Interv. *2*, 392.

Ren, W., and Neugebauer, V. (2010). Pain-related increase of excitatory transmission and decrease
of inhibitory transmission in the central nucleus of the amygdala are mediated by mGluR1. Mol.
Pain *6*, 93.

Ressler, R.L., and Maren, S. (2019). Synaptic encoding of fear memories in the amygdala. Curr.
Opin. Neurobiol. *54*, 54–59.

Rosen, J.B., Asok, A., and Chakraborty, T. (2015). The smell of fear: innate threat of 2,5-dihydro-
2,4,5-trimethylthiazoline, a single molecule component of a predator odor. Front. Neurosci. *9*.

Rubins, J.L., and Friedman, E.D. (1948). ASYMBOLIA FOR PAIN. Arch. Neurol. Psychiatry *60*,
554–573.

Russell, F.A., King, R., Smillie, S.-J., Kodji, X., and Brain, S.D. (2014). Calcitonin Gene-Related
Peptide: Physiology and Pathophysiology. Physiol. Rev. *94*, 1099–1142.

Russo, A.F. (2015). Calcitonin Gene-Related Peptide (CGRP): A New Target for Migraine. Annu.
Rev. Pharmacol. Toxicol. *55*, 533–552.

Salay, L.D., Ishiko, N., and Huberman, A.D. (2018). A midline thalamic circuit determines
reactions to visual threat. Nature *557*, 183.

Sanz, E., Yang, L., Su, T., Morris, D.R., McKnight, G.S., and Amieux, P.S. (2009). Cell-type-
specific isolation of ribosome-associated mRNA from complex tissues. Proc. Natl. Acad. Sci. *106*,
13939–13944.

Sato, M., Ito, M., Nagase, M., Sugimura, Y.K., Takahashi, Y., Watabe, A.M., and Kato, F. (2015).
The lateral parabrachial nucleus is actively involved in the acquisition of fear memory in mice.
Mol. Brain *8*, 22.

Shi, C., and Davis, M. (1999). Pain Pathways Involved in Fear Conditioning Measured with Fear-
Potentiated Startle: Lesion Studies. J. Neurosci. *19*, 420–430.

Shinohara, K., Watabe, A.M., Nagase, M., Okutsu, Y., Takahashi, Y., Kurihara, H., and Kato, F.
(2017). Essential role of endogenous calcitonin gene-related peptide in pain-associated plasticity
in the central amygdala. Eur. J. Neurosci. *46*, 2149–2160.

Silva, B.A., Mattucci, C., Krzywkowski, P., Murana, E., Illarionova, A., Grinevich, V., Canteras,
N.S., Ragozzino, D., and Gross, C.T. (2013). Independent hypothalamic circuits for social and
predator fear. Nat. Neurosci. *16*, 1731–1733.

Silva, B.A., Gross, C.T., and Gräff, J. (2016). The neural circuits of innate fear: detection, integration, action, and memorization. Learn. Mem. *23*, 544–555.

Simons, L.E., Moulton, E.A., Linnman, C., Carpino, E., Becerra, L., and Borsook, D. (2014). The human amygdala and pain: Evidence from neuroimaging. Hum. Brain Mapp. *35*, 527–538.

Stamatakis, A.M., and Stuber, G.D. (2012). Activation of lateral habenula inputs to the ventral midbrain promotes behavioral avoidance. Nat. Neurosci. *15*, 1105–1107.

Tanimoto, S., Nakagawa, T., Yamauchi, Y., Minami, M., and Satoh, M. (2003). Differential contributions of the basolateral and central nuclei of the amygdala in the negative affective component of chemical somatic and visceral pains in rats. Eur. J. Neurosci. *18*, 2343–2350.

Thompson, J.M., and Neugebauer, V. (2017). Amygdala Plasticity and Pain. Pain Res. Manag. *2017*.

Todd, A.J. (2010). Neuronal circuitry for pain processing in the dorsal horn. Nat. Rev. Neurosci. *11*, 823–836.

Tong, W.H., Abdulai-Saiku, S., and Vyas, A. (2020). Medial Amygdala Arginine Vasopressin Neurons Regulate Innate Aversion to Cat Odors in Male Mice. Neuroendocrinology.

Veinante, P., Yalcin, I., and Barrot, M. (2013). The amygdala between sensation and affect: a role in pain. J. Mol. Psychiatry *1*.

Wang, L., Gillis-Smith, S., Peng, Y., Zhang, J., Chen, X., Salzman, C.D., Ryba, N.J.P., and Zuker, C.S. (2018). The coding of valence and identity in the mammalian taste system. Nature *558*, 127–131.

Wei, P., Liu, N., Zhang, Z., Liu, X., Tang, Y., He, X., Wu, B., Zhou, Z., Liu, Y., Li, J., et al. (2015). Processing of visually evoked innate fear by a non-canonical thalamic pathway. Nat. Commun. *6*, 6756.

Willis, W.D., Zhang, X., Honda, C.N., and Giesler, G.J. (2002). A critical review of the role of the proposed VMpo nucleus in pain. J. Pain *3*, 79–94.

Xiu, J., Zhang, Q., Zhou, T., Zhou, T., Chen, Y., and Hu, H. (2014). Visualizing an emotional valence map in the limbic forebrain by TAI-FISH. Nat. Neurosci. *17*, 1552–1559.

Yasui, Y., Saper, C.B., and Cechetto, D.F. (1991). Calcitonin gene-related peptide (CGRP) immunoreactive projections from the thalamus to the striatum and amygdala in the rat. J. Comp. Neurol. *308*, 293–310.

Yeh, L.-F., Watanabe, M., Sulkes-Cuevas, J., and Johansen, J.P. (2018). Dysregulation of aversive signaling pathways: a novel circuit endophenotype for pain and anxiety disorders. Curr. Opin. Neurobiol. *48*, 37–44.

793 Yokota, S., Kaur, S., VanderHorst, V.G., Saper, C.B., and Chamberlin, N.L. (2015). Respiratory-
794 related outputs of glutamatergic, hypercapnia-responsive parabrachial neurons in mice. J. Comp.
795 Neurol. *523*, 907–920.

796 Yu, L.-C., Hou, J.-F., Fu, F.-H., and Zhang, Y.-X. (2009). Roles of calcitonin gene-related peptide
797 and its receptors in pain-related behavioral responses in the central nervous system. Neurosci.
798 Biobehav. Rev. *33*, 1185–1191.

799 Zhang, X., and Giesler, G.J. (2005). Response Characterstics of Spinothalamic Tract Neurons That
800 Project to the Posterior Thalamus in Rats. J. Neurophysiol. *93*, 2552–2564.

801 Zhou, Z., Liu, X., Chen, S., Zhang, Z., Liu, Y., Montardy, Q., Tang, Y., Wei, P., Liu, N., Li, L., et
802 al. (2019). A VTA GABAergic Neural Circuit Mediates Visually Evoked Innate Defensive
803 Responses. Neuron *103*, 473-488.e6.

Figure Legends

Figure 1. CGRP^SPFp and CGRP^PBel neurons form spino-thalamo-amygdaloid and spino-parabrachio-amygdaloid pathways.

(**A**) Schematic and representative images of Cre-dependent expression of EYFP in the SPFp and mCherry in the PBel of a *Calca^Cre* mouse. Scale bars, 200 µm.

(**B**) The axonal projections from the CGRP^SPFp and CGRP^PBel neurons are mutually exclusive in the amygdala sub-regions. Scale bars, 500 µm.

(**C**) Schematic diagrams and images of Cre-dependent expression of TVA and G in SPFp or PBel neurons of *Calca^Cre* mice for the rabies tracing.

(**D**) Representative images of the SPFp and PBel five days after EnvA-RVdG-mCherry injection. Yellow indicates the starter cells. Scale bars, 200 µm.

(**E**) A schematic diagram of identifying presynaptic neurons by monosynaptic rabies tracing.

(**F**) Representative images of superficial and deep layer dorsal horn neurons that project to CGRP^SPFp neurons.

(**G**) The number of spinal dorsal horn neurons in the superficial (Sup) and deep (deep) layers project to the CGRP^SPFp neurons.

(**H**) The number of spinal cord neurons in different spinal segments that project to the CGRP^{SPFp} neurons.

(**I**) Representative images of the superficial dorsal horn and lateral spinal nucleus neurons project to

CGRP^{PBel} neurons.

(**J**) The number of spinal dorsal horn neurons in the superficial (Sup) and deep (deep) layers project to the

CGRP^{PBel} neurons.

(**K**) The number of spinal cord neurons in different spinal segments that project to the CGRP^{PBel} neurons.

Statistics

(G) Superficial: 10.20 ± 1.53, Deep: 16.60 ± 2.62 (n = 5). Paired t-test (two-tailed), p = 0.0723.

(H) Cervical: 17.40 ± 2.91, Thoracic: 5.80 ± 2.91, Lumbar: 2.80 ± 1.11, Sacral: 0.40 ± 0.24 (n = 5). Repeated

measure one-way ANOVA, p =0.0017. Cervical group was significantly different with lumbar ($p < 0.05$)

and sacral ($p < 0.05$, Tukey's multiple comparisons).

(J) Superficial: 1.00 ± 0.53, Deep: 1.43 ± 0.43 (n = 8). Paired t-test (two-tailed), p = 0.6286.

(K) Cervical: 0.63 ± 0.38, Thoracic: 0.13 ± 0.13, Lumbar: 1.00 ± 0.27, Sacral: 0.38 ± 0.18 (n = 8). Repeated

measure one-way ANOVA, p =0.1615. Thoracic group was significantly different with lumbar ($p < 0.05$,

Tukey's multiple comparisons).

Figure 2. CGRP^{SPFp} and CGRP^{PBel} neurons are activated by multimodal nociceptive stimuli.

(**A**) A diagram of the fiber photometry calcium imaging experiment in an anesthetized mouse.

(**B and C**) A schematic and representative image of the Cre-dependent expression of GCaMP6m in the *Calca^{Cre}* mice with an optical fiber implanted (red dashed line in the overlay image indicates fiber track) above the SPFp and PBel. Scale bars, 200 µm.

(**D-F**) Intensity-dependent calcium activity increase in CGRP^{SPFp} neurons in response to a pressure meter (0, 50, 100, 200, and 300 g).

(**G-I**) Intensity-dependent calcium activity increase in CGRP^{PBel} neurons to a pressure meter.

(**J-L**) Intensity-dependent calcium increase in CGRP^{SPFp} neurons in response to a temperature-controlled rod (25, 35, 45, or 55 °C).

(**M-O**) Intensity-dependent calcium activity increase in CGRP^{PBel} neurons in response to a temperature-controlled rod.

(**P**) Calcium signal increases in CGRP^{SPFp} neurons following subcutaneous injection of formalin (4%) into the paw.

(**Q**) Area under curve (A.U.C.) quantification of the CGRP^{SPFp} neuronal activity during the acute phase (5–10 min) of formalin response.

(**R**) A.U.C. quantification of the CGRP^{SPFp} neuronal activity during the inflammatory phase (15–45 min)
of formalin injection.
(**S**) Calcium signal increases in CGRP^{PBel} neurons in response to subcutaneous injection of formalin (4%)
into the paw.
(**T**) A.U.C. quantification of the CGRP^{PBel} neuronal activity during the acute phase (5–10 min) of formalin
response.
(**U**) A.U.C. quantification of the CGRP^{PBel} neuronal activity during the inflammatory phase (15–45 min) of
formalin response.
865
866
Statistics
(F) Repeated measure two-way ANOVA showed significance in interaction ($F_{(4, 152)} = 9.838$, $p < 0.0001$),
intensity ($F_{(4, 152)} = 12.28$, $p < 0.0001$) and group ($F_{(1, 38)} = 16.01$, $p = 0.0003$). 100 ($p < 0.05$), 200 (p
< 0.0001) and 300 g ($p < 0.0001$) points were significantly different between EYFP and GCaMP6m with
Sidak's multiple comparisons test.
(I) Repeated measure two-way ANOVA showed significance in interaction ($F_{(4, 248)} = 15.08$, $p < 0.0001$),
intensity ($F_{(4, 248)} = 15.17$, $p < 0.0001$) and group ($F_{(1, 62)} = 64.40$, $p < 0.0001$). 50 ($p < 0.0001$), 100
($p < 0.0001$) and 200 g ($p < 0.0001$) points were significantly different between EYFP and GCaMP6m with
Sidak's multiple comparisons test.
(L) Repeated measure two-way ANOVA showed significance in interaction ($F_{(3, 114)} = 10.97$, $p < 0.0001$),
intensity ($F_{(3, 114)} = 9.61$, $p < 0.0001$) and group ($F_{(1, 38)} = 11.12$, $p = 0.0019$). 55°C ($p < 0.0001$) was
significantly different between EYFP and GCaMP6m with Sidak's multiple comparisons test.
(M) Repeated measure two-way ANOVA showed significance in interaction ($F_{(3, 186)} = 23.96$, $p <$
0.0001), intensity ($F_{(3, 186)} = 22.67$, $p < 0.0001$) and group ($F_{(1, 62)} = 46.05$, $p < 0.0001$). 45 and 55°C
(both $p < 0.0001$) points were significantly different between EYFP and GCaMP6m with Sidak's multiple
comparisons test.
(Q) EYFP: 2.86 ± 8.00 (n = 4 mice), GCaMP6m: 471.9 ± 85.53 (n = 6 mice). Unpaired t-test (two-tailed),
$p = 0.0024$.
(R) EYFP: 100.60 ± 31.89 (n = 4), GCaMP6m: 894.80 ± 203.40 (n = 6). Unpaired t-test (two-tailed), $p =$
0.0145.
(K) EYFP: 6.18 ± 4.05 (n = 4), GCaMP6m: 77.00 ± 26.83 (n = 4). Unpaired t-test (two-tailed), $p = 0.0381$.
(L) EYFP: 100.60 ± 31.89 (n = 4), GCaMP6m: 443.50 ± 92.46 (n = 4). Unpaired t-test (two-tailed), $p =$
0.0127.
890

Figure 3. Pain-induced synaptic plasticity changes in CGRP^{SPFp} and CGRP^{PBel} neurons.

(**A**) A representative image of a brain slice including SPFp used for the whole-cell patch-clamp experiment. Enlarged images show the fluorescence from Cre:GFP-expressing CGRP neurons. Scale bars, 100 µm in left images, and 10 µm in enlarged images.

(**B**) Example traces of AMPA and NMDA EPSCs in control (blue) and formalin (red) injected groups. Scale bars, 20 ms, and 50 pA.

(**C**) The AMPA/NMDA ratio was increased in the formalin injected group compared with the control group.

(**D and E**) No significant differences in the paired-pulse ratio were observed between the formalin-injected group and the control group. Scale bars, 20 ms, and 50 pA.

(**F**) Example traces of mEPSCs in the control (blue) and formalin (red) injected group. Scale bars, 1 s, and 20 pA.

(**G**) mEPSC amplitude was increased in the formalin injected group compared with the control group.

(**H**) mEPSC interval was not changed in the formalin injected group compared with the control group.

(**I**) A representative image of a brain slice including PBel for the whole-cell patch-clamp experiment. Enlarged images are the PBel cells with and without fluorescence. Cells with fluorescence are CGRP neurons. Scale bars, 100 µm in left images, and 10 µm in enlarged images.

(**J**) Example traces of AMPA and NMDA EPSCs in control (blue) and formalin (red) injected groups. Scale bars, 20 ms, and 20 pA.

(**K**) The AMPA/NMDA ratio was increased in the formalin injected group compared with the control group.

(**L and M**) The formalin injected group display an increased paired-pulse ratio compared to the control group. Scale bars, 20 ms, and 50 pA.

(**N**) Example traces of mEPSCs in the control (blue) and formalin (red) injected group. Scale bars. 1 s and 50 pA.

(**O**) mEPSC amplitude was not changed in the formalin injected group compared with the control group.

(**P**) mEPSC interval increased in the formalin injected group compared with the control group.

Statistics

(C) SPFp; Control: 2.38 ± 0.31 (n = 11 cells), formalin: 4.62 ± 0.61 (n = 14 cells). Unpaired t test (two-tailed), p = 0.0063.

(E) SPFp; Control: 0.88 ± 0.09 (n = 10 cells), formalin: 0.77 ± 0.05 (n = 9 cells). Unpaired t test (two-tailed), p = 0.2957.

(G) SPFp; Control: 13.81 ± 0.90 (n = 11 cells), formalin: 17.83 ± 1.56 (n = 12 cells). Unpaired t test (two-tailed), p = 0.0390.

(H) SPFp; Control: 227.4 ± 40.90 (n = 11 cells), formalin: 154.2 ± 30.80 (n = 12 cells). Unpaired t test (two-tailed), p = 0.1687.

(K) PBel; Control: 2.49 ± 0.56 (n = 7 cells), formalin: 4.38 ± 0.65 (n = 11 cells). Unpaired t test (two-tailed), p = 0.0437.

(M) PBel; Control: 0.84 ± 0.04 (n = 8 cells), formalin: 1.20 ± 0.77 (n = 11 cells). Unpaired t test (two-tailed), p = 0.001.

(O) PBel; Control: 16.93 ± 2.08 (n = 7 cells), formalin: 18.85 ± 2.04 (n = 13 cells). Unpaired t test (two-tailed), p = 0.5196.

(P) PBel; Control: 173.2 ± 13.89 (n = 7 cells), formalin: 396.5 ± 80.66 (n = 13 cells). Unpaired t test (two-tailed), p = 0.0175.

939
Figure 4. Active transcriptome profiling of CGRP^{SPFp} and CGRP^{PBel} neurons.

(**A**) Correlation plot showing expression (transcript per million, TPM in log 10 scale) of genes enriched in

CGRP^{SPFp} neurons compared with the total SPFp inputs using RiboTag transcriptome profiling. Up and

down-regulated genes were separated based on 2.5 or -2.5-fold enrichment.

(**B**) Correlation plot of the transcriptome profiles of CGRP^{PBel} neurons versus total PBel inputs.

(**C**) Heatmaps showing fold changes of genes in the SPFp and PBel that encode membrane proteins.

(**D**) Heatmaps showing fold changes of genes in the SPFp and PBel that encode neuropeptides,

(**E**) Heatmaps showing fold changes of genes in the SPFp and PBel that encode markers of inhibitory neuron

or glia.

(**F-H**) Co-expression of Na$_V$1.7 (**F**), Ca$_V$2.1 (**G**), or FAAH (**H**) with CGRP by double IHC. Scale bars, 50

μm.

Figure 5. CGRP^{SPFp} and CGRP^{PBel} neurons are activated by multimodal sensory threat stimuli.

(**A and B**) A schematic and representative images of Cre-dependent expression of GCaMP7s in the CGRP^{SPFp} (**A**) and CGRP^{PBel} (**B**) neurons for fiber photometry. Scale bars, 200 µm.

(**C**) Cued fear conditioning test with low volume tone (72 dB) was performed to examine the responses of CGPR neurons to the somatosensory aversive stimulus (2-s, 0.6 mA electric foot shock).

(**D and E**) CGRP^{SPFp} (**D**) and CGRP^{PBel} (**E**) neurons were activated by the electric foot shock during the conditioning period, but not habituation, nor the cued retrieval period. Left, averaged calcium traces. Right, A.U.C. quantification.

(**F**) Intense sound (85 dB) was used as an aversive auditory stimulus, with a70-dB low-intensity sound as a control.

(**G and H**) CGRP^{SPFp} (**G**) and CGRP^{PBel} (**H**) neurons were activated by the intense sound. Left, averaged calcium traces. Right, A.U.C. quantification.

(**I**) A large looming disk was used as an aversive visual stimulus, with a small disk as a control.

(**J and K**) CGRP^{SPFp} (**J**) and CGRP^{PBel} (**K**) neurons were activated by the large looming disk. Left, averaged calcium traces. Right, A.U.C. quantification.

(**L**) TMT-soaked cotton was used as an aversive olfactory stimulus, with water as a control.

(**M**) There was no activity change in CGRP^{SPFp} neurons when the animal approached the TMT-soaked cotton. Left, averaged calcium traces. Right, A.U.C. quantification.

(**N**) CGRP^{PBel} neurons were activated when the animal approached the TMT-soaked cotton. Left, averaged calcium traces. Right, A.U.C. quantification.

(**O**) Quinine solution (0.5 mM) was introduced as an aversive gustatory stimulus (water was the control).

(**P**) Quinine did not affect CGRP^{SPFp} neurons. Left, averaged calcium traces. Right, A.U.C. quantification.

(**Q**) CGRP^{PBel} neurons were activated by quinine solution (0.5 mM). Left, averaged calcium traces. Right, A.U.C. quantification.

Statistics

(D) SPFp; Habituation: 3.63 ± 4.45, conditioning: 37.31 ± 7.18, cue test: -0.92 ± 2.57 (n = 6). Repeated measure one-way ANOVA, p = 0.0029. Conditioning was significantly different with habituation (p < 0.05) and cue test (p <0.01, Tukey's multiple comparisons).

(E) PBel; SPFp; Habituation: 1.85 ± 1.33, conditioning: 20.73 ± 4.25, cue test: 1.49 ± 1.35 (n = 6). Repeated measure one-way ANOVA, p = 0.0022. Conditioning was significantly different with habituation (p < 0.05) and cue test (p <0.01, Tukey's multiple comparisons).

(G) SPFp; 70 dB: 1.226 ± 1.27, 85 dB: 17.14 ± 2.38 (n = 6). Paired t test (two-tailed), p = 0.0018.

(H) PBel; 70 dB: 0.78 ± 1.46, 85 dB: 14.06 ± 5.86 (n = 7). Paired t test (two-tailed), p = 0.0437.

(J) SPFp; Small disk: 4.41 ± 1.11, large disk: 29.30 ± 5.81 (n = 6). Paired t test (two-tailed), p = 0.0091.

(K) PBel; Small disk: 1.08 ± 0.69, large disk: 6.93 ± 2.23 (n = 7). Paired t test (two-tailed), p = 0.0357.

(M) SPFp; Water: 1.93 ± 4.88, TMT: -1.82 ± 3.78 (n = 6). Paired t test (two-tailed), p = 0.4411.

(N) PBel; Water: -2.72 ± 2.84, TMT: 8.28 ± 3.57 (n = 6). Paired t test (two-tailed), p = 0.0145.

(P) SPFp; Water: 1.10 ± 3.09, quinine: -0.01 ± 1.22 (n = 6). Paired t test (two-tailed), p = 0.7707.

(Q) PBel; Water: 0.2209 ± 0.83, quinine: 4.96 ± 2.01 (n = 7). Paired t test (two-tailed), p = 0.0377.

Figure 6. Silencing CGRP^{SPFp} or CGRP^{PBel} neuronal activities attenuates perception of affective pain

and multisensory threat stimuli.

(**A**) Schematics and representative images of Cre-dependent TetTox expression targeting CGRP^{SPFp} or

CGRP^{PBel} neurons. Scale bars, 200 μm.

(**B**) A schematic diagram of the Formalin assay to test inflammatory pain responses.

(**C**) The CGRP^{SPFp} and CGRP^{PBel} silenced groups displayed significantly alleviated inflammatory pain

responses.

(**D**) Both the CGRP^{SPFp} and CGRP^{PBel} silenced groups froze less in response to electric foot shock (2-s, 0.6

mA) compared to controls.

(**E**) Both the CGRP^{SPFp} and CGRP^{PBel} silenced groups froze less in response to 85-dB intense sound

compared with controls.

(**F**) In response to a looming aversive visual stimulus, the CGRP^{SPFp} silenced group showed less freezing,

whereas there was no difference between EYFP and CGRP^{PBel} silenced groups.

(**G**) A two-chamber choice test with TMT- and water-soaked cotton placed at each corner of the chamber

was used to test animals' responses to aversive olfactory stimulus. The CGRP^{SPFp} silenced group, and EYFP

controls both avoided the TMT chamber, while the CGRP^{PBel} TetTox group exhibited less aversion to TMT,

spending equal amounts of time in the water and TMT chambers.

(**H**) A quinine-water choice test was performed to test the animals' responses to the aversive gustatory

stimulus (0.5 mM quinine solution). There was no change in water preference in the CGRP^{SPFp} silenced

group, but the CGRP^{PBel} silenced group showed a lower difference index than the control group.

1014

1015

Statistics

(C) SPFp: Repeated measure two-way ANOVA showed no significance in interaction ($F_{(11, 187)} = 1.32$,

$p = 0.2148$), but in time ($F_{(11, 187)} = 9.05$, $p < 0.0001$) and group ($F_{(1, 17)} = 9.57$, $p = 0.0066$). 20-25 (p

< 0.05) and 25-30 ($p < 0.01$) min point were significantly different between EYFP and TetTox with Sidak's

multiple comparisons test.

PBel: Repeated measure two-way ANOVA showed significance in interaction ($F_{(11, 187)} = 3.45$, $p =$

0.0002), time ($F_{(11, 187)} = 5.97$, $p < 0.0001$) and group ($F_{(1, 17)} = 53.24$, $p < 0.0001$). 15-20 ($p < 0.05$),

25-30 (p < 0.0001), 30-35 (p < 0.0001), 35-40 (p < 0.0001), 40-45 (p < 0.01), 45-50 (p < 0.01) min point

were significantly different between EYFP and TetTox with Sidak's multiple comparisons test.

(D) SPFp; EYFP: 56.62 ± 6.21 % (n = 9), TetTox: 34.65 ± 6.11 % (n = 8). Unpaired t test (two-tailed), p =

0.0240.

PBel; EYFP: 33.43 ± 6.66 % (n = 9), TetTox: 12.04 ± 4.88 % (n = 9). Unpaired t test (two-tailed), p =

0.0197.

(E) SPFp; EYFP: 45.74 ± 10.26 % (n = 9), TetTox: 17.32 ± 5.58 % (n = 8). Unpaired t test (two-tailed), p

= 0.0332.

PBel; EYFP: 35.39 ± 5.29 % (n = 7), TetTox: 18.14 ± 4.66 % (n = 6). Unpaired t test (two-tailed), p =

0.0350.

(F) SPFp; EYFP: 43.12 ± 9.82 % (n = 9), TetTox: 5.73 ± 1.57 % (n = 8). Unpaired t test (two-tailed), p =

0.0029.

PBel; EYFP: 56.20 ± 10.33 % (n = 8), TetTox: 51.22 ± 13.21 % (n = 7). Unpaired t test (two-tailed), p =

0.0029.

(G) SPFp; EYFP: 20.12 ± 4.91 % (n = 9), TetTox: 15.55 ± 7.75 % (n = 8). Unpaired t test (two-tailed), p =

0.6175.

PBel; EYFP: 13.87 ± 3.94 % (n = 11), TetTox: -1.53 ± 5.89 % (n = 10). Unpaired t test (two-tailed), p =

0.0396.

(H) SPFp; EYFP: 69.96 ± 4.64 % (n = 9), TetTox: 72.66 ± 4.48 % (n = 8). Unpaired t test (two-tailed), p =

0.6823.

PBel; EYFP: 86.87 ± 1.77 % (n = 8), TetTox: 31.66 ± 9.45 % (n = 7). Unpaired t test (two-tailed), p <

0.0001.

1045

1046

Figure 7. Activating the CGRP$^{SPFp \to LA}$ or CGRP$^{PBel \to CeA}$ pathways encodes negative valence.

(**A**) Schematics and representative image of Cre-dependent expression of ChR2 in the SPFp of *Calca^{Cre}* mice and optic fiber placement (red dotted line).

(**B**) A schematic diagram and heatmap of the real-time place aversion (RTPA) test with CGRP^{SPFp} cell body stimulation.

(**C**) Time spent in the stimulated zone during the RTPA test.

(**D**) Representative image of optic fiber placement in the LA for terminal stimulation (red dotted line).

(**E**) Time spent in the stimulated zone during the RTPA test with CGRP$^{SPFp \to LA}$ terminal stimulation.

(**F**) Time spent freezing during the context test.

(**G**) Time spent freezing during cue test after pairing optic stimulation and tone for cued fear conditioning.

(**H**) Schematic and representative image of Cre-dependent expression of ChR2 in the PBel of *Calca^{Cre}* mice and optic fiber placement (red dotted line).

(**I**) Schematic diagram and heatmap of real-time place aversion (RTPA) test with CGRP^{PBel} cell body stimulation.

(**J**) Time spent in the stimulated zone during the RTPA test with CGRP^{PBel} cell body stimulation.

(**K**) Representative image of optic fiber placement in the CeA (red dot).

(**L**) Time spent in the stimulated zone during the RTPA test with CGRP$^{PBel \rightarrow CeA}$ terminal stimulation.

(**M**) Time spent freezing during the context test.

(**N**) Time spent freezing during cue test after pairing optogenetic stimulation and tone for cued fear conditioning. Scale bars indicate 200 µm.

Statistics

(C) Repeated measure two-way ANOVA showed significance in laser x group interaction ($F_{(1, 17)} = 17.69$, $p = 0.0006$) and laser ($F_{(1, 17)} = 20.19$, $p = 0.0003$) but not group ($F_{(1, 17)} = 4.03$, $p = 0.0608$). Laser ON and OFF in ChR2 ($p < 0.0001$), and EYFP and ChR2 during ON ($p < 0.001$) had statistically significant difference with Sidak's multiple comparisons test.

(E) Repeated measure two-way ANOVA showed significance in only laser x group interaction ($F_{(1, 18)} = 8.718$, $p = 0.0085$) but not laser ($F_{(1, 18)} = 3.159$ $p = 0.0924$) and group ($F_{(1, 18)} = 4.343$, $p = 0.0517$). Laser ON vs OFF in ChR2 ($p < 0.05$), and EYFP vs ChR2 during ON ($p < 0.01$) have statistically significant difference with Sidak's multiple comparisons test.

(F) EYFP: 3.842 ± 0.89 (n = 13), ChR2: 31.92 ± 7.806 (n = 10). Unpaired t test (two-tailed), $p = 0.0005$.

(G) Repeated measure two-way ANOVA showed significance in CS x group interaction ($F_{(1, 21)} = 12.41$, $p = 0.0020$), CS ($F_{(1, 21)} = 38.26$, $p < 0.0001$) and group ($F_{(1, 21)} = 8.392$, $p = 0.0086$). CS- vs CS+ in ChR2 ($p < 0.0001$), and EYFP vs ChR2 during CS+ ($p < 0.001$) have statistically significant difference with Sidak's multiple comparisons test.

(J) Repeated measure two-way ANOVA showed significance in laser x group interaction ($F_{(1, 9)} = 10.49$, $p = 0.0102$) and laser ($F_{(1, 9)} = 8.959$, $p = 0.0151$) but not group ($F_{(1, 9)} = 1.159$, $p = 0.3096$). Laser ON and OFF in ChR2 ($p < 0.01$), and EYFP and ChR2 during ON ($p < 0.05$) had statistically significant difference with Sidak's multiple comparisons test.

(L) Repeated measure two-way ANOVA showed significance in laser x group interaction ($F_{(1, 9)} = 12.91$, $p = 0.0058$) and laser ($F_{(1, 9)} = 6.094$, $p = 0.0357$) but not in group ($F_{(1, 9)} = 4.405$, $p = 0.0652$). Laser ON and OFF in ChR2 ($p < 0.01$), and EYFP and ChR2 during ON ($p < 0.01$) had statistically significant difference with Sidak's multiple comparisons test.

(M) EYFP: 6.278 ± 1.244 (n = 5), ChR2: 28.93 ± 5.133 (n = 6). Unpaired t test (two-tailed), $p = 0.0035$.

(N) Repeated measure two-way ANOVA showed significance in CS x group interaction ($F_{(1, 9)} = 12.67$, $p = 0.0061$), CS ($F_{(1, 9)} = 8.226$, $p = 0.0185$) and group ($F_{(1, 9)} = 6.314$, $p = 0.0332$). CS- vs CS+ in ChR2 ($p < 0.01$), and EYFP vs ChR2 during CS+ ($p < 0.01$) have statistically significant difference with Sidak's multiple comparisons test.

1097 **Supplementary Information**
1098
Materials and methods
Animals
All protocols for animal experiments were approved by the IACUC of the Salk Institute for Biological
Studies according to NIH guidelines for animal experimentation. The *Calca^{Cre}*, and *Tacr1^{Cre}* transgenic
mouse lines used in this study were generated from the Richard Palmiter's laboratory (Han et al., 2015 or
Carter et al., 2013) *Cdx2^{FlpO}* line was generated from the Martyn Goulding's laboratory. *Calca^{CreER}* mouse
line was generated from the Pao-Tien Chuang's laboratory. RiboTag *Rpl22$^{HA/HA}$* (Stock No. 011029) and
Ai65 (Stock No: 021875) mouse line was obtained from the Jackson Laboratory. All mouse lines are
backcrossed with C57Bl/6J for > 6 generations. Male and female mice were used in all studies. Animals
were randomized to experimental groups and no sex differences were noted. Mice were maintained on a
normal 12 hours light/dark cycle and provided with food and water *ad libitum*.
1110
Stereotaxic surgery for virus injection and optic fiber implantation
Mice were anesthetized by isoflurane gas anesthesia (induction at 3.5%, and maintenance at 1.5-2%, the
Dräger Vapor® 2000; Draeger, Inc., USA). Mice were then placed on a stereotaxic frame (David Kopf
Instruments, USA). Holes were drilled with a micromotor handpiece drill (Foredom, USA) after the
exposure of the skull. The virus was injected using a syringe (65458-01, Hamilton, USA) connected to an
ultra-micropump (UMP-3, World Precision Instruments, USA). Unilateral (right side) and bilateral
injections were made for the following target regions: SPFp (antero-posterior (AP), -3.1 mm; medio-lateral
(ML), 2.0 mm; dorso-ventral (DV) -3.6 mm from bregma) or PBel (AP, -5.1 mm; ML, 1.35 mm; DV, -3.5
mm). Viruses were injected at a rate of 0.08 µl/min (total volume of 0.75 µl for optogenetic projection
studies and 0.5 µl for all the others) and the syringe needle was slowly removed from the injection site
seven-minute after injection. To determine the inputs to CGRP^{SPFp} and CGRP^{PBel} neurons, 0.5 µl of AAV8-
hSyn-FLEX-TVA-P2A-GFP-2A-oG (Salk Institute viral vector core, USA) was injected into the SPFp or
PBel of *Calca^{Cre}* transgenic mice. After three weeks, 0.5 µl of EnvA-ΔG-rabies-mCherry (Salk Institute
viral vector core, USA) was injected, and the mice were sacrificed five days after the injection. To silence
the CGRP^{SPFp} and CGRP^{PBel} neurons, 0.5 µl of AAV-DIO-TetTox-GFP or AAV-DIO-EYFP was injected
into the SPFp or PBel of *Calca^{Cre}* transgenic mice, and experiments were performed two weeks after
injection. For fiber photometry experiments, mice were injected with 0.5 µl of either AAV-DIO-GCaMP6m,
AAV-DIO-GCaMP7s or AAV-DIO-EYFP into the SPFp or PBel of *Calca^{Cre}* mice. Stainless-steel mono
fiberoptic cannulas (400 um diameter, 0.37 NA, Doric Lenses) were implanted above the SPFp or PBel.
For electrophysiology, mice were injected with 0.5 or 0.75 µl of AAV1-DIO-ChR2-EYFP into the SPFp or
PBel of *Calca^{Cre}* mice. Experiments were performed two weeks after viral injection for recording SPFp and
PBel neurons or four weeks after injection for recording cells in terminal regions. For optogenetics, mice
were injected with 0.5 or 0.75 µl of AAV1-DIO-ChR2-EYFP or AAV-DIO-EYFP into the SPFp or PBel
of *Calca^{Cre}* mice and custom made mono fiberoptic cannula (200 um diameter, 0.22 NA) were implanted
above SPFp (0.5 mm above the injection site), PBel (0.5 mm above the injection site), Astr (AP, -1.8 mm;
ML, 3.3 mm, DV, -3.8 mm from bregma), lateral amygdala (LA; AP, -1.8 mm; ML, 3.6 mm, DV, -4.0 mm
from bregma), pIC (AP, -1.5 mm; ML, 4.6 mm, DV, -3.0 mm from bregma) or central amygdala (CeA; AP,
1.2 mm; ML, 2.7 mm; DV, -4.2 mm from bregma). Experiments were executed two weeks after injection
to manipulate SPFp neurons or four weeks later for terminal stimulations.
1140

Histology and quantification of rabies tracing experiment

Mice were intracardially perfused with 4% paraformaldehyde in PBS 5 days after the rabies virus injection. Spinal cords were post-fixed at 4°C for 1 h and dehydrated with 30% sucrose at 4°C overnight. 40 um transverse sections were obtained with a cryostat (CM 1950, Leica, USA) throughout the spinal cord. Spinal cord slices were directly dry mounted on superfrost plus microscope slide glasses (12-550-15, Fisher Scientific, USA). The labeled neurons were counted manually by dividing the transverse spinal sections into four groups (cervical, thoracic, lumbar, and sacral) or different dorsal horn layers. Brains were kept in 4% PFA overnight for post-fixation and dehydrated in 30% sucrose for 1-2 days before sectioning. Frozen brains were cut into 50 µm coronal slices with a cryostat and stored in Phosphate buffered saline before mounting. Both spinal cord and brain tissues were mounted on a slide glass with a DAPI containing mounting solution (0100-20, SouthernBiotech, USA).

Fiber photometry

Bulk calcium signals from the CGRP^{SPFp} and CGRP^{PBel} neurons were monitored using a custom-built fiber photometry system based on the open-source pyPhotometry platform (https://pyphotometry.readthedocs.io/en/latest/). 465 nm LED was used to induce Ca^{2+} dependent fluorescence signals, and 405 nm LED was used for Ca^{2+} independent (isosbestic) fluorescence signals. Motion corrected ΔF/F was calculated by a post-hoc analysis ($\Delta F/F = F_{465} - F_{405fit}/F_{405fit}$). The least-squares polynomial function was used to calculate F_{405fit}, and the area under the curve was used to analyze the data.

Multi-modal aversive stimuli experiments

Mechanical and thermal stimuli

Mechanical and thermal stimuli were applied to mice forepaw, hind paw, or tail. Mechanical pressure was applied using a dial tension gauge (ATG-300-1, Vetus Instruments, USA) with stimulus strength 0, 50, 100, 200 g. The thermal stimulus was applied using a custom-made temperature-controlled hot-rod (TA4-SNR+K, Mypin, China) at 25, 35, 45, and 55°C. A stable baseline was recorded first for 10 s, and stimuli were applied immediately after 5 seconds.

Formalin test

For fiber photometry, lightly anesthetized mice were placed in the stereotaxic frame head fixed to minimize movement. 10 µl of 4% formalin (1.6% Paraformaldehyde, 19210, Electron Microscopy Sciences, USA) was injected subcutaneously on the contralateral forepaw after at least 5 min of stable baseline. Calcium transients were recorded for 45 min. For the loss of function experiment, 10 µl of 4 % formalin was injected subcutaneously on one side of the forepaw. Mice were then placed in a Plexiglas chamber (10 x 10 x 13 cm) with a mirror placed behind. Behaviors were recorded for an hour, and licking behaviors were manually counted throughout the experiment.

Auditory and visual stimuli

For both auditory and visual stimuli experiments, mice were placed in a cylinder-shaped arena (11 cm diameter with 15 cm height) with homecage bedding and were habituated for 30-120 min. For auditory experiments, after a stable 10s baseline, an intense sound (85 dB, 2 s) or a control sound (70 dB) was played. For the loss-of-function experiments, mice were placed inside an open field chamber and were habituated for 10 min. After 1 min baseline, an intense sound (85 dB, 2 s) was delivered three times with an inter-stimulus interval of 28 s. All the trials were recorded by a USB camera (DFK 33GX236, Imagine Source, Germany) attached to a computer, and freezing behavior was analyzed using video-tracking software (Ethovision XT, Noldus, Netherlands). For visual looming experiments, after a stable 10 s baseline, an

expanding looming stimulus (2 s) was delivered three times with 10 s inter-stimulus interval with a LED
screen facing the arena from above. For the loss-of-function experiment, mice were placed in a cage with
bedding and were positioned under the same LED screen. Mice were habituated for 20-30 min. When mice
were in the center, the expanding looming stimulus (2 s) was delivered three times with 10 s inter-stimulus
interval. All the trials were recorded by a USB camera (DFK 33GX236, Imagine Source, Germany)
attached to a computer, and freezing behavior was analyzed using video-tracking software (Ethovision XT,
Noldus, Netherlands) with manual counting for the duration of tail rattling behaviors.

Gustatory stimulus
For fiber photometry, mice were placed in the same arena for auditory and visual stimulus experiments with
an additional 2 cm drilled hole. The water bottle spout was inserted into the hole, and the calcium signal
was measured when the mice were licking. The bottle was filled with water or quinine (0.5 mM, QU109,
Spectrum Chemical, USA). For the loss-of-function experiment, mice were water-deprived overnight. The
next day, mice were placed in a homecage with a water-, and 0.5 mM quinine-containing bottle inserted
into the water valve slot. Mice were allowed to drink for 10 min without habituation. All the trials were
recorded by a USB camera (DFK 33GX236, Imagine Source, Germany) attached to a computer, and the
licking behaviors were counted manually.

Olfactory stimulus
For fiber photometry, mice were placed in the same arena for gustatory stimulus experiments. Water- or
Trimethylthiazoline (TMT, 97%, 5 µl, 1G-TMT-97, BioSRQ, USA)-soaked cotton swab was introduced
into the hole. Calcium signals were measured when mice smelled the cotton swab. For the loss-of-function
experiment, mice movement was tracked in a two-chamber arena (30 x 60 x 30 cm) with a USB camera
(DFK 33GX236, Imagine Source, Germany) using video-tracking software (EthoVision XT 12, Noldus,
Netherlands). Two Petri dishes with small holes were placed in each chamber (one at the corner of the left
chamber, and the other to the corner of the right chamber). On day 1, mice were able to habituate and
explore the arena for 10 min. The next day, a water-soaked cotton swab, or TMT-soaked cotton swab were
placed in each dish. Mice were first placed at the center and monitored for 10 minutes as they interreacted
with the two dishes.

Foot shock
A fear-conditioning chamber (26 x 30 x 33 cm, ENV-007CT, MED Associates INC, USA) with a metal
grid floor (ENV-005, MED Associates INC, USA) connected to a standalone aversive electric shock
stimulator (ENV-414S, MED Associates INC, USA) was used for foot shock delivery. A USB camera
(DFK 33GX236, Imagine Source, Germany) was connected to a computer, and the video tracking software
(Ethovision XT, Noldus, Netherlands) was used for shock delivery and behavioral analysis. The chamber
was enclosed in a light- and sound-attenuating cubicle (ENV-018MD, MED Associates INC, USA). The
chamber was cleaned with 70% ethanol and double distilled water between each trial.
For fiber photometry and the loss-of-function experiment, mice were placed inside the chamber without
habituation. After 2 min of baseline, an electric shock (2 s, 0.6 mA) was delivered, and the behavior was
recorded for an extra 2 min. Freezing behavior was monitored before (habituation), after (conditioning),
and one day after (post-test) the shock.

Elevated plus maze test
A custom-built elevated plus maze with two transparent closed arms (77 x 7 x 30 cm) and two open arms
(77 x 7 x 2 cm) was used to monitor the anxiety-like behaviors of test mice. This maze was elevated 70 cm
above ground for all tests. Mice were placed to the tip of the open arm by facing towards the center of the
maze. The behavior was video recorded for 10 min and tracked with a video-tracking software (EthoVision

XT 12, Noldus, Netherlands). Both 70% ethanol solution and deionized water were used to clean the maze
immediately after each trial.

Hot plate test
Mice were placed into a cylinder-shaped transparent Plexiglas chamber (11 cm diameter with 15 cm length)
on a heated hot plate (48 or 55°C, PE34, IITC Life Science, USA). The latency of various pain responses
(hind paw shake, lick, or jump) was measured manually.

Electronic von Frey test
A Dynamic Plantar Aesthesiometer (37450, Ugo Basile, Italy) was used to measure the mechanical pain
thresholds. Mice were placed inside a Plexiglas chamber (10 x 10 x 13 cm) on a metal mesh floor and were
habituated for 30 min. The max force of the system was set to reach 50 g at 20 s. The blunt metal rod of the
aesthesiometer was placed under the hind paw and gradually protruded as the mice were immobile but
awake. The latency and force delivered were automatically recorded as the mouse withdraw hind paw from
the metal rod. The measurement was performed 5 times with 5-10 min intervals in between trials and
averaged for a final mechanical threshold value.

1243

Optogenetics
A 470 nm laser (LRD-0470-PFFD-00100-05, LaserGlow Tech., Canada) was used for all optogenetic
experiments in this study. Optic fibers were bilaterally connected to pre-implanted optic ferrules on the
mice. All mice were optogenetically stimulated 90 min before sacrifice for cFos immunohistochemistry.

Hot plate test
The experiments were performed as described in the '*Hot plate test'* section above with minor modification
for optogenetic stimulation. The order of laser ON or OFF was counterbalanced, and the interval time
between each experiment was more than 30 min.

Electronic von Frey test
The experiments were performed as described above in the '*Electronic von Frey test'* with minor
modification for optogenetic stimulation. The laser was on immediately before the metal rod touched the
paw pad and turned off right after paw withdrawal.

Real-time place aversion (RTPA)
A two-chamber arena (30 x 60 x 30) was used for the RTPA test. The behavior was tracked with a USB
camera (DFK 33GX236, Imagine Source, Germany) using video-tracking software (EthoVision XT 12,
Noldus, Netherlands). After connecting the optic fiber, mice were placed in one side of the chamber. No
stimulation was given for 10 min baseline. Afterward, one side of the chamber was randomly selected, and
the mouse was photostimulated (20 Hz for cell body stimulation, and 40 Hz for terminal stimulation, 8-9
mW) for 20 min. The stimulated side was counterbalanced between animals. Mice showing over 15%
preference to one side during baseline were excluded.

Context-dependent optogenetic conditioning
An open field arena (40 x 40 x 30 cm) was used for context-dependent threat conditioning. After 10-min
habituation to head-attached optic fibers in the home cage, mice were placed in the novel open field area
and received photostimulation (20 Hz, and 8-9 mW) throughout the experiment. After 24 h, the mouse was
re-introduced in the same context to test whether the photo-stimulation produced aversive memory. All the
trials were recorded by a USB camera (DFK 33GX236, Imagine Source, Germany) attached to a computer,
and freezing behavior was analyzed by a video-tracking software (EthoVision XT 12, Noldus, Netherlands).

Auditory cue dependent optogenetic conditioning

The same fear-conditioning chamber and the settings as described in the '*Foot shock'* section above were
used. Two speakers (AX210, Dell, USA) were placed beside the chamber for CS. On day 1, the test mouse
was habituated with the conditioning chamber, which was cleaned with 70% ethanol and DW immediately
after each test. During habituation, optic fibers were connected bilaterally to the optic ferrules on the
mouse's head, and the CS+ (30 s, 3 kHz pure tone, 75 dB) was delivered to the test mouse six times with
random inter-event intervals. On day 2, the test mouse was returned to the same context with optic fibers
connected and received 10-s photostimulation (20 Hz frequency for cell body and 40Hz for terminal
stimulation, 8-9 mW intensity) as the US, which was co-terminated with CS+ six times with random inter-
event intervals. On day 3, the conditioned mouse without the optic fiber connected was returned to the same
context for 2 min. On day 4, a conditioned mouse without the optic fiber connected was introduced to a
new context (a glass cylinder wrapped with a non-transparent material), and the CS+ was delivered without
the US. All the trials were recorded by a USB camera (DFK 33GX236, Imagine Source, Germany) attached
to the computer, and freezing behavior was analyzed by a video-tracking software (EthoVision XT 12,
Noldus, Netherlands).
1286
Immunohistochemistry
Mice were perfused intracardially with % PFA solution in PBS, and the brain was extracted. The brain was
kept in 4% PFA overnight for post-fixation and dehydrated in 30 % sucrose for 1-2 days before sectioning.
Frozen brains were cut into 40 µm coronal slices with a cryostat (CM 1950, Leica, USA) and washed with
PBST (Phosphate buffered saline with 0.1% Tween-20 (BP337-500, Fisher BioReagents, USA)). Initial
blocking was performed by 1hr incubation with 3% normal donkey serum (NDS, 017-000-121, Jackson
ImmunoResearch Laboratories, Inc., USA). After another round of washing with PBST, the slices were
incubated with anti-GFP (diluted 1:100 in 3% NDS, GFP-1020, Aves, USA), anti-fos (1:10000, rabbit
polyclonal), anti-Na$_V$1.7 (1:200, ASC-008, Alomone), anti-Ca$_V$2.1 (1:100, ACC-001, Alomone, Isreal), or
anti-FAAH (1:250, 101600, Cayman, USA) antibody at 4 ℃ overnight. The next day, brain tissues were
rinsed with PBST, then incubated with anti-rabbit Alexa Fluor® 647-secondary antibody (1:500, 711-605-
152, Jackson ImmunoResearch Laboratories, Inc., USA), and / or anti-chicken Alexa Fluor® 488-secondary
antibody (1:500, Jackson ImmunoResearch Laboratories) for 1 h. After washing these brain slices with PBS,
they were mounted on slide glass (12-550-143, Fisher Scientific, USA) with DAPI containing mounting
solution.
1302
Preparation of acute brain slices and electrophysiology
Mice were anesthetized with isoflurane and perfused via the vascular system using ice-cold cutting solution
(110.0 mM choline chloride, 25.0 mM NaHCO$_3$, 1.25 mM NaH$_2$PO$_4$, 2.5 mM KCl, 0.5 mM CaCl$_2$, 7.0 mM
MgCl$_2$, 25.0 mM glucose, 5.0 mM ascorbic acid and 3.0 mM pyruvic acid, bubbled with 95% O$_2$ and %
CO$_2$). After decapitation, brains were quickly removed and chilled in an ice-cold cutting solution. Coronal
slices containing the SPFp, PBel (250 µm) or the amygdaloid complex (300 µm) were cut by using a Leica
VT 1200S Vibratome (Leica Biosystems Inc.), and subsequently transferred to a storage chamber
containing artificial cerebrospinal fluid (aCSF; 124 mM NaCl, 2.5 mM KCl, 26.2 mM NaHCO$_3$, 1.2 mM
NaH$_2$PO$_4$, 13 mM glucose, 2 mM MgSO$_4$ and 2 mM CaCl$_2$, at 32 °C, pH 7.4, bubbled with 95% O$_2$ and 5%
CO$_2$). After at least 30 min recovery time, slices were transferred to room temperature (22–24°C) for at
least 60 min before use. Slices were transferred into the recording chamber, perfused with aCSF (flow rate
around 2 ml/ min). The temperature of aACSF was held constant at 32°C by TC-324C temperature
controller (Warner Instruments). Since CGRP-positive neurons express EGFP under the *Calca* promoter,

they were visualized under Scientifica Microscope equipped with epifluorescence illumination at 490 nm LED. The Astr, LA, and IC neurons were visualized under trans-illumination. Whole-Cell patch clamp was performed with Multiclamp 700B amplifiers (Molecular Devices). Signals were digitized at 10 kHz with Digidata 1550B (Molecular Devices). For evoked EPSCs, synaptic responses were evoked with a broken glass pipette positioned 100 µm away from the recording glass electrode (3.0~5.0 MOm, back filled with internal solution: $CsMeSO_3$ 130 mM, CsCl 5, HEPES 10 mM, $MgCl_2$ 2.5 mM, EGTA 0.6 mM, Sodium phosphocreatine 10 mM, Na_2ATP 4 mM and Na_3GTP 0.4 mM, pH 7.23, 285 Osm). The stimulus was given at 0.1 Hz. AMPA EPSC was recorded holding at -70 mV for 10 to 30 sweeps to get a stable response. NMDA EPSC was recorded at +40 mV for 10 – 15 sweeps. To ensure that the EPSCs were stable, the holding potential was set to -70 mv to check the AMPA EPSC change after NMDA EPSC recording. Evoked EPSCs were recorded with picrotoxin (100 μM) in the aCSF. mEPSCs were recorded in the presence of tetrodotoxin (1 μM) and picrotoxin (100 μM). To record optogenetically evoked EPSC and IPSC, slices were harvested from the AAV-DIO-ChR2-EYFP injected $Calca^{Cre}$ mice brain. 2-ms 470 nm LED light (TTL from Clampex to Cool Led pE-300) was illuminated through 40X NA 0.8 objective lens at 0.1 Hz to evoke optogenetically evoked postsynaptic current. The internal solution was calculated to make chloride reversal potential at -70 mV. EPSCs were recorded at -70 mV, and IPSCs were recorded at 0mV. CNQX (10 μM) was perfused to check the glutamatergic synapse. EPSCs and IPSCs were analyzed using pCLAMP 10 software (Molecular Devices). NMDA EPSCs were defined as signals 100 ms apart from stimulus artifacts. mEPSCs were analyzed using Mini Analysis Program (Synaptosoft).

Imaging
The images were taken with an automatic fluorescence microscope (BZ-X710, Keyence, USA) using included imaging software (BZ-X viewer, Keyence, USA) or with a scanning confocal microscope (FV 1000, Olympus, Japan) using with Fluoview software (Olympus, Japan). For quantification purposes, images were processed with the same gain, offset, and exposure time. Cell counting for retrograde tracing was done manually.

RiboTag Transcriptomic Profiling
To label the active transcriptome of CGRP^{SPFp} and CGRP^{PBel} neurons, we crossed $Calca^{CreER}$ with RiboTag $Rpl22^{HA/HA}$ mice. To induce gene expression, 200 µl of tamoxifen freshly prepared with 20 mg/ml in corn oil and dissolved overnight with continuous agitation was administered intraperitoneally for five consecutive days in each mouse. Experiments were performed two weeks after the final tamoxifen injection. 250 µm thick slices containing the PBel and the SPFp were obtained using a VT 1200S Vibratome (Leica, Germany). The region of interest was further dissected using surgical scissors under the stereoscope. Tissues of interest from four $Calca^{CreER}$; RiboTag crossed mice (10-12 weeks old) were collected into 1.5 mL microcentrifuge tubes containing homogenization buffer and were mechanically dissociated and lysed using pellet pestles (Cat.no.7495211500-DS, DWK Life Sciences LLC, USA). Total RNA was extracted from 15% of cleared lysate for input samples. The remaining lysate was incubated with mouse anti-HA antibody (Cat.no.MMS-101R, Biolegend, USA) and was rocked for 4 hours at 4 °C. Afterward, magnetic beads (Cat.no.88803, Thermo Fisher Scientific, USA) were added, and the solution was incubated overnight, rocking at 4 °C. The beads were washed three times in high salt solution. The bound ribosomes and RNA were separated from the beads by 30 s of vortexing in RLT lysis buffer as IP. All RNA samples were purified from the IP and corresponding input samples (Qiagen RNeasy Mini Kit, cat.no. 74104), then quantified with the Qubit RNA Assay Kit (Invitrogen, USA) and analyzed with the RNA 6000 Pico Kit

(Agilent, USA). Isolated RNA was prepared using the Trio RNA-Seq (Cat. No. 0507-08; NuGEN, USA).
Briefly, cDNA was synthesized from the total RNA using reverse transcriptase with oligo dT and
resynthesized to produce double-stranded cDNA. After amplifying double-stranded cDNA, cDNA was
purified with AMPure XP Beads (Cat. No. A63881; Beckman Coulter, USA), fragmented to the library,
and classified using a barcoded adaptor. All libraries were quantified by qPCR and analyzed with the RNA
6000 Pico Kit. RNA library quality was checked using the 2100 Bioanalyzer (Agilent, USA). Barcoded
samples were pooled and sequenced on the NextSeq500 (Illumina, USA) with the 75 bp read length single-
end. Image analysis and base calling were conducted using the Illumina CASAVA-1.8.2 software. The
FastQC package was utilized to evaluate the sequencing read quality. Fastq reads were then aligned to the
reference genome (GRCm38.p6) using the STAR tool (version 2.7.2) in a pair-end mode. The quantification
package RSEM (version 1.2.28) was employed to calculate gene expression from BAM files using the
default setting changed to pair-end mode. In doing so, estimated count and TPM (Transcripts Per Million)
were generated. Fold changes were calculated from TPM values (estimated counts, > 20) between HA-tag
and HA negative controls., The ggplot2 package from R was utilized to visualize fold changes. UP (> 2.5-
fold change) and DOWN ($< - 2.5$-fold change) were highlighted with orange and blue colors, respectively.
1375
1376 *Statistical analysis*
1377 All data are shown as mean $\pm$ s.e.m. and analyzed using Student's t-test, one-way ANOVA with Tukey's
1378 post hoc comparison, and two-way ANOVA with Sidak's post hoc comparison. All the statistical analyses
1379 were done using Prism 6 (GraphPad Software Inc., USA). NS p>0.05, * $p < 0.05$, ** $p < 0.01$, *** $p <$
1380 0.001
1381

 Supplementary Figure Legends

1383

Figure S1. Identification of direct spino-recipient areas in the brain by specific genetic labeling of
***Tacr1*-expressing spinal projection neurons.**
(**A**) Schematics of the triple cross strategy to specifically label *Tacr1*-expressing neurons in the spinal dorsal
horn. Scale bar indicates 200 µm.
(**B**) Spinal *Tacr1*-expressing neurons send projections to the posterior complex of the thalamus (PO),
ventral posterolateral nucleus of the thalamus (VPL), the ventral posteromedial nucleus of the thalamus
(VPM), SPFp, superior colliculus (SC), periductal gray (PAG), PB, dorsal column nuclei (DCN) and
ventrolateral medulla (VLM). Scale bar indicates 500 µm.
1392

1393

Figure S2. Projections from CGRP^SPFp and CGRP^PBel neurons.

(**A**) Additional images from Figure 1A, B. Projections are prominent in the amygdala regions and have distinct patterns along the anterior-posterior axis; AP: -1.1, -1.5 and -2.0 mm from bregma. Scale bar indicates 500 µm.

(**B**) The CGRP^PBel neurons also project to BNST, VPMpc, PSTN, and ventral pIC. The CGRP^SPFp neurons project to the auditory cortex, and dorsal pIC. Scale bars indicate 500 µm for BNST, and VPMpc; 200 µm for pIC.

1401

1402

Figure S3. Retrograde tracing from CGRP^SPFp and CGRP^PBel neurons.

(**A** and **B**) Example schematic of brain regions that send inputs to CGRP^SPFp (**A**) and CGRP^PBel neurons (**B**).

Scale bars indicate 100 μm.

(**C**) Percentage of total projections from brain regions to CGRP^SPFp neurons.

(**D**) Percentage of total projections from brain regions to CGRP^PBel neurons.

(**E** and **F**) Diagram of the projection (%) from other brain regions to CGRP^SPFp (**E**) and CGRP^PBel neurons (**F**).

CTX: cortex, Amy: amygdala, CP: striatum, GP: globus pallidus, BNST: bed nuclei of the stria terminalis,
SI: substantia innominate, HPF: hippocampus, HY: hypothalamus, TH: thalamus, VTA: ventral tegmental
area, SNr: substantia nigra, PRT: pretectal region, PAG: periaqueductal gray, SC: superior colliculus, IC:
inferior colliculus, MRN: midbrain reticular nucleus, Pons: including the nucleus of the lateral lemniscus,
pontine central gray, PBN, pontine reticular nucleus, VN: vestibular nucleus, NTS: nucleus tractus solitarius,
SpV: trigeminal spinal nucleus, CBX: cerebellum.
1416

1417

Figure S4. CGRP^{SPFp} and CGRP^{PBel} neurons are bilaterally stimulated by noxious stimuli.

(**A and B**) Maximum calcium responses of the CGRP^{SPFp} and CGRP^{PBel} neurons to ipsilateral (**A**) and contralateral (**B**) mechanical stimulation.

(**C**) CGRP^{SPFp} displayed faster increase of the calcium responses (bigger initial slope) to the contralateral mechanical stimulation (300 g for CGRP^{SPFp} and 100 g for CGRP^{PBel} neurons).

(**D-G**) Calcium responses of CGRP^{SPFp} (**D** and **E**) neurons by ipsilateral or contralateral 300 g stimulation and CGRP^{PBel} (**F** and **G**) neurons by 100 g stimulation.

(**H and I**) Maximum calcium responses of the CGRP^{SPFp} and CGRP^{PBel} neurons to ipsilateral (**H**) and contralateral (**I**) thermal stimulation.

(**J**) Comparison of the initial slope of both neurons to contralateral 55°C stimulation.

(**K-N**) Calcium responses of CGRP^{SPFp} (**K** and **L**) neurons and CGRP^{PBel} (**M** and **N**) neurons by ipsilateral or contralateral 55°C stimulation.

Statistics

(A) Repeated measure two-way ANOVA showed significance in intensity X region interaction ($F_{(4, 136)}$ = 11.80, $p < 0.0001$), intensity ($F_{(4, 136)}$ = 17.66, $p < 0.0001$), but not in region ($F_{(1, 34)}$ = 2.075, p = 0.1589). SPFp and PBel were significantly different in 50 ($p < 0.05$), 100 ($p < 0.01$) and 300 g ($p < 0.001$) with Sidak's multiple comparisons test.

(B) Repeated measure two-way ANOVA showed significance in intensity X region interaction ($F_{(4, 136)}$ = 5.468, p = 0.0004), intensity ($F_{(4, 136)}$ = 18.46, $p < 0.0001$), but not in region ($F_{(1, 34)}$ = 1.528, p = 0.2249).

(C) SPFp: 4.82 ± 1.08 (n = 6 mice, 12 trial), PBel: 1.68 ± 0.26 (n = 6 mice, 24 trial). Unpaired t test (two-tailed), p = 0.0007.

(E) SPFp; Ipsi: 8.51 ± 3.00 (n = 6 mice, 12 trial), Contra: 15.06 ± 4.51 (n = 6 mice, 12 trial). Paired t-test (two-tailed), p = 0.0498.

(G) PBel; Ipsi: 8.79 ± 0.95 (n = 6 mice, 24 trial), Contra: 15.26 ± 1.61 (n = 6 mice, 24 trial). Paired t-test (two-tailed), p = 0.0012.

(H) Repeated measure two-way ANOVA showed significance in intensity X region interaction $(F_{(3, 102)}$
$= 8.995$, $p < 0.0001$), intensity $(F_{(3, 102)} = 17.79$, $p < 0.0001$), and region $(F_{(1, 34)} = 11.19$, $p = 0.002$).

SPFp and PBel were significantly different in 55 ^{o}C ($p < 0.0001$) with Sidak's multiple comparisons test.

(I) Repeated measure two-way ANOVA showed significance in intensity X region interaction $(F_{(3, 102)} =$
10.16, $p < 0.0001$), intensity $(F_{(3, 102)} = 40.99$, $p < 0.0001$), but not in region $(F_{(1, 34)} = 2.885$, $p = 0.0986$).

SPFp and PBel were significantly different in 45 ($p < 0.05$) and 55 ^{o}C ($p < 0.0001$) with Sidak's multiple
comparisons test.

(J) SPFp: 5.55 ± 1.59 (n = 6 mice, 12 trial), PBel: 2.72 ± 0.26 (n = 6 mice, 24 trial). Unpaired t test (two-
tailed), $p = 0.0210$.

(L) SPFp; Ipsi: 1.74 ± 1.12 (n = 6 mice, 12 trial), Contra: 10.15 ± 3.08 (n = 6 mice, 12 trial). Paired t-test
(two-tailed), $p = 0.0146$.

(N) PBel; Ipsi: 15.55 ± 2.71 (n = 6 mice, 24 trial), Contra: 21.24 ± 2.63 (n = 6 mice, 24 trial). Paired t-test
(two-tailed), $p = 0.1385$.

1459

1460

Figure S5. Bilateral activation of CGRP^{SPFp} and CGRP^{PBel} neurons by unilateral formalin injection in the same animal.

(**A**) Number of CGRP positive cells in the SPFp and PBel.

(**B** and **C**) The number of CGRP positive cells in each side of the SPFp (**B**) and PBel (**C**).

(**D** and **E**) The percentage of CGRP neurons co-expressing c-Fos in the SPFp (**D**) and PBel (**E**).

(**F** and **G**) Representative images of the SPFp (**F**) and PBel (**G**). Scale bars indicate 200 µm.

1467

1468

Statistics

(A) SPFp: 1255 ± 137.5, PBel 1545 ± 112.2 (n = 6 mice). Paired t-test (two-tailed), p = 0.0188.

(B) SPFp; Ipsi: 315.5 ± 44.94, Contra: 311.8 ± 27.85 (n = 6 mice). Paired t-test (two-tailed), p = 0.9057.

(C) PBel; Ipsi: 373.0 ± 35.34, Contra 399.3 ± 24.06 (n = 6 mice). Paired t-test (two-tailed), p = 0.2958.

(D) SPFp; Repeated measure two-way ANOVA showed significance in treatment X side interaction (F(1, 4) = 17.42, p = 0.0140), treatment (F(1, 4) = 13.28, p = 0.0219), and side (F(1, 4) = 33.97, p = 0.0043). Ipsi vs contra was significant in formalin group (p < 0.01). Control vs formalin was significantly different in both ipsi (p < 0.05) and contra (p < 0.01) with Sidak's multiple comparisons test.

(E) PBel; Repeated measure two-way ANOVA showed significance in treatment X side interaction (F(1, 4) = 55.27, p = 0.0017), treatment (F(1, 4) = 19.57, p = 0.0115), and side (F(1, 4) = 58.66, p = 0.0016). Ipsi vs contra was significant in formalin group (p < 0.001). Control vs formalin was significantly different in both ipsi (p < 0.05) and contra (p < 0.01) with Sidak's multiple comparisons test.

1481

1482

1483 **Figure S6. Cued fear conditioning with mice for CGRP[SPFp] and CGRP[PBel] fiber photometry.**

1484 (**A**) Behavioral scheme for cued fear conditioning. Low intensity (70 dB, 3kHz) tone was used as CS in

1485 order not to induce calcium activity by sound.

1486 (**B** and **C**) Freezing was induced in both CGRP[SPFp] (**B**) and CGRP[PBel] (**C**) group.

1487

1488

1489 **Statistics**

1490 (B) SPFp; CS-: 2.47 ± 2.04 %, CS+: 38.96 ± 4.02 % (n = 6). Paired t test (two-tailed), p < 0.0001.

1491 (C) PBel; CS-: 11.27 ± 4.14 %, CS+ 51.31 ± 11.00 % (n = 6). Paired t test (two-tailed), p = 0.01.

1492

1493

Figure S7. The CGRP^{SFFp} and CGRP^{PBel} neurons are differentially activated by multiple sensory threat cues.

(**A**-**E**) Calcium response (peak amplitude) in the CGRP^{SFFp} and CGRP^{PBel} neurons by 2-s electric foot shock (0.6 mA) (**A**), 85-dB intense sound (**B**), rapidly expanding looming disk (**C**), TMT (**D**), and 0.5 mM quinine solution (**E**).

Statistics

(A) SPFp: $9.17 \pm 2.25\%$ (n = 6), PBel: $5.70 \pm 1.03\%$ (n = 6). Unpaired t test (two-tailed), p = 0.1917.

(B) SPFp: $5.84 \pm 1.17\%$ (n = 6), PBel: $3.82 \pm 1.39\%$ (n = 7). Unpaired t test (two-tailed), p = 0.2987.

(C) SPFp: $6.48 \pm 1.31\%$ (n = 6), PBel: $2.81 \pm 0.96\%$ (n = 7). Unpaired t test (two-tailed), p = 0.0418.

(D) SPFp: $-2.80 \pm 1.68\%$ (n = 6), PBel: $2.68 \pm 1.07\%$ (n = 6). Unpaired t test (two-tailed), p = 0.0206.

(E) SPFp: $-0.27 \pm 0.66\%$ (n = 6), PBel: $2.24 \pm 0.79\%$ (n = 7). Unpaired t test (two-tailed), p = 0.0363.

1507
Figure S8. Manipulation of CGRP^{SPFp} neurons in behavior assays.
(**A**, and **B**) Hot plate assay (55°C) with CGRP^{SPFp} silenced mice.
(**C**, and **D**) Automatic von Frey assay with CGRP^{SPFp} silenced mice.
(**E**) Experimental design for contextual fear conditioning.

(**F**) Quantification of freezing 24 hr after contextual fear conditioning.

(**G**) Schematic diagram of the elevated plus maze (EPM) test.

(**H**) The EPM tests in CGRP^{SPFp} (left panel)- or CGRP^{PBel} neurons (right panel)-silenced mice.

(**I**, and **J**) Optogenetic stimulation during hot plate assay (55°C).

(**K**, and **L**) Optogenetic stimulation during electric von Frey assay.

(**M**) Schematic diagram of optogenetic fear conditioning (photostimulation was used instead of foot shock).

(**N**) Quantified freezing level 24 hr after optogenetic conditioning.

(**O**) Schematic diagram of optogenetic cue-dependent fear conditioning (photostimulation paired with 3

kHz tone).

(**P**) Context-dependent freezing 24 hr after optogenetic conditioning.

(**Q**) Cue-dependent freezing 24 hr after optogenetic conditioning.

1523

1524

Statistics

(B) EYFP: 5.95 ± 0.33 s (n = 15), TetTox: 7.82 ± 0.73 s (n = 16). Unpaired t test (two-tailed), p = 0.0306

(D) EYFP: 4.75 ± 0.24 g (n = 15), TetTox: 5.66 ± 0.33 g (n = 16). Unpaired t test (two-tailed), p = 0.0113

(F) EYFP: 35.61 ± 3.18 % (n = 9), TetTox: 23.84 ± 3.17 % (n = 8). Unpaired t test (two-tailed), p = 0.0196

(H) SPFp; EYFP: 26.96 ± 3.66 % (n = 7), TetTox: 39.83 ± 4.05 % (n = 9). Unpaired t test (two-tailed), p =

0.0383.

PBel; EYFP: 34.56 ± 7.37 % (n = 7), TetTox: 57.58 ± 5.15 % (n = 7). Unpaired t test (two-tailed), p =

0.0250.

(I) Repeated measure two-way ANOVA showed no significance in interaction (F (1, 16) = 1.79,   p   =

0.2002), Laser (F (1, 16) = 3.36, p = 0.0857) and group (F (1, 16) = 1.49, p = 0.2393).

(L) Repeated measure two-way ANOVA showed no significance in interaction (F (2, 36) = 1.72, p =

0.1938), Laser (F (2, 36) = 0.06, p = 0.9465) and group (F (1, 18) = 1.26, p = 0.2768).

(N) EYFP: 25.31 ± 2.18 % (n = 8), ChR2: 47.95 ± 5.42 % (n = 12). Unpaired t test (two-tailed), p = 0.0042.

(P) EYFP: 15.06 ± 3.20 (n = 8), ChR2: 37.55 ± 8.13 (n = 12). Unpaired t test (two-tailed), p = 0.044.

(Q) Repeated measure two-way ANOVA showed significance in CS x group interaction (F (1, 18) = 8.072,

p = 0.0108), CS (F (1, 18) = 39.57, p < 0.0001) and group (F (1, 18) = 6.827, p = 0.0176). Freezing at CS-

and CS+ in ChR2 group (p < 0.0001) and CS+ in EYFP and ChR2 (p < 0.01) were significantly different

with Sidak's multiple comparisons test.

1543

Figure S9. Mapping the functional downstream of CGRP^SPFp projection.

(**A**) Schematics of the experiment.

(**B**) Representative image of the projection regions from CGRP^SPFp neurons. Scale bar indicates 500 µm.

(**C**) Example traces of an optically induced EPSC (blue) and IPSC (red) of Astr and LA without or with CNQX to confirm glutamatergic synapse. Scale bars indicate 10 ms and 50 pA.

(**D**) Examples of Astr EPSC (blue) and IPSC (red) traces by optogenetic terminal activation. Scale bars indicate 10 ms and 50 pA.

(**E**) Proportion of Astr cells with 'Both' EPSC and IPSC, 'EPSC only', 'IPSC only', or 'No-Response'.

(**F**) Onset of each Astr EPSC and IPSC to optogenetic stimulation.

(**G-L**) Results of the same experiments with LA (**G-I**) and pIC (**J-L**) neurons. Scale bars indicate 10 ms and 50 pA.

(**M**) Result of RTPA with terminal photo-stimulation.

(**N**) Context-dependent freezing at 24 h after terminal photo-stimulation conditioning.

(**O**) Cue-dependent freezing at 24 h after terminal photo-stimulation conditioning.

Statistics

(F) Astr; EPSC: 3.40 ± 0.25 ms (n = 12 cells), Astr IPSC: 7.74 ± 1.21 ms (n = 7 cells). Unpaired t test (two-tailed), p = 0.0003.

(I) LA; EPSC: 3.30 ± 0.16 ms (n = 21 cells), LA IPSC: 11.24 ± 1.13 ms (n = 17 cells). Unpaired t test (two-tailed), p < 0.0001.

(L) pIC; EPSC: 3.63 ± 0.49 ms (n = 15 cells), IC IPSC: 8.49 ± 0.83 ms (n = 8 cells). Unpaired t test (two-tailed), p < 0.0001.

(M) Repeated measure two-way ANOVA showed significance in laser X group interaction ($F_{(2, 29)}$ = 7.215, p = 0.0029) and laser ($F_{(1, 29)}$ = 13.24, p = 0.0011, but not group ($F_{(2, 29)}$ = 2.607, p = 0.0910). Laser ON and was significantly larger than OFF in AStr (p < 0.001) and pIC (p < 0.05). Additionally, difference between EYFP vs Astr (p < 0.01) and EYFP vs pIC (p < 0.05) during ON period were significant.

(N) EYFP: 3.842 ± 0.89 (n = 13), AStr: 14.82 ± 3.32 (n = 8), pIC: 17.94 ± 6.40 % (n = 9). One-way ANOVA showed significant (p = 0.0161). EYFP vs IC was significantly different (p < 0.05) with Tukey's multiple comparison test.

(O) Repeated measure two-way ANOVA showed significance in CS X group interaction ($F_{(2, 27)}$ = 3.647, p = 0.0396) and CS ($F_{(1, 27)}$ = 39.74, p < 0.0001) but not in group ($F_{(2, 27)}$ = 2.948, p = 0.0695). CS+ was significantly larger than CS- in AStr (p < 0.05) and pIC (p < 0.0001). Moreover, difference between EYFP and pIC in CS+ was significant (p < 0.01) with Sidak's multiple comparison test.

1577
1578 **Figure S10. Summary illustration**
1579