An Integrated Workflow for Enhanced Taxonomic and Functional Coverage of the Mouse Faecal Metaproteome

Nicolas Nalpas[1], Lesley Hoyles[2,3], Viktoria Anselm[1], Tariq Ganief[1], Laura Martinez-Gili[2], Cristina Grau[4], Irina Droste-Borel[1], Laetitia Davidovic[5], Xavier Altafaj[4,6], Marc-Emmanuel Dumas[2,7,8], Boris Macek[1]

[1] Proteome Center Tuebingen, University of Tuebingen, Germany; [2] Biomolecular Medicine Section, Division of systems Medicine, Department of Metabolism, Digestion and Reproduction, Imperial College London, Sir Alexander Fleming building, London SW7 2AZ, UK; [3] Department of Biosciences, Nottingham Trent University, UK; [4] Bellvitge Biomedical Research Institute, Spain; [5] Université Côte d'Azur, CNRS, Inserm, IPMC, France; [6] Neurophysiology Unit, University of Barcelona - IDIBAPS, Spain; [7] Genomic and Environmental Medicine, National Heart & Lung Institute, Faculty of Medicine, Imperial College London, London, SW3 6KY, United Kingdom. [8] European Genomic Institute for Diabetes, INSERM UMR 1283, CNRS UMR 8199, Institut Pasteur de Lille, Lille University Hospital, University of Lille, 59045 Lille, France.

Short title: Enhanced workflow for metaproteomics.

Keywords: Metaproteomics; Microbiome; *Mus musculus*; Mass spectrometry; Proteogenomics.

*To whom correspondence should be addressed:
Prof. Dr. Boris Macek
Proteome Center Tuebingen
Interfaculty Institute for Cell Biology
Auf der Morgenstelle 15
72076 Tuebingen
Germany
Phone: +49/(0)7071/29-70558
Fax: +49/(0)7071/29-5779
E-Mail: boris.macek@uni-tuebingen.de

Abstract

Intestinal microbiota plays a key role in shaping host homeostasis by regulating metabolism,

immune responses and behaviour. Its dysregulation has been associated with metabolic,

immune and neuropsychiatric disorders and is accompanied by changes in bacterial metabolic

regulation. Although proteomics is well suited for analysis of individual microbes,

metaproteomics of faecal samples is challenging due to the physical structure of the sample,

presence of contaminating host proteins and coexistence of hundreds of microorganisms.

Furthermore, there is a lack of consensus regarding preparation of faecal samples, as well as

downstream bioinformatic analyses following metaproteomics data acquisition. Here we assess

sample preparation and data analysis strategies applied to mouse faeces in a typical mass

spectrometry-based metaproteomic experiment. We show that subtle changes in sample

preparation protocols may influence interpretation of biological findings. Two-step database

search strategies led to significant underestimation of false positive protein identifications.

Unipept software provided the highest sensitivity and specificity in taxonomic annotation of

the identified peptides of unknown origin. Comparison of matching metaproteome and

metagenome data revealed a positive correlation between protein and gene abundances.

Notably, nearly all functional categories of detected protein groups were differentially

abundant in the metaproteome compared to what would be expected from the metagenome,

highlighting the need to perform metaproteomics when studying complex microbiome

samples.

Introduction

The prokaryotic component of the gut microbiota has multiple roles, contributing to carbohydrate fermentation and maintenance of gut barrier integrity, as well as antimicrobial and immunomodulation activities [1,2]. In metabolically healthy humans and mice, the gut microbiota is predominated by two to three bacterial enterotypes [3-5]. These enterotypes display significant heterogeneity in terms of species number, composition and relative abundances depending on the location of the sample (upper vs lower gastroinstestinal tract) or the timing (circadian variations) [6,7]. The gut microbiota has recently been associated with conditions ranging from inflammatory bowel syndrome to Parkinson's disease [8-11]. An increasing number of studies have reported associations between the gut microbiota and neurodevelopmental disorders [12-14]. This includes changes in the gut microbiota of Down syndrome individuals in comparison to non-trisomic individuals [15]. Given the established interaction between the host and the gut microbiota, a functional analysis of the gut microbiome may help in understanding its contribution to pathophysiology.

In this context, approaches relying on nucleotide sequencing have so far been preferred by the scientific community due to lower experimental costs, higher data throughput and proven analytical workflows. While metagenomics assesses the genetic potential, metaproteomics investigates gene products (and therefore functions). However, metagenomics usually provides more in-depth information in comparison to metaproteomics, for example due to the higher dynamic range of detection. In particular, microbiome functional analysis can be performed using high-resolution mass spectrometry (MS), to measure either protein abundance or metabolite production [16-18]. Although bacterial MS-based proteomic approaches are well established, metaproteomic sample preparation is hindered by many challenges, such as physical structure of the sample, the presence of contaminating proteins and the coexistence of hundreds of microorganisms.

Many studies have reported increased protein identification due to laboratory optimisation for

the analysis of metaproteome samples [19-23]. In humans, different sample preparation

methodologies have been shown to result in significant changes in the taxonomic composition

and functional activities represented [19,24,25]. Beyond sample preparation, the

bioinformatic processing of metaproteomic data remains challenging, due to the choice of

representative protein sequence database, elevated false discovery rate for peptide

identification and the redundancy in protein functional annotation. Some of these challenges

have already been addressed by published software packages, such as MetaProteomeAnalyzer

[26] and MetaLab [27], which are all-in-one metaproteomic analytical workflows, or UniPept

[28], which allows peptide-based taxonomic representation. In addition, the choice of protein

sequence database has been shown to play a major role in protein identification from

metaproteome samples, with notably matching metagenome-derived protein sequence

databases displaying the best identification rate performance [29-32]. Previous studies have

also investigated ways to determine taxonomic representation from metaproteome samples,

which has been shown to differ between metagenome (bacterial presence) and metaproteome

(bacterial activity) [32,33].

Here, we present a state-of-the-art MS-based workflow for the optimal metaproteome

characterisation of murine faecal samples. We focused on a number of aspects that remain

under-investigated in murine stool samples: (1) the impact of sample preparation methods,

namely low speed centrifugation (LSC) and no LSC (nLSC), on protein identification and

taxonomic representation; (2) the high false positive rates in searches involving very large

databases; (3) the differences in taxonomic annotation of MS-identified peptides based on

different software; and (4) the lack of assessment of the functional enrichment provided by the

metaproteome compared to its matching metagenome potential.

Results

Low-speed centrifugation increases peptide identification rates

Our initial experiment involved the establishment of an optimal sample preparation workflow applied to the mouse faecal metaproteome. In this context, we assessed two critical steps within our sample preparation method: 1) the usage of LSC; and 2) in-solution digestion versus filter-aided sample preparation (FASP) (**Figure S1A**, **Table S1**).

The number of peptide spectral match (PSM) identified per MS raw file in the LSC group was significantly higher with 26 % more identifications (**Figure 1A**). This was also observed at the peptide and protein group level, but to a lower extent for the latter. Approximately 15 % of protein groups were identified by a single peptide, while the median protein sequence coverage was 18.7 %. Such metrics are usually indicative of highly complex samples that are not completely covered by a single MS measurement under the stated parameters.

In-solution digestion consistently outperformed FASP based on PSMs, peptides and protein groups identification (**Figure 1B**). Compared to other methods, in-solution digestion combined with the LSC procedure provided nearly twice as many PSM or peptide identifications and 30 % more protein groups. Furthermore, there was much less variability in the number of peptides and protein groups identified across samples with this method.

LSC aids in recovery of *Bacteroidetes* proteins, whereas nLSC favours *Firmicutes* and *Deferribacteres* proteins

Peptides identified after LSC and nLSC were analysed to identify their phylogenetic origin. The lowest common ancestor was determined using the Unipept interface [34], which assigns peptide sequences to taxa. The most abundant superkingdom consisted of bacteria, among which two taxa were highly represented in both LSC and nLSC, namely *Bacteroidetes* and *Firmicutes* (**Figure 1C**, **Table S1**). However, there were large differences in the number of

peptides assigned to these two main bacterial phyla when comparing LSC and nLSC methods.

Bacteroidetes accounted for 66 % and 37 % of peptides, whereas *Firmicutes* amounted to 18 %

and 47 % of peptides in LSC and nLSC procedures, respectively. In addition, *Actinobacteria*

and *Deferribacteres* showed a higher taxonomic representation in nLSC compared to LSC,

whereas *Verrucomicrobia* showed an opposite trend.

Based on peptides identification, Eukaryota was the second most abundant superkingdom and

consisted mostly of metazoan hits. Under the assumption that these eukaryotic peptide

sequences originated from the host, the proportion of *Mus musculus* proteins was investigated

further using intensity-based absolute quantification (iBAQ) values. The LSC samples

contained on average nearly two-fold more murine proteins (20.4 %) in comparison to nLSC

samples (14.6 %) (**Figure S1B**). Such findings were surprising since the use of the LSC method

was reported in a previous study to help with the removal of human cells [24]. We also

investigated the presence of peptides from host diet and found very low levels of dietary

peptides contamination (approximately 2 %), which was higher among LSC-prepared samples

(**Figure S1C**). As previously reported, we show that the majority of dietary proteins are absent

or depleted during the initial solubilisation step of the faecal pellet, a step common to both

procedures [35]. Overall, our results show that LSC and nLSC methods favour the recovery of

different taxa, suggesting that both methods have merits and may be used in combination.

LSC and nLSC methods are characterised by different protein abundance profiles

We further investigated the overlap between the peptides or protein groups identified following

either LSC and nLSC procedures (**Figure 1D**). In terms of peptides, only 27.7 % were

identified with both procedures, the rest of the peptides being split equally into unique to LSC

and nLSC methods. Similar results were observed at the protein groups level with 38.7 % of

protein groups being identified in both procedures. This was illustrated further through a

principal component analysis (PCA), showing separation of samples based on centrifugation

methods, as well as clustering of technical replicates (from cell lysis step) (**Figure S1D**). Label-

free quantitative (LFQ) comparison between LSC and nLSC procedures revealed an

intermediate correlation ($\rho = 0.44$) (**Figure S1E** and **F**). Besides, LFQ correlation among the

samples prepared via LSC was superior to samples prepared with nLSC (**Figure S1G**). Our

findings indicate that while the two procedures have a poor identification overlap, the main

differences may still result from biological variations.

Using LFQ intensities, we then performed a *t*-test to identify which protein groups have

different abundances between the two procedures. Out of 2,589 quantified protein groups, 365

and 267 showed a significant increase and decrease in abundance between LSC and nLSC

samples, respectively (FDR ≤ 0.01 and absolute fold-change ≥ 2.5) (**Figure 1E**, **Table S1**). We

gained functional insights into these differences by performing an over-representation analysis

of KEGG pathways. The over-represented pathways based on the up- or down-regulated

protein groups were mostly similar (FDR ≤ 0.05) and were associated with core microbial

functions, such as ribosome, carbon metabolism and carbon fixation pathways (**Figure 1F**,

Table S1). The protein groups unique to LSC or nLSC showed over-representation of protein

export in the LSC samples, whereas biosynthesis of amino acid, fatty acid degradation and

bacterial chemotaxis were over-represented in the nLSC samples (**Figure S1H**). Protein

differential abundance testing confirmed the divergence between LSC and nLSC procedures

and was suggestive of broad taxonomic changes, rather than variation in functional activities.

Two-step database search strategy shows a dramatic increase in false positive rate

After measurement via liquid chromatography coupled to tandem mass spectrometry (LC-

MS/MS) and acquisition of LC-MS/MS raw data, the MS/MS spectra are searched against a

protein sequence database. One aspect of database search is the controversial use of a two-step

search strategy, whereby LC-MS/MS measurements are processed initially against a large

protein sequence database with no FDR control (FDR $\leq 1\%$). Subsequently, the original

database is filtered to retain only protein sequences that were identified during the first search.

During the second database search, the measurements are processed against the reduced

database with FDR control (e.g. FDR ≤ 0.01) [36]. To assess these search strategies, we

searched a single HeLa cell LC-MS/MS file using MaxQuant software against a *Homo sapiens*

protein sequence database supplemented with different number of bacterial protein sequences

(**Figure S2A**). The HeLa measurement is used here as a proxy for a complex microbiome

measurement, with the exception that the sample composition is known and from a single

organism.

We initially established a benchmarked standard by processing the HeLa measurement only

against an *H. sapiens* database, which resulted in approximately 5,000 human (eukaryota)

protein groups identified for the single-step search at FDR ≤ 0.01 (**Figure 2A**, **Table S2**).

Notably, the same database used in a two-step search identified less than 1 % additional protein

groups in comparison to a single-step search, despite nearly twice as much processing time.

We then processed our HeLa measurement against the *H. sapiens* database supplemented with

1×, 2×, 5×, 10× and 20× bacterial protein sequences, resulting in increasingly large databases

(**Figure S2A**, **Table S2**). For the single-step database search against the 1:20 database, we

observed a 10 % decline in the number of human protein groups identified, while 132 bacterial

protein groups were identified (false positives). On the contrary, the 1:20 two-step database

search resulted only in a 1 % decrease compared to the benchmarked standard. This processing

also revealed a large number of bacterial protein groups identification (980 protein groups).

Furthermore, the two-step search led to large number of MS/MS spectra to be assigned to

different sequences (or newly assigned) in comparison to the benchmarked standard (**Figure**

S2B, **Table S2**); this phenomenon was much less pronounced when performing the single-step

search.

We then calculated the factual FDR for each processing approach using either the reverse hits

or the reverse hits plus the bacterial hits (which in our case are false positives). For both the

single-step and the two-step search, we obtained an FDR of 2.6 % when using only the reverse

hits for FDR calculation (**Figure 2B**). However, when using the reverse hits plus the bacterial

hits, we calculated a factual FDR of 8 % and 34 % for the single- and two-step search with

1:20 database, respectively. This represents a dramatic increase in the rate of false positive

identification when using two-step search, despite controlling for 1 % FDR. Notably, these

false positive hits would remain unnoticed in a microbiome sample of unknown composition,

thus highlighting the inherent problem associated with the two-step database search.

Optimisations of the two-step database search cancels out its higher sensitivity

To further assess database search strategies in a metaproteomic context, we retrieved a

metaproteome dataset of known taxonomic composition that was published by Kleiner and

colleagues [32]. This dataset consisted of 32 organisms of uneven abundances, including

bacteria (25), archaea (1), eukaryotes (1) and viruses (5). We processed eight LC-MS/MS

measurements against a database containing the proteomes of these 32 organisms, which we

supplemented with 0.5×, 1×, 2×, 5×, 10× and 20× bacterial protein sequences, resulting in

increasingly large databases. We then compared the results obtained from single-step search

strategy against: (1) "two-step protein" search to keep identified proteins [36]; (2) "two-step

taxa" search to keep identified taxa [30]; and (3) "two-step two sections" search to keep

identified proteins after sectioned search [37]. While all search strategies resulted in similar

accuracies, the "two-step protein" search maintained a high sensitivity even when using large

databases (i.e. 20×) (**Figure 2C** and **S2C**). However, upon investigation of the factual FDR

(reverse hits plus the false bacterial hits), the "two-step protein" search resulted in twice as

many false positive identifications compared to the single-step search (**Figure 2D**, **Table S2**).

Similar results were also observed when focusing on the precision (**Figure S2D**). Our

investigations revealed that the "two-step taxa" search behaved nearly identically to the single-

step search, whereas the "two-step two sections" search displayed performance in-between the

first-step and "two-step protein" searches.

Because, all assessed search strategies underestimated the real FDR, we attempted to identify

any particularity of the false positive protein groups identification and thus focused on

processings against the largest database (20×). We show that the median number of unique

peptides (i.e. peptides that are uniquely assigned to a protein group) are 1 and 2 for the false

and true positive hits, respectively (**Figure S2E**). We then compared results obtained using a

post-processing filtering step requiring a minimum of 1 or 2 unique peptides per protein groups.

Our results show that requiring a minimum of 2 unique peptides would efficiently control the

FDR ($\leq 1\%$) at the expense of a significant drop in protein identification (**Figure 2E** and **S2F**,

Table S2). This investigation of different database search strategies applied to metaproteome

samples further highlighted the limitations (i.e. factual FDR) of two-step searches, even

following optimisation (i.e. sectioned search) or filtering.

Unipept software provides the most accurate and precise taxonomic annotation

Another important aspect of metaproteomic studies is the determination of taxonomic activity

(protein biomass), which has been reported to differ from taxonomic representation derived

from metagenomic studies [32]. While it is straightforward to compute taxonomic activity from

the abundance of peptides (or proteins) of known taxonomic origin, there has not been an

exhaustive assessment of software that can taxonomically annotate MS-identified peptides.

Here, we assessed three software—i.e. Kraken2 [38], Diamond [39] and Unipept [28]—with

regards to their taxonomic annotation performance on the dataset from Kleiner and colleagues

[32]. The Kraken2 software provided consistently higher percentage of peptides that could be

taxonomically annotated (c.a. 18% peptides annotated to species level), followed by Unipept

(5%) and Diamond (1%) (**Figure 3A**). However, Kraken2 also identified a very large number

of taxa that were not present in the artificial samples from Kleiner and colleagues (**Table S3**)

and thus would be false positive hits. Unsurprisingly, these false positive hits were

characterised by low PSM counts in comparison to true positives (**Figure S3A**). This led us to

assess these software packages in terms of accuracy, precision, sensitivity, specificity and F-

measure for taxonomic identification using a range of PSM count thresholds (**Figure 3B** and

S3B-E). In this context, the Unipept software significantly outperformed Kraken2 and

Diamond, especially with regard to the F-measure and precision. Notably, the implementation

of a minimum PSM count threshold (i.e. between 1 and 5) resulted in accuracy, precision and

specificity improvements for all software, but at the cost of a reduced sensitivity.

Thus, without a PSM count threshold, we correlated the taxonomic abundance derived from

each software annotation against the known input protein from Kleiner and colleagues'

artificial samples (**Figure 3C**). Overall, the Unipept software provided the highest correlation

(Spearman $\rho = 0.83$), as well as at most taxonomic levels (including species). Interestingly, the

dynamic range of taxon detection by MS spanned two orders of magnitude, with *Salmonella*

enterica being approximately 230 times more abundant than *Nitrosomonas europaeae* (**Figure**

3D, **Table S3**). Unipept was also the only software allowing identification of *Nitrosomonas*

ureae, *Paraburkholderia xenovorans* and *Nitrosospira multiformis*. Importantly, none of the

software could identify the five viral organisms present in the samples, the reason being

technical since no peptide coming from those viral proteins was detected by MS. Finally, we

assessed the impact of different database search strategies on taxonomic abundance derived by

the Unipept software (**Figure S3F**). Similarly to our findings from the previous section, the F-

measure metric highlighted the superiority of single-step strategy when it comes to taxonomic

identification. Taken together, we show that, based on different metrics and samples of known

composition, the Unipept software provides better taxonomic annotation in comparison to

Kraken2 and Diamond.

The complex microbial composition of faecal samples is best recapitulated by the Unipept software

To check whether our results are also applicable to the microbial composition of faecal samples, we prepared samples using the LSC method from faeces collected in a cohort of 38 mice. The resulting LC-MS/MS data were processed using a single-step search strategy against a matching metagenome protein database (with no knowledge of taxonomic composition). We initially annotated the MS-identified peptides using Kraken2, Diamond and Unipept, which revealed an overlap of 232 taxon (1.9%) between all three software. Such low overlap was largely driven by the suspected large number of false positive hits identified by Kraken2 (10,203 uniquely identified taxon), as seen in the previous section. We then performed pairwise correlation between every samples combination within each software using taxonomic abundance (**Figure S3G**). While Diamond displayed a higher correlation (median spearman ρ = 0.71), this is likely driven by the small number of identified taxa, most of which at the taxonomic levels closer to the root (e.g. superkingdom, phylum) and is thus a poor performance estimate.

To determine which taxa are likely true or false positive hits, we made use of the taxonomic composition foreknowledge (at the species level only) from the mouse microbiome catalogue [40]. With this approach, the Kraken2 software showed the best sensitivity (median = 0.17) compared to Unipept (0.13) and Diamond (0.03) (**Figure S3H**, **Table S3**). Based on precision and F-measure, Kraken2 performance collapsed, whereas Unipept software had median precision and F-measure superior to 0.1 (**Figure 3F** and **S3I**). Using taxonomic foreknowledge, our findings suggest that the Unipept software provides superior predictive power for taxonomic annotation of faecal samples.

Metaproteome to metagenome correlation highlights an over-representation in the core microbiome functions

Due to the availability of matching metagenomic and metaproteomic data for our cohort of 38 mice, we assessed the correlation between gene and protein abundances. To deal with the intrinsic differences between the two datasets, the gene entries were grouped in a similar fashion as the protein groups (i.e. based on peptide identification) and the maximum expression was calculated per gene group. Here, we show that a majority of gene-protein pairs (91 %) have a positive correlation, with a median of 0.39, the rest having a median negative correlation of -0.09 (**Figure 4A**, **Table S4**). Notably, 3,519 gene-protein pairs displayed a significant positive correlation. In addition, we compared the distribution in gene abundances depending on whether the corresponding protein was identified by MS (**Figure S4A**). As expected, it shows that MS-based proteomics only identifies a subset of proteins towards the higher abundance.

To identify the core pathways within our mice cohort, we performed an over-representation analysis of the significantly correlated gene-protein pairs (**Figure 4B**, **Table S4**). Among these pairs, there was an over-representation in carbon fixation, glycolysis-gluconeogenesis, citrate cycle and carbon metabolism pathways (KEGG) [41]. We further characterised the correlating genes and proteins and identified 20 over-represented gene ontology molecular functions (GOMF) that were involved in ADP, ribosome, carbohydrate and electron transfer (**Figure S4B**, **Table S4**). Our results confirm the central role of carbon fixation and general metabolism, which are associated with bacterial energy production, in the murine faecal microbiome under the analysed conditions.

The metaproteome is enriched in functionally active pathways compared to the matching potential encoded in the metagenome

The metagenome corresponds to the microbiome's genetic potential, whereas the metaproteome represents its truly expressed functional activities. Thereby, we compared the functional abundance derived from the metagenomic versus metaproteomic datasets within our cohort of 38 mice. To allow comparison, the KEGG level 2 categories were quantified and normalised separately for each omic datasets (**Figure S4C**, **Table S4**). Out of 55 KEGG categories, we found 15 and 37 to be significantly increased and decreased in abundance at the metaproteome level in comparison to the metagenome (FDR $\leq$ 0.05). In general, the metagenome-based quantification of KEGG categories was stable across categories, whereas large differences were observed for the metaproteome.

To prioritise the KEGG categories, we selected eight categories differing significantly in terms of gene-protein correlation in comparison to the overall correlation (**Figure 4C** and **S4D**). Among the KEGG categories displaying higher abundance in the metaproteome compared to the metagenome were the membrane transport, translation, signalling and cellular processes, and genetic information processing. Conversely, transcription, carbohydrate metabolism and antimicrobial drug resistance exhibited lower abundance. The KEGG Orthology (KO) entries differing significantly in abundance between the metagenomes and metaproteomes were identified via *t*-test and used for gene set enrichment analysis (GSEA). GSEA revealed an enrichment of a number of overlapping KEGG pathways, with 19 and 6 pathways positively and negatively enriched, respectively (**Figure 4D**, **Table S4**). Interestingly, we found the ribosome pathway enriched in protein with increased abundance (between metaproteome and metagenome datasets), therefore highlighting the functional activation of this pathway (**Figure 4E** and **S4E**). Conversely, homologous recombination, DNA replication and mismatch repair were enriched in protein with decreased abundance, suggesting no or low activation of these

pathways. Overall, our findings highlight the critical importance of metaproteomics to

characterise microbiome samples particularly when it comes to their functional activity.

Discussion

Here, we investigate some key aspects of metaproteomic workflow applied to murine faecal

samples in order to enhance protein identification, taxonomic and functional coverage. We

focused on the assessment of (1) different sample preparation methods, (2) strategies to control

for false positive rates during database search, (3) taxonomic annotation software for accurate

MS-derived taxonomic representation and (4) the importance of metaproteomics to determine

functionally enriched pathways. Our results led to an overview of the strengths and weaknesses

of each assessed methods (**Table 1**) in the context of murine faecal metaproteomics.

To the best of our knowledge this is one of the largest and most extensive comparisons

undertaken to date, comprising over 40 different biological samples and over 50 LC-MS/MS

runs. Overall, we reached identification rates that are similar to bacterial shotgun proteomics

(ca. 20-40 %). In comparison to previous murine faecal metaproteomic studies, we identified

more non-redundant peptides per samples (approximately 20,000 non-redundant peptides on a

60 min gradient) [42,43]. Several parameters may have influenced such performance, among

which are the use of a faster and more sensitive Orbitrap instrument (i.e. Q Exactive HF)

[44,45] and a more representative protein sequence database (i.e. mouse metagenome catalogue

or mouse matching metagenome) [40]. Importantly, the impact of mass spectrometer speed and

sensitivity should not be overlooked in a typical metaproteomic measurements. Indeed, the

type and model of MS instrument was among the parameters with the greatest impact on

identification rates. Some of our initial investigation showed significant increase in peptide and

protein identification rate when using the Q Exactive HF (faster scanning, improved sensitivity)

versus the Orbitrap Elite (data not shown, but downloadable from ProteomeXchange).

Both LSC and nLSC methods have merits for the metaproteomic analysis of murine

faecal samples

Our study confirms previous observation with regard to the depletion or enrichment of several

major bacterial phyla, which is dependent on laboratory preparation method and specifically

the usage of differential centrifugation [24]. In this context our results do not match with the

study from Tanca and colleagues, who reached opposite conclusions. There are several possible

explanations for these discrepancies, such as the host organism under study (*i.e. Mus musculus*

versus *Homo sapiens*) and different protein sequence database construction (*i.e.* mouse

microbiome catalogue versus UniProtKB custom microbiome). The LSC approach also leads

to more consistent identifications and as a result fewer missing values, which is a general and

extensive problem in metaproteomic datasets. Regarding the topic of reproducible protein

identification and quantification, a recent metaproteomic study demonstrated the use of

Tandem Mass Tag (TMT) approach in human stool samples [22].

Investigation of the differences in taxonomic composition between LSC and nLSC revealed

broad changes already at the phylum level. Notably, *Bacteroidetes* and *Verrucomicrobia* were

enriched within LSC-prepared samples, whereas *Firmicutes, Actinobacteria* and

Deferribacteres phyla were over-represented in nLSC samples. Additional comparison to the

phyla detected by metagenomics in the mouse microbiome catalogue study tends to agree more

with the nLSC approach [40]. However, the samples from that study were also prepared using

a nLSC approach, which may explain the similarity. Importantly, it has been reported that the

removal of faecal particles may also lead to exclusion of proteins or organisms attached to these

faecal debris [24], thus leading to a bias in the LSC approach. Our results at the protein level

showed significant changes in abundance, which were indicative of broad taxonomic changes,

more so than variation in functional activities. Importantly, recent studies have reported

considerable changes in rodent microbiota depending on suppliers or on shipping batch, even

for mice housed in identical environments [46,47]. Murine gut microbiota is also significantly

different from other mammals, such as human [40]. In this context, our results on

metaproteomic sample preparation may not translate to other of murine faecal pellets (e.g.

young vs. old individuals) or other mammalian faeces (e.g. *H. sapiens*) and would suggest a

required sample preparation optimisation for each cohort. To conclude, both sample

preparation approaches have advantages, and the choice may ultimately come down to which

bacterial organism is under-investigation [25].

Single-step database search allows optimal control of false discovery rate

Currently, many metaproteomic studies use two-step database searches as a way to boost

identification rates [36]. However, we demonstrate that this type of search dramatically

underrepresents the number of false positives, due to the use of a decoy search strategy that is

unsuitable in this context. Our results elaborate on a previous study by Muth and co-workers,

who also emphasised the drawbacks of using a two-step search together with decoy strategy

[48]. Using a single human LC-MS/MS measurement, our findings were so extreme that the

number of false positives was equal or greater to the number of false negatives, with FDR

outside of any accepted range (i.e. factual FDR > 0.1).

Using metaproteome samples of known composition, we expanded our investigation of search

strategies by including "two-step taxa" and "two-step two sections". The "two-step two

sections" approach, implemented according to Kumar and colleagues [37], provided a middle

ground in performance between the "two-step protein" and single-step search strategies, but at

the expanse of much longer processing time. Nonetheless, our results confirmed the inability

of two-step searches to control the FDR, including in context of metaproteomic samples. We

argue that the use of a two-step search should be avoided whenever possible and replaced by

alternative strategies, such as taxonomic foreknowledge or using matching metagenomes [49].

Accurate taxonomic annotation of murine faecal samples can be generated by the Unipept software

Previous studies have shown that it is possible to derive taxonomic representation from MS-identified peptides of known taxonomic origin [32,33,50]. However, to the best of our knowledge, there has not been a comparison of software for the taxonomic annotation of peptides with unknown origin. Here, we compared three software packages, namely Kraken2 [38], Diamond [39] and Unipept [28], which use different algorithms to perform such taxonomic annotation. Using metaproteome samples of known composition, as well as metaproteome samples from 38 murine faeces, we determined that the Unipept software provided superior performance (i.e. precision, sensitivity). Notably, Unipept is very user-friendly, fast and was designed to work on MS-identified peptides [28]. Whereas, Diamond and Kraken2 have both been designed to work on full protein/gene sequences or nucleotide sequencing reads (as opposed to peptides), which may have contributed to their lower performance [38,39]. Our assessments (i.e. sensitivity, specificity) were based on exact taxonomic identity and ignored hits from closely related taxon, which may have negatively affected the performance estimates of Kraken2 [38]. While, Unipept was clearly the optimal taxonomic annotation software for MS-identified peptides, it is currently limited to UniProt proteins, NCBI taxonomic hierarchy and trypsin cleavage.

The metaproteome shows an enrichment in functionally-active pathways compared to the matching metagenomic potential

Here, we observed an overall positive correlation between gene and protein abundances derived from metaproteome and matching-metagenome analysis. This was previously reported in a longitudinal study of metaproteome/metagenome fluctuations from one individual with Crohn's Disease [51]. In our case the significantly correlated entries were associated with core

bacterial metabolic functions, such as carbon and energy metabolism or electron transfer

activity [52]. Despite such correlations, we also reported extensive differences in quantified

functions between metagenomics and metaproteomics. Notably, with regard to genetic

information processing (KEGG level 2), the ribosome pathway was over-represented in entries

with higher abundance in metaproteomes, whereas pathways associated with DNA repair,

replication or recombination were over-represented in entries with increased abundance in

metagenomes. This greatly highlights the main advantage of metaproteomics, which captures

functionally active pathways, as opposed to the genetic potential represented by metagenomics

[53]. Thus, these approaches are complementary to each other and can provide a more

comprehensive understanding of a biological system.

Conclusion

To conclude, in this study we present an integrated analytical and bioinformatic workflow to

improve protein identification, taxonomic and functional coverage of the murine faecal

metaproteome. LSC combined with in-solution digestion provided the highest identification

rates, although leading to a potential enrichment in specific taxa. We also show that fast and

accurate MS data processing can be achieved using a single-step database search. Taxonomic

annotation can be generated directly from MS-based peptide identification using the Unipept

software. While protein and gene abundances displayed an overall positive correlation, the

metaproteome showed a significant functional enrichment compared to its metagenomic

potential; thus, emphasizing the need for more metaproteomic studies for adequate functional

characterisation of the microbiome.

Methods

Animals and faecal samples collection

Mouse faecal pellets obtained from a small cohort of six wild-type B6EiC3SnF1/J mice were used to compare sample purification and protein extraction methodologies from faeces (**Figure S1A**). A larger cohort of 38 mice (euploid and trisomic Ts65Dn) was used to obtain mouse faeces, for further assessment of the data analysis workflow. Mice were housed and faeces were collected following the experimental procedures evaluated by the local Ethical Committee (Barcelona Biomedical Research Park, Spain). Faecal pellets were collected fresh, placed at -20 °C and stored at -80 °C until analysis.

DNA extraction and whole-genome sequencing

Whole genome analysis was performed on the mouse cohort used for data analysis assessment. In brief, DNA was extracted from faecal samples using the FastDNA SPIN Kit (MP Biochemicals) and following manufacturer's instructions. DNA concentration was measured using a Qubit fluorometer (Invitrogen) and samples were shipped frozen to the Quantitative Biology Centre (QBiC) at the University of Tuebingen for whole genome sequencing. Sequence data were generated on an Illumina HiSeq 2500 instrument (chemistry SBS v3 plus ClusterKit cBot HS) and processed as described previously [54] but with minor modifications that follow. Supplied sequence data were checked using fastQC v0.11.5 [55]. Data were trimmed with Trim Galore! (--clip_R1 10 --clip_R2 10 --three_prime_clip_R1 10 --three_prime_clip_R2 10 --length 50; Babraham Bioinformatics). Mouse DNA within samples was detected by mapping reads against the mouse genome (GRCm38). Mouse-filtered read files (with an average of 3.58 ± 0.08 Gb sequence data per sample) were used for all subsequent analyses. Kraken2 2.0.8-beta [38] with the pre-compiled Genome Taxonomy Database [56] Functional annotation was achieved by mapping centroid protein sequences generated as

described before [38,54] using the eggNOG-mapper software (v.1.0.3) [57] and associated

database (v.4.5).

Sample treatment, cell lysis and protein extraction

Mouse faecal pellets obtained from wild-type B6EiC3SnF1/J mice were used to compare

sample initial preparation methodologies (**Figure S1A**).

For the LSC procedure, faeces (~50 mg) were resuspended in phosphate buffer (50 mM

Na_2HPO_4/NaH_2PO_4, pH 8.0, 0.1 % Tween 20, 35x volume per mg) by vortexing vigorously

for 5 min using 4 mm glass beads (ColiRollers[TM] Plating beads, Novagen), followed by

incubation in a sonication bath for 10 min and shaking at 1,200 rpm for 10 min in a

Thermomixer with a thermo block for reaction tubes. Insoluble material was removed by

centrifugation at $200 \times g$ at 4 °C for 15 min. The supernatant was removed and the remaining

pellet was subjected to two additional rounds of microbial cell extraction. After merging

supernatants, microbial cells were collected by centrifugation at $13,000 \times g$ at 4 °C for 30 min.

The pellet was resuspended in 80 µL sodium dodecyl sulfate (SDS) buffer (2 % SDS, 20 mM

Tris, pH 7.5; namely pellet extraction buffer) and heated at 95 °C for 30 min in a Thermomixer.

The resulting suspension was divided into two parts to obtain technical replicates for the rest

of the sample preparation workflow. Protein extraction was performed by cell homogenization

using 0.1 mm glass beads (100 mg, SartoriusTM Glass Beads) for each replicate and the

FastPrep-24 5G instrument (MP) at 4 m/s or BeadBug microtube homogenizer (BeadBug) at

4,000 rpm. Three cycles of homogenization including 1 min bead beating, 30 sec incubation at

95 °C, and 30 sec centrifugation at $13,000 \times g$ were performed. The homogenate was diluted

with 800 µL $MgCl_2$ buffer (0.1 mg/mL $MgCl_2$, 50 mM Tris, pH 7.5) and centrifuged at

13,000 rpm for 15 min. Proteins from the supernatant were precipitated overnight in acetone

and methanol at -20 °C (acetone:methanol:sample with 8:1:1 ratio). Protein pellets were

resuspended in 120 µL denaturation buffer (6 M urea, 2 M thiourea, 10 mM Tris, pH 8.0) for

downstream use.

For the nLSC procedure, mouse faeces (~25 mg) were homogenised directly in 150 µL pellet

extraction buffer as described above with the following changes. A bead mixture of 0.1 mm

glass beads (100 mg), 5 × 1.4 mm ceramic beads (Biolab products), and 1 × 4 mm glass bead

was used for five cycles of homogenisation to break-up the faecal material.

Protein digestion

Following extraction, protein amount was quantified using Bradford assay (Bio-Rad, Munich,

Germany) [58] and two methods were compared to digest proteins extracted from LSC or nLSC

procedures.

The in-solution digestion method was performed as follows. Proteins (20 µg starting material)

were reduced in 1 mM dithiothreitol (DTT) and alkylated in 5.5 mM iodoacetamide at room

temperature (RT) for 1 h each. Proteins were pre-digested with LysC at RT for 3 h using a

protein to protease ratio of 75:1. Samples were diluted nine-fold with 50 mM ammonium

bicarbonate and digested overnight with trypsin (Sequencing Grade Modified Trypsin,

Promega) at pH 8.0 using a protein to protease ratio of 75:1.

Filter-aided sample preparation (FASP) was performed as previously published [59]. Briefly,

proteins (10 µg starting material) were reduced in 0.1 M DTT for 40 min at RT. The reduced

samples were added to the filter units (30 kDa membrane cut off) and centrifuged at 14,000 × g

for 15 min. All further centrifugation steps were performed similarly unless otherwise noted.

Samples were then washed with 2X 200 µL urea buffer (100mM Tris/HCl, pH 8.5, 8M urea)

and centrifuged. Proteins were incubated in 50 mM IAA for 20 min at RT in the dark. After

alkylation, samples were centrifuged and washed three times with 100 µL urea buffer. This was

followed by three wash steps with 50 mM ammonium bicarbonate (ABC) for 10 min. Proteins

were digested overnight at 37 °C using trypsin digestion (Sequencing Grade Modified Trypsin,

Promega) at pH 8.0 using a protein to protease ratio of 100:1. On the following day, the peptides

were centrifuged into fresh tubes at 14,000 $\times$ g for 10 min. An additional 40 μL ABC buffer

was added to the filter units and this solution was also centrifuged to increase the peptide yield.

To stop the digestion from either in-solution or FASP workflows, the samples were acidified

to pH 2.5 with formic acid and cleaned for LC-MS/MS measurement using Empore C18 disks

in StageTips [60].

LC-MS/MS measurements

Samples were measured on an EASY-nLC 1200 (Thermo Fisher Scientific) coupled to a Q

Exactive HF mass spectrometer (Thermo Fisher Scientific). The samples prepared for the

sample purification and protein extraction methodologies assessment were all measured in

duplicates to assess instrument reproducibility. Peptides were chromatographically separated

using 75 μm (ID), 20 cm packed in-house with reversed-phase ReproSil-Pur 120 C18-AQ

1.9 μm resin (Dr. Maisch GmbH).

Peptide samples generated as part of the laboratory method optimisation (LSC vs. nLSC, FASP

vs. in-solution) were eluted over 43 min using a 10 to 33 % gradient of solvent B (80 % ACN

in 0.1 % formic acid) followed by a washout procedure. Peptide samples generated as part of

the data analysis assessment (metaproteome vs. metagenome) were eluted over 113 min using

a 10 to 33 % gradient of solvent B (80 % ACN in 0.1 % formic acid) followed by a washout

procedure.

MS1 spectra were acquired between 300-1,650 Thompson at a resolution of 60,000 with an

AGC target of 3×10^6 within 25 ms. Using a dynamic exclusion window of 30 sec, the top 12

most intense ions were selected for HCD fragmentation with an NCE of 27. MS2 spectra were

acquired at a resolution of 30,000 and a minimum AGC of 4.5×10^3 within 45 ms.

LC-MS/MS data processing

Raw data obtained from the instrument were processed using MaxQuant (version 1.5.2.8) [61]. The protein sequence databases used for database search consisted of the complete *Mus musculus* Uniprot database (54,506 sequences) and frequently observed contaminants (248 entries), as well as the mouse microbiome catalogue (~2.6 million proteins) [40] for the raw data from laboratory method optimisation samples or the matching metagenome gene translation (~1.5 million proteins) for the raw data from data analysis assessment samples. A FDR of 1 % was required at the peptide and protein levels. A maximum of two missed cleavages was allowed and full tryptic enzyme specificity was required. Carbamidomethylation of cysteines was defined as fixed modification, while methionine oxidation and N-terminal acetylation were set as variable modifications. Match between runs was enabled where applicable. Quantification was performed using label-free quantification (LFQ) [62] and a minimum peptide count of 1. All other parameters were left to MaxQuant default settings.

Comparison of sample preparation methods

Unless stated otherwise, the analyses described below were performed in the R environment [63]. To compare the different centrifugation, digestion and lysis methods, we counted for each sample the number of peptide and protein groups with intensities and LFQ intensities superior to zero, respectively. We tested for significant differences between methods using unpaired t-tests via the ggplot2 package [64]. Quantified peptides and protein groups were checked for overlap between the centrifugation methods using the VennDiagram package. The proportion of host (*Mus musculus*) proteins was computed by summing up all host proteins iBAQ values and then dividing by the total iBAQ per sample. The centrifugation methods were evaluated using an unpaired t-test.

The taxonomy representation, for the centrifugation methods, was done via the Unipept online software (v. 4.5.1) [34]. The quantified peptides (intensity superior to zero) were imported into

Unipept with I-L not equal. The Unipept result were used to count the number of non-redundant

peptides assigned to each taxonomic node.

For the differential protein abundance analysis (between LSC and nLSC), the MSnBase

package was used as organisational framework for the protein groups LFQ data [65]. Host

proteins, reverse hit and potential contaminant proteins were filtered out. Protein groups were

retained for further analysis only if more than 90 % of samples within either LSC or nLSC

group had an LFQ superior to the first quartile overall LFQ. Significantly changing proteins

were identified using paired t-test. Significance was set at an adjusted p-value of 0.01 following

Benjamini-Hochberg multiple correction testing, as well as a minimum LSC/nLSC fold-change

of ±1.5. The over-representation and GSEA testing of KEGG pathways were done for the

significantly up- and down-regulated proteins as well as for the proteins uniquely identified per

group via the clusterProfiler package based on hypergeometric distribution (p-adj. ≤ 0.05) [66].

Single- versus two-step search assessment using HeLa cell line sample

HeLa cells were prepared for LC-MS/MS measurements using published method [67]. Briefly,

cells were grown in DMEM medium and harvested at 80 % confluence. Proteins were

precipitated using acetone and methanol. Proteins were reduced with DTT and digested with

Lys-C and trypsin. Peptides were purified on Sep-Pak C18 Cartridge.

Sample was measured as described in the LC-MS/MS measurements section but for a few

changes. Peptide sample was eluted over 213 min using a 7 % (0 min), 15 % (140 min) and

33 % (213 min) gradient of solvent B (80 % ACN in 0.1 % formic acid) followed by a washout

procedure. The top 10 most intense ions were selected for HCD fragmentation.

Raw data were processed as described in the LC-MS/MS data processing section with a few

alterations. Match between runs was disabled. The protein sequence databases used for

database search consisted of the complete *Homo sapiens* Uniprot database (93,799 sequences),

frequently observed contaminants (248 entries), as well as the mouse microbiome catalogue

(~2.6 million proteins) [40]. Several processings were performed differing in the number of

microbiome catalogue entries included, which led to an increase in database size of 0×, 1×, 2×,

5×, 10× and 20× compared to the *H. sapiens* database alone. These processings also differed

in the database search strategies used, namely single- or two-step search [36].

Identified MS/MS, peptides and protein groups were assigned to kingdom of origin (conflicts

were resolved to Eukaryota by default). To compare the different database search strategies,

we counted the number of identified MS/MS, non-redundant peptides and protein groups

associated to each kingdom (as well as reverse hits and potential contaminants). We also

calculated the FDR based solely on reverse hits or together with bacterial hits (factual FDR) in

order to investigate the true number of false positives.

Database search strategies assessment using known microbiome samples

We used the samples generated by Kleiner and colleagues, specifically the uneven organisms

preparation described in the earlier publication [32]. This dataset contained LC-MS/MS

measurements (N = 8) that we processed as described in the LC-MS/MS data processing

section with a few alterations. Match between runs was disabled. The protein sequence

databases used for database search consisted of the proteome of all 32 organisms present in the

synthetic samples ("uneven database" = 122,972 sequences), frequently observed contaminants

(248 entries), as well as the mouse microbiome catalogue (~2.6 million proteins) [40]. Several

processings were performed differing in the number of microbiome catalogue entries included,

which led to an increase in database size of 0×, 0.5×, 1×, 2×, 5×, 10× and 20× compared to the

"uneven database" alone. These processings also differed in the database search strategies used,

namely single-step search, "two-step protein" search to keep identified proteins [36], "two-step

taxa" search to keep identified taxa [30], and "two-step two sections" search to keep identified

proteins after sectioned search [37].

Identified protein groups were assigned to database of origin, namely "uneven database" or

mouse microbiome catalogue database. For each sample, this allowed computation of the

number of (1) true positive hits, must be hits from the "uneven database"; (2) false positive

hits, must be hits from the mouse microbiome catalogue; (3) false negative hits, the total

identified protein count in the "uneven database" (total from 8 samples) minus the true

positives; and (4) true negative hits, the total protein count in the mouse microbiome catalogue

minus the false positives. This allowed calculation of the accuracy, precision and sensitivity

for each increase in the database size. We also calculated the factual FDR based on reverse hits

together with mouse microbiome catalogue hits in order to investigate the true number of false

positives.

Using only the processings against the largest database (20×), we filtered our data for protein

groups with a minimum of one or two unique peptides. The true positive count and factual FDR

were calculated (and compared) for each combination of search strategy and filtering, as

described in the previous paragraph.

Taxonomic representation of known microbiome samples

We also used the uneven samples generated by Kleiner and colleagues [32] to investigate the

taxonomic representation derived from MS-identified peptides. The gold-standard processing

was used, with single-step database search against the proteome of all 32 organisms present in

the synthetic samples ("uneven database" = 122,972 sequences). MS-identified peptides were

submitted to (1) Kraken2 (v. 2.1.1) [38], (2) Diamond (v. 2.0.9) [39], or (3) Unipept online (v.

4.5.1) [28] software for taxonomic assignments. The protein sequences from Uniprot

(swissprot and trembl) were used as database for each software. The Diamond alignment was

performed using sensitive and taxonomic classification mode. The Unipept online analysis was

done via the metaproteome analysis function with I-L not equal. The Kraken2 k-mer analysis

was carried out in translated mode using back-translated peptide sequences (back-translation

done with EMBOSS backtranseq). For each software approach, the complete taxonomic

lineage (NCBI) was retrieved per peptide and the lowest common ancestor was determined.

For each sample, we determined and computed the number of taxa that are (1) true positive

hits, must be an identified taxon used for the preparation of the synthetic samples; (2) false

positive hits, must be an identified taxon not used for the preparation of the synthetic samples;

(3) false negative hits, the total number of taxa used for the preparation of the synthetic samples

minus the true positives; and (4) true negative hits, the total number of taxa (with at least one

Uniprot protein) minus the true and false positives. This allowed calculation of the accuracy,

precision, specificity, sensitivity and F-measure for different PSM count thresholds. Taxa were

then quantified per sample based on the different software approaches by summing the peptide

intensities and then normalised to percentage of total peptide intensities. At each taxonomic

level, the Spearman's rank correlation was calculated between the expected taxon

representation in the uneven samples and the taxa representation determined from each

software.

To investigate the taxonomic identification in context of different database search strategies,

we performed the taxonomic annotation via Unipept for all uneven data processings described

in the above section. We then carried out all steps described in the previous paragraph in order

to compute the F-measure per search strategy and database size.

Taxonomic representation of faecal microbiome samples

All subsequent sections use the faecal samples from a 38 mice cohort. These were prepared via

LSC and in-solution protein digestion, as described above. The resulting peptide mixtures were

measured on a Q Exactive HF mass spectrometer and processed against the matching

metagenome gene translation, as described above.

The MS-identified peptides in this dataset were taxonomically annotated with Kraken2,

Diamond and Unipept, as described above. Taxa were quantified as described above (sum of

peptide intensities). The Spearman's rank correlation in taxon representation was calculated for

each pairwise combination of samples within software.

For each sample, we determined and computed the number of species that are (1) true positive

hits, must be an identified species reported in the mouse microbiome catalogue; (2) false

positive hits, must be an identified species not reported in the mouse microbiome catalogue;

and (3) false negative hits, the total number of species reported in the mouse microbiome

catalogue minus the true positives. This allowed calculation of the precision, sensitivity and F-

measure for each samples and annotation software.

Metagenome to metaproteome correlation

All subsequent sections use the faecal samples from a 38 mice cohort. These were prepared via

LSC and in-solution protein digestion, as described above. The resulting peptide mixtures were

measured on a Q Exactive HF mass spectrometer and processed against the matching

metagenome gene translation, as described above. For direct comparison between metagenome

and metaproteome, the identified genes were collapsed into groups identical to protein groups

composition from mass spectrometry. Each gene groups abundance was calculated as the

highest gene abundance within that group. Each gene groups and corresponding protein groups

abundances were correlated across samples using Spearman's rank correlation from the stats

package. Significance was set at an adjusted p-value of 0.05 following Benjamini-Hochberg

multiple correction testing. The GSEA testing of KEGG pathways and Gene ontologies were

performed via the clusterProfiler package based on hypergeometric distribution (p-adj. ≤ 0.05)

[66] following z-scoring of Spearman rho estimate per KEGG orthologies.

Functional KEGG categories representation

For each sample, the protein groups iBAQ values were summed per KEGG category (level 2)

on the basis of KEGG orthology annotation. The same approach was also undertaken for gene

count. The KEGG category abundance were normalised for differing number of KO entries per category and for variation between samples; this was done separately for metagenome and metaproteome. Differences in KEGG category abundance between metagenome and metaproteome were tested using paired t-tests from the stats package. Significance was set at an adjusted p-value of 0.01 following Benjamini-Hochberg multiple correction testing. Significantly changing KEGG categories were prioritised based on gene groups to protein groups correlation (see section Metagenome to metaproteome correlation), whereby the Wilcoxon rank-sum test was used to identify KEGG category containing KO entries whose correlation differ from overall distribution (adjusted p-value ≤ 0.05).

To investigate further these selected KEGG categories, the protein groups iBAQ and gene count were used as described in the previous paragraph to derive KO normalised abundance and t-test results. Using the KO entries from each selected KEGG categories, separate GSEA testing of KEGG pathways were performed via the clusterProfiler package based on hypergeometric distribution (p-adj. ≤ 0.05).

Acknowledgments

MED, BM, XA and LD are grateful to the European Community 7[th] Framework Program under Coordinated Action NEURON-ERANET (grant agreement 291840). BM was supported by grants from the Deutsche Forschungsgemeinschaft (German Research Foundation Cluster of Excellence EXC 2124). BM and NN acknowledge support by the High Performance and Cloud Computing Group at the Center for Data Processing of the University of Tübingen, the state of Baden-Wuerttemberg through bwHPC. The metagenomic work detailed herein used the computing resources of the UK MEDical BIOinformatics partnership – aggregation, integration, visualization and analysis of large, complex data (UK Med-Bio) – which was

737   supported by the Medical Research Council (grant number MR/L01632X/1). XA acknowledge

738   support from the MINECO, Spain (grant number PCIN-2014-105).

Author contributions

LD, XA, MD and BM designed the study. XA and CG generated the mouse cohorts and

collected the murine faecal material. LMG extracted the DNA from all faecal samples prior to

metagenomic sequencing. VA, TG and ID prepared the murine faecal samples for proteomic

measurement by mass spectrometry. LH processed the metagenomic data, generating the

taxonomic and gene abundance outputs. NN processed the metaproteomic datasets and

performed the proteogenomic integration. NN wrote the manuscript with the input from all

authors.

Data Access

The complete metaproteomic bioinformatic workflow is available online [68]. The mass

spectrometry proteomic data have been deposited to the ProteomeXchange Consortium via the

PRIDE [69] partner repository with the dataset identifiers PXD020695, PXD020738,

PXD021928, PXD021932 and PXD027306. Trimmed whole genome sequence data with

mouse reads removed have been deposited with GenBank, EMBL and DDBJ databases under

the BioProject accession PRJNA473429.

References

1.   Jandhyala SM, Talukdar R, Subramanyam C, et al. Role of the normal gut microbiota.
World J Gastroenterol. 2015;21(29):8787-8803.
2.   Valdes AM, Walter J, Segal E, et al. Role of the gut microbiota in nutrition and health.
BMJ. 2018;361:k2179.
3.   Arumugam M, Raes J, Pelletier E, et al. Enterotypes of the human gut microbiome.
Nature. 2011 2011/05//;473(7346):174-180.

4.   Vieira-Silva S, Falony G, Belda E, et al. Statin therapy is associated with lower
prevalence of gut microbiota dysbiosis. Nature. 2020 May;581(7808):310-315.
5.   Wang J, Linnenbrink M, Künzel S, et al. Dietary history contributes to enterotype-like
clustering and functional metagenomic content in the intestinal microbiome of wild
mice. Proceedings of the National Academy of Sciences. 2014;111(26):E2703.
6.   Ladau J, Eloe-Fadrosh EA. Spatial, Temporal, and Phylogenetic Scales of Microbial
Ecology. Trends in Microbiology. 2019 2019/08/01/;27(8):662-669.
7.   Parfrey LW, Knight R. Spatial and temporal variability of the human microbiota.
Clinical Microbiology and Infection. 2012 2012/07/01/;18:5-7.
8.   Sampson TR, Debelius JW, Thron T, et al. Gut Microbiota Regulate Motor Deficits and
Neuroinflammation in a Model of Parkinson's Disease. Cell. 2016 Dec 1;167(6):1469-
1480 e12.
9.   Ley RE, Turnbaugh PJ, Klein S, et al. Microbial ecology: human gut microbes
associated with obesity. Nature. 2006 Dec 21;444(7122):1022-3.
10.   Xiao L, Sonne SB, Feng Q, et al. High-fat feeding rather than obesity drives
taxonomical and functional changes in the gut microbiota in mice. Microbiome. 2017
Apr 8;5(1):43.
11.   Zhang X, Deeke SA, Ning Z, et al. Metaproteomics reveals associations between
microbiome and intestinal extracellular vesicle proteins in pediatric inflammatory
bowel disease. Nat Commun. 2018 Jul 20;9(1):2873.
12.   Hsiao EY, McBride SW, Hsien S, et al. Microbiota modulate behavioral and
physiological abnormalities associated with neurodevelopmental disorders. Cell. 2013
Dec 19;155(7):1451-63.
13.   Tengeler AC, Dam SA, Wiesmann M, et al. Gut microbiota from persons with
attention-deficit/hyperactivity disorder affects the brain in mice. Microbiome. 2020
2020/04/01;8(1):44.
14.   Wang L, Christophersen CT, Sorich MJ, et al. Increased abundance of Sutterella spp.
and Ruminococcus torques in feces of children with autism spectrum disorder. Mol
Autism. 2013 Nov 4;4(1):42.
15.   Biagi E, Candela M, Centanni M, et al. Gut Microbiome in Down Syndrome. PLOS
ONE. 2014;9(11):e112023.
16.   Hettich RL, Pan C, Chourey K, et al. Metaproteomics: harnessing the power of high
performance mass spectrometry to identify the suite of proteins that control metabolic
activities in microbial communities. Anal Chem. 2013 May 7;85(9):4203-14.
17.   Li X, LeBlanc J, Truong A, et al. A metaproteomic approach to study human-microbial
ecosystems at the mucosal luminal interface. PloS one. 2011;6(11):e26542-e26542.
18.   Ram RJ, Verberkmoes NC, Thelen MP, et al. Community proteomics of a natural
microbial biofilm. Science. 2005 Jun 24;308(5730):1915-20.
19.   Zhang X, Li L, Mayne J, et al. Assessing the impact of protein extraction methods for
human gut metaproteomics. J Proteomics. 2018 May 30;180:120-127.
20.   Tanca A, Palomba A, Pisanu S, et al. A straightforward and efficient analytical pipeline
for metaproteome characterization. Microbiome. 2014;2(1):49.
21.   Heyer R, Schallert K, Budel A, et al. A Robust and Universal Metaproteomics
Workflow for Research Studies and Routine Diagnostics Within 24 h Using Phenol
Extraction, FASP Digest, and the MetaProteomeAnalyzer. Front Microbiol.
2019;10:1883.
22.   Gonzalez CG, Wastyk HC, Topf M, et al. High-Throughput Stool Metaproteomics:
Method and Application to Human Specimens. mSystems. 2020 Jun 30;5(3).

23. Xiong W, Giannone RJ, Morowitz MJ, et al. Development of an enhanced
metaproteomic approach for deepening the microbiome characterization of the human
infant gut. J Proteome Res. 2015 Jan 2;14(1):133-41.
24. Tanca A, Palomba A, Pisanu S, et al. Enrichment or depletion? The impact of stool
pretreatment on metaproteomic characterization of the human gut microbiota.
Proteomics. 2015 Oct;15(20):3474-85.
25. Gavin PG, Wong J, Loo D, et al. Metaproteomic sample preparation methods bias the
recovery of host and microbial proteins according to taxa and cellular compartment. J
Proteomics. 2021 May 30;240:104219.
26. Muth T, Behne A, Heyer R, et al. The MetaProteomeAnalyzer: a powerful open-source
software suite for metaproteomics data analysis and interpretation. J Proteome Res.
2015 Mar 6;14(3):1557-65.
27. Cheng K, Ning Z, Zhang X, et al. MetaLab: an automated pipeline for metaproteomic
data analysis. Microbiome. 2017 Dec 2;5(1):157.
28. Mesuere B, Devreese B, Debyser G, et al. Unipept: tryptic peptide-based biodiversity
analysis of metaproteome samples. J Proteome Res. 2012 Dec 7;11(12):5773-80.
29. Timmins-Schiffman E, May DH, Mikan M, et al. Critical decisions in metaproteomics:
achieving high confidence protein annotations in a sea of unknowns. ISME J. 2017
Feb;11(2):309-314.
30. Tanca A, Palomba A, Fraumene C, et al. The impact of sequence database choice on
metaproteomic results in gut microbiota studies. Microbiome. 2016 Sep 27;4(1):51.
31. Kunath BJ, Minniti G, Skaugen M, et al. Metaproteomics: Sample Preparation and
Methodological Considerations. Adv Exp Med Biol. 2019;1073:187-215.
32. Kleiner M, Thorson E, Sharp CE, et al. Assessing species biomass contributions in
microbial communities via metaproteomics. Nat Commun. 2017 Nov 16;8(1):1558.
33. Pible O, Allain F, Jouffret V, et al. Estimating relative biomasses of organisms in
microbiota using "phylopeptidomics". Microbiome. 2020;8(1):30-30.
34. Mesuere B, Van der Jeugt F, Willems T, et al. High-throughput metaproteomics data
analysis with Unipept: A tutorial. J Proteomics. 2018 Jan 16;171:11-22.
35. Bäckhed F, Ding H, Wang T, et al. The gut microbiota as an environmental factor that
regulates fat storage. Proceedings of the National Academy of Sciences of the United
States of America. 2004;101(44):15718-15723.
36. Jagtap P, Goslinga J, Kooren JA, et al. A two-step database search method improves
sensitivity in peptide sequence matches for metaproteomics and proteogenomics
studies. Proteomics. 2013 Apr;13(8):1352-7.
37. Kumar P, Johnson JE, Easterly C, et al. A Sectioning and Database Enrichment
Approach for Improved Peptide Spectrum Matching in Large, Genome-Guided Protein
Sequence Databases. J Proteome Res. 2020 Jul 2;19(7):2772-2785.
38. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome
Biol. 2019 Nov 28;20(1):257.
39. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.
Nat Methods. 2015 Jan;12(1):59-60.
40. Xiao L, Feng Q, Liang S, et al. A catalog of the mouse gut metagenome. Nature
biotechnology. 2015 Oct;33(10):1103-8.
41. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids
Res. 2000 Jan 1;28(1):27-30.
42. Zhang X, Ning Z, Mayne J, et al. MetaPro-IQ: a universal metaproteomic approach to
studying human and mouse gut microbiota. Microbiome. 2016 Jun 24;4(1):31.

43. Tanca A, Manghina V, Fraumene C, et al. Metaproteogenomics Reveals Taxonomic
and Functional Changes between Cecal and Fecal Microbiota in Mouse. Front
Microbiol. 2017;8:391.
44. Michalski A, Damoc E, Hauschild JP, et al. Mass spectrometry-based proteomics using
Q Exactive, a high-performance benchtop quadrupole Orbitrap mass spectrometer. Mol
Cell Proteomics. 2011 Sep;10(9):M111 011015.
45. Williamson JC, Edwards AV, Verano-Braga T, et al. High-performance hybrid
Orbitrap mass spectrometers for quantitative proteome analysis: Observations and
implications. Proteomics. 2016 Mar;16(6):907-14.
46. Ericsson AC, Hart ML, Kwan J, et al. Supplier-origin mouse microbiomes significantly
influence locomotor and anxiety-related behavior, body morphology, and metabolism.
Communications Biology. 2021 2021/06/10;4(1):716.
47. Randall DW, Kieswich J, Swann J, et al. Batch effect exerts a bigger influence on the
rat urinary metabolome and gut microbiota than uraemia: a cautionary tale.
Microbiome. 2019 Sep 2;7(1):127.
48. Muth T, Kolmeder CA, Salojarvi J, et al. Navigating through metaproteomics data: A
logbook of database searching. Proteomics. 2015 Oct;15(20):3439-53.
49. Heyer R, Schallert K, Zoun R, et al. Challenges and perspectives of metaproteomic data
analysis. J Biotechnol. 2017 Nov 10;261:24-36.
50. Grassl N, Kulak NA, Pichler G, et al. Ultra-deep and quantitative saliva proteome
reveals dynamics of the oral microbiome. Genome medicine. 2016;8(1):44.
51. Mills RH, Vazquez-Baeza Y, Zhu Q, et al. Evaluating Metagenomic Prediction of the
Metaproteome in a 4.5-Year Study of a Patient with Crohn's Disease. mSystems. 2019
Jan-Feb;4(1).
52. Edirisinghe JN, Weisenhorn P, Conrad N, et al. Modeling central metabolism and
energy biosynthesis across microbial life. BMC Genomics. 2016 Aug 8;17:568.
53. Sidoli S, Kulej K, Garcia BA. Why proteomics is not the new genomics and the future
of mass spectrometry in cell biology. J Cell Biol. 2017 Jan 2;216(1):21-24.
54. Hoyles L, Fernandez-Real JM, Federici M, et al. Molecular phenomics and
metagenomics of hepatic steatosis in non-diabetic obese women. Nat Med. 2018
Jul;24(7):1070-1080.
55. Andrews S. FastQC A Quality Control tool for High Throughput Sequence Data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/; 2010.
56. Parks DH, Chuvochina M, Chaumeil PA, et al. A complete domain-to-species
taxonomy for Bacteria and Archaea. Nature biotechnology. 2020 Sep;38(9):1079-1086.
57. Huerta-Cepas J, Forslund K, Coelho LP, et al. Fast Genome-Wide Functional
Annotation through Orthology Assignment by eggNOG-Mapper. Molecular Biology
and Evolution. 2017;34(8):2115-2122.
58. Bradford MM. A rapid and sensitive method for the quantitation of microgram
quantities of protein utilizing the principle of protein-dye binding. Anal Biochem. 1976
May 7;72:248-54.
59. Wisniewski JR, Zougman A, Nagaraj N, et al. Universal sample preparation method for
proteome analysis. Nat Methods. 2009 May;6(5):359-62.
60. Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-
fractionation and storage of peptides for proteomics using StageTips. Nat Protoc.
2007;2(8):1896-906.
61. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nature
biotechnology. 2008 Dec;26(12):1367-72.

62.   Cox J, Hein MY, Luber CA, et al. Accurate proteome-wide label-free quantification by
delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol Cell
Proteomics. 2014 Sep;13(9):2513-26.
63.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing; 2018.
64.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016.
65.   Gatto L, Lilley KS. MSnbase-an R/Bioconductor package for isobaric tagged mass
spectrometry data visualization, processing and quantitation. Bioinformatics. 2012 Jan
15;28(2):288-9.
66.   Yu G, Wang LG, Han Y, et al. clusterProfiler: an R package for comparing biological
themes among gene clusters. OMICS. 2012 May;16(5):284-7.
67.   Schmitt M, Sinnberg T, Nalpas NC, et al. Quantitative Proteomics Links the
Intermediate Filament Nestin to Resistance to Targeted BRAF Inhibition in Melanoma
Cells. Mol Cell Proteomics. 2019 Jun;18(6):1096-1109.
68.   Nalpas N, Macek B. Integrated metaproteomics workflow. 1.0. Zenodo; 2020.
69.   Perez-Riverol Y, Csordas A, Bai J, et al. The PRIDE database and related tools and
resources in 2019: improving support for quantification data. Nucleic Acids Res. 2019
Jan 8;47(D1):D442-D450.

Table 1: Performance comparison of different sample preparation and data analysis steps. In bold are the best methods according to assessed criteria: peptide/protein count, host/dietary contamination, *Firmicutes* or *Bacteroidetes* representation, time efficiency, FDR, identification rate, taxon-assigned peptides and number of taxonomic identification precision. The performance status is displayed using minus sign for poor, equal sign for similar/no difference or plus sign for good performance.

		Peptide/protein count	Host/dietary contamination	Firmicutes	Bacteroidetes	Time efficiency	FDR	Identification rate	Taxon assigned peptides	Precision
Centrifugation	**LSC**	+	-	-	+	-				
	nLSC	-	+	+	-	+				
Digestion	**In-solution**	+				-				
	FASP	-				+				
Search strategy	**Single-step**	-				+	+	-		
	Two-step protein	+				-	--	+		
	Two-step sections	+				--	-	+		
	Two-step taxa	-				-	+	-		
Taxon quantification	Kraken2					+			+	--
	Diamond					--			-	-
	Unipept					+			+	+

933 Figures

Figure 1: Low speed centrifugation impacts protein identification and taxonomic representation. A) Number of MS/MS spectra, peptides and protein groups per samples for the comparison between LSC (red) and nLSC (blue) methods. B) Number of identified MS/MS

spectra, peptides and protein groups per samples for the comparison between LSC-in solution

digestion (red), LSC-FASP (grey), nLSC-in solution digestion (blue) and nLSC-FASP (orange)

methods. A-B) Represented significance results correspond to t-test on N = 12 (A) or N = 6

(B): * p- value $\leq$ 0.05, ** $\leq$ 0.01, *** $\leq$ 0.001. C) Hierarchical representation of Unipept-

derived taxonomy (down to phylum level) for the peptide identified in the LSC and nLSC. The

barplot represent the taxonomic abundance for LSC (red) and nLSC (blue) methods based on

peptide counts (only for taxon identified with 3 or more peptides). D) Overlap in the overall

identified peptides or protein groups between the LSC and nLSC methods. (E) Volcano plot of

the protein abundance comparison between LSC and nLSC approaches. Significant protein

groups based on paired t-test from N = 12 with FDR $\leq$ 0.01 and absolute fold-change $\geq$ 2.5. (F)

KEGG pathways over-representation testing for the protein groups that significantly increase

(red) or decrease (blue) in abundance between LSC and nLSC sample preparation approaches.

Fisher exact-test threshold (gold dotted line) set to adjusted p-value $\leq$ 0.05.

Figure 2: Two-step database search in combination with target-decoy strategy leads to a

dramatic increase in false positive rate. A) The protein groups count is shown for single- or

two-step search strategies across increasingly large protein sequence databases. Counts are

colour-coded per category, with eukaryote (grey), bacteria (red), contaminant (blue) and

reverse (orange) hits. B) The FDR is calculated for single- or two-step search strategies across

increasingly large protein sequence databases. The FDR is calculated based on reverse hits only

(circle shape) or reverse plus bacterial hits (triangle shape). C & D) The sensitivity (C) and

factual FDR (D) based on protein groups identification across increasingly large protein

sequence databases. The compared database search strategies are single-step (blue), two-step

taxon filtering (grey) and two-step protein filtering without (red) or with (orange) database

sectioning. Lines represent the median (and the shading corresponds to the standard error) from

N = 8 LC-MS/MS runs. E) The true positive count based on protein groups identified with a

minimum of one (shaded colouring) or two (unshaded colouring) unique peptides for the largest database (i.e. 20). The compared database search strategies are single-step (blue), two-step taxon filtering (grey) and two-step protein filtering without (red) or with (orange) database sectioning. Bars and numbers indicate the median count, while error bars correspond to the standard deviation, from N = 8 LC-MS/MS runs. The overall maxima of true positive count based on single-step search is indicated as a horizontal dotted line (gold).

Figure 3: Unipept software provides the most precise taxonomic annotation of MS-based peptide identification. A) Percentage of taxon-annotated peptides at each taxonomic level for the comparison between Kraken2 (red), Diamond (grey) and Unipept (blue) software. B) Assessment of the impact of the minimum number of PSM count per taxon onto the F-measure for taxonomic annotation. The F-measure was compared between Kraken2 (red), Diamond (grey) and Unipept (blue) software. C) Heatmap representing the correlation (Spearman ρ) in taxonomic abundance between sample input protein (expectation) and different taxonomic annotation software (i.e. Kraken2, Diamond and Unipept). The correlation was performed overall, as well as for each taxonomic level. D) Organisms pooled in artificial samples are ranked based on the protein material input, as displayed in the left-most barplots (x-axis in $\log_{10}$ scale). The proteome size (ORFs) for these organisms on UniProt web resource is displayed in the right-most barplot (x-axis in $\log_{10}$ scale). The heatmap compares the taxon identification across samples between Kraken2, Diamond and Unipept. A-D) Samples from the study by Kleiner and colleagues, with N = 8. E) Overlap in the overall identified taxa between the Kraken2 (red), Diamond (grey) and Unipept (blue) software. F) A comparison of the F-measure distribution for taxonomic annotation between the Kraken2 (red), Diamond (grey) and Unipept (blue) software. Each point represents an individual mouse. E-F) Samples from this study using mouse faecal material, with N = 38.

Figure 4: Functionally active pathways derived from the metaproteome differs from the

metagenome potential. A) Correlation is shown between each protein groups (metaproteome)

and corresponding gene "groups" (metagenome) abundances. Correlation was tested using

Spearman's rank correlation and p-value was adjusted for multiple testing using Benjamini-

hochberg correction. Significantly correlating protein/gene groups are in red colours, while

significantly anti-correlating protein/gene groups are in green colours (adjusted p-value $\leq$

0.05). B) GSEA of KEGG pathways based on ranking of the protein/gene groups correlation.

Pathway node colour corresponds to GSEA results adjusted p-value and node size matches the

number of protein/gene group assigned to the pathway. C) Comparison in the proportion of

selected KEGG functional categories (level 2) between metaproteome (red) and metagenome

(grey). Paired t-test p-values are indicated (N = 38). D) GSEA of KEGG pathways based on

ranking of t-test results from KEGG orthology proportion between metaproteome and

metagenome. KEGG pathways are colour-coded based on KEGG functional categories (level

2). Only significantly over-represented KEGG pathways are shown with adjusted p-value $\leq$

0.05. E) Interaction network between KEGG orthologies and KEGG pathways for the KEGG

functional category "Protein families: genetic information processing". Pathway node size

corresponds to number of KEGG orthologies associated to it. KEGG orthologies are colour-

coded based on directional adjusted p-value from the t-test comparison between metaproteome

and metagenome.

1005