AbaM Regulates Quorum Sensing, Biofilm Formation and Virulence in

Acinetobacter baumannii

Mario López-Martín[1], Jean-Frédéric Dubern[1], Morgan R. Alexander[2], Paul Williams[1]*

[1]Biodiscovery Institute and School of Life Sciences, University of Nottingham, Nottingham, United Kingdom.

[2]Advanced Materials and Healthcare Technologies, School of Pharmacy, University of Nottingham, Nottingham, United Kingdom

Running title: AbaM and quorum sensing in *Acinetobacter*

Keywords**:** *Acinetobacter*, quorum sensing, *N*-acylhomoserine lactones, RsaM, virulence, biofilms

Corresponding author email: paul.williams@nottingham.ac.uk

ABSTRACT

Acinetobacter baumannii possesses a single divergent *luxR/luxI*-type quorum sensing (QS) locus named *abaR/abaI*. This locus also contains a third gene located between *abaR* and *abaI* which we term *abaM* that codes for an uncharacterized member of the RsaM protein family known to regulate *N*-acylhomoserine lactone (AHL) dependent QS in other β- and γ-proteobacteria. Here we show that disruption of *abaM* via a T26 insertion in *A. baumannii* strain AB5075 resulted in increased production of *N*-(3-hydroxydodecanoyl)-L-homoserine lactone (OHC12) and enhanced surface motility and biofilm formation. In contrast to the wild type and *abaI*::T26 mutant, the virulence of the *abaM::*T26 mutant was completely attenuated in a *Galleria mellonella* infection model. Transcriptomic analysis of the *abaM*::T26 mutant revealed that AbaM differentially regulates at least 76 genes including the *csu* pilus operon and the acinetin 505 lipopeptide biosynthetic operon, that are involved in surface adherence, biofilm formation and virulence. A comparison of the wild type, *abaM*::T26 and *abaI::*T26 transcriptomes, indicates that AbaM regulates ~21% of the QS regulon including the *csu* operon. Moreover, the QS genes (*abaI/abaR*) were among the most upregulated in the *abaM*::T26 mutant. *A. baumannii lux*-based *abaM* reporter gene fusions revealed that *abaM* expression is positively regulated by QS but negatively auto-regulated. Overall, the data presented in this work demonstrates that AbaM plays a central role in regulating *A. baumannii* QS, virulence, surface motility and biofilm formation.

IMPORTANCE

Acinetobacter baumanni is a multi-antibiotic resistant pathogen of global healthcare importance. Understanding *Acinetobacter* virulence gene regulation could aid the development of novel anti-infective strategies. In *A. baumannii,* the *abaR* and *abaI* genes that code for the receptor and synthase components of an *N*-acylhomoserine (AHL) lactone-dependent quorum sensing system (QS) are separated by *abaM.* Here we show that although mutation of *abaM* increased AHL production, surface motility and biofilm development, it resulted in the attenuation of virulence. AbaM was found to control both QS-dependent and QS-independent genes. The significance of this work lies in the identification of AbaM, an RsaM ortholog known to control virulence in plant pathogens, as a modulator of virulence in a human pathogen.

INTRODUCTION

Acinetobacter baumannii is a Gram-negative opportunistic nosocomial pathogen that causes a wide range of infections in humans, most commonly pneumonia, but also bacteremia, skin, soft tissue and urinary tract infections, meningitis and endocarditis (1). The rise of multi-drug resistant strains has limited the treatment options for this pathogen which has become a major threat to hospital patients worldwide (2). Indeed, the WHO classified *A. baumannii* as a critical pathogen for which new antibiotics are urgently required (3). For this reason, a better understanding of the virulence of *A. baumannii* should aid the development of new therapeutic strategies for preventing and treating *Acinetobacter* infections. Several virulence factors and regulators involved in *A. baumannii* pathogenesis have been characterized to date. These include outer membrane proteins (e.g. OmpA), pili, capsular polysaccharide, iron acquisition systems, outer membrane vesicles, secretion systems and phospholipases (4–9), as well as regulators such as H-NS and two component systems (10–13). Some *A. baumannii* strains also undergo phase variation where opaque colony variants exhibit greater motility and virulence but reduced biofilm formation compared with the translucent variants (11). A detailed review of *A. baumannii* virulence can be found in Morris *et al*, 2019 (14).

One well-established mechanism of virulence gene regulation in diverse pathogens is quorum sensing (QS) (15). This cell-cell communication system is employed by bacteria to coordinate the expression of specific genes as a function of population density. QS is mediated via the synthesis, release and detection of diffusible signalling molecules such as the *N*-acyl-homoserine lactones (AHLs) (16). *A. baumannii* and related pathogenic *Acinetobacter* spp. possess a LuxR/LuxI QS system, consisting of an AHL synthase (AbaI) and

a transcriptional regulator (AbaR) that is activated on binding an AHL. Most pathogenic

Acinetobacter spp. produce AHLs with acyl side chains of 10 to 12 carbons length with *N*-(3-

hydroxydodecanoyl)-L-homoserine lactone (OHC12) being most commonly encountered.

Many strains are however capable of producing other AHLs (17). Several reports have linked

QS to biofilm formation and surface motility (18–20) while others have suggested that it

plays a role in virulence in a strain and animal model dependent manner (21, 22). However,

our current knowledge of the role of QS in the virulence of pathogenic *Acinetobacter* spp. is

limited.

Located adjacent to the *abaI* gene in *Acinetobacter* there is an ortholog of the RsaM protein

family. These are found in diverse β- and γ-proteobacteria*,* including *Burkholderia* spp.,

Pseudomonas fuscovaginae, Halothiobacillus neapolitanus and *Acidithiobacillus ferrooxidans*

(23). The first ortholog to be characterized was RsaM in the plant pathogen *P. fuscovaginae*.

This was shown to negatively regulate AHL production and was required for full virulence in

rice plants (24). Transcriptomic analysis revealed that RsaM partially regulates the QS

regulon, as well as modulating the expression of diverse genes in a QS-independent manner

(25). Similarly, TofM, the RsaM ortholog found in *Burkholderia glumae*, represses AHL

production while positively regulating toxoflavin and virulence in rice (26). The two RsaM

orthologs present in *Burkholderia thailandensis* are both negatively auto-regulated while

being positively controlled by their cognate QS systems (27). In *Burkholderia cenocepacia*

H111, *Bc*RsaM downregulates AHL biosynthesis and modulates swarming motility, biofilm

formation, protease and siderophore production (28). Structural and biochemical analysis of

*Bc*RsaM showed that it forms dimers in solution and does not appear to bind DNA or AHLs,

suggesting that RsaM family proteins act as post-transcriptional or post-translational

regulators (23).

RsaM orthologs clearly play a central role in the regulation of QS-dependent and QS-

independent gene expression and virulence in plant pathogens. Here we investigated the

role and regulation of *abaM* in *A. baumannii* AB5075, a comparatively recently isolated

multi-antibiotic resistant, hypervirulent strain (29). We show that AbaM, in the opaque

variant of *A. baumanni* 5075, controls AHL production, surface motility and biofilm

formation and is required for virulence in a *Galleria mellonella* infection model. QS

positively regulates *abaM* expression which is turn is negatively autoregulated.

Transcriptomic analysis of the *abaM* and *abaI* mutants indicate that the AbaM and QS

regulons overlap. These data are consistent with a central role for AbaM in regulating gene

expression and the pathobiology of *A. baumannii*.

RESULTS

Organization of the QS locus in *A. baumannii* 5075. The genome of *A. baumannii* AB5075 possesses a single QS locus comprised of two divergently transcribed genes: an AHL synthase (*abaI/ABUW_3776/ABUW_RS18385*) and a response regulator (*abaR/ABUW_3774/ ABUW_RS18375*) gene. Between *abaR* and *abaI* a third gene is located which here we term *abaM* (*ABUW_3775*) (**Fig. 1A**). The chromosomal organization of these three QS genes is well conserved among *Acinetobacter* spp (**Fig. S1**). Despite the location of *abaM* adjacent to *abaI* and transcribed in the same direction, AbaM has only low (~20-30 %) sequence identity to orthologs present in *Pseudomonas fuscovaginae* and *Burkholderia spp* (**Fig. 1B**). However, it retains the well-conserved regions shared by other RsaM orthologues, including most of the protein secondary structural elements and the characteristic hydrophobic core cluster, consisting of four tryptophan residues (Trp60, Trp75, Trp77 and Trp125) (23).

AHL production is enhanced in an *abaM* mutant AHL production in the opaque variants of the AB5075 wild type, *abaM*::T26 and *abaI*::T26 mutants respectively was quantified via LC MS/MS during growth under static conditions since it appears to be enhanced by surface attachment in other *A. baumannii* strains (30). OHC12 (**Fig. 2A**) was the major AHL produced by AB5075 under these conditions. Compared with the wild type, the *abaM*::T26 mutant produced significantly greater amounts of OHC12 at each time point (between 100 and 875-fold difference) sampled (**Fig. 2B**). *N*-(3-hydroxydecanoyl)-L-homoserine (OHC10), was also detected at much lower concentrations in the *abaM*::T26 mutant throughout growth, but only in the 24 h sample in the wild type (**Fig. 2C**). No AHLs were detected in any of the

abaI::T26 samples (**Fig. 2B and C**). These data suggest that AbaM is a negative regulator of

AHL production.

Contribution of QS and *abaM* to surface motility, biofilm formation and virulence. The

surface motility of all three strains on 0.3% Eiken agar LS-LB plates was examined.

Compared with the wild-type (59.6 ± 0.7 mm), the *abaI*::T26 mutant exhibited significantly

reduced (36.5 ± 1.4 mm) whereas the *abaM*::T26 mutant was significantly more motile (76.7

± 2.4 mm) (**Fig. 3A**). The provision of exogenous OHC12 increased the surface motility of

both wild type and *abaI* mutant to a similar level to that of the *abaM* mutant (**Fig. S2A**)

The ability of AB5075, *abaI*::T26 and *abaM*::T26 to attach to abiotic surfaces was evaluated

on propylene tubes. The *abaM* mutant formed $\sim$3 fold more biofilm than the wild-type (**Fig.**

3B). Under these growth conditions, the biofilm produced by the *abaI* mutant (opaque

variant; **Fig. 3B**) was not significantly different from than the wild type but increased

following the exogenous provision of OHC12 (**Fig. S2B**).

The contribution of *abaM* and QS to AB7075 virulence was assessed using a

G. mellonella larvae infection model (**Fig. 3C**). No differences in killing were observed

between the wild-type and the *abaI*::T26 mutant when injecting either

2×10^4 or 2×10^5 CFU/larva. However, the *abaM*::T26 mutant was significantly less virulent

than the parental strain. Larvae injected with 2×10^5 CFU of the *abaM*::T26 mutant also

showed a lower death rate than the larvae injected with 2×10^4 CFU of the wild-type or the

abaI::T26 mutant. Exogenously supplied OHC12 did not affect the virulence of the wild type

(**Fig. S2C**).

Overall, these results suggest that *abaM* is a negative regulator of surface motility and

biofilm formation and required for full virulence in *G. mellonella*.

Genetic complementation of the *abaM* mutant phenotypes. Complementation of the

abaM::T26 mutant with the *abaM* gene in *trans* (pMQ_*abaM*) restored surface motility

(**Fig.S3A**), biofilm formation (**Fig. S3B**) and reduced both OHC12 and OHC10 production by

approximately 50% (**Fig. S3C and D**). However, complementation of the *abaM* mutation did

not restore *abaM*::T26 virulence to wild-type (**Fig. S3E**).

Transcriptomic analysis of *abaI*::T26 and *abaM*::T26. To characterize the AbaM and QS

regulon of *A. baumannii* AB5075, we performed transcriptomic profiling of AB5075 in

comparison with the *abaM*::T26 and *abaI*::T26 mutants using RNA sequencing (RNA-seq),

which was then validated for two key target genes via quantitative real-time PCR. For these

analyses we used total RNA extractions from cells grown for 18 h in static conditions when

maximum OHC12 levels are produced by the *abaM* mutant (**Fig. 2**).

When compared with the wild-type strain, 88 genes were upregulated and 9 downregulated

in the *abaI*::T26 mutant, while 52 were upregulated and 24 downregulated in the

abaM::T26 mutant (log$_2$(fold change) $\geq$ 1; **Fig.4** and **Tables S3** and **S4**). Moreover, 21 of the

upregulated genes were shared between *abaI*::T26 and *abaM*::T26 (**Fig. 4, Tables S3** and

S4), while none of the downregulated genes were co-regulated. Among the genes

upregulated in both mutants there were all the genes of the *csu* operon, a putative TetR

family   transcriptional   regulator   (*ABUW_1486/ABUW_RS07245)*   located   immediately

upstream of the *csu* operon as well as   genes coding for a flavohemoprotein, an

uncharacterized transcriptional regulator, a thermonuclease, a sulfate permease, a toxic

anion resistance protein and 7 hypothetical proteins (**Tables S3 and S4**).

Moreover, genes of the biosynthetic operon involved in the synthesis of acinetin 505, the

QS response regulator, *abaR* and the AHL synthase *abaI* were both upregulated in the

abaM::T26 mutant, which also differentially expressed genes encoding for proteins involved

in the stress response, iron-acquisition, diverse metabolism and energy production,

chaperones, protein folding and antibiotic resistance e.g. class D beta-lactamase OXA-23

involved in resistance to carbapenems (**Table S4**). Similarly, differentially regulated genes in

the *abaI* mutant included diverse metabolic and energy production-related genes, as well as

diverse genes coding for transcriptional regulators, stress response-related proteins and

membrane transport proteins (**Tables S3)**.

To validate the transcriptomic data, qPCR was performed with the same RNA samples used

in the RNA-seq for the *csuA/B* and the *ABUW_3773* (the first gene of the acinetin 505

biosynthetic operon) genes (**Fig. 5**). When compared with the wild-type, the data obtained

showed a significant increase in *csuA/B* expression in both the *abaI*::T26 and *abaM*::T26

mutants, while *ABUW_3773* expression was significantly higher only in the *abaM*::T26

mutant. These results correlate with the data obtained from the RNA-seq experiments.

Regulation of *abaM*. To further elucidate the regulation of *abaM* expression, an *abaM*

promoter – *luxCDABE* operon fusion was constructed. This was introduced via a miniTn*7*

transposon into AB5075 and both *abaI*::T26 and *abaM*::T26 mutants, and the activity of the

predicted promoter was measured by luminescence output in the presence or absence of

OHC12. The activity of the *abaM* promoter significantly varied between the strains. In the

abaM::T26, luminescence was approximately 40% higher than in the wild-type, while the

abaI::T26 mutant showed a 75% reduction when compared with the parental strain (**Fig.**

6A). Moreover, exogenous provision of OHC12 increased the *abaM* promoter activity in all

three strains (**Fig. 6B**). Similarly, **Fig. 6C** shows that expression of an *abaI*::*lux* promoter

fusion which is reduced in the *abaI* mutant compared with the wild type strain, is strongly

stimulated by OHC12. These data suggest that *abaM* expression is negatively autoregulated

but, in common with *abaI* is positively regulated by QS which in turn is negatively controlled

via AbaM (**Fig. 7**).

DISCUSSION

In this study we have established that the RsaM ortholog, AbaM plays a major role in

regulating QS-dependent and QS-independent genes in *A. baumannii* 5075*.* Disruption of

abaM substantially increased AHL production, indicating that AbaM negatively regulates

AHL biosynthesis. In *A. baumannii* 5075, the concentration of OHC12 produced was very low

(< 1 nM), suggesting that, at least under the static growth conditions used in this study,

AbaM exerts tight control over QS. This raises the question of when and under what

conditions QS is active in AB5075, especially since the half-maximal responses for LuxR

proteins activated by long chain AHLs is in the 5-10 nM range (31). In *A. baumannii*, the QS

locus has an *RXI* topological arrangement previously described in other bacteria, where the

X gene between the QS regulator (*R*) and the synthase (*I*) genes is a negative regulator of

QS. Examples of three different classes of X include *rsaL*, *rsaM* and *mupX* (32). In this

context, AbaM appears to behave similarly to RsaL in the *Pseudomonas aeruginosa*

lasR/rsaL/lasI QS system, despite their functional and structural differences. The *rsaL* gene

which is divergent to *lasI* is positively regulated by LasR and antagonizes LasR-mediated

activation of *lasI* so counteracting the QS positive feedback autoinduction and providing

AHL homeostasis (33). This type of regulatory circuit (**Fig. 7**), termed an incoherent feed

forward loop (IFFL) in contrast to simple feed forward loops, can display complex behaviours

that include stabilization of output signals and bounded output which ensures robustness

against fluctuations in the input signal levels (32, 34). Hence the large increase in AHL

production following deletion of *abaM*, *rsaL* and *rsaM* genes that result in less virulent

mutants is indicative of the importance of a stabilizing negative regulatory pathway in AHL-

dependent QS systems (32, 34).

Consistent with this model, *abaM* expression was found to be negatively auto-regulated but

positively regulated by QS. *In silico* analysis of the DNA sequence between *abaR* and *abaI*

using Bprom (www.softberry.com) and BDGP (www.fruitfly.org) as well as our RNAseq data

all predict the presence of putative -10 and -35 regions for *abaI* and *abaM* respectively

suggesting that these genes are not co-transcribed. Similar findings have been reported for

rsaM1 and *rsaM2* from *Burkholderia thailandensis* (27). Reverse transcriptase (RT)-PCR (**Fig.**

S4) confirm that *abaM* and *abaI* in *A. baumannii* A5075 do not form an operon. Moreover,

the *abaM* gene has a predicted *lux*-box (CTGGTTAAATATAACAG) 68 bp upstream of the start

codon and 178 bp downstream of the -35 and -10 promoter elements. This *lux*-box is similar

to those found upstream of *abaI* (CTGTAAATTCTTACAG) in both *A. baumannii* 5075 and *A.*

nosocomialis M2. These results are consistent with an IFFL circuit although further work will

be required to fully characterize its properties and control of genes co-regulated by QS and

AbaM.

Phenotypic characterization of the *A. baumanni abaM* mutant revealed enhanced surface

motility and biofilm formation but reduced virulence compared with the wild type. Previous

studies have shown that *rsaM* orthologues are required for full virulence in plants (24, 26),

but this is, to the best of our knowledge, the first time that an *rsaM*-like gene has been

reported to be required for full virulence in a human pathogen albeit in an insect infection

model. The contribution of *Bc*rsaM in *B. cenocepacia* to swarming motility and biofilm

formation has also been reported (28). However, deletion of *bcrsaM* reduced both

swarming and surface attachment, the opposite to that observed for the *abaM* mutant.

Interestingly, both biofilm formation and surface motility in *Acinetobacter* have been

associated with increased virulence (10, 35).

Genetic complementation of the *abaM* mutant was achieved for surface motility, biofilm

formation and AHL production but not for virulence in *Galleria mellonella.* Similar

observations have been previously reported for other *abaM* orthologues, most notably

tofM (26) and *bcrsaM* (28), leading to the suggestion that RsaM-like proteins may be *cis-*

acting regulators (23).

To further define the role of QS in *A. baumannii* 5075, phenotypic characterization of the

AHL synthase mutant, *abaI*::T26, was performed. The mutant did not produce any

detectable AHLs, consistent with previous studies and bioinformatic analysis indicating that

A. baumannii 5075 possesses a single QS locus that is responsible for AHL production (18,

36). Similarly, disruption of *abaI* negatively affected surface motility and responded to

exogenous 3OHC12, as previously noted for other *Acinetobacter* strains/species (20). In

AB5075, biofilm formation in the opaque variant of the *abaI*::T26 mutant was not

significantly different from with wild type. However, it increased well above wild type in

response to exogenous OHC12. consistent with other work on the *Acinetobacter* AHL

synthase (18, 37).   Previous studies on QS and biofilm formation in *Acinetobacter* have been

performed with strains that were, in contrast to AB5075, either not phase variable or not

known to be phase variable. Since the experiments performed in this study were all carried

out with the opaque *Acinetobacter* variant, we also investigated biofilm formation by the

translucent variant. **Fig S5** shows that the translucent variant produced less biofilm that the

wild type. However, biofilm formation increased for both opaque and translucent variants

in response to exogenous OHC12. Furthermore, our results suggest that QS does not play

an important role in virulence in the *G. mellonella*. While this is not unprecedented (38), the

role of *Acinetobacter* QS in virulence is still not well defined, and other studies suggest that

QS may play an important role, depending on the strain and infection model used (21, 22).

Overall, our data suggest that for strain AB5075, QS is important in surface motility and

biofilm formation, but not virulence. However, further work is required to fully elucidate the

role of QS in the pathogenesis of *Acinetobacter* infection and any cross-talk with other

regulatory networks.

Here we performed RNAseq on *A. baumannii* AB5075 grown in static conditions where AHL

production was elevated in order to identify genes regulated via AbaM and QS and likely to

be involved in surface attachment and biofilm formation. A comparison of the genes

differentially expressed in the *abaM* and *abaI* mutants with the wildtype revealed that

AbaM has both QS-dependent (~22% of the QS regulon) and QS independent gene targets.

A similar overlap has been noted for the RsaM regulon and its cognate QS system in *P.*

fuscovaginae (25). Among the most upregulated genes in both *abaI* and *abaM* mutants

when compared with the wild-type were those belonging to the *csu* operon (*ABUW_1487-*

ABUW_1492/ABUW_RS07250-ABUW_RS07275). This operon encodes the proteins

responsible for the synthesis of the Csu pilus, a type I chaperone-usher pilus involved in

attachment and biofilm formation (39–41). Moreover, the *abaM*::T26 mutant also showed

higher expression of some genes of the acinetin 505 biosynthetic operon, which has also

been linked to biofilm formation in *A. baumannii* ATCC17978 (42).

A previous comparison of the transcriptomes of the multi-drug resistant clinical *A.*

baumanni strain 863 with an isogenic *abaI* deletion mutant highlighted the differential

regulation of 352 genes involved in carbon source metabolism, energy production, stress

response and translation (49). However, apart from *abaI*, no other common differentially

regulated genes could be identified when the A5075 *abaI* and the 863 *abaI* mutant

transcriptomes are compared. This may be because of the different strains and growth

conditions and sampling times used. In addition, the *abaI* mutant reported by Ng et al (49)

exhibited a growth defect (49). This raises the possibility of a secondary mutation

contributing to the transcriptome data which was not validated by chemical or genetic

complementation with OHC12 or *abaI* respectively.

The work described here establishes that AbaM plays a central role in regulating QS, surface

motility, biofilm formation and virulence. The apparently contradictory regulatory impact of

abaM and *abaI* mutations that result in either increased or no AHL production respectively

on the expression of genes such as the *csu* cluster can be explained as follows. In an *abaI*

mutant (no AHLs), *abaM* expression is reduced and hence *csu* expression is increased. In an

abaM mutant *csu* expression is also increased as AbaM is absent (**Fig. 7**). Further work will

be required to elucidate the biochemical function and mechanism of action of AbaM and

the RsaM protein family in general.

MATERIALS AND METHODS

Strains and growth conditions. The strains and plasmids used are listed in **Table S1.** *A.*

baumannii AB5075 (29) and the isogenic *abaI*::T26 and *abaM*::T26 mutants were obtained

from the transposon library available from the University of Washington (44). *A. baumannii*

was routinely grown in low-sodium chloride (5g/L) lysogeny broth (LS-LB). OHC12 was

synthesized as described before (46). The opaque and translucent phases of the wild type,

and mutants were separated as described by Tipton *et al* (11) after growth on phase-

observation LB (PO-LB) plates and observation of colonies under light microscopy using

oblique indirect illumination.

Construction of a genetically complemented *abaM*::T26 strain. Plasmid pMQ557M (**Table**

S1) was obtained by digesting pMQ557 with *Pml*I (to remove the genes required for yeast

replication) and re-ligating the resulting large linear product. The *abaM* gene plus 768 bp

from its upstream region (containing the predicted native promoter) were amplified by PCR

using primers listed in **Table S2**. The PCR fragments were digested with *Bam*HI and *Kpn*I and

ligated in the multiple cloning site (MCS) of both pMQpMQ557M and introduced into

abaM::T26 by electroporation. The stability of the vector pMQ557M and *abaM*

complementing plasmid pMQ_*abaM* in both A5075 and the *abaM* mutant mutant were

confirmed by repeated daily subculture and plating out on LB agar with and without

hygromycin (125μg/ml) to determine viable counts as cfu/ml (**Fig. S6**).

Construction of *abaM::luxCDABE* and *abaI::luxCDABE* promoter fusions. The *abaR* gene

and the intergenic region between *abaR* and *abaM* (for the *abaM* fusion) or the region

between *abaR* and the *abaI* (for the *abaI* fusion) were amplified by PCR and ligated in

pGEM®-T Easy using the pGEM®-T Easy Vector System (Promega). The resulting plasmids

and the promoterless *luxCDABE* operon were digested with *Kpn*I and *Bam*HI and ligated in

order to introduce the *lux* operon downstream of the predicted promoter of *abaM* or *abaI*.

These constructs were transferred into the MCS of the miniTn*7*T in pUC18T-miniTn*7*T-Hyg[R]

plasmid (**Table S1**) after digesting with *Not*I and *Pst*I and ligating the corresponding

fragments.

MiniTn*7*T-based constructs were inserted into *A. baumannii* through four-parental

conjugation. Briefly, PBS-washed overnight cultures of the *E. coli* DH5α donor strain

(containing pUC18T-mini-Tn*7*T_Hyg[R]_*abaR*_P*abaM*::*lux* or the pUC18T-mini-

Tn*7*T_Hyg[R]_*abaR*_P*abaI*::*lux*), the *E. coli* DH5α helper strain (containing pUX-B13), the *E. coli*

DH5α mobilizable strain (containing pRK600) and the *A. baumanii* recipient strain were

mixed in a 1:1:1:1 ratio and grown on LB agar prior to counterselection with hygromycin

(125 µg/ml for miniTn*7* selection) and gentamicin (100 µg/ml).

The miniTn*7* transposon (45) carrying the *abaM* promoter-*lux* operon fusion was inserted

into *A. baumannii* AB5075 and the isogenic *abaI*::T26 and *abaM*::T26 mutants; while the

abaI promoter-*lux* fusion was inserted into AB5075 and the isogenic *abaI*:T26 mutant.

Bioluminescence output from the reporter fusions as a function of bacterial growth was

quantified using an Infinite 200 PRO (Tecan Diagnostics) plate-reader over 24 h and OD$_{600}$

and relative light units RLUs were recorded every 30 min. When required OHC12 was added

at 200 nM unless otherwise stated.

Biofilm assays. Strains to be tested were inoculated into 1.5 ml polypropylene

microcentrifuge tubes in LS-LB with or without OHC12 and incubated under static conditions

at 37°C for 24 h. Biofilms were quantified by staining with 0.25% crystal violet, extracting

with ethanol and recording the absorbance at A585.

Surface motility assays. Surface motility was quantified as previously described (11) on LS-

LB plates with or without OHC12 and containing 0.3% Eiken agar. Plates were incubated at

30°C for 16 h.

AHL extraction and detection. Cell-free supernatants from cultures grown in LS-LB under

static conditions at 37°C were sterile-filtered and extracted with acidified ethyl acetate.

Extracts were evaporated to dryness and subjected to by liquid chromatography with

tandem mass spectrometry (LC-MS/MS) as previously described (46).

G. mellonella **killing assays.** *G. mellonella* larvae (Trularv[TM]) were obtained from

BioSystems Technology Ltd, U.K. Assays were performed as described previously (11).

Briefly, *Acinetobacter* (2×10^4 or 2×10^5 cfu) were injected into the larval hemolymph,

incubated at 37°C and the larvae monitored for viability. At least 10 larvae were used for

each strain and assay.

Total RNA extraction and RNA-seq. Bacteria were cultured in LS-LB under static conditions

at 37°C for 18h. Cells were resuspended in RNAprotect (Qiagen) prior to extracting total

RNA using an RNeasy minikit (QiAgen). After treatment with DNA-free (Invitrogen) the

absence of DNA contamination was conformed using by PCR and the quality and quantity of

the RNA samples was established using a 2100 Bioanalyzer (Agilent). Samples were sent for

150 bp paired-end sequencing via an Illumina platform and bioinformatic analysis, to

NovoGene (Hong Kong, China). The data have been deposited in NCBI's Gene Expression

Omnibus (50) and are accessible through GEO Series accession number GSE151925

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE151925;   token   for

reviewers: **wxqxkwygphkbliv**).

Quantitative real-time PCR (qPCR). Complementary DNA (cDNA) synthesis and qPCR were

carried out using LunaScript RT Supermix and Luna Universal qPCR Master Mix (New

England BioLabs), respectively. The oligonucleotides used for qPCR are listed in **Table S2** and

qPCR was carried out in triplicate using a 7500 Real-Time PCR System (Thermofisher).

Negative controls lacking template or RNA incubated without reverse transcriptase were

included. The housekeeping gene *rpoB* was used as endogenous control for normalization.

Reverse transcription PCR (RT_PCR). cDNA was amplified using Q5 High-Fidelity polymerase

(New England Biolabs) using specific primers annealing in the coding region of each gene.

Genomic DNA, extracted using DNeasy Blood and Tissue kit (QIAgen), was used as a positive

control. The PCR products were run in a 1.5% agarose electrophoresis gel before imaging

under UV light using a Gel Doc XR+ Imager (Bio-Rad).

ACKNOWLEDGEMENTS

We thank Nigel M. Halliday for AHL quantification and Alex Truman for AHL synthesis,

Robert M.Q. Shanks (University of Pittsburgh) for kindly providing the pMQ557 plasmid and

Prof. Philip N. Rather from Emory University for his guidance and advice on *Acinetobacter*

phase variation. This work was supported via Wellcome Trust joint senior investigator

awards to MRA and PW (grant nos. 103882 and 103884). MLM was supported by a

University of Nottingham postgraduate doctoral studentship.

REFERENCES

1.   Peleg AY, Seifert H, Paterson DL. 2008. *Acinetobacter baumannii*: Emergence of a

successful pathogen. Clin Microbiol Rev 21:538–582.

2.   Lin M-F, Lan C-Y. 2014. Antimicrobial resistance in *Acinetobacter baumannii*: From

bench to bedside. World J Clin cases 2:787–814.

3.   World Health Organization. 2017. WHO publishes list of bacteria for which new

antibiotics are urgently needed. Saudi Med J 38(4): 444–445.

4.   Gaddy JA, Tomaras AP, Actis LA. 2009. The *Acinetobacter baumannii* 19606 OmpA

protein plays a role in biofilm formation on abiotic surfaces and in the interaction of

this pathogen with eukaryotic cells. Infect Immun 77:3150–3160.

5.   Russo TA, Luke NR, Beanan JM, Olson R, Sauberan SL, MacDonald U, Schultz LW,

Umland TC, Campagnari AA. 2010. The K1 capsular polysaccharide of *Acinetobacter*

baumannii strain 307-0294 is a major virulence factor. Infect Immun 78:3993–4000.

6.   Jin JS, Kwon S-O, Moon DC, Gurung M, Lee JH, Kim S Il, Lee JC. 2011. *Acinetobacter*

baumannii secretes cytotoxic outer membrane protein a via outer membrane

vesicles. PLoS One 6:e17027.

7.   Runci F, Gentile V, Frangipani E, Rampioni G, Leoni L, Lucidi M, Visaggio D, Harris G,

Chen W, Stahl J, Averhoff B, Visca P. 2019. Contribution of active iron uptake to

Acinetobacter baumannii pathogenicity. Infect Immun 87(4):e00755-18

8.   Jacobs AC, Hood I, Boyd KL, Olson PD, Morrison JM, Carson S, Sayood K, Iwen PC,

Skaar EP, Dunman PM. 2010. Inactivation of phospholipase D diminishes

Acinetobacter baumannii pathogenesis. Infect Immun 78:1952–1962.

9. Bentancor L V., Camacho-Peiro A, Bozkurt-Guzel C, Pier GB, Maira-Litran T. 2012. Identification of Ata, a multifunctional trimeric autotransporter of *Acinetobacter baumannii*. J Bacteriol 194:3950–3960.

10. Eijkelkamp BA, Stroeher UH, Hassan KA, Elbourne LDH, Paulsen IT, Brown MH. 2013. H-NS plays a role in expression of *Acinetobacter baumannii* virulence features. Infect Immun 81:2574.

11. Tipton KA, Dimitrova D, Rather PN. 2015. Phase-variable control of multiple phenotypes in *Acinetobacter baumannii* strain AB5075. J Bacteriol 197:2593–2599.

12. Liou M-L, Soo P-C, Ling S-R, Kuo H-Y, Tang CY, Chang K-C. 2014. The sensor kinase BfmS mediates virulence in *Acinetobacter baumannii*. J Microbiol Immunol Infect 47:275–281.

13. Cerqueira GM, Kostoulias X, Khoo C, Aibinu I, Qu Y, Traven A, Peleg AY. 2014. A global virulence regulator in *Acinetobacter baumannii* and Its control of the phenylacetic acid catabolic pathway. J Infect Dis 210:46–55.

14. Morris FC, Dexter C, Kostoulias X, Uddin MI, Peleg AY. 2019. The mechanisms of disease caused by *Acinetobacter baumannii*. Front Microbiol 10:1601.

15. Rutherford ST, Bassler BL. 2012. Bacterial quorum sensing: its role in virulence and possibilities for its control. Cold Spring Harb Perspect Med 2:a012427.

16. Williams P, Winzer K, Chan WC, Cámara M. 2007. Look who's talking: communication and quorum sensing in the bacterial world. Philos Trans R Soc Lond B Biol Sci 362:1119–34.

17. Kamaraju S, Ch S, Ratnakar KS, Sritharan V. 2019. Quorum sensing system in *Acinetobacter baumannii* : a potential target for new drug development. J Appl

Microbiol jam.14330.

18.   Niu C, Clemmer KM, Bonomo RA, Rather PN. 2008. Isolation and characterization of

an autoinducer synthase from *Acinetobacter baumannii*. J Bacteriol 190:3386–92.

19.   Anbazhagan D, Mansor M, Yan GOS, Md Yusof MY, Hassan H, Sekaran SD. 2012.

Detection of quorum sensing signal molecules and identification of an autoinducer

synthase gene among biofilm forming clinical isolates of *Acinetobacter* spp. PLoS One

7:e36696.

20.   Clemmer KM, Bonomo RA, Rather PN. 2011. Genetic analysis of surface motility in

Acinetobacter baumannii. Microbiology 157:2534.

21.   Bhuiyan MS, Ellett F, Murray GL, Kostoulias X, Cerqueira GM, Schulze KE, Mahamad

Maifiah MH, Li J, Creek DJ, Lieschke GJ, Peleg AY. 2016. *Acinetobacter baumannii*

phenylacetic acid metabolism influences infection outcome through a direct effect on

neutrophil chemotaxis. Proc Natl Acad Sci U S A 113:9599–604.

22.   Fernandez-Garcia L, Ambroa A, Blasco L, Bleriot I, López M, Alvarez-Marin R,

Fernández-Cuenca F, Martinez-Martinez L, Vila J, Rodríguez-Baño J, Garnacho-

Montero J, Cisneros JM, Pascual A, Pachón J, Bou G, Smani Y, Tomás M. 2018.

Relationship between the quorum network (sensing/quenching) and clinical features

of pneumonia and bacteraemia caused by *A. baumannii*. Front Microbiol 9:3105.

23.   Michalska K, Chhor G, Clancy S, Jedrzejczak R, Babnigg G, Winans SC, Joachimiak A.

2014. RsaM: a transcriptional regulator of *Burkholderia* spp. with novel fold. FEBS J

281:4293–4306.

24.   Mattiuzzo M, Bertani I, Ferluga S, Cabrio L, Bigirimana J, Guarnaccia C, Pongor S,

Maraite H, Venturi V. 2011. The plant pathogen *Pseudomonas fuscovaginae* contains

two conserved quorum sensing systems involved in virulence and negatively

regulated by RsaL and the novel regulator RsaM. Environ Microbiol 13:145–162.

25.   Uzelac G, Patel HK, Devescovi G, Licastro D, Venturi V. 2017. Quorum sensing and

RsaM regulons of the rice pathogen *Pseudomonas fuscovaginae*. Microbiology

163:765–777.

26.   Chen R, Barphagha IK, Karki HS, Ham JH. 2012. Dissection of quorum-sensing genes in

Burkholderia glumae reveals non-canonical regulation and the new regulatory gene

tofM for toxoflavin production. PLoS One 7:e52150.

27.   Le Guillouzer S, Groleau M-C, Déziel E. 2018. Two *rsaM* homologues encode central

regulatory elements modulating quorum sensing in *Burkholderia thailandensis*. J

Bacteriol 200(14):e00727-17.

28.   Silja Inhülsen. 2011. Investigations of the quorum sensing circuitry in *Burkholderia*

cenocepacia H111. PhD thesis. University of Zürich, Zürich.

29.   Jacobs AC, Thompson MG, Black CC, Kessler JL, Clark LP, McQueary CN, Gancz HY,

Corey BW, Moon JK, Si Y, Owen MT, Hallock JD, Kwak YI, Summers A, Li CZ, Rasko DA,

Penwell WF, Honnold CL, Wise MC, Waterman PE, Lesho EP, Stewart RL, Actis LA,

Palys TJ, Craft DW, Zurawski D V. 2014. AB5075, a highly virulent isolate of

Acinetobacter baumannii, as a model strain for the evaluation of pathogenesis and

antimicrobial treatments. MBio.

30.   Mayer C, Muras A, Romero M, López M, Tomás M, Otero A. 2018. Multiple quorum

quenching enzymes are active in the nosocomial pathogen *Acinetobacter baumannii*

ATCC17978. Front Cell Infect Microbiol 8:310.

31.   Ahlgren NA, Harwood CS, Schaefer AL, Giraud E, Greenberg EP. 2011. Aryl-

homoserine lactone quorum sensing in stem-nodulating photosynthetic

bradyrhizobia. Proc Natl Acad Sci U S A 108:7183–7188.

32. Gelencsér Z, Galbáts B, Gonzalez JF, Choudhary KS, Hudaiberdiev S, Venturi V, Pongor

S. 2012. Chromosomal arrangement of ahl-driven quorum sensing circuits in

Pseudomonas. ISRN Microbiol 2012:1–6.

33. Rampioni G, Schuster M, Greenberg EP, Bertani I, Grasso M, Venturi V, Zennaro E,

Leoni L. 2007. RsaL provides quorum sensing homeostasis and functions as a global

regulator of gene expression in *Pseudomonas aeruginosa*. Mol Microbiol 66:1557–

1565.

34. Bondí R, Longo F, Messina M, D'Angelo F, Visca P, Leoni L, Rampioni G. 2017. The

multi-output incoherent feedforward loop constituted by the transcriptional

regulators LasR and RsaL confers robustness to a subset of quorum sensing genes in

Pseudomonas aeruginosa. Mol Biosyst 13:1080–1089.

35. Wand ME, Bock LJ, Turton JF, Nugent PG, Sutton JM. 2012. *Acinetobacter baumannii*

virulence is enhanced in *Galleria mellonella* following biofilm adaptation. J Med

Microbiol 61:470–477.

36. Bhargava N, Sharma P, Capalash N. 2010. Quorum sensing in *Acinetobacter* : an

emerging pathogen. Crit Rev Microbiol 36:349–360.

37. Kang Y-S, Park W. 2010. Contribution of quorum-sensing system to hexadecane

degradation and biofilm formation in *Acinetobacter* sp. strain DR1. J Appl Microbiol

109:no-no.

38. Peleg AY, Jara S, Monga D, Eliopoulos GM, Moellering RC, Jr., Mylonakis E. 2009.

Galleria mellonella as a model system to study *Acinetobacter baumannii* pathogenesis

and therapeutics. Antimicrob Agents Chemother 53:2605.

39. Tomaras AP, Dorsey CW, Edelmann RE, Actis LA. 2003. Attachment to and biofilm

formation on abiotic surfaces by *Acinetobacter baumannii*: involvement of a novel

chaperone-usher pili assembly system. Microbiology 149:3473–3484.

40. Rumbo-Feal S, Gómez MJ, Gayoso C, Álvarez-Fraga L, Cabral MP, Aransay AM,

Rodríguez-Ezpeleta N, Fullaondo A, Valle J, Tomás M, Bou G, Poza M. 2013. Whole

transcriptome analysis of *Acinetobacter baumannii* assessed by RNA-sequencing

reveals different mRNA expression profiles in biofilm compared to planktonic cells.

PLoS One 8:e72968.

41. Pakharukova N, Tuittila M, Paavilainen S, Malmi H, Parilova O, Teneberg S, Knight SD,

Zavialov A V. 2018. Structural basis for *Acinetobacter baumannii* biofilm formation.

Proc Natl Acad Sci U S A 115:5558–5563.

42. Rumbo-Feal S, Pérez A, Ramelot TA, Álvarez-Fraga L, Vallejo JA, Beceiro A, Ohneck EJ,

Arivett BA, Merino M, Fiester SE, Kennedy MA, Actis LA, Bou G, Poza M. 2017.

Contribution of the *A. baumannii A1S_0114* gene to the interaction with eukaryotic

cells and virulence. Front Cell Infect Microbiol 7:108.

43. Gerdt JP, Wittenwyler DM, Combs JB, Boursier ME, Brummond JW, Xu H, Blackwell

HE. 2017. Chemical interrogation of LuxR-type quorum sensing receptors reveals new

insights into receptor selectivity and the potential for interspecies bacterial signaling.

ACS Chem Biol 12:2457–2464.

44. Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, Huse HK, Zurawski

D V, Brittnacher MJ, Manoil C. 2015. Resources for genetic and genomic analysis of

emerging pathogen *Acinetobacter baumannii*. J Bacteriol 197:2027–35.

45. Choi K-H, Gaynor JB, White KG, Lopez C, Bosio CM, Karkhoff-Schweizer RR, Schweizer

HP. 2005. A Tn*7*-based broad-range bacterial cloning and expression system. Nat

Methods 2:443–448.

46.   Ortori CA, Dubern JF, Chhabra SR, Cámara M, Hardie K, Williams P, Barrett DA. 2011.

Simultaneous quantitative profiling of N-acyl-L-homoserine lactone and 2-alkyl-4(1H)-

quinolone families of quorum-sensing signaling molecules using LC-MS/MS. Anal

Bioanal Chem. 399:839-50.

47.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. 32:1792-1797

48.   Robert X, Gouet P. 2014. Deciphering key features in protein structures with the new

ENDscript server. Nucleic Acids Res;42(Web Server issue):W320-W324.

49.   Ng CK, How KY, Tee KK, Chan KG. 2019. Characterization and transcriptome studies of

autoinducer synthase gene from multidrug resistant *Acinetobacter baumannii* strain

863.*Genes (Basel)* 10(4):282

50.   Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene

expression and hybridization array data repository. *Nucleic Acids Res* 30(1):207-10

FIGURE LEGENDS

FIG 1 (A) Schematic of the *abaRMI* quorum sensing locus in *A. baumannii* AB5075 showing the organization of the three QS genes and their orientation. Green boxes represent predicted *lux* boxes. Curved arrows represent the predicted *abaM* and *abaI* promoters. **(B)** Multiple sequence alignment of *A. baumannii* AB5075 AbaM (Abau) with previously characterized orthologs in other bacterial species: Bcen: *Burkholderia cenocepacia* J2315 *Bc*RsaM, Pfus: *Pseudomonas fuscovaginae* UPB0736 RsaM, Btha-2: *Burkholderia thailandensis* E264 RsaM-2, Bglu: *Burkholderia glumae* BGR1 TofM, *Burkholderia thailandensis* E264 RsaM-1. The MUSCLE algorithm (47) was used to create the alignment and ESPript (48) to render residue similarities and generate the final figure. White characters in a red background indicate conserved residues. Red residues indicate conservative substitutions. Blue frames indicate highly conserved regions. The secondary structures in *B. cenocepacia Bc*RsaM (PDB entry 4O2H) is displayed above the alignment. η: 3_{10}-helix, α: α-helices, β: β-strands, **TT**: strict β-turns, **TTT**: strict α-turns.

FIG 2 AHL production in wild-type, *abaI*::T26 and *abaM*::T26. **(A)** Chemical structures of the AHLs produced by *A. baumannii* AB5075**.** Quantification of OHC12 **(B)** and OHC10 **(C)** production throughout growth. Error bars represent the standard deviation between three biological replicates. Asterisks indicate statistically significant differences: **, $p \leq 0.01$; ***, $p \leq 0.001$

FIG 3 Phenotypic characterization of the *Acinetobacter abaM* and *abaI* mutants**. (A)** Surface motility on agar **(B)** Biofilm formation on polypropylene. For both biofilm and surface

motility assays, error bars indicate standard deviation. Asterisks indicate statistically

significant differences compared with the wild-type AB5075 strain: ***, $p \leq 0.001$; ****, $p \leq$

0.0001. **(C)** *Galleria mellonella* larvae killing after inoculation of approximately 2×10^4 (left)

or 2×10^5 (right) CFU/larva. Each graph represents data from 3 independent biological

replicates together. At least 30 larvae were used for each strain and assay. None of the

control larvae died after 5 days. Asterisks indicate statistically significant differences

compared with the wild-type AB5075 strain: ****, $p \leq 0.0001$.

FIG 4 Comparison of the transcriptomes of the *abaI*::T26 and *abaM*::T26 mutants. Genes

differentially expressed in **(A)** *abaM*::T26 **(B)** *abaI*::T26 compared with wild-type. Blue

circles indicate upregulated genes. Red circles indicate downregulated genes. Grey circles

represent genes where changes in expression are unlikely to be biologically significant. **(C)**

Venn diagram showing that AbaM regulates genes that are both QS-dependent and QS-

independent.

FIG 5 Validation ABUW_3773 and *csu* expression by quantitative real-time PCR**.** Relative

expression of *csuA/B* and *ABUW_3773* in   mutants compared with the wild-type AB5075

strain. The expression was normalized in relation to an endogenous control gene (*rpoB*).

Error bars indicate standard deviation between 3 independent biological replicates.

Hashtags (#) indicate a biologically significant difference ($|\log_2(\text{fold change})| \geq 1$)

compared to the wild-type.

FIG 6 Expression of *abaM* and *abaI*. **(A)** *abaM* promoter activity in *A. baumannii* A5075 wild

type, *abaM*::T26 and *abaI*::T26 relative to the wild-type strain. **(B)** *abaM* promoter activity

in response to exogenous OHC12 as a function of growth (RLU/OD$_{600}$). **(C)** *abaI* promoter

activity in wild type, *abaI* mutant and in response to exogenous OHC12 as a function of

growth (RLU/OD$_{600}$). Error bars indicate standard deviation between 3 independent

biological replicates. Asterisks indicate statistically significant differences compared to the

wild-type AB5075 strain: ***, $p \leq 0.001$; ****, $p \leq 0.0001$.

FIG 7. Proposed model for the QS/AbaM incoherent feed forward loop (IFFL) in *A.

baumannii 5075.** AbaR activated by OHC12 positively activates expression of *abaM* and *abaI*

and hence OHC12 production. AbaM is negatively autoregulated and also represses

expression of both *abaR* and *abaI*. Under the growth conditions used here, AbaM negatively

regulates surface motility, biofilm and Csu pili but is required for virulence as *abaM* mutants

are avirulent in *Galleria mellonella* larvae. Green arrows and red blunt ended lines represent

positive and negative regulation respectively

FIG 1 (A) Schematic of the *abaRMI* quorum sensing locus in *A. baumannii* AB5075 showing the organization of the three QS genes and their orientation. Green boxes represent predicted *lux* boxes. Curved arrows represent the predicted *abaM* and *abaI* promoters. **(B)** Multiple sequence alignment of *A. baumannii* AB5075 AbaM (Abau) with previously characterized orthologs in other bacterial species: Bcen: *Burkholderia cenocepacia* J2315 *Bc*RsaM, Pfus: *Pseudomonas fuscovaginae* UPB0736 RsaM, Btha-2: *Burkholderia thailandensis* E264 RsaM-2, Bglu: *Burkholderia glumae* BGR1 TofM, *Burkholderia thailandensis* E264 RsaM-1. The MUSCLE algorithm (47) was used to create the alignment

and ESPript (48) to render residue similarities and generate the final figure. White

characters in a red background indicate conserved residues. Red residues indicate

conservative substitutions. Blue frames indicate highly conserved regions. The secondary

structures in *B. cenocepacia Bc*RsaM (PDB entry 4O2H) is displayed above the alignment. η:

3_{10}-helix, α: α-helices, β: β-strands, **TT**: strict β-turns, **TTT**: strict α-turns.

FIG 2 AHL production in wild-type, *abaI*::T26 and *abaM*::T26. **(A)** Chemical structures of the

AHLs produced by *A. baumannii* AB5075**.** Quantification of OHC12 **(B)** and OHC10 **(C)**

production throughout growth. Error bars represent the standard deviation between three

biological replicates. Asterisks indicate statistically significant differences: **, $p \leq 0.01$; ***,

$p \leq 0.001$**.**

FIG 3 Phenotypic characterization of the *Acinetobacter abaM* and *abaI* mutants. **(A)** Surface

motility on agar **(B)** Biofilm formation on polypropylene. For both biofilm and surface

motility assays, error bars indicate standard deviation. Asterisks indicate statistically

significant differences compared with the wild-type AB5075 strain: ***, $p \leq 0.001$; ****, $p \leq$

0.0001. **(C)** *Galleria mellonella* larvae killing after inoculation of approximately 2×10^4 (left)

or 2×10^5 (right) CFU/larva. Each graph represents data from 3 independent biological

replicates together. At least 30 larvae were used for each strain and assay. None of the

control larvae died after 5 days. Asterisks indicate statistically significant differences

compared with the wild-type AB5075 strain: ****, $p \leq 0.0001$.

FIG 4 Comparison of the transcriptomes of the *abaI*::T26 and *abaM*::T26 mutants. Genes

differentially expressed in **(A)** *abaM*::T26 or **(B)** *abaI*::T26 compared with wild-type. Blue

circles indicate upregulated genes. Red circles indicate downregulated genes. Grey circles

represent genes where changes in expression are unlikely to be biologically significant. **(C)**

Venn diagram showing that AbaM regulates genes that are both QS-dependent and QS-

independent.

FIG 5 Validation ABUW_3773 and *csu* expression by quantitative real-time PCR. Relative

expression of *csuA/B* and *ABUW_3773* in *abaI*::T26 and *abaM*::T26 mutants compared with

the wild-type AB5075 strain. The expression was normalized in relation to an endogenous

control gene (*rpoB*). Error bars indicate standard deviation between 3 independent

biological replicates. Hashtags (#) indicate a biologically significant difference (|log2(fold

change)| ≥ 1) compared to the wild- type.

674
675
FIG 6 Expression of *abaM* and *abaI*. **(A)** *abaM* promoter activity in *A. baumannii* A5075 wild

type, *abaM*::T26 and *abaI*::T26 relative to the wild-type strain. **(B)** *abaM* promoter activity

in response to exogenous OHC12 (200 nM) as a function of growth (RLU/OD$_{600}$). **(C)** *abaI*

promoter activity in wild type, *abaI* mutant and in response to exogenous OHC12 as a

function of growth (RLU/OD$_{600}$). Acn, acetonitrile solvent control. Error bars indicate

standard deviation between 3 independent biological replicates. Asterisks indicate

statistically significant differences compared to the wild-type AB5075 strain: ***, $p \leq 0.001$;

****, $p \leq 0.0001$.

FIG 7 Proposed model for the QS/AbaM incoherent feed forward loop (IFFL) in *A.*
baumannii **5075.** AbaR activated by OHC12 positively activates expression of *abaM* and *abaI*
and hence OHC12 production*.* AbaM is negatively autoregulated and also represses
expression of both *abaR* and *abaI*. Under the growth conditions used here, AbaM negatively
regulates surface motility, biofilm and Csu pili but is required for virulence as *abaM* mutants
are avirulent in *Galleria mellonella* larvae. Green arrows and red blunt ended lines represent
positive and negative regulation respectively.