Full title: Substrate traits govern the assembly of microbial community engaged in

metabolic division of labor

Running title: Substrate traits govern community assembly

Miaoxiao Wang[1, 2, 3, 4], Xiaoli Chen[1, 5], Yong Nie[1#] and Xiao-Lei Wu[1, 5, 6#]

[1] College of Engineering, Peking University, Beijing 100871, China

[2] Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland

[3] Department of Environmental Microbiology, Eawag, Dübendorf, Switzerland

[4] College of Architecture and Environment, Sichuan University, Chengdu, China

[5] Institute of Ocean Research, Peking University, Beijing 100871, China

[6] Institute of Ecology, Peking University, Beijing 100871, China

[#]Corresponding author: Research Scientist, College of Engineering, Peking University.

Tel: +86 10-62759047; Fax: +86 10-62759047; E-mail: nieyong@pku.edu.cn

[#]Corresponding author: Professor, College of Engineering, Peking University.

Tel: +86 10-62759047; Fax: +86 10-62759047; E-mail: xiaolei_wu@pku.edu.cn

Abstract

Metabolic division of labor (MDOL) is widespread in nature, whereby a complex metabolic pathway is shared between different strains within a community for mutual benefit. However, little is known about how microbial populations engaged in MDOL assemble into mutually beneficial communities. We hypothesized that when degradation of an organic compound is carried out via MDOL, substrate concentration and its toxicity modulate the benefit allocation between the two microbial populations, thus governing the assembly of this community. We tested this hypothesis by combining mathematical modelling with experiments using an engineered synthetic microbial community. Numerous modelling analyses suggested that the proportion of the population executing the first metabolic step can be simply estimated by Monod-like formulas governed by substrate traits. The model and the proposed formula quantitatively predicted the assembly of our synthetic community composed of two strains degrading salicylate through MDOL. Individual-based modelling and colony pattern formation assays further indicated that our rule is also applicable to estimating community assembly in spatially structured environments. Our results demonstrate that the structure of the microbial communities can be quantitatively predicted from simple environmental factors, such as substrate concentration and its toxicity, which provides novel perspectives on understanding the assembly of natural communities, as well as insights into how to manage artificial microbial systems.

Introduction

In natural environments, microorganisms rarely live autonomously; instead, they interact with other individuals to form complex communities, in which they secrete a variety of toxins to compete with each other, or share metabolites to mutually benefit their survival. Among diverse modes of microbial interaction, metabolic division of labor (MDOL) is one of the most widespread phenomena, where distinct populations perform different but complementary steps of the same metabolic pathway[1-4]. MDOL controls numerous ecologically and environmentally important biochemical processes. For example, the nitrification pathway is often operated by the collaboration between two populations, in which the first group oxidizes ammonia to nitrite, before being converted to nitrate by another bacterial group[5]. Similarly, during the Deepwater Horizon oil spill Gulf of Mexico (2010), complete degradation of polycyclic aromatic hydrocarbons (PAHs) required the partitioning of key pathway steps into several bacterial groups[6]. Since each population only contains a subset of genetic components required for the overall pathway, MDOL is thought to be a key evolutionary strategy to reduce metabolic burden of each individual[7, 8]. In addition, bacteria have been found to use this strategy to reduce the inter-enzyme competition and the inhibitory effects of intermediates, thus improving the systematic metabolic efficiency[9]. Furthermore, MDOL vastly enhances functional differentiation of microbial communities, contributing to genetic diversity in complex ecosystem[7, 8]. Nevertheless, how microbial communities engaged in MDOL assemble in the first place remains ill-defined. Understanding the assembly rules of communities engaged in

MDOL is fundamental for better evaluating their contribution to the natural biochemical cycle.

One important aspect of microbial metabolism implemented by MDOL is the degradation of a variety of organic compounds, including PAHs [10, 11], pesticide [12-15], plastics [16], antibiotics [17], or lignocellulose [18]. Bacterial degradation of these complex substrates is usually mediated by long metabolic pathways via a number of intermediates. While these pathways often remain intact within one population, they are frequently found segregated across different members within a community, whereby the intermediates are being secreted by one species, before being degraded as substrate by another species for further processing along the same pathway. Typical examples include syringate degradation via sequential cross-feeding between *Acetobacterium woodii* and *Pelobacter acidigallici* [10], phenanthrene degradation between *Marinobacter* sp. N4 and other PAH-degrading microbes in marine environments [11], as well as atrazine degradation through MDOL within four bacterial species[14]. In most of these cases, resource allocation among different community members hold one typical feature: direct carbon sources that support bacterial growth, such as small organic acids or coenzyme A, are normally present as the final product of a pathway (Supplementary Figure 1); therefore, the population performing the last steps can preferentially acquire and privatize these nutrients (which we henceforth call *product privatization*), thus acquiring the greater benefit, while other members have to collect nutrients released from this population (Figure 1; left of the first row). We thus hypothesized that this uneven allocation of limiting resource largely determines the

relative fitness of strains engaged in MDOL, and thus govern the assembly of these

communities.

The findings in recent studies on MDOL communities are largely consistent with our

hypothesis. For instance, one earlier study developed a dual-species community for

biodegradation of the insecticide parathion [15]. In this consortium, an *Escherichia coli*

strain SD2 was responsible for hydrolyzing parathion, yielding two intermediates

including p-nitrophenol, while another *Pseudomonas putida* strain KT2440 was

responsible for metabolizing p-nitrophenol. That study found that the strain KT2440

that largely dominated the biofilm formed by this community. Another study

investigated the interactions among five bacterial species in a cellulose-degrading

community, which also found that the strains responsible for the last step of the

cellulose degradation constituted more than 60 percent of the total population [19]. In

addition, one study used a mathematical model to capture the dynamics of a MDOL

community degrading fenitrothion, in which the final product, methylhydroquinone

served as the common resource for the two community members [12]. That study found

that the strain executing the last step overwhelmingly dominated the community in most

of the simulations. On the basis of this finding, the study proposed that the coexistence

of two members requires the second member excreting sufficient amounts of

methylhydroquinone in order to support the growth of another member. Overall, these

findings suggest that our hypothesized 'Embezzler behavior' may be a common

phenomenon that occurs when a microbial community adopt MDOL strategy to degrade

organic compounds. Since 'product privatization' is a purely selfish trait, we named the

final strain at the end of a MDOL-enabled pathway the 'Embezzler', analogous to a

human worker responsible for the final step of an assembly line, who pockets the final

product and fails to share profits with other workers.

We also derived two corollaries based on this hypothesis. Firstly, increasing

concentration of the substrate improves the flux of the pathway. Because the Embezzler

only have a limited capacity of consuming the final product, more end product is

released from the 'Embezzler' strain, in turn facilitating the growth of the other

population (Figure 1; Right of the first row). In natural environments, the concentration

of the initial substrate, such as PAHs and other hydrocarbons, is highly fluctuating over

time and different sites in the marine[20], soil[21] and wastewater[22] environments.

Therefore, it is crucial to understand how this variable environmental factor affect the

relative fitness of strains engaged in MDOL, and thus govern the assembly of these

communities. Secondly, many initial substrates that can be degraded through MDOL,

such as PAHs [23, 24], are toxic to bacterial cells, inhibiting their growth. We proposed

that the population performing the first step is favored when the initial substrate is toxic.

Because its ability of transforming the initial substrate to the intermediates will help it

possess a lower intracellular concentration of the toxic substrate (Figure 1; The second

row), resulted in that the toxic substrate is generally less harmful to Detoxifier than to

Embezzler. Also because of this feature, we named the population executing the first

metabolic step 'Detoxifier'. In summary, in presence of the resource privatization by

the 'Embezzler' population, concentration and toxicity of the substrate can be two key

factors that govern the assembly of organic compound degradation community engaged

in MDOL. The goal of this study is to test this prediction.

Microorganisms often live in spatially structured environments, forming well-

organized biofilms[25, 26]. In these spatially structured habitats, mass diffusion is

largely limited, driving to heterogeneous distribution of the substrates and metabolites,

which have large impacts on the outcomes of the metabolic exchanges between

different microbial cells [27], and further govern the assembly and productivity of such

communities [26, 28-30]. Particularly, since the intermediates and the product are

required to be exchanged between different community members, the assembly of

community engaged in MDOL should be also affected by the limited mass diffusion.

For example, a recent study investigated how a well-defined community executing

toluene degradation through MDOL spatially self-organized on surface, which

suggested that MDOL generally favored strong intermixing between the two strains

involved to facilitate their metabolic exchange [31]. Therefore, it is also important to

investigate how levels of substance diffusion level in the spatially structured

environment affect the assembly of MDOL communities.

Here, we set out to investigate how substrate traits drive assembly of microbial

community engaged in MDOL by combining mathematical modelling and

experimentation using a synthetic microbial community. We also tested whether these

effects are different when the community grows in spatially well-mixed and structured

environments.

Results

Testing of our prediction in well-mixed system

 An ODE model for the assembly of microbial community engaged in MDOL

To test our hypothesis about Embezzler behavior, as well as the effects of substrate

concentration and its toxicity, we first built a mathematical model to simulate the

assembly of a community engaged in metabolic division of labor (MDOL) in a well-

mixed batch culture system. The dimensionless form of this model is composed of 11

ordinary differential equations (ODEs; Eqn [3]-Eqn [13] in Methods). As summarized

in Figure 2A, we considered the degradation of an organic substrate (S) into an

intermediate metabolite (I) by one enzyme (E1), before being degraded to the final

product (P) by a second enzyme (E2). We assumed that two strains carried out this

pathway via MDOL, with the first strain only expressing E1, and the second only

expressing E2. Initially, only S was supplied and the initial concentration was

parameterized by s_0 (nondimensional). In addition, transport of S, I, and P across the

cell membrane was assumed to occur via passive diffusion. Importantly, based on our

hypothesis of 'Embezzler behavior', we assumed that, P, which was synthesized by the

second strain, was the sole available resource for the growth of both strains, while

neither S nor I could be directly used for growth. As a result, the second strain possessed

the advantage of preferentially acquiring the resource, while the first strain only

obtained those growth-limiting resource that were secreted from the second strain.

Therefore, the second strain behaved as an 'Embezzler'. Moreover, biotoxicity of the

substrate was imposed by adding three different toxic terms (Supplementary Table

3;[32]) to the growth function, and the toxic strength was mediated by parameter θ.

Thus, for the scenarios where substrate is assumed to be toxic, the strain executing the

first step behaved as a 'Detoxifier'. Details about the model construction are described

in Supplementary Information S1.

Analysis of the ODE model indicates initial substrate concentration affects the

assembly of MDOL community.

To test our basic hypothesis about Embezzler behavior, as well as our corollary 1 stating

that substrate concentration affects the assembly of the community, we analyzed our

ODE model omitting substrate toxicity (Figure 2A). As the dimensionless model

contains 11 independent parameters (Supplementary Table 4) that may affect the

outcomes of community assembly, we performed a first round of numerical simulations

using 885,735 parameter sets considering realistic value ranges of all the 11 parameters

(Supplementary Information S1.3; Supplementary Table 4). Our analysis showed that

the Embezzler population dominated the steady-state community in all these

simulations (Supplementary Figure 2; no toxic scenarios, i.e., steady-state frequencies

of Detoxifier are lower than 0.5), which was in agreement with our basic hypothesis of

'Embezzler behavior'. We then assessed the relative importance of the involved

parameters by performing multivariate regression analyses, and found that six key

parameters played vital roles in governing the assembly of MDOL community

(Supplementary Table 4; Supplementary Figure 3A; $p < 0.01$ and the fitting coefficient

values over 0.01). Notably, we found that s_0 was one of these key parameters (second

most important according to the absolute value of the fitting coefficient). As shown in

Supplementary Figure 3A and Supplementary Table 4, s_0 positively correlated with

the steady-state proportion of the Detoxifier population, suggesting that a higher initial

substrate concentration favors the Detoxifier, consistent with our corollary 1.

To investigate how initial substrate concentration affects the assembly of a MDOL

community, we performed second round of simulations using newly-designed 163,296

parameter sets that included more testing values of the six key parameters

(Supplementary Information S1.3). We found that when all other five key parameters

were kept constant, the steady-state proportion of the Detoxifier population increased

with an increase of the initial substrate concentration (Figure 2B and 2C), and can be

estimated by a Monod-like formula using s_0 as the function argument (Figure 2C),

$$DF = \frac{Fd_{max} \cdot s_0}{ks + s_0} \tag{1}$$

Here, DF represents the steady-state proportion of the Detoxifier population; Fd_{max}

represents the maximum proportion of the Detoxifier populations when substrate is

non-toxic; ks represents the half-saturation constant. Our analysis indicated that the

simulation results of all tested parameter sets can be accurately fitted to Eqn. [1] (Figure

2D, values of Adjusted R^2 mostly over 0.95), although the best fitting of Fd_{max} and ks

were affected by the values of other five key parameters (Figure 2E and 2F;

Supplementary Information S1.3; Supplementary Table 5; Supplementary Figure 4-5).

Together, these results suggest that, in the absence of substrate toxicity, the proportion

of the Detoxifier population increases nonlinearly with the increase of the initial

substrate concentration, and maintains a maximum value (Fd_{max}).

To investigate why substrate concentration governs the community assembly, we next

analyzed the dynamics of intracellular and extracellular product concentration of the

two populations.

We found that with the increase of initial substrate concentration, the fraction of final

product released by the Embezzler population (that is, the ratio of extracellular to

intracellular concentration) increased (Supplementary Figure 6A-H; Supplementary

Figure 6I, Red dots). As a consequence, the Detoxifier obtained more product from the

environment, resulting in a higher intracellular product concentration, which gradually

approached that of the Embezzler (Supplementary Figure 6A-H; Supplementary Figure

6I, blue dots). Therefore, consistent with our corollary 1, increasing the initial substrate

concentration increases the amount of the final product released by the Embezzler

population, thus favoring the Detoxifier. Moreover, we also found that even if the

substrate concentration was elevated to high levels, the intracellular product

concentration of the Detoxifier never exceeded that of the Embezzler, even if the

substrate concentration was elevated to high levels (Supplementary Figure 6A-H;

Supplementary Figure 6I, blue dots). As a result, Embezzler cells still maintained their

growth advantage from privatizing final product. This result suggests that in the absence

of substrate toxicity, the benefit from product privatization obtained by the Embezzler

strain cannot be completely eliminated by simply increasing initial substrate

concentrations. This observation also matched with our result that the maximum

proportion of the Detoxifier population (Fd_{max}) never exceeded 0.5 (Figure 2F;

Supplementary Figure 5). In addition, based on this mechanism, increasing the P

consuming capacity of Embezzler may cause less release of P to environment, thus

harming the Detoxifier. This corollary is supported by our observation that the value of

Fd_{max} was strongly negatively correlated with the consuming rate of the final product

(Cp), and ks was positively correlated with Cp (Supplementary Table 5;

Supplementary Figure 4A-B; Supplementary Figure 5A-B). In summary, these results

suggested that substrate concentration governs the assembly of microbial community

engaged in MDOL by affecting the amount of the final product supplying from

Embezzler to Detoxifier (Figure 1; the first row).

Analysis of the ODE model indicates that substrate toxicity also affects the assembly of

MDOL community.

To test our corollary 2, we next employed an ODE model that included the parameter

of substrate toxicity to test our corollary 2 (Figure 3A). Applying similar simulation

and analysis method as used in the above section (Supplementary Information S1.3),

we found that in addition to the six key parameters identified above, the toxic strength

(θ) of substrate also played a significant role in governing the assembly of our MDOL

community, exhibiting a significantly positive relationship with the final proportion of

the Detoxifier population (Figure 3B; Supplementary Figure 2-3; Supplementary Table

4), which is in agreement with our corollary 2. We then upgraded Eqn. [1] to

collectively consider the effects of substrate concentration and its toxicity (Figure 3C)

$$DF = \frac{Fd_{max} \cdot s_0}{ks + s_0} \cdot (1 + \frac{Ts_{max} \cdot \theta s_0}{kt + \theta s_0})$$ [2]

In Eqn. [2], we use term $1 + \frac{Ts_{max} \cdot \theta s_0}{kt + \theta s_0}$ to describe the effect of substrate toxicity on the

proportion of the Detoxifier populations, in which Ts_{max} represents the maximum fold

increase of Detoxifier proportion benefiting from the substrate toxicity and ks

represents the half-saturation constant of this toxic effect. This term is positively

affected by the toxic strength (θ) and substrate concentration (s_0), since increasing

either toxic strength or substrate concentration harms population growth, according to

our model assumptions (see Eqn. [12]-[13] in Methods and Supplementary Table 3).

We next tested the applicability of Eqn. [2] on prediction of assembly of MDOL

community by performing numerical simulations using 8,328,096 parameter sets

(Supplementary Information 1.3). These sets included 7,776 different combinations of

other five key parameters and considered three different equation forms

(Supplementary Table 3) that quantified how substrate toxicity affects population

growth. The results indicated that the simulation results also accurately fitted to Eqn.

[2] (Figure 3D; values of Adjusted R^2 mostly over 0.90). We also assessed the main

factors that affect the best fitting values of the parameters in Eqn. [2] (Supplementary

Information S1.3; Supplementary Table 5; Supplementary Figure 7-10). In agreement

with our results from the model omitting substrate toxicity, we found that the value of

Fd_{max} still negatively correlated with the consuming rate of the final product (Cp), and

ks was positively correlated with Cp (Supplementary Table 5; Supplementary Figure

8-9). These results suggest that in the when substrate toxicity was taken into account,

the proportion of the Detoxifier population increased with both the initial concentration

and the toxic strength of the substrate.

To address the mechanisms of how substrate toxicity affects the community assembly,

we next analyzed the dynamics of intracellular and extracellular S and P concentration

of the two populations. As shown in Supplementary Figure 11 , the dynamics of P

largely agrees with the result derived from those non-toxic scenarios, suggesting that

increasing the substrate concentration affects the amount of final product that was

exchanged between Detoxifier and Embezzler in these simulations, thus governing

community assembly. Our analysis of the S dynamics clearly showed that the Detoxifier

population generally maintained a lower intracellular concentration level of S than that

of the Embezzler (Supplementary Figure 12), due to its conversion of S, thus possessing

a growth advantage over the Embezzler population. Based on this mechanism, higher

speed of the first reaction, or lower S transport rate, appears to favor the Detoxifier

population since these two conditions assist Detoxifier in maintaining a lower

intracellular S concentration. Consistent with this corollary, Ts_{max} was significantly

positively correlated with a_1 and significantly negatively correlated with

γ_s (Supplementary Table 5; Supplementary Figure 10). Overall, these results indicated

that the difference in intracellular concentration of substrate is the main reason why

substrate toxicity favors the Detoxifier population (Figure 1; second row).

When we assessed the community structure at different conditions of initial substrate

concentration and substrate toxicity, we found that Detoxifier population are able to

dominate the community when the initial substrate concentration and substrate toxicity

are sufficiently high (that is, its relative proportion exceeded 50% of the community;

Figure 3C; Supplementary Figure 2), suggesting that the benefit from product

privatization of the Embezzler can be neutralized by higher substrate concentration and

higher substrate toxicity. This phenomenon is quantitively characterized by Eqn. [2]:

the maximum Detoxifier proportion in the absence of substrate toxicity (meaning $\theta =$

0), Fd_{max}, never exceed 0.5 (Supplementary Figure 9), but substrate toxicity can assist

Detoxifier in breaking through this constraint, as quantified by the term $1 + \frac{Ts_{max} \cdot \theta s_0}{kt + \theta s_0}$.

In summary, our simulations clearly showed that when a compound degradation pathway is executed through MDOL in a community, both increasing substrate concentration and increasing the toxicity of the substrate govern the favor the Detoxifier population, thus resulting in substrate traits to govern the assembly of the community.

Experimental evaluation of our rule using a liquid culture of a synthetic microbial community engaged in MDOL

To experimentally test the prediction from our ODE model, we engineered a synthetic community composed of two *P. stutzeri* strains, both of which degrade an organic compound, salicylate, via MDOL (Figure 4A). In this synthetic community, strain *P. stutzeri* AN0010 only retained its ability to convert toxic substrate, salicylate [33], behaving as the 'Detoxifier'; the second strain, *P. stutzeri* AN0001, was only able to metabolize the intermediate catechol, but possessed the preferential access to the final product, i.e., pyruvate and acetyl-CoA (Figure 4A), the direct carbon source of both strains, thus behaving as the 'Embezzler'. Details about the strain construction and identification are described in Supplementary Information S3. For simplicity, we henceforth refer to our community as 'SMC-mdol'.

We first used our model to predict the assembly of SMC-mdol. We imposed the experimentally measured or previous reported parameters to our ODE model (Supplementary Table 6), and performed mathematical simulations across 21 different toxic strength of the substrate as well as twelve different initial substrate concentration. We fitted the simulation data to Eqn. [2], thus generating a predictive function that matches with our synthetic system (Figure 4B). To evaluate the toxicity of salicylate,

we quantitively measured the relationship between strain growth rate and concentrations of salicylate (see Supplementary Information S3.4 for measurement details). As shown in Supplementary Figure 13, this relationship mostly follows a reciprocal-form equation, and the measured dimensionless value of its toxic strength (θ) was 0.0032. After inserting this value into the predictive function, we mathematically predicted the effect of substrate concentration on the assembly of SMC-mdol, as indicated by the red line in Figure 4B and 4C. We then cultured the SMC-mdol community in liquid minimal medium supplemented with different concentrations of salicylate, and analyzed the salicylate-dependent changes in structures of this synthetic community. As shown in Supplementary Figure 14, the community dynamics of SMC-mdol is largely consistent with the dynamics of our corresponding ODE simulations (Supplementary Figure 14). In addition, the proportion of Detoxifier population increased from 25.6% ± 2.5% to 61.1% ± 2.6% as a function of initial salicylate concentration (Figure 4C). This shift exhibited the same trends as obtained from our models, suggesting that substrate concentration governs the assembly of MDOL community.

Importantly, when the substrate concentration reached high levels, the Detoxifier population dominated the community (i.e., its relative fraction over 50 %), suggesting that substrate toxicity considerably affected the assembly of the community. Moreover, our predictive function accurately estimated the assembly of SMC-mdol, with a predictive power (Adjusted R^2) of 0.983. Together, these experiments clearly demonstrated that the assembly of microbial community engaged in MDOL can be

estimated by our proposed simple rules regarding concentration and toxicity of the

substrate.

Testing the prediction in spatially structured environments

In the above modeling and experiments, we investigated the assembly rules governing

a MDOL community, principally by assuming that the substances and cells were well-

mixed in the system, resulting in a spatially unstructured environment. However,

microorganisms are frequently observed to live in spatially structured environments,

such as biofilm [34-36]. Previous studies reported that different physical characteristics

between these two habitat types significantly affected the assembly of microbial

communities involved [26, 30, 37, 38]. Therefore, we set out to test whether our rule

derived from the assumption of a well-mixed system can be expanded to estimate the

assembly of a MDOL community in spatially structured environments.

Individual-based modelling of the assembly of MDOL community.

To develop a mathematical framework regarding the assembly of MDOL community

in spatially structured environment, we built an individual-based (IB) model to simulate

the colonization and range expansion of a MDOL community on a surface. The basic

configuration regarding reaction kinetics, microbial growth, and metabolic interactions

was identical to the framework of our ODE model. Moreover, we assumed that the

diffusion of S, I, and P was limited in the IB model, and mediated by their diffusion

coefficients (D_s, D_i, and D_p). Details about the IB model construction and simulations

are described in Supplementary Information S2.

To test our hypothesis, we ran the IB model using the parameters consistent with our

experimental system, but varied the toxic strength (θ) and initial concentration of the

substrate (s_0). We found that when toxicity was not considered and substrate

concentration was low (e.g., $\theta = 0$ and $S_0 = 2.5$ C-mmol/L), the model successfully

reproduced our hypothesis of 'Embezzler behavior'. In this case, 'Embezzler' cells

grew much faster than Detoxifier cells and occupied a large proportion of the available

space (Figure 5A; Supplementary video 1). In addition, either increasing substrate

concentration or introducing substrate toxicity favored the 'Detoxifier' population

(Figure 5A; Supplementary video 1-4). This finding is similar to our observations in

the spatially unstructured system, and is consistent with our two corollaries.

Furthermore, we found that during the colony range expansion, the two cell lineages

(Detoxifier and Embezzler) segregated at frontiers, forming adjacent red and green cell

sectors (Figure 5A; Supplementary video 1-4). Analysis of the distribution of substrate

and metabolites suggested that the development of this colony characteristic was mainly

attributed to the 'active layer effect' reported previously [28]. S is generally supplied

from the outside of the colony and the mass diffusion is limited, resulted in thin active

cell layers depending on the penetration of S, I and P (Supplementary video 1-4).

Consequently, community assembly in the inoculating and expanding regions may

follow different rules. To better quantify how community assembles in these spatially

structured colonies, we separately analyzed the community structures in the inoculating

region and expanding region of the colonies (Supplementary Figure 15). We found that

with the growth of colony, community structures in the inoculating region changed little,

while the community structures in the expanding region shifted over time, gradually

approaching a steady-state (Supplementary Figure 16). Therefore, we next investigated

how substrate traits affect the steady-state structures of the MDOL community in the

expanding regions.

As shown in Figure 5B, the community assembly in the expanding region was

significantly affected by substrate traits, and can be well estimated by the rule (Eqn. [2])

that we proposed for a spatially unstructured system. The assembly process was also

affected by the consuming rate of P, as it significantly affected the values of fitted

parameters, Fd_{max} and ks (Supplementary Figure 17; correlation analysis $p<0.0001$).

Further analysis suggested that increasing initial substrate concentrations assisted

Detoxifier to obtain more product from the environment, thus retaining higher

intracellular product concentrations (Supplementary Figure 18). These results indicate

that in spatially structured environment, substrate concentration governs the assembly

of MDOL community as well by affecting the amount of the final product supplying

from Embezzler cells to Detoxifier cells. Furthermore, we found that the Detoxifier

cells possessed a lower intracellular concentration level of S than that of the Embezzler

cells in our IB simulations (Supplementary Figure 19); higher speed of the first reaction,

or lower S transport rate, also significantly increased the maximum benefit (Ts_{max}) that

Detoxifier cells can obtained from substrate toxicity (Supplementary Figure 20;

correlation analysis $p<0.0001$), same as our results from ODE modelling. Therefore,

lower intracellular S concentration also benefits Detoxifier population when substrate

toxicity was taken into account in a spatially structured environment. Together, our IB

modelling demonstrated that the assembly of our MDOL community in spatially

structured environments can also be estimated by the proposed simple formula

governed by substrate traits.

Experimental evaluation of our rule by culturing our synthetic microbial community in

spatially structured environment.

We next experimentally tested whether substrate traits affect the assembly of a MDOL

community in spatially structured environments. Several studies have reported that type

IV pilus may affected the microbial colony patterns [39-41]. To directly focus on the

effects of substrate traits and avoid the effects of pili, we deleted the *pilA* and *pilB* genes

of the both strains involved in our synthetic community. This design follows other

studies that performed patterning experiments using non-motile strains [42-45]. The

derived community was named as SMC-mdol△*pilAB*. As shown in Fig. 4C, this strain

modification did not change the effect of substrate traits on community assembly in

spatially unstructured system (Figure 4C), as well as the toxic effects of salicylate on

the strains (Supplementary Figure 13).

To test our hypothesis, we cultured SMC-mdol△*pilAB* on an agarose surface to which

salicylate was added at different concentrations. The experimentally observed patterns

of the colonies were very similar to those observed for the simulated patterns (Figure

5C). We next separately assessed the community structures of the entire colonies, as

well as the inoculating region and expanding region of the colonies using image

analysis method. We found that, in the inoculating region, the proportion of Detoxifier

population slightly shifted from $40.9\% \pm 3.5\%$ to $60.0\% \pm 6.0\%$ (Supplementary Figure

21), but in the expanding region, it largely varied from $17.4\% \pm 1.5\%$ to $69.0\% \pm 7.0\%$

(Figure 5D). Importantly, the experimental results of expanding region accurately fitted

to our predictive function, which was derived from Eqn. [2] and IB simulations (using

the same protocol as that in spatially unstructured system; Figure 5D) with a predicting

power (Adjusted R^2) of 0.982. Together, our simulations and experiments demonstrated

that our rules on how substrate traits govern the assembly of MDOL community were

applicable even when this community grew in a spatially structured environment.

The effects of substance diffusivity on the assembly of MDOL community

Our results suggested that the effects of substrate traits on the assembly of MDOL

community in spatially structured or unstructured environments can both be estimated

by Eqn. [2]. However, the estimating values of the involved parameters (Fd_{max}, ks,

Ts_{max}, and kt) in the predictive functions from ODE and IB model are slightly different

(Figure 4 and 5), even when applying identical parameters, as well as equations to

model reaction kinetics, microbial growth and metabolic interactions (Supplementary

Information S2). The difference between the two predictive functions is likely to be

caused by the differences in the assumptions of the two model forms. One main

difference between the two models is in the mass diffusion: the IB model assumes a

spatially structured environment with a limiting diffusion of S, I and P in the

extracellular space, while the ODE model are constructed based on the assumptions

that all the substances are rapidly and thoroughly mixed.

To eliminate the effects of other model settings and directly test whether the diffusivity

of S, I and P affects the assembly of the MDOL community, we modified our IB model

and performed additional simulations. We simultaneously modulated the diffusion level

of S, I, and P (that is, simultaneously changing D_s, D_i, and D_p), as well as constructed

a 'well-mixed' IB model, in which S, I and P were assumed to be evenly mixed during

the simulations (Supplementary Information S2.2). As shown in Fig. 6A, the predictive

functions derived from the scenarios with higher level of diffusivity, as well as the

'well-mixed' IB model, are more similar to the function derived from 'well-mixed'

ODE model. This result indicated that limited mass diffusion plays a significant role in

governing the assembly of the MDOL community.

How the diffusion level affects the assembly of MDOL community? Firstly, limited

diffusion of P promoted the product privatization by the Embezzler cells, since product

was distributed more around its producers when its diffusion is limited (Supplementary

Figure 22). From this perspective, limited diffusion of P favors the Embezzler. As

expected, when substrate toxicity is not considered ($\theta = 0$), the proportion of Detoxifier

is generally lower when the community grows in spatially structured environments

(Figure 6A; first row). We further tested this hypothesis by modulating the diffusion

level of P (solely changing D_p) in our IB model that did not include substrate toxicity.

As shown in Supplementary Figure 23, we found that higher level of P diffusion indeed

favors the Detoxifier. In conclusion, diffusivity of final product governs the assembly

of MDOL community by affecting the level of product privatization.

Secondly, Embezzler cells require intermediate metabolite I to produce final product P

for growth, but we found that I was largely distributed around the Detoxifier cells

(Supplementary Figure 24). Thus, when I diffused at a lower speed, more Detoxifier

cells are required to fulfill the I supply (that is, to ensure sufficient high level of I

concentration around Embezzler cells). Therefore, we hypothesized that limited diffusion of I favored the growth of Detoxifier. As shown in Supplementary Figure 25, with an increase of D_i, I was distributed more around the Embezzler cells, and the relative proportion of Detoxifier cells decreases. Therefore, intermediate diffusivity affects the assembly of MDOL community by influencing the efficiency of intermediate exchange.

Finally, to investigate how substrate diffusivity affects the assembly of MDOL community, we performed IB simulations initialized at different diffusion levels of the substrate (solely changing D_s). As shown Figure 6A (the second row, $\theta = 0$), when substrate toxicity is omitted, the proportion of Detoxifier cells increased with an increase of diffusion levels of the substrate. This finding is likely to be due to the fact that higher diffusion levels of S thicken the cell's 'active layer', thus increasing production and secretion of the final product by Embezzler cells (Figure 6B; [45]). However, limited diffusion of S will also help to maintain a lower local extracellular S concentration around the Detoxifier cells (Supplementary Figure 26). When substrate toxicity was taken into account, this local gradient caused the substrate toxicity to be less harmful to Detoxifier cells compared to Embezzler cells. Thus, in addition to thickening cell 'active layer' that benefits Detoxifier cells, higher diffusion levels of S oppose the development of local substrate gradient around the Detoxifier cells. Subsequently, this process also decreases the fitness of Detoxifier cells.

Based on our findings for the combination of these two effects, we hypothezised that with the increase of substrate toxicity and its concentration, Detoxifier should gradually

turn to be favored in an intermediate diffusion level of substrate. This hypothesis was

successfully verified by our IB simulations (Figure 6A; second row. For instance, when

$\theta = 0.3$ and $S_0 > 10$ C-mM, Detoxifier was favored at intermediate levels of D_s, 2

$\mu m^2/s$). We also formulated a term to estimate how the ability of the substrate to diffuse

affects the assembly of MDOL community, and thus obtained a new formula

$$DF = \frac{Fd_{max} \cdot s_0}{ks + s_0} \cdot (1 + \frac{Ts_{max} \cdot \theta s_0}{kt + \theta s_0}) \cdot (\frac{s_0 D_s}{kd_1 + s_0 D_s} - \frac{\theta D_s}{kd_2 + \theta D_s}) \qquad [3]$$

In this formula, $\frac{s_0 D_s}{kd_1 + s_0 D_s}$ represents an estimate of the positive effect of increasing

substrate diffusion level via thickening cell 'active layer', which is related to the initial

substrate concentration (s_0; Figure 6B; [45]); $\frac{\theta D_s}{kd_2 + \theta D_s}$ represents an estimate of the

negative effect of increasing substrate diffusion level, which is influenced by toxic

strength of the substrate (Figure 6A; the second row). Our fitting analysis suggested

that Eqn. [3] can well estimate the assembly of MDOL community in our IB simulations

(Figure 6C; R^2=0.994). Therefore, substrate diffusivity is another trait of substrate that

governs the assembly of MDOL community in spatially structured environment.

Discussion

Here we show how substrate traits drive the assembly of the microbial communities engaged in metabolic division of labor (MDOL) when degrading organic compounds. During this process, the population performing the first step is favored by both higher substrate concentration and its toxicity. We found that this rule is applicable both when the community grow in either a well-mixed or a spatially structured environment. Previous studies based on meta-omics approached demonstrated that MDOL was utilized in many microbial communities specialized in degrading organic compounds[6, 46]. Inspired by these natural MDOL systems, numerous studies have recently explored strategies to divide metabolic roles across different populations in a consortium toward removal of organic pollutants[13, 47-51]. While most of these studies focused on the efficiency in pollutant degradation of these communities, a small number of studies also measured their community structures. For instance, one recent study reported that a bacterial consortium composed of *Leucobacter* sp. GP and *Achromobacter denitrificans* PR1 efficiently degrades an antibiotic, sulfamethoxazole, in which the strain GP is responsible for the initial metabolism of the sulfamethoxazole (Detoxifier), and the strain PR1 carries out the subsequent conversion (Embezzler)[17]. This study measured the structures of the community across a gradient of initial substrate concentrations, and found that the proportion of the GP is positively correlated with the initial sulfamethoxazole concentration. This observation largely agrees with the idea derived from our model and experiments, namely that increasing concentration of the substrate favors the Detoxifier population. Therefore, the proposed rule in our study

may be expanded to forecast the assembly of other communities engaged in MDOL.

Our study also compared the assembly of MDOL community when it grew in a well-

mixed system and spatially structured environment. Our data indicated that limited

mass diffusion in spatially structured environments is one key factor that cause the

difference. This finding is reminiscent of other studies proposing that limited mass

diffusion plays significant role on governing the assembly of microbial communities

containing other diffusion-based interaction modes (for example, community with

syntrophic exchange [30, 37, 52], community with cross-protection [53], and

community with 'rock-paper-scissors' interaction [54, 55]. Here, in our study, we

proved that this is also the case in the community engaged in MDOL. In addition, we

reconfirmed that limited mass diffusion is one possible way to privatize public benefit

[30, 37, 56]. On the one hand, it helps the Embezzler population to privatize the final

product for its own growth. On the other hand, it helps the Detoxifier population to

privatize its benefit from toxicant degradation (i.e., detoxification). Therefore, limited

mass diffusion in spatially structured environments may be a universally used avenue

for microorganisms to maintain their private benefit.

Nevertheless, our study also found that predictive function derived from the individual-

based (IB) model with the assumption of mass well-mixing was not completely

identical to the function from our ordinary differential equation (ODE) model,

suggesting that other physical features of spatially structured environment may also

contribute to the assembly of MDOL community. For example, the spatial distributions

of different cell types in the colony are thought to represent another vital factor that

governs community assembly [27, 56-58]. In agreement with previous studies [31, 38,

59], our IB simulations suggested that when two populations engaged in MDOL, cells

from the two populations are spatially more proximal to each other than the scenario

when the two populations did not exhibit defined interactions (Supplementary Figure

27). In addition, we also found that this trait of the colony pattern (spatial proximity of

the two populations) was governed by substrate traits (Supplementary Figure 27).

Particularly, when the strength of substrate toxicity was higher, the Detoxifier cells

occupied the periphery of the growing colony, forming a clearly 'ring' around the

colony (Figure 5; Supplementary video 3; Supplementary Figure 28). The formation of

this ring might be due to the fact that the substrate was present at much higher

concentrations at the colony edge, and hence more toxic, thus largely favoring

Detoxifier cells. These results suggest that substrate traits also govern the spatial

distributions of different cells in the colony developed by MDOL community, which

may in turn, affect the assembly of such community. This featured cell distribution was

not observed in the experiments in this study, presumably because the toxicity of

salicylate was insufficiently high. However, one recent study found that during the

spatial organization of a MDOL community that collectively degrades toluene, the *P.*

putida cells responsible for the first metabolic step formed a monolayer structure in the

periphery of the growing colony [31]. Therefore, such cell distribution may represent a

critical feature of the spatial patterns developed by a MDOL community that degrades

substrates characterized by higher toxic strengths.

While our study provides critical new insights into how the community engaged in

MDOL assembles, a number of limitations need to be taken into consideration. First, our model analysis showed that substrate toxicity is vital to govern the assembly of communities engaged in MDOL. However, due to the difficulties in manipulating the toxicity of the substrate (salicylate) *in vitro*, we were unable to experimentally compare the impact of the different toxic strengths on community assembly. Nevertheless, our model correctly predicts that simply increasing the initial substrate concentration is unlikely to shape a community dominated by the Detoxifier population, while the presence of substrate toxicity renders the 'Detoxifier' population in the community to become dominant. Therefore, the observation that Detoxifier population was able to dominate the synthetic community when high concentration of substrate was provided, and the measured biotoxicity of salicylate strongly suggested that substrate toxicity should be included in the assembly of the synthetic microbial community. In agreement with this idea, our predictive functions involved in salicylate toxic strength fits the experiment results very well. To further examine this idea, it is necessary to design a better system in which the toxicity of the substrate can be modulated. For example, several studies applied defined co-culturing of multiple bacterial species for degrading antibiotics [17, 60, 61]. In these systems, antibiotics are toxic to all strains involved, but their toxicity may be modulated by genetically modifying the expression of the genes encoding the corresponding antibiotic resistance. Experiments using these well-defined systems should assist in more thoroughly investigating the effects of substrate toxicity on the assembly of community.

Second, our ODE model suggests that apart from substrate traits, five other key

parameters exist that exhibit considerable effects on the assembly of a MDOL

community. Here, we primarily focused on the effects of substrate traits, without

analyzing in detail how all the seven key factors collectively determine the process of

community assembly. Nonetheless, our analysis presented here suggests that biotic

factors such as speed of the first reaction (a_1), mass transport rate (γ_s, γ_i, γ_p), as well

as consumtion rate of P (Cp), affected the assembly of the community, namely by

determining the value of parameters in Eqn. [2] (i.e., Fd_{max}, ks, Ts_{max}, and kt).

However, due to the difficulties in analytically solving non-linear ODEs, as well as the

low efficiency of individual-based simulations [62], detailed quantitative understanding

of how all these factors affect the assembly of MDOL community remains limited. We

thus suggest further studies to use more simplified models that combine these elements

to provide a more general description of the principles governing the assembly of a

MDOL community.

To engineer stable and high-efficient microbial systems for bioproduction or

biodegradation, it will be critical to predict how the communities assembled by a given

set of strains exhibiting modularized functions. Our results demonstrate that, for a given

community engaged in MDOL, its assembly can be quantitatively estimated from the

abiotic factors, such as the traits of its substrate, suggesting that it is feasible to manage

microbial communities through manipulation of specific environmental factors, to

address grand challenges facing human society in agriculture, degradation of the

environment, and human health.

Methods

Formulation and analyses of the ODE model

Formulation of the ODE model

To investigate the assembly of MDOL community in well-mixed system, a

mathematical model was formulated using ordinary differential equations (ODEs).

Here, the dimensionless forms of the models were presented. The detailed derivations

of the models, and choices of parameter values are described in Supplementary

Information S1.

As described in the Results section, a two-step pathway was assumed to be

implemented by MDOL between two populations (Figure 2A and Figure 3A). For

simplicity, the basic model was built based on five simple assumptions: (1) The systems

are well mixed in each compartment (inside a cell or in the extracellular space). (2)

transport of substrate (S), intermediate (I) and final product (P) is mediated by passive

diffusion; (3) P was assumed to be the sole and limited resource for the growth of the

two populations and its consumption was calculated following Monod equations; (4)

Basic biological properties (the coefficients in Monod equations) regarding the growth

of the two populations are identical, since we only focused on the effects of abiotic

factors; (5) when applicable, substrate toxicity was introduced by adding three different

toxic terms to the growth equation (Supplementary Table 3), dependent on intracellular

S concentration of the corresponding population. The dynamics of intracellular and

extracellular I and P are given by

651
$$\frac{ds_{1,in}}{d\tau} = -\frac{a_1}{1+s_{1,in}} s_{1,in} + \gamma_s \cdot \left(s_{out} - s_{1,in}\right) \qquad [4]$$

$$652 \qquad \frac{ds_{2,in}}{d\tau} = \gamma_s \cdot \left(s_{out} - s_{2,in}\right) \qquad [5]$$

$$653 \qquad \frac{di_{1,in}}{d\tau} = \frac{a_1}{1 + s_{1,in}} s_{1,in} - \gamma_i \cdot \left(i_{1,in} - i_{out}\right) \qquad [6]$$

$$654 \qquad \frac{di_{2,in}}{d\tau} = -\frac{a_2}{\beta_2 + i_{2,in}} i_{2,in} + \gamma_i \cdot \left(i_{out} - i_{2,in}\right) \qquad [7]$$

$$655 \qquad \frac{dp_{1,in}}{d\tau} = -\frac{Cp}{\beta_g + p_{1,in}} p_{1,in} + \gamma_p \cdot \left(p_{out} - p_{1,in}\right) \qquad [8]$$

$$656 \qquad \frac{dp_{2,in}}{d\tau} = \frac{a_2}{\beta_2 + i_{2,in}} i_{2,in} - \frac{Cp}{\beta_g + p_{2,in}} p_{2,in} + \gamma_p \cdot \left(p_{out} - p_{2,in}\right) \qquad [9]$$

$$657 \qquad \frac{ds_{out}}{d\tau} = -x_1 \cdot \gamma_s \cdot \left(s_{out} - s_{1,in}\right) - x_2 \cdot \gamma_s \cdot \left(s_{out} - s_{2,in}\right) \qquad [10]$$

$$658 \qquad \frac{di_{out}}{d\tau} = x_1 \cdot \gamma_i \cdot \left(i_{out} - i_{1,in}\right) - x_2 \cdot \gamma_i \cdot \left(i_{out} - i_{2,in}\right) \qquad [11]$$

$$659 \qquad \frac{dp_{out}}{d\tau} = x_2 \cdot \gamma_p \cdot \left(p_{out} - p_{1,in}\right) - x_1 \cdot \gamma_p \cdot \left(p_{out} - p_{1,in}\right) \qquad [12]$$

660   The growth of the two populations was modeled using a general logistic function with

661   first-order cell death:

$$662 \qquad \frac{dx_1}{d\tau} = \frac{Cp}{bg + p_{1,in}} p_{1,in} yt_1 x_1 \left(1 - \frac{x_1 + x_2}{\rho}\right) \qquad [13]$$

$$663 \qquad \frac{dx_2}{d\tau} = \frac{Cp}{bg + p_{2,in}} p_{2,in} yt_2 x_2 \left(1 - \frac{x_1 + x_2}{\rho}\right) \qquad [14]$$

The definitions and dimensionless methods of all variables are listed in Supplementary

Table 1. The definitions and dimensionless methods, as well as the value ranges of all

the parameters involved in these equations are listed in Supplementary Table 2.

Simulation and analyzing protocol of the ODE model

Details of the simulation and analysis protocols of our ODE model and the downstream

analyses are described in Supplementary Information S1.3. Briefly, to solve the

community dynamics of the MDOL community with given parameter sets, numerical

simulations of our ODE model were performed using *NDsolve* function of *Wolfram*

Mathematica. The numerical solutions of all the variables, including the dynamics of

mass (S, I, P) concentration and biomass, were recorded for further analyses. To

perform simulations with numerous parameter sets, as well as the downstream analysis,

custom *Mathematica* scripts were wrote mainly based on the *Do* loop function.

Individual-based modeling

Our individual based (IB) model was constructed based on *gro* platform (https:

https://github.com/liaupm/GRO-LIA), a simulator designed by Gutiérrez and

colleagues aiming to describe multicellular bacterial behavior [63]. The model aims to

simulate the growth of a microbial colony composed of two populations who execute

substrate degradation via MDOL on a surface. The model was formulated mainly using

the same equations as our dimensional ODE model (Supplementary Information S1.1,

Eqns. [S1]-[S13]) to characterize the intra- and extracellular dynamics of mass (S, I, P)

concentration, as well as to calculate the rate of cell growth. Four main differences exist

between our IB model and the ODE model: (1) The IB model was formulated on a

spatially structured surface, and the diffusion of S, I, and P was limited; (2) Mass

dynamics was modelled at single-cell level; (3) The growth of both populations was

modelled at single-cell level, and passive cell shoving during the cell growth was

included; (4) cells were inoculated in the center of the surface, and the entire community

underwent 'colony range expansion', a process whereby the community immigrate

outwards as a whole, driven by the force generated from cell growth and division

(Supplementary Figure 15). The mathematical framework formulating these four points

is described in Supplementary Information S2.1. To implement our design of the IB

model, custom codes were written in *gro* language. Variables and Parameters in the IB

model are summarized in Supplementary Table 7. Details of the IB simulation

workflow are described in Supplementary Information S2.

Experimental verification of our model prediction

Genetic manipulation of the P. stutzeri strains

All *P. stutzeri* strains were engineered from a naphthalene-degrading bacterial strain *P.*

stutzeri AN10 [64]. Genes that encode the key enzymes responsible for corresponding

metabolic steps in salicylate degradation pathway were knocked out to generate the *P.*

stutzeri strains. The details of the genetic manipulation of are described in

Supplementary information S3.

Liquid cultivation of our synthetic microbial communities

Liquid cultivation of our synthetic microbial communities was performed in 96-well

plates that contains 120 μL fresh minimum medium. Proportions of the two populations

in the community were estimated by measuring the fluorescent intensity of the two

strains involved using a microplate reader (Molecular Devices, Sunnyvale, America).

Detailed protocols are described in Supplementary information S4.

Colony pattern formation assays

Colony pattern formation assays were performed on the agarose surface in a Petri dish

(60 mm in diameter). Images of the colony patterns were taken under a 5× objective

using a Leica DM6000B fluorescence microscope (Leica Corporation, Wetzlar,

Germany) equipped with a LED fluorescence illuminator (Leica Corporation). The

relative fraction of each population in the colonies was measured by image analysis, as

well as similar fluorescence-measurement method as performed in liquid cultivation

experiments. Detailed protocols are described in Supplementary information S5.

Statistical analysis

Unless indicated otherwise, the number of replicates was three for each simulation, and six for each experiment. For comparative statistics, unpaired, two-tailed, Student's t-test was performed in Wolfram Mathematica (version 12.4). To fit the data to the proposed function, Nonlinearmodelfit function of the Wolfram Mathematica (version 12.4) was applied.

Code availability

All custom *Mathematica* codes used for ODE simulation and data analyses, as well as the source *gro* codes used for our IB simulations are available at Github: https://github.com/RoyWang1991/MDOLcode/tree/master/MDOL-spatial.

Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgments

We wish to thank Professor Ping Xu (Shanghai Jiao Tong University, Shanghai, P.R. China) for supplying plasmid pMMPc-Gm, used for fluorescence labeling in this study; Dr. Min Lin (Chinese Academy of Agricultural Sciences, Beijing, P.R. China) for providing plasmid pK18mobsacB and pRK2013, used for genetic engineering in this work; Professor Martin Ackermann (ETH Zurich, Zurich, Switzerland), Dr. David Johnson (Eawag, Dübendorf, Switzerland) and Yinyin Ma (Eawag, Dübendorf, Switzerland) for constructive inputs on the design of this study; Professor Martín Gutiérrez (Universidad Politécnica de Madrid, Madrid, Spain) for his kindly guidance for the set-up of the *gro* platform for the individual-based simulations; Dr. T. Juelich

740   (UCAS, Beijing) for linguistic assistance during the preparation of this manuscript.

741   This work was supported by National Key R&D Program of China (2018YFA0902100

742   and 2018YFA0902103), and National Natural Science Foundation of China (91951204,

743   31761133006, 31770120, and 31770118).

References

1. Kreft JU, Griffin BM, Gonzalez-Cabaleiro R. Evolutionary causes and consequences of metabolic division of labour: why anaerobes do and aerobes don't. Curr Opin Biotechnol. 2020;62:80-7; doi:10.1016/j.copbio.2019.08.008.

2. Tsoi R, Wu F, Zhang C, Bewick S, Karig D, You L. Metabolic division of labor in microbial systems. Proc Natl Acad Sci U S A. 2018;115:10:2526-31; doi:10.1073/pnas.1716888115.

3. Harvey E, Heys J, Gedeon T. Quantifying the effects of the division of labor in metabolic pathways. J Theor Biol. 2014;360:222-42; doi:10.1016/j.jtbi.2014.07.011.

4. Thommes M, Wang T, Zhao Q, Paschalidis IC, Segre D. Designing Metabolic Division of Labor in Microbial Communities. mSystems. 2019;4:2; doi:10.1128/mSystems.00263-18.

5. Seelke R. The Prokaryotes: a handbook on the biology of bacteria. Choice: Current Reviews for Academic Libraries. 2007;45:1:68-.

6. Dombrowski N, Donaho JA, Gutierrez T, Seitz KW, Teske AP, Baker BJ. Reconstructing metabolic pathways of hydrocarbon-degrading bacteria from the Deepwater Horizon oil spill. Nat Microbiol. 2016;1:7:16057; doi:10.1038/nmicrobiol.2016.57.

7. West SA, Cooper GA. Division of labour in microorganisms: an evolutionary perspective. Nat Rev Microbiol. 2016;14:11:716-23; doi:10.1038/nrmicro.2016.111.

8. van Gestel J, Vlamakis H, Kolter R. Division of Labor in Biofilms: the Ecology of Cell Differentiation. Microbiol Spectr. 2015;3:2:MB-0002-2014;

doi:10.1128/microbiolspec.MB-0002-2014.

9. Lilja EE, Johnson DR. Segregating metabolic processes into different microbial cells

accelerates the consumption of inhibitory substrates. Isme Journal. 2016;10:7:1568-78;

doi:10.1038/ismej.2015.243.

10. Schink B, Pfennig N. Fermentation of Trihydroxybenzenes by Pelobacter-

Acidigallici Gen-Nov Sp-Nov a New Strictly Anaerobic, Non-Sporeforming Bacterium.

Arch Microbiol. 1982;133:3:195-201; doi:Doi 10.1007/Bf00415000.

11. Wang CY, Huang Y, Zhang ZT, Hao H, Wang H. Absence of the nahG-like gene

caused the syntrophic interaction between *Marinobacter* and other microbes in PAH-

degrading process. J Hazard Mater. 2020;384; doi:ARTN 121387

10.1016/j.jhazmat.2019.121387.

12. Katsuyama C, Nakaoka S, Takeuchi Y, Tago K, Hayatsu M, Kato K.

Complementary cooperation between two syntrophic bacteria in pesticide degradation.

Journal of Theoretical Biology. 2009;256:4:644-54; doi:10.1016/j.jtbi.2008.10.024.

13. Yang C, Li Y, Zhang K, Wang X, Ma C, Tang H, Xu P. Atrazine degradation by a

simple consortium of *Klebsiella* sp. A1 and *Comamonas* sp. A2 in nitrogen enriched

medium. Biodegradation. 2010;21:1:97-105; doi:10.1007/s10532-009-9284-9.

14. Billet L, Devers M, Rouard N, Martin-Laurent F, Spor A. Labour sharing promotes

coexistence in atrazine degrading bacterial communities. Sci Rep. 2019;9:1:18363;

doi:10.1038/s41598-019-54978-2.

15. Gilbert ES, Walker AW, Keasling JD. A constructed microbial consortium for

biodegradation of the organophosphorus insecticide parathion. Appl Microbiol Biot.

2003;61:1:77-81; doi:10.1007/s00253-002-1203-5.

16. Meyer-Cifuentes IE, Werner J, Jehmlich N, Will SE, Neumann-Schaal M, Ozturk B. Synergistic biodegradation of aromatic-aliphatic copolyester plastic by a marine microbial consortium. Nature Communications. 2020;11:1; doi:ARTN 579010.1038/s41467-020-19583-2.

17. Reis AC, Cvancarova M, Liu Y, Lenz M, Hettich T, Kolvenbach BA, Corvini PFX, Nunes OC. Biodegradation of sulfamethoxazole by a bacterial consortium of *Achromobacter denitrificans* PR1 and *Leucobacter* sp GP. Appl Microbiol Biot. 2018;102:23:10299-314; doi:10.1007/s00253-018-9411-9.

18. Peng XF, Wilken S, Lankiewicz TS, Gilmore SP, Brown JL, Henske JK, Swift CL, Salamov A, Barry K, Grigoriev IV, Theodorou MK, Valentine DL, O'Malley MA. Genomic and functional analyses of fungal and bacterial consortia that enable lignocellulose breakdown in goat gut microbiomes. Nat Microbiol. 2021;6:4:499-+; doi:10.1038/s41564-020-00861-0.

19. Kato S, Haruta S, Cui ZJ, Ishii M, Igarashi Y. Stable coexistence of five bacterial strains as a cellulose-degrading community. Appl Environ Microb. 2005;71:11:7099-106; doi:10.1128/Aem.71.11.7099-7106.2005.

20. Koudryashova Y, Chizhova T, Tishchenko P, Hayakawa K. Seasonal Variability of Polycyclic Aromatic Hydrocarbons (PAHs) in a Coastal Marine Area in the Northwestern Region of the Sea of Japan/East Sea (Possiet Bay). Ocean Sci J. 2019;54:4:635-55; doi:10.1007/s12601-019-0031-9.

21. Jiang Y, Hu X, Yves UJ, Zhan H, Wu Y. Status, source and health risk assessment

of polycyclic aromatic hydrocarbons in street dust of an industrial city, NW China. Ecotoxicol Environ Saf. 2014;106:11-8; doi:10.1016/j.ecoenv.2014.04.031.

22. Ozaki N, Takamura Y, Kojima K, Kindaichi T. Loading and removal of PAHs in a wastewater treatment plant in a separated sewer system. Water Res. 2015;80:337-45; doi:10.1016/j.watres.2015.05.002.

23. Peng RH, Xiong AS, Xue Y, Fu XY, Gao F, Zhao W, Tian YS, Yao QH. Microbial biodegradation of polyaromatic hydrocarbons. Fems Microbiol Rev. 2008;32:6:927-55; doi:10.1111/j.1574-6976.2008.00127.x.

24. Park W, Jeon CO, Cadillo H, DeRito C, Madsen EL. Survival of naphthalene-degrading Pseudomonas putida NCIB 9816-4 in naphthalene-amended soils: toxicity of naphthalene and its metabolites. Appl Microbiol Biot. 2004;64:3:429-35; doi:10.1007/s00253-003-1420-6.

25. Hall-Stoodley L, Costerton JW, Stoodley P. Bacterial biofilms: from the natural environment to infectious diseases. Nat Rev Microbiol. 2004;2:2:95-108; doi:10.1038/nrmicro821.

26. Nadell CD, Drescher K, Foster KR. Spatial structure, cooperation and competition in biofilms. Nat Rev Microbiol. 2016;14:9:589-600; doi:10.1038/nrmicro.2016.84.

27. Dal Co A, van Vliet S, Kiviet DJ, Schlegel S, Ackermann M. Short-range interactions govern the dynamics and functions of microbial communities. Nat Ecol Evol. 2020;4:3:366-75; doi:10.1038/s41559-019-1080-2.

28. Nadell CD, Foster KR, Xavier JB. Emergence of Spatial Structure in Cell Groups and the Evolution of Cooperation. Plos Computational Biology. 2010;6:3; doi:ARTN

e100071610.1371/journal.pcbi.1000716.

29. Frost I, Smith WPJ, Mitri S, San Millan A, Davit Y, Osborne JM, Pitt-Francis JM, MacLean RC, Foster KR. Cooperation, competition and antibiotic resistance in bacterial colonies. Isme Journal. 2018;12:6:1582-93; doi:10.1038/s41396-018-0090-4.

30. Momeni B, Waite AJ, Shou W. Spatial self-organization favors heterotypic cooperation over cheating. Elife. 2013;2:e00960; doi:10.7554/eLife.00960.

31. Tecon R, Or D. Cooperation in carbon source degradation shapes spatial self-organization of microbial consortia on hydrated surfaces. Sci Rep-Uk. 2017;7; doi:ARTN 4372610.1038/srep43726.

32. Quintas C, Leyva JS, Sotoca R, Loureiro-Dias MC, Peinado JM. A model of the specific growth rate inhibition by weak acids in yeasts based on energy requirements. Int J Food Microbiol. 2005;100:1-3:125-30; doi:10.1016/j.ijfoodmicro.2004.10.009.

33. Lanfranconi MP, Christie-Oleza JA, Martin-Cardona C, Suarez-Suarez LY, Lalucat J, Nogales B, Bosch R. Physiological role of NahW, the additional salicylate hydroxylase found in *Pseudomonas stutzeri* AN10. Fems Microbiol Lett. 2009;300:2:265-72; doi:10.1111/j.1574-6968.2009.01787.x.

34. Nadell CD, Drescher K, Foster KR. Spatial structure, cooperation and competition in biofilms. Nat Rev Microbiol. 2016;14:9:589-600; doi:10.1038/nrmicro.2016.84.

35. Mitri S, Xavier JB, Foster KR. Social evolution in multispecies biofilms. P Natl Acad Sci USA. 2011;108:10839-46; doi:10.1073/pnas.1100292108.

36. Xavier JB, Foster KR. Cooperation and conflict in microbial biofilms. P Natl Acad Sci USA. 2007;104:3:876-81; doi:10.1073/pnas.0607651104.

37. Pande S, Kaftan F, Lang S, Svatos A, Germerodt S, Kost C. Privatization of

cooperative benefits stabilizes mutualistic cross-feeding interactions in spatially

structured environments. ISME J. 2016;10:6:1413-23; doi:10.1038/ismej.2015.212.

38. Goldschmidt F, Regoes RR, Johnson DR. Successive range expansion promotes

diversity and accelerates evolution in spatially structured microbial populations. Isme

Journal. 2017;11:9:2112-23; doi:10.1038/ismej.2017.76.

39. Klausen M, Heydorn A, Ragas P, Lambertsen L, Aaes-Jorgensen A, Molin S, Tolker-

Nielsen T. Biofilm formation by *Pseudomonas aeruginosa* wild type, flagella and type

IV pili mutants. Mol Microbiol. 2003;48:6:1511-24; doi:10.1046/j.1365-

2958.2003.03525.x.

40. Barken KB, Pamp SJ, Yang L, Gjermansen M, Bertrand JJ, Klausen M, Givskov M,

Whitchurch CB, Engel JN, Tolker-Nielsen T. Roles of type IV pili, flagellum-mediated

motility and extracellular DNA in the formation of mature multicellular structures in

Pseudomonas aeruginosa biofilms. Environ Microbiol. 2008;10:9:2331-43;

doi:10.1111/j.1462-2920.2008.01658.x.

41. O'Toole GA, Kolter R. Flagellar and twitching motility are necessary for

Pseudomonas aeruginosa biofilm development. Mol Microbiol. 1998;30:2:295-304;

doi:10.1046/j.1365-2958.1998.01062.x.

42. Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding

frontiers promotes gene segregation. P Natl Acad Sci USA. 2007;104:50:19926-30;

doi:10.1073/pnas.0710150104.

43. Hallatschek O, Nelson DR. Life at the Front of an Expanding Population. Evolution.

2010;64:1:193-206; doi:10.1111/j.1558-5646.2009.00809.x.

44. Korolev KS, Xavier JB, Nelson DR, Foster KR. A Quantitative Test of Population

Genetics Using Spatiogenetic Patterns in Bacterial Colonies. American Naturalist.

2011;178:4:538-52; doi:10.1086/661897.

45. Mitri S, Clarke E, Foster KR. Resource limitation drives spatial organization in

microbial groups. Isme Journal. 2016;10:6:1471-82; doi:10.1038/ismej.2015.208.

46. Yu K, Yi S, Li B, Guo F, Peng X, Wang Z, Wu Y, Alvarez-Cohen L, Zhang T. An

integrated meta-omics approach reveals substrates involved in synergistic interactions

in a bisphenol A (BPA)-degrading microbial community. Microbiome. 2019;7:1:16;

doi:10.1186/s40168-019-0634-5.

47. Li Z, Yoshida N, Wang A, Nan J, Liang B, Zhang C, Zhang D, Suzuki D, Zhou X,

Xiao Z, Katayama A. Anaerobic mineralization of 2,4,6-tribromophenol to CO2 by a

synthetic microbial community comprising *Clostridium*, *Dehalobacter*, and

Desulfatiglans.   Bioresour   Technol.   2015;176:225-32;

doi:10.1016/j.biortech.2014.10.097.

48. Che S, Men Y. Synthetic microbial consortia for biosynthesis and biodegradation:

promises and challenges. J Ind Microbiol Biotechnol. 2019;46:9-10:1343-58;

doi:10.1007/s10295-019-02211-4.

49. Hudcova T, Halecky M, Kozliak E, Stiborova M, Paca J. Aerobic degradation of

2,4-dinitrotoluene by individual bacterial strains and defined mixed population in

submerged   cultures.   J   Hazard   Mater.   2011;192:2:605-13;

doi:10.1016/j.jhazmat.2011.05.061.

50. Roell GW, Zha J, Carr RR, Koffas MA, Fong SS, Tang YJ. Engineering microbial

consortia by division of labor. Microb Cell Fact. 2019;18:1:35; doi:10.1186/s12934-

019-1083-3.

51. Jia XQ, He Y, Jiang DW, Liu C, Lu WY. Construction and analysis of an engineered

Escherichia coli-Pseudomonas aeruginosa co-culture consortium for phenanthrene

bioremoval. Biochem Eng J. 2019;148:214-23; doi:10.1016/j.bej.2019.05.010.

52. Momeni B, Brileya KA, Fields MW, Shou WY. Strong inter-population cooperation

leads to partner intermixing in microbial communities. Elife. 2013;2; doi:ARTN

e0023010.7554/eLife.00230.

53. Frost I, Smith WPJ, Mitri S, Millan AS, Davit Y, Osborne JM, Pitt-Francis JM,

MacLean RC, Foster KR. Cooperation, competition and antibiotic resistance in

bacterial colonies. ISME J. 2018;12:6:1582-93; doi:10.1038/s41396-018-0090-4.

54. Kerr B, Riley MA, Feldman MW, Bohannan BJ. Local dispersal promotes

biodiversity in a real-life game of rock-paper-scissors. Nature. 2002;418:6894:171-4;

doi:10.1038/nature00823.

55. Nahum JR, Harding BN, Kerr B. Evolution of restraint in a structured rock-paper-

scissors   community.   P   Natl   Acad   Sci   USA.   2011;108:10831-8;

doi:10.1073/pnas.1100296108.

56. Wang M, Liu X, Nie Y, Wu XL. Selfishness driving reductive evolution shapes

interdependent patterns in spatially structured microbial communities. ISME J.

2021;15:5:1387-401; doi:10.1038/s41396-020-00858-x.

57. Kim W, Racimo F, Schluter J, Levy SB, Foster KR. Importance of positioning for

microbial evolution. Proc Natl Acad Sci U S A. 2014;111:16:E1639-47; doi:10.1073/pnas.1323632111.

58. Liu X, Wang M, Nie Y, Wu X-L. Successful microbial colonization of space using an anti-aggregation strategy. bioRxiv. 2021:2021.07.09.451734; doi:10.1101/2021.07.09.451734.

59. Goldschmidt F, Regoes RR, Johnson DR. Metabolite toxicity slows local diversity loss during expansion of a microbial cross-feeding community. Isme Journal. 2018;12:1:136-44; doi:10.1038/ismej.2017.147.

60. Liu CX, Xu QM, Yu SC, Cheng JS, Yuan YJ. Bio-removal of tetracycline antibiotics under the consortium with probiotics *Bacillus clausii* T and *Bacillus amyloliquefaciens* producing biosurfactants. Sci Total Environ. 2020;710; doi:ARTN 13632910.1016/j.scitotenv.2019.136329.

61. Kong XX, Jiang JL, Qiao B, Liu H, Cheng JS, Yuan YJ. The biodegradation of cefuroxime, cefotaxime and cefpirome by the synthetic consortium with probiotic Bacillus clausii and investigation of their potential biodegradation pathways. Sci Total Environ. 2019;651:271-80; doi:10.1016/j.scitotenv.2018.09.187.

62. Hellweger FL, Clegg RJ, Clark JR, Plugge CM, Kreft JU. Advancing microbial sciences by individual-based modelling. Nat Rev Microbiol. 2016;14:7:461-71; doi:10.1038/nrmicro.2016.62.

63. Gutierrez M, Gregorio-Godoy P, Perez Del Pulgar G, Munoz LE, Saez S, Rodriguez-Paton A. A New Improved and Extended Version of the Multicell Bacterial Simulator gro. ACS Synth Biol. 2017;6:8:1496-508; doi:10.1021/acssynbio.7b00003.

64. Brunet-Galmes I, Busquets A, Pena A, Gomila M, Nogales B, Garcia-Valdes E,

Lalucat J, Bennasar A, Bosch R. Complete Genome Sequence of the Naphthalene-

Degrading Bacterium *Pseudomonas stutzeri* AN10 (CCUG 29243). J Bacteriol.

2012;194:23:6642-3; doi:10.1128/Jb.01753-12.

Figures

Figure 1 Hypothesis for how substrate concentration and toxicity govern the assembly of community engaged in MDOL. In a community

degrading an organic compound through metabolic division of labor (MDOL), final product was assumed to be the sole resource and was

synthesized by the strain performing the second step. Therefore, this strain will obtain more nutrients (denoted as bigger 'smiling face'), while the

other strain has to collect product released from this population (denoted as smaller 'smiling face'). Thus, the last population was named

'Embezzler'. However, increasing the concentration of the substrate (vertical axis) improves the flux of the pathway. Since the P consuming ability

of Embezzler cells is limited (dashed box), increasing the concentration will lead to higher final product leakiness, favoring the growth of the first

population. Moreover, introducing substrate biotoxicity (horizontal axis) also favors the first population, because it converts this toxic substrate

(denoted as smaller sad face), resulting in lower intracellular substrate concentration compared to that of the Embezzler cells (denoted as bigger

sad face). Thus, the first population was named 'Detoxifier'.

Figure 2 Simulation of the ordinary differential equation (ODE) model excluding substrate toxicity. (A) Schematic diagram showing the basic

assumptions of our ODE model without including substrate toxicity. (B-C) A representative case shows how substrate concentration affects the

assembly of our? a? MDOL community. The simulation dynamics of the fraction of Detoxifier population with the conditions of different initial

substrate concentrations are shown in (B). The relationship between substrate concentration and stead-state fraction of Detoxifier is shown in (C).

964   In (C), the green dots denote the simulated stead-state fraction of Detoxifier, and the red dashed line shows the plot of the best fitting function

965   using Eqn. [1]. Parameter values used in these simulations: $y = 10^{-4}$, $Cp = 10$, $bg = 1$, $a_1 = 10000$, $a_2 = 1000$, $\beta_2 = 1$, $\gamma_s = 1$, $\gamma_i = 1$, $\gamma_p = 1$, $\rho = 10^{-2}$.

966   The best fitting value of ks in this case is 35.3, and that of Fd_{max} is 0.417. (D-F) Distributions of Adjusted R^2 (D) of the fitting functions, best fitting

967   value of ks (E) and Fd_{max} (F) in the second-round simulations that does not include substrate toxicity, using 7776 parameter value combinations of

968   the five key parameters (a_1, γ_s, γ_i, γ_p, and Cp).

Figure 3 Simulation of the ordinary differential equation (ODE) model that includes

substrate toxicity, suggesting that both substrate concentration and its toxicity

collectively affect the assembly of a community engaged in MDOL. (A) Schematic

diagram showing the basic assumptions of our ODE model that includes substrate

toxicity. (B) Multiple linear regression analysis of the simulation results of the ODE

model showed how the parameters included in the model affect the assembly of the

MDOL community. Left: results from the first-round simulations that considered all

the twelve parameters are shown. Blue font denotes the identified key parameters. Right:

results from the second-round simulations that only considered the seven key

parameters. The axis of the radar plot denotes the values of fitting coefficients of the

parameters from multiple linear regression analyses. Red dots denote the steady-state

fraction of Detoxifier is positively correlated with corresponding parameter, while the

green dots represent the negative correlation. The origin axis (0) is highlighted by dash

line to emphasize the fact that the closer a value is to zero, the smaller the effect on the

community assembly of the corresponding parameter. The data are also listed in

Supplementary Table 4 and Supplementary Table 5. In this analysis, the toxic effects

of substrate on population growth were assumed to follow a reciprocal relationship.

Results considering other relationships are shown in Supplementary Figure 3. (C) A

representative case shows how both substrate concentration and its toxicity collectively

affect the stead-state proportion of Detoxifier cells. The green dots denote the simulated

stead-state fraction of Detoxifier, and the surface shows the plot of the best fitting

function using Eqn. [2]. Parameter values used in these simulations: $y = 10^{-4}$, $Cp = 10$,

$bg = 1$, $a_1 = 10000$, $a_2 = 1000$, $\beta_2 = 1$, $\gamma_s = 1$, $\gamma_i = 1$, $\gamma_p = 1$, $\rho = 10^{-2}$. The best fitting

value of ks, Fd_{max}, kt, and TS_{max} in this case are 48.9, 0.423, 0.848, 3.39, respectively.

(D) Distributions of Adjusted R^2 of the fitting functions in the second-round simulations

that includes substrate toxicity, using 7776 parameter value combinations of the five

key parameters (a_1, γ_s, γ_i, γ_p, and Cp).

$$DF = \frac{0.481 \cdot s_0}{22.9 + s_0} \cdot \left(1 + \frac{0.886 \cdot \theta s_0}{2.66 + \theta s_0}\right) \xrightarrow{\theta = 0.0032} DF = \frac{0.481 \cdot s_0}{22.9 + s_0} \cdot \left(1 + \frac{0.886 \cdot s_0}{831 + s_0}\right)$$

Figure 4 Community assembly of SMC-mdol in a spatially unstructured system across

different substrate traits. (A) Design of the SMC-mdol. Shown are the pathway of

salicylate degradation in 'Superman' strain *P. stutzeri* AN0011, as well as partial

pathways carried out by Detoxifier strain AN0010 and Embezzler strain AN0001. Skull

marks that salicylate is toxic. (B) Predicting the assembly of the synthetic community

using our ODE model, as well as the derived predictive function using Eqn. [2]. The

relationship between the steady-state fraction of the Detoxifier population and substrate

concentration (s_0), as well as substrate toxic strength (θ), was built from our

mathematical model using parameters consistent with our experiemental system. Each

green dot shows the steady-state fraction of Detoxifier obtained by one simulation

accociated with the specific parameter set. The surface diagram shows distribution of

the steady-state fraction of Detoxifier predicted by our proposed simple formula. The

Red line in the surface denotes the scenarios θ=0.0032, which is the toxic strength of

salicylate obtained from experiemental measurements. (C) The experimental measured

steady-state fractions of Detoxifier in cultures with different salicylate concentrations

is consistent with those from mathematical predictions. Note that in the plots, substrate

concentrations are shown in dimentional form (S_0, Cmmol/L), but in the predictive

functions, the fitting analysis was performed using its dimensionless form (s_0).

1017

$$DF_O = \frac{0.444 \cdot s_0}{37.2+s_0} \cdot (1+\frac{1.10 \cdot \theta s_0}{1.49+\theta s_0}) \xrightarrow[\text{Experiment}]{\theta=0.0032} DF_O = \frac{0.444 \cdot s_0}{37.2+s_0} \cdot (1+\frac{1.10 \cdot s_0}{464+s_0})$$

1018

1019 **Figure 5** Substrate traits governing the assembly of a microbial community engaged in

1020 metabolic division of labor (MDOL) in a spatially structured environment. (A)

1021 Representative colony patterns from Individual-based (IB) modelling initialized with

different substrate traits. Detoxifier cells are shown in red, while Embezzler cells are shown in green. (B) Analysis of community composition in the expanding region of the colonies from IB simulations across eight kinds of initial substrate concentrations and five different toxic strength. Plot shows how both substrate concentration and its toxicity collectively affect the stead-state proportion of Detoxifier. The green dots denote the simulated stead-state fraction of Detoxifier. The surface shows the plot of the best fitting function using Eqn. [2]. The Red line in the surface denotes the scenarios θ=0.0032, which is the toxic strength of salicylate obtained from experimental measurements. (C) Representative colony patterns from the pattern formation assays of SMC-mdol$\triangle$*pilAB*, as well as the IB simulations using the parameters matched with our synthetic system (Supplementary Table 7), across eight different initial substrate concentrations. (D) The experimental measured steady-state fractions of Detoxifier in the expanding region of these colonies is consistent with those from mathematical predictions. Note that in the plots, substrate concentrations are shown in dimensional form (S_0, Cmmol/L), but in the predictive functions, the fitting analyses were performed using its dimensionless form (s_0).

1039

Figure 6 The effects of the diffusion level of substrate, intermediate and product on the

assembly of MDOL community. (A) The relationship between initial substrate

concentration (S_0) with the steady-state proportion of Detoxifier cells in the expanding

region of the colonies, across different substance diffusion level (denoted by different

curve colors) and different strength of substrate toxicity (θ, denoted by five subgraphs).

First row: diffusion levels of S, I and P (that is D_i, D_i, and D_p) were set to be identical

and simultaneously modulated in the simulations. Second row: Diffusion levels of I and

P (D_i and D_p) were set as default values shown in Supplementary Table 7, while

diffusion levels of S were solely modulated. Other parameters in these simulations were

initialized with the default values shown in Supplementary Table 7. The simulation data

were then fitted to Eqn. [2] to obtain the curves shown in the plot. The Adjust R^2 values

for these fitting analyses range from 0.994 to 0.997. (B) Diffusion levels of substrate

affected the thickness of cell 'active layer'. Representative colony images (first row),

the corresponding distributions of final product (second row), as well as the

distributions of cell growth rates (third row) in the 2D plane at steady-state, obtained

from individual-based simulations initialized with different diffusion level of substrate.

Shown are the results in which S_0 was set to 10 C-mol/L and θ was 0 (not include

substrate toxicity). In the colony images, Detoxifier cells are shown in red, while

Embezzler cells are shown in green. Thickness of cell 'active layer' is reflected by

thickness of the cell layer that possessing positive growth rate (third row). (C) The

linear correlation between the steady-state frequencies of Detoxifier predicted by Eqn.

[4] and those frequencies obtained by our Individual-based simulations. The dashed

line shows the linear curve in which the predicting results is completely identical to

simulated results. The best fitting value of ks, Fd_{max}, kt, TS_{max}, kd_1, and kd_2 in this case

are 30.8, 0.446, 1.46, 1.05, 14000, and 44.8 respectively.