A bacterial prophage small peptide counteracts DnaA activities in *B. subtilis*.

Magali Ventroux and Marie-Francoise Noirot-Gros*

Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78350, Jouy-en-Josas, France.

* Correspondence:
marie-francoise.noirot-gros@inrae.fr

Keywords: small ORF, SEP, skin element, DnaA, Spo0A, *B. subtilis*

Abstract

Bacteriophages are able to hijack host essential machineries to benefit their fitness and assemble their own progeny. Phage proteins targeting major bacterial pathways can be powerful tools to understand cell functions and have possible applications in human health and industry. Bacterial genomes also harbor cryptic prophages carrying genes that may contribute to their host fitness and properties. The cryptic prophages are mostly transcriptionally silent and most of the functions they encode are not annotated. In *B. subtilis,* the 48 kb-long *skin* element is a prophage carrying the *yqaF-yqaN* operon, which is tightly regulated by the Xre-like repressor *sknR*. The small *yqaH* gene potentially encodes the protein *YqaH* in absence of SknR. It was previously reported that YqaH interacts with the replication initiator DnaA in yeast two-hybrid assay and its expression in *B. subtilis* causes defects in the chromosomal cycle. In this study, we report that, in addition to DnaA, YqaH interacts with Spo0A, a master regulator of sporulation. To decipher *yqaH* mode of action, we used the yeast two-hybrid to isolate single mutations in *yqaH* that separate interactions with DnaA and Spo0A. We isolated mutations that caused loss-of-interaction (LOI) with DnaA but not Spo0A. However, all mutations disrupting the interaction with Spo0A were also DnaA-LOI functions, suggesting that these functions could not be separated. We found that expression YqaH carrying DnaA-LOI mutations affects both chromosome integrity and DnaA-mediated transcription, leading to growth inhibition as well as preventing bacterial development such as sporulation and biofilm formation. These results show that YqaH acts as an antimicrobial peptide in *B. subtilis* and pave the way for the structural design of mutants with improved antibacterial action.

1   Introduction

Small proteins encoded by short open reading frames (smORFs) has emerged as a new class of micropeptides widespread in both in eukaryotic and prokaryotic genomes (Albuquerque et al., 2015; Couso and Patraquim, 2017; Hellens et al., 2016; Samayoa et al., 2011; Storz et al., 2014). Due to their small size, smORFs have been overlooked for a long time. However, with advanced computation and ribosome profiling-based biochemical methods, small proteins are being now more

widely identified and some has been functionally characterized (Chu et al., 2015; He et al., 2018;
Samayoa et al., 2011; Straub and Wenkel, 2017; VanOrsdel et al., 2018). Growing evidence indicates
they often encode bioactive peptides (Chu et al., 2015; Saghatelian and Couso, 2015). However, their
contribution to cellular functions remains largely unexplored.

Micropeptides (for the smallest <30aa or microProteins up to 100 aa) were identified to act as
regulators of diverse cellular processes in eukaryotes (Staudt and Wenkel, 2011). In plants,
characterized micro-proteins were found to regulate transcription factors by sequestering them into
nonfunctional states, preventing DNA binding or transcriptional activation (Dolde et al., 2018; Graeff
and Wenkel, 2012). In Drosophila melanogaster, smORF-encoded peptides (SEPs) represent about
5% of the transcriptome and play an important role in controlling drosophila development by
triggering post-translational processing of transcriptional regulators (Albuquerque et al., 2015; Zanet
et al., 2015). In human, SEPs has been discovered with specific subcellular localization, suggesting
they can fulfill biological functions (Slavoff et al., 2013). As example, a 69 aa long SEP, MRI-2,
has been described to play a role in stimulating DNA repair through binding to the DNA end-binding
protein complex Ku (Slavoff et al., 2014). SEPs represent about 2% of the genome of *S. cerevisiae*
(Erpf and Fraser, 2018). Their function remains largely elusive but few has been identified to play
regulatory roles in diverse physiological processes such as iron homeostasis (An et al., 2015) or DNA
synthetis (Chabes et al., 1999; Erpf and Fraser, 2018; Lee et al., 2008). Genomic analysis of the *S.*
cerevisiae sORFs revealed that a significant part are conserved in other eukaryotes even as
phylogenetically distantly related such as humans, thus emphasizing their biological significance
(Kastenmayer et al., 2006).

In Prokaryotes, small proteins are encoded by 10 to 20% of sRNA in average, and are often species-
specific (Friedman et al., 2017; Miravet-Verde et al., 2019; VanOrsdel et al., 2018; Yang et al., 2016;
Zuber, 2001). SEPs with characterized functions were found involved in various cellular processes
(Storz et al., 2014). In *P. aeruginosa*, PtrA (63-aa long) and PtrB (59-aa long) repress the type III
secretion system in response to DNA stress (Ha et al., 2004; Wu and Jin, 2005). In *E. coli*, the 43-aa
long peptide SrgT interferes with the PTS glucose transport system allowing cells to utilize
alternative non-PTS carbon sources to rapidly adapt to environmental changes in nutrient availability
(Lloyd et al., 2017). In *B. subtilis*, SEPs have been found to participate in regulating cell division and
stress responses (Ebmeier et al., 2012; Handler et al., 2008; Schmalisch et al., 2010). A compelling
example is the recently characterized developmental regulator MciZ (mother cell inhibitor of FtsZ), a
40-aa long peptide which prevents cytokinesis in the mother cell during sporulation (Araujo-Bazan et
al., 2019; Bisson-Filho et al., 2015). In this bacteria, about 20% of the total core protein of the mature
spores is composed by the small acid-soluble spore proteins (SASPs) playing an important role in
protecting DNA in the dormant spores (Moeller et al., 2008; Setlow, 2007). Notably, several
smORFs identified in intergenic regions were reported to be expressed during sporulation
(Schmalisch et al., 2010). The sporulation inhibitor *sda* encodes a 52aa long protein, which acts as a
checkpoint system coordinating DNA replication with sporulation initiation (Burkholder et al., 2001;
Cunningham and Burkholder, 2009; Rowland et al., 2004). Its mode of action has been extensively
characterized at molecular level. Sda binds to the primary sporulation kinase KinA, preventing its
activation and subsequent phosphorelay-mediated activation of the master sporulation regulator

Spo0A (Cunningham and Burkholder, 2009; Veening et al., 2009). By linking DNA replication to a
phosphorylation-dependent signaling cascade that triggers cellular development, this system
illustrates an important biological role played by a SEP in blocking sporulation in response to DNA
stress in *bacillus* (Veening et al., 2009).

Small proteins encoded by phage or by prophage-like regions of bacterial genomes have been
identified to hijack the host cellular machineries, as part of a strategy to shift host resources toward
the production of viral progeny (Liu et al., 2014; Duval and Cossart, 2017). The bacteriophage T7
gene 2 encodes the 64 aa-long gp2 protein essential for infection of *E. coli*. Studies of it biological
role revealed that gp2 inhibits transcription by binding to RNA polymerase (RNAP), promoting a
host-to-viral RNAP switch (Nechaev and Severinov, 2003; Savalia et al., 2010). Another illustration
is the 52-aa long protein ORF104 of phage 77 infecting *S. aureus*. ORF104 is able to interfere with
the host chromosome replication by binding to the ATPase domain of the helicase loader protein
DnaI, thus preventing the loading of the DNA helicase DnaC (Liu et al., 2004; Hood and Berger,
2016).

DNA replication is an essential process in all living organisms. Owing to their essentiality, proteins
that compose the orisome and the replisome machineries are potential targets for the development of
antimicrobials. In bacteria, DNA replication is initiated by the conserved initiator protein DnaA that
assembles to the chromosomal replication origin to elicit local DNA strand opening (Hwang and
Kornberg, 1992; Leonard and Grimwade, 2011; Mott and Berger, 2007; Ozaki and Katayama, 2009).
This step triggers the coordinated assembly of the proteins that will further built a functional
replication fork, from the DNA helicase, unwinding the DNA duplex, to the many components of the
replisome that form the replication machinery (Messer, 2002). In addition to its initiator activity,
DnaA acts as a transcription factor repressing or activating genes (Messer and Weigel, 2003). The
activity of the initiator DnaA is tightly controlled to coordinate chromosomal replication initiation
with other cellular processes during the bacterial cell cycle (Katayama et al., 2010; Scholefield and
Murray, 2013). Part of this control is mediated by protein-protein interactions and involves various
protein regulators that bind DnaA and affect his activity (Felicori et al., 2016; Jameson and
Wilkinson, 2017; Katayama et al., 2017; Riber et al., 2016; Skarstad and Katayama, 2013). In *B.*
subtilis, four proteins SirA, Soj, DnaD and YabA have been identified to regulate DnaA activity or
its assembly at OriC through direct interaction (Bonilla and Grossman, 2012; Felicori et al., 2016;
Martin et al., 2019; Murray and Errington, 2008).

Phage SEPs targeting important functions in bacteria are regarded as promising antimicrobial
peptides, and ignited a strong interest in their identification and characterization (Hood and Berger,
2016; Liu et al., 2004). The phage-like element skin of *B. subtilis* encodes about 60 proteins. The
skin element is repressed under most physiological conditions (Nicolas et al., 2012), silenced by the
skin repressor SknR (Figure 1A) (Kimura et al., 2010). Excision of skin from the genome restore
the integrity of sigK gene encoding the late sporulation σK factor (Kunkel et al., 1990). Among the
skin ORFs, *yqaH* encodes the 85 aa-long polypeptide YqaH that has been previously identified to
bind to the replication initiator protein DnaA in a yeast two-hybrid genomic screen (Noirot-Gros et
al., 2002; Marchadier et al., 2011). When *yqaH* is overexpressed, bacillus cells exhibits aberrant

nucleoid morphological defects suggestive of replication deficiency (Kimura et al., 2010). In this study, we further characterized the function of *yqaH* in antagonizing DnaA. In addition, we found that YqaH also interacts with the master regulator Spo0A involved in developmental transitions to sporulation and biofilm formation (Dubnau et al., 2016). As DnaA, Spo0A is a DNA-binding protein, which, under its activated phosphorylated form (Spo0A-P) controls the expression of numerous genes during the early stages of sporulation (Molle et al., 2003). To further understand the biological role of this sORF in *B. subtilis* we performed the functional dissection of YqaH. Using a reverse yeast two-hybrid system, we selected *yqaH* alleles that selectively disrupted the YqaH/DnaA complex. This approach allowed us to link specific DnaA loss-of-interaction with loss-of-function phenotypes

2   Material and Methods

2.1   Strains, plasmids and primers

Experiments were performed in strains 168 trp+ or TF8A a phage-cured strain lacking the prophage-like element 'skin' element as well as the prophages SPβ and PBSX (Nicolas et al., 2012; Westers et al., 2003) (Table S1A). Saccharomyces cerevisiae PJ69-4a or α strains used for yeast-two-hybrid experiments (James et al., 1996). Escherichia coli strain DH10B (Durfee et al., 2008) was used as a cloning host. Plasmids constructs are listed in Table S1B. Primers are listed in Table S2. Sequences of interest cloned or mutated in this study were verified by DNA sequencing.

2.2   Bacterial culture conditions

Bacteria strains were grown at 37°C in LB medium containing ampicillin 100 µg/ml for plasmid selection (in *E. coli*), spectinomycin 60 µg/ml, kanamycin 5 µg/ml or chloramphenicol 5 µg/ml (for *B. subtilis*), and inducer molecules IPTG 0.5 mM, or D-xylose 0.5%, when necessary. *B. subtilis* strains containing pDG148 and pDG148-*yqaH* constructions were grown overnight in LB complemented with kanamycin. ON cultures were then diluted at OD600 0.01on fresh media and grown to mid-exponential phase (OD600 0.3-0.4) prior to addition of IPTG. The effect of *yqaH* expressing on growth has been performed in 96 microwells plates in a final volume of 200 µL.

2.3   Strains constructions

yqaH expression:

The *yqah* wild type and mutated gene derivatives, were PCR-amplified using the *yqaH-HindIII-RBS-F/pYqaH-R* primer pair and inserted in plasmid pDG148 between HindIII and SalI restriction sites to be placed under control of the IPTG-inducible *Pspac* promoter (Stragier et al., 1988). The plasmid constructs were extracted from *E.coli* and transformed in *B. subtilis*. Expression of YqaH wild-type and mutated proteins were assessed using 3xflag-fusions.

Construction of yqaH-gfp fusion:

yqaH WT and mutated gene were PCR-amplified from the pGAD-*yqaH* yeast vector derivatives
carrying either the WT or LOI mutations (see below) using *yqaH-apa* and *pYqaH-R* primer pair.
DNA fragments were digested by ApaI and SalI and inserted in between ApaI and XhoI restriction
sites of pSG1154m, a version of the pSG1154 vector (Lewis and Marston, 1999) encoding a
monomeric version of the GFP (GFPm). Final constructions, consisting of translational fusions of
the *gfpm* gene to the C-terminal region encoded *yqaH* under the control of the P*xyl* promoter, were
transferred at the *amyE* locus of the *B. subtilis* chromosome by transforming competent *B. subtilis*
cells with plasmid DNA. The transfer of single mutations in the *spo0A* chromosomal locus of *B.*
subtilis were performed by gene replacement using the "pop-in pop-out" gene replacement method
(Fabret et al., 2002; Tanaka et al., 2013).
162

Yeast two-hybrid plasmid constructs:

Genes encoding for full size YqaH, Spo0A, YabA and DnaA, as well as truncated DnaA (boundaries
as illustrated figure 1D) were translationally fused to the activating domain (AD) or the binding
domain (BD) of the transcriptional factor Gal4 by cloning into pGAD and pGBDU vectors,
respectively (James et al., 1996). DNA fragments were amplified by PCR using appropriated primer
sets (Table S2), double digested by EcoR1 and SalI and ligated to corresponding pGAD or pGBDU
linearized vector to generate a translational fusion with Gal4-AD or BD domains. Plasmid constructs
were first transformed into *E. coli* prior to be introduced pJ69-4α (pGAD-derivatives) or pJ69-4a
(pGBDU-derivatives) haploid strain. pGAD- and pGBDU- plasmid derivatives were selected onto
SD media lacking leucine (SD-L) or Uracyl (SD-U), respectively (James et al., 1996).
173

2.4   Sporulation conditions

Sporulation of *B. subtilis* was induced by nutrient limitation by the re-suspension in Sterlini-
Mandelstam medium (SM) (Sterlini and Mandelstam, 1969). The beginning of sporulation (t0) is
arbitrarily defined as the moment of re-suspension of the cells in SM. To study the effects of *yqaH*
overexpression on *B. subtilis* sporulation, ON cultures of strains containing the pDG148
constructions were then diluted in CH medium at starting OD600 of 0.05. When cultures reached an
OD600=1, cells were re-suspended in an equivalent volume of SM medium complemented with
0.5M IPTG (t0).To monitored sporulation efficiency, cells were collected at different times after
sporulation induction (t0) until t6 (6 hours after induction, defined as a stage of production of mature
spores) and t18. Asymmetric septa and spores were enumerated by microscopic observations from all
samples.

2.5   Spores survival assay

Cells were spread and grown ON at 30°C on DSM agar medium (Schaeffer et al., 1965), prior to be
inoculated in preheated liquid DSM at OD600 0.1 and incubated at 37°C until reaching OD600 1.5.
From this point (taken as T0) cultures were grown for 20 h to produce matured resistant spores.
Samples were collected and half of the material was heated at 80°C (10 min) before plating to kill
potential vegetative cells prior to be plated on LB. Colony counts were performed after 36 hours at

37°C. The percentage of spores was calculated as the ratio of colonies forming units (cfu) from
heated and unheated samples.

2.6   Yeast-two-hybrid assay

The yeast-two-hybrid assays were performed as described (Marchadier et al., 2011; Noirot-Gros et
al., 2002; Noirot-Gros et al., 2006). PJ69-4A and α haploid yeast strains transformed by pGAD- and
pGBDU- plasmid derivatives were mixed onto YPD-rich media plates to allow formation of diploids.
Diploids containing both pGAD and pGBDU type of plasmids were then selected on SD-UL and
interacting phenotypes were monitored by the ability of diploids to grow on SD-LUHA medium
further lacking histidine (H) or adenine (A).

2.7   Generation of Loss of Interaction (LOI) mutation

YqaH_LOI mutants were identified using a yeast two-hybrid-based assay as described elsewhere
(Natrajan et al., 2009; Noirot-Gros et al., 2006; Quevillon-Cheruel et al., 2012). Random mutagenesis
of the targeted genes were achieved by PCR amplification under mutagenic conditions that promotes
less that one miss incorporation per amplification cycle (Noirot-Gros et al., 2006). For *yqaH*, a
library of mutated pGAD-*yqaH** were constructed by gap-repair recombination into yeast (PJ69–4α
strain). About 1000 individual transformants were organized in 96-well format on plates containing a
defined medium lacking leucine (-L) to form an arrayed collection of AD-*yqaH** gene fusion
mutants. This organized library was then mated with PJ69–4a strains containing either pGBDU-
dnaA or pGBDU-*spo0A*, or an empty pGBDU plasmid as a negative control. Selective pressure for
interacting phenotypes was then applied on media lacking –LUH or –LUA. Diploids that failed to
grow on interaction selective media are considered as potentially expressing a loss-of-interaction
(LOI) mutant of *yqaH*. Importantly, any particular AD-YqaH*_LOI proteins unable to trigger
interacting phenotypes in the presence of BD-DnaA while still producing interacting phenotypes
when expressed in the presence of BD-Spo0A is defined as a DnaA-specific LOI mutant. The
corresponding *yqaH_LOI* genes were retrieved from the initially organized haploid library and the
mutations identified by DNA sequencing. Only mutations resulting from single substitutions were
considered.

2.8   Ori/Ter ratios measurements by qPCR

To determine the ratio of amounts of origin-proximal and terminus-proximal DNA sequences, ON
cultures of *B. subtilis* strains containing pDG148 or pDG148-*yqaH* plasmids were freshly diluted
(OD600=0.01) in LB supplemented with kanamycin 5 µg/mL in the presence of IPTG 0.5 mM, and
grown at 37°C in up to mid-exponential phase (OD600 = 0.3 to 0.4). A determined volume of culture
was taken, mixed with similar volume of sodium azide solution (1%, 0.5% final) to stop all metabolic
activities, and subjected to total lysis. Total genomic DNA extracts were kept at -80°C and thawed
aliquots were used only once. Quantitative real time PCR were performed on a Mastercycler® ep
realplex (Eppendorf) thermocycler device using ABsolute[TM] Blue QPCR SYBR® Green ROX Mix
(ABgene), to amplify specific origin or terminus proximal sequences. Primers used for sequence
amplification were chosen using Primer3Plus program (http://www.bioinformatics.nl/cgi-

bin/primer3plus/primer3plus.cgi/). Amplification using the ORI pair of primers (oriL3F and oriL3R,
Table S2) targeting the 4212889-4211510 region of the *B. subtili*s chromosome, yields a 128 bp size
product corresponding to sequence at the left side of the origin. The terminus sequence is a 122 bp
long fragment obtained from the TER pair of primers (terR3F and terR3R, Table S2) amplifying the
region 2016845-2017711 at the right side of the terminus. The two primer pairs ORI and TER
exhibited ≥ 95% of amplification efficiency. Data analysis was performed using the software
Realplex (Eppendorf) and the quantification with the ΔΔCt method. The formula reflects the efficacy
(Eff) of the primers (Rate = (1+Eff)^(-ΔΔCt)).

2.9   Gene expression quantifications

To quantify the expression rates of genes regulated by DnaA, sample of cells (as collected for Ori/Ter
ratios measurements) were also harvested in exponential growth at OD600=0.3-0.4, and RNA
extractions were performed. To quantify the expression rates of genes regulated by Spo0A,
sporulating cells were harvested when reaching stages T2-T3 of sporulation ($t_{2,5}$) in SM medium. In
both case, a determined volume of culture was mix with equivalent volume of centrifuged of sodium
azide solution, and subjected to lysis followed by total RNA extraction as described (Nicolas et al.,
2012). Total RNA was reversely transcribed and quantitative real time PCR were performed on
cDNA. Primers used for sequence amplification were chosen using Primer3Plus program
(http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi/) and are detailed in Table S2.
The couples of primers exhibited ≥ 95% amplification efficiency. Data analysis was performed using
the software Realplex (Eppendorf) and the quantification with the ΔΔCt method. The formula reflects
the efficacity (Eff) of the primers (Rate = (1+Eff)^(-ΔΔCt)).

2.10   Fluorescence microscopy

The cytolocalization of proteins fused to GFP were observed by epifluorescence microscopy on
vegetative or sporulating cells. The expression of *yqaH-gfp* gene fusions from the *B. subtilis amyE*
chromosomal locus was trigered by supplementing exponential growth or sporulation cultures with
xylose 0.5%. Sample of cells were harvested and rinsed in a minimal transparent media to avoid LB
auto-fluorescence prior to be mounted onto 1.2% agarose pads. When necessary, bacterial
membranes were stained with FM4-64 dye and nucleoids with DAPI. Fluorescence microscopy was
performed on a Leica® DMR2A (100X UplanAPO objective with an aperture of 1.35.) coupled with
CoolSnap HQ camera (Roper Scientific). System control and image processing were achieved using
Metamorph software (Molecular Devices, Sunnyvale, CA, USA). Counts of cells, spores, foci or
nucleoids were determined with the ImageJ® software, from at least 500 cells.

2.11   Biofilm assay

Production and analysis of air-to-liquid biofilm pellicles were performed as already described (Garcia
Garcia et al., 2018). Briefly, strains expressing yqaH, wild type or K17E mutant derivative as well as
control strain were grown in LB to OD600 of 1.0 and inoculated in 12-well culture plates containing

3.5 ml of MSgg media at starting OD600 = 0.1. Cultures were maintained at 28°C and 70% humidity,
with no agitation. After 48 hours, wells were filled out with MSgg media (slowly added at the edge)
to lift the biofilm pellicles up to the top of the wells. The pellicles were then peeled-off onto a 2.5 cm
diameter circular cover slide. The cover slides with intact biofilm pellicles were mounted onto an
Attofluor Cell Chamber and stained with the Film Tracer FM 1-43 Green Biofilm dye (Thermo
Fisher Scientific). Stained biofilms were observed using a spinning disk confocal microscope [Nikon
Eclipse Ti-E coupled with CREST X-LightTM confocal imager; objectives Nikon CFI Plan Fluor
10X, DIC, 10x/0.3 NA (WD = 16 mm); excitation was performed at 470 nm and emission recorded
at 505 nm]. Images were processed using IMARIS software (Bitplane, South Windsor, CT, United
States). Biofilm images were quantified using the surface function in IMARIS (XTension biofilm).
Biovolumes were calculated based on total volume (μm3) per area (μm2) from n ≥ 4 samples.

2.12 Protein immunodectection

Production of 3FLAG-YqaH mutated derivatives was determined from total protein extract of *B.*
subtilis by immuno-detection using monoclonal antibody anti-FLAG® M2 (Sigma). An IgG goat
secondary antibody (Sigma) peroxidase conjugated was used at 1/10000e to detect the anti-FLAG®
M2. Proteins immuno-detections were performed using the Clarity Western ECL kit (Biorad)
according to the supplier's indications followed by chemiluminescence detection (ChemiDoc imager,
Biorad). Images were analysed with the software Image Lab™.

2.13 Structure predictions

Secondary structures predictions were performed with the computer server Jpred 3
(http://www.compbio.dundee.ac.uk/www-jpred/) and 3D structures predictions with the server Phyre[2]
(http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index).

3   Results

3.1   The Skin element SPORF YqaH interacts with DnaA and Spo0A

YqaH was originally identified as interacting with the replication initiator DnaA in a yeast two-
hybrid screen of a *B. subtilis* genomic library (Noirot-Gros et al., 2002). Interestingly, we also
identified YqaH as binding partner to Spo0A in a subsequent screen targeting the sporulation
transcriptional factor (Figure 1B). Other DnaA-binding proteins SirA, Soj, DnaD and YabA also
exert regulatory functions by interacting with different functional domains of DnaA (Figure 1B). To
gain insights into YqaH mode of action, we characterized the minimal interacting domain of DnaA.
We defined the C-terminal domain IV as necessary and sufficient to trigger the interaction
phenotypes in a yeast two-hybrid binary assay (Figure 1C). This 118 AA long fragment spans residue
328 to 446, and carries the signature motif for binding to double stranded DNA. This result
distinguishes YqaH from other regulators, found to inhibit DnaA binding to *oriC* through interacting
with domains I, responsible for self-oligomerization (SirA), and/or the central AAA+ ATPase
domain III, (SirA, Soj, YabA and DnaD).

3.2   YqaH expression triggers *dnaA*-related mutant phenotypes

The interaction of YqaH with DnaA hints at a potential role in replication initiation control. We first
investigated the effect of *yqaH* expression on cell growth and viability. *B. subtilis* strains carrying a
plasmid expressing *yqaH* under control of an IPTG-inducible promoter or carrying a control plasmid
with no expressed gene were treated under identical conditions Addition of IPTG at early exponential
stage specifically halted cell growth after two hours in cells expressing *yqaH* (Figure 2A). We found
that cell viability was affected almost instantly in the presence of *yqaH*, leading to about 30x
decrease after three hours (Figure 2A). Expression of *yqaH* also affects cell morphology leading to
significant filamentation (Figure 2B). Closer examination of nucleoids revealed aberrant segregation
of chromosomes with both diffused and compacted nucleoids unequally distributed within filaments,
as well as a large portion of cells with no DNA. Most importantly, septum-entrapped nucleoids were
also observed indicative of impaired nucleoid occlusion (NO) (Wu and Errington, 2011). These
observations highlighted that the expression of YqaH triggered a large panel of chromosomal
disorders suggestive of replication and segregation stress, and compatible with replication initiation
deficiency, in agreement to earlier described (Kimura et al., 2010). We further investigate a role of
YqaH by investigating DnaA-dependant transcriptional regulation. Among genes of the DnaA-
regulon, active DnaA-ATP negatively regulate its own expression by binding to *dnaA* promoter
region (Goranov et al., 2005). We monitored the *dnaA* mRNA levels in the presence or absence of
YqaH and shown that expression of *yqaH* led to a 2 fold increase in *dnaA* mRNA, indicating a
regulatory defect. Taking all together, these results are in agreement with a direct role of YqaH in
counteracting DnaA activity.
325

3.3   YqaH expression impairs sporulation

We investigated the possible role of YqaH in Spo0A functions, by examining the effect of *yqaH*
expression on sporulation. Cells carrying either the control or the *yqaH*-inducible plasmids were put
to sporulate by the re-suspension method. To prevent an inhibitory effect of YqaH during vegetative
growth, the *yqaH* gene expression was induced only at the onset of sporulation (T0). The formation
of spores was monitored over time. Appearance of asymetric septa at early stage (T2,5) was imaged
by fluorescent microscopy after staining by a red-fluorescent membrane-dye, while the engulfed
forespore (T6 and T18) was revealed using bright field (Figure. 3A). We observed that the counts of
spore forming bacteria was drastically affected in the presence of YqaH (Figure 3B). We also
examined the effect of *yqaH* expression on Spo0A-mediated transcriptional regulation. Among the
genes under the direct control of Spo0A are *spoIIE*, encoding the protein serine phosphatase SpoIIE
and *spoIIGA*, encoding a pro- σE processing protease (Molle et al., 2003). These two genes are
involved in the activation of the alternative sporulation sigma factors σF and σE in the forespore and
mother cell compartments, respectively (Baldus et al., 1994; Bradshaw et al., 2017; Errington and
Wu, 2017; Fujita et al., 2005). Examination of their expression at the onset of sporulation revealed a
significant downregulation (about 7 and 8 fold, respectively) in the presence of *yqaH* (Figure 3C).
Altogether, these results pointed to a negative effect of YqaH in Spo0A-mediated function.
343

3.4   Functional dissection of YqaH

To further decipher YqaH mechanism, we looked for *yqaH* separation of function mutations that selectively affect its ability to interact to one protein partner while preserving its interaction with the other. Using a yeast two-hybrid based assay, we screened a *yqaH* mutant library specific loss-of-interaction (LOI) phenotypes (Figure S1A). This approached is based on the selection of single amino acid changes in the protein of interest that selectively disrupt the interaction with one partner without affecting the interaction with the other partner. Such screening favors the substitution of residues located at the interacting surface, leading to a LOI phenotype with the targeted partner, while preserving the overall 3D-structure of the protein that remains proficient for interaction with another partner (Natrajan et al., 2009; Noirot-Gros et al., 2006; Quevillon-Cheruel et al., 2012). We identify three single residue substitutions K17E, E38K and K48E that elicited a complete loss-of-interaction phenotype with DnaA without affecting interaction with Spo0A (Figure S1B, Table 1). Six additional substitutions affecting residues R16G, M27T, Y37H, E38V A44V and R56W were only partially abolishing interaction phenotypes with DnaA (Figure S1B, Table 1). Two substitutions, S25T and A40T were found to partially affect both interaction with DnaA and Spo0A, and 4 substitutions, D10G, S20L, L28P, L43P and L57P were totally abolishing interaction phenotypes with both DnaA and Spo0A (Figure S1B, Table1). In the latter case, these substitutions could have affected the overall YqaH 3D-structure integrity. Is it worthy of note that, while several specific DnaA-LOI mutants were obtained, no substitution that specifically prevent interaction with Spo0A were identified in our screen. The YqaH protein is predicted to fold into three successive alpha helix (Figure S1B). The residues important for interaction with DnaA mapped within the two main helix presumably involved in a coiled-coil structure.

3.5   DnaA-LOI mutants of yqaH restore replication and transcriptional regulation defects caused by YqaH expression

We investigated the effect of two DnaA-LOI mutants carrying substitutions K17E and R56W, exhibiting total (K17E) or partial (R56W) loss of interaction phenotypes with YqaH in our yeast two-hybride assay. (Table 1). Both substitutions abolished the nucleoid segregation defects resulting from *yqaH* expression (Figure 4A). To rule out that this phenotypic compensation could potentially result from expression of the gene copy from the silent SknR operon, we confirmed that the only source of YqaH in the cell arisen from plasmidic expression (Figure S3). No YqaH protein was detected in cells carrying a control plasmid devoid of yqaH coding sequence, indicating that the chromosomal copy remained completely silent during our experimental conditions.

We further examined the effect of YqaH on replication initiation by measuring the ratio between chromosomal origin-proximal and terminus-proximal sequences (Figure 4B). The average of origin-to-terminus (*ori:ter*) ratio was determined by qPCR on genomic DNA harvested from exponentially growing cells as previously described (Soufo et al., 2008). The *ori:ter* ratio was about 4 in control cells that do not expressed *yqaH*, indicating that two events of replication initiation has taken place in most cells, in agreement to already observed in similar experimental conditions (Murray and Koh, 2014). In the presence of *yqaH*, the *ori:ter* ratio dropped to 2, indicating that only one replication

initiation event per cell has taken place in average (Figure 4B). This observation suggests that the
YqaH protein could exerts a negative effect on replication initiation by counteracting DnaA activity.
We observed that this effect was abolished upon expression of the *yqaH*_DnaA LOI variants carrying
either the K17E or the R56W substitutions (Figure 4B). In cells, expressing YqaH-R56W the
replication initiation rate was similar to control cells while expression of YqaH-K17E restored about
75% of the initiation rate indicating a significant but partial complementation. In another assay, we
analyzed the number of chromosomal origin in individual cells using the *lacO*/LacI-GFP system.
Cells carrying a *lacO* repeat array near the replication origin and expressing the LacI-GFP fusion
were observed by fluorescence microscopy, in the presence of absence of YqaH (Figure 4C). In the
absence of YqaH, the number of lacI foci per nucleoid was found to be 3.6 in average, in agreement
with the *ori-to-ter* ratio (Figure 4D). This number dropped to 1 upon expression of YqaH, suggestive
of replication deficiency. As observed earlier, cell were also exhibiting aberrant nucleoids and
segregation defects. Conversely, expression of YqaH-K17E mutant fully restored nucleoid and cell
morphology and partially restored replication defects with an average of LacI-GFP foci about 2.3.
Together these observations indicate that YqaH binds to DnaA and antagonized its activity in
replication initiation.

To substantiate the role of YqaH/ DnaA interaction, we also examined the effect of the DnaA-LOI
mutant on *dnaA* transcription (Figure 4E). We found that the loss of DnaA-mediated control of *dnaA*
expression in the presence of wild type YqaH was restored by the introduction of both K17E and
R56W substitutions. This result further supports the conclusion that the binding of YqaH to DnaA
antagonizes DnaA functions.

3.6   The YqaH-mediated defects in sporulation and biofilm formation requires interaction
with DnaA.

Although we did not identified any YqaH_SpoA-LOI mutant in our screen, we thought that the
YqaH_K17E DnaA-LOI mutant provided for a convenient separation-of-function for a more reliable
investigation of the potential role of *yqaH* in *spo0A*-mediated phenotypes. Indeed, this K17E variant
remains fully proficient for interacting with Spo0A in our yeast 2HB assays, allowing to investigate
the role of *yqaH* during sporulation while circumventing DnaA-related defects during vegetative
cycle. We compared the sporulation efficiencies of strains expressing the wild-type YqaH or the
YqaH_K17E variant in both 168 and TF8A (devoid of Skin element) backgrounds. The percent of
cells ongoing sporulation was determined at t6 (i.e stage VI, refers as the spore maturation stage,
Figure 5A). Surprisingly, we found that the K17E-LOI substitution abolished the defect elicited by
the wild type YqaH (Figure 5). This observation indicates that the sporulation deficiency mediated by
YqaH requires its ability to interact with DnaA. To further investigate the potential role of YqaH in
Spo0A-related processes, we also examined its effect on biofilm formation (Figure 5B, Figure S4).
We observed that the production of biofilm pellicles at the air-medium interface was impaired in
strains expressing YqaH, leading to a loss of biomass and cohesion. Damaged biofilm pellicles were
not observed in cells expressing the YqaH_K17E DnaA-LOI mutant (Figure 5B, Figure S4B,C) thus
pointing to a role of DnaA during biofilm formation.

4   Discussion

Our study sheds light on the mechanism of action of YqaH, a small protein with antimicrobial
activities encoded by the *B. subtilis* skin element. By physically interacting with two master
regulators DnaA and Spo0A, the YqaH protein holds the promise to inhibit two essential cellular
processes under two different life styles of the bacteria. Upon nutrient stress and other environmental
conditions, B. subtilis cells switch from vegetative growth to sporulation or biofilm formation. DnaA
is essential during vegetative growth, acting as both replication initiator and transcriptional factor,
while the response regulator Spo0A controls developmental transitions.

In Bacillus, several regulatory proteins inhibit DnaA by targeting different functional domains of the
protein. In cells committed to sporulation, the protein SirA prevents DnaA from binding to the
replication origin OriC by interacting with its structural domains I and III (Jameson et al., 2014).
During vegetative growth, the regulatory proteins YabA interacts with DnaA during most of the cell
cycle (Felicori et al., 2016; Noirot-Gros et al., 2006; Soufo et al., 2008). YabA as well as the
primosomal protein DnaD affect DnaA cooperative binding to OriC by interacting with DnaA
structural domain III (Merrikh and Grossman, 2011; Scholefield and Murray, 2013). Finally, the
ATPase protein Soj negatively regulates DnaA by also interacting with the structural domain III,
preventing oligomerization (Scholefield et al., 2011). Our results indicate that YqaH controls DnaA
activity by a mode distinct from the other known regulators, specifically by contacting the structural
domain IV of DnaA that is responsible for DnaA-box sequence recognition (Fujikawa et al., 2003).

B. subtilis cells expressing YqaH exhibited various DnaA-related phenotypes that spanned from a
general growth defect, aberrant nucleoid morphology, impaired replication initiation and loss of
transcriptional control (Figure 2 and 4). These cells also exhibited Spo0A-related phenotypes such as
a dramatic reduction of sporulation efficiency (Figure 3A, B), an inhibition of the expression of
Spo0A-dependent genes (Figure 3C) and a strong impairment of biofilm formation (Figure 5). These
observations are in agreement with a role of YqaH in counteracting both DnaA and Spo0A activities
during vegetative growth and sporulation. However, it is well documented that initiation of
sporulation is closely coupled to the cell cycle and DNA replication in order to ensure that
sporulation occurs only in cells containing two fully replicated chromosomes (Veening et al., 2009).
Because of this intricate relationship between DNA replication and sporulation, it is difficult to
separate the phenotypes of YqaH related to its interactions with DnaA on one hand and Spo0A on the
other hand. By using YqaH single point mutants unable to interact with DnaA while remaining
proficient for interaction with Spo0A, we identified mutational pattern on YqaH (Figure S2) and
demonstrated their direct involvement in DnaA-related a loss-of-function phenotype. In our screens,
we did not obtain YqaH mutants that affected only the interaction with Spo0A. However, the YqaH
DnaA_LOI mutants revealed that sporulation and biofilm formation phenotypes of YqaH expression
required interaction with DnaA. Thus, we could not establish whether interaction with Spo0A plays a
role in these phenotypes. Our results could be explained by the role of DnaA as a pleiotropic
transcriptional regulator controlling the expression of several sporulation genes, including the
checkpoint histidine kinase Sda (Washington et al., 2017). Sda is known to bind to KinA, inhibiting
its phosphorylation activity required to activate Spo0A (Burkholder et al., 2001). A recent study

highlighted that a large part of the role of DnaA into gene expression is indirect and mediated by sda
(Washington et al., 2017). Sda is positively regulated by both DnaA and Spo0A, and a coupling
between expression of Sda and the replication cycle results in cyclic bursts of Spo0A activation
(Narula et al., 2015). Many genes that appear regulated by DnaA are involved in sporulation and
biofilm formation, illustrating the complexity of regulatory circuits that control transitions in
lifestyle. In conclusion, our study characterizes the role of small peptide YqaH that negatively
interfers with a broad range of cellular processes by counteracting the activity of DnaA. Controlling
YqaH expression could be part of a strategy to prevent the dissemination of engineered B. subtilis
strains in a controlled environment.

5   Acknowledgements

6   Author contributions

MV and MFNG designed the experiments. MV performed all experiments. MV and MFNG
interpreted the results. MFNG wrote the manuscript. Both authors contributed to data interpretation
and reviewed the final version of the manuscript.

7   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

8   Supplementary material

Figure S1: Yeast 2HB screening of a yqaH gene mutant library for specific LOI phenotypes.

Figure S2: Mapping of the YqaH single LOI mutations with DnaA

Figure S3: Immunodetection of YqaH in cell extracts after induction

Figure S4: DnaA-dependant YqaH-mediated defects in biofilm formation

Figure S5: ORF conservation within bacillus species

Table S1: Strains and Plasmids

Table S2: Primers list

9   References

Albuquerque, J.P., Tobias-Santos, V., Rodrigues, A.C., Mury, F.B., and da Fonseca, R.N. (2015).
small ORFs: A new class of essential genes for development. Genetics and molecular biology *38*,
278-283.

An, X., Zhang, C., Sclafani, R.A., Seligman, P., and Huang, M. (2015). The late-annotated small
ORF LSO1 is a target gene of the iron regulon of Saccharomyces cerevisiae. MicrobiologyOpen *4*,
941-951.

Araujo-Bazan, L., Huecas, S., Valle, J., Andreu, D., and Andreu, J.M. (2019). Synthetic
developmental regulator MciZ targets FtsZ across Bacillus species and inhibits bacterial division.
Mol Microbiol *111*, 965-980.

Baldus, J.M., Green, B.D., Youngman, P., and Moran, C.P., Jr. (1994). Phosphorylation of Bacillus
subtilis transcription factor Spo0A stimulates transcription from the spoIIG promoter by enhancing
binding to weak 0A boxes. J Bacteriol *176*, 296-306.

Bisson-Filho, A.W., Discola, K.F., Castellen, P., Blasios, V., Martins, A., Sforca, M.L., Garcia, W.,
Zeri, A.C., Erickson, H.P., Dessen, A.*, et al.* (2015). FtsZ filament capping by MciZ, a
developmental regulator of bacterial division. Proc Natl Acad Sci U S A *112*, E2130-2138.

Bonilla, C.Y., and Grossman, A.D. (2012). The primosomal protein DnaD inhibits cooperative DNA
binding by the replication initiator DnaA in Bacillus subtilis. J Bacteriol *194*, 5110-5117.

Bradshaw, N., Levdikov, V.M., Zimanyi, C.M., Gaudet, R., Wilkinson, A.J., and Losick, R. (2017).
A widespread family of serine/threonine protein phosphatases shares a common regulatory switch
with proteasomal proteases. Elife *6*.

Burkholder, W.F., Kurtser, I., and Grossman, A.D. (2001). Replication initiation proteins regulate a
developmental checkpoint in Bacillus subtilis. Cell *104*, 269-279.

Chabes, A., Domkin, V., and Thelander, L. (1999). Yeast Sml1, a protein inhibitor of ribonucleotide
reductase. J Biol Chem *274*, 36679-36683.

Chu, Q., Ma, J., and Saghatelian, A. (2015). Identification and characterization of sORF-encoded
polypeptides. Crit Rev Biochem Mol Biol *50*, 134-141.

Couso, J.P., and Patraquim, P. (2017). Classification and function of small open reading frames.
Nature reviews Molecular cell biology *18*, 575-589.

Cunningham, K.A., and Burkholder, W.F. (2009). The histidine kinase inhibitor Sda binds near the
site of autophosphorylation and may sterically hinder autophosphorylation and phosphotransfer to
Spo0F. Mol Microbiol *71*, 659-677.

Dolde, U., Rodrigues, V., Straub, D., Bhati, K.K., Choi, S., Yang, S.W., and Wenkel, S. (2018).
Synthetic MicroProteins: Versatile Tools for Posttranslational Regulation of Target Proteins. Plant
physiology *176*, 3136-3145.

Dubnau, E.J., Carabetta, V.J., Tanner, A.W., Miras, M., Diethmaier, C., and Dubnau, D. (2016). A
protein complex supports the production of Spo0A-P and plays additional roles for biofilms and the
K-state in Bacillus subtilis. Mol Microbiol *101*, 606-624.

Durfee, T., Nelson, R., Baldwin, S., Plunkett, G., Burland, V., Mau, B., Petrosino, J.F., Qin, X.,
Muzny, D.M., Ayele, M.*, et al.* (2008). The complete genome sequence of Escherichia coli DH10B:
Insights into the biology of a laboratory workhorse. Journal of Bacteriology *190*, 2597-2606.

Ebmeier, S.E., Tan, I.S., Clapham, K.R., and Ramamurthi, K.S. (2012). Small proteins link coat and
cortex assembly during sporulation in Bacillus subtilis. Mol Microbiol *84*, 682-696.

Erpf, P.E., and Fraser, J.A. (2018). The Long History of the Diverse Roles of Short ORFs: sPEPs in
Fungi. Proteomics *18*, e1700219.

Errington, J., and Wu, L.J. (2017). Cell Cycle Machinery in Bacillus subtilis. Sub-cellular
biochemistry *84*, 67-101.

Fabret, C., Ehrlich, S.D., and Noirot, P. (2002). A new mutation delivery system for genome-scale
approaches in Bacillus subtilis. Mol Microbiol *46*, 25-36.

Felicori, L., Jameson, K.H., Roblin, P., Fogg, M.J., Garcia-Garcia, T., Ventroux, M., Cherrier, M.V.,
Bazin, A., Noirot, P., Wilkinson, A.J.*, et al.* (2016). Tetramerization and interdomain flexibility of
the replication initiation controller YabA enables simultaneous binding to multiple partners. Nucleic
Acids Res *44*, 449-463.

Friedman, R.C., Kalkhof, S., Doppelt-Azeroual, O., Mueller, S.A., Chovancova, M., von Bergen, M.,
and Schwikowski, B. (2017). Common and phylogenetically widespread coding for peptides by
bacterial small RNAs. BMC genomics *18*, 553.

Fujita, M., Gonzalez-Pastor, J.E., and Losick, R. (2005). High- and low-threshold genes in the Spo0A
regulon of Bacillus subtilis. J Bacteriol *187*, 1357-1368.

Graeff, M., and Wenkel, S. (2012). Regulation of protein function by interfering protein species.
Biomolecular concepts *3*, 71-78.

Ha, U.H., Kim, J., Badrane, H., Jia, J., Baker, H.V., Wu, D., and Jin, S. (2004). An in vivo inducible
gene of Pseudomonas aeruginosa encodes an anti-ExsA to suppress the type III secretion system.
Mol Microbiol *54*, 307-320.

Handler, A.A., Lim, J.E., and Losick, R. (2008). Peptide inhibitor of cytokinesis during sporulation
in Bacillus subtilis. Mol Microbiol *68*, 588-599.

He, C., Jia, C., Zhang, Y., and Xu, P. (2018). Enrichment-Based Proteogenomics Identifies
Microproteins, Missing Proteins, and Novel smORFs in Saccharomyces cerevisiae. J Proteome Res
17, 2335-2344.

Hellens, R.P., Brown, C.M., Chisnall, M.A.W., Waterhouse, P.M., and Macknight, R.C. (2016). The
Emerging World of Small ORFs. Trends in plant science *21*, 317-328.

Hood, I.V., and Berger, J.M. (2016). Viral hijacking of a replicative helicase loader and its
implications for helicase loading control and phage replication. Elife *5*.

Hwang, D.S., and Kornberg, A. (1992). Opening of the replication origin of Escherichia coli by
DnaA protein with protein HU or IHF. J Biol Chem *267*, 23083-23086.

James, P., Halladay, J., and Craig, E.A. (1996). Genomic libraries and a host strain designed for
highly efficient two-hybrid selection in yeast. Genetics *144*, 1425-1436.

Jameson, K.H., and Wilkinson, A.J. (2017). Control of Initiation of DNA Replication in Bacillus
subtilis and Escherichia coli. Genes *8*.

Kastenmayer, J.P., Ni, L., Chu, A., Kitchen, L.E., Au, W.C., Yang, H., Carter, C.D., Wheeler, D.,
Davis, R.W., Boeke, J.D.*, et al.* (2006). Functional genomics of genes with small open reading
frames (sORFs) in S. cerevisiae. Genome research *16*, 365-373.

Katayama, T., Kasho, K., and Kawakami, H. (2017). The DnaA Cycle in Escherichia coli:
Activation, Function and Inactivation of the Initiator Protein. Front Microbiol *8*, 2496.

Katayama, T., Ozaki, S., Keyamura, K., and Fujimitsu, K. (2010). Regulation of the replication
cycle: conserved and diverse regulatory systems for DnaA and oriC. Nat Rev Microbiol *8*, 163-170.

Kimura, T., Amaya, Y., Kobayashi, K., Ogasawara, N., and Sato, T. (2010). Repression of sigK
intervening (skin) element gene expression by the CI-like protein SknR and effect of SknR depletion
on growth of Bacillus subtilis cells. J Bacteriol *192*, 6209-6216.

Lee, Y.D., Wang, J., Stubbe, J., and Elledge, S.J. (2008). Dif1 is a DNA-damage-regulated facilitator
of nuclear import for ribonucleotide reductase. Molecular cell *32*, 70-80.

Leonard, A.C., and Grimwade, J.E. (2011). Regulation of DnaA assembly and activity: taking
directions from the genome. Annual review of microbiology *65*, 19-35.

Lewis, P.J., and Marston, A.L. (1999). GFP vectors for controlled expression and dual labelling of
protein fusions in Bacillus subtilis. Gene *227*, 101-110.

Liu, J., Dehbi, M., Moeck, G., Arhin, F., Bauda, P., Bergeron, D., Callejo, M., Ferretti, V., Ha, N.,
Kwan, T.*, et al.* (2004). Antimicrobial drug discovery through bacteriophage genomics. Nature
biotechnology *22*, 185-191.

Lloyd, C.R., Park, S., Fei, J., and Vanderpool, C.K. (2017). The Small Protein SgrT Controls
Transport Activity of the Glucose-Specific Phosphotransferase System. J Bacteriol *199*.

Marchadier, E., Carballido-Lopez, R., Brinster, S., Fabret, C., Mervelet, P., Bessieres, P., Noirot-
Gros, M.F., Fromion, V., and Noirot, P. (2011). An expanded protein-protein interaction network in
Bacillus subtilis reveals a group of hubs: Exploration by an integrative approach. Proteomics *11*,
2981-2991.

Martin, E., Williams, H.E.L., Pitoulias, M., Stevens, D., Winterhalter, C., Craggs, T.D., Murray, H.,
Searle, M.S., and Soultanas, P. (2019). DNA replication initiation in Bacillus subtilis: structural and
functional characterization of the essential DnaA-DnaD interaction. Nucleic Acids Res *47*, 2101-
2112.

Messer, W. (2002). The bacterial replication initiator DnaA. DnaA and oriC, the bacterial mode to
initiate DNA replication. FEMS microbiology reviews *26*, 355-374.

Messer, W., and Weigel, C. (2003). DnaA as a transcription regulator. Methods in enzymology *370*,
338-349.

Miravet-Verde, S., Ferrar, T., Espadas-Garcia, G., Mazzolini, R., Gharrab, A., Sabido, E., Serrano,
L., and Lluch-Senar, M. (2019). Unraveling the hidden universe of small proteins in bacterial
genomes. Molecular systems biology *15*, e8290.

Moeller, R., Setlow, P., Horneck, G., Berger, T., Reitz, G., Rettberg, P., Doherty, A.J., Okayasu, R.,
and Nicholson, W.L. (2008). Roles of the major, small, acid-soluble spore proteins and spore-specific
and universal DNA repair mechanisms in resistance of Bacillus subtilis spores to ionizing radiation
from X rays and high-energy charged-particle bombardment. J Bacteriol *190*, 1134-1140.

Molle, V., Fujita, M., Jensen, S.T., Eichenberger, P., Gonzalez-Pastor, J.E., Liu, J.S., and Losick, R.
(2003). The Spo0A regulon of Bacillus subtilis. Mol Microbiol *50*, 1683-1701.

Mott, M.L., and Berger, J.M. (2007). DNA replication initiation: mechanisms and regulation in
bacteria. Nat Rev Microbiol *5*, 343-354.

Murray, H., and Errington, J. (2008). Dynamic control of the DNA replication initiation protein
DnaA by Soj/ParA. Cell *135*, 74-84.

Murray, H., and Koh, A. (2014). Multiple regulatory systems coordinate DNA replication with cell
growth in Bacillus subtilis. PLoS Genet *10*, e1004731.

Natrajan, G., Noirot-Gros, M.F., Zawilak-Pawlik, A., Kapp, U., and Terradot, L. (2009). The
structure of a DnaA/HobA complex from Helicobacter pylori provides insight into regulation of
DNA replication in bacteria. Proc Natl Acad Sci U S A *106*, 21115-21120.

Nicolas, P., Mader, U., Dervyn, E., Rochat, T., Leduc, A., Pigeonneau, N., Bidnenko, E., Marchadier,
E., Hoebeke, M., Aymerich, S.*, et al.* (2012). Condition-dependent transcriptome reveals high-level
regulatory architecture in Bacillus subtilis. Science *335*, 1103-1106.

Noirot-Gros, M.F., Dervyn, E., Wu, L.J., Mervelet, P., Errington, J., Ehrlich, S.D., and Noirot, P.
(2002). An expanded view of bacterial DNA replication. Proc Natl Acad Sci U S A *99*, 8342-8347.

Noirot-Gros, M.F., Velten, M., Yoshimura, M., McGovern, S., Morimoto, T., Ehrlich, S.D.,
Ogasawara, N., Polard, P., and Noirot, P. (2006). Functional dissection of YabA, a negative regulator
of DNA replication initiation in Bacillus subtilis. Proc Natl Acad Sci U S A *103*, 2368-2373.

Ozaki, S., and Katayama, T. (2009). DnaA structure, function, and dynamics in the initiation at the
chromosomal origin. Plasmid *62*, 71-82.

Quevillon-Cheruel, S., Campo, N., Mirouze, N., Mortier-Barriere, I., Brooks, M.A., Boudes, M.,
Durand, D., Soulet, A.L., Lisboa, J., Noirot, P.*, et al.* (2012). Structure-function analysis of
pneumococcal DprA protein reveals that dimerization is crucial for loading RecA recombinase onto
DNA during transformation. Proc Natl Acad Sci U S A *109*, E2466-2475.

Riber, L., Frimodt-Moller, J., Charbon, G., and Lobner-Olesen, A. (2016). Multiple DNA Binding
Proteins Contribute to Timing of Chromosome Replication in E. coli. Frontiers in molecular
biosciences *3*, 29.

Rowland, S.L., Burkholder, W.F., Cunningham, K.A., Maciejewski, M.W., Grossman, A.D., and
King, G.F. (2004). Structure and mechanism of action of Sda, an inhibitor of the histidine kinases
that regulate initiation of sporulation in Bacillus subtilis. Molecular cell *13*, 689-701.

Saghatelian, A., and Couso, J.P. (2015). Discovery and characterization of smORF-encoded bioactive
polypeptides. Nat Chem Biol *11*, 909-916.

Samayoa, J., Yildiz, F.H., and Karplus, K. (2011). Identification of prokaryotic small proteins using a
comparative genomic approach. Bioinformatics *27*, 1765-1771.

Schaeffer, P., Millet, J., and Aubert, J.P. (1965). Catabolic repression of bacterial sporulation. Proc
Natl Acad Sci U S A *54*, 704-711.

Schmalisch, M., Maiques, E., Nikolov, L., Camp, A.H., Chevreux, B., Muffler, A., Rodriguez, S.,
Perkins, J., and Losick, R. (2010). Small genes under sporulation control in the Bacillus subtilis
genome. J Bacteriol *192*, 5402-5412.

Scholefield, G., and Murray, H. (2013). YabA and DnaD inhibit helix assembly of the DNA
replication initiation protein DnaA. Mol Microbiol *90*, 147-159.

Setlow, P. (2007). I will survive: DNA protection in bacterial spores. Trends in microbiology *15*,
172-180.

Skarstad, K., and Katayama, T. (2013). Regulating DNA replication in bacteria. Cold Spring Harbor
perspectives in biology *5*, a012922.

Slavoff, S.A., Heo, J., Budnik, B.A., Hanakahi, L.A., and Saghatelian, A. (2014). A human short
open reading frame (sORF)-encoded polypeptide that stimulates DNA end joining. J Biol Chem *289*,
10950-10957.

Slavoff, S.A., Mitchell, A.J., Schwaid, A.G., Cabili, M.N., Ma, J., Levin, J.Z., Karger, A.D., Budnik,
B.A., Rinn, J.L., and Saghatelian, A. (2013). Peptidomic discovery of short open reading frame-
encoded peptides in human cells. Nat Chem Biol *9*, 59-64.

Soufo, C.D., Soufo, H.J., Noirot-Gros, M.F., Steindorf, A., Noirot, P., and Graumann, P.L. (2008).
Cell-cycle-dependent spatial sequestration of the DnaA replication initiator protein in Bacillus
subtilis. Dev Cell *15*, 935-941.

Staudt, A.C., and Wenkel, S. (2011). Regulation of protein function by 'microProteins'. EMBO
reports *12*, 35-42.

Sterlini, J.M., and Mandelstam, J. (1969). Commitment to sporulation in Bacillus subtilis and its
relationship to development of actinomycin resistance. The Biochemical journal *113*, 29-37.

Storz, G., Wolf, Y.I., and Ramamurthi, K.S. (2014). Small proteins can no longer be ignored. Annu
Rev Biochem *83*, 753-777.

Stragier, P., Bonamy, C., and Karmazyn-Campelli, C. (1988). Processing of a sporulation sigma
factor in Bacillus subtilis: how morphological structure could control gene expression. Cell *52*, 697-
704.

Straub, D., and Wenkel, S. (2017). Cross-Species Genome-Wide Identification of Evolutionary
Conserved MicroProteins. Genome biology and evolution *9*, 777-789.

Tanaka, K., Henry, C.S., Zinner, J.F., Jolivet, E., Cohoon, M.P., Xia, F., Bidnenko, V., Ehrlich, S.D.,
Stevens, R.L., and Noirot, P. (2013). Building the repertoire of dispensable chromosome regions in
Bacillus subtilis entails major refinement of cognate large-scale metabolic model. Nucleic Acids Res
41, 687-699.

VanOrsdel, C.E., Kelly, J.P., Burke, B.N., Lein, C.D., Oufiero, C.E., Sanchez, J.F., Wimmers, L.E.,
Hearn, D.J., Abuikhdair, F.J., Barnhart, K.R.*, et al.* (2018). Identifying New Small Proteins in
Escherichia coli. Proteomics *18*, e1700064.

Veening, J.W., Murray, H., and Errington, J. (2009). A mechanism for cell cycle regulation of
sporulation initiation in Bacillus subtilis. Genes & development *23*, 1959-1970.

Westers, H., Dorenbos, R., van Dijl, J.M., Kabel, J., Flanagan, T., Devine, K.M., Jude, F., Seror, S.J.,
Beekman, A.C., Darmon, E.*, et al.* (2003). Genome engineering reveals large dispensable regions in
Bacillus subtilis. Mol Biol Evol *20*, 2076-2090.

Wu, L.J., and Errington, J. (2011). Nucleoid occlusion and bacterial cell division. Nat Rev Microbiol
10, 8-12.

Wu, W., and Jin, S. (2005). PtrB of Pseudomonas aeruginosa suppresses the type III secretion system
under the stress of DNA damage. J Bacteriol *187*, 6058-6068.

Yang, X., Jensen, S.I., Wulff, T., Harrison, S.J., and Long, K.S. (2016). Identification and validation
of novel small proteins in Pseudomonas putida. Environmental microbiology reports *8*, 966-974.

Zanet, J., Benrabah, E., Li, T., Pelissier-Monier, A., Chanut-Delalande, H., Ronsin, B., Bellen, H.J.,
Payre, F., and Plaza, S. (2015). Pri sORF peptides induce selective proteasome-mediated protein
processing. Science *349*, 1356-1358.

Zuber, P. (2001). A peptide profile of the Bacillus subtilis genome. Peptides *22*, 1555-1577.

10   Legends to figures

Figure 1

A) SknR transcriptionally repress the yqaF-yqaN operon of the skin element. **B-C) Yeast two hybrid**
interaction assay. Haploid yeast strains expressing yqaH, dnaA and yabA infusion with the BD and
AD domains of gal4 are separately introduced into haploid yeast strains. Binary interactions are
tested by the ability of diploids to grow onto selective media. **D) DnaA interaction with YqaH and**
regulators. Numbers refer to amino acid boundaries (see also Figure S1**).** Schematic representation
of the architecture of the four functional domains of DnaA with associated functions as illustrated by
colors. The binding of negative regulators to their targeted DnaA functional domain is illustrated
accordingly. The YqaH interacting domains of DnaA with associated yeast 2HB interacting
phenotypes (IP) are indicated. (+ and -) refers growth or absence of growth on selective media,
reflecting interacting or loss of interaction phenotype, respectively.

Figure 2:

YqaH triggers *dnaA*-related phenotypes. A) Effect of *yqaH* expression on growth. Cells
carrying plasmids pDG148 (control, plain lines) or pDG148-yqaH (dashed lines) were examined in
the presence of IPTG (0.5 mM) over 5 hours. Growth was monitored either by OD600 (left x-axe,
black) or by cells viability, measured as the number of colony forming units per ml, normalized by
OD600 (right x-axe, red). **B) Nucleoid morphological defects**. Samples of living cells (OD or
time?) were examined by fluorescent microscopy after staining with of FM4-64 (membrane dye
false-colored in red) and DAPI (DNA dye, false-colored in blue). White and yellow arrows indicate
guillotined chromosomes resulting from septal closing over nucleoids and aberrant nucleoid
segregation patterns, respectively and typical example of chromosomal segregation defects is
magnified. Scale bares are 5 μm. **C) YqaH affects dnaA expression**. Cells harboring either the
pDG148 or pDG148-yqaH were grown in LB in the presence of IPTG. RNAs from exponentially
grown cells (OD600~0.3) were extracted and expression levels of the dnaA gene were monitored by
qPCR in the presence (+) or absence (-) of YqaH (*** P<0.001).

Figure 3:

YqaH triggers sporulation and Spo0A-related phenotypes. A-B) YqaH expression inhibits
sporulation. Cells carrying plasmids pDG148 (control) or pDG148-*yqaH* were grown in DSM
media in the presence of IPTG (0.5 mM) added at the onset (t$_0$) of sporulation. Sporulant cells were
observed at different time after initiation of sporulation. **A)** Snapshot captures of light and
fluorescence microscopy at indicated t$_{hrs}$ time. Cells were stained with fluorescent membrane dye
FM4-64 (left panel). **B)** Sporulating cells were quantified by monitoring asymmetric septa, engulfed
forespore and free spores, in the presence (+) or absence (-) of YqaH. Ratio were determined from
observation of > 500 cells over 2 independent experiments and 3 biological replicates per
experiment. **C) YqaH affects expression Spo0A-regulated genes**. Cells harboring either the

pDG148 or pDG148-yqaH were grown in DSM. IPTG was. Expression levels of *spoIIE* and *spoIIGA*
genes from the Spo0A regulon were monitored by real-time qPCR in the presence (+) or absence (-)
of YqaH and expressed as relative expression ratio compared to control (-) (*** P <0.001, n≥6).
738

Figure 4:

Restoration of replication initiation defects by YqaH_DnaA LOI mutation. Exponentially
growing *B. subtilis* cells carrying either pDG148 (-), pDG148-*yqaH* (WT), pDG148-*yqaH* LOI
mutant derivatives K17E or R56W were grown in the presence of IPTG and harvested at similar
OD600~0.3 and assessed for various *dnaA*-related phenotypes: **(A) Nucleoid segregation and**
morphology. Cells were treated with DAPI to reveal nucleoids (false-colored blue) and with the
membrane dye FM4-64 (false colored red). **B-C-D) Analysis of DnaA-dependent replication**
initiation phenotypes. B) Ori:ter ratio; Origin-proximal and Terminus proximal DNA sequences
were quantified by qPCR. **C)** Visualization of origins foci in living cells. Origins are tagged through
binding of the GFP-lacI repressor to LacO operator sequences inserted at proximal location from
OriC. **D**) Averaged number of replication origins per cell determined as the number of LacI-GFP foci
upon induced condition in control (-), WT and K17E- mutant derivative of YqaH. Statistical
significance are illustrated by stars (t-test; . Ori:ter: n=6; *dnaA* mRNA: n=12; P< 0.01 *; P<0.001**).
E) **Effect of YqaH LOI mutant on dnaA expression**. Cells harboring either the pDG148, pDG148-
yqaHWT or K17E and R56W mutated derivatives were grown in LB in the presence of IPTG. RNAs
from exponentially grown cells (OD600~0.3) were extracted and expression levels of the dnaA gene
were monitored by qPCR in the presence (+) or absence (-) of YqaH (*** P<0.001).

Figure 5:

Role of K17E DnaA_LOI substitution in sporulation and biofilm formation. A) 168 and Tf8
strains carrying plasmids pDG148 (control) or pDG148-*yqaH* and K17E mutant were grown in DSM
media in the presence of IPTG (0.5 mM) added at the onset (t$_0$) of sporulation. Sporulant cells were
observed - hours after initiation of sporulation (as described in figure 3). **B**) Surface rendering of
biofilms pellicles of 168 strain expressing yqaH or yqaH-K17E mutant after 48h in Mgss media.
Biomass was averaged from 8 samples. Pairwise comparisons were performed using the Tukey
Method (*p <= 0.05 **p <= 0.01 ***p <= 0.001).

Table 1:

YqaH_LOI mutational screen. Aminoacid substitution affecting interaction with DnaA and/or
DnaN are indicated with their associated interaction phenotype. (-) refers as a total loss of interaction
phnotype leading to absence of growth on both –LUH and –LUA media. (+/-) refers as a partial loss
of interaction leading to some growth on the –LUH but not –LUA; see also Figure S1 for additional
explanation.

Figure 1

Figure 2

Figure 3

788   **Figure 5**

791 **Table 1**

		DnaA	Spo0A
YqaH_LOI	D10G	-	-
	R16G	+/-	+
	K17E	-	+
	S20L	-	-
	S25T	+/-	+/-
	M27T	+/-	+
	L28P	-	-
	Y37H	+/-	+
	E38K	-	+
	E38V	+/-	+
	A40T	+/-	+/-
	L43P	-	-
	A44V	+/-	+
	K48E	-	+
	R56W	-	+
	L57P	-	-