Repeated blast mild traumatic brain injury and oxycodone self-administration produce

interactive effects on neuroimaging outcomes

Matthew J. Muelbl[1,3], Breanna Glaeser[1,3], Alok S. Shah[4,5], Rachel Chiariello[4,5], Natalie N.

Nawarawong[1,3], Brian D. Stemper[2,3,4,5], Matthew D. Budde[3,4,5], Christopher M. Olsen[1,3]*

[1]Department of Pharmacology and Toxicology, Medical College of Wisconsin, 8701 Watertown

Plank Rd., Milwaukee, WI 53226, USA; [2]Joint Department of Biomedical Engineering,

Marquette University, 1515 W. Wisconsin Ave, Milwaukee WI, 53233, USA and Medical

College of Wisconsin, 8701 Watertown Plank Rd., Milwaukee, WI 53226, USA; [3]Neuroscience

Research Center, Medical College of Wisconsin, 8701 Watertown Plank Rd., Milwaukee, WI

53226, USA; [4]Department of Neurosurgery, Medical College of Wisconsin, 8701 Watertown

Plank Rd., Milwaukee, WI 53226, USA; [5]Clement J. Zablocki Veterans Affairs Medical Center,

5000 W National Ave, Milwaukee, WI 53295, USA

*Corresponding author:

Christopher M. Olsen, PhD

Medical College of Wisconsin

8701 Watertown Plank Road, Milwaukee, WI 53226, USA

Phone: (414) 955-7629

Email: colsen@mcw.edu

Running title: Neuroimaging of mild TBI and oxycodone

Keywords: opioid, mild traumatic brain injury, extinction, drug seeking, MRI

Abstract

Traumatic brain injury (TBI) and drug addiction are common comorbidities, but it is unknown if the neurological sequelae of TBI contribute to this relationship. We have previously reported elevated oxycodone seeking after drug self-administration in rats that received repeated blast TBI (rbTBI). TBI and exposure to drugs of abuse can each change structural and functional neuroimaging outcomes, but it is unknown if there are interactive effects of injury and drug exposure. To determine the effects of TBI and oxycodone exposure, we subjected rats to rbTBI and oxycodone self-administration and measured drug seeking and several neuroimaging measures. We found interactive effects of rbTBI and oxycodone on fractional anisotropy (FA) in the nucleus accumbens (NAc), and that FA in the medial prefrontal cortex (mPFC) was correlated with drug seeking. We also found an interactive effect of injury and drug on widespread functional connectivity and regional homogeneity of the BOLD response, and that interhemispheric functional connectivity in the infralimbic medial prefrontal cortex positively correlated with drug seeking. In conclusion, rbTBI and oxycodone self-administration had interactive effects on structural and functional MRI measures, and correlational effects were found between some of these measures and drug seeking. These data support the hypothesis that TBI and opioid exposure produce neuroadaptations that contribute to addiction liability.

Introduction

Traumatic brain injury (TBI) and opioid abuse have each reached epidemic proportions. An estimated 2.4 million Americans suffer a TBI each year [1], and in 2017 there were over 47,000 deaths due to opioid overdose [2]. The incidence of TBI among veterans of the military has been approximated at 30% [3], with blast injury causing the overwhelming majority of TBIs in combat settings. Co-morbidity between non-severe TBI and substance abuse has been found in numerous civilian and military epidemiological studies [4-10], and there is evidence that opioid abuse may be particularly prevalent following TBI [11, 12]. For example, in a study of >40,000 Air Force personnel, those that sustained mild TBI (compared to non-brain injuries) had elevated hazard ratios for several drugs of abuse [5]. Furthermore, the hazard ratio for opioids was higher than all other drugs of abuse in the 18 months following injury (including alcohol) [5]. Despite extensive evidence of increased substance abuse disorder following TBI and high rates of opioid prescriptions to veterans and civilians that sustain TBI, little is known about how neurological damage associated with TBI may exacerbate opioid addiction liability.

We have previously demonstrated that our blast model of mild TBI leads to enduring microstructural changes in the medial prefrontal cortex [13], a brain region critically involved in drug seeking in humans and rodent models (reviewed in [14-16]). We have also found that repeated blast mild TBI (rbTBI) persistently elevated levels of oxycodone seeking, despite similar levels of prior drug self-administration [17]. In the current study, we hypothesized that rbTBI and oxycodone self-administration would lead to persistent changes in structure and/or function of the medial prefrontal cortex which may also extend to other aspects of mesocorticolimbic circuitry that would be associated with elevated drug seeking. To test this hypothesis, rats underwent rbTBI, oxycodone self-administration, and drug seeking sessions. Structural and functional MRI were performed at acute and chronic timepoints (1- and 7- weeks post-injury, respectively).

Methods

An overview and timeline of experimental procedures is found in Fig 1A.

Animals

Male Sprague Dawley rats (Harlan/Envigo, Madison, WI, USA) arrived at the Zablocki Veterans Affairs Medical Center (ZVAMC) and were housed in pairs. Three days after the final blast or sham exposure, rats were transferred from the ZVAMC to Medical College of Wisconsin (MCW, approximately 5 miles away). Upon arrival at MCW, rats were singly housed for the remainder of the study. Food and water were provided ad libitum throughout the study, except when rats were in the behavioral apparatus. Animals were housed under a reverse light cycle (lights off 0730-1930), and experiments were conducted during the dark period. Experiments were conducted 5 days/week by experimenters blinded to the treatment condition. All experiments were conducted in compliance with the Institutional Animal Care and Use Committee (IACUC) at MCW and ZVAMC (protocol number: AUA 2816).

Repeated blast exposure

Rats were exposed to repeated blast overpressure or repeated sham conditions as described in [13, 17-19]. During blast or sham treatment, rats were maintained under anesthesia with a continuous delivery of 1.5% isoflurane and administered Rimadyl (carprofen, 5 mg/kg, s.c., Zoetis Inc, Kalamazoo, MI, USA). The torso of the rat was shielded with a metal cylinder, earplugs were inserted, and the head was restrained to prevent injury from rotational acceleration [13]. Blast-exposed rats were positioned 17 cm from the end of a custom shock tube (3.6 cm inner diameter, 3.0 m driven section and 0.3 m driver section with Mylar membrane between the driver and

driven sections) and 18 deg off axis from the shock tube to prevent interaction of the head with exhaust gases. Blast overpressure (450 kPa, 80 kPa*ms) was produced by pressurizing the driver section with helium until the membrane ruptured. Sham-exposed rats were subjected to the same procedures but were placed outside the overpressure shockwave. Following exposure, rats were placed on a heating pad and observed until return of righting reflex. Time from initiation of anesthesia prior to blast exposure until return of the righting reflex was quantified as 'recovery time'. Following recovery, rats were transferred back to their home cage, and periodically observed for 6 h. This procedure was conducted daily for 3 days as a repeated blast injury model. Sham-exposed rats received sham procedures daily for 3 days. Experimenters were blind to group assignment during subsequent phases of experiments.

Jugular catheterization surgery

Jugular catheterization was performed as described [17, 20, 21]. Rats were anesthetized with isoflurane (3–5% induction, 1–3% maintenance), and a silicone catheter (Silastic, ID: 0.51 mm, OD: 0.94 mm; Dow Corning, Auburn, MI, USA) was implanted into the right jugular vein and exited via the interscapular region. The catheter was connected to a cannula assembly: a 22-gauge stainless steel cannula (Plastics One, Roanoke, VA, USA) mounted on a base made from dental acrylic and nylon mesh, which was implanted subcutaneously. At the start of the surgery and the following day, Rimadyl (carprofen, 5 mg/kg, s.c. Zoetis Inc) was given. Catheters were flushed daily with 0.2 mL of heparinized saline (30 units/mL) and cefazolin (100 mg/mL). Rats recovered for at least 1 week prior to the start of the drug self-administration experiments.

Food training

Food training and oxycodone self-administration were performed as described in [17]. Food and oxycodone self-administration and the seeking test all took place in operant conditioning chambers (31.8 × 25.4 × 26.7 cm, Med Associates, Fairfax, VT, USA) which were enclosed in sound-attenuating boxes. Chambers were equipped with two retractable levers on either side of a central food receptacle on the right wall, cue lights above each lever and a house light located on the left wall near the ceiling. Computer-controlled infusion pumps held syringes that were connected to tubing and liquid swivels. Sound attenuating boxes had a software-controlled exhaust fan.

Rats began by learning to self-administer food (fruit punch flavored 45 mg sucrose tablets, LabDiet, St. Louis, MO, USA) without food restriction. Each session lasted for 2 h/day unless the maximum number of rewards (64 sucrose pellets) was met prior to the conclusion of the 2 h.

At the beginning of each session, the fan was turned on, both levers were extended, and a sucrose

pellet was delivered. Rats were first trained on a fixed ratio (FR)-1 schedule of reinforcement,

with a single press on the active lever resulting in sucrose pellet delivery. After meeting criteria

(three consecutive days with $\geq$ 50 rewards earned and $\geq$ 70% of total lever presses on the active

lever, 4 day minimum), rats performed a single food self-administration session on an FR-2

schedule of reinforcement (2 active lever presses required per reward).

The active lever was assigned to be on either the left or right side, and lever assignment was

counterbalanced between rats within each group. After reward delivery, a 10 s timeout began,

and the cue light above the active lever was illuminated for 5 s. During the timeout period, active

and inactive lever responses were recorded but had no programmed consequence. Throughout

the entire session, responses on the inactive lever were recorded but also had no programmed

consequence.

Oxycodone self-administration and seeking

After completion of food training, rats in the rbTBI and sham groups were further divided into

those that self-administered oxycodone or received yoked infusions of saline. Each yoked saline

rat was matched to a "master" rat from the same cohort. When the "master" rat earned an

infusion of oxycodone, the yoked rat received an infusion of saline at the exact same time

(computer controlled). Thus, the infusion history between the "master" and "yoke" rats is

identical, controlling for the potential influence of infusions. Yoked saline rats still received cue

light presentation in response to completing the response requirements. The infusion volume and

time varied for each rat to ensure the appropriate dose was delivered from the stock solution

(e.g., a 300 g rat received a 40 μL infusion over 1.6 s) and yoked saline rats received weight-

corrected volumes of saline. This phase consisted of 10 daily 2-h sessions (maximum 96

infusions/session): 5 d each on FR-2 and FR-4 schedules of reinforcement. Catheter patency was

determined after completion of the self-administration sessions using Brevital (methohexital, 9

mg/kg, i.v.; JHP Pharmaceuticals, Rochester, MI, USA) and any rat that did not meet criteria for

patency (sedation within 10 s) was removed from the study. Three days after completion of the

10-day oxycodone self-administration (or yoked saline) phase, all rats were placed back into the

operant conditioning chambers for a drug seeking test under extinction conditions. Completion

of the FR-4 schedule resulted in saline infusion and cue delivery for all animals (i.e., yoked

saline animals were no longer yoked).

Magnetic Resonance Imaging (MRI)

In vivo magnetic resonance imaging was performed on a Bruker 9.4 T Biospec System. A Bruker Biospin 4-channel surface coil array was used for signal reception and transmission was achieved via a 72 mm diameter quadrature volume coil. Heart rate (ECG), core temperature, and respiration were monitored continuously. Anesthesia was induced with isoflurane (3.0%), which was tapered to 0% after initiation of constant tail vain infusion of dexmedetomidine. Dexmedetomidine was delivered via constant tail vain infusion (initial concentration: 50 µg/kg/hr) and was titrated to maintain anesthetic depth as determined by ECG, body temperature, and respiratory rate. Anesthetized rats were secured to a custom-made temperature-controlled cradle to minimize motion. A high-resolution anatomical image was acquired with an isotropic resolution of 134 µm^3. For diffusion imaging, a 4-shot spin-echo echo planar imaging (DW-EPI) sequence was used as described previously [22] with TR=2000 and TE=23 ms, including fat saturation using both chemical shift-selective saturation and gradient reversal of the refocusing pulse slice. Thirty slices covered the entire brain at an in-plane resolution of 0.30×0.30 mm^2, slice thickness of 0.8 mm and 0.3 mm slice gap. Two signal averages were used to acquire 60 unique diffusion directions [23] at each b-value of 250, 1000, and 2000 s/mm^2 with 8 non-diffusion-weighted images. For resting state functional MRI, 200 images were acquired at a temporal resolution of 3 sec using single-shot gradient echo EPI at the identical spatial resolution. Three images were acquired for each repetition at echo times of 16.5, 33.1, and 49.6 ms, and each run was repeated 3 times for a total of 30 minutes for fMRI data acquisition. To examine for any evidence of overt injury or hemorrhage, a 3D multiple echo acquisition with 0.175 mm^3 isotropic voxels (TR=50 ms) was acquired with 10 echo times between 2.2 and 24.9 ms. Across all animals, no hemorrhage was evident based on visual inspection.

Statistical Analysis

Statistical tests were performed using Prism 8 (GraphPad, San Diego, CA, USA) or SPSS Statistics 24 (IBM, Inc.) software. Data were analyzed using two-way mixed model ANOVAs (session: within-subjects factors, experimental group: between-group factor) with Holm-Sidak post hoc tests. ANOVAs were performed using ranked data if data did not fit a normal distribution as determined by D'Agonstino & Pearson tests. Correlational analyses were performed using linear regressions (Spearman rank correlations if data did not fit a normal distribution) followed F tests to determine if slopes were significantly different from zero. Differences were considered significant when $p \leq 0.05$.

Results

Behavioral measures

Repeated blast TBI resulted in shorter recovery time (36.6 ± 8.1 sec faster) from anesthesia

compared to sham treated rats ($F_{(1, 38)}=20.26$, $\eta^2=0.20$ $p<0.0001$), but there was no effect of

blast exposure number or interaction of blast with exposure number (both $\eta^2<0.10$, $p>0.39$). On

week three following rbTBI or sham treatment, rats underwent food self-administration training.

Food self-administration did not differ between rats exposed to rbTBI and sham (main and

interaction effects of rbTBI all $\eta^2<0.045$, $p>0.39$; Fig 1B). These data indicate that there was no

impact of injury on the ability to acquire and perform this task, a critical consideration in TBI

research that utilizes operant conditioning [24]. Next, rats either self-administered oxycodone or

received yoked infusions of saline. Specifically, rats were split into pairs: when the "master" rat

earned an infusion of oxycodone, this triggered a time-matched infusion of saline into the

"yoked" rat. There was no significant difference between rbTBI and sham rats in oxycodone

self-administration as measured by active lever presses during each FR phase (main and

interaction effects of rbTBI all $\eta^2<0.065$, $p>0.21$; Fig 2A) or number of infusions earned during

the FR-2 phase (main and interaction effects of rbTBI both $\eta^2<0.045$, $p>0.26$; Fig 2C) or the FR-

4 phase (main and interaction effects of rbTBI all $\eta^2<0.083$, $p>0.16$; Fig 2D). There was also no

significant difference in lever pressing between rbTBI and sham rats in the yoked saline groups

(main and interaction effects of rbTBI both $\eta^2<0.016$, $p>0.67$; Fig 2B). This demonstrates that

rbTBI does not lead to generalized deficits in extinction learning (i.e., extinction of sucrose pellet

seeking); a significant finding considering that we have previously reported that rbTBI resulted

in elevated oxycodone seeking in extinction conditions [17] and several studies have found that

mild TBI impaired extinction of conditioned fear [25-28]. Despite the lack of significant difference

in oxycodone self-administration, there was a trend for elevated drug seeking by rbTBI rats

relative to sham rats ($t_{(20)}=1.5$, $d=0.64$, $p=0.15$). This trend was similar in effect size to our

previous report on the first day of oxycodone seeking using the same injury and oxycodone self-

administration model ($d=0.80$) [17].

A

Week	Experiment Phase
0	Repeated Blast Injury
1	Handling
2	Jugular Catheterization Surgery
3	Food Self-Administration
4	Oxycodone Self-Administration (FR-2)
5	Oxycodone Self-Administration (FR-4)
6	Oxycodone Seeking
7	MRI

B Food Self-Administration

Figure 2: Oxycodone *self-administration* and seeking. A) Active lever responses for oxycodone. B) Active lever responses in rats that received yoked saline infusions. C, D) Oxycodone infusions earned during FR-2 (C) and FR-4 (D) sessions. E) Active presses during extinction trial. Boxes represent median and quartiles, whiskers represent range. Bars represent mean $\pm$ SEM.

MRI

To examine neuroimaging correlates of injury and oxycodone self-administration, rats underwent structural and functional MRI at 7-weeks post-injury, approximately 10 days following cessation of oxycodone self-administration.

<u>Structural MRI</u>

Structural assessment was performed by examining T2, T2*3D, fractional anisotropy (FA), and

mean diffusivity (MD).

Factorial analysis of fractional anisotropy and mean diffusivity

We examined main and interactive effects of rbTBI and oxycodone self-administration on

microstructure of the prefrontal cortex and nucleus accumbens. These brain regions are engaged

during drug craving in human opioid users and drug seeking behaviors in rodent models of

opioid addiction [15, 29, 30]. These areas are also subject to damage in human blast and non-blast

mild TBI [31-34]. There were no significant main effects of injury or oxycodone, but a significant

interaction was found in the FA of both hemispheres of the nucleus accumbens ($p \leq 0.05$).

Relationship between fractional anisotropy and drug seeking

We have previously demonstrated that bTBI lead to changes in hippocampal FA that correlated

with spatial learning deficits [13]. To examine the possibility that mPFC FA was associated with

drug seeking, we performed correlational analyses between FA in the subregions of the

prefrontal cortex (PFC) and nucleus accumbens (NAc) and active lever presses during the drug

seeking session. We examined sham and rbTBI rats separately and together to determine if

associations were specific to injured rats or if associations generalized to both populations.

Significant associations were only found in the subregions of the mPFC ipsilateral to injury (Fig

3). In the prelimbic PFC (PrL), there were significant linear correlations in the rbTBI group

(r^2=41, p=0.047) and across injury groups (r^2=23, p=0.020), but not in sham injured rats (r^2=11,

p=0.37). In the infralimbic PFC (IL), there was only a significant association between FA and

drug seeking when the sham and injury groups were combined (r^2=22, p=0.046). To determine if

this association was explained by the amount of oxycodone intake during prior self-

administration sessions, we examined the correlations between the PFC/NAc and drug intake

(Table 1). There was no association between either ipsilateral subregion and drug intake

(Pearson's r and Spearman's rho all <0.48), however, there was a positive linear association

between the contralateral IL and prior oxycodone intake in rbTBI (rho=0.65, p=0.049) and in

combined sham and injury rats (rho=0.59, p=0.0084). These data suggest that the association

between reduced FA ipsilateral to injury and elevated drug seeking is not due to the amount of

oxycodone self-administered, and that there is a positive relationship between total oxycodone

intake and FA in the contralateral IL mPFC.

281 *Figure 3: Correlations between FA and active lever presses in ipsilateral PrL and IL. *p≤0.05*

285 *Table 1: Correlations between structural neuroimaging measures and seeking or intake. FA: fractional anisotropy, MD: mean*
286 *diffusivity. Ipsilateral: left, contralateral: right. Italicized values represent Spearman's rho instead of Pearson's r. *p≤0.05*
287 ***p≤0.01.*

FA v. Seeking		sham	rbTBI	sham+rbTBI
Ipsilateral	PrL	-0.34	**-0.64***	**-0.53***
	IL	-0.46	-0.46	**-0.46***
	NAc	*0.30*	*-0.42*	*-0.19*
Contralateral	PrL	0.10	-0.03	-0.03
	IL	0.35	0.30	0.30
	NAc	-0.26	-0.37	-0.26

MD v. Seeking		sham	rbTBI	sham+rbTBI
Ipsilateral	PrL	*0.42*	*-0.05*	*0.12*
	IL	*0.25*	*-0.12*	*0.02*
	NAc	*0.10*	*-0.12*	*-0.01*
Contralateral	PrL	*0.03*	*-0.07*	*0.11*
	IL	*-0.10*	*-0.36*	*-0.16*
	NAc	0.26	0.01	*0.05*

FA v. Intake		sham	rbTBI	sham+rbTBI
Ipsilateral	PrL	-0.03	*-0.32*	*-0.31*
	IL	-0.48	*-0.36*	*-0.36*
	NAc	*0.48*	*0.04*	*0.03*
Contralateral	PrL	0.15	*0.05*	*0.16*
	IL	0.39	*0.65***	*0.59****
	NAc	0.11	*0.07*	*0.01*

MD v. Intake		sham	rbTBI	sham+rbTBI
Ipsilateral	PrL	*-0.03*	*-0.30*	*-0.19*
	IL	*-0.10*	***-0.67****	*-0.36*
	NAc	*-0.27*	*-0.40*	*-0.31*
Contralateral	PrL	*-0.32*	*-0.29*	*-0.17*
	IL	*-0.15*	***-0.67****	***-0.55****
	NAc	*-0.15*	*-0.52*	*-0.30*

<u>Relationship between mean diffusivity and drug seeking</u>

We examined mean diffusivity (MD) to determine if this additional measure of local

microstructure was associated with oxycodone seeking or intake during self-administration

(Table 1). No significant associations were found between MD and drug seeking in any region,

although the MD in both hemispheres of the IL mPFC had a significant negative association with

oxycodone intake in rbTBI rats (rho=-0.67, p=0.039 for each hemisphere). Similarly, there was a

significant negative association between MD and oxycodone intake when injury groups were

combined (rho=-0.55, p=0.014).

Functional MRI

Regional homogeneity (REHO)

Figure 4: Regional homogeneity (REHO) of BOLD signal. Red/yellow voxels represent main or interaction effects at p≤0.05.

Global functional connectivity

Regions of interest were represented as nodes and edges were calculated as the temporal Pearson

correlation coefficient in the BOLD response between pairwise nodes for each of the four

treatment groups (Fig 7). rbTBI/saline and sham/oxycodone self-administration rats showed

comparable patterns of connectivity to sham/saline rats, while rbTBI/oxycodone self-

administration rats had a larger number of connected nodes, as defined by Pearson r≥0.6 (Fig 7).

Edge strengths were tested in a factorial design, which also yielded a significant interaction

(p≤0.05), but no significant main effects (both p>0.62).

Figure 5: Network-based statistics. A) Each node (blue sphere) represents a region of interest, each edge represents a significant
correlation between nodes. B) Nodes connected by edges had a significant interaction between rbTBI and oxycodone self-
administration. Edges are colored based on Pearson r values (A) or F values (B).

<u>Relationship between functional connectivity and drug seeking</u>

Prefrontal-accumbens neurotransmission has been implicated in drug seeking for several drugs

of abuse, including opioids (reviewed in [15, 35, 36]). We tested the hypothesis that functional

connectivity between regions of the PFC and nucleus accumbens would be positively correlated

with oxycodone seeking, and this would be relationship would be independent of injury.

Pairwise correlations were performed on each hemisphere of the PrL, IL, and NAc. Analysis of

the relationship between functional connectivity and drug seeking (active lever presses during

extinction session) found that only the inter-hemispheric connectivity of the IL was significantly

correlated with drug seeking in injured and uninjured rats (Fig 6C). Prefrontal-accumbens

functional connectivity was significantly associated with drug intake on the left hemisphere

(ipsilateral to injury in rbTBI rats), and this was observed in rbTBI and the combined population

of rats (Fig 6E,F).

Figure 6: Relationship between functional connectivity and drug seeking (A-C) or drug intake (D-F). Values are Pearson r except where denoted by ρ. Significant correlations are identified in bold text. *p≤0.05. I, C: ispi- or contra-lateral to injury, corresponding to left or right hemisphere, respectively in all rats.

Discussion

We have previously shown that rbTBI-exposed rats had elevated drug seeking following oxycodone self-administration despite similar levels of drug intake [17]. In this study, we used neuroimaging to explored potential interactive effects and correlates of behavior on structural and functional brain networks. We focused on the prefrontal cortex and nucleus accumbens due to their known involvement in persistent drug seeking behavior [15, 37, 38]. We found several neuroimaging metrics that were associated with drug seeking behavior, prior drug intake, and interactive effects of injury and oxycodone self-administration. Microstructural analysis identified significant negative associations between FA and drug seeking in the left (but not right) PrL and IL PFC, the side ipsilateral to injury in rbTBI rats. There was also a significant positive association between FA and drug intake during self-administration in the right hemisphere, while the association between MD and prior oxycodone intake was bilateral. The effect was largest in rbTBI rats but was also significant in the combined population of injured and un-injured rats. Within the NAc, there was a significant interaction between injury and oxycodone self-administration on FA, but neither FA nor MD were associated with the amount of oxycodone self-administered or subsequent drug seeking. All of the microstructural correlations with behavior had larger effects in injured relative to uninjured rats, suggesting that

rbTBI enabled or enhanced the emergence of a relationship between structural parameters of the PFC and prior drug seeking and/or drug intake. Resting state functional connectivity (rsfc) analyses found a widespread increase in network connectivity only in rats that received injury and oxycodone SA. There were also significant correlations between IL intra-hemispheric connectivity and drug seeking, and between ipsilateral IL – NAc connectivity and oxycodone intake during self-administration. As observed in the structural measures, these effects were largest in injured animals, suggesting that similar to our structural outcomes, the interaction of injury and oxycodone exposure enabled the emergence of a relationship between functional connectivity and prior drug seeking and/or drug intake. An interaction was also measured in local connectivity: regional homogeneity was minimally affected by injury or oxycodone self-administration alone, but animals receiving both had marked increases in voxelwise homogeneity of the medial PFC, NAc, and other areas, such as motor cortex, hippocampus, deep layers of the superior colliculus, and external cortex of the inferior colliculus.

The present study adds to a growing body of preclinical work that has largely demonstrated that experimental TBI is associated with worse outcomes in models of substance abuse. For example, studies across different species and TBI models have found elevated alcohol intake following injury [18, 39-42] (although see [43]). Experimental TBI has also been shown to increase cocaine conditioned reward [44, 45] and self-administration [46] (although see [20]). In the same model of repeated blast mild TBI used here, we found that injured animals had reduced oxycodone self-administration, but drug seeking was elevated during extinction testing [17]. Thus, consistent with clinical studies that report associations between TBI and substance abuse [4-12], there is a growing body of preclinical work that supports a causative role for TBI in elevated addiction liability. In the current study, we found that drug seeking was elevated after injury and oxycodone SA. We also found structural and functional changes in the brain, with some of these alterations associated with prior drug seeking or intake.

The significance of injury and opioid exposure has been examined in peripheral and central injury models. Mu opioid receptor agonists administered after injury lead to longer recovery times in models of spinal cord injury and neuropathic pain ([47-50]), while kappa opioid receptor agonists may be protective [51]. In regard to TBI, the impact of opioid exposure on injury outcomes is less clear. Fentanyl administered immediately following controlled cortical impact exacerbated motor and spatial learning deficits within three weeks of injury [52]. In a comparison of multiple anesthetic agents delivered immediately after controlled cortical impact, morphine was associated with the worst outcomes on motor performance in the first five days after injury,

while fentanyl was associated with the worst performance in the Morris water maze two weeks

following injury [53]. Conversely, there is also evidence that opioids may be protective in

experimental TBI. Morphine given immediately after injury was found to be beneficial in weight

drop TBI-associated deficits in spatial learning ability 30-90 days post-injury [54]. The mu opioid

receptor selective agonist DAMGO delivered i.c.v. immediately prior to injury protected against

beam walk and beam balance deficits in a rat fluid percussion injury model, and these outcomes

were worsened by the mu antagonist beta-funaltrexamine [55]. In the present study, we observed

several interactive effects between rbTBI and oxycodone self-administration consistent with

worse outcomes. In addition to injury exacerbating oxycodone seeking, injury x oxycodone self-

administration interactions on FA were found in the NAc. Functional measures were more

sensitive to interaction effects, with significant effects identified in the mPFC, NAc, and several

other regions in local connectivity (REHO) and widespread functional connectivity.

Furthermore, functional connectivity between the left and right IL mPFC positively correlated

with both oxycodone seeking and intake in the combined analysis of sham and injured rats, and

the left IL-NAc (ipsilateral to injury) connectivity positively correlated with drug intake in

analysis of rbTBI rats alone and when combined with shams. These measurements we taken

following measures of drug seeking – future studies will need to be done in order to determine if

these metrics are predictive of future drug seeking.

Neuroimaging studies of TBI

Diffusion tensor imaging (DTI) is a sensitive measure to identify microstructural damage

associated with TBI (reviewed in [56-58]). Experimental blast TBI has been shown to lead to

widespread decreases in FA[13], with closely-spaced repeated blast exposures exacerbating these

effects [59]. DTI measurements also tend to be exacerbate with time post-injury: we have reported

an increase in the volume of microstructural damage between post-injury day 4 and 30 [13], and

Badea et al. have reported greater changes in DTI at 90 days relative to 7 days post-injury [60].

Another study found region-specific changes in DTI between 1 and 14 days after injury,

reflecting regional differences in time to vasogenic or cytotoxic edema and demyelination [61].

In the present study, we did not observe a significant effect of rbTBI on FA in the prefrontal

cortex or nucleus accumbens, but we did observe a significant interaction between rbTBI and

oxycodone self-administration in the nucleus accumbens. Despite a main or interactive effect in

the PFC, we did observe correlations between FA and oxycodone seeking that were largely

driven by animals that also experienced rbTBI.

Changes in rsfc have been observed in human TBI and in experimental TBI. Increased intra-hemispheric connectivity of the anterior cingulate cortex was found in veterans and athletes that had experienced a mild TBI [62, 63], an effect we observed in the PrL (immediately ventral of the anterior cingulate cortex in the rat). In collegiate athletes, elevated local connectivity (REHO) in the right middle and superior frontal gyri was associated with clinical symptoms following sports concussion [64]. Widespread disruptions in functional connectivity have also been associated with long-term effects of TBI (reviewed in [65-67]). In a study of high school and collegiate football players, increased widespread rsfc was observed only in athletes that remained symptomatic one week following injury [68]. Similarly, elevated widespread rsfc was found in adolescent hockey players at three months following injury, although this effect was greater in recovered athletes relative to those with fewer symptoms [69]. Combat veterans that sustained blast mild TBI and had persistent post-concussive symptoms had increased rsfc (assessed by magnetoencephalography) in several brain regions, including the ventromedial cortex and anterior cingulate cortices [70]. Similarly, increases in functional connectivity in moderate and severe TBI were found for up to a year after injury, and the changes were greatest in areas that were already highly connected [71]. Conversely, other studies using BOLD MRI have reported lower rsfc and/or local connectivity in injured patients [72, 73]. In a study of TBI across different injury mechanisms, increased local connectivity was observed, and in a subset of patients, areas of highest connectivity were colocalized with high [^{18}F]AV-1451 binding, a marker of tau hyperphosphorylation [74]. The authors speculate that this may be associated with vascular neuropathology. Consistent with many of these studies, we found increased widespread and local connectivity. Similar to our observation of increased inter-hemispheric connectivity within the mPFC, Kulkarni et al. reported increased rsfc within PrL and IL after a single closed head momentum exchange injury [75].

Neuroimaging studies of opioid use

Clinical studies have consistently identified structural and functional changes associated with opioid abuse. A voxel-based meta-analysis of studies employing DTI found that bilateral reductions in FA in frontal subgyral regions, including areas impacting the communication between cingulate and limbic association cortices with prefrontal association cortices [76]. Heroin dependent individuals were also found to have lower FA in the white matter tract connecting the mPFC and postcentral gyrus [77] and a comparison of heroin users that relapsed to those that remained abstinent 6 months after initiation of methadone maintenance therapy found lower FA in several white matter tracts in the relapse group [78]. In a neuroimaging study of over 150 subjects, increased widespread structural connectivity was reported in abstinent heroin users

relative to non-heroin using controls [79]. In this study, white matter FA was increased in heroin

users, and FA within frontal networks was correlated with daily heroin intake [79].

Opioid use and abstinence have also been associated with altered functional neuroimaging

measures in human populations. In a study of former heroin users undergoing methadone

maintenance, those that relapsed had elevated REHO in medial orbital cortex and the right

caudate nucleus, and elevated REHO in the caudate was correlated with relapse rates [80]. In

contrast, methadone maintained former heroin users had lower REHO in the medial orbital

cortex compared to control subjects without a history of heroin or methadone use [81]. In the

present study, we found that rats receiving rbTBI and oxycodone self-administration had

elevated REHO in several prefrontal regions, including medial and lateral orbital cortex. A trace

of this effect was evident in the orbital cortex from blast alone, but we observed no indication of

this in animals exposed to oxycodone only. Differences between our preclinical data and the

aforementioned clinical data may arise from the interactive effects of the oxycodone and injury,

the lack of opioid maintenance therapy in our rodent study, or other experimental differences.

Zhou et al. found increased functional connectivity between the ventromedial PFC and nucleus

accumbens in former heroin users with $>=$ 3 years of abstinence [82], similar to findings of

increased functional connectivity between ventral/rostral anterior cingulate cortex and the

nucleus accumbens [83]. Increased widespread functional connectivity has been also been reported

in heroin use during abstinence [84], although there are also findings of decreased rsfc in opioid

addiction, especially when examining connectivity within an executive network (reviewed in [85,

86]). In conclusion, we find evidence that repeated mild TBI and oxycodone interact in such a way

that manifests as elevated drug seeking, increased microstructural damage, and widespread

aberrant functional connectivity. Repeated blast mild TBI also increased the effect sizes of many

relationships between the neuroimaging findings and prior drug seeking and/or intake.

Acknowledgements

This research was supported by National Institute on Drug Abuse Grant DA039276, United

States Department of Veteran Affairs grant RX002931and the Research and Education Initiative

Fund, a component of the Advancing a Healthier Wisconsin Endowment at the Medical College

of Wisconsin. The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health.

References

1. Taylor, C.A., Bell, J.M., Breiding, M.J. and Xu, L. (2017). Traumatic Brain Injury-Related Emergency Department Visits, Hospitalizations, and Deaths - United States, 2007 and 2013. MMWR Surveillance Summaries 66, 1-16.

2. Scholl, L., Seth, P., Kariisa, M., Wilson, N. and Baldwin, G. (2019). Drug and Opioid-Involved Overdose Deaths — United States, 2013–2017. MMWR Morb Mortal Wkly Rep 67, 1419-1427.

3. Tanielian, T. and Jaycox, L.H. (eds) 2008. Invisible wounds of war: Psychological and cognitive injuries, their consequences, and services to assist recovery. Rand Corportation

4. Fann, J.R., Burington, B., Leonetti, A., Jaffe, K., Katon, W.J. and Thompson, R.S. (2004). Psychiatric illness following traumatic brain injury in an adult health maintenance organization population. Archives of general psychiatry 61, 53-61.

5. Miller, S.C., Baktash, S.H., Webb, T.S., Whitehead, C.R., Maynard, C., Wells, T.S., Otte, C.N. and Gore, R.K. (2013). Risk for addiction-related disorders following mild traumatic brain injury in a large cohort of active-duty U.S. airmen. The American journal of psychiatry 170, 383-390.

6. West, S.L. (2011). Substance use among persons with traumatic brain injury: a review. NeuroRehabilitation 29, 1-8.

7. McKinlay, A., Grace, R., Horwood, J., Fergusson, D. and MacFarlane, M. (2009). Adolescent psychiatric symptoms following preschool childhood mild traumatic brain injury: evidence from a birth cohort. The Journal of head trauma rehabilitation 24, 221-227.

8. Kreutzer, J.S., Wehman, P.H., Harris, J.A., Burns, C.T. and Young, H.F. (1991). Substance abuse and crime patterns among persons with traumatic brain injury referred for supported employment. Brain injury : [BI] 5, 177-187.

9. Corrigan, J.D. (1995). Substance abuse as a mediating factor in outcome from traumatic brain injury. Archives of physical medicine and rehabilitation 76, 302-309.

10. Bjork, J.M. and Grant, S.J. (2009). Does traumatic brain injury increase risk for substance abuse? Journal of neurotrauma 26, 1077-1082.

11. Adams, R.S., Corrigan, J.D. and Dams-O'Connor, K. (2019). Opioid Use among Individuals with Traumatic Brain Injury: A Perfect Storm? Journal of neurotrauma.

12. Corrigan, J.D. and Adams, R.S. (2019). The intersection of lifetime history of traumatic brain injury and the opioid epidemic. Addict Behav 90, 143-145.

13. Budde, M.D., Shah, A., McCrea, M., Cullinan, W.E., Pintar, F.A. and Stemper, B.D. (2013). Primary blast traumatic brain injury in the rat: relating diffusion tensor imaging and behavior. Frontiers in neurology 4, 154.

14. Koob, G.F. and Volkow, N.D. (2016). Neurobiology of addiction: a neurocircuitry analysis. Lancet Psychiatry 3, 760-773.

15. Reiner, D.J., Fredriksson, I., Lofaro, O.M., Bossert, J.M. and Shaham, Y. (2019). Relapse to opioid seeking in rat models: behavior, pharmacology and circuits. Neuropsychopharmacology 44, 465-477.

16. Wilson, S.J., Sayette, M.A. and Fiez, J.A. (2004). Prefrontal responses to drug cues: a neurocognitive analysis. Nat Neurosci 7, 211-214.

17. Nawarawong, N.N., Slaker, M., Muelbl, M., Shah, A.S., Chiariello, R., Nelson, L.D., Budde, M.D., Stemper, B.D. and Olsen, C.M. (2018). Repeated blast model of mild traumatic brain injury alters oxycodone self-administration and drug seeking. The European journal of neuroscience.

18. Lim, Y.W., Meyer, N.P., Shah, A.S., Budde, M.D., Stemper, B.D. and Olsen, C.M. (2015). Voluntary Alcohol Intake following Blast Exposure in a Rat Model of Mild Traumatic Brain Injury. PLoS One 10, e0125130.

19. Stemper, B.D., Shah, A.S., Budde, M.D., Olsen, C.M., Glavaski-Joksimovic, A., Kurpad, S.N.,
McCrea, M. and Pintar, F.A. (2016). Behavioral Outcomes Differ between Rotational
Acceleration and Blast Mechanisms of Mild Traumatic Brain Injury. Frontiers in neurology 7.
20. Muelbl, M.J., Slaker, M.L., Shah, A.S., Nawarawong, N.N., Gerndt, C.H., Budde, M.D.,
Stemper, B.D. and Olsen, C.M. (2018). Effects of Mild Blast Traumatic Brain Injury on Cognitive-
and Addiction-Related Behaviors. Sci Rep 8, 9941.
21. Ikegami, A., Olsen, C.M., Fleming, S.M., Guerra, E.E., Bittner, M.A., Wagner, J. and
Duvauchelle, C.L. (2002). Intravenous ethanol/cocaine self-administration initiates high intake
of intravenous ethanol alone. Pharmacology, biochemistry, and behavior 72, 787-794.
22. Skinner, N.P., Kurpad, S.N., Schmit, B.D., Tugan Muftuler, L. and Budde, M.D. (2017). Rapid
in vivo detection of rat spinal cord injury with double-diffusion-encoded magnetic resonance
spectroscopy. Magnetic resonance in medicine : official journal of the Society of Magnetic
Resonance in Medicine / Society of Magnetic Resonance in Medicine 77, 1639-1649.
23. Basser, P.J., Mattiello, J. and LeBihan, D. (1994). Estimation of the effective self-diffusion
tensor from the NMR spin echo. J Magn Reson B 103, 247-254.
24. Vonder Haar, C. and Winstanley, C.A. (2016). Minor Functional Deficits in Basic Response
Patterns for Reinforcement after Frontal Traumatic Brain Injury in Rats. Journal of neurotrauma
33, 1892-1900.
25. Schneider, B.L., Ghoddoussi, F., Charlton, J.L., Kohler, R.J., Galloway, M.P., Perrine, S.A. and
Conti, A.C. (2016). Increased Cortical Gamma-Aminobutyric Acid Precedes Incomplete
Extinction of Conditioned Fear and Increased Hippocampal Excitatory Tone in a Mouse Model
of Mild Traumatic Brain Injury. Journal of neurotrauma 33, 1614-1624.
26. Zhao, J., Huynh, J., Hylin, M.J., O'Malley, J.J., Perez, A., Moore, A.N. and Dash, P.K. (2018).
Mild Traumatic Brain Injury Reduces Spine Density of Projection Neurons in the Medial
Prefrontal Cortex and Impairs Extinction of Contextual Fear Memory. Journal of neurotrauma
35, 149-156.
27. Cheng, W.H., Martens, K.M., Bashir, A., Cheung, H., Stukas, S., Gibbs, E., Namjoshi, D.R.,
Button, E.B., Wilkinson, A., Barron, C.J., Cashman, N.R., Cripton, P.A. and Wellington, C.L.
(2019). CHIMERA repetitive mild traumatic brain injury induces chronic behavioural and
neuropathological phenotypes in wild-type and APP/PS1 mice. Alzheimer's research & therapy
11, 6.
28. Davies, D.R., Olson, D., Meyer, D.L., Scholl, J.L., Watt, M.J., Manzerra, P., Renner, K.J. and
Forster, G.L. (2016). Mild Traumatic Brain Injury with Social Defeat Stress Alters Anxiety,
Contextual Fear Extinction, and Limbic Monoamines in Adult Rats. Frontiers in behavioral
neuroscience 10, 71.
29. Langleben, D.D., Ruparel, K., Elman, I., Loughead, J.W., Busch, E.L., Cornish, J., Lynch, K.G.,
Nuwayser, E.S., Childress, A.R. and O'Brien, C.P. (2014). Extended-release naltrexone modulates
brain response to drug cues in abstinent heroin-dependent patients. Addict Biol 19, 262-271.
30. Li, Q., Li, W., Wang, H., Wang, Y., Zhang, Y., Zhu, J., Zheng, Y., Zhang, D., Wang, L., Li, Y., Yan,
X., Chang, H., Fan, M., Li, Z., Tian, J., Gold, M.S., Wang, W. and Liu, Y. (2015). Predicting
subsequent relapse by drug-related cue-induced brain activation in heroin addiction: an event-
related functional magnetic resonance imaging study. Addict Biol 20, 968-978.
31. Hoffman, S.W. and Harrison, C. (2009). The interaction between psychological health and
traumatic brain injury: a neuroscience perspective. The Clinical neuropsychologist 23, 1400-
1415.
32. Taylor, P.A. and Ford, C.C. (2009). Simulation of blast-induced early-time intracranial wave
physics leading to traumatic brain injury. Journal of biomechanical engineering 131, 061007.
33. Mu, W., Catenaccio, E. and Lipton, M.L. (2017). Neuroimaging in Blast-Related Mild
Traumatic Brain Injury. The Journal of head trauma rehabilitation 32, 55-69.

34. Eierud, C., Craddock, R.C., Fletcher, S., Aulakh, M., King-Casas, B., Kuehl, D. and LaConte,
S.M. (2014). Neuroimaging after mild traumatic brain injury: review and meta-analysis.
NeuroImage: Clinical 4, 283-294.
35. Kalivas, P.W. and Volkow, N.D. (2005). The neural basis of addiction: a pathology of
motivation and choice. The American journal of psychiatry 162, 1403-1413.
36. Dong, Y., Taylor, J.R., Wolf, M.E. and Shaham, Y. (2017). Circuit and Synaptic Plasticity
Mechanisms of Drug Relapse. Journal of Neuroscience 37, 10867-10876.
37. Kalivas, P.W., Peters, J. and Knackstedt, L. (2006). Animal models and brain circuits in drug
addiction. Mol Interv 6, 339-344.
38. Peters, J., Kalivas, P.W. and Quirk, G.J. (2009). Extinction circuits for fear and addiction
overlap in prefrontal cortex. Learning & memory 16, 279-288.
39. Mayeux, J.P., Teng, S.X., Katz, P.S., Gilpin, N.W. and Molina, P.E. (2015). Traumatic brain
injury induces neuroinflammation and neuronal degeneration that is associated with escalated
alcohol self-administration in rats. Behavioural brain research 279, 22-30.
40. Weil, Z.M., Karelina, K., Gaier, K.R., Corrigan, T.E. and Corrigan, J.D. (2016). Juvenile
Traumatic Brain Injury Increases Alcohol Consumption and Reward in Female Mice. Journal of
neurotrauma 33, 895-903.
41. Weil, Z.M., Corrigan, J.D. and Karelina, K. (2016). Alcohol abuse after traumatic brain injury:
Experimental and clinical evidence. Neuroscience & Biobehavioral Reviews 62, 89-99.
42. Karelina, K., Nicholson, S. and Weil, Z.M. (2018). Minocycline blocks traumatic brain injury-
induced alcohol consumption and nucleus accumbens inflammation in adolescent male mice.
Brain, behavior, and immunity 69, 532-539.
43. Lowing, J.L., Susick, L.L., Caruso, J.P., Provenzano, A.M., Raghupathi, R. and Conti, A.C.
(2014). Experimental traumatic brain injury alters ethanol consumption and sensitivity. Journal
of neurotrauma 31, 1700-1710.
44. Merkel, S.F., Andrews, A.M., Lutton, E.M., Razmpour, R., Cannella, L.A. and Ramirez, S.H.
(2017). Dexamethasone Attenuates the Enhanced Rewarding Effects of Cocaine Following
Experimental Traumatic Brain Injury. Cell Transplant 26, 1178-1192.
45. Merkel, S.F., Razmpour, R., Lutton, E.M., Tallarida, C.S., Heldt, N.A., Cannella, L.A., Persidsky,
Y., Rawls, S.M. and Ramirez, S.H. (2017). Adolescent Traumatic Brain Injury Induces Chronic
Mesolimbic Neuroinflammation with Concurrent Enhancement in the Rewarding Effects of
Cocaine in Mice during Adulthood. Journal of neurotrauma 34, 165-181.
46. Vonder Haar, C., Ferland, J.N., Kaur, S., Riparip, L.K., Rosi, S. and Winstanley, C.A. (2019).
Cocaine self-administration is increased after frontal traumatic brain injury and associated with
neuroinflammation. The European journal of neuroscience 50, 2134-2145.
47. Grace, P.M., Strand, K.A., Galer, E.L., Urban, D.J., Wang, X., Baratta, M.V., Fabisiak, T.J.,
Anderson, N.D., Cheng, K., Greene, L.I., Berkelhammer, D., Zhang, Y., Ellis, A.L., Yin, H.H.,
Campeau, S., Rice, K.C., Roth, B.L., Maier, S.F. and Watkins, L.R. (2016). Morphine paradoxically
prolongs neuropathic pain in rats by amplifying spinal NLRP3 inflammasome activation. Proc
Natl Acad Sci U S A 113, E3441-3450.
48. Grace, P.M., Galer, E.L., Strand, K.A., Corrigan, K., Berkelhammer, D., Maier, S.F. and
Watkins, L.R. (2019). Repeated Morphine Prolongs Postoperative Pain in Male Rats. Anesthesia
and analgesia 128, 161-167.
49. Gaudet, A.D., Ayala, M.T., Schleicher, W.E., Smith, E.J., Bateman, E.M., Maier, S.F. and
Watkins, L.R. (2017). Exploring acute-to-chronic neuropathic pain in rats after contusion spinal
cord injury. Exp Neurol 295, 46-54.
50. Loram, L.C., Grace, P.M., Strand, K.A., Taylor, F.R., Ellis, A., Berkelhammer, D., Bowlin, M.,
Skarda, B., Maier, S.F. and Watkins, L.R. (2012). Prior exposure to repeated morphine

potentiates mechanical allodynia induced by peripheral inflammation and neuropathy. Brain,
behavior, and immunity 26, 1256-1264.
51. Hall, E.D., Wolf, D.L., Althaus, J.S. and Von Voigtlander, P.F. (1987). Beneficial effects of the
kappa opioid receptor agonist U-50488H in experimental acute brain and spinal cord injury.
Brain research 435, 174-180.
52. Statler, K.D., Kochanek, P.M., Dixon, C.E., Alexander, H.L., Warner, D.S., Clark, R.S.,
Wisniewski, S.R., Graham, S.H., Jenkins, L.W., Marion, D.W. and Safar, P.J. (2000). Isoflurane
improves long-term neurologic outcome versus fentanyl after traumatic brain injury in rats.
Journal of neurotrauma 17, 1179-1189.
53. Statler, K.D., Alexander, H., Vagni, V., Dixon, C.E., Clark, R.S., Jenkins, L. and Kochanek, P.M.
(2006). Comparison of seven anesthetic agents on outcome after experimental traumatic brain
injury in adult, male rats. Journal of neurotrauma 23, 97-108.
54. Zohar, O., Getslev, V., Miller, A.L., Schreiber, S. and Pick, C.G. (2006). Morphine protects for
head trauma induced cognitive deficits in mice. Neurosci Lett 394, 239-242.
55. Lyeth, B.G., Jiang, J.Y., Gong, Q.Z., Hamm, R.J. and Young, H.F. (1995). Effects of mu opioid
agonist and antagonist on neurological outcome following traumatic brain injury in the rat.
Neuropeptides 29, 11-19.
56. Barkhoudarian, G., Hovda, D.A. and Giza, C.C. (2011). The molecular pathophysiology of
concussive brain injury. Clin Sports Med 30, 33-48, vii-iii.
57. Niogi, S.N. and Mukherjee, P. (2010). Diffusion tensor imaging of mild traumatic brain
injury. The Journal of head trauma rehabilitation 25, 241-255.
58. Dodd, A.B., Epstein, K., Ling, J.M. and Mayer, A.R. (2014). Diffusion tensor imaging findings
in semi-acute mild traumatic brain injury. Journal of neurotrauma 31, 1235-1248.
59. Calabrese, E., Du, F., Garman, R.H., Johnson, G.A., Riccio, C., Tong, L.C. and Long, J.B. (2014).
Diffusion tensor imaging reveals white matter injury in a rat model of repetitive blast-induced
traumatic brain injury. Journal of neurotrauma 31, 938-950.
60. Badea, A., Kamnaksh, A., Anderson, R.J., Calabrese, E., Long, J.B. and Agoston, D.V. (2018).
Repeated mild blast exposure in young adult rats results in dynamic and persistent
microstructural changes in the brain. NeuroImage: Clinical 18, 60-73.
61. Venkatasubramanian, P.N., Keni, P., Gastfield, R., Li, L., Aksenov, D., Sherman, S.A., Bailes, J.,
3rd, Sindelar, B., Finan, J.D., Lee, J., Bailes, J.E. and Wyrwicz, A.M. (2020). Diffusion Tensor
Imaging Detects Acute and Subacute Changes in Corpus Callosum in Blast-Induced Traumatic
Brain Injury. ASN Neuro 12, 1759091420922929.
62. Sheth, C., Rogowska, J., Legarreta, M., McGlade, E. and Yurgelun-Todd, D. (2021). Functional
connectivity of the anterior cingulate cortex in Veterans with mild traumatic brain injury.
Behavioural brain research 396, 112882.
63. Czerniak, S.M., Sikoglu, E.M., Liso Navarro, A.A., McCafferty, J., Eisenstock, J., Stevenson,
J.H., King, J.A. and Moore, C.M. (2015). A resting state functional magnetic resonance imaging
study of concussion in collegiate athletes. Brain Imaging Behav 9, 323-332.
64. Meier, T.B., Giraldo-Chica, M., Espana, L.Y., Mayer, A.R., Harezlak, J., Nencka, A.S., Wang, Y.,
Koch, K.M., Wu, Y.C., Saykin, A.J., Giza, C.C., Goldman, J., DiFiori, J.P., Guskiewicz, K.M., Mihalik,
J.P., Brooks, A., Broglio, S.P., McAllister, T. and McCrea, M.A. (2020). Resting-State fMRI Metrics
in Acute Sport-Related Concussion and Their Association with Clinical Recovery: A Study from
the NCAA-DOD CARE Consortium. Journal of neurotrauma 37, 152-162.
65. Caeyenberghs, K., Verhelst, H., Clemente, A. and Wilson, P.H. (2017). Mapping the
functional connectome in traumatic brain injury: What can graph metrics tell us? NeuroImage
160, 113-123.
66. Hayes, J.P., Bigler, E.D. and Verfaellie, M. (2016). Traumatic Brain Injury as a Disorder of
Brain Connectivity. J Int Neuropsychol Soc 22, 120-137.

67. Sharp, D.J., Scott, G. and Leech, R. (2014). Network dysfunction after traumatic brain injury.
Nat Rev Neurol 10, 156-166.
68. Kaushal, M., Espana, L.Y., Nencka, A.S., Wang, Y., Nelson, L.D., McCrea, M.A. and Meier, T.B.
(2019). Resting-state functional connectivity after concussion is associated with clinical
recovery. Human brain mapping 40, 1211-1220.
69. Manning, K.Y., Schranz, A., Bartha, R., Dekaban, G.A., Barreira, C., Brown, A., Fischer, L.,
Asem, K., Doherty, T.J., Fraser, D.D., Holmes, J. and Menon, R.S. (2017). Multiparametric MRI
changes persist beyond recovery in concussed adolescent hockey players. Neurology 89, 2157-
2166.
70. Huang, M.X., Harrington, D.L., Robb Swan, A., Angeles Quinto, A., Nichols, S., Drake, A.,
Song, T., Diwakar, M., Huang, C.W., Risbrough, V.B., Dale, A., Bartsch, H., Matthews, S., Huang,
J.W., Lee, R.R. and Baker, D.G. (2017). Resting-State Magnetoencephalography Reveals
Different Patterns of Aberrant Functional Connectivity in Combat-Related Mild Traumatic Brain
Injury. Journal of neurotrauma 34, 1412-1426.
71. Hillary, F.G., Rajtmajer, S.M., Roman, C.A., Medaglia, J.D., Slocomb-Dluzen, J.E., Calhoun,
V.D., Good, D.C. and Wylie, G.R. (2014). The rich get richer: brain injury elicits hyperconnectivity
in core subnetworks. PLoS One 9, e104021.
72. Vakhtin, A.A., Calhoun, V.D., Jung, R.E., Prestopnik, J.L., Taylor, P.A. and Ford, C.C. (2013).
Changes in intrinsic functional brain networks following blast-induced mild traumatic brain
injury. Brain injury : [BI] 27, 1304-1310.
73. Robinson, M.E., Lindemer, E.R., Fonda, J.R., Milberg, W.P., McGlinchey, R.E. and Salat, D.H.
(2015). Close-range blast exposure is associated with altered functional connectivity in Veterans
independent of concussion symptoms at time of exposure. Human brain mapping 36, 911-922.
74. Wooten, D.W., Ortiz-Teran, L., Zubcevik, N., Zhang, X., Huang, C., Sepulcre, J., Atassi, N.,
Johnson, K.A., Zafonte, R.D. and El Fakhri, G. (2019). Multi-Modal Signatures of Tau Pathology,
Neuronal Fiber Integrity, and Functional Connectivity in Traumatic Brain Injury. Journal of
neurotrauma 36, 3233-3243.
75. Kulkarni, P., Morrison, T.R., Cai, X., Iriah, S., Simon, N., Sabrick, J., Neuroth, L. and Ferris, C.F.
(2019). Neuroradiological Changes Following Single or Repetitive Mild TBI. Front Syst Neurosci
13, 34.
76. Wollman, S.C., Alhassoon, O.M., Stern, M.J., Hall, M.G., Rompogren, J., Kimmel, C.L. and
Perez-Figueroa, A.M. (2015). White matter abnormalities in long-term heroin users: a
preliminary neuroimaging meta-analysis. Am J Drug Alcohol Abuse 41, 133-138.
77. Ma, X., Qiu, Y., Tian, J., Wang, J., Li, S., Zhan, W., Wang, T., Zeng, S., Jiang, G. and Xu, Y.
(2015). Aberrant default-mode functional and structural connectivity in heroin-dependent
individuals. PLoS One 10, e0120861.
78. Li, W., Zhu, J., Li, Q., Ye, J., Chen, J., Liu, J., Li, Z., Li, Y., Yan, X., Wang, Y. and Wang, W.
(2016). Brain white matter integrity in heroin addicts during methadone maintenance
treatment is related to relapse propensity. Brain Behav 6, e00436.
79. Sun, Y., Wang, G.B., Lin, Q.X., Lu, L., Shu, N., Meng, S.Q., Wang, J., Han, H.B., He, Y. and Shi,
J. (2017). Disrupted white matter structural connectivity in heroin abusers. Addict Biol 22, 184-
195.
80. Chang, H., Li, W., Li, Q., Chen, J., Zhu, J., Ye, J., Liu, J., Li, Z., Li, Y., Shi, M., Wang, Y. and
Wang, W. (2016). Regional homogeneity changes between heroin relapse and non-relapse
patients under methadone maintenance treatment: a resting-state fMRI study. BMC Neurol 16,
145.
81. Qiu, Y.W., Han, L.J., Lv, X.F., Jiang, G.H., Tian, J.Z., Zhuo, F.Z., Su, H.H., Lin, C.L. and Zhang,
X.L. (2011). Regional homogeneity changes in heroin-dependent individuals: resting-state
functional MR imaging study. Radiology 261, 551-559.

82. Zou, F., Wu, X., Zhai, T., Lei, Y., Shao, Y., Jin, X., Tan, S., Wu, B., Wang, L. and Yang, Z. (2015).
Abnormal resting-state functional connectivity of the nucleus accumbens in multi-year
abstinent heroin addicts. Journal of neuroscience research 93, 1693-1702.
83. Ma, N., Liu, Y., Li, N., Wang, C.X., Zhang, H., Jiang, X.F., Xu, H.S., Fu, X.M., Hu, X. and Zhang,
D.R. (2010). Addiction related alteration in resting-state brain connectivity. NeuroImage 49,
738-744.
84. Liu, J., Liang, J., Qin, W., Tian, J., Yuan, K., Bai, L., Zhang, Y., Wang, W., Wang, Y., Li, Q., Zhao,
L., Lu, L., von Deneen, K.M., Liu, Y. and Gold, M.S. (2009). Dysfunctional connectivity patterns in
chronic heroin users: an fMRI study. Neurosci Lett 460, 72-77.
85. Pandria, N., Kovatsi, L., Vivas, A.B. and Bamidis, P.D. (2018). Resting-state Abnormalities in
Heroin-dependent Individuals. Neuroscience 378, 113-145.
86. Ieong, H.F. and Yuan, Z. (2017). Resting-State Neuroimaging and Neuropsychological
Findings in Opioid Use Disorder during Abstinence: A Review. Front Hum Neurosci 11, 169.
750