Optimal density of bacterial cells

Tin Yau Pang[1], Martin J. Lercher[1,*]

[1] Institute for Computer Science & Department of Biology, Heinrich Heine University, 40225 Düsseldorf, Germany

* To whom correspondence should be addressed

Keywords: macromolecular crowding, Michaelis Menten kinetics, metabolic reaction, ribosome, translation

Abstract

A substantial fraction of the bacterial cytosol is occupied by catalysts and their substrates. While a higher volume density of catalysts and substrates might boost biochemical fluxes, the resulting molecular crowding can slow down diffusion, perturb the reactions' Gibbs free energies, and reduce the catalytic efficiency of proteins. Due to these tradeoffs, dry mass density likely possesses an optimum that facilitates maximal cellular growth and that is interdependent on the cytosolic molecule size distribution. Here, we analyze the balanced growth of a model cell, accounting systematically for crowding effects on reaction kinetics. Its optimal cytosolic volume occupancy depends on the nutrient-dependent resource allocation into large ribosomal vs. small metabolic macromolecules, reflecting a tradeoff between the saturation of metabolic enzymes, favoring larger occupancies with higher encounter rates, and the inhibition of the ribosomes, favoring lower occupancies with unhindered diffusion of tRNAs. Our predictions across growth rates are quantitatively consistent with the experimentally observed reduction in volume occupancy on rich media compared to minimal media in *E. coli*. Strong deviations from optimal cytosolic occupancy only lead to minute reductions in growth rate, which are nevertheless evolutionarily relevant due to large bacterial population sizes. In sum, cytosolic density variation in bacterial cells appears to be consistent with an optimality principle of cellular efficiency.

Introduction

The dry mass dissolved in the major compartment of bacterial cells, the cytosol, comprises hundreds of molecular species, including proteins, metabolites, polysaccharides, and nucleic acids. These molecules can be roughly classified into two sectors: the ribosomal sector, dominated by ribosomes and tRNA; and the non-ribosomal sector, comprising mostly metabolites, enzymes, and other proteins (Scott et al., 2010). The molecules in these two sectors have very different size distributions: the ribosome is 65 times larger than the median enzyme size (2,600kDa (Rabl et al., 2011) vs. 40kDa (Neidhardt and Curtiss, 1996)), and tRNAs are about 300 times larger than typical metabolites (26kDa (Nierlich, 1972) vs. 89Da, the mass of alanine). The allocation of dry mass between the two sectors of the cellular economy can be summarized by a single parameter, the growth rate μ, with the dry mass fraction of the ribosomal sector increasing almost linearly with μ (Hu et al., 2020; Scott et al., 2010). Accordingly, the ribosome-rich cytosol at fast growth in nutrient-rich environments and the ribosome-meager cytosol at slow growth in nutritionally poor environments exhibit very different distributions of molecule sizes.

Experiments have found an approximately constant dry mass density of the cytosol at 300g/L across minimal nutrient conditions with low growth rates, but a roughly 10% less dense cytosol in a rich medium supporting high growth rates (Oldewurtel et al., 2021). We hypothesized that the observed difference in cytosolic density between slow and fast growth may be an evolutionary consequence of the differences in molecular composition. Cellular physiology has evolved under natural selection; thus, if the level of molecular crowding—the dry mass density of the cytosol—affects cellular efficiency and hence fitness, we expect density regulation to have evolved to a near-optimal, possibly condition-dependent state. While previous work has explored the influence of (macro-)molecular crowding on bacterial physiology and growth rates, these analyses assumed a constant, hard limit on the total cytosolic protein (Beg et al., 2007; Weiße et al., 2015) or dry mass (Dourado and Lercher, 2020; Molenaar et al., 2009) concentration. These works do not justify the existence and magnitude of the density limit based on physicochemistry,

55   and they cannot explain differences in the level of crowding (dry mass density) across conditions.

Biochemical reaction fluxes typically increase with increasing encounter rates of the molecule species involved; ignoring crowding effects on diffusion, encounter rates increase with increasing density of the respective molecules. At the same time, the molecular crowding

60   caused by other molecule species in the background (volume-excluding co-solutes) affects fluxes in at least three distinct ways. (i) Crowding slows down the diffusion of a catalyst and its substrates, thereby reducing their encounter rates (Minton, 1981; Zimmerman and Minton, 1993). (ii) Crowding limits the available volume and thus reduces a solute's entropy (the "excluded volume phenomenon"), thereby changing the free energies of the molecules involved

65   in the reaction and consequently shifting the equilibrium concentrations of substrates and products (Ralston, 1990). (iii) Crowding can affect the structure of the protein catalyst, the process of its folding, and its conformational stability (Kuznetsova et al., 2014); these structural changes may disturb the reaction flux if they affect the active site (Kuznetsova et al., 2014; Norris and Malys, 2011; Olsen, 2006; Pozdnyakova and Wittung-Stafshede, 2010; Totani et al.,

70   2008). Due to these opposing effects, there may be an optimal cytosolic density where cellular efficiency and hence fitness are maximal.

The effect of crowding depends on the size of the catalyst and its substrate: in the presence of other volume-excluding cosolutes, the larger the size of a solute, the stronger the reduction of its diffusion (Kalwarczyk et al., 2012) and the perturbation of its free energy (Kihara, 1953; Minton,

75   1998). Hence, when the size distribution of the molecules changes, the cell needs to adjust its cytosolic density in order to optimize its physiological efficiency.

In a pioneering theoretical study, Vazquez (Vazquez, 2010) considered how the objective flux of a metabolic network is affected by crowding, assuming that the enzymes are also the crowders in their own right. The corresponding model accounts for the slowdown of diffusion due to

80   crowding, but ignores the growth-rate dependent role of the ribosomal sector and the corresponding changes in the molecular size distribution. This study found that there exists an

optimal cytosolic density, which maximizes reaction fluxes and depends on details of the network, such as the ratio between diffusion limited and transition-state limited reactions. Using an alternative modeling approach, Dill et al. also arrive at a similar conclusion (Dill et al., 2011).

85 While these studies demonstrated the existence of a flux-optimizing cytosolic density, natural selection maximizes fitness, not metabolic reaction fluxes. For non-interacting cells in a uniform environment, fitness is closely related to the growth rate (Fisher, 1930)**.** The cellular physiology at maximal growth rate—and hence maximal fitness—can be described mathematically through growth balance analysis (GBA) (Dourado and Lercher, 2020).This modeling framework

90 simulates the balanced growth of a self-replicating bacterial cell while accounting for the major physicochemical constraints on cellular growth: mass balance of metabolism and protein production, non-linear reaction kinetics that depend on the concentrations of catalysts and their substrates, and the effects of molecular crowding.

Standard GBA assumes a given, hard limit on dry mass density (Dourado and Lercher, 2020;

95 Molenaar et al., 2009) or on total protein concentration (Weiße et al., 2015). Here, to assess the effects of molecular crowding on fitness, we apply a generalization of GBA that instead describes the kinetics of metabolic reactions and protein translation through crowding-adjusted Michaelis-Menten kinetics (Aoki et al., 2013; Minton, 1998). We maximize the balanced growth rate while varying the concentrations of transporters, catalytic proteins, and metabolites; these

100 molecular species also form the volume excluding co-solutes in the background of each reaction, affecting diffusion, free energies, and hence reaction kinetics through molecular crowding. Consistent with experimental observations, we find that the cytosolic density of optimal growth strategies depends on the cellular growth rate.

 Results

Crowding-adjusted reaction kinetics

The effects of molecular crowding on biochemical reaction kinetics are due to volume exclusion

effects. Thus, the relevant parameter for their quantification is not the total mass density (dry

mass per volume) but the total volume occupancy ρ, i.e., the fraction of cytosolic volume

110   occupied by dry mass. However, as molecular mass and volume are approximately proportional

(Lee, 1983) and as experimental measurements for the cytosolic occupancy are not available,

we treat density and occupancy as interchangeable, subject to a scaling coefficient that

quantifies the mass/volume ratio of the cytosolic dry mass.

To model biochemical reaction kinetics as a function of molecular crowding, we use a modified

115   description of irreversible Michaelis-Menten kinetics (Aoki et al., 2013; Minton, 1998) (see

Methods for details),

$$v = k_{\text{cat}} \frac{[s][E]}{K_{\text{M}}^* + [s]} \ ,$$

(1)

where [S] and [E] are the concentrations of the substrate and catalyst, respectively; k_{cat} is the

catalytic rate constant (turnover number); and K_{M}^* is the crowding-adjusted Michaelis parameter

(Aoki et al., 2013; Minton, 1998). In a typical catalytic reaction, a substrate molecule S

120   encounters a catalyst molecule E to form a catalyst-substrate complex ES, which either

proceeds forward to convert the substrate into the product, or reverts back to release the

substrate. To derive K_{M}^*, we assume that a catalytic reaction can be divided into two

subsequent, independent steps that both depend on the cytosolic volume occupancy: (1) S and

E diffuse until their encounter (Fig. 1B) and thereafter (2) they bind and unbind reversibly until

125   the reaction proceeds forward and product P is released (Fig. 1A). In step (2), we further

assume that S and E will stay in close proximity and do not diffuse away from each other. These

approximations simplify the model derivation and make the reaction times of the two steps

additive (Minton, 1998). Combining the rate laws of the two steps provides an estimate for the

effective Michaelis parameter K_{M}^* (Aoki et al., 2013; Minton, 1998):

Figure 1. Limiting cases of crowding effects on kinetics. (A) Transition state limit, valid if the conversion of enzyme (E) plus substrate (S) to the complex ES is much faster than the conversion of ES to product (P). **(B)** Diffusion limit, where the conversion of ES to P is much faster than the formation of ES.

$$K_M^* = K_M^0 \frac{\Gamma + \theta e^{-g\rho}}{(1+\theta) \times \Gamma \times e^{-g\rho}}.$$ (2)

Here, K_M^0 is the Michaelis parameter in the low-crowding limit; Γ is the correction term for the shift in Gibbs free energy ΔG due to molecular crowding; the exponential term exp(-$g\rho$) accounts for the slow-down of diffusion, where the scaling factor g can be estimated from the size of the substrate, i.e.,   for a spherical substrate with radius r_S; and θ is the relative weight

140   between the rate laws of steps (1) and (2), or in other words the relative ratio between time spent on step (1) and on step (2) at low cytosolic occupancy. The best available estimate for θ is 2.3, obtained for the ERK MAP kinase phosphorylation reaction (Aoki et al., 2013); we employ this value throughout our simulations.

Protein translation favors lower occupancy compared to metabolic pathways

145   To understand the effect of crowding on catalytic reactions, we first consider a simple, linear biochemical pathway model consisting of N=20 consecutive enzyme-catalyzed reactions at steady state, i.e., with identical flux through each reaction (Fig. 2A). The kinetics of each step in the pathway depend not only on the concentrations of the catalyzing enzyme and the substrate of the reaction, but—through the crowding effects on K_M^*—also on the concentrations of the

150   molecules involved in the remaining N-1 reactions. We identified the combination of enzyme and metabolite concentrations that maximizes the pathway output per dry mass, calculated from crowding-adjusted kinetics; adding these concentrations, weighted by the respective molecular

Original Model **Simplified Model** **Model Equations**

A Linear Model

B Parallel Model

$$\begin{cases} \mu = \dfrac{k_{cat}}{N}\dfrac{[s][E]}{NK_M^*+[s]}\dfrac{1}{\rho} \\ \rho = N_A\dfrac{4\pi}{3}([s]r_s^3 + [E]r_E^3) \end{cases}$$

C Bacterial Cell Model

$$\begin{cases} \mu = k_{cat}^{R}\dfrac{[p][R]}{K_M^{R^*}+[p]}\dfrac{1}{N_A(l_M[M]+l_R[R]+l_T[T])} \\ \rho = N_A\dfrac{4\pi}{3}([s]r_s^3 + [M]r_M^3 + [p]r_p^3 + [R]r_R^3) \\ 0 = k_{cat}^{T}\dfrac{[s_{ext}][T]}{K_M^{T}+[s_{ext}]} - \dfrac{k_{cat}^{M}}{N}\dfrac{[s][M]}{NK_M^{M^*}+[s]} - \mu[s] \\ 0 = \dfrac{k_{cat}^{M}}{N}\dfrac{[s][M]}{NK_M^{M^*}+[s]} - k_{cat}^{R}\dfrac{[p][R]}{K_M^{R^*}+[p]} - \mu[p] \end{cases}$$

Figure 2. Illustration of the models. (A) Reaction system in an *N*-steps linear pathway,
representing a metabolic system. The initial substrate, s_1, is replenished by a transport process not
included in the model, and is converted by the pathway into the product p. **(B) Reaction system of**
***N* parallel reactions**, representing ribosomes presenting N distinct anticodons; s_1, ... , s_N are the
corresponding ternary complexes, p_1, …, p_N are the extending amino acid chains. Assuming that (i)
all reactions follow identical kinetics, (ii) all substrate concentrations s_i are identical, and (iii) all
enzyme concentrations are identical, the fluxes of the models in (A) and (B) are both mathematically
identical to the simplified model to the right of the two panels, with re-scaled concentrations
$[s]=[s_1]+[s_2]+...+[s_N]$ and $[E]=[E_1]+[E_2]+...+[E_N]$. **(C) GBA model simulating the balanced growth of**
a bacterial cell. Transporter T imports nutrient s_1, which is converted to the precursor for protein
production p by a metabolic pathway with N consecutive enzymes, M_1, …, M_N, via intermediate
substrates s_2, …, s_N. The ribosome R synthesizes the $N+2$ proteins (T, R, M_1, …, M_N) from p.
Assuming identical concentrations of metabolic enzymes, $[M_1]=...[M_N]=:[M]$, and of metabolites,
$[s_1]=...[s_N]=:[s]$, the solution space of this model cell is spanned by the five concentrations
$[s],[p],[T],[M],[R]$.

volumes, resulted in an estimate of the optimal cytosolic occupancy ρ_{opt} for this system. To

restrict the number of model parameters, we assumed identical enzyme and substrate sizes as

well as identical crowding-adjusted kinetics (Eq. (1)) for all reactions in the pathway. We fixed

the physicochemical parameter K_M^0=130μM, which is the median value for metabolic enzymes

and their substrates (Bar-Even et al., 2011) and is also close to the value estimated for the

175   binding of ternary complexes to the ribosome (Klumpp et al., 2013), 120µM. The second

physicochemical parameter in this model, k_{cat}, appears merely as a scaling factor of the

pathway flux, and for simplicity we set $k_{cat}=1s^{-1}$ without loss of generality.

We varied the sizes of the catalyst and its substrate, which we assumed to be both spherical.

We considered (i) a metabolic pathway, where substrates have sizes typical for metabolites

180   (r=0.34nm) and catalyst have sizes typical for globular proteins (r=2.4nm); and (ii) a ribosomal

system with sizes resembling those of tRNAs (r=2.4nm) and ribosomes (r=13nm). We also

assumed the catalyst-substrate complex to be spherical, with a volume equal to that of the

catalyst plus the substrate. Note that while cells do not contain pathways of consecutive

ribosomes, the model for the 20-step linear pathway depicted in Fig. 2A is mathematically

185   identical to the model for 20 parallel reactions shown in Fig. 2B if all metabolite concentrations

are assumed to be identical; this latter model can be interpreted as a molecular snapshot

showing 20 types of ribosomes, each presenting an anticodon for a different tRNA species.

Fig. 3 shows that the optimal occupancy is ρ_{opt}=0.30 for the metabolic system and ρ_{opt}=0.12 for

the ribosomal system. The effect of the occupancy on the pathway output is substantial in the

190   ribosomal system, where a 50% drop in occupancy from ρ_{opt} decreases pathway flux per dry

mass by 21%, while it is small for the metabolic system, where a 50% drop in occupancy incurs

a flux decrease per dry mass of only 1.3%. Increasing the number of steps in the pathway from

20 to 100 has almost no effects on the optimal occupancies ρ_{opt} for both systems; however, the

reduction of flux per dry mass when ρ deviates from ρ_{opt} is substantially increased for longer

195   pathways or more parallel reactions (Fig. 3, dashed lines).

We conclude that the results from the pathway model are consistent with the experimentally

observed trend that the cytosol of fast-growing bacteria, which is dominated by the larger

molecules of the ribosomal sector, favors a lower occupancy than the cytosol of a slowly growing

cell dominated by smaller metabolites and enzymes. Moreover, our results suggest that

200   compared to cells dominated by the ribosomal sector, cells dominated by the smaller molecules

of the metabolic sector may suffer a smaller reduction in biochemical efficiency when the cytosol shifts away from ρ_{opt}.

Figure 3. The cytosolic occupancy that facilitates maximal biochemical reaction fluxes is lower for ribosomal than for metabolic systems. Blue lines represent metabolic systems with small catalysts and substrates, red lines represent ribosomal systems with much larger molecules. Solid lines are for systems of N=20 consecutive (metabolic, blue) or parallel (ribosomal, red) reactions, dashed lines are for larger systems of N=100 reactions.

Optimal occupancy for a self-replicating cell at balanced growth

Can the single-pathway results be generalized to more realistic models of cellular growth, which combine metabolic and ribosomal activities, and where we can directly assess the effect of concentration changes on cellular growth rates? How does the cytosolic occupancy optimal for growth, ρ_{opt}, change when the number of active metabolic reactions changes, e.g., when switching from a minimal medium, where all biomass components have to be synthesized from a single carbon source through multi-step biochemical pathways, to a rich medium that provides many cellular building blocks through simple transport processes?

To answer these questions, we considered a schematic GBA model of a bacterial cell (Dourado and Lercher, 2020), the cytosol of which comprises interdependent ribosomal and metabolic sectors (**Fig. 2C**). In this model cell, a nutrient s_{ext} is imported into the cell by transport protein T;

the nutrient is then converted into precursor p by an N-steps metabolic pathway; finally, the ribosome R uses p to synthesize all catalytic proteins, including T, R, and the metabolic enzymes M_1 to M_N. For simplicity, all molecules are again assumed to be spherical in shape. All macromolecules, except for the transporter protein T, are located in the cytosol and are thus
225   crowders in their own right; T is assumed to be fully integrated into the membrane and does not contribute to crowding. As before, the molecules of the metabolic sector are small (metabolites s_i with r_s=0.34nm and metabolic enzymes M_i with r_M=2.4nm), while the constituents of the ribosomal sector are much larger (precursor p with r_p=2.4nm and ribosome R with r_R=13nm).

To estimate the appropriate number of active enzymatic reactions in the metabolic sector, N, we
230   used flux balance analysis constrained by enzyme concentration (Adadi et al., 2012); simulations were performed with an improved implementation parameterized for a genome-scale model of *Escherichia coli* metabolism (Desouki, 2016). We found 259 active metabolic enzymes for growth in a minimal medium with glucose as the sole carbon source; 206 active enzymes for the same medium supplemented with amino acids; and 174 active enzymes for growth in a rich
235   medium (Methods).

To facilitate the numerical determination of the state with maximal growth rate, we approximate the metabolic pathway through a single, lumped reaction with catalyst M and substrate s, scaling kinetic parameters and molecular masses to account for the pathway length N; this approximation neglects the dilution of intermediate metabolite concentrations through volume
240   growth (Methods). The solution space of the model with N metabolic enzyme-catalyzed reactions is spanned by 5 concentration variables (**Fig. 2C**), describing the transporter protein with concentration $[T]$, the cytosolic substrates with identical concentrations $[s_1]=...=[s_N]=:[s]$, the protein precursor ("ternary complex") with concentration $[p]$, the metabolic enzymes with identical concentrations $[M_1]=...=[M_N]=:[M]$, and the ribosome with concentration $[R]$.

245   The metabolic and ribosomal reactions are described by crowding-adjusted irreversible Michaelis Menten kinetics (Eq. (1)), whereas the transporter reaction is described by conventional irreversible Michaelis-Menten kinetics with constant K_M^T. We investigated how the

cytosolic occupancy that allows the fastest growth varies with two parameters, the number of

enzyme species in the metabolic pathway, N, inversely related to the richness of the nutrient

250 composition and entering the model through scaling the lumped metabolic reaction; and the

concentration of the external nutrient, $[s_{ext}]$, which parameterizes the degree to which the

available nutrients are limiting growth.

Optimal occupancy is lower at faster growth due to higher protein translation demands

255 In the whole-cell model, the optimal occupancy ρ_{opt} generally falls between 0.1 and 0.3 (Fig. 4).

ρ_{opt} increases when the number of simultaneously active metabolic reactions N increases or

when the external nutrient concentration s_{ext} (and consequently the growth rate) decreases (Fig.

5). These effects are due to shifts in the relative dry mass fractions of the metabolic sector

(favoring higher occupancy) and the ribosomal sector (favoring lower occupancy). At increasing

260 N and constant s_{ext} , the fraction of the cytosolic volume occupied by the metabolic sector

expands at the expense of the ribosomal sector (Fig. S2A), and the saturation of the ribosome

with its substrate drops, whereas the saturation of the metabolic enzymes with their substrates

varies within a small range (Fig. S2B). At constant N=250 and improving nutrient conditions

(increasing s_{ext}), the fraction of the cytosol occupied by the metabolic sector also expands

265 slightly at the expense of the ribosomal sector (Fig. S4A), while the saturation of both metabolic

enzymes and ribosomes increases (Fig. S4B). Note that while the cytosolic concentration of

ribosomes decreases with increasing growth rate (Fig. S3B), the ribosomal proteome fraction

increases (Fig. S3C), mirroring experimental observations (Hui et al., 2015; Scott et al., 2010).

At optimal occupancy ρ_{opt} for different N and s_{ext} , the crowding-adjusted Michaelis parameter K_M^*

270 of the ribosomal reaction shows a marked increase with ρ_{opt} (Fig. 6; two-sided Spearman rank

correlation coefficient r=0.995, P<10^{-15}), consistent with our observations of a strong

dependence of the ribosomal flux on ρ in the simple pathway model (Fig. 3). In contrast, the K_M^*

Figure 4. The growth rate μ of the crowding-adjusted whole-cell model constrained at

275   **different cytosolic occupancies ρ.** The optimal cytosolic occupancy ρ_{opt} **(A)** increases with N (the
number of enzymes in the metabolic pathway) and **(B)** decreases with external nutrient
concentration $[s_{ext}]$. μ_{max} is the maximal growth rate for each curve across occupancies.

280   **Figure 5. The optimal cytosolic occupancy increases with metabolic pathway length N and
decreasing external nutrient concentration s_{ext}.**

Figure 6. The optimal occupancy strongly influences the effective Michaelis parameter K_M^* of the ribosomal (red) but not of the metabolic (blue) reactions in the whole-cell model. Each data point corresponds to a different combination of external nutrient concentration s_{ext} and number of active metabolic reactions N. While the K_M^* of the metabolic reactions does not correlate with ρ_{opt} (blue markers; two-sided Spearman's rank correlation coefficient $r=0.034$, $P=0.85$), the K_M^* of the ribosomal reactions correlates with ρ_{opt} (red markers; $r=0.998$, $P<10^{-15}$). The discrete distribution of points along the x-axis reflects the step size used for ρ in the simulations.

Figure 7. Opposing effects of molecular crowding on the Michaelis parameter K_M^* through perturbations of diffusion and of Gibbs free energies. Simulations considered only the respective effect for systems of (linear, blue) metabolic and (parallel, red) ribosomal reactions with pathway sizes of (A) $N=20$ and (B) $N=100$.

of the metabolic reactions is almost invariant when plotted against ρ_{opt} (r=0.034, P=0.85). These observations support the intuitive notion that the optimal occupancy in the whole-cell model reflects a tradeoff between the saturation of metabolic enzymes, favoring larger occupancies with higher encounter rates, and the inhibition of the ribosomes, favoring lower occupancies with

300　　unhindered diffusion of tRNAs. In agreement with our findings for the simple pathway model, the dependence of the growth rate on ρ appears to be moderate: e.g., at N=150 and s_{ext}=1μM, a 50% reduction of ρ from ρ_{opt} reduces μ by only 1.1% (Fig. 4).

The slow-down of diffusion and the perturbation of Gibbs free energies have opposing effects on reaction efficiencies. To consider these two effects separately, we define the transition state-

305　　perturbation only Michaelis parameter as given by Eq. (2) when setting the diffusion scaling exponent to g=0, and we define the diffusion-perturbation only Michaelis parameter as given by Eq. (2) when setting the Gibbs perturbation term to Γ=1. We used these hypothetical Michaelis parameters in renewed simulations of the pathway models of the metabolic and ribosomal systems (Fig. 2A,B). In the diffusion-perturbation only model, crowding increases K_M^* and

310　　reduces the reaction fluxes (Fig. 7, dotted lines); in contrast, crowding in the transition state-perturbation only model decreases K_M^* and boosts the fluxes (Fig. 7, dashed lines). At high occupancies (ρ≥0.5), the transition state-perturbation effects reach a plateau, whereas the

diffusion-perturbation effects continue to increase. Thus, at larger occupancies, the flux-reducing slow-down of diffusion always dominates over the flux-enhancing perturbation of Gibbs free

315　　energy when considering their joint effect (Fig. 7, solid lines). Comparison of Fig. 7A and B shows that these trends are largely independent of N, the number of reactions in the system. While the trend for the slow-down of diffusion depends on the sizes of the reacting molecules, the trend for the perturbation of Gibbs free energies is largely independent of molecule sizes.

Optimal occupancy is quantitatively consistent with the observed E. coli dry mass
320   *density*

To compare our predictions of optimal occupancy to experimental data from *E. coli*, we first consider the transition from slow to fast growth on minimal media (simulated extracellular nutrient concentration s_{ext}=0.1µM vs. 1µM, at a constant number of active metabolic reactions N=250). Here, the optimal cytosolic volume occupancy ρ_{opt} predicted by our whole-cell model
325   decreases from 0.250 down to 0.234 (a 7% drop, Fig. 5).

While no direct experimental observations of the occupancy ρ are available, occupancy is expected to scale approximately in proportion to the cytosolic dry mass density ρ_{DM} (Lee, 1983). Empirical observations indicate that ρ_{DM} does not vary noticeably with μ across minimal media, i.e., ρ_{DM}=0.31g/mL (with 5% measurement error) when $\mu \leq 0.7h^{-1}$ (see Fig. 2B of Ref (Oldewurtel
330   et al., 2021)). As cellular resources are shifted from protein to ribosomal RNA and tRNA with increasing growth rate, and because RNA is denser than protein, the conversion factor between dry mass and volume changes with growth rate. Taking this into account, we find that the experimental data indicates a decrease of ρ from 0.221 to 0.208, a 6% reduction that is highly consistent with our estimate (Methods).

335   Second, we consider the transition from fast growth on minimal media to ultrafast growth in rich media (simulated extracellular nutrient concentration s_{ext} of 1µM vs. 10µM, number of active metabolic reactions N=250 vs. N=150). Here, the predicted optimal cytosolic volume occupancy ρ_{opt} decreases by 15%, from 0.234 to 0.204 (Fig. 5). Empirical observation shows that ρ_{DM} in *E. coli* decreases from 0.31g/mL to 0.28g/mL as μ increases from 0.7h^{-1} to 1.2h^{-1} (Oldewurtel et al.,
340   2021), corresponding to a reduction in occupancy from ρ=0.208 to ρ=0.184, an 11% decrease. Thus, the predicted change in ρ_{opt} is also consistent with the empirically observed reduction in occupancy in the transition to ultrafast growth.

Our simulations also show that even if the cell remains at the state of optimal occupancy, a large reduction in the nutrient level results in only a small decrease of the ribosome's saturation with

345 its substrate (0.81 to 0.72, Fig. S4B); at the same time, the ribosome's Michaelis parameter K_M^*

increases by 51% (Fig. S4C). This finding is consistent with experimental observations that

changes in translation rate per ribosome in *E. coli*, a direct consequence of ribosomal substrate

saturation, are much smaller than the simultaneous changes in growth rate (Dai et al., 2016). At

increasing nutrient levels, a higher growth rate μ is facilitated by relocating a substantial amount

350 of the ribosome's synthesis capacity from transporter proteins to metabolic enzymes and

additional ribosomes (Fig. S3C).

Discussion

The linear pathway model shows that reactions with larger catalysts and substrates favour a

lower occupancy than reactions with smaller molecules (Fig. 3). This effect explains the

355 observation of lower optimal occupancies for decreasing pathway length N in the GBA model

cell: the decrease in pathway length simulates the switch from minimal media, requiring on the

order of 260 metabolic enzyme species to convert a small number of nutrients to the full range of

cellular building blocks, to increasingly richer media, where progressively more biomass

components can be taken up directly from the environment, requiring as few as 140 metabolic

360 enzyme species. With decreasing numbers of active metabolic reactions N, the ribosomal sector

expands at the expense of the metabolic sector, pushing ρ_{opt} to lower values (Fig. S2A). Given

the minimalistic nature of the whole-cell model, which assumes that all metabolic reactions

follow identical kinetics and approximates protein production through a single Michaelis-Menten

type reaction, it is striking that the model not only predicts differences across physiological

365 states, but predicts experimentally observed values (Oldewurtel et al., 2021) quantitatively with

an error below 12%.

The whole-cell model predicts that the growth rate μ decreases only mildly when ρ deviates from

ρ_{opt}; for example, μ decreases by only 3% when ρ increases to twice the optimal occupancy ρ_{opt}

(Fig. 4; N=150, s_{ext}=1.0μM). This observation is consistent with an experiment that arrested the

370 volume growth of a yeast cell while cytosolic dry mass continued to accumulate, increasing the

concentration of a fluorescent protein—a proxy for dry mass density—to twice the wildtype value

(Knapp et al., 2019). These results suggest that the dry mass accumulation rate largely remains constant throughout the volume growth arrest, even when the cytosol density reaches twice the wildtype level.

375 In the whole-cell model, a 10% deviation of ρ from ρ_{opt} results in a 0.02% drop in μ (Fig. 4); using growth rate as a proxy for fitness, this corresponds to a selection coefficient $s=2\text{x}10^{-4}$. The effective population size of most bacterial species is on the order of $N_e=10^8$ (Bobay and Ochman, 2018), and we thus have $s \gg 1/N_e = 10^{-8}$; accordingly, natural selection would be sufficiently strong to explain the difference in average ρ_{DM} observed between the two

380 physiological states. Experiments show substantial between-cell variation in each nutritional condition (Oldewurtel et al., 2021): the observed dry mass densities show coefficients of variation (standard deviation / mean) of around 5%. According to the whole-cell model, the corresponding difference in occupancy corresponds to a selection coefficient of $s = 10^{-4} \gg 1/N_e$, indicating that this large cell-to-cell variation is not selectively neutral but persists despite

385 negative selection.

Questions about the optimal allocation of protein resources can be addressed by maximizing the growth rate (or flux) in computational models of cellular growth with fixed, crowding-unaware kinetic parameters. While such simulations produce meaningful predictions for the relative amounts of the proteins, they cannot limit absolute protein concentrations (Sezer et al., 2020).

390 To solve this problem, existing models (Dourado and Lercher, 2020; Molenaar et al., 2009) implement hard, phenomenological constraints on the total concentration of protein or cellular dry mass, based on experimental observations that found these to be approximately constant across growth conditions (Hui et al., 2015; Oldewurtel et al., 2021; Schmidt et al., 2015; Woldringh and Nanninga, 1985). Our results elucidate the biophysical origin of these

395 observations, showing that the cellular dry mass density represents a compromise between the saturation of metabolic enzymes with their substrates and the effects of reduced diffusion on the effective affinity of the ribosome for its much larger substrate, the ternary complex.

Methods

Crowding-adjusted Michaelis-Menten kinetics

400   Macromolecular crowding affects the flux of a metabolic reaction in multiple ways. It can (i) slow

down diffusion; (ii) affect the free energy of substrate, catalyst, and the substrate-catalyst

complex and thereby change their relative equilibrium ratios; and (iii) disturb the folding of a

protein and affect the shape of the active site. In our modelling framework, we followed the

derivation proposed in Ref (Minton, 1998) that systematically accounts for the effects of

405   crowding on metabolic fluxes caused by effects (i) and (ii).

In this section, let us consider the metabolic reaction carried out by an enzyme E that converts

substrate S into product P, in the presence of other volume-excluding co-solutes that,

collectively, constitute the dry mass of the solution. The metabolic reaction is described by the

chemical equation following Michaelis-Menten kinetics:

$$\text{E} + \text{S} \underset{k_{\text{off}}}{\overset{k_{\text{on}}}{\rightleftharpoons}} \text{ES} \xrightarrow{k_{\text{cat}}} \text{E} + \text{P}$$

It is described by two parameters: the enzyme-substrate dissociation (or Michaelis) parameter

$K_M \sim k_{\text{off}}/k_{\text{on}}$, and the catalytic rate constant (or turnover number) k_{cat}. Note that while K_M is

usually assumed to be invariable and is hence referred to as "Michaelis constant", we here

examine crowding-dependent changes in K_M und hence refer to it as the "Michaelis parameter".

415   We assume that all reactions follow effectively irreversible Michaelis Menten kinetics and thus

free energy changes are irrelevant to the transition rate k_{cat}; moreover, we ignore crowding

effects on the enzyme structure, and thus k_{cat} is not affected by crowding. To derive the effect of

crowding on the Michaelis parameter K_M, we divide a catalytic reaction into two independent,

consecutive steps: (1) S and E diffuse until they meet, and then (2) S and E bind and unbind

420   reversibly until the reaction proceeds forward to make P.

Step (1): the substrate-catalyst encounter

The encounter rate between S and E depends on the cytosolic occupancy. Its rate law is similar

to that of a diffusion-limited catalytic reaction, in which the ES encounter rate is low and their

encounter is the rate determining step; ys soon as the *ES* complex is formed, the reaction

425   quickly proceeds, converting *S* into *P* and releasing *E*. At given concentrations of *E* and *S*, the

rate of formation of *ES* is thus mostly determined by the rate of encounter, which is proportional

to the sum of *E*'s and *S*'s diffusion coefficients. A reduction of the diffusion rate shifts the

equilibrium between the concentrations of the individual molecules and the complex *ES*; this can

be accounted for in the following way (Minton, 1998):

$$\frac{[ES]}{[E][S]} = \frac{1}{K_M^{\text{diff}}} \simeq \frac{1}{K_M^0} \exp(-g(r_S)\rho) \quad ; \tag{S1}$$

430   here, K_M^{diff} is the hypothetical diffusion limited Michaelis parameter, while K_M^0 is the Michaelis

parameter in the low-crowding limit; ρ is the volume occupancy of the volume-excluding co-

solutes (dry mass) of the solution (with range $0 < \rho < 1$), and g is a function that depends on the

shape of E, S, and other volume-excluding co-solutes. Since S is typically much smaller than E,

the diffusion coefficient of S in a crowded solution is in general much higher than that of E.

435   Hence, we estimate this scaling term $exp\,(-g\rho)$ solely from the diffusion coefficient of S.

Approximating S as a sphere of radius r_S, we can write it as $exp\,(-g(r_S)\rho)$.

The bacterial cytosol is crowded, slowing down the diffusion of all molecular species. The extent

of this slow-down, however, is non-uniform and depends largely on the size of the affected

molecule: the larger the molecule, the more it is slowed down. The slow-down of diffusion in the

440   *E. coli* cytosol is summarized by the following empirical scaling law, which was inferred from

molecular dynamics simulations (Kalwarczyk et al., 2012):

$$ln\,(\frac{D_0(r_h)}{D_{cyto}(r_h)}) = (\frac{\xi^2}{R^2} + \frac{\xi^2}{r_h{}^2})^{-\frac{a}{2}} \tag{S2}$$

where $r_h = 1.3(r + 1.4\text{Å})$ is the hydrodynamic radius of a molecule with radius *r* (Trovato and

Tozzini, 2014), *i.e.*, its effective radius including the attached water molecules; $D_0(r_h)$ is the

diffusion coefficient in the low crowding limit, while $D_{cyto}(r_h)$ is the diffusion coefficient in the

445   crowded cytosol condition; *ξ*=0.51nm is the average distance between the surfaces of volume-

excluding co-solutes in the cytosol of *E. coli*; *R*=42nm is the radius of the largest common

crowders in the cytosol; and *a*=0.53 is an empirical scaling factor (Kalwarczyk et al., 2012). Note

that in general, the parameter ξ depends on ρ; however, as we are only interested in the relationship between reaction fluxes, growth rate, and ρ in a small range centered around the native *E. coli* cytosolic density, we can approximate ξ by a constant in our analysis. As the cytosolic dry mass density is ~0.3g/mL (Oldewurtel et al., 2021) and the mass-to-volume-ratio of protein is 1.35g/mL (Fischer et al., 2004), the cytosolic volume occupancy ρ is approximately 0.22=0.3/1.35. If the reaction rate is proportional to the rate of encounter between *E* and *S*, and this rate of encounter is in turn approximately proportional to the diffusion coefficient of *S*, then Eq. (S2) can be used to calculate the scaling factor $g(r_S)$:

$$g(r_S) = \left(\frac{\xi^2}{R^2} + \frac{\xi^2}{(1.3(r_S + 1.4\text{Å}))^2}\right)^{-\frac{a}{2}}/0.22$$

and the rate of this reaction is denoted as k_{diff}:

$$k_{\text{diff}} = k_{\text{cat}}[\text{ES}] = \frac{k_{\text{cat}}[\text{E}][\text{S}]}{K_{\text{M}}^{\text{diff}}}$$

Step (2): repeated binding and unbinding of substrate and catalyst

The rate law of this step is related to a transition-state limited catalytic reaction, in which the rate of encounter of E and S is much larger than the rate of conversion of the *ES* complex into the product P. In this case, the complex *ES* exists in near equilibrium with *E* and *S*, and macromolecular crowding affects the reaction rate parameter mainly through shifting this equilibrium. To assess the magnitude of this effect, we consider the reversible reaction

$$\text{E} + \text{S} \underset{k_{\text{off}}}{\overset{k_{\text{on}}}{\rightleftharpoons}} \text{ES}$$

and denote the adjusted equilibrium Michaelis parameter in this hypothetical transition-limited case as K_{M}^{ts} (Minton, 1998):

$$K_M{}^{ts} = \frac{\gamma_E \gamma_S [S][S]}{\gamma_{ES}[ES]} = \Gamma \frac{[E][S]}{[ES]} = \Gamma K_M{}^0,$$

465   where $\square_i$ denotes the activity coefficient of molecular species i, $\Gamma = \frac{\gamma_E \gamma_S}{\gamma_{ES}}$, and $K_M{}^0 = k_{off}/k_{on}$ is

the enzyme-substrate dissociation parameter in the low-crowding limit. The activity coefficients

are defined as

$$\square_i = exp\frac{\mu_i - \mu_i{}^{ideal}}{RT}$$

with

$$\mu_i = \frac{\partial G}{\partial [i]};$$

here, G is the Gibbs free energy of the system; [i] is the concentration of molecular species i; μ_i

470   is the chemical potential of i , and $\mu_i{}^{ideal}$ is the chemical potential in an idealized situation, i.e.,

without intermolecular interactions and in the absence of other volume-excluding co-solutes. In

other words, the equilibrium of the reaction will be identical to the dissociation parameter $K_M{}^0$ if

the system is ideal. The Γ term, therefore, accounts for the deviation of the Gibbs free energy

from the idealized situation.

475   The $\square_i$ of each molecular species i can be written as an expansion in terms of the concentrations

of all molecular species (McMillan and Mayer, 1945):

$$ln\,\gamma_i = \sum_j B_{ij}[j] + \sum_{j,k} B_{ijcok}[j][k]+...$$

The coefficients B_{ij} (B_{ijk}, ...) reflect the interaction between 2 (3, ...) molecular species. For

example, the coefficient B_{ij} is given as (Hirschfelder et al., 1964):

$$B_{ij} = 4\pi N_A \int_0^{inf} \{1 - exp\,[-\frac{U_{ij}(r)}{kT}]\}r^2\,dr$$

where N_A is Avogadro's number, r is the distance between the center of mass of molecular

480   species i and j, and $U_{ij}(r)$ is the potential of average force acting between the two molecular

species. While the interaction potential among multiple molecular species is complex, it has

been found that the colligative properties of solutions of globular proteins can be well accounted

for over a wide range of concentrations by using a simple hard sphere potential, where two

molecules cannot overlap but do not interact otherwise (see Ref (Minton, 1998) for a review):

$$U_{ij}(r) = \begin{cases} \infty & r \leq r_i + r_j \\ 0 & r > r_i + r_j \end{cases}$$

The scaled particle theory applies this rigid sphere potential to calculate the activity coefficient γ_i of molecular species i (Boublík, 1974):

$$\ln \gamma_i = -\ln(1 - \langle\langle V \rangle\rangle) + \frac{r_i \langle\langle S \rangle\rangle + S_i \langle\langle r \rangle\rangle + V_i \langle\langle 1 \rangle\rangle}{1 - \langle\langle V \rangle\rangle}$$
$$+ \frac{r_i^2 \langle\langle S \rangle\rangle^2 + 2 V_i \langle\langle r \rangle\rangle \langle\langle S \rangle\rangle}{2(1 - \langle\langle V \rangle\rangle)^2} + \frac{V_i \langle\langle r^2 \rangle\rangle \langle\langle S \rangle\rangle^2}{3(1 - \langle\langle V \rangle\rangle)^3} \qquad \text{(S3)}$$

Here, w_i is the concentration (number density) of molecular species i;   is a

weighted sum of property X, with   ;   is the surface area and

490   –   is the volume of molecular species i; in addition,   and

            . As in the hypothetical transition-state limited case, the reaction rate parameter is proportional to the concentration of the enzyme-substrate complex, and a shift of its equilibrium constant by   leads to a corresponding shift of the reaction rate, quantified by the rate parameter k_{ts} (Minton, 1998):

$$k_{ts} = k_{cat}[ES] = \frac{k_{cat}[E][S]}{K_M^{ts}} \qquad \text{(S4)}$$

495 **Combining step (1) and (2) for the general formulation**

We assume that a catalytic reaction is divided into two independent steps that occur in tandem: (1) the encounter of S and E, whose rate law is proportional to that of a diffusion limited reaction, and thereafter (2) their reversible binding and unbinding until S is converted into P, whose rate law is proportional to a transition-state limited reaction. The crowding-adjusted reaction rate k is

500 obtained by adding the inverse rate parameters (i.e., the reaction times) of the two steps (Aoki et al., 2013; Minton, 1990, 1998):

.

Given Eq. (S1) and (S4), this leads to (Aoki et al., 2013):

$$k = k_0 \frac{(1 + \theta)\Gamma \, exp \, (-g\rho)}{\Gamma + \theta \, exp \, (-g\rho)}$$

Here, $\theta \equiv \frac{k_{ts}}{k_{diff}}|_{\rho=0}$ quantifies the relative contributions of step (1) and (2) to the overall reaction, or in other words the ratio between times spent on step (1) and on (2) at low cytosolic

505   occupancy; the reaction tends towards diffusion limited if $\theta \rightarrow 0$, or transition-state limited if $\theta \rightarrow \infty$.

k_0 is the overall rate parameter in the absence of crowding. Remember that $[E_{free}] = [E_{total}] - [ES] = [E_{total}]/(1 + [S]/K_M)$ is the concentration of free enzymes only, and thus the reaction rate is $k = k_{cat}[ES] = k_{cat}[E_{free}][S]/K_M$; from now on, we describe the reaction rate as the flux of the reaction, and denote it as *v*. As we assume that the reaction is irreversible and hence

510   crowding does not influence k_{cat}, to arrive at crowding-adjusted Michaelis-Menten kinetics, we have to scale the Michaelis parameter, which arises from considerations on the equilibrium between the enzyme-substrate complex and its constituents. Thus, $v/v^0 = K_M^0/K_M^*$, with K_M^0 the Michaelis parameter in the low crowding limit; then the crowding-adjusted Michaelis parameter is

$$K_M^* = K_M^0 \frac{\Gamma + \theta \, exp \, (-g\rho)}{(1 + \theta)\Gamma \, exp \, (-g\rho)} \qquad\qquad \text{(S5)}$$

and so the flux can be written as

$$v = k_{cat} \frac{[E_{free}][S]}{K_M^*} \simeq k_{cat} \frac{[E_{total}][S]}{K_M^* + [S]} \; . \qquad\qquad \text{(S6)}$$

515   The θ of ERK MAP kinase phosphorylation reaction was estimated to be 2.3 (Aoki et al., 2013); we assume that this value is representative for cellular enzymes and use it for all modeled reactions.

The concentration of the enzyme-substrate complex, [*ES*], depends on K_M^* through $[ES] = $

520    $[E_{total}][S]/([S] + K_M^*)$; at the same time, K_M^* depends on [*ES*] through Eq. (S2). To find a self-consistent solution for these two quantities, we iterate these two equations until convergence. In each iteration, we first calculate the concentration of the catalyst-substrate complexes of different reactions; we then update their K_M^* and proceed to the next iteration. We stop the interaction when all K_M^* values changed by less than 0.001% compared to the previous iteration.

 Single-pathway model

We considered a simple model of a linear pathway to investigate how the size of the substrate

and catalyst of a metabolic reaction affect the optimal cytosolic occupancy; here, optimality is

defined as a maximal pathway flux per unit dry mass, calculated from crowding-adjusted

kinetics. The pathway is divided into N steps, where E_n is the catalyst of step n, which converts

530   its substrate s_n into s_{n+1}, the substrate of the next step (Fig. 2A). We assume that all internal

metabolite concentrations are in steady state (i.e., producing and consuming fluxes cancel

exactly), and we ignore the dilution of intermediates. Thus, all reaction fluxes have the same

value, v. We assume that s_1 is replenished by a flux $v_{s_1} = v$, which is not modeled explicitly.

535   We assume that the N reactions are described by crowding-adjusted Michaelis Menten kinetics

with identical k_{cat} and $K_M{}^0$. We further assume that the N catalyst species and the N substrate

species are spherical, with radius r_E (volume V_E) for the catalysts and radius r_s (volume V_s) for

the substrate species. These assumptions simplify the solution space of the model, as in the

optimal steady state, all catalysts have equal total concentrations ($[E_1]=[E_2]=...=[E_N]$, and so do

540   all the substrates ($[s_1]=[s_2]=...=[s_N]$). We define the total substrate concentration $[s] \equiv$

$\sum_n \quad [s_n] = N[s_1]$ and the total catalyst concentration $[E] \equiv \sum_n \quad [E_n] = N[E_1]$. The two

variables $[s]$ and $[E]$ span the solution space of this model. Substrates and catalysts, as well as

the substrate-catalyst complexes, are crowders in their own right; they slow down diffusion and

perturb Gibbs free energies. The cytosolic volume occupancy of dry mass in the solution, ρ

545    $(0 \leq \ \leq 1)$, is

$$= N_A \frac{4\pi}{3} ([s]r_s{}^3 + [E]r_E{}^3) \tag{S7}$$

with the Avogadro number N_A. The flux through the pathway per unit volume is

$$v = k_{cat} \frac{[s]/N\ [E]/N}{K_m{}^* + [s]/N} = \frac{k_{cat}}{N} \frac{[s][E]}{NK_m{}^* + [s]}$$

Accordingly, the flux per unit dry mass is

$$\mu = \frac{k_{cat}}{N} \frac{[s][E]}{NK_M{}^*+[s]} \frac{1}{\rho} \ .$$

(S8)

As we ignore crowding effects on the turnover number, k_{cat} acts only as a scaling factor, and we thus set k_{cat}=1 for simplicity. We set the Michaelis parameter at the low crowding limit to

550 $K_M{}^0$=130µM, which is the median K_M of metabolic enzymes (Bar-Even et al., 2011), and is also very close to the Michaelis constant of the ribosome estimated from the diffusion limit without molecular crowding (Klumpp et al., 2013).

Because we assume identical kinetics of all reactions and ignore the dilution of intermediates

555 through growth, the whole pathway is equivalent to a single reaction with re-scaled kinetics. We still describe it as an *N*-steps pathway, as this more faithfully reflects the situation in the real cell, allowing us to use realistic parameter values. Moreover, while mathematically, *N* represents a scaling factor, it has an intuitive biological interpretation.

560 The same equations can be used to describe a system of *N* parallel enzymatic reactions with identical fluxes, with only an additional multiplication by *N* in Eq. (S8), $\mu_{parallel}=\mu$ x *N* (Fig. 2B). Here, catalyst E_n of reaction *n* converts substrate s_n into the end product, and the consumption of s_n is compensated by a flux v_{s_n} that supplies s_n at an equal rate.

565 We consider two systems of substrate and catalyst sizes: metabolic and ribosomal. In the metabolic system, we use r_s=0.34nm for metabolites (the approximate radius of the amino acid alanine (Chwastyk et al., 2015)) and r_E=2.4nm for globular proteins (an "average" protein in *E. coli* has a mass 40kDa (Neidhardt, 1996), while a globular protein with mass 50kDa has a radius r=2.4nm (Erickson, 2009); in an alternative estimation, the radius of a typical globular protein is

570 approximately r=2.5nm (Roskams and Rodgers, 2002)). In the ribosomal system, we use r_s=2.4nm for the ternary complexes (the gyroscopic radius of tRNA is estimated to range from 2.33nm to 2.46nm based on Eq. (7) of (Hyeon et al., 2006)). We use a radius of r_E=13nm for the ribosome, as the diameter of a ribosome is reported to be 20nm - 30nm (Freitas, 2004; Kurland,

1960; Roskams and Rodgers, 2002). We assume that the catalyst-substrate complex is also

575 spherical, with a volume equal to the sum of the substrate's volume and the catalyst's volume.

For a fixed value of the total occupancy ρ, we calculated the specific flux μ using MATLAB, while (i) varying the occupancy ρ in steps of 0.01 from 0.01 to 0.8, with additional, finer-grained steps of 0.001 from 0.100 to 0.360, and (ii) varying the ratio of the volume occupied by the substrates

580 s, from 0.1% to 97.7%, with an increase by a factor of 1.0023 at each step.

Model cell with a metabolic and a ribosomal pathway

To more faithfully represent a complete cell and to study the tradeoff between metabolic and ribosomal reactions, we also consider a balanced growth model with a metabolic sector and a ribosomal sector (Fig. 2C). As seen from the results for the pathway models, the two types of

585 reactions have very different optimal conditions: the metabolic sector involves smaller catalysts and substrates than the ribosomal sector, and hence has maximal specific fluxes at a higher cytosolic occupancy.

In this model (Fig. 2C), the transporter T imports the external substrate s_{ext} into the cytosol,

590 where it is now labeled s_1. The transport reaction is described by ordinary, crowding-unaware irreversible Michaelis Menten kinetics, with flux

$$v_T = k_{cat}{}^T[s_{ext}][T]/([s_{ext}] + K_M{}^T).$$

We set $k_{cat}{}^T$=13.7s^{-1} (the median k_{cat} of enzyme reactions (Bar-Even et al., 2011)), and $K_M{}^T$=1µM (close to the growth limiting glucose concentration of *E. coli* (Shehata and Marr, 1971)).

The metabolic sector comprises an *N*-steps linear pathway of metabolic reactions, identical to the one studied in the simple pathway model: the enzyme of metabolic reaction n ($1 \leq n \leq N$), denoted as M_n, converts substrate s_n into s_{n+1}; in reaction $n=N$, M_N converts one s_N into one precursor p (Weiße et al., 2015). These *N* reactions follow crowding-adjusted irreversible

600 Michaelis Menten kinetics with identical rate parameters $K_M{}^{M0}$=130µM and $k_{cat}{}^M$=13.7s^{-1} (the

median K_M and k_{cat} of enzyme reactions (Bar-Even et al., 2011)). As before, all N substrates

have radius r_s=0.34nm, while all N enzymes have radius r_M=2.4nm; as before, we assume that

the enyzme-substrate complex is spherical and occupies as much volume as one substrate plus

one enzyme molecule. To facilitate the numerical solution of the model, we assume that all

605   metabolite concentrations $[s_n]$ and also all enzyme concentrations $[M_n]$ are identical. This

corresponds to the optimal balanced growth solution when the differential dilution of intermediate

metabolites is ignored (Bar-Even et al., 2011); thus, we treat the dilution of intermediate

metabolites only approximately here. In the balanced growth condition, the production rate of

each substrate is equal to its consumption rate plus its (approximate) rate of dilution through

610   growth, so that its concentration remains stable. We define the total substrate concentration

$[s] \equiv \sum_n \ [s_n]$ and total enzyme concentration $[M] \equiv \sum_n \ [M_n]$. The flux of each metabolic

reaction is $\frac{k_{cat}{}^M}{N} \frac{[s][M]}{N K_M{}^{M*}+[s]}$.

The ribosomal sector comprises the ribosome (R) and the protein precursor (p): R converts p

615   into the N+2 types of protein in the model, N metabolic enzymes (M_n), the ribosome (R), and the

transporter (T). As before, the radius of p is r_p=2.4nm and the radius of R is r_R=13nm; their

complex is assumed to be spherical and to occupy as much volume as one precursor plus one

ribosome molecule. The ribosomal conversion rate is described by crowding-adjusted

irreversible Michaelis Menten kinetics with parameters $K_M{}^{R0}$=120µM (Klumpp et al., 2013) and

620    $k_{cat}{}^R$=22.0s^{-1} (Klumpp et al., 2013; Weiße et al., 2015), and the consumption rate of p by R to

make proteins is $k_{cat}{}^R \frac{[p][R]}{K_M{}^{R*}+[p]}$. The ribosome converts l_T=300 precursors into one transporter,

l_M=300 precursors into one metabolic enzyme, and l_R=7459 precursors into one ribosome

(Weiße et al., 2015).

625   Note that the reactions that produce and consume the precursor p do not conserve volume. The

reason is that to model a realistic cell, we envision the precursor as a charged tRNA, only the

amino acid part of which is (i) produced by the metabolic pathway and (ii) integrated into the

growing protein. The metabolic pathway only provides the amino acid, while the pool of free

tRNAs is not explicitly modeled. For this reason, the size of p is substantially larger than the size

630   of the metabolite s_N consumed in its production. Conversely, the ribosome consumes 300

precursors to produce a single transporter or enzyme, which are both substantially smaller than

the combined volume of the 300 precursors. Here, we envision that the tRNA part of p is set free

again and can be re-charged through M_N. This treatment assumes that the concentration of free

tRNA is so low that we can ignore its dilution through growth and its contribution to the cytosolic

635   crowding.

The solution space of this model cell spans five dimensions: $[s]$, $[p]$, $[T]$, $[M]$, and $[R]$. The

corresponding molecules, along with their complexes, are also the crowders that slow down

diffusion and disturb Gibbs free energies. The cytosolic occupancy ρ, $0 \leq\ \leq 1$, is

$$ = N_A \times \frac{4\pi}{3}([s]r_s^3 + [M]r_M^3 + [p]r_p^3 + [R]r_R^3) \tag{S9}$$

640   with the Avogadro number N_A. The growth rate μ of this model cell can be expressed as the flux

through the ribosome reaction divided by the total protein concentration,

$$\mu = k_{cat}^R \frac{[p][R]}{K_M^{R*} + [p]} \frac{1}{N_A(l_M[M] + l_R[R] + l_T[T])} \tag{S10}$$

In the balanced growth state, the production of s and p is offset by their consumption and dilution

by growth,

$$k_{cat}^T \frac{[s_{ext}][T]}{K_M^T + [s_{ext}]} - \frac{k_{cat}^M}{N} \frac{[s][M]}{NK_M^{M*} + [s]} - \mu[s] = 0 \tag{S11}$$

$$\frac{k_{cat}^M}{N} \frac{[s][M]}{NK_M^{M*} + [s]} - k_{cat}^R \frac{[p][R]}{K_M^{R*} + [p]} - \mu[p] = 0$$

645   For a given number of enzymes N and occupancy ρ, we solved this model numerically. As a

preliminary step, we used the BARON algorithm (Sahinidis, 1996) implemented in Pyomo (Hart

et al., 2017, 2011) and assumed normal crowding-unaware irreversible MIchaelis Menten

kinetics, maximizing the growth rate μ over the space of concentrations ($[s],[p],[T],[M],[R]$),

subject to the constraints of Eq. (S9) and (S11). Using the BARON solution as a starting point,

650 we then applied the SLSQP algorithm within the function "*minimize*" in SciPy (Virtanen et al.,

2020), now using crowding-adjusted Michaelis-Menten kinetics. It is not clear if the optimization

problem has a unique solution; to increase the probability that the solution is a global maximum,

we repeatedly ran the SLSQP algorithm at least 20 times for each (N,ρ) combination, and picked

the solution with the highest μ. For each simulation, at least half of the independent runs

655 supported the same, maximal optimum.

Estimating the number of enzymes in the metabolic pathway

To obtain a realistic estimate of the number of simultaneously active metabolic reactions in a

bacterial cell, we performed flux balance analysis simulations accounting for molecular crowding

in terms of a hard limit on the total cellular protein concentration. Simulations were run using

660 "sybil", an R library for efficient constraint-based analyses (Desouki, 2016; Desouki et al., 2015;

Gelius-Dietrich et al., 2013). We used sybilccFBA (Alzoubi et al., 2019; Heckmann et al., 2018),

a re-implementation of the MOMENT algorithm with an improved treatment of multifunctional

enzymes, which maximizes the biomass production rate while constraining the sum of cytosolic

metabolic enzyme concentrations at the experimentally observed level.

We used a sybilccFBA implementation of the iAF1260 stoichiometric model (Feist et al., 2007),

parameterized with turnover numbers for *E. coli* (Alzoubi et al., 2019; Heckmann et al., 2018).

We considered four different nutritional environments. In each case, we counted the number of

active metabolic reactions with both substrates and products located in the cytosol, and with a

670 flux $>10^{-6}$ mmol/(gram Dry Weight)/h to filter out numerical noise; we enumerated the enzymes

supporting these reactions, filtering out those with proteome fraction lower than a cutoff (10^{-9}; for

reference, the most abundant enzyme has a density $\sim 10^{-3}$).

The estimated numbers of active metabolic enzymes and reactions in the different conditions are

675 as follows: (i) 259 active enzymes and 349 active reactions in a minimal medium with glucose as

the sole carbon source, corresponding to slow to intermediate growth with a cytosol dominated

by the metabolic sector (Alzoubi et al., 2019; Fuhrer et al., 2017) (Supplementary Table S1); (ii) 206 active enzymes and 288 active reactions in the same minimal glucose medium supplemented with 20 amino acids, simulating intermediate growth (Alzoubi et al., 2019; Fuhrer

680   et al., 2017) (Supplementary Table S2); (iii) 174 active enzymes and 250 active reactions in a rich medium, corresponding to fast growth and a cytosol dominated by the ribosome and its substrates (Alzoubi et al., 2019; Takeuchi et al., 2014) (Supplementary Table S3); and (iv) 140 active enzymes and 234 active reactions in an extremely rich medium, where all exchange reactions in the iAF1260 *E. coli* model are allowed to be active.

685   ***Conversion of experimental dry mass density to occupancy***

An empirical growth law connects the RNA/protein mass ratio (*r*) of an *E. coli* cell with its growth rate μ: $r = 0.087 + \mu/(4.5\text{h}^{-1})$ (Eq. (1) of Ref (Scott et al., 2010)). When μ increases from 0 to 0.7h[-1], *r* increases from 0.087 to 0.243. If we ignore all molecular species in the dry mass except protein and RNA (which constitute ~75% of the dry mass in an *E. coli* cell (Milo et al., 2010; Milo

690   and Phillips, 2015; Neidhardt et al., 1990)), and given the average density of protein (1.35 g/mL) (Fischer et al., 2004) and RNA (2.0g/mL) (Anderson et al., 1966), we find that the overall density (mass per volume of dry mass) increases from $1.35\frac{1}{1.087} + 2.0\frac{0.087}{1.087} = 1.40$ (g/mL) to $1.35\frac{1}{1.243} +$ $2.0\frac{0.243}{1.243} = 1.49$ (g/mL) when μ increases from ~0h[-1] to 0.7h[-1]. Oldewurthel et al. (Oldewurtel et al., 2021) found that up to these growth rates, the observed cellular dry mass density in *E. coli* is

695   approximately constant at ρ_{DM}=0.31g/mL. Accordingly, the empirical ρ decreases from $0.31 \div 1.36 = 0.221$ to $0.31 \div 1.49 = 0.208$, a 6% reduction.

The same set of experiments also showed that ρ_{DM} in *E. coli* decreases from 0.31g/mL to 0.28g/mL as μ increases further, from 0.7h[-1] to 1.2h[-1] (Oldewurtel et al., 2021). Within this range of μ, *r* increases from 0.243 to 0.354 (Scott et al., 2010), and the overall density of cytosolic dry

700   mass changes from 1.49g/mL to $1.35\frac{1}{1.354} + 2.0\frac{0.354}{1.354} = 1.52$ (Anderson et al., 1966; Fischer et al., 2004; Milo et al., 2010). Accordingly, the empirical occupancy ρ decreases from $0.31 \div 1.49 = 0.208$ to $0.28 \div 1.52 = 0.184$, an 11% decrease.

Acknowledgements

We thank Deya Alzoubi and David Heckmann for providing the environments used in the ccFBA simulations. We thank Hugo Dourado and Deniz Sezer for helpful discussions. This work was supported by DFG Grants CRC 1310 to T.Y.P. and M.J.L. and by a grant of the Volkswagen Foundation in the "Life?" initiative to M.J.L..

Author contributions

T.Y.P. developed the methodology and performed all analyses. M.J.L. and T.Y.P. conceived of the study, interpreted the results, and wrote the manuscript.

References

Adadi R, Volkmer B, Milo R, Heinemann M, Shlomi T. 2012. Prediction of Microbial Growth Rate versus Biomass Yield by a Metabolic Network with Kinetic Parameters. *PLOS Comput Biol* **8**:e1002575. doi:10.1371/journal.pcbi.1002575

Alzoubi D, Desouki AA, Lercher MJ. 2019. Flux balance analysis with or without molecular crowding fails to predict two thirds of experimentally observed epistasis in yeast. *Sci Rep* **9**:11837. doi:10.1038/s41598-019-47935-6

Anderson NG, Harris WW, Barber AA, Rankin CT, Candler EL. 1966. Separation of subcellular components and viruses by combined rate- and isopycnic-zonal centrifugation. *Natl Cancer Inst Monogr* **21**:253–283.

Aoki K, Takahashi K, Kaizu K, Matsuda M. 2013. A quantitative model of ERK MAP kinase phosphorylation in crowded media. *Sci Rep* **3**.

Bar-Even A, Noor E, Savir Y, Liebermeister W, Davidi D, Tawfik DS, Milo R. 2011. The Moderately Efficient Enzyme: Evolutionary and Physicochemical Trends Shaping Enzyme Parameters. *Biochemistry* **50**:4402–4410. doi:10.1021/bi2002289

Beg QK, Vazquez A, Ernst J, de Menezes MA, Bar-Joseph Z, Barabási A-LL, Oltvai ZN. 2007. Intracellular crowding defines the mode and sequence of substrate uptake by Escherichia coli and constrains its metabolic activity. *Proc Natl Acad Sci U S A* **104**:12663–12668. doi:10.1073/pnas.0609845104

Bobay L-M, Ochman H. 2018. Factors driving effective population size and pan-genome evolution in bacteria. *BMC Evol Biol* **18**:153. doi:10.1186/s12862-018-1272-4

Boublík T. 1974. Statistical thermodynamics of convex molecule fluids. *Mol Phys* **27**:1415–1427. doi:10.1080/00268977400101191

Chwastyk M, Bernaola AP, Cieplak M. 2015. Statistical radii associated with amino acids to determine the contact map: fixing the structure of a type I cohesin domain in the Clostridium thermocellum cellulosome. *Phys Biol* **12**:046002. doi:10.1088/1478-3975/12/4/046002

Dai X, Zhu M, Warren M, Balakrishnan R, Patsalo V, Okano H, Williamson JR, Fredrick K, Wang Y-P, Hwa T. 2016. Reduction of translating ribosomes enables Escherichia coli to maintain elongation rates during slow growth. *Nat Microbiol* **2**:1–9. doi:10.1038/nmicrobiol.2016.231

Desouki A. 2016. Algorithms for Improving the Predictive Power of Flux Balance Analysis

(Dissertation). Düsseldorf: Heinrich Heine University Düsseldorf.

Desouki AA, Jarre F, Gelius-Dietrich G, Lercher MJ. 2015. CycleFreeFlux: efficient removal of thermodynamically infeasible loops from flux distributions. *Bioinformatics* **31**:2159–2165. doi:10.1093/bioinformatics/btv096

Dill KA, Ghosh K, Schmit JD. 2011. Physical limits of cells and proteomes. *Proc Natl Acad Sci* **108**:17876–17882. doi:10.1073/pnas.1114477108

Dourado H, Lercher MJ. 2020. An analytical theory of balanced cellular growth. *Nat Commun* **11**:1226. doi:10.1038/s41467-020-14751-w

Erickson HP. 2009. Size and Shape of Protein Molecules at the Nanometer Level Determined by Sedimentation, Gel Filtration, and Electron Microscopy. *Biol Proced Online* **11**:32–51. doi:10.1007/s12575-009-9008-x

Feist AM, Henry CS, Reed JL, Krummenacker M, Joyce AR, Karp PD, Broadbelt LJ, Hatzimanikatis V, Palsson BØ. 2007. A genome-scale metabolic reconstruction for Escherichia coli K-12 MG1655 that accounts for 1260 ORFs and thermodynamic information. *Mol Syst Biol* **3**:121. doi:10.1038/msb4100155

Fischer H, Polikarpov I, Craievich AF. 2004. Average protein density is a molecular-weight-dependent function. *Protein Sci Publ Protein Soc* **13**:2825–2828. doi:10.1110/ps.04688204

Fisher RA. 1930. The Genetical Theory Of Natural Selection. Oxford, UK: Clarendon Press.

Freitas RA. 2004. Kinematic Self-Replicating Machines, 1st ed. Georgetown, Tex: CRC Press.

Fuhrer T, Zampieri M, Sévin DC, Sauer U, Zamboni N. 2017. Genomewide landscape of gene–metabolome associations in Escherichia coli. *Mol Syst Biol* **13**:907. doi:10.15252/msb.20167150

Gelius-Dietrich G, Desouki AA, Fritzemeier CJ, Lercher MJ. 2013. sybil – Efficient constraint-based modelling in R. *BMC Syst Biol* **7**:125. doi:10.1186/1752-0509-7-125

Hart WE, Laird CD, Watson J-P, Woodruff DL, Hackebeil GA, Nicholson BL, Siirola JD. 2017. Pyomo — Optimization Modeling in Python, 2nd ed, Springer Optimization and Its Applications. Springer International Publishing. doi:10.1007/978-3-319-58821-6

Hart WE, Watson J-P, Woodruff DL. 2011. Pyomo: modeling and solving mathematical programs in Python. *Math Program Comput* **3**:219. doi:10.1007/s12532-011-0026-8

Heckmann D, Lloyd CJ, Mih N, Ha Y, Zielinski DC, Haiman ZB, Desouki AA, Lercher MJ, Palsson BO. 2018. Machine learning applied to enzyme turnover numbers reveals protein structural correlates and improves metabolic models. *Nat Commun* **9**:5252. doi:10.1038/s41467-018-07652-6

Hirschfelder JO, Curtiss CF, Bird RB. 1964. The Molecular Theory of Gases and Liquids, Revised Auflage. ed. New York,NY: Wiley-Interscience.

Hu X-P, Dourado H, Schubert P, Lercher MJ. 2020. The protein translation machinery is expressed for maximal efficiency in Escherichia coli. *Nat Commun* **11**:5260. doi:10.1038/s41467-020-18948-x

Hui S, Silverman JM, Chen SS, Erickson DW, Basan M, Wang J, Hwa T, Williamson JR. 2015. Quantitative proteomic analysis reveals a simple strategy of global resource allocation in bacteria. *Mol Syst Biol* **11**:784. doi:10.15252/msb.20145697

Hyeon C, Dima RI, Thirumalai D. 2006. Size, shape, and flexibility of RNA structures. *J Chem Phys* **125**:194905. doi:10.1063/1.2364190

Kalwarczyk T, Tabaka M, Holyst R. 2012. Biologistics–diffusion coefficients for complete proteome of Escherichia coli. *Bioinforma Oxf Engl* **28**:2971–2978.

Kihara T. 1953. Virial Coefficients and Models of Molecules in Gases. *Rev Mod Phys* **25**:831–843. doi:10.1103/RevModPhys.25.831

Klumpp S, Scott M, Pedersen S, Hwa T. 2013. Molecular crowding limits translation and cell growth. *Proc Natl Acad Sci* **110**:16754–16759. doi:10.1073/pnas.1310377110

Knapp BD, Odermatt P, Rojas ER, Cheng W, He X, Huang KC, Chang F. 2019. Decoupling of Rates of Protein Synthesis from Cell Expansion Leads to Supergrowth. *Cell Syst* **9**:434-445.e6. doi:10.1016/j.cels.2019.10.001

Kurland CG. 1960. Molecular characterization of ribonucleic acid from Escherichia coli
ribosomes: I. Isolation and molecular weights. *J Mol Biol* **2**:83–91. doi:10.1016/S0022-2836(60)80029-0

Kuznetsova IM, Turoverov KK, Uversky VN. 2014. What macromolecular crowding can do to a
protein. *Int J Mol Sci* **15**:23090–23140.

Lee B. 1983. Calculation of volume fluctuation for globular protein models. *Proc Natl Acad Sci U
S A* **80**:622–626.

McMillan WG, Mayer JE. 1945. The Statistical Thermodynamics of Multicomponent Systems. *J
Chem Phys* **13**:276–305. doi:10.1063/1.1724036

Milo R, Jorgensen P, Moran U, Weber G, Springer M. 2010. BioNumbers—the database of key
numbers in molecular and cell biology. *Nucleic Acids Res* **38**:D750–D753.
doi:10.1093/nar/gkp889

Milo R, Phillips R. 2015. Cell Biology by the Numbers, Illustrated edition. ed. New York, NY:
Taylor & Francis.

Minton A. 1990. Structural and Organizational Aspects of Metabolic Regulation; Chapter
Holobiochemistry: An integrated Approach to the Understanding of Biochemical
Mechanisms That Emerges from the Study of Proteins and Protein Associations in
Volume-Occupied Solutions. Wiley: Hoboken, NJ, USA.

Minton AP. 1998. Molecular crowding: analysis of effects of high concentrations of inert
cosolutes on biochemical equilibria and rates in terms of volume exclusion. *Methods
Enzymol* **295**:127–149.

Minton AP. 1981. Excluded volume as a determinant of macromolecular structure and reactivity.
Biopolym Orig Res Biomol **20**:2093–2120.

Molenaar D, van Berlo R, de Ridder D, Teusink B. 2009. Shifts in growth strategies reflect
tradeoffs in cellular economics. *Mol Syst Biol* **5**. doi:10.1038/msb.2009.82

Neidhardt FC, editor. 1996. Escherichia coli and Salmonella: Cellular and Molecular Biology, 2nd
ed. Washington, D.C: ASM Press.

Neidhardt FC, Curtiss R. 1996. Escherichia Coli and Salmonella (2 Volume Set: Cellular and
Molecular Biology, 2nd ed. Washington, D.C: American Society for Microbiology.

Neidhardt FC, Ingraham JL, Schaechter M. 1990. Physiology of the Bacterial Cell: A Molecular
Approach. Sunderland, Mass: Sinauer Associates Inc.

Nierlich DP. 1972. Regulation of ribonucleic acid synthesis in growing bacterial cells: II. Control
over the composition of the newly made RNA. *J Mol Biol* **72**:765–777. doi:10.1016/0022-2836(72)90190-8

Norris MG, Malys N. 2011. What is the true enzyme kinetics in the biological system? An
investigation of macromolecular crowding effect upon enzyme kinetics of glucose-6-phosphate dehydrogenase. *Biochem Biophys Res Commun* **405**:388–392.

Oldewurtel ER, Kitahara Y, Teeffelen S van. 2021. Robust surface-to-mass coupling and turgor-dependent cell width determine bacterial dry-mass density. *Proc Natl Acad Sci* **118**.
doi:10.1073/pnas.2021416118

Olsen SN. 2006. Applications of isothermal titration calorimetry to measure enzyme kinetics and
activity in complex solutions. *Thermochim Acta* **448**:12–18. doi:10.1016/j.tca.2006.06.019

Pozdnyakova I, Wittung-Stafshede P. 2010. Non-linear effects of macromolecular crowding on
enzymatic activity of multi-copper oxidase. *Biochim Biophys Acta BBA - Proteins
Proteomics* **1804**:740–744. doi:10.1016/j.bbapap.2009.11.013

Rabl J, Leibundgut M, Ataide SF, Haag A, Ban N. 2011. Crystal Structure of the Eukaryotic 40S
Ribosomal Subunit in Complex with Initiation Factor 1. *Science* **331**:730–736.
doi:10.1126/science.1198308

Ralston GB. 1990. Effects of" crowding" in protein solutions. *J Chem Educ* **67**:857.

Roskams J, Rodgers L, editors. 2002. Lab Ref: A Handbook of Recipes, Reagents, and Other
Reference Tools for Use at the Bench. Cold Spring Harbor, NY: Cold Spring Harbor
Laboratory Press.

Sahinidis NV. 1996. BARON: A general purpose global optimization software package. *J Glob Optim* **8**:201–205. doi:10.1007/BF00138693

Schmidt A, Kochanowski K, Vedelaar S, Ahrne E, Volkmer B, Callipo L, Knoops K, Bauer M, Aebersold R, Heinemann M. 2015. The quantitative and condition-dependent Escherichia coli proteome. *Nat Biotechnol* **34**:104–110. doi:10.1038/nbt.3418

Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T. 2010. Interdependence of Cell Growth and Gene Expression: Origins and Consequences. *Science* **330**:1099–1102. doi:10.1126/science.1192588

Sezer D, Schubert P, Lercher MJ. 2020. Structure of autosynthetic models of balanced cell growth and numerical optimization of their growth rate. *bioRxiv* 2020.09.19.304998. doi:10.1101/2020.09.19.304998

Shehata TE, Marr AG. 1971. Effect of Nutrient Concentration on the Growth of Escherichia coli. *J Bacteriol* **107**:210–216.

Takeuchi R, Tamura T, Nakayashiki T, Tanaka Y, Muto A, Wanner BL, Mori H. 2014. Colony-live — a high-throughput method for measuring microbial colony growth kinetics— reveals diverse growth effects of gene knockouts in Escherichia coli. *BMC Microbiol* **14**:171. doi:10.1186/1471-2180-14-171

Totani K, Ihara Y, Matsuo I, Ito Y. 2008. Effects of macromolecular crowding on glycoprotein processing enzymes. *J Am Chem Soc* **130**:2101–2107. doi:10.1021/ja077570k

Trovato F, Tozzini V. 2014. Diffusion within the cytoplasm: a mesoscale model of interacting macromolecules. *Biophys J* **107**:2579–2591.

Vazquez A. 2010. Optimal cytoplasmatic density and flux balance model under macromolecular crowding effects. *J Theor Biol* **264**:356–359. doi:10.1016/j.jtbi.2010.02.024

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, Polat İ, Feng Y, Moore EW, VanderPlas J, Laxalde D, Perktold J, Cimrman R, Henriksen I, Quintero EA, Harris CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P. 2020. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* **17**:261–272. doi:10.1038/s41592-019-0686-2

Weiße AY, Oyarzún DA, Danos V, Swain PS. 2015. Mechanistic links between cellular trade-offs, gene expression, and growth. *Proc Natl Acad Sci* **112**:E1038–E1047. doi:10.1073/pnas.1416533112

Woldringh CL, Nanninga N. 1985. Structure of the Nucleoid and Cytoplasm in the Intact CellMolecular Cytology of Escherichia Coli. London; Orlando: Academic Press Inc. pp. 161–197.

Zimmerman SB, Minton AP. 1993. Macromolecular crowding: biochemical, biophysical, and physiological consequences. *Annu Rev Biophys Biomol Struct* **22**:27–65.