MLL1 is required for maintenance of intestinal stem cells and the

expression of the cell adhesion molecule JAML

Neha Goveas[1], Claudia Waskow[2,#a,#b], Kathrin Arndt[2], Julian Heuberger[3,#c],

Qinyu Zhang[1], Dimitra Alexopoulou[4], Andreas Dahl[4], Walter Birchmeier[3],

Konstantinos Anastassiadis[5], A. Francis Stewart[1,6*], Andrea Kranz[1*]

[1] Genomics, Center for Molecular and Cellular Bioengineering, Biotechnology

Center, Technische Universität Dresden, Dresden, Germany

[2] Institute for Immunology and Department of Medicine III, Technische

Universität Dresden, Dresden, Germany

[3] Laboratory of Signal Transduction in Development and Cancer, Max

Delbrück Center for Molecular Medicine, Berlin, Germany

[4] DRESDEN-concept Genome Center, Center for Molecular and Cellular

Bioengineering, Technische Universität Dresden, Dresden, Germany

[5] Stem Cell Engineering, Center for Molecular and Cellular Bioengineering,

Biotechnology Center, Technische Universität Dresden, Dresden, Germany

[6] Max Planck Institute of Molecular Cell Biology and Genetics, Dresden,

Germany

[#a]Current Address: Regeneration in Hematopoiesis, Leibniz-Institute on

Aging, Fritz-Lipmann-Institute (FLI), Jena, Germany

[#b]Current Address: Institute of Biochemistry and Biophysics, Faculty of

Biological Sciences, Friedrich-Schiller-Universität Jena, Jena, Germany

#cCurrent Address: Department of Hepatology and Gastroenterology, Charité

University Medicine, Berlin, Germany

* Corresponding authors

E-mail: francis.stewart@tu-dresden.de (AFS)

E-mail: andrea.kranz@tu-dresden.de (AK)

Abstract

Epigenetic control is crucial for lineage-specific gene expression that creates cellular identity during mammalian development and in adult organism. Histone 3 lysine 4 methylation (H3K4) is a universal epigenetic mark. Mixed lineage leukemia (MLL1) is the founding member of the mammalian family of H3K4 methyltransferases. It was originally discovered as the main gene mutated in early onset leukemias and then found to be required for hematopoietic stem cell development and maintenance. However, the roles of MLL1 in non-hematopoietic tissues remain largely unexplored. To bypass hematopoietic lethality, we used bone marrow transplantation and conditional mutagenesis to discover that the most overt phenotype in *Mll1*-mutant mice is intestinal failure. Loss of MLL1 is accompanied by a differentiation bias towards the secretory lineage with increased numbers of goblet cells. MLL1 is expressed in intestinal stem cells (ISCs) and transit amplifying (TA) cells but at reduced levels in Paneth cells and not in the villus. MLL1 is required for the maintenance of intestinal stem cells (ISCs) and proliferation in the crypt. Transcriptome analysis implicate MLL1-dependent expression in ISCs of several transcription factors including *Pitx2, Gata4, Foxa1* and *Onecut2*, and also a cell adhesion molecule, *Jaml*. Reactive transcriptome changes in Paneth cells and organoids imply that JAML plays a key role in the crypt stem cell niche. All known postnatal functions of MLL1 relate to stem cell maintenance and lineage decisions thereby highlighting the suggestion that MLL1 is a master stem cell regulator.

Author Summary

The ability of adult stem cells to produce functional progenies through differentiation is critical to maintain function and integrity of organs. A fundamental challenge is to identify factors that control the transition from self-renewal to the differentiated state. Epigenetic factors amongst others can fullfill such a role. Methylation of histone 3 on lysine 4 (H3K4) is a posttranslational epigenetic modification that is associated with actively transcribed genes. In mammals, this epigenetic mark is catalyzed by one of six H3K4 methyltransferases, including the founding member of the family, MLL1. MLL1 is important for the precise functioning of the hematopoietic stem cell compartment. This raises the possibility of similar functions in other adult stem cell compartments. Due to its intense self-renewal kinetics and its simple repetitive architecture, the intestinal epithelium serves as a prime model for studying adult stem cells. We demonstrate that MLL1 controls intestinal stem cell proliferation and differentiation. Additionally, transcriptome analysis suggests a pertubation in the close interaction between intestinal stem cells and neighbouring Paneth cells through loss of junction adhesion molecule like (JAML). Our work sheds new light on the function of MLL1 for the control of intestinal stem cell identity.

Introduction

Stem cells are cornerstones of tissue biology, ensuring homeostasis and regeneration in many organs, including epithelial tissues such as skin, intestine and mammary gland [1]. Stem cells are characterized by multipotency, which is the ability to differentiate into a restricted number of defined cell types, and self-renewal, which is the capacity to undergo infinite replicative cycles without losing stem cell identity [2]. The remarkable capacities of stem cells, particularly the restricted specificities of multipotency, rely on interplays between specific transcription factors and distinct epigenetic landscapes. Whereas the transcription factors involved in stem cell maintenance and differentiation have been clearly defined, epigenetic contributions are proving more elusive. For example, the transcription factor hierarchies in the stem cell paradigm, hematopoiesis, have been elegantly dissected [3]. However the contributions of DNA and histone methyltransferases to hematopoiesis are still emerging and indicate both specificities and the deeper complexities of epigenetic regulation[4-8].

Methylation of histone 3 on lysine 4 (H3K4) is one of the most conserved and widespread epigenetic systems [9]. H3K4 is methylated in euchromatic regions, with trimethylated H3K4 (H3K4me3) on nucleosomes surrounding active promoters, H3K4me2 marking transcribed regions and H3K4me1 relating to enhancers and active chromatin in general [10-14]. Mammals have six Set1/Trithorax-related methyltransferases that are encoded by three pairs of paralogous sister genes namely, *Mll1* (*Kmt2a*) and *Mll2* (*Kmt2b*), *Mll3* (*Kmt2c*) and *Mll4* (*Kmt2d*), *Setd1a* (*Kmt2f*) and *Setd1b*

(*Kmt2g*). Each of the six methyltransferases reside in their own, large, protein

complex. However all six complexes are based on a four membered scaffold

termed WRAD for the subunits WDR5, RBBP5, ASH2L and DPY30 [15].

Functional differences between the six complexes potentially arise from the

presence of additional subunits, which are usually shared by paralogous pairs

or sometimes uniquely found in one of the six complexes.

Mixed lineage leukemia (*MLL1*) was the first mammalian gene

identified as a Trithorax homologue and subsequently found to encode a

mammalian Set1/Trithorax-type H3K4 methyltransferase (HMT) [16, 17]. In

mice, MLL1 is first required at embryonic day 12.5 (E12.5) for definitive

hematopoiesis [18, 19] and also required for the maintenance of adult

hematopoietic stem cells (HSCs) [20, 21]. *MLL1*, but not its paralogue, *MLL2,*

is a proto-oncogene because it can be activated by chromosomal

translocations to promote leukemias without additional mutagenesis [22, 23].

Over 80 translocation partners have been identified including AF6 and AF9

[24]. Notably, *MLL1-AF6* and *-AF9* leukemias rely on *Mll2* expression [6].

Mouse studies also indicated that *MLL1-AF9* leukemiogenesis is entirely

conveyed by overexpression of *Hoxa9* [25]. Conditional mutagenesis has also

revealed MLL1 functions in satellite cells [26] and postnatal neural stem cells

(NSCs) [27]. These observations raise the possibility that MLL1 regulates

specific functions in stem cell compartments.

Due to its high turnover and hierarchical architecture, intestinal stem

cells (ISCs) in the intestinal epithelium have become an adult stem cell

paradigm. ISCs have been identified as either actively cycling crypt base

columnar cells (CBCs) or quiescent label-retaining cells (LRCs) located at the

+4 position from the crypt base [28]. Leucine-rich repeat-containing G protein-

coupled receptor 5 (LGR5) is one of the best-characterized markers for the

CBC class ISCs [29]. They generate transit amplifying (TA) daughter cells that

give rise to the terminally differentiated progenies: absorptive enterocytes,

secretory goblet, enteroendocrine and Paneth cells [30]. Except for Paneth

cells, these cell types take 3 to 5 days to migrate up the villi and are shed into

the intestinal lumen. Paneth cells reside at the base of the crypts in close

association with ISCs and turn over at a slower rate [30].

The quiescent LRCs are active only during stress or injury. They

represent a stem cell reserve to replace damaged ISCs [31]. Enterocytes,

preterminal enteroendocrine cells, goblet cell precursors and Dll1[+] secretory

progenitors are also notable for their plasticity and can serve as a reservoir for

lost stem cells [32, 33] [34]. Mature Paneth cells also show an injury-activated

conversion to a stem cell like state [35]. Together with new perceptions in

hematopoiesis [4, 6, 36], the dynamic plasticity of the intestinal crypt has

expanded the stem cell paradigm [37], especially during replenishment after

damage or inflammation.

To examine whether MLL1 plays additional roles in the adult, we

employed ligand-induced conditional mutagenesis using a tamoxifen-inducible

Rosa26-CreERT2 line (*RC*) for near-ubiquitous Cre recombination to discover

that MLL1 is required for the maintenance of the ISC compartment and the

balance between secretory and absorptive cell lineages in the adult intestine.

Results

Intestinal functions collapse after loss of MLL1 in adult mice

To explore MLL1 functions, we utilized a multipurpose allele that can be converted from one state to another using FLP and Cre recombination (S1A-S1C Figs) [38, 39]. Homozygous embryos carrying the targeted allele, $Mll1^{A/A}$, developed normally until E12.5 when they displayed pallor of the liver and were smaller (S1D Fig). After E13 no live $Mll1^{A/A}$ embryos were found (S1 Table). After FLP and Cre recombination, $Mll1^{FC/FC}$ embryos displayed the same phenotype indicating that both A and FC are true null alleles in concordance with the loss of MLL1 protein [40]. Furthermore the $Mll1^{A/A}$ and $Mll1^{FC/FC}$ phenotype recapitulated another likely null allele [6] with all three homozygous mice presenting the same embryonic lethality due to the failure to engage definitive hematopoiesis [20].

To identify postnatal roles of MLL1, conditional mutagenesis using *Rosa26-CreERT2* was applied to 2 month old adults. As expected, mice lacking MLL1 developed severe bone marrow cytopenia and died or had to be sacrificed on average within two weeks (Figs 1A and 1B). As previously reported using the same conditional allele, *in vitro* deletion of *Mll1* in KSL-enriched HSCs from $Mll1^{F/F; RC/+}$ mice resulted in significant downregulation of *Hoxa9*, *Meis1*, *Mecom/Evi1* and *Prdm16* [6].

To bypass the bone marrow related lethality and thereby uncover non-hematopoietic phenotypes, bone marrow from wild type (wt) B6.SJL mice was transplanted into lethally irradiated $Mll1^{F/+; RC/+}$ or $Mll1^{F/F; RC/+}$ mice. After stable engraftment tamoxifen gavage induced widespread Cre-mediated excision of

Mll1 (Fig 1C). Examination of the bone marrow confirmed the successful and

near-complete reconstitution of the hematopoietic stem cell compartment by

wt donor cells. FACS analysis for KSL-Slam enriched hematopoietic stem

cells showed comparable frequencies in bone marrow transplanted (BMTx)

Mll1$^{FC/+;\ RC/+}$ and *Mll1*$^{FC/FC;\ RC/+}$ mice with the hematopoietic compartment

comprised only of wt cells of CD45.1 origin (Fig 1C). Notably, the BMTx

Mll1$^{FC/FC;\ RC/+}$ mice suffered from diarrhea and wasting (Fig 1D). These data

indicate that MLL1 is not only required in the hematopoietic compartment but

also elsewhere and this additional requirement is similarly critical for survival

as the hematopoietic requirement.

The small and large intestine harbor their stem cell compartments at

the base of the crypt. MLL1 is strongly expressed at the crypt bottom and in

the TA compartment whereas it is absent in differentiated cells above the TA

compartment (Figs 1E and S1E). RNA profiling of sorted ISCs and Paneth

cells confirmed expression of *Mll1* and the other family members in both cell

types however more strongly in ISCs (S1F Fig). In the mutant small intestine

of BMTx *Mll1*$^{FC/FC;\ RC/+}$ mice, expression of MLL1 was efficiently ablated (Fig

2A). Loss of stem cell markers olfactomedin (OLFM4) and SOX9 suggested

depletion of ISCs in the small intestine of BMTx *Mll1*$^{FC/FC;\ RC/+}$ mice (Fig 2A).

Consistent with this, the mutant showed decreased proliferation in the crypt as

demonstrated by strongly reduced expression of the mitotic marker Ki67 (Fig

2A). However, the intestinal epithelium of *Mll1*$^{FC/FC;\ RC/+}$ mice revealed no

apparent change in global H3K4 mono-, di- and trimethylation (S1G Fig).

Shortened villi with distorted morphology including vacuolar structures

at the tip indicated diminished replenishment of cells into the villus (Fig 2B).

Furthermore we observed increased numbers of enlarged goblet cells

distributed irregularly along the villus and also ectopically in the crypt (Fig 2B).

However, Paneth cells appeared unchanged possibly due to their longer life

span (Fig 2C). Similarly, as evaluated by chromogranin A and alkaline

phosphatase, the enteroendocrine and absorptive lineages appeared to be

unaffected (Fig 2C).

Without bone marrow rescue, $Mll1^{FC/FC; RC/+}$ mice showed the same

defects with depletion of ISCs, decreased proliferation and a distortion of the

secretory lineage (S2A-S2C Figs). Differentiation into the enteroendocrine

and absorptive lineage was also apparently unaffected (S2D Fig). Notably the

hallmark of Wnt signalling, nuclear β-catenin, was also unaffected (S2E Fig).

Intestinal epithelium-specific *Mll1* conditional mutagenesis recapitulates

ubiquitous deletion

To delete *Mll1* exclusively in the adult intestine we employed the

tamoxifen-inducible gut epithelium-specific *Villin-CreERT2* strain [41]. After

tamoxifen administration $Mll1^{FC/FC; Vil-CreERT2/+}$ mice lost weight compared to

control mice (Fig 3A). In agreement with our observations after ubiquitous

deletion of *Mll1,* OLFM4, SOX9 and proliferation were markedly decreased,

goblet cells were increased whereas Paneth and enteroendocrine cell

numbers were unchanged (Figs 3B and 3C).

Transcriptome analysis identifies key intestine specific transcription

factors and *Jaml* as central to the crypt stem cell niche

For transcriptome analysis, ISCs and Paneth cells were isolated 4 and

10 days after tamoxifen administration from *Mll1*$^{FC/+;\ Lgr5-eGFP-CreERT2/+}$ and

Mll1$^{FC/FC;\ Lgr5-eGFP-CreERT2/+}$ littermates (S3 Fig). Using 75 base-pair reads, 20-

37 million reads per sample with high levels of uniqueness (70-77% in Lgr5$^+$

ISCs and 60-76% in Paneth cells; S4A-S4C Figs) and comparable

mappability (99%) were obtained. Principal component analysis (PCA)

revealed that our datasets are in good agreement with published datasets

obtained from sorted ISCs and Paneth cells [35, 42] (S4D Fig).

We applied DESeq2 to analyze differentially expressed genes (DEGs).

For the 4 day Lgr5$^+$ stem cell profile, only 87 and 49 genes were up- or

downregulated at a 5% false discovery rate (FDR) after removal of MLL1 (Fig

4A, Supplementary excel file 1). However, none of the upregulated transcripts

were increased by more than log2-fold and by DAVID analysis were mainly

related to diverse terms such as 'response to metal ion', 'organic acid

metabolic process' and 'regulation of lipid metabolic process' (Fig 4B). In

contrast the most significant terms associated with the downregulated mRNAs

were 'regulation of gene expression', 'epithelial cell differentiation' and 'cell

proliferation' (Fig 4B). The 10 day profile revealed 179 DEGs, of which 105

were upregulated, 74 were downregulated with significant overlaps to the 4

day profile (Supplementary excel file 1, Table 1).

Table 1. Downregulated mRNAs common to both 4 and 10 day stem cell profiles.

Gene		4d C	KO	FC	10d C	KO	FC
Jaml/Amica1	Junction Adhesion Molecule Like	4124	119,9	**34,4**	3779	113,4	**33,3**
Pitx2	Paired Like Homeodomain 2	1399,6	162,9	**8,6**	854,8	34,7	**24,6**
Far1	FattyAcyl-CoA Reductase 1	402,3	229,1	**1,8**	231,8	47,4	**4,9**
Foxa1/ Hnf3α	Forkhead Box A1	356,8	105	**3,4**	428,4	93,9	**4,6**
Onecut2/ Hnf6β	Onecut Homeobox 2	2334,5	1177,1	**2,0**	2411,9	583	**4,1**
Gata4	GATA Binding Protein 4	676,4	356,9	**1,9**	508,2	136,7	**3,7**
Ces2g	Carboxyl esterase 2	275,2	177	**1,6**	209,9	58,6	**3,6**
Pla2g2a	Phospholipase A2 group IIA	291,3	119	**2,5**	215,7	68,2	**3,2**
Hspa8	Hsp 70 family	12104	4655,8	**2,6**	7654,5	25856	**3,0**
E230029C05	ncRNA	221,9	115,8	**1,9**	203,4	70,9	**2,9**
Casc4	Cancer Susceptible 4	120,5	61,4	**2,0**	255,1	93,9	**2,7**
Anpep	Alanyl amino peptidase N	855,8	562,5	**1,5**	393,1	150,8	**2,6**
Defa29	Defensinα 29	859,4	556,3	**1,5**	860,3	336,2	**2,6**
Lyz1	Lysozyme	10861	7181,7	**1,5**	8334,4	3429	**2,4**
Defa3	Defensinα 3	342,6	239,5	**1,4**	521,2	214,7	**2,4**
Lcp1	Lymphocyte Cytosolic P1	824,4	576	**1,4**	662	291	**2,3**
Defa35	Defensinα 35	831,4	555,4	**1,5**	938,4	431,4	**2,2**
Gm10925	pseudogene?	18872	14115	**1,3**	30453	14252	**2,1**
Afap1l1	Lnc-Afap1/1	485,5	354,5	**1,4**	420,6	197	**2,1**
Phlpp1	PH Domain Leucine Rich Phosphatase	468,7	302,8	**1,6**	480,5	227	**2,1**
Fap1	Familial Adenomatous Polyposis 1	990,5	768,4	**1,3**	1459,8	690,9	**2,1**
Casp6	Caspase 6	4727,6	3356,6	**1,4**	3861,6	1835	**2,1**
mt-Atp6	Mitochondrial encoded ATP Synthase S6	49048	40226	**1,2**	61873	29894	**2,1**
Fam149a	Toll-like receptor 3?	170,7	97,6	**1,8**	193,9	94,3	**2,1**
Zfpm1	FOG1 (Friend of Gata 1)	198,5	146	**1,4**	216,6	106,2	**2,0**

mRNAs expressed more than 150 reads in wt and downregulated more than 2 fold in the 10 day profile are shown. FC, fold change.

Gene set enrichment analysis (GSEA) of both 4 and 10 day ISC

profiles revealed that overall ISC signature genes were downregulated whereas goblet cell signature genes were upregulated (Figs 4C-4G; S5A and S5B Figs), which concords with our immunohistochemical analyses. We focused on the overlap between the 4 and 10 day downregulated mRNAs. The transcription factors *Pitx2, Foxa1*, *Onecut2* and *Gata4* are prominent (Table 1; Fig 4E). The expression of *Foxa1* and *Pitx2* and was evaluated by qRT-PCR and good agreement to the RNA-sequencing mRNA reads was found (S5C Fig). These data suggest a role for MLL1 in maintaining the transcriptional identity of ISCs.

However the top downregulated gene in both stem cell profiles (34.7 and 33.3 fold down) was junction adhesion molecule like (*Jaml* or *Amica1*; Figs 4A, 4F, S5C) previously identified as an ISC signature gene [43, 44]. It is highly expressed in Lgr5$^+$ ISCs, where it appears to rely completely on MLL1, but not in Paneth cells (at least 55 fold lower expressed in Paneth cells; S5D Fig). JAML is a 65 kDa type I transmembrane glycoprotein in the JAM subset of the immunoglobulin superfamily. JAML mediates adhesion of monocytes to endothelial cells and neutrophil migration across epithelial cell monolayers through interaction with Coxsackie and adenovirus receptor (CXADR or CAR) in tight junctions [45]. However, the cognate receptor in the intestinal epithelium is unknown. Notably the transcript for CAR-like soluble protein (*Clsp*) (GM1123) is upregulated in both 4 and 10 day profiles (Supplementary excel file 1). CLSP is closely related to CXADR however it lacks a transmembrane domain [46].

Loss of MLL1 in ISCs provokes transcriptional changes in Paneth cells

ISCs are anchored in the crypt in close association with Paneth cells.

In order to elucidate whether the transcriptional changes in ISCs influenced

the neighboring Paneth cells, we also analyzed Paneth cell transcriptional

profiles 4 days after deletion of *Mll1* in Lgr5$^+$ ISCs, with 198 and 72 transcripts

up- and downregulated respectively (Fig 5A, Supplementary excel file 1). The

most significant terms associated with downregulated mRNAs relate to

perturbation of protein folding and homeostasis in the endoplasmic reticulum

(Figs 5B-5D). In contrast upregulated mRNAs associate with metabolic

changes. Strikingly, transcripts of genes belonging to all five of the respiratory

chain complexes were upregulated (Figs 5B-5D). Paneth cells normally run on

glycolysis with lactate as the end product whereas ISCs depend on

mitochondrial oxidative phosphorylation [47]. These transcriptional changes

suggest that loss of MLL1 in the stem cell compartment provokes changes in

Paneth cells and indeed expression of Paneth cell marker genes was

downregulated (Figs 5C and 5D).

Skewed differentiation of organoids after loss of MLL1

To evaluate the cell-intrinsic requirement of MLL1 in the small intestine,

we isolated crypts from *Mll1*$^{F/+; RC/+}$ and *Mll1*$^{F/F; RC/+}$ mice and cultured them to

form organoids. After passaging, organoids were induced with 4-OH

tamoxifen for 24 hours on day 2. After further passages, the *Mll1*$^{FC/FC; RC/+}$

organoids increasingly formed round, less differentiated cyst-like spheres

(Figs 6A-6C). qRT-PCR revealed mRNA downregulation of *Jaml* and

transcription factors such as *Pitx2 and Foxa1* (Fig 6D) indicating the expected

loss of ISCs and elevation of the goblet cell marker, *Muc2*. Notably the cyst-

like organoids kept proliferating after loss of MLL1 and the Lgr5$^+$ ISC marker,

Olfm4, was elevated indicating differences between events in the crypt and in

culture. Differences are also indicated by the elevation of the Paneth cell

markers *Mmp7*, *Wnt3* and the elevation of the putative JAML receptor, *Cxadr*,

which might enable the transition from organoids to spheroids (Fig 6D). These

data establish a cell intrinsic requirement for MLL1 and indicate that it is

required to maintain ISCs in organoid cultures.

Discussion

Here we add a fourth stem cell to the known MLL1 repertoire of (i) HSCs [20, 21], (ii) skeletal muscle satellite cells [26] and (iii) postnatal neural stem cells (NSCs) [27]. The complete concordance between known MLL1 functions in postnatal stem cells suggests that MLL1 conveys an essential stem cell property. This possibility is enhanced by comparison to the MLL1 paralogue, MLL2, whose known functions in adult mice do not relate to stem cells rather macrophages (to respond to lipopolysaccharides) or fertility [48-51].

To explore the idea that MLL1 conveys a key stem cell property, we inspected the transcriptome profiles after conditional loss of MLL1 in the four adult/postnatal stem cells [26, 27, 52] (Supplementary excel file 1). However no shared candidate regulators or gene expression programs were identified. Although deeper, more systematic, transcriptome or cell biology approaches may reveal a shared MLL1 stem cell property, the lack of concordance between MLL1 regulation of these four stem cell transcriptomes is not unexpected. Previous work with MLL1 noted that direct target genes are not shared between different cell types [52] and a similar observation was made for MLL2 [48]. That is, the regulation of gene expression by the Trithorax homologues, MLL1 and MLL2, varies depending on the cell type and is not universal.

As again documented here for ISCs, the strongest relationship between the loss of MLL1 and cellular processes involves the downregulation of mRNAs that regulate transcription. Upon loss of MLL1, downregulation of

transcription factor mRNAs include - (i) in HSCs; *Mecom, Prdm16, Pbx1,*

Eya1, Meis1 and *Hoxa9*; (ii) in postnatal NSCs; *Nkx2.1, Nkx2.3*; (iii) in satellite

cells; *Pax7* and (iv) in ISCs; *Pitx2, Foxa1, Gata4* and *Onecut2*.

How does MLL1 regulate key lineage specific transcription factors

differently in different lineages? MLL1 and MLL2 are amongst the few proteins

that include the CxxC zinc finger that binds unmethylated CpG dinucleotides

[23, 53] as well as a PHD finger that binds H3K4me3 [54]. Hence, as

suggested before [48], MLL1 and 2 have the potential ability to bind CpG

island promoters without the need for recruitment by sequence specific DNA

binding transcription factors. This potential accords with the observation that

both MLL1 and 2 appear to be bound at almost all active promoters [40, 55].

Consequently additional factors are required to explain the restricted

transcriptional specificities of the MLLs. Notable in this regard, PAX7 is bound

to MLL1 when satellite cells are activated, and enhanced transcriptional

activation from both the *Myf5* promoter, to initiate skeletal muscle

replenishment, and the *Pax7* promoter itself, depends on MLL1 [26]. This

suggests that key transcription factors can either acquire the ability to interact

with MLL1 bound at target promoters or recruit MLL1 to target promoters, or

both.

Amongst the transcription factor mRNAs identified after loss of MLL1 in

ISCs, *Pitx2* is prominent. *Pitx2* was previously identified as a direct target of

MLL1 in ESCs and HSCs/hematopoietic progenitor cells [56, 57]. PITX2 is a

homeodomain protein responsible for left-right asymmetric morphogenesis in

the gut and proper positioning of the small intestine in the body cavity [58].

Also notably identified in ISCs are *Foxa1* and *Onecut2*. Both genes,

previously known as *Hnf3α* and *Hnf6α*, are expressed in all epithelia of the

gastrointestinal tract from its embryonic origin into adulthood. Together with

Math1, they are critical for goblet cell differentiation and function [59, 60].

Gata4, which has previously been described as an MLL1 target gene [61], is

also amongst the top downregulated mRNAs after loss of MLL1. Some

aspects of the intestine specific deletion of *Gata4* in the adult mouse resemble

the MLL1 phenotype described here including decreased proliferation in the

crypts with increased numbers of goblet cells [62].

In addition to the central relationship between MLL1 and transcription

factor expression, by far the most dramatically downregulated mRNA in both 4

and 10 day ISC transcriptomes was *Jaml*, which was previously included in

the transcriptome profile that characterizes ISCs [43, 44]. As a prominent cell

adhesion molecule, investigations by Tetteh and Clevers found that *Jaml* is

expressed in the base of the crypt in ISCs but not Paneth cells and used

Villin-CreERT2 to conditionally knock it out [63]. Loss of *Jaml* resulted in loss

of both *Olfm4*[+] ISCs and proliferation in the crypt without loss of Paneth cells.

These observations support the conclusion that MLL1 contributes to ISC

function mainly by expression of *Jaml*. However in contrast to the loss of

MLL1, Tetteh and Clevers did not observe an increase in goblet cells after

loss of JAML. Imbalanced commitment in the secretory lineage may indicate a

second aspect of MLL1 function that does not operate through *Jaml*

expression, possibly including the regulation of *Gata4* expression [62] and

other transcription factors. Upon loss of MLL1 *Jaml* is downregulated and the

close association between stem cells and Paneth cells is probably

destabilized (Fig 7). ISCs are anchored in the crypt surrounded by Paneth

cells, which confers positional identity. The central ISC at the bottom of the

crypt flanked by two Paneth cells has long-term self-renewal potential

compared to border ISCs [64]. Interestingly positional identity of NSCs seems

to be regulated in an MLL1 dependent fashion [27].

Once again we report the remarkably specific dependency of the

expression of one or two genes on a Trithorax homologue. As described here,

loss of MLL1 resulted in loss of *Jaml* expression with subsequent significant

functional consequences in the crypt. In ESCs, the expression of one gene,

Magoh2, entirely depends on MLL2. Removal of MLL2 resulted in the

suppression of *Magoh2* expression by H3K27 methylation, followed by DNA

methylation [48], thereby providing more evidence supporting the conclusion

that a primary function of Trithorax action is to prevent Polycomb-Group

repression [65]. These observations and conclusions with *Magoh2* in ESCs

were recently confirmed [66]. Selective gene specific anti-repression could

also explain the action of MLL2 on *Pigp* in macrophages [49] and MLL1 on

Hoxa9 in HSCs [25].

Despite a high degree of evolutionary conservation and near-

ubiquitous expression, the Trithorax homologues appear to regulate a small

number of genes in cell-type specific patterns, one or two of which are entirely

dependent on one homologue for expression due to anti-repression. In ISCs

this extraordinary specificity is focused on *Jaml* with functional consequences

for the crypt niche. As observed in other adult stem cells, MLL1 also regulates

the expression of transcription factors in ISCs that likely influence lineage

commitment decisions. These observations lead to the attractive proposition

that MLL1 is a master stem cell regulator. However a unifying molecular basis

for this proposition remains to be identified.

Methods and materials

Targeting constructs

The targeting construct for *Mll1* was generated using recombineering (S1A Fig) employing an engrailed-intron-splice-acceptor-IRES-LacZ-Neomycin-polyA cassette flanked by FRT sites [39]. The critical exon 2, which upon deletion results in a frameshift and a premature stop codon in exon 3, was flanked by loxP sites.

Gene targeting and generation of conditional knockout mice

Gene targeting in R1 embryonic stem cells (ESCs) was performed as described [67]. Correct integration in the *Mll1* locus was confirmed by Southern blot analysis using an internal probe and 5' and 3' external probes (S1B and S1C Figs). Two correctly targeted ESC clones were injected into blastocysts and gave rise to several chimeras, which were able to establish germ line transmission. *Mll1$^{A/+}$* mice were crossed to the *hACTB-Flpe* line to generate *Mll1$^{F/+}$* mice. *Mll1$^{A/+}$* and *Mll1$^{F/+}$* mice were backcrossed at least six generations to *C57BL/6JOlaHsd* mice. Subsequently, those mice were crossed to the *Rosa26-Cre-ERT2* (*RC*) line [5] to generate conditional, tamoxifen-inducible *Mll1$^{F/+; RC/+}$* mice. *Mll1$^{F/+}$* mice were bred with *Lgr5-eGFP-CreERT2* and *Villin-CreER^{T2}* (*Vil-Cre-ERT2*) [29, 41] mice. Primers for genotyping are provided in S2 Table. Experiments were performed in accordance with German animal welfare legislation, and were approved by the relevant authorities, the Landesdirektion Dresden.

Tamoxifen

Tamoxifen (Sigma Aldrich, T5648) was given to at least 10-week old mice by

gavage (4.5 mg per day) for six days with three days break in between [68].

For RNA-Sequencing experiments, *Mll1*$^{F/+;}$ and *Mll1*$^{F/F;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$ mice

received 1 mg tamoxifen via intraperitoneal (IP) injection for 3 consecutive

days. Intestinal organoids were induced on day 2 after splitting using 800 nM

4-OH tamoxifen (Sigma H7904) for 24 h.

Bone marrow transplantation

Mll1$^{F/+;\ RC/+}$ and *Mll1*$^{F/F;\ RC/+}$ recipient (CD45.2) mice were lethally irradiated

with 8.5 Gy (X-ray source MaxiShot from Yxlon). Bone marrow cells from

B6.SJL mice (CD45.1) were prepared by crushing with a mortar and pestle in

ice-cold PBS supplemented with 5% fetal bovine serum. Red blood cells were

removed with ACK lysis buffer (Thermo Fisher Scientific). 1 x 10^6 lineage

depleted (Lin$^-$) bone marrow cells were injected into the retro-orbital venous

plexus. Animals were maintained on water containing 1,17 mg/ml Neomycin

(Merck) for three weeks after irradiation. Complete donor cell engraftment of

wt CD45.1$^+$ cells was confirmed by flow cytometry on peripheral blood with

antibodies directed against the following murine antigens (clones given in

brackets): CD45.1 (A20), CD45.2 (104), CD11b (M1/70), Gr-1 (RB6-8C5).

Stably engrafted mice were fed six times with tamoxifen 30 weeks after

transplantation. FACS analysis for KSL-Slam enriched HSCs was done with

antibodies directed against the following murine antigens: CD3 (2C11; 17A2),

CD11b (M1/70), CD16/32 (93), CD19 (eBio 1D3), CD34 (RAM34), CD45.1

(A20), CD45.2 (104), CD45R (RA3-6B2), CD117 (2B8), CD135 (A2F10), Gr-1

(RB6-8C5), Nk1.1 (PK136), Ter119 (Ter119, all eBioscience), CD11b (M1/70)

and CD45.1 (A20, all BD Pharmingen), Sca-1 (D7), CD48 (HM48-1) and

CD150 (TC15-12F1, all BioLegends). Lin$^-$ cells were identified by lack of CD3,

CD11b, CD19, CD45R, Gr-1, Nk1.1 and Ter119 expression.

5-Bromo-2-deoxyuridine (BrdU) assay

459 Mice were injected IP with BrdU (0.6 mg/10 g body weight in sterile PBS) and

460 sacrificed 2 h later. The jejunum was dissected in cold PBS and processed for

461 immunohistochemistry.

Histochemistry and immunohistochemistry

Mouse intestine was flushed gently with cold PBS. Embryos were dissected

from plugged mice on the respective gestational stage and placed in PBS.

Intestine and embryos were fixed in 4% paraformaldehyde overnight.

Dehydration and paraffin infiltration utilized the Paraffin-Infiltration-Processor

(STP 420, Zeiss). Dehydrated tissues were embedded in paraffin (Paraffin

Embedding Center EG1160, Leica) and 5 μm sections were prepared.

Sections were deparaffinized in xylene, rehydrated through a series of

alcohols, stained, dehydrated, and mounted. Femur sections were stained

with Giemsa according to standard protocols. For the basic evaluation of the

intestine hematoxylin and eosin (H&E) stain was performed. Periodic acid-

Schiff (PAS) stain and alcian blue were used to identify goblet cells. For

enterocytes, alkaline phosphatase stain was performed using two different

methods. Sections were either incubated with Red Alkaline Phosphatase

Substrate (Vector red, Vector Laboratories) for 10 minutes (min) or with

nitroblue tetrazolium/5-bromo-4-chloro-3-indolyl phosphate solution for 30 min

at room temperature (RT). For immunohistochemistry, antigen retrieval was

performed by microwaving slides in 10 mM citrate buffer (pH 6.0) for 12 min

(Microwave RHS 30, Diapath). Endogenous peroxidases were quenched with

0.3% H_2O_2 in methanol. Sections were incubated in blocking serum (5% goat

serum) for 1 hr at RT followed by overnight incubation with primary antibodies

(S3 Table) at 4°C. Following incubation with the secondary antibody (S3

Table) the immune peroxidase was detected using a Vectastain ELITE ABC

kit (Vector) and visualized with a solution of diaminobenzidine (Sigma Aldrich)

in the presence of 0.01% H_2O_2. All sections were counterstained with

hematoxylin or alcian blue. Images were collected with an Olympus WF

upright microscope and analyzed using the MetaMorph® Microscopy

Automation and Image Analysis Software.

In situ **hybridization**

The protocol for *in situ* hybridization was modified from [69]. Briefly, 8 µm

thick sections were rehydrated. The sections were treated with 0.1 N HCl and

proteinase K. Slides were postfixed and sections were then demethylated with

acetic anhydride and prehybridized. Hybridization was done with 2 µg/ml

digoxigenin (DIG)-labeled *Olfm4* RNA probe for 24 h at 65°C. Slides were

washed and incubated with blocking solution for 1 hr. The sections were

incubated with anti-DIG-alkaline phosphatase conjugate overnight at 4°C.

Slides were washed and developed with BM purple.

Crypt isolation

The jejunum was harvested from mice, flushed with ice cold PBS to remove

any faecal content and cut open longitudinally. The tissue was placed lumen

side up on a petri dish and the villi were removed by gently scraping the tissue

using a glass cover slip. The tissue was cut into 2-4 cm pieces and was

washed several times with ice-cold PBS to remove residual villi fragments.

Tissues were transferred into a fresh tube containing 15 ml of 2 mM

EDTA/PBS chelation buffer and placed on a rotating wheel for 30 min at 4°C.

The crypts were then detached from the basal membrane by vigorous shaking

in 5% FCS/PBS solution. The suspension was filtered with a 100 µm cell

strainer followed by a 70 µm cell strainer. Isolated crypts were centrifuged at

800 rpm for 5 min at 4°C. The final fraction consisted of pure crypts and was

used for cell culture or single cell dissociation.

Organoid culture

Purified crypts were resuspended in 10 ml DMEM/F12 (Life Technologies). 10

µl of the crypt suspension was used to count the number of crypts under the

microscope. The pelleted crypts were resuspended in Matrigel® matrix

(Corning) at desired crypt density. Approximately, 400 crypts in 25 µl

Matrigel® matrix were seeded per well in a pre-warmed 24-well plate and

incubated for 15 min at 37°C until the Matrigel® matrix solidified. Then, 400 µl

of IntestiCult™ Organoid Growth Medium (STEMCELL Technologies) was

added to each well. Organoids were cultured at 37°C in a 5% CO_2 incubator

and maintained in culture for 5 days before being passaged and split for

experimental procedures. The growth medium was replaced every 2-3 days

Flow cytometry

ISCs were initially characterized and identified with the use of an *Lgr5-eGFP-*

CreERT2 knockin allele [29]. *Mll1^{F/+}* and *Mll1^{F/F; Lgr5-eGFP-CreERT2/+}* littermates

(n=4) were given tamoxifen via IP injections for 3 consecutive days and were

dissected 4 and 10 days later. Crypts were dissociated into single cells with

TrypLE Express (Thermo Fisher Scientific) for 30 min at 37°C. Dissociated

cells were passed through 70 µm cell strainer and washed with 5% FCS/PBS.

Cells were stained with antibodies with the following antibodies for 45 min on

ice: anti-mouse CD24 PE (clone M1/69), anti-mouse CD326 (EpCAM) APC

(clone G8.8), anti-mouse CD45 Alexa-Fluor 700 (clone 104). Sorting was

performed on a FACS Aria™ III cell sorter (BD). After scatter discrimination to

remove doublets the cell suspension was negatively selected with SYTOX

blue dead cell stain and anti-CD45 to remove dead and hematopoietic cells,

respectively. The cells where then positively selected with anti-EpCAM to

enrich for intestinal epithelial cells. According to [70] CD45$^-$, EpCAM^{high},

CD24^{med} and GFP^{high} characterized ISCs and CD45$^-$, EpCAM^{high}, CD24^{high}

and GFP$^-$ characterized Paneth cells (S3A and S3B Figs). *Mll1* recombination

via PCR on the sorted populations confirmed deletion of *Mll1* solely in the

stem cell compartment (S3C Fig).

RNA sequencing

300 intestinal stem and Paneth cells were sorted into 2 µl of nuclease free

water with 0.2% Triton-X 100 and 4 U murine RNase Inhibitor (NEB). RNA

was reverse transcribed (Invitrogen) and cDNA amplified using Kapa HiFi

HotStart Readymix (Roche). The cDNA quality and concentration was

determined with the Fragment Analyzer (Agilent). Samples were subjected to

library preparation (TruePrep DNA library Prep Kit V2 for Illumina, Vazyme).

Libraries were purified followed by Illumina sequencing on a Nextseq500 with

a sample sequencing depth of 30 million reads on average. The short reads

were aligned to the mm10 transcriptome with GSNAP (2018-07-04) and a

table of read counts per gene was created based on the overlap of the

uniquely mapped reads with the Ensembl Gene annotation version 92, using

featureCounts (version 1.6.3). Normalization of the raw read counts based on

the library size and testing for differential gene expression between the

different genotypes was performed using the DESeq2 R package (version

1.24.0). Genes with an adjusted p-value (padj)≤ 0.05 were considered as

significantly differentially expressed accepting a 5% FDR. To identify

enrichment for particular biological processes and pathways associated with

the DEGs, the DAVID GO/BP/FAT and KEGG database [71] was used. Gene

set enrichment analysis was performed using GSEA software from the Broad

Institute [72].

Reverse transcription and quantitative PCR (qRT-PCR) analysis

RNA from sorted cells and organoids was extracted using Trizol (Sigma-

Aldrich) and reverse transcribed using AffinityScript Multiple Temperature

cDNA Synthesis Kit (Agilent Technologies). Real-time quantitative PCR was

performed with GoTaq qPCR Master Mix (Promega) by Mx3000P QPCR

System (Agilent Technologies). Ct values were normalized against *Rpl19*.

Primer sequences and length of the amplified products are given in S2 Table.

Fold differences in expression levels were calculated according to the $2^{-\Delta Ct}$

method [73].

Quantification and Statistical analysis

Data is presented as mean and error bars indicate standard deviation (s.d.)

unless otherwise indicated. Statistical details of the experiments can be found

in the figure legends. Graphs and statistics were generated with GraphPad

Prism software (v6.0) and Microsoft excel. Significance (p-values) for Kaplan-

Meier graphs was determined by Mantel-Cox test. Significance (p-values <

0.05) was determined with Wald test or two-tailed Student's *t* test.

Significance for the breeding statistics was determined with Chi-square test. N

indicates the numbers of independent biological replicates per experiment

unless otherwise indicated.

Data availability: RNA sequencing data have been deposited in the Gene

Expression Omnibus under accession number GSE 157285.

Acknowledgements

We thank Mandy Obst, Isabell Kolbe, Heike Petzold and Stefanie

Weidlich for excellent technical assistance. We also thank the Biomedical

Services (BMS) of the Max Planck Institute of Molecular Cell Biology and

Genetics, Dresden for the excellent service and technical assistance. We are

grateful to Prof. Sebastian Zeissig (CRTD, Dresden) for helpful advice. We

thank the core facilities of the Biotechnology Center for providing assistance

with flow cytometry (Katja Schneider). The Advanced Imaging Facility, a core

facility of the CMCB Technology Platform at TU Dresden, http://biotp.tu-

dresden.de/facilities/advanced-imaging/ assisted this research.

References

1.   Blanpain C, Fuchs E. Stem cell plasticity. Plasticity of epithelial stem cells in tissue regeneration. Science. 2014;344(6189):1242281. Epub 2014/06/14. doi: 10.1126/science.1242281. PubMed PMID: 24926024; PubMed Central PMCID: PMCPMC4523269.

2.   Simons BD, Clevers H. Strategies for homeostatic stem cell self-renewal in adult tissues. Cell. 2011;145(6):851-62. doi: 10.1016/j.cell.2011.05.033. PubMed PMID: 21663791.

3.   Stadhouders R, Filion GJ, Graf T. Transcription factors and 3D genome conformation in cell-fate decisions. Nature. 2019;569(7756):345-54. Epub 2019/05/17. doi: 10.1038/s41586-019-1182-7. PubMed PMID: 31092938.

4.   Arndt K, Kranz A, Fohgrub J, Jolly A, Bledau AS, Di Virgilio M, et al. SETD1A protects HSCs from activation-induced functional decline in vivo. Blood. 2018;131(12):1311-24. Epub 2018/01/20. doi: 10.1182/blood-2017-09-806844. PubMed PMID: 29348130.

5.   Schmidt K, Zhang Q, Tasdogan A, Petzold A, Dahl A, Arneth BM, et al. The H3K4 methyltransferase Setd1b is essential for hematopoietic stem and progenitor cell homeostasis in mice. Elife. 2018;7. Epub 2018/06/20. doi: 10.7554/eLife.27157. PubMed PMID: 29916805; PubMed Central PMCID: PMCPMC6025962.

6.   Chen Y, Anastassiadis K, Kranz A, Stewart AF, Arndt K, Waskow C, et al. MLL2, Not MLL1, Plays a Major Role in Sustaining MLL-Rearranged Acute Myeloid Leukemia. Cancer Cell. 2017;31(6):755-70 e6. Epub 2017/06/14. doi: 10.1016/j.ccell.2017.05.002. PubMed PMID: 28609655; PubMed Central PMCID: PMCPMC5598468.

7.   Di Carlo V, Mocavini I, Di Croce L. Polycomb complexes in normal and malignant hematopoiesis. J Cell Biol. 2019;218(1):55-69. Epub 2018/10/21. doi: 10.1083/jcb.201808028. PubMed PMID: 30341152; PubMed Central PMCID: PMCPMC6314559.

8.   Lyko F. The DNA methyltransferase family: a versatile toolkit for epigenetic regulation. Nat Rev Genet. 2018;19(2):81-92. Epub 2017/10/17. doi: 10.1038/nrg.2017.80. PubMed PMID: 29033456.

9.   Piunti A, Shilatifard A. Epigenetic balance of gene expression by Polycomb and COMPASS families. Science. 2016;352(6290):aad9780. Epub 2016/06/04. doi: 10.1126/science.aad9780. PubMed PMID: 27257261.

10.   Liang G, Lin JC, Wei V, Yoo C, Cheng JC, Nguyen CT, et al. Distinct localization of histone H3 acetylation and H3-K4 methylation to the transcription start sites in the human genome. Proc Natl Acad Sci U S A. 2004;101(19):7357-62. Epub 2004/05/05. doi: 10.1073/pnas.0401866101. PubMed PMID: 15123803; PubMed Central PMCID: PMCPMC409923.

11.   Soares LM, He PC, Chun Y, Suh H, Kim T, Buratowski S. Determinants of Histone H3K4 Methylation Patterns. Mol Cell. 2017;68(4):773-85 e6. Epub 2017/11/14. doi: 10.1016/j.molcel.2017.10.013. PubMed PMID: 29129639; PubMed Central PMCID: PMCPMC5706784.

12.   Choudhury R, Singh S, Arumugam S, Roguev A, Stewart AF. The Set1 complex is dimeric and acts with Jhd2 demethylation to convey symmetrical H3K4 trimethylation. Genes Dev. 2019;33(9-10):550-64. Epub 2019/03/08. doi: 10.1101/gad.322222.118. PubMed PMID: 30842216; PubMed Central PMCID: PMCPMC6499330.

13.   Suganuma T, Workman JL. Chromatin and signaling. Curr Opin Cell
Biol. 2013;25(3):322-6. doi: 10.1016/j.ceb.2013.02.016. PubMed PMID:
23498660.
14.   Heintzman ND, Hon GC, Hawkins RD, Kheradpour P, Stark A, Harp
LF, et al. Histone modifications at human enhancers reflect global cell-type-
specific   gene   expression.   Nature.   2009;459(7243):108-12.   doi:
10.1038/nature07829. PubMed PMID: 19295514; PubMed Central PMCID:
PMC2910248.
15.   Ernst P, Vakoc CR. WRAD: enabler of the SET1-family of H3K4
methyltransferases.   Brief   Funct   Genomics.   2012;11(3):217-26.   Epub
2012/06/02. doi: 10.1093/bfgp/els017. PubMed PMID: 22652693; PubMed
Central PMCID: PMCPMC3388306.
16.   Roguev A, Schaft D, Shevchenko A, Pijnappel WW, Wilm M, Aasland
R, et al. The Saccharomyces cerevisiae Set1 complex includes an Ash2
homologue and methylates histone 3 lysine 4. EMBO J. 2001;20(24):7137-48.
Epub 2001/12/18. doi: 10.1093/emboj/20.24.7137. PubMed PMID: 11742990;
PubMed Central PMCID: PMCPMC125774.
17.   Milne TA, Briggs SD, Brock HW, Martin ME, Gibbs D, Allis CD, et al.
MLL targets SET domain methyltransferase activity to Hox gene promoters.
Mol   Cell.   2002;10(5):1107-17.   Epub   2002/11/28.   doi: 10.1016/s1097-
2765(02)00741-4. PubMed PMID: 12453418.
18.   Yagi H, Deguchi K, Aono A, Tani Y, Kishimoto T, Komori T. Growth
disturbance   in   fetal   liver   hematopoiesis   of   Mll-mutant   mice.   Blood.
1998;92(1):108-17. Epub 1998/06/25. PubMed PMID: 9639506.
19.   Ernst P, Fisher JK, Avery W, Wade S, Foy D, Korsmeyer SJ. Definitive
hematopoiesis   requires   the   mixed-lineage   leukemia   gene.   Dev   Cell.
2004;6(3):437-43. Epub 2004/03/20. PubMed PMID: 15030765.
20.   Jude CD, Climer L, Xu D, Artinger E, Fisher JK, Ernst P. Unique and
independent roles for MLL in adult hematopoietic stem cells and progenitors.
Cell Stem Cell. 2007;1(3):324-37. doi: 10.1016/j.stem.2007.05.019. PubMed
PMID: 18371366; PubMed Central PMCID: PMC2234224.
21.   Gan T, Jude CD, Zaffuto K, Ernst P. Developmentally induced Mll1 loss
reveals defects in postnatal haematopoiesis. Leukemia. 2010;24(10):1732-41.
doi: 10.1038/leu.2010.171.   PubMed   PMID: 20724987;   PubMed   Central
PMCID: PMC2954260.
22.   Slany RK. The molecular biology of mixed lineage leukemia.
Haematologica.   2009;94(7):984-93.   doi: 10.3324/haematol.2008.002436.
PubMed PMID: 19535349; PubMed Central PMCID: PMC2704309.
23.   Bach C, Mueller D, Buhl S, Garcia-Cuellar MP, Slany RK. Alterations of
the CxxC domain preclude oncogenic activation of mixed-lineage leukemia 2.
Oncogene. 2009;28(6):815-23. doi: 10.1038/onc.2008.443. PubMed PMID:
19060922.
24.   Slany RK. MLL fusion proteins and transcriptional control. Biochim
Biophys Acta Gene Regul Mech. 2020;1863(3):194503. Epub 2020/02/18.
doi: 10.1016/j.bbagrm.2020.194503. PubMed PMID: 32061883.
25.   Milne TA, Kim J, Wang GG, Stadler SC, Basrur V, Whitcomb SJ, et al.
Multiple interactions recruit MLL1 and MLL1 fusion proteins to the HOXA9
locus   in   leukemogenesis.   Mol   Cell.   2010;38(6):853-63.   doi:
10.1016/j.molcel.2010.05.011. PubMed PMID: 20541448; PubMed Central
PMCID: PMC2902588.

26. Addicks GC, Brun CE, Sincennes MC, Saber J, Porter CJ, Francis
Stewart A, et al. MLL1 is required for PAX7 expression and satellite cell self-
renewal in mice. Nat Commun. 2019;10(1):4256. Epub 2019/09/20. doi:
10.1038/s41467-019-12086-9. PubMed PMID: 31534153; PubMed Central
PMCID: PMCPMC6751293.
27. Delgado RN, Mansky B, Ahanger SH, Lu C, Andersen RE, Dou Y, et
al. Maintenance of neural stem cell positional identity by mixed-lineage
leukemia 1. Science. 2020;368(6486):48-53. Epub 2020/04/04. doi:
10.1126/science.aba5960. PubMed PMID: 32241942.
28. Clevers H. The intestinal crypt, a prototype stem cell compartment.
Cell. 2013;154(2):274-84. doi: 10.1016/j.cell.2013.07.004. PubMed PMID:
23870119.
29. Barker N, van Es JH, Kuipers J, Kujala P, van den Born M, Cozijnsen
M, et al. Identification of stem cells in small intestine and colon by marker
gene Lgr5. Nature. 2007;449(7165):1003-7. Epub 2007/10/16. doi:
10.1038/nature06196. PubMed PMID: 17934449.
30. Gehart H, Clevers H. Tales from the crypt: new insights into intestinal
stem cells. Nat Rev Gastroenterol Hepatol. 2019;16(1):19-34. Epub
2018/11/16. doi: 10.1038/s41575-018-0081-y. PubMed PMID: 30429586.
31. Tian H, Biehs B, Warming S, Leong KG, Rangell L, Klein OD, et al. A
reserve stem cell population in small intestine renders Lgr5-positive cells
dispensable. Nature. 2011;478(7368):255-9. doi: 10.1038/nature10408.
PubMed PMID: 21927002; PubMed Central PMCID: PMC4251967.
32. Tetteh PW, Basak O, Farin HF, Wiebrands K, Kretzschmar K, Begthel
H, et al. Replacement of Lost Lgr5-Positive Stem Cells through Plasticity of
Their Enterocyte-Lineage Daughters. Cell Stem Cell. 2016;18(2):203-13.
Epub 2016/02/03. doi: 10.1016/j.stem.2016.01.001. PubMed PMID:
26831517.
33. van Es JH, Sato T, van de Wetering M, Lyubimova A, Yee Nee AN,
Gregorieff A, et al. Dll1+ secretory progenitor cells revert to stem cells upon
crypt damage. Nat Cell Biol. 2012;14(10):1099-104. Epub 2012/09/25. doi:
10.1038/ncb2581. PubMed PMID: 23000963; PubMed Central PMCID:
PMCPMC3789123.
34. Jadhav U, Saxena M, O'Neill NK, Saadatpour A, Yuan GC, Herbert Z,
et al. Dynamic Reorganization of Chromatin Accessibility Signatures during
Dedifferentiation of Secretory Precursors into Lgr5+ Intestinal Stem Cells. Cell
Stem Cell. 2017;21(1):65-77 e5. Epub 2017/06/27. doi:
10.1016/j.stem.2017.05.001. PubMed PMID: 28648363; PubMed Central
PMCID: PMCPMC5505276.
35. Yu S, Tong K, Zhao Y, Balasubramanian I, Yap GS, Ferraris RP, et al.
Paneth Cell Multipotency Induced by Notch Activation following Injury. Cell
Stem Cell. 2018;23(1):46-59 e5. Epub 2018/06/12. doi:
10.1016/j.stem.2018.05.002. PubMed PMID: 29887318; PubMed Central
PMCID: PMCPMC6035085.
36. Yang W, Ernst P. Distinct functions of histone H3, lysine 4
methyltransferases in normal and malignant hematopoiesis. Curr Opin
Hematol. 2017;24(4):322-8. Epub 2017/04/05. doi:
10.1097/moh.0000000000000346. PubMed PMID: 28375985; PubMed
Central PMCID: PMCPMC5603181.

37. Tetteh PW, Farin HF, Clevers H. Plasticity within stem cell hierarchies
in mammalian epithelia. Trends Cell Biol. 2015;25(2):100-8. doi:
10.1016/j.tcb.2014.09.003. PubMed PMID: 25308311.
38. Fu J, Teucher M, Anastassiadis K, Skarnes W, Stewart AF. A
recombineering pipeline to make conditional targeting constructs. Methods
Enzymol. 2010;477:125-44. Epub 2010/08/12. doi: 10.1016/s0076-
6879(10)77008-7. PubMed PMID: 20699140.
39. Testa G, Schaft J, van der Hoeven F, Glaser S, Anastassiadis K,
Zhang Y, et al. A reliable lacZ expression reporter cassette for multipurpose,
knockout-first alleles. Genesis. 2004;38(3):151-8. doi: 10.1002/gene.20012.
PubMed PMID: 15048813.
40. Denissov S, Hofemeister H, Marks H, Kranz A, Ciotta G, Singh S, et al.
Mll2 is required for H3K4 trimethylation on bivalent promoters in embryonic
stem cells, whereas Mll1 is redundant. Development. 2014;141(3):526-37.
Epub 2014/01/16. doi: 10.1242/dev.102681. PubMed PMID: 24423662.
41. el Marjou F, Janssen KP, Chang BH, Li M, Hindie V, Chan L, et al.
Tissue-specific and inducible Cre-mediated recombination in the gut
epithelium. Genesis. 2004;39(3):186-93. Epub 2004/07/30. doi:
10.1002/gene.20042. PubMed PMID: 15282745.
42. Yan KS, Gevaert O, Zheng GXY, Anchang B, Probert CS, Larkin KA, et
al. Intestinal Enteroendocrine Lineage Cells Possess Homeostatic and Injury-
Inducible Stem Cell Activity. Cell Stem Cell. 2017;21(1):78-90 e6. Epub
2017/07/08. doi: 10.1016/j.stem.2017.06.014. PubMed PMID: 28686870;
PubMed Central PMCID: PMCPMC5642297.
43. Munoz J, Stange DE, Schepers AG, van de Wetering M, Koo BK,
Itzkovitz S, et al. The Lgr5 intestinal stem cell signature: robust expression of
proposed quiescent '+4' cell markers. EMBO J. 2012;31(14):3079-91. Epub
2012/06/14. doi: 10.1038/emboj.2012.166. PubMed PMID: 22692129;
PubMed Central PMCID: PMCPMC3400017.
44. Haber AL, Biton M, Rogel N, Herbst RH, Shekhar K, Smillie C, et al. A
single-cell survey of the small intestinal epithelium. Nature.
2017;551(7680):333-9. Epub 2017/11/17. doi: 10.1038/nature24489. PubMed
PMID: 29144463; PubMed Central PMCID: PMCPMC6022292.
45. Zen K, Liu Y, McCall IC, Wu T, Lee W, Babbin BA, et al. Neutrophil
migration across tight junctions is mediated by adhesive interactions between
epithelial coxsackie and adenovirus receptor and a junctional adhesion
molecule-like protein on neutrophils. Mol Biol Cell. 2005;16(6):2694-703.
Epub 2005/04/01. doi: 10.1091/mbc.e05-01-0036. PubMed PMID: 15800062;
PubMed Central PMCID: PMCPMC1142417.
46. Kawabata K, Tashiro K, Sakurai F, Osada N, Kusuda J, Hayakawa T,
et al. Positive and negative regulation of adenovirus infection by CAR-like
soluble protein, CLSP. Gene Ther. 2007;14(16):1199-207. Epub 2007/06/01.
doi: 10.1038/sj.gt.3302975. PubMed PMID: 17538635.
47. Rodríguez-Colman MJ, Schewe M, Meerlo M, Stigter E, Gerrits J,
Pras-Raves M, et al. Interplay between metabolic identities in the intestinal
crypt supports stem cell function. Nature. 2017;543(7645):424-7. Epub
2017/03/09. doi: 10.1038/nature21673. PubMed PMID: 28273069.
48. Glaser S, Lubitz S, Loveland KL, Ohbo K, Robb L, Schwenk F, et al.
The histone 3 lysine 4 methyltransferase, Mll2, is only required briefly in
development and spermatogenesis. Epigenetics Chromatin. 2009;2(1):5. doi:

10.1186/1756-8935-2-5. PubMed PMID: 19348672; PubMed Central PMCID:
PMC2674429.
49.   Austenaa L, Barozzi I, Chronowska A, Termanini A, Ostuni R,
Prosperini E, et al. The histone methyltransferase Wbp7 controls macrophage
function through GPI glycolipid anchor synthesis. Immunity. 2012;36(4):572-
85. doi: 10.1016/j.immuni.2012.02.016. PubMed PMID: 22483804.
50.   Andreu-Vieyra CV, Chen R, Agno JE, Glaser S, Anastassiadis K,
Stewart AF, et al. MLL2 is required in oocytes for bulk histone 3 lysine 4
trimethylation and transcriptional silencing. PLoS Biol. 2010;8(8). doi:
10.1371/journal.pbio.1000453. PubMed PMID: 20808952; PubMed Central
PMCID: PMC2923083.
51.   Hanna CW, Taudt A, Huang J, Gahurova L, Kranz A, Andrews S, et al.
MLL2 conveys transcription-independent H3K4 trimethylation in oocytes. Nat
Struct Mol Biol. 2018;25(1):73-82. Epub 2018/01/13. doi: 10.1038/s41594-
017-0013-5. PubMed PMID: 29323282.
52.   Artinger EL, Mishra BP, Zaffuto KM, Li BE, Chung EK, Moore AW, et
al. An MLL-dependent network sustains hematopoiesis. Proc Natl Acad Sci U
S A. 2013;110(29):12000-5. doi: 10.1073/pnas.1301278110. PubMed PMID:
23744037; PubMed Central PMCID: PMC3718146.
53.   Carlone DL, Hart SR, Ladd PD, Skalnik DG. Cloning and
characterization of the gene encoding the mouse homologue of CpG binding
protein. Gene. 2002;295(1):71-7. Epub 2002/09/21. doi: 10.1016/s0378-
1119(02)00820-x. PubMed PMID: 12242013.
54.   Chang PY, Hom RA, Musselman CA, Zhu L, Kuo A, Gozani O, et al.
Binding of the MLL PHD3 finger to histone H3K4me3 is required for MLL-
dependent gene transcription. J Mol Biol. 2010;400(2):137-44. doi:
10.1016/j.jmb.2010.05.005. PubMed PMID: 20452361; PubMed Central
PMCID: PMC2886590.
55.   Guenther MG, Jenner RG, Chevalier B, Nakamura T, Croce CM,
Canaani E, et al. Global and Hox-specific roles for the MLL1
methyltransferase. Proc Natl Acad Sci U S A. 2005;102(24):8603-8. doi:
10.1073/pnas.0503072102. PubMed PMID: 15941828; PubMed Central
PMCID: PMC1150839.
56.   Ernst P, Mabon M, Davidson AJ, Zon LI, Korsmeyer SJ. An Mll-
dependent Hox program drives hematopoietic progenitor expansion. Curr Biol.
2004;14(22):2063-9. doi: 10.1016/j.cub.2004.11.012. PubMed PMID:
15556871.
57.   Degar BA, Baskaran N, Hulspas R, Quesenberry PJ, Weissman SM,
Forget BG. The homeodomain gene Pitx2 is expressed in primitive
hematopoietic stem/progenitor cells but not in their differentiated progeny. Exp
Hematol. 2001;29(7):894-902. Epub 2001/07/05. doi: 10.1016/s0301-
472x(01)00661-0. PubMed PMID: 11438212.
58.   Welsh IC, Thomsen M, Gludish DW, Alfonso-Parra C, Bai Y, Martin JF,
et al. Integration of left-right Pitx2 transcription and Wnt signaling drives
asymmetric gut morphogenesis via Daam2. Dev Cell. 2013;26(6):629-44.
Epub 2013/10/05. doi: 10.1016/j.devcel.2013.07.019. PubMed PMID:
24091014; PubMed Central PMCID: PMCPMC3965270.
59.   Ye DZ, Kaestner KH. Foxa1 and Foxa2 control the differentiation of
goblet and enteroendocrine L- and D-cells in mice. Gastroenterology.

2009;137(6):2052-62. Epub 2009/09/10. doi: 10.1053/j.gastro.2009.08.059.
PubMed PMID: 19737569; PubMed Central PMCID: PMCPMC2789913.
60.   van der Sluis M, Vincent A, Bouma J, Korteland-Van Male A, van
Goudoever JB, Renes IB, et al. Forkhead box transcription factors Foxa1 and
Foxa2 are important regulators of Muc2 mucin expression in intestinal
epithelial cells. Biochem Biophys Res Commun. 2008;369(4):1108-13. Epub
2008/03/14. doi: 10.1016/j.bbrc.2008.02.158. PubMed PMID: 18336786.
61.   Wang P, Lin C, Smith ER, Guo H, Sanderson BW, Wu M, et al. Global
analysis of H3K4 methylation defines MLL family member targets and points
to a role for MLL1-mediated H3K4 methylation in the regulation of
transcriptional initiation by RNA polymerase II. Mol Cell Biol.
2009;29(22):6074-85. doi: 10.1128/MCB.00924-09. PubMed PMID:
19703992; PubMed Central PMCID: PMC2772563.
62.   Beuling E, Baffour-Awuah NY, Stapleton KA, Aronson BE, Noah TK,
Shroyer NF, et al. GATA factors regulate proliferation, differentiation, and
gene expression in small intestine of mature mice. Gastroenterology.
2011;140(4):1219-29 e1-2. doi: 10.1053/j.gastro.2011.01.033. PubMed PMID:
21262227; PubMed Central PMCID: PMC3541694.
63.   Tetteh PW. Plasticity of intestinal epithelial cells in regeneration and
cancer. Utrecht University; 2015.
64.   Ritsma L, Ellenbroek SIJ, Zomer A, Snippert HJ, de Sauvage FJ,
Simons BD, et al. Intestinal crypt homeostasis revealed at single-stem-cell
level by in vivo live imaging. Nature. 2014;507(7492):362-5. Epub 2014/02/18.
doi: 10.1038/nature12972. PubMed PMID: 24531760; PubMed Central
PMCID: PMCPMC3964820.
65.   Klymenko T, Muller J. The histone methyltransferases Trithorax and
Ash1 prevent transcriptional silencing by Polycomb group proteins. EMBO
Rep. 2004;5(4):373-7. doi: 10.1038/sj.embor.7400111. PubMed PMID:
15031712; PubMed Central PMCID: PMC1299022.
66.   Douillet D, Sze CC, Ryan C, Piunti A, Shah AP, Ugarenko M, et al.
Uncoupling histone H3K4 trimethylation from developmental gene expression
via an equilibrium of COMPASS, Polycomb and DNA methylation. Nat Genet.
2020;52(6):615-25. Epub 2020/05/13. doi: 10.1038/s41588-020-0618-1.
PubMed PMID: 32393859.
67.   Bledau AS, Schmidt K, Neumann K, Hill U, Ciotta G, Gupta A, et al.
The H3K4 methyltransferase Setd1a is first required at the epiblast stage,
whereas Setd1b becomes essential after gastrulation. Development.
2014;141(5):1022-35. doi: 10.1242/dev.098152. PubMed PMID: 24550110.
68.   Anastassiadis K, Glaser S, Kranz A, Berhardt K, Stewart AF. A
practical summary of site-specific recombination, conditional mutagenesis,
and tamoxifen induction of CreERT2. Methods Enzymol. 2010;477:109-23.
Epub 2010/08/12. doi: 10.1016/s0076-6879(10)77007-5. PubMed PMID:
20699139.
69.   Gregorieff A, Clevers H. In Situ Hybridization to Identify Gut Stem
Cells. Curr Protoc Stem Cell Biol. 2015;34:2F 1 -2F 1 11. doi:
10.1002/9780470151808.sc02f01s34. PubMed PMID: 26237569.
70.   Nefzger CM, Jarde T, Rossello FJ, Horvay K, Knaupp AS, Powell DR,
et al. A Versatile Strategy for Isolating a Highly Enriched Population of
Intestinal Stem Cells. Stem Cell Reports. 2016;6(3):321-9. Epub 2016/03/01.

doi: 10.1016/j.stemcr.2016.01.014. PubMed PMID: 26923822; PubMed
Central PMCID: PMCPMC4788784.
71. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. Nat Protoc.
2009;4(1):44-57. doi: 10.1038/nprot.2008.211. PubMed PMID: 19131956.
72. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL,
Gillette MA, et al. Gene set enrichment analysis: a knowledge-based
approach for interpreting genome-wide expression profiles. Proc Natl Acad
Sci U S A. 2005;102(43):15545-50. Epub 2005/10/04. doi:
10.1073/pnas.0506580102. PubMed PMID: 16199517; PubMed Central
PMCID: PMCPMC1239896.
73. Livak KJ, Schmittgen TD. Analysis of relative gene expression data
using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method.
Methods. 2001;25(4):402-8. doi: 10.1006/meth.2001.1262. PubMed PMID:
11846609.
907

   Figure captions

   Fig 1. Loss of MLL1 in adult mice. (A) Giemsa-stained sections from the

   femur of $Mll1^{FC/+; RC/+}$ and $Mll1^{FC/FC; RC/+}$ mice. Two weeks after the last

   tamoxifen gavage mice were sacrificed and the femurs were dissected,

   decalcified, sectioned and stained. Bone marrow cellularity was severely

   decreased in $Mll1^{FC/FC; RC/+}$ mice. Scale bar 100 µm. (B) Kaplan-Meier survival

   curve. The first day of tamoxifen gavage was day zero. All $Mll1^{FC/+; RC/+}$ mice

   (n=25) survived whereas all $Mll1^{FC/FC; RC/+}$ mice (n=33) died within 33 days

   after tamoxifen induction with a median survival of 11 days. (C) Scheme of the

   experimental setup for bone marrow transplantation. Donor bone marrow from

   B6.SJL (CD45.1$^+$) mice was transplanted into lethally irradiated $Mll1^{F/+; RC/+}$

   and $Mll1^{F/F; RC/+}$ recipients (CD45.2$^+$). Blood chimerism (BC) was measured

   three times. After 30 weeks $Mll1$ deletion was achieved by administrating

   tamoxifen (TAM). FACS analysis for KSL-Slam enriched HSCs (Kit$^+$ Sca1$^+$

   Lin$^-$ CD48$^-$ CD150$^+$ CD34$^-$ CD135$^-$) showed comparable numbers in BMTx

   $Mll1^{FC/+; RC/+}$ and $Mll1^{FC/FC; RC/+}$ mice. Dot plots show Lin$^-$ CD48$^-$ CD150$^+$ CD34$^-$

   CD135$^-$ gated bone marrow (BM) cells of indicated genotypes resolved for the

   expression of c-Kit and Sca-1. Donor and host cells are distinguished by

   surface markers CD45.1 and CD45.2. (D) Kaplan-Meier analysis for the onset

   of diarrhea. Tamoxifen was given by gavage for 6 days to $Mll1^{F/+; RC/+}$ (n=6)

   and $Mll1^{F/F; RC/+}$ (n=8). The first day of tamoxifen gavage was day zero. While

   all BMTx mice with the genotype $Mll1^{FC/+; RC/+}$ remained healthy, all BMTx

   $Mll1^{FC/FC; RC/+}$ mice developed diarrhea with a median of 16.5 days. (E)

   Antibody staining (brown) showed that MLL1 is expressed in crypts of the

small and large intestine but absent in the villus (hematoxylin, purple). Scale

bars are 50 µm.

Fig 2. ***Mll1* deletion in BMTx mice leads to loss of stem and proliferating**

cells and increased goblet cells. (A) Antibody stainings showed reduced

MLL1, OLFM4 and SOX9 expression in BMTx *Mll1*$^{FC/FC; \ RC/+}$ intestine

compared to controls. Hematoxylin was used as a counterstain for MLL1 and

OLFM4 immunohistochemistry (IHC). Expression of Ki67, a marker for

proliferating cells was reduced in BMTx *Mll1*$^{FC/FC; \ RC/+}$ mutant intestinal

sections. Arrowheads point toward proliferating ISCs. Alcian blue was used as

a counterstain for SOX9 and Ki67 IHC, which also revealed enlarged goblet

cells (turquoise) in the crypts of BMTx *Mll1*$^{FC/FC; RC/+}$ sections. Scale bars are

50 µm. (B) PAS stain to examine goblet cells in villi and crypts of BMTx

intestine. Black arrowheads point towards vacuolar structures. Yellow

arrowhead points to a mislocalized goblet cell in BMTx *Mll1*$^{FC/FC; \ RC/+}$ crypt.

Left panels scale bar 100 µm; middle panels scale bar 50 µm. GOB5 antibody

staining of BMTx *Mll1*$^{FC/FC; RC/+}$ intestinal sections. Right panels scale bar 100

µm. (C) Left panels; lysozyme antibody staining reveals that Paneth cell

numbers remain unchanged. Arrowheads point at Paneth cells. Alcian blue

was used as a counterstain and marks goblet cells (turquoise). Middle panels;

chromogranin A antibody stain, arrowheads point to the sparse

enteroendocrine cells (dark brown) in the villi. Right panels; red enterocytes

covering the villi were visualized by alkaline phosphatase staining.

Hematoxylin was used as a counterstain for chromogranin A IHC and alkaline

phosphatase histochemical staining. Scale bars are 50 µm.

Fig 3. Decreased ISCs and increased goblet cells after intestinal specific loss of MLL1. (A) Percent of weight loss of *Mll1*$^{FC/+; Vil-Cre-ERT2/+}$ (n=5) and of *Mll1*$^{FC/FC; Vil-Cre-ERT2/+}$ (n=9) mice. Mean ± s.d. is shown (p=0.029, Student's *t*-test). (B) Decrease in ISC markers, OLFM4 and SOX9 in *Mll1*$^{FC/FC; Vil-Cre-ERT2/+}$ intestinal sections. SOX9$^+$ ISCs in *Mll1*$^{FC/+; Vil-Cre-ERT2/+}$ intestinal sections are marked with red arrowheads. Proliferating cells in the TA compartment as well as proliferative ISCs (arrowhead) are reduced in *Mll1*$^{FC/FC; Vil-Cre-ERT2/+}$ sections. Alcian blue was used as a counterstain after staining for SOX9 and Ki67 and marks goblet cells (turquoise). Scale bars are 100 μm for OLFM4 and 50 μm for SOX9 and Ki67. (C) Left panels; alcian blue staining of goblet cells with nuclear fast red (NFR) to stain nuclei. Middle panels; Paneth cells visualized by staining of granules containing lysozyme (arrowheads). Right panels; enteroendocrine cells stained with chromogranin A antibody (brown; arrowheads) in the villi. Scale bars are 100 μm.

Fig 4. RNA profiling of *Mll1*$^{FC/FC; Lgr5-eGFP-CreERT2/+}$ and control ISCs. (A) ISCs were sorted from control (*Mll1*$^{FC/+; Lgr5-eGFP-CreERT2/+}$) (n=4) and *Mll1*$^{FC/FC; Lgr5-eGFP-CreERT2/+}$ (n=4) mice 4 days after tamoxifen induction was completed and subjected to RNA profiling. MA plot visualizing the log2-fold change differences according to expression levels of ISCs. Red dots represent significant DEGs at a 5% FDR. *Jaml* is the top downregulated gene. (B) Plots show biological processes (BP) that are enriched in genes up- or downregulated in *Mll1*$^{FC/FC; Lgr5-eGFP-CreERT2/+}$ compared to control ISCs. Analysis was performed using the gene ontology (GO)/BP/FAT database of DAVID 6.8. (C) (D) GSEA shows significant negative or positive correlation of

genes from the stem (C) and goblet cell (D) signature gene set in $Mll1^{FC/FC;}$

$^{Lgr5\text{-}eGFP\text{-}CreERT2/+}$ compared to control ISCs (4 days after tamoxifen). The

signature gene sets originate from [44]. NES: normalized enrichment score.

(E) DESeq normalized counts for genes coding for transcription factors

downregulated in $Mll1^{FC/FC;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$ compared to control ISCs (4 days

after tamoxifen). Mean+s.d. is shown; n=4; p<0.05, Wald test. (F) DESeq

normalized counts for genes coding for ISC markers downregulated in

$Mll1^{FC/FC;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$ compared to control ISCs (4 days after tamoxifen).

Mean+s.d. is shown; n=4; p<0.05, Wald test (G) DESeq normalized counts for

genes coding for goblet cell markers upregulated in $Mll1^{FC/FC;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$

compared to control ISCs (4 days after tamoxifen). Mean+s.d. is shown; n=4;

p<0.05, Wald test.

Fig 5. RNA profiling of wt Paneth cells after deletion of *Mll1* in the

neighboring ISCs. (A) Wt Paneth cells neighboring either $Mll1^{FC/+;\ Lgr5\text{-}eGFP\text{-}}$

$^{CreERT2/+}$ or $Mll1^{FC/FC;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$ ISCs were sorted 4 days after tamoxifen

induction was completed and were subjected to RNA profiling. MA plot

visualizing the log2-fold change differences according to expression levels of

Paneth cells. Red dots represent significant DEGs at a 5% FDR. (B) Enriched

terms of biological processes and pathways down- and upregulated using

DAVID GO/BP/FAT and KEGG database. (C) GSEA shows significant

negative or positive correlation of genes from the GO ERAD pathway, GO

oxidative phosphorylation and Paneth cell signature gene set in wt Paneth

cells neighboring either $Mll1^{FC/FC;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$ or $Mll1^{FC/+;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$

ISCs. The Paneth cell signature gene set originates from [44]. NES:

normalized enrichment score. (D) DESeq normalized counts for selected

genes differentially regulated in the ERAD pathway, oxidative phosphorylation

and Paneth cell signature gene set. Mean+s.d. is shown; n=4; $p<0.05$, Wald

test.

1013

Fig 6. *Mll1* deletion in organoids results in formation of spheres. (A)

Differential interference contrast (DIC) images of $Mll1^{FC/+; \ RC/+}$ and $Mll1^{FC/FC;}$

$^{RC/+}$ organoids. Organoids were induced with 4-OH tamoxifen for 24 h. Upon

passaging (P), the mutant starts to loose its budding morphology giving rise to

an undifferentiated cyst-like appearance. Scale bar 100 µm. (B) Genotyping of

organoids/spheres for *Mll1* at passage 2. After tamoxifen induction the $Mll1^F$

allele recombines and results in the $Mll1^{FC}$ allele. PCR primers located

upstream of the 5' FRT site and downstream of the 3' loxP site (see S1A Fig)

identify the respective band. Consequently, the $Mll1^F$ band is 1084 bp, the wt

band is 933 bp and the $Mll1^{FC}$ band is 186 bp. An additional PCR was

performed with primers flanking the 3' loxP site. Consequently, the $Mll1^F$ band

is 297 bp and the wt band is 251 bp. (C) Quantification of organoids and

spheres shown in A. (D) qRT-PCR was performed for selected genes on

cDNA from $Mll1^{FC/+; \ RC/+}$ and $Mll1^{FC/FC; \ RC/+}$ organoid culture. Mean+s.d. is

shown; n=3; *$p<0.05$, **$p<0.01$, ***$p<0.001$, Student's *t* test.

1029

1030   **Fig 7. Cell-cell contacts are essential for cell identity.** JAML is an integral

1031   transmembrane protein expressed on ISCs and interacts with unknown

1032   proteins on Paneth cells. Loss of MLL1 in ISCs causes loss of JAML mediated

1033   interactions, transcriptional changes and subsequently loss of stem and

1034   Paneth cell identity.

 Supporting information

1036

S1 Fig. *Mll1* gene targeting, embryonic phenotype and aspects of

expression. (A) Diagram of the *Mll1* gene with numbered exons and the

multipurpose allele (*Mll1^A*). This allele is converted to *Mll1^F* upon FLP

recombination. Cre recombination leads to excision of the frameshifting exon

2 generating the conditional mutant allele (*Mll1^{FC}*). Genotyping primers are

depicted for the downstream loxP site (loxP1 – loxP2) and for Flp

recombination (Flp se – loxP2). SA = splice acceptor, IRES = internal

ribosome entry site, pA = polyadenylation signal, lacZ-neo = β-galactosidase

and neomycin resistance gene, * depicts premature stop codon. **(B)**

Schematic representation of the Southern blot strategy. For identifying correct

targeted events in the *Mll1* locus, Southern blot analysis employed 5' (blue

box) and 3' (red box) probes. **(C)** Southern blot analysis using 5' and 3'

external probes. **(D)** Dissected embryos from *Mll1$^{A/+}$* intercrosses at E12.5.

Mll1$^{A/A}$ embryos had a pale liver (marked by arrow). **(E)** Antibody staining

(brown) shows that MLL1 is expressed in crypts and TA compartment of the

small intestine but is absent in the villus (hematoxylin, purple). Scale bar 50

μm. **(F)** Normalized RNA-sequence counts for *Mll1/Kmt2a, Mll2/Kmt2b,*

Mll3/Kmt2c, Mll4/Kmt2d, Setd1a/Kmt2f and Setd1b/Kmt2g in ISCs (eGFP^{high})

and Paneth cells sorted from *Lgr5-eGFP-CreERT2* mice. Mean+s.d. is shown;

n=4; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, Student's *t* test. **(G)**

Antibody stainings of H3K4me1, H3K4me2 and H3K4me3 are comparable in

Mll1$^{FC/+; RC/+}$ and *Mll1$^{FC/FC; RC/+}$* intestinal sections. Scale bars are 100 μm.

1059

S2 Fig. Without bone marrow transplantation, *Mll1* deletion recapitulates

the BMTx *Mll1* mutant phenotype. (A) Antibody stain (left panels) and *in situ*

hybridization (right panels) to visualize OLFM4/*Olfm4* in intestinal sections.

Arrowheads point towards ISCs. Scale bar 100 µm. **(B)** Proliferative activity

visualized by both Ki67 stain and BrdU incorporation in intestinal sections.

Arrowheads point towards proliferative ISCs. Scale bars are 50 µm. **(C)** PAS

staining and GOB5 antibody stain to visualize goblet cells in intestinal

sections. Scale bars are 100 µm. **(D)** Chromogranin A and alkaline

phosphatase staining to visualize enteroendocrine cells and enterocytes

respectively. Arrows point to enteroendocrine cells (brown cytoplasmic stain)

in the villi. Blue enterocytes covering the villi are marked by arrowheads.

Scale bars are 100 µm for chromogranin A and 50 µm for alkaline

phosphatase. **(E)** Nuclear β-catenin is comparable between the two different

genotypes. Arrowheads point at β-catenin positive nuclei. Scale bar is 50 µm.

1074

S3 Fig. FACS gating strategy to sort ISCs and Paneth cells. Flow sorting

on **(A)** *Mll1$^{FC/+; Lgr5-eGFP-CreERT2/+}$* and **(B)** *Mll1$^{FC/FC; Lgr5-eGFP-CreERT2/+}$* single cell

suspension of crypts. Briefly, the consecutive gating steps were applied: (i) –

(iii) Definition of the population of interest by exclusion of debris based on size

(FSC), granularity (SSC) and the selection for single cells; (iv) Exclusion of

dead cells that incorporated the nucleic acid stain SYTOX blue; (v) Depletion

of CD45^{pos} population; (vi) Definition of Paneth (EpCAM^{high}/CD24^{high}) cell

population by plotting EpCAM vs CD24 fluorescence; (vii) EpCAM^{high}/CD24^{med}

cell population was gated to discriminate the stem cell population (GFP^{high}).

(C) Stem cells (SC) and Paneth cells (PC) from *Mll1$^{FC/+; Lgr5-eGFP-CreERT2/+}$* and

Mll1^(FC/FC; Lgr5-eGFP-CreERT2/+) mice 4 days after tamoxifen induction were checked

for recombination. Left panel; PCR genotyping was using primers upstream of

the 5' FRT site and downstream of the 3' loxP site identified the *Mll1^F* band at

1084 bp, the wild type band a 933 bp and the *Mll1^(FC)* band at 186 bp. Right

panel; primers flanking the 3' loxP site identified the *Mll1^F* band at 297 bp and

the wild type band at 251 bp.

1091

S4 Fig. Alignment and quality of the sequenced data. (A) ISCs and

Paneth cells were analyzed from control (*Mll1^(FC/+; Lgr5-eGFP-CreERT2/+)*) (ctrl) (n=4)

and *Mll1^(FC/FC; Lgr5-eGFP-CreERT2/+)* (n=4) (KO) mice. Mappability of reads for sorted

ISCs 4 days after tamoxifen induction was completed. **(B)** Mappability of

reads for sorted Paneth cells 4 days after tamoxifen induction was completed.

(C) Mappability of reads for sorted ISCs 10 days after tamoxifen induction was

completed. ISCs were analyzed from control (*Mll1^(FC/+; Lgr5-eGFP-CreERT2/+)*) (ctrl)

(n=4) and *Mll1^(FC/FC; Lgr5-eGFP-CreERT2/+)* (n=3) (KO) mice. **(D)** Principal-component

analysis (PCA) was performed on Paneth cell and ISC samples sorted 4 days

after tamoxifen. PCA is based on mRNA changes for the top 500 most diverse

genes of stem cell (SC) and Paneth cell (PC) samples in comparison to

published datasets for Lgr5^+ SC [42] and CD24^+ PC [35].

1104

S5 Fig. ISCs lacking MLL1 loose their cellular identity. (A) (B) GSEA

shows significant negative or positive correlation of genes from the stem (A)

and goblet cell (B) signature gene set in *Mll1^(FC/FC; Lgr5-eGFP-CreERT2/+)* ISCs

compared to control ISCs 10 days after tamoxifen induction was completed.

The signature gene sets originate from [44]. NES: normalized enrichment

score. **(C)** To validate RNA-seq results qRT-PCR was performed for selected

genes on cDNA from *Mll1*$^{FC/+;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$ and *Mll1*$^{FC/FC;\ Lgr5\text{-}eGFP\text{-}CreERT2/+}$

sorted stem cells 4 days after tamoxifen induction was completed. Mean+s.d.

is shown; n=3; *$p<0.05$, **$p<0.01$, Student's *t* test. **(D)** DESeq normalized

counts demonstrate *Jaml* being highly expressed in Lgr5^{+} ISCs but not in

Paneth cells. Mean+s.d. is shown; n=4; ***$p<0.001$, Student's *t* test.

1116

Fig 1.

+ Goblet cell   ◣TA cell   * Paneth cell   ◣Stem cell   ►Border between crypt and villus

Fig 2.

Fig 3.

Fig 4.

Fig 5.

Fig 6.

A

	P0 uninduced	P1 induced	P2 induced	P3 induced

Mll1^FC/+; RC/+

Mll1^FC/FC; RC/+

B

F →
wt →
— 1000 bp
— 500 bp
FC →
F →
wt →
— 300 bp
Mll1^FC/+   Mll1^FC/FC

C

Ratio of organoids vs spheres

P0 uninduced   P1 induced   P2 induced   P3 induced

⬚ Spheres
▪ Organoids

Mll1^F/+ Mll1^F/F Mll1^FC/+ Mll1^FC/FC Mll1^FC/+ Mll1^FC/FC Mll1^FC/+ Mll1^FC/FC

D

▪ Mll1^FC/+; RC/+ ▪ Mll1^FC/FC; RC/+

Relative to *Rpl19* expression

Olfm4 **   Pitx2 *   Foxa1 *   Jaml **   Cxadr ***   Mmp7 *   Wnt3 *   Muc2 ***

Fig 7.