Phylogenomics of a new fungal phylum reveals multiple waves of reductive evolution across Holomycota

Luis Javier Galindo[1], Purificación López-García[1], Guifré Torruella[1], Sergey Karpov[2,3] and David Moreira[1]

[1]Ecologie Systématique Evolution, CNRS, Université Paris-Saclay, AgroParisTech, Orsay, France.

[2]Zoological Institute, Russian Academy of Sciences, St. Petersburg, 199034, Russia.

[3]St. Petersburg State University, St. Petersburg, 199034, Russia.

* Corresponding authors

 E-mails: david.moreira@u-psud.fr; luisjagg92@gmail.com

Abstract

Compared to multicellular fungi and unicellular yeasts, unicellular fungi with free-living flagellated stages (zoospores) remain poorly known and their phylogenetic position is often unresolved. Recently, rRNA gene phylogenetic analyses of two atypical parasitic fungi with amoeboid zoospores and long kinetosomes, the sanchytrids *Amoeboradix gromovi* and *Sanchytrium tribonematis*, showed that they formed a monophyletic group without close affinity with known fungal clades. Here, we sequence single-cell genomes for both species to assess their phylogenetic position and evolution. Phylogenomic analyses using different protein datasets and a comprehensive taxon sampling result in an almost fully-resolved fungal tree, with Chytridiomycota as sister to all other fungi, and sanchytrids forming a well-supported, fast-evolving clade sister to Blastocladiomycota. Comparative genomic analyses across fungi and their allies (Holomycota) reveal an atypically reduced metabolic repertoire for sanchytrids. We infer three main independent flagellum losses from the distribution of over 60 flagellum-specific proteins across Holomycota. Based on sanchytrids' phylogenetic position and unique traits, we propose the designation of a novel phylum, Sanchytriomycota. In addition, our results indicate that most of the hyphal morphogenesis gene repertoire of multicellular fungi had already evolved in early holomycotan lineages.

Key words: Fungi, Holomycota, phylogenomics, Sanchytriomycota, flagella, hyphae

Introduction

Within the eukaryotic supergroup Opisthokonta, multicellularity evolved independently in Fungi and Metazoa from unicellular ancestors along their respective branches, Holomycota and Holozoa[1]. The ancestor of this major supergroup originated ~1-1,5 Ga ago[2–4] and likely possessed one posterior flagellum for propulsion in aquatic environments[1]. This character has been retained in many modern fungal lineages at least during some life cycle stages[5,6]. Along the holomycotan branch, the free-living, non-flagellated nucleariid amoebae were the first to diverge, followed by the flagellated, phagotrophic, endoparasitic Rozellida (Cryptomycota)[7–9] and Aphelida[10], and the highly reduced, non-flagellated Microsporidia[11,12]. Aphelids branch as sister lineage to *bona fide*, osmotrophic, Fungi[13] (i.e., the zoosporic Blastocladiomycota and Chytridiomycota, and the non-zoosporic Zoopagomycota, Mucoromycota, Glomeromycota, and Dikarya). Within fungi, except for the secondary flagellar loss in the chytrid *Hyaloraphydium curvatum* [14], all known early divergent taxa are zoosporic (having at least one flagellated stage).

Zoosporic fungi are widespread across ecosystems, from soils to marine and freshwater systems, from tropical to Artic regions[15,16]. They include highly diverse saprotrophs and/or parasites, participating in nutrient recycling through the "mycoloop"[17–20]. Initially considered monophyletic, zoosporic fungi were recently classified into Blastocladiomycota and Chytridiomycota[21], in agreement with multigene molecular phylogenies[13,21,22]. These two lineages appear sister to the three main groups of non-flagellated fungi, Zoopagomycota, Mucoromycota and Dikarya, for which a single ancestral loss of the flagellum has been proposed[23]. Characterizing the yet poorly-known zoosporic fungi is important to understand the evolutionary changes (e.g., flagellum loss, hyphae development) that mediated land colonization and the adaptation of fungi to plant-dominated terrestrial ecosystems[24,25]. This requires a well-resolved phylogeny of fungi including all zoosporic lineages. Unfortunately, previous phylogenomic analyses did not resolve which zoosporic group, either Blastocladiomycota or Chytridiomycota, is sister to non-flagellated fungi[21,26–28]. This lack of resolution may result from the old age of these splits (~0,5-1 Ga)[3,29,30] and the existence of several radiations of fungal groups, notably during their co-colonization of land with plants[25,31], which would leave limited phylogenetic signal to resolve these deep nodes[22].

Because of this phylogenetic uncertainty, the number and timing of flagellum losses in fungi remain under debate, with estimates ranging between four and six for the whole

Holomycota[32]. Improving taxon sampling with new, divergent zoosporic fungi can help resolving

these deep nodes. One of such lineages is the sanchytrids, a group of chytrid-like parasites of algae

with uncertain phylogenetic position represented by the genera *Amoeboradix* and *Sanchytrium*,

which exhibit a highly reduced flagellum with an extremely long kinetosome[33,34] (Fig. 1).

Determining the phylogenetic position of this zoosporic lineage remains decisive to infer the

history of flagellum losses and the transition to hyphal-based multicellularity.

In this work we generate the first genome sequences for the sanchytrids *Amoeboradix*

gromovi and *Sanchytrium tribonematis* and analyse them together with available genomic and

transcriptomic data for Chytridiomycota and Blastocladiomycota. We obtain an almost fully-

resolved phylogeny of fungi which shows sanchytrids as a new fast-evolving lineage sister to

Blastocladiomycota. Contrasting with previous weakly-supported analyses[13,22,28,35,36], we robustly

place the root of the fungal tree between chytrids and all other fungi. Our new phylogenomic

framework of Fungi supports a conservative model of three flagellum losses in Holomycota and

highlights the importance of early-diverging unicellular Holomycota in the evolution of hyphal-

based multicellularity.

Results and discussion

The new zoosporic fungal phylum Sanchytriomycota

We isolated individual sporangia of *Amoeboradix gromovi* and *Sanchytrium tribonematis* (Fig. 1)

by micromanipulation and sequenced their genomes after whole genome amplification. After

thorough data curation (see Methods), we assembled two high-coverage genome sequences

(123.9X and 45.9X, respectively) of 10.5 and 11.2 Mbp, encoding 7,220 and 9,638 proteins,

respectively (Table 1). Comparison with a fungal dataset of 290 near-universal single-copy

orthologs[37] indicated very high completeness for the two genomes (92.41% for *A. gromovi*;

91.72% for *S. tribonematis*). Only half of the predicted sanchytrid proteins could be functionally

annotated using EggNOG[38] (3,838 for *A. gromovi*; 4,772 for *S. tribonematis*) (Supplementary Data

1). This could be partly due to the fact that, being fast-evolving parasites[33,34,39], many genes have

evolved beyond recognition by annotation programs[40,41]. However, low annotation proportions are

common in Holomycota, including fast-evolving parasites (e.g., only 20% and 52% of the genes

of the microsporidia *Nosema parisii* and *Encephalitozoon cuniculi* could be assigned to Pfam

domains and GO terms[40,42]) but also the less fast-evolving metchnikovellids (*Amphiamblys* sp.,

45.6%), rozellids (*Rozella allomycis*, 64.9%; *Paramicrosporidium saccamoebae*, 66.7%) and

blastocladiomycetes (*Catenaria anguillulae*, 47.5%). Many of the non-annotated genes were

unique to sanchytrids as deduced from orthologous gene comparison with 57 other species,

including representatives of the other major fungal lineages, and several outgroups

(Supplementary Data 2). After clustering of orthologous proteins with OrthoFinder [43], we

identified 1217 that were only present in both *A. gromovi* and *S. tribonematis*. Their analysis using

eggNOG[38] resulted in only 93 proteins annotated. The remaining (93.4%) sanchytrid-specific

proteins lack functional annotation (Supplementary Data 2).

The two sanchytrid genomes yielded similar sequence statistics (Table 1) but showed

important differences with genomes from other well-known zoosporic fungi. They are 4-5 times

smaller than those of blastocladiomycetes (40-50 Mb) and average chytrids (~20 to 101 Mb), an

observation that extends to the number of protein-coding genes. Their genome G+C content

(~35%) is much lower than that of blastocladiomycetes and most chytrids (40-57%, though some

chytrids, like *Anaeromyces robustus*, may have values down to 16.3%[44,45]). Low G+C content

correlates with parasitic lifestyle in many eukaryotes[46]. In Holomycota, low G+C is observed in

microsporidian parasites and Neocallimastigomycota, both anaerobic and exhibiting reduced

mitochondrion-derived organelles[45,47], and in the aerobic parasite *R. allomycis*[9]. Although

sanchytrids are aerobic parasites with similar life cycles to those of blastocladiomycetes and

chytrids[33,34,39], their smaller genome size and G+C content suggest that they are more derived

parasites. This pattern is accompanied by a global acceleration of evolutionary rate (see below), a

trend also observed, albeit at lower extent, in *R. allomycis*[9,48–50].

The mitochondrial genomes of *S. tribonematis* and *A. gromovi* showed similar trends. Gene

order was highly variable (Supplementary Fig. 1), as commonly observed in Fungi[51], and their size

(24,749 and 27,055 bp, respectively) and G+C content (25.86% and 30.69%, respectively) were

substantially smaller than those of most other Fungi. However, despite these signs of reductive

evolution, most of the typical core mitochondrial genes were present, indicating that they have

functional mitochondria endowed with complete electron transport chains.

To resolve the previously reported unstable phylogenetic position of sanchytrids based on

rRNA genes[33], we carried out phylogenomic analyses on a manually curated dataset of 264

conserved proteins (93,743 amino acid positions)[13,49,52] using Bayesian inference (BI) and

maximum likelihood (ML) with the CAT[53] and PMSF[54] models of sequence evolution,

respectively. Both mixture models are known to alleviate homoplasy and long-branch attraction

(LBA) artefacts[22,53]. We used the same 59 species as above (dataset GBE59), including a wide

representation of holomycota plus two holozoa, two amoebae and one apusomonad as outgroups.

BI and ML phylogenomic analyses yielded the same tree topology for major fungal groups with

only minor changes in the position of terminal branches (Fig. 2a; Supplementary Fig. 2a-b). We

recovered maximum statistical support for both the monophyly of sanchytrids (*A. gromovi + S.*

tribonematis) and their sister position to Blastocladiomycota. Thus, sanchytrids form a new deep-

branching zoosporic fungal clade. Given their divergence and marked genomic differences with

their closest relatives (Blastocladiomycota), we propose to create the new phylum

Sanchytriomycota to accommodate these fungal species (see description below).

141 An updated phylogeny of Fungi

In addition to the monophyly of Sanchytriomycota and Blastocladiomycota, our analysis retrieved

chytrids as sister group to all other fungi with full Bayesian posterior probability (PP=1) and

smaller ML bootstrap support (BS=92%). However, this position of the root of fungi on the chytrid

branch became fully supported (BS=100%) after removing the fast-evolving sanchytrids in a new

dataset of 57 species and 93,421 amino acid positions (dataset GBE57; Supplementary Fig. 2c),

suggesting an LBA artefact on the tree containing the sanchytrids. In fact, as observed in rRNA

gene-based phylogenies[33], sanchytrids exhibited a very long branch (Fig. 2a), indicating a fast

evolutionary rate. Long branches associated with fast-evolving genomes, well-known in

Microsporidia[48] and other Holomycota, can induce LBA artefacts in phylogenetic analyses[55–57].

To further study if LBA affected the position of the fast-evolving sanchytrids and other fungi in

our tree, we carried out several tests. First, we introduced in our dataset additional long-branched

taxa, metchnikovellids and core Microsporidia, for a total of 74 species and 86,313 conserved

amino acid positions (dataset GBE74). Despite the inclusion of these LBA-prone fast-evolving

taxa, we still recovered the monophyly of sanchytrids and Blastocladiomycota (Supplementary

Fig. 2d). However, they were dragged to the base of Fungi, although with low support (BS 78%).

To confirm the possible impact of LBA on this topology, we removed the fast-evolving sanchytrids

(dataset GBE72; 72 species and 84,949 amino acid positions), which led to chytrids recovering

their position as sister group of all other fungi with higher support (BS 91%; Supplementary Fig.

2e), corroborating the LBA induced by the long sanchytrid branch.

Second, we tested the influence of fast-evolving sites by applying a slow-fast approach[58]

that progressively removed the fastest-evolving sites (in 5% steps) in the 59-species alignment.

The monophyly of sanchytrids and Blastocladiomycota obtained maximum support (BS >99%) in

all steps until only 20% of the sites remained, when the phylogenetic signal was too low to resolve

any deep-level relationship (Fig. 2b). This relationship was as strongly supported as a well-

accepted relationship, the monophyly of Dikarya. Similarly, the root of the fungal tree between

chytrids and the rest of fungi was supported (>90% bootstrap) until only 40% of the sites remained.

By contrast, a root between sanchytrids+Blastocladiomycota and the rest of fungi always received

very weak support (<26% bootstrap).

We then further tested the robustness of the position of chytrids using alternative topology

(AU) tests. For the dataset GBE59, these tests did not reject alternative positions for the divergence

of Chytridiomycota and Blastocladiomycota+Sanchytriomycota (p-values >0.05; Supplementary

Data 3), which likely reflected the LBA due the long sanchytrid branch. In fact, the position of

Blastocladiomycota at the base of fungi was significantly rejected (p-values <0.05; Supplementary

Data 3) after removing the fast-evolving sanchytrid clade (dataset GBE57).

Finally, we compared the results based on the GBE dataset with those based on a different

dataset. We decided to use the BMC dataset[59] (which includes 53 highly conserved proteins and

14,965 amino acid positions) because it was originally designed to study fast-evolving holomycota

(e.g. Microsporidia), which made it appropriate to deal with the fast-evolving sanchytrids. Using

the same taxon sampling of 59 species (BMC59), we recovered the same topology, particularly for

the deeper nodes, with full ML ultrafast and conventional bootstrap (BS=100) and Bayesian

posterior probability (PP=1) supports for both the monophyly of sanchytrids+blastocladiomycetes

and chytrids as the sister lineage to all other fungi (Fig. 2c; Supplementary Fig. 2f-h).

The origin of the conspicuous long branch exhibited by sanchytrids is unclear. It has been

shown that fast-evolving organisms, including those within Holomycota, tend to lack part of the

machinery involved in genome maintenance and DNA repair[50,60]. To verify if it was also the case

in sanchytrids, we searched in the two sanchytrid genomes 47 proteins involved in genome

maintenance and DNA repair that have been observed to be missing in several fast-evolving

budding yeasts[60]. Our results confirm that most of these genes are also absent in both sanchytrids

(Supplementary Fig. 3). However, they are also missing in the closely related short-branching

blastocladiomycete *A. macrogynus*, suggesting that the long sanchytrid branch is not only due to

the absence of these genes.

The relative position of Chytridiomycota or Blastocladiomycota as first branch to diverge

within Fungi has remained a major unresolved question[61]. If the earliest fungal split occurred ~1

billion years ago[30], the phylogenetic signal to infer it may have been largely eroded over time.

Likewise, if evolutionary radiations characterized early fungal evolution[22], the accumulation of

sequence substitutions during early diversification would have been limited. Both factors would

explain the difficulty to resolve the deepest branches of the fungal tree so far. Our results, based

on an improved gene and taxon sampling, provided strong support for the placement of chytrids

as sister clade to all other Fungi. This solid position of the root on the chytrid branch is additionally

consistent with the distribution of so-considered derived characters in Blastocladiomycota,

including sporic meiosis, relatively small numbers of carbohydrate metabolism genes and, in some

species, hyphal-like apical growing structures (*Allomyces*) and narrow sporangia exit tubes (e.g.,

Catenaria spp.)[5,62–64]. Despite the use of a large dataset, some branches remained unresolved, in

particular the position of Glomeromycota, sister either to Mucoromycota or Dikarya (Fig. 2a),

which has important implications related to their symbiotic adaptation to land plants[24,65,66].

Macroevolutionary trends in primary metabolism

To assess if sanchytrid metabolic capabilities are as reduced as suggested by their small genome

sizes, we inferred their metabolic potential in comparison with other major fungal clades as well

as other opisthokonts and amoebozoa as outgroup (43 species). We compared the EggNOG[38]-

annotated metabolic repertoires of holomycotan phyla by focusing on 1,158 orthologous groups

(Supplementary Data 4) distributed in eight primary metabolism categories. Unexpectedly, cluster

analyses based on the presence/absence of these genes did not group sanchytrids with canonical

fungi and their closest aphelid relatives (i.e. *Paraphelidium*), but with non-fungal parasites (*R.*

allomycis, Mitosporidium daphniae and *P. saccamoebae*) which show evidence of reductive

genome evolution[9,11,67] (Fig. 3a), suggesting gene loss-related convergence. An even further

metabolic reduction was observed in Neocallimastigomycota, gut-inhabiting symbiotic anaerobic

chytrids[68–70]. A principal coordinate analysis of the same gene matrix confirmed this result (Fig.

3b).

At a more detailed level, the main differences in the metabolic complement of sanchytrids
and of canonical fungi (+*Paraphelidium*) concerned the carbohydrate and lipid transport and
metabolism categories, for which sanchytrids clustered with rozellids (Supplementary Fig. 4). We
further pairwise-compared KEGG[71] orthologs of sanchytrids against *R. allomycis* and the
blastocladiomycete *A. macrogynus* (as representative of the sanchytrid closest canonical fungal
relatives). The KEGG metabolic maps of *A. gromovi* and *S. tribonematis* contained 1,222 and
1,418 orthologous groups, respectively, whereas those of *R. allomycis* and *A. macrogynus*
contained 845 and 4,860, respectively (Supplementary Fig. 5a-c). Blastocladiomycetes and
sanchytrids shared more similarities, including the maintenance of amino acid and nucleotide
metabolism and energy production with a complete electron transport chain, which were largely
lost in *Rozella*[9,13]. Nonetheless, a reductive trend in energy production pathways could be observed
in sanchytrid mitochondria, including the loss of ATP8, one F-type ATP synthase subunit that is
also absent or highly modified in several metazoans, including chaetognaths, rotifers, most bivalve
molluscs, and flatworms[72,73]. *S. tribonematis* also lacked the NADH dehydrogenase subunit
NAD4L (Supplementary Fig. 1), although this loss is unlikely to impact its capacity to produce
ATP since *R. allomycis*, which lacks not only ATP8 but also the complete NADH dehydrogenase
complex, still seems to be able to synthesize ATP[9].

Most carbohydrate-related metabolic pathways were retained in sanchytrids and canonical
fungi except for the galactose and inositol phosphate pathways, absent in both sanchytrids and
Rozella. Nonetheless, sanchytrids displayed a rich repertoire of carbohydrate-degrading enzymes
(Supplementary Figs. 6-10), most of them being likely involved in the degradation of algal cell
walls required for penetration into the host cells[13,74,75]. The most important difference with
canonical fungi concerned lipid metabolism, with the steroids and fatty acid metabolism missing
in sanchytrids and also in *Rozella*[9] (Supplementary Fig. 5d-i). Collectively, our data suggest that,
compared to Blastocladiomycota and other fungal relatives, sanchytrids have undergone a
metabolic reduction that seems convergent with that observed in the phylogenetically distinct
rozellid parasites.

Convergent reductive flagellum evolution in Holomycota
The loss of the ancestral opisthokont single posterior flagellum in terrestrial fungi[76,77] is thought
to have been involved in their adaptation to land environments[78]. The number and timing of

flagellum losses along the holomycotan branch remains to be solidly established. The flagellum is
completely absent in nucleariids[12,52] but is found in representatives of all other major holomycotan
clades to the exclusion of the highly derived Microsporidia. They include rozellids[79], aphelids[10],
and various canonical fungal groups, namely chytrids[70], blastocladiomycetes[80], *Olpidium*[21,26], and
sanchytrids[33,34,39], although the latter are atypical. Sanchytrid amoeboid zoospores have never been
observed swimming but gliding on solid surfaces via two types of pseudopods: thin filopodia
growing in all directions and a broad hyaline pseudopodium at the anterior end (Fig. 1). The
posterior flagellum, often described as a pseudocilium, drags behind the cell without being
involved in active locomotion[33,34]. Its basal body (kinetosome) and axoneme ultrastructure differs
from that of most flagellated eukaryotes. Instead of the canonical kinetosome with 9 microtubule
triplets and axonemes with 9 peripheral doublets + 2 central microtubules[81], sanchytrids exhibit
reduced kinetosomes (9 singlets in *S. tribonematis*; 9 singlets or doublets in *A. gromovi*) and
axonemes (without the central doublet and only 4 microtubular singlets)[33,34]. Despite this
substantial structural simplification, sanchytrid kinetosomes are among the longest known in
eukaryotes, up to 2.2 μm[33,34]. Such long but extremely simplified kinetosomes have not been
reported in any other zoosporic fungi, including Blastocladiomycota[33,34]. Some of them, including
P. sedebokerense[82], display amoeboid zoospores during the vegetative cycle, flagellated cells
being most likely gametes[83–85].

To better understand flagellar reduction and loss across Holomycota, we analysed 61
flagellum-specific proteins on a well-distributed representation of 43 flagellated and non-
flagellated species. Sanchytrids lacked several functional and maintenance flagellar components
(Fig. 4a), namely axonemal dyneins, single- and double-headed inner arm dyneins, all
intraflagellar transport proteins (IFT) of the group IFT-A and several of the group IFT-B.
Sanchytrid kinetosomes have also lost several components of the centriolar structure and tubulins,
including Centrin2, involved in basal body anchoring[86], and Delta and Epsilon tubulins, essential
for centriolar microtubule assembly and anchoring[87]. These losses (Fig. 4b) explain why
sanchytrids lack motile flagella. Cluster analyses based on the presence/absence of flagellar
components (Supplementary Fig. 11) showed sanchytrids at an intermediate position between
flagellated and non-flagellated lineages. Therefore, sanchytrids are engaged in an unfinished
process of flagellum loss, thereby providing an interesting model to study intermediate steps of
this reductive process.

In addition to sanchytrid reduction, between four and six independent flagellar losses have been inferred in Holomycota[32]. Our new, more robust phylogenetic framework (Fig. 2) allowed to infer three large independent flagellum losses, plus the ongoing one in sanchytrids (Fig. 4c). These losses occurred at the base of high-rank taxa: nucleariids, Microsporidia, and the Zoopagomycota+Mucoromycota+Dikarya clade. A possible fourth loss event occurred in *Hyaloraphidium curvatum*, an atypical non-flagellated fungus originally classified as a colourless green alga[14] and later reclassified within the Monoblepharidomycota[26,88]. Further analysis will be needed to confirm loss of flagellar components in this species. In addition, a putative fifth independent loss might have occurred in the Nephridiophagida, a clade of fungal parasites of insects and myriapods[89,90] without clear affinity with established fungal clades[90]. Recently, a possible relationship to chytrids has been suggested[91], though genomic or transcriptomic data to clarify their phylogenetic position are still missing.

Fungal "vision" and flagellum exaptation

Why do sanchytrids retain a non-motile flagellum with a simplified but very long kinetosome? Since the primary flagellar function has been lost in favour of amoeboid movement, other selective forces must be acting to retain this atypical structure for a different function in zoospores. In bacteria, the exaptation of the flagellum for new roles in mechanosensitivity[92,93] and wetness sensing[94] has been documented. Microscopy observations of sanchytrid cultures showed that the flagellum is rather labile and can be totally retracted within the cell cytoplasm, the long kinetosome likely being involved in this retraction capability[33,34]. Interestingly, a conspicuous curved rosary chain of lipid globules has been observed near the kinetosome in *A. gromovi* zoospores, often also close to mitochondria[33,34]. In the blastocladiomycete *B. emersonii*, similar structures tightly associated with mitochondria are known as "side-body complexes"[95]. *B. emersonii* possesses a unique bacterial type-1-rhodopsin+guanylyl cyclase domain fusion (BeGC1, 626 amino acids) which, together with a cyclic nucleotide-gated channel (BeCNG1), controls zoospore phototaxis in response to cGMP levels after exposure to green light[96]. BeGC1 was localized by immunofluorescence on the external membrane of the axoneme-associated lipid droplets, which function as an eyespot at the base of the flagellum and control its beating[96–98]. The BeGC1 fusion and the channel BeCNG1 proteins have also been found in other blastocladiomycetes (*A. macrogynus* and *C. anguillulae*).

Both *A. gromovi* and *S. tribonematis* possessed the BeGC1 fusion (532 and 535 amino

acids, respectively) and the gated channel BeCNG1 (Supplementary Fig. 12a-c). Therefore, this

fusion constitutes a shared trait in Blastocladiomycota and Sanchytriomycota. Despite some

ultrastructural differences and the need of functional studies to confirm their role, the presence of

lipid threads in the vicinity of the kinetosome and mitochondria, together with the BeGC1 and

BeCNG1 homologs, suggest the existence of a comparable light-sensing organelle in *Amoeboradix*

and *Sanchytrium*. We hypothesize that, as in *B. emersonii,* the sanchytrid reduced flagellum could

be involved in phototactic response, at least as a structural support for the lipid droplets.

Interestingly, sanchytrids showed considerably shorter branches in rhodopsin and guanylyl cyclase

domain phylogenetic trees (Supplementary Fig. 12a-b) than in multi-gene phylogenies (Fig. 2),

indicating that these proteins (and their functions) are subjected to strong purifying selection as

compared to other proteins encoded in their genomes.

Since rhodopsins capture light by using the chromophore retinal[99], we looked for the

carotenoid (β-carotene) biosynthesis enzymes[96,100] necessary for retinal production. Surprisingly,

the enzymes involved in the classical pathway (bifunctional lycopene cyclase/phytoene synthase,

phytoene dehydrogenase and carotenoid oxygenase)[96,100] were missing in both sanchytrid

genomes, suggesting that they are not capable to synthesize their own retinal (Supplementary Data

5). We only detected two enzymes (isopentenyl diphosphate isomerase and farnesyl diphosphate

synthase) that carry out early overlapping steps in biosynthesis of both sterol and carotenoids. By

contrast, the β-carotene biosynthesis pathway is widely distributed in Fungi, including chytrids

and blastocladiomycetes (*Allomyces* and *Blastocladiella*[96,101]). Therefore, sanchytrids, like most

heterotrophic eukaryotes, seem unable to synthesize β-carotene and must obtain carotenoids

through their diet[100]. Indeed, we detected all carotenoid and retinal biosynthesis genes in the

transcriptome of the yellow-brown alga *Tribonema gayanum* (Supplementary Data 5), their host

and likely retinal source during infection.

Evolution of multicellularity in Holomycota

Fungal multicellularity results from connected hyphae[102]. Diverse genes involved in hyphal

multicellularity were present in the ancestors of three lineages of unicellular fungi

(Blastocladiomycota, Chytridiomycota and Zoopagomycota; BCZ nodes)[103]. To ascertain whether

they were also present in other deep-branching Holomycota with unicellular members, we

reconstructed the evolutionary history of 619 hyphal morphogenesis proteins[103], grouped into 10

functional categories (see Methods; Supplementary Data 6; Supplementary Fig. 13). Our results

showed that most hyphal morphogenesis genes were not only present in the last common fungal

ancestor but also in other unicellular holomycotan relatives, indicating that they evolved well

before the origin of fungal multicellularity (Fig. 5a). This pattern could be observed for all

functional categories with the clear exception of the adhesion proteins, most of which only occur

in Dikarya (Supplementary Fig. 13), reinforcing previous conclusions that adhesion proteins

played a marginal role in the early evolution of hyphae[103] but highlighting their crucial role in

current (fruiting body-producing) hyphal-based multicellularity.

The common ancestor of sanchytrids and blastocladiomycetes possessed a high percentage

of hyphae-related proteins (88.4%, node B in Fig. 5a), though this ancestral repertoire became

secondarily reduced in sanchytrids (66.6%). Likewise, many yeasts, which are also secondarily

reduced organisms, retained most of the genetic repertoire needed for hyphal development. Many

of these proteins were also present in nucleariids, Rozellida-Microsporidia and Aphelida (nodes

N, R, and A in Fig. 5a and Supplementary Data 7). Consequently, clustering analysis based on the

presence/absence of hyphal morphogenesis proteins did not clearly segregate unicellular and

multicellular lineages (Fig. 5b) and retrieved very weak intragroup correlation (Fig. 5c). Our

results extend previous observations of hyphal morphogenesis genes from Fungi[103] to much more

ancient diversifications in the Holomycota. The holomycotan ancestor already possessed a rich

repertoire of proteins, notably involved in 'actin cytoskeleton' and 'microtubule-based transport',

that were later recruited for hyphal production (Supplementary Data 7). Most innovation

concerned the proteins involved in the 'cell wall biogenesis/remodeling' and 'transcriptional

regulation' functional categories, which expanded since the common ancestor of Aphelida and

Fungi. This pattern is consistent with the enrichment of gene duplications in these two categories

in all major fungal lineages[103]. Nevertheless, genome and transcriptome data remain very scarce

for Aphelida and we expect that part of these duplications will be inferred to be older when more

data for this sister lineage of Fungi become available.

Conclusions

We generated the first genome sequence data for the two known species of sanchytrids, a group of

atypical fungal parasites of algae. The phylogenetic analysis of two independent datasets of

conserved proteins showed that they form a new fast-evolving fungal phylum, the Sanchytriomycota, sister to the Blastocladiomycota. Our phylogenetic analyses also provided strong support for Chytridiomycota being the sister group to all other fungi. Sanchytrids have a complex life cycle that includes a flagellated phase (zoospores) with non-motile flagella that are engaged in an ongoing reductive process. The inclusion of sanchytrids and a wide taxon sampling of fungi in our multi-gene phylogeny allowed the inference of three large independent flagellum loss events across Holomycota. Interestingly, the sanchytrid residual flagellum endowed with a long kinetosome might represent an exaptation of this structure as support for a lipid organelle probably involved in light sensing. Our taxon-rich dataset of deep-branching Holomycota also provided evidence for a very ancient origin of most genes related to hyphal morphogenesis, well before the evolution of the multicellular fungal lineages.

Taxonomic appendix

Sanchytriomycota phyl. nov.

Monocentric thallus, epibiotic; usually amoeboid zoospores with longest-known kinetosome in fungi (1-2 μm) and immobile pseudocilium; centrosome in sporangium with two centrioles composed by nine microtubular singlets.

Index Fungorum ID: IF558519

Class **Sanchytriomycetes** (Tedersoo et al. 2018)[104] emend.

Diagnosis as for the phylum.

Order **Sanchytriales** (Tedersoo et al. 2018)[104] emend.

Diagnosis as for the phylum.

Family **Sanchytriaceae** (Karpov & Aleoshin, 2017)[39] emend.

Amoeboid zoospores with anterior lamellipodium producing subfilopodia, and lateral and posterior filopodia; with (rarely without) posterior pseudocilium; kinetosome composed of nine microtubule singlets or singlets/doublets, 1–2 μm in length. Zoospores attach to algal cell wall, encyst, and penetrate host wall with a short rhizoid. Interphase nuclei in sporangia have a

centrosome of two centrioles composed of nine microtubular singlets. Predominantly parasites of
freshwater algae.

Type: *Sanchytrium* (Karpov et Aleoshin 2017[39]) emend. Karpov, 2019[34].

Parasite of algae. Epibiotic, spherical to ovate, sporangia with one (rarely more) discharge papillae.
Amoeboid zoospores with anterior lamellipodium; with (rarely without) pseudocilium; contains
kinetosome composed of nine single microtubules 1–1.2 µm in length. Interphase nuclei in
sporangia have a centrosome with two orthogonal centrioles composed of nine microtubular
singlets and with an internal fibrillar ring.

Type: *Sanchytrium tribonematis* (Karpov et Aleoshin, 2017[39]) emend. Karpov, 2019[34].

Sanchytrium tribonematis (Karpov et Aleoshin, 2017[39]) emend. Karpov, 2019[34].

Round to ovate smooth sporangium, ~10 µm diameter, without or with one discharge papilla;
sessile on algal surface. Slightly branched rhizoid, almost invisible inside host. Amoeboid
zoospores 5.4 - 3.3 µm (maximum) with anterior lamellipodium producing subfilopodia, lateral
and posterior filopodia; normally with posterior pseudocilium up to 5 µm in length supported by
up to four microtubules.

Amoeboradix Karpov, López-García, Mamkaeva & Moreira, 2018[33].

Zoosporic fungus with monocentric, epibiotic sporangia and amoeboid zoospores having posterior
pseudocilium that emerges from long kinetosome (ca. 2 µm) composed of microtubular singlets
or doublets.

Type species: *A. gromovi* Karpov, López-García, Mamkaeva & Moreira, 2018[33].

A. gromovi Karpov, López-García, Mamkaeva & Moreira, 2018[33].

Amoeboid zoospores of 2-3×3-4 µm with posterior pseudocilium up to 8 µm in length, which can
be totally reduced; kinetosome length, 1.8-2.2 µm. Sporangia of variable shape, from pear-shaped
with rounded broad distal end and pointed proximal end to curved asymmetrical sac-like 8-10 µm
width 16-18 µm long with prominent rhizoid and 2-3 papillae for zoospore discharge; spherical
cysts about 3-4 µm in diameter; resting spore ovate and thick-walled 12×6 µm. Parasite of
Tribonema gayanum, T. vulgare and *Ulothrix tenerrima.*

Methods

Biological material. *Sanchytrium tribonematis* strain X-128 and *Amoeboradix gromovi* strain X-113, isolated from freshwater sampling locations in Russia[33,34], were maintained in culture with the freshwater yellow-green alga *Tribonema gayanum* Pasch. strain 20 CALU as host [39]. The algal host was grown in mineral freshwater medium at room temperature under white light. After inoculation with *Sanchytrium* or *Amoeboradix*, cultures were incubated for 2 weeks to reach the maximum infection level. We then collected both individual zoospores and sporangia full of moving zoospores by micromanipulation with an Eppendorf PatchMan NP2 micromanipulator using 19 μm VacuTip microcapillaries (Eppendorf) on an inverted Leica Dlll3000 B microscope. Sporangia were separated from the algal host cells using a microblade mounted on the micromanipulator. Zoospores and sporangia were washed 2 times in clean sterile water drops before storing them into individual tubes for further analyses.

Whole genome amplification and sequencing. DNA extraction from single zoospores and sporangia was done with the PicoPure kit (Thermo Fisher Scientific) according to the manufacturer's protocol. Whole genome amplification (WGA) was carried out by multiple displacement amplification with the single-cell REPLI-g kit (QIAGEN). DNA amplification was quantified using a Qubit fluorometer (Life Technologies). We retained WGA products that yielded high DNA concentration. As expected, WGA from sporangia (many zoospores per sporangium) yielded more DNA than individual zoospores and were selected for sequencing (K1-9_WGA for *A. gromovi*; SC-2_WGA for *S. tribonematis*). TruSeq paired-end single-cell libraries were prepared from these samples and sequenced on a HiSeq 2500 Illumina instrument (2 x 100 bp) with v4 chemistry. We obtained 121,233,342 reads (26,245 Mbp) for *A. gromovi* and 106,922,235 reads (21,384 Mbp) for *S. tribonematis*.

Genome sequence assembly, decontamination and annotation. Paired-end read quality was assessed with FastQC v0.11.9[105] before and after quality trimming. Illumina adapters were removed with Trimmomatic v0.32 in Paired-End mode[106], with the following parameters: ILLUMINACLIP:adapters.fasta:2:30:10LEADING:28 TRAILING:28 SLIDINGWINDOW:4:30. Trimmed paired-end reads were assembled using SPAdes v3.9.1 in single-cell mode[107]. This

produced assemblies of 48.7 and 37.1 Mb with 8,420 and 8,015 contigs for *A. gromovi* and *S.*

tribonematis, respectively. Elimination of contaminant contigs was carried out by a three-step

process. First, genome sequences were subjected to two rounds of assembly, before and after

bacterial sequence removal with BlobTools v0.9.19[108] (Supplementary Fig. 14). Second, open-

reading frames were predicted and translated from the assembled contigs using Transdecoder v2

(http:transdecoder.github.io) with default parameters and Cd-hit v4.6[109] with 100% identity to

produce protein sequences for *A. gromovi* and *S. tribonematis*. Finally, to remove possible

eukaryotic host (*Tribonema*) contamination, the predicted protein sequences were searched by

BLASTp[110] against two predicted yellow-green algae proteomes inferred from the *Tribonema*

gayanum transcriptome [13] and the *Heterococcus* sp. DN1 genome (PRJNA210954; also a member

of the Tribonematales). We excluded sanchytrid hits that were 100% or >95% identical to them,

respectively. Statistics of the final assembled genomes were assessed with QUAST v4.5[111] and

Qualimap v2.2.1[112] for coverage estimation. In total, we obtained 7,220 and 9,368 protein

sequences for *A. gromovi* and *S. tribonematis*, respectively (Table 1). To assess genome

completeness, we used BUSCO v2.0.1[37] on the decontaminated predicted proteomes with the

fungi_odb9 dataset of 290 near-universal single-copy orthologs. The inferred proteins were

functionally annotated with eggNOG mapper v2[38] using DIAMOND as the mapping mode and the

eukaryotic taxonomic scope. This resulted in 3,757 (*A. gromovi*) and 4,670 (*S. tribonematis*)

functionally annotated peptides for the predicted proteomes (Supplementary Data 1). To determine

the presence of sanchytrid-specific proteins we used OrthoFinder[43] to generate orthologous groups

using the proteomes of the 59 species used in the GBE59 dataset (Supplementary Data 2). After

extracting the orthogroups that were present only in both sanchytrid species, we functionally

annotated them with eggNOG mapper v2[38] (Supplementary Data 2). The mitochondrial genomes

of the two sanchytrid species were identified in single SPAdes-generated contigs using BLAST[113]

and annotated with MITOS v1[114]. We confirmed the quality of the obtained mitochondrial genome

assemblies by reassembling them with NOVOPlasty[115]. Additional BLAST searches were made

to verify missing proteins.

Phylogenomic analyses and single-gene phylogenies. A revised version of a 264 protein dataset

(termed GBE)[49] was used to reconstruct phylogenomic trees[13,52]. This dataset was updated with

sequences from the two sanchytrid genomes and all publicly available Blastocladiomycota

sequences. Sequences were obtained mainly from GenBank
(http://www.ncbi.nlm.nih.gov/genbank, last accessed November, 2019) and, secondarily, the Joint
Genome Institute (http://www.jgi.doe.gov/; last accessed May 2017). For details on the origin of
sequence data see Supplementary Data 8. Our updated taxon sampling comprises a total of 71
Opisthokonta (2 Holozoa and 69 Holomycota), 2 Amoebozoa and 1 Apusomonadida. Two datasets
with two different taxon samplings were prepared, one with all 74 species (GBE74) and one
without the long-branching core Microsporidia and metchnikovellids for a total of 59 species
(GBE59). In addition, these datasets were also constructed without the sanchytrids to test if they
induced LBA artefacts, for a total of 72 (GBE 72) and 57 (GBE57) species. Additionally, for the
taxon sampling of 59 species we used a second phylogenomic dataset of 53 highly conserved
proteins[59]; the BMC dataset (BMC59).

The proteins for these datasets were searched in the new species with tBLASTn[110],
incorporated into the individual protein datasets, aligned with MAFFT v7 [116] and trimmed with
TrimAl v1.2 with the automated1 option[117]. Alignments were visualized, manually edited and
concatenated with Geneious v6.0.6[118] and single-gene trees obtained with FastTree v2.1.7[119] with
default parameters. Single-gene trees were manually checked to identify and remove paralogous
and/or contaminating sequences. The concatenation of the clean trimmed 264 proteins of the GBE
dataset resulted in alignments containing 93,743 (GBE59) and 86,313 (GBE74) amino acid
positions. The alternative alignment without the sanchytrids contained 93,421 (GBE57) and
84,949 (GBE72) amino acid positions. The concatenation of the 53 proteins of the BMC59 dataset
resulted in an alignment of 14,965 amino acid positions. The Bayesian inference (BI) phylogenetic
trees of the GBE59 and BMC59 datasets were reconstructed using PhyloBayes-MPI v1.5[120] under
the CAT-Poisson and CAT-GTR models, respectively, with two MCMC chains and run for more
than 15,000 generations, saving one every 10 trees. Analyses were stopped once convergence
thresholds were reached (i.e. maximum discrepancy <0.1 and minimum effective size >100,
calculated using bpcomp) and consensus trees constructed after a burn-in of 25%. Maximum
likelihood (ML) phylogenetic trees were inferred with IQ-TREE v1.6 under the PMSF
approximation of the LG+C20+R8 (GBE59 and GBE57), LG+C20+F+R9 (GBE72),
LG+C20+F+R10 (GBE74) or LG+C60+R7 (BMC59) models with guide trees inferred with the
LG+R8, LG+F+R9, LG+F+R10 or LG+R7 models, respectively, selected with the IQ-TREE
TESTNEW algorithm as per the Bayesian information criterion (BIC). Statistical support was

generated with 1,000 ultrafast bootstraps[121] and 1,000 replicates of the SH-like approximate

likelihood ratio test[122]. All trees were visualized with FigTree[123]. Additionally, we reconstructed

an ML tree for BMC59 under the same model with 100 conventional bootstrap replicates.

To test alternative tree topologies we used Mesquite[124] to constrain the following

topologies: 1) chytrids as sister lineage of all other fungi (i.e., monophyly of

Blastocladiomycota+Sanchytriomycota+Zygomycota+Dikarya),   and   2)

Blastocladiomycota+Sanchytriomycota as sister lineage of all other fungi (i.e., monophyly of

Chytridiomycota+Zygomycota+Dikarya). The constrained topologies without branch lengths

were reanalysed with the -g option of IQ-TREE and the best-fitting model. AU tests were carried

out on the resulting trees for each taxon sampling with the -z and -au options of IQ-TREE.

Additionally, to minimize possible systematic bias due to the inclusion of fast-evolving sites in

our GBE59 dataset, we progressively removed the fastest evolving sites, 5% of sites at a time. For

that, among-site substitution rates were inferred using IQ-TREE under the -wsr option and the

best-fitting model for a total of 19 new data subsets (Supplementary Data 3). We then reconstructed

phylogenetic trees for all these subsets using IQ-TREE with the same best-fitting model as for the

whole dataset. To assess the support of the alternative topologies in the bootstrapped trees, we

used CONSENSE from the PHYLIP v3.695 package[125] and interrogated the UFBOOT file using

a Python script (M. Kolisko, pers. comm). Single-protein phylogenies were reconstructed by ML

using IQ-TREE with automatic best fit model selection.

Comparative analysis of primary metabolism. We carried out statistical multivariate analyses

to get insights into the metabolic capabilities of sanchytrids in comparison with other Holomycota.

We searched in both sanchytrids 1,206 eggNOG orthologous groups[38] corresponding to 8 primary

metabolism categories (Gene Ontology, GO). The correspondence between GO terms and primary

metabolism COGs used are the following: [C] Energy production and conversion (227 orthologs);

[G] Carbohydrate transport and metabolism (205 orthologs); [E] Amino acid transport and

metabolism (200 orthologs); [F] Nucleotide transport and metabolism (87 orthologs); [H]

Coenzyme transport and metabolism (94 orthologs); [I] Lipid transport and metabolism (201

orthologs); [P] Inorganic ion transport and metabolism (153 orthologs); and [Q] Secondary

metabolites biosynthesis, transport and catabolism (70 orthologs). From these categories, we

identified 1,158 orthologs (non-redundant among categories) in the sanchytrid genomes which

were shared among a set of 43 species, including 38 Holomycota, 2 Holozoa, 2 Amoebozoa, and 1 apusomonad (for the complete list, see Supplementary Data 4). We annotated the protein sets of these 43 species using eggNOG-mapper v2[38] with DIAMOND as mapping mode and the eukaryotic taxonomic scope. All ortholog counts were transformed into a presence/absence matrix (encoded as 0/1) and analysed with the R script[126] detailed in Torruella et al. (2018) in which similarity values between binary COG profiles of all species were calculated to create a complementary species-distance matrix. We then analysed this distance matrix using a Principal Coordinate Analysis (PCoA) and plotted binary COG profiles in a presence/absence heatmap. Clustering of the species and orthologs was done by Ward hierarchical clustering (on Euclidean distances for orthologs) of the interspecific Pearson correlation coefficients. The raw species clustering was also represented in a separate pairwise correlation heatmap colour-coded to display positive Pearson correlation values (0–1). Finally, COG categories of each primary metabolism were also analysed separately (categories C, E, F, G, H, I, P, and Q) using the same workflow (for more details see Torruella et al. 2018 and https://github.com/xgrau/paraphelidium2018). We compared in more detail the inferred metabolism of a subset of species (the two sanchytrids, *Rozella allomycis*, and *Allomyces macrogynus*). The annotation of these proteomes was done using BlastKOALA v2.2[127], with eukaryotes as taxonomy group and the genus_eukaryotes KEGG GENES database, and the annotations were uploaded in the KEGG Mapper Reconstruct Pathway platform[71] by pairs. First, we compared the two sanchytrid proteomes to confirm their similarity and, second, we compared them with the proteomes of *R. allomycis* and *A. macrogynus* to study their metabolic reduction. To explore if the long branch observed in sanchytrids could be explained by the absence of DNA repair and genome maintenance genes, we used a dataset of 47 proteins involved on these functions from a previous study on fast-evolving budding yeasts[60]. These proteins where blasted[113] against the proteomes of the two sanchytrid species, the blastoclad *A. macrogynus* and four species of fast- and slow-evolving yeasts from that study.

Homology searches and phylogenetic analysis of specific proteins. To assess the evolution of the flagellum in holomycotan lineages we used a dataset of over 60 flagellum-specific proteins [77,128–130] to examine a total of 43 flagellated and non-flagellated species within and outside the Holomycota. The flagellar toolkit proteins were identified using *Homo sapiens* sequences as Blast queries. Candidate proteins were then blasted against the non-redundant GenBank database to

confirm their identification and submitted to phylogenetic analysis by multiple sequence alignment with MAFFT[116], trimming with TrimAl[117] with the automated1 option, and tree reconstruction with FastTree[119]. After inspection of trees, we removed paralogs and other non-orthologous protein sequences. We excluded the proteins with no identifiable presence in any of the 43 species used in the analysis and encoded the presence/absence of the remaining ones in a 1/0 matrix. The native R heatmap function[126] was used to plot the flagellar proteome comparison between all species according to their presence/absence similarity profiles. To study the presence or absence of the fusion of the BeGC1 and BeCNG1 light-sensing proteins, we blasted them against the proteomes of *S. tribonematis* and *A. gromovi* using the *Blastocladiella emersonii* sequences (BeGC1: AIC07007.1; BeCNG1: AIC07008.1) as queries. We used the protein dataset of Avelar et al. (2014)[96] in which the authors blasted both the guanylyl-cyclase GC1 and the rhodopsin domains of the BeGC1 fusion protein of *B. emersonii* and the BeCNG1 proteins against a database of more than 900 genomes from eukaryotic and prokaryotic species all across the tree of life (see Table S1 of their manuscript). We used MAFFT to include the new sequences in a multiple sequence alignment for the BeCNG1 protein channel and separately for both domains of the BeGC1 fusion protein[96]. After trimming with TrimAl we reconstructed phylogenetic trees for the three datasets using IQ-TREE with the best-fitting models: LG+F+I+G4 for the rhodopsin domain and BeCNG1, and LG+G4 model for GC1 guanylyl cyclase domain. The resulting trees were visualized with FigTree[123]. To study the possible presence of a carotenoid synthesis pathway for retinal production in sanchytrids, we combined previous datasets for carotenoid biosynthesis and cleavage enzymes from one giant virus (ChoanoV1), two choanoflagellates and two haptophytes[100], together with the carotenoid biosynthesis enzymes found in *B. emersonii*[96]. This dataset also included three enzymes for early sterol and carotenoid biosynthesis (isoprenoid biosynthesis steps). The protein sequences from this dataset where blasted (BLASTp) against the proteomes of the two sanchytrids and their host *Tribonema gayanum*. Results were confirmed by blasting the identified hits to the NCBI non-redundant protein sequence database. To study the presence of proteins involved in cell-wall penetration in sanchytrids (including cellulases, hemicellulases, chitinases, and pectinases), we used the complete mycoCLAP v1 database of carbohydrate-degrading enzymes (https://mycoclap.fungalgenomics.ca/mycoCLAP/). We blasted this database against our two sanchytrid proteomes and other two representatives of Blastocladiomycota and four Chytridiomycota. We then screened for canonical cellulose, hemicellulose, pectin and chitin

degrading enzymes identified in Fungi from previous studies[13,131,132]. The identified hits were

blasted back to both the NCBI non-redundant protein sequence database and the CAZy database

(cazy.org) to create a dataset for phylogenetic reconstruction for each enzyme. We added the

identified sequences to specific protein datasets from previous studies in the case of cellulases[13]

and of the chitin degradation proteins GH20 β -N-acetylhexosaminidase (NAGase)[75] and GH18

chitinase[74]. Proteins were aligned with MAFFT[116] and trimmed from gaps and ambiguously

aligned sites using TrimAl[117] with the automated1 option. ML trees were inferred using IQ-

TREE[133] with the best-fitting model selected with the IQ-TREE TESTNEW algorithm as per BIC.

The best-scoring tree was searched for up to 100 iterations, starting from 100 initial parsimonious

trees; statistical supports were generated from 1000 ultra-fast bootstrap replicates and 1000

replicates of the SH-like approximate likelihood ratio test. Trees were visualized with FigTree[123].

Analyses of hyphae multicellularity-related genes. We used a dataset of 619 hyphal

multicellularity-related proteins belonging to 362 gene families[103]. These gene families were

grouped into 10 functional categories: actin cytoskeleton (55 proteins), adhesion (30), polarity

maintenance (107), cell wall biogenesis/remodeling (92), septation (56), signaling (82),

transcriptional regulation (51), vesicle transport (103), microtubule-based transport (32) and cell

cycle regulation (11). The 619 proteins were searched by BLAST in the GBE59 dataset proteome

and later incorporated into individual gene family protein alignments with MAFFT v7[116]. After

trimming with TrimAl[117] with the automated1 option, alignments were visualized with Geneious

v6.0.6[118] and single gene trees obtained with FastTree[119] with default parameters. Single protein

trees were manually checked to identify paralogous sequences and confirm the presence of genes

within large gene families. Presence/absence of genes was binary coded (0/1) for each species and

the resulting matrix processed with a specific R script[13]. Finally, we reconstructed the

multicellularity-related gene gain/loss dynamics by applying the Dollo parsimony method

implemented in Count v10.04[134]. These analyses were done on both the complete dataset of all

hyphae multicellularity-related genes and on 10 separate datasets corresponding to the functional

categories mentioned above.

Data availability

The raw sequence data and assembled genomes generated in this study have been deposited at the National Center for Biotechnology Information (NCBI) sequence databases under Bioproject accession codes PRJNA668693 [https://www.ncbi.nlm.nih.gov/bioproject/PRJNA668693] and PRJNA668694 [https://www.ncbi.nlm.nih.gov/bioproject/PRJNA668694]. Additional data generated in this study (including alignments, phylogenetic trees and assembled genomes) are available in the Figshare repository project 91439 [https://figshare.com/projects/Sanchytriomycota_Galindo_etal/91439]. DNA and protein sequences of the species used in this study were downloaded from the GenBank public databases nr [https://www.ncbi.nlm.nih.gov/nucleotide/ and https://www.ncbi.nlm.nih.gov/protein/], genome [https://www.ncbi.nlm.nih.gov/genome/], SRA [https://www.ncbi.nlm.nih.gov/sra/], the JGI genome database [https://genome.jgi.doe.gov/portal/], the CAZy database [cazy.org], and the mycoCLAP database [https://mycoclap.fungalgenomics.ca/mycoCLAP/], for more details see Supplementary Data 8.

Acknowledgments

We thank the UNICELL single-cell genomics platform (https://www.deemteam.fr/en/unicell). This work was funded by the European Research Council Advanced Grants ProtistWorld (No. 322669, P.L.-G.) and Plast-Evol (No. 787904, D.M.) and the Horizon 2020 research and innovation program under the Marie Skłodowska-Curie ITN project SINGEK (http://www.singek.eu/; grant agreement No. H2020-MSCA-ITN-2015-675752, L.J.G. & P.L.-G.). S.A.K. contribution was supported by the RSF grant No. 16-14-10302. We thank Christina Cuomo and the Broad Institute for allowing the use of unpublished fungal genome sequences in this study.

Author contributions

P.L.-G. and D.M. conceived and supervised the study. S.A.K. and D.M. prepared the biological material; L.J.G., G.T. and D.M. analysed the sequence data; L.J.G., P.L.-G. and D.M. supervised the study and wrote the manuscript, which was edited and approved by all authors.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to L.J.G or D.M.

References

1. Cavalier-Smith, T. A revised six-kingdom system of life. *Biol. Rev. Camb. Philos. Soc.* **73**, 203–266 (1998).

2. Douzery, E. J. P., Snell, E. A., Bapteste, E., Delsuc, F. & Philippe, H. The timing of eukaryotic evolution: Does a relaxed molecular clock reconcile proteins and fossils? *Proc. Natl. Acad. Sci. U. S. A.* **101**, 15386–15391 (2004).

3. Parfrey, L. W., Lahr, D. J. G., Knoll, A. H. & Katz, L. A. Estimating the timing of early eukaryotic diversification with multigene molecular clocks. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 13624–13629 (2011).

4. Eme, L., Sharpe, S. C., Brown, M. W. & Roger, A. J. On the Age of Eukaryotes: Evaluating Evidence from Fossils and Molecular Clocks. *Cold Spring Harb. Perspect. Biol.* **6**, 1–16 (2014).

5. Berbee, M. L., James, T. Y. & Strullu-Derrien, C. Early Diverging Fungi: Diversity and Impact at the Dawn of Terrestrial Life. *Annu. Rev. Microbiol.* **71**, 41–60 (2017).

6. Spatafora, J. W. *et al.* The Fungal Tree of Life: from Molecular Systematics to Genome-Scale Phylogenies. *Microbiol. Spectr.* **5**, 1–32 (2017).

7. Lara, E., Moreira, D. & López-García, P. The Environmental Clade LKM11 and Rozella Form the Deepest Branching Clade of Fungi. *Protist* **161**, 116–121 (2010).

8. Jones, M. D. M. *et al.* Discovery of novel intermediate forms redefines the fungal tree of life. *Nature* **474**, 200–203 (2011).

9. James, T. Y. *et al.* Shared signatures of parasitism and phylogenomics unite cryptomycota and microsporidia. *Curr. Biol.* **23**, 1548–1553 (2013).

10. Karpov, S. A. *et al.* Morphology, phylogeny, and ecology of the aphelids (Aphelidea, Opisthokonta) and proposal for the new superphylum Opisthosporidia. *Frontiers in Microbiology* **5,** 112 (2014) doi:10.3389/fmicb.2014.00112.

11. Haag, K. L. *et al.* Evolution of a morphological novelty occurred before genome compaction in a lineage of extreme parasites. *Proc. Natl. Acad. Sci.* **111**, 15480–15485 (2014).

12. Bass, D. *et al.* Clarifying the Relationships between Microsporidia and Cryptomycota. *J.*

Eukaryot. Microbiol. **65**, 773–782 (2018).

13.   Torruella, G. *et al.* Global transcriptome analysis of the aphelid Paraphelidium tribonemae

supports the phagotrophic origin of fungi. *Commun. Biol.* **1**, 1–11 (2018).

14.   Ustinova, I., Krienitz, L. & Huss, V. A. R. Hyaloraphidium curvatum is not a green alga but

a lower fungus; Amoebidium parasiticum is not a fungus, but a member of the DRIPs.

Protist **151**, 253–262 (2000).

15.   Powell, M. J. Looking at mycology with a Janus face: a glimpse at Chytridiomycetes active

in the environment. *Mycologia* **85**, 1–20 (1993).

16.   Freeman, K. R. *et al.* Evidence that chytrids dominate fungal communities in high-elevation

soils. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 18315–18320 (2009).

17.   Kagami, M., Miki, T. & Takimoto, G. Mycoloop: Chytrids in aquatic food webs. *Front.*

Microbiol. **5**, 166 (2014).

18.   Frenken, T. *et al.* Warming accelerates termination of a phytoplankton spring bloom by

fungal parasites. *Glob. Chang. Biol.* **22**, 299–309 (2016).

19.   Frenken, T. *et al.* Integrating chytrid fungal parasites into plankton ecology: research gaps

and needs. *Environmental Microbiology* **19,** 3802–3822 (2017) doi:10.1111/1462-

2920.13827.

20.   Tedersoo, L., Bahram, M., Puusepp, R., Nilsson, R. H. & James, T. Y. Novel soil-inhabiting

clades fill gaps in the fungal tree of life. *Microbiome* **5**, 42 (2017).

21.   James, T. Y. *et al.* A molecular phylogeny of the flagellated fungi (Chytridiomycota) and

description of a new phylum (Blastocladiomycota). *Mycologia* **98**, 860–871 (2006).

22.   Chang, Y. *et al.* Phylogenomic analyses indicate that early fungi evolved digesting cell

walls of algal ancestors of land plants. *Genome Biol. Evol.* **7**, 1590–1601 (2015).

23.   Liu, Y. J., Hodson, M. C. & Hall, B. D. Loss of the flagellum happened only once in the

fungal lineage: Phylogenetic structure of Kingdom Fungi inferred from RNA polymerase II

subunit genes. *BMC Evol. Biol.* **6**, 1–13 (2006).

24.   Bidartondo, M. I. *et al.* The dawn of symbiosis between plants and fungi. *Biol. Lett.* **7**, 574–

577 (2011).

25.   Lutzoni, F. *et al.* Contemporaneous radiations of fungi and plants linked to symbiosis. *Nat.*

Commun. **9**, 1–11 (2018).

26.   Sekimoto, S., Rochon, D., Long, J. E., Dee, J. M. & Berbee, M. L. A multigene phylogeny

of Olpidium and its implications for early fungal evolution. *BMC Evol. Biol.* **11**, 331 (2011).

27.   Spatafora, J. W. *et al.* A phylum-level phylogenetic classification of zygomycete fungi
based on genome-scale data. *Mycologia* **108**, 1028–1046 (2016).

28.   Li, Y. *et al.* A genome-scale phylogeny of the kingdom Fungi. *Curr. Biol.* **31**, 1653–1665
(2021).

29.   Taylor, J. W. & Berbee, M. L. Dating divergences in the Fungal Tree of Life: Review and
new analyses. *Mycologia* **98,** 838–849 (2006) doi:10.3852/mycologia.98.6.838.

30.   Loron, C. C. *et al.* Early fungi from the Proterozoic era in Arctic Canada. *Nature* **570,** 232–
235 (2019) doi:10.1038/s41586-019-1217-0.

31.   Ebersberger, I. *et al.* A consistent phylogenetic backbone for the fungi. *Mol. Biol. Evol.* **29**,
1319–1334 (2012).

32.   James, T. Y. *et al.* Reconstructing the early evolution of Fungi using a six-gene phylogeny.
Nature **443**, 818–822 (2006).

33.   Karpov, S. A. *et al.* The Chytrid-like Parasites of Algae Amoeboradix gromovi gen. et sp.
nov. and Sanchytrium tribonematis Belong to a New Fungal Lineage. *Protist* **169**, 122–140
(2018).

34.   Karpov, S. A., Vishnyakov, A. E., Moreira, D. & López-García, P. The Ultrastructure of
Sanchytrium tribonematis (Sanchytriaceae, Fungi incertae sedis) Confirms its Close
Relationship to Amoeboradix. *J. Eukaryot. Microbiol.* **66**, 892–898 (2019).

35.   McCarthy, C. G. P. & Fitzpatrick, D. A. *Multiple Approaches to Phylogenomic
Reconstruction of the Fungal Kingdom. Advances in Genetics* vol. 100 (Elsevier Inc., 2017).

36.   Ahrendt, S. R. *et al.* Leveraging single-cell genomics to expand the fungal tree of life. *Nat.
Microbiol.* **3**, 1417–1428 (2018).

37.   Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.
BUSCO: Assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

38.   Huerta-Cepas, J. *et al.* Fast genome-wide functional annotation through orthology
assignment by eggNOG-mapper. *Mol. Biol. Evol.* **34**, 2115–2122 (2017).

39.   Karpov, S. A. *et al.* Monoblepharidomycetes diversity includes new parasitic and
saprotrophic species with highly intronized rDNA. *Fungal Biol.* **121**, 729–741 (2017).

40.   Cuomo, C. A. *et al.* Microsporidian genome analysis reveals evolutionary strategies for

obligate intracellular growth. *Genome Res.* **22**, 2478–2488 (2012).

41.   Nakjang, S. *et al.* Reduction and expansion inmicrosporidian genome evolution: New
insights from comparative genomics. *Genome Biol. Evol.* **5**, 2285–2303 (2013).

42.   Vivarès, C. P., Gouy, M., Thomarat, F. & Méténier, G. Functional and evolutionary analysis
of a eukaryotic parasitic genome. *Current Opinion in Microbiology* **5,** 499–505 (2002)
doi:10.1016/S1369-5274(02)00356-9.

43.   Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157
(2015).

44.   Billon-Grand, G., Fiol, J. B., Breton, A., Bruyère, A. & Oulhaj, Z. DNA of some anaerobic
rumen fungi: G + C content determination. *FEMS Microbiol. Lett.* **82**, 267–270 (1991).

45.   Youssef, N. H. *et al.* The genome of the anaerobic fungus Orpinomyces sp. strain c1a
reveals the unique evolutionary history of a remarkable plant biomass degrader. *Appl.*
Environ. Microbiol. **79**, 4620–4634 (2013).

46.   Videvall, E. Plasmodium parasites of birds have the most AT-rich genes of eukaryotes.
Microb. genomics. **4**, e000150 (2018).

47.   Chen, Y. ping *et al.* Genome sequencing and comparative genomics of honey bee
microsporidia, Nosema apis reveal novel insights into host-parasite interactions. *BMC*
Genomics **14**, 451 (2013).

48.   Thomarat, F., Vivarès, C. P. & Gouy, M. Phylogenetic analysis of the complete genome
sequence of Encephalitozoon cuniculi supports the fungal origin of microsporidia and
reveals a high frequency of fast-evolving genes. *J. Mol. Evol.* **59**, 780–791 (2004).

49.   Mikhailov, K. V., Simdyanov, T. G. & Aleoshin, V. V. Genomic survey of a hyperparasitic
microsporidian Amphiamblys sp. (Metchnikovellidae). *Genome Biol. Evol.* **9**, 454–467
(2017).

50.   Galindo, L. J. *et al.* Evolutionary genomics of metchnikovella incurvata
(metchnikovellidae): An early branching microsporidium. *Genome Biol. Evol.* **10**, 2736–
2748 (2018).

51.   Aguileta, G. *et al.* High variability of mitochondrial gene order among fungi. *Genome Biol.*
Evol. **6**, 451–465 (2014).

52.   Galindo, L. J. *et al.* Combined cultivation and single-cell approaches to the phylogenomics

| of nucleariid amoebae, close relatives of fungi. *Philos. Trans. R. Soc. B Biol. Sci.* **374**,
| 20190094 (2019).

| 53. Lartillot, N. & Philippe, H. A Bayesian mixture model for across-site heterogeneities in the
| amino-acid replacement process. *Mol. Biol. Evol.* **21**, 1095–1109 (2004).

| 54. Wang, H. C., Minh, B. Q., Susko, E. & Roger, A. J. Modeling site heterogeneity with
| posterior mean site frequency profiles accelerates accurate phylogenomic estimation. *Syst.*
| *Biol.* **67**, 216–235 (2018).

| 55. Leipe, D. D., Gunderson, J. H., Nerad, T. A. & Sogin, M. L. Small subunit ribosomal RNA+
| of Hexamita inflata and the quest for the first branch in the eukaryotic tree. *Molecular And*
| *Biochemical Parasitology*. **59,** 41–48 (1993).

| 56. Kamaishi, T. *et al.* Complete nucleotide sequences of the genes encoding translation
| Elongation Factors 1 and 2 from a microsporidian parasite, Glugea plecoglossi: Implications
| for the deepest branching of eukaryotes. *J. Biochem.* **120**, 1095–1103 (1996).

| 57. Philippe, H. *et al.* Early-branching or fast-evolving eukaryotes? An answer based on slowly
| evolving positions. *Proc. R. Soc. B Biol. Sci.* **267**, 1213–1221 (2000).

| 58. Brinkmann, H. & Philippe, H. Archaea sister group of Bacteria? Indications from tree
| reconstruction artifacts in ancient phylogenies. *Mol. Biol. Evol.* **16**, 817–825 (1999).

| 59. Capella-Gutiérrez, S., Marcet-Houben, M. & Gabaldón, T. Phylogenomics supports
| microsporidia as the earliest diverging clade of sequenced fungi. *BMC Biol.* **10**, 47 (2012).

| 60. Steenwyk, J. L. *et al.* Extensive loss of cell-cycle and DNA repair genes in an ancient
| lineage of bipolar budding yeasts. *PLoS Biol.* **17**, e3000255 (2019).

| 61. Naranjo-Ortiz, M. A. & Gabaldón, T. Fungal evolution: diversity, taxonomy and phylogeny
| of the Fungi. *Biol. Rev.* **94**, 2101–2137 (2019).

| 62. Vargas, M. M., Aronson, J. M. & Roberson, R. W. The cytoplasmic organization of hyphal
| tip cells in the fungus Allomyces macrogynus. *Protoplasma* **176**, 43–52 (1993).

| 63. Stajich, J. E. *et al.* The Fungi. *Current Biology* **19,** 840–845 (2009)
| doi:10.1016/j.cub.2009.07.004.

| 64. Archibald, J. M., Simpson, A. G. B. & Slamovits, C. H. *Handbook of the Protists*. *Handbook*
| *of the Protists* (Springer International Publishing, 2017). doi:10.1007/978-3-319-32669-6.

| 65. Field, K. J. *et al.* First evidence of mutualism between ancient plant lineages
| (Haplomitriopsida liverworts) and Mucoromycotina fungi and its response to simulated

Palaeozoic changes in atmospheric CO2. *New Phytol.* **205**, 743–756 (2015).

66.   Feijen, F. A. A., Vos, R. A., Nuytinck, J. & Merckx, V. S. F. T. Evolutionary dynamics of

mycorrhizal symbiosis in land plant diversification. *Sci. Rep.* **8**, 10698 (2018).

67.   Corsaro, D. *et al.* Microsporidia-like parasites of amoebae belong to the early fungal lineage

Rozellomycota. *Parasitol. Res.* **113**, 1909–1918 (2014).

68.   Ho, Y. W. & Barr, D. J. S. Classification of Anaerobic Gut Fungi from Herbivores with

Emphasis on Rumen Fungi from Malaysia. *Mycologia* **87**, 655–677 (1995).

69.   Powell, M. J. & Letcher, P. M. Chytridiomycota, Monoblepharidomycota, and

Neocallimastigomycota. in *Systematics and Evolution: Part A: Second Edition* 141 (2014).

doi:10.1007/978-3-642-55318-9.

70.   Powell, M. J. Chytridiomycota. in *Handbook of the Protists: Second Edition* (2017).

doi:10.1007/978-3-319-28149-0_18.

71.   Kanehisa, M. & Goto, S. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids*

Research **28,** 27–30 (1999) doi:10.1093/nar/27.1.29.

72.   Gissi, C., Iannelli, F. & Pesole, G. Evolution of the mitochondrial genome of Metazoa as

exemplified by comparison of congeneric species. *Heredity* **101,** 301–320 (2008)

doi:10.1038/hdy.2008.62.

73.   Egger, B., Bachmann, L. & Fromm, B. Atp8 is in the ground pattern of flatworm

mitochondrial genomes. *BMC Genomics* **26**, 414 (2017).

74.   Agrawal, Y., Khatri, I., Subramanian, S. & Shenoy, B. D. Genome sequence, comparative

analysis, and evolutionary insights into chitinases of entomopathogenic fungus Hirsutella

thompsonii. *Genome Biol. Evol.* **7**, 916–930 (2015).

75.   de Oliveira, E. S. *et al.* Molecular evolution and transcriptional profile of GH3 and GH20

β-n-acetylglucosaminidases in the entomopathogenic fungus Metarhizium anisopliae.

Genet. Mol. Biol. **41**, 843–857 (2018).

76.   Cavalier-Smith, T. & Chao, E. E. Y. Phylogeny of choanozoa, apusozoa, and other protozoa

and early eukaryote megaevolution. *J. Mol. Evol.* **56**, 540–563 (2003).

77.   Torruella, G. *et al.* Phylogenomics Reveals Convergent Evolution of Lifestyles in Close

Relatives of Animals and Fungi. *Curr. Biol.* **25**, 2404–2410 (2015).

78.   Naranjo-Ortiz, M. A. & Gabaldón, T. Fungal evolution: major ecological adaptations and

evolutionary transitions. *Biol. Rev.* **94**, 1443–1476 (2019).

79. Letcher, P. M. & Powell, M. J. A taxonomic summary and revision of Rozella (Cryptomycota). *IMA Fungus* **9**, 383–399 (2018).

80. Hibbett, D. S. *et al.* A higher-level phylogenetic classification of the Fungi. *Mycol. Res.* **111**, 509–547 (2007).

81. Mitchell, D. R. Speculations on the evolution of 9+2 organelles and the role of central pair microtubules. *Biology of the Cell* **96,** 691–696 (2004) doi:10.1016/j.biolcel.2004.07.004.

82. Strittmatter, M., Guerra, T., Silva, J. & Gachon, C. M. M. A new flagellated dispersion stage in Paraphysoderma sedebokerense, a pathogen of Haematococcus pluvialis. *J. Appl. Phycol.* **28**, 1553–1558 (2016).

83. Sparrow, F. K. Aquatic Phycomycetes. *Univ. Michigan Press* **132** (1960).

84. Letcher, P. M. *et al.* An ultrastructural study of Paraphysoderma sedebokerense (Blastocladiomycota), an epibiotic parasite of microalgae. *Fungal Biol.* **120**, 324–337 (2016).

85. Powell, M. J. Blastocladiomycota. in *Handbook of the Protists: Second Edition* 1497–1521 (2017). doi:10.1007/978-3-319-28149-0_17.

86. Aubusson-Fleury, A. *et al.* Centrin diversity and basal body patterning across evolution: New insights from Paramecium. *Biol. Open* **6**, 765–776 (2017).

87. Dupuis-Williams, P. *et al.* Functional role of ε-tubulin in the assembly of the centriolar microtubule scaffold. *J. Cell Biol.* **158**, 1183–1193 (2002).

88. Forget, L., Ustinova, J., Wang, Z., Huss, V. A. R. & Lang, B. F. Hyaloraphidium curvatum: A linear mitochondrial genome, tRNA editing, and an evolutionary link to lower fungi. *Mol. Biol. Evol.* **19**, 310–319 (2002).

89. Fabel, P., Radek, R. & Storch, V. A new spore-forming protist, Nephridiophaga blaberi sp. nov., in the Death's head cockroach Blaberus craniifer. *Eur. J. Protistol.* **36**, 387–395 (2000).

90. Radek, R. *et al.* Morphologic and molecular data help adopting the insect-pathogenic nephridiophagids (Nephridiophagidae) among the early diverging fungal lineages, close to the Chytridiomycota. *MycoKeys* **25**, 31–50 (2017).

91. Strassert, J. F. H. *et al.* Long rDNA amplicon sequencing of insect-infecting nephridiophagids reveals their affiliation to the Chytridiomycota and a potential to switch between hosts. *Sci. Rep.* **11**, 396 (2021).

92. Belas, R., Simon, M. & Silverman, M. Regulation of lateral flagella gene transcription in
Vibrio parahaemolyticus. *J. Bacteriol.* **167**, 210–218 (1986).

93. Belas, R. Biofilms, flagella, and mechanosensing of surfaces by bacteria. *Trends Microbiol.*
22, 517–527 (2014).

94. Wang, Q., Suzuki, A., Mariconda, S., Porwollik, S. & Harshey, R. M. Sensing wetness: A
new role for the bacterial flagellum. *EMBO J.* **24**, 2034–2042 (2005).

95. Lovett, J. S. Growth and differentiation of the water mold Blastocladiella emersonii:
cytodifferentiation and the role of ribonucleic acid and protein synthesis. *Bacteriol. Rev.* **39**,
345–404 (1975).

96. Avelar, G. M. *et al.* A Rhodopsin-Guanylyl cyclase gene fusion functions in visual
perception in a fungus. *Curr. Biol.* **24**, 1234–1240 (2014).

97. Avelar, G. M. *et al.* A cyclic GMP-dependent K+ channel in the blastocladiomycete fungus
Blastocladiella emersonii. *Eukaryot. Cell* **14**, 958–963 (2015).

98. Richards, T. A. & Gomes, S. L. How to build a microbial eye. *Nature* **523**, 166–167 (2015).

99. Ernst, O. P. *et al.* Microbial and animal rhodopsins: Structures, functions, and molecular
mechanisms. *Chemical Reviews* . 114, 126-163 (2014) doi:10.1021/cr4003769.

100. Needham, D. M. *et al.* A distinct lineage of giant viruses brings a rhodopsin photosystem to
unicellular marine predators. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 20574–20583 (2019).

101. Wöstemeyer, J., Grünler, A., Schimek, C. & Voigt, K. Genetic regulation of carotenoid
biosynthesis in fungi. in *Applied Mycology and Biotechnology*. **5,** 257-274 (2005).
doi:10.1016/S1874-5334(05)80013-9.

102. Nagy, L. G., Kovács, G. M. & Krizsán, K. Complex multicellularity in fungi: evolutionary
convergence, single origin, or both? *Biol. Rev.* **93**, 1778–1794 (2018).

103. Kiss, E. *et al.* Comparative genomics reveals the origin of fungal hyphae and
multicellularity. *Nat. Commun.* **10**, (2019).

104. Tedersoo, L. *et al.* High-level classification of the Fungi and a tool for evolutionary
ecological analyses. *Fungal Divers.* **90**, 135–159 (2018).

105. Andrews, S. FastQC: A quality control tool for high throughput sequence data.
Http://Www.Bioinformatics.Babraham.Ac.Uk/Projects/Fastqc/
http://www.bioinformatics.babraham.ac.uk/projects/   (2010)   doi:citeulike-article-
id:11583827.

106. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

107. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

108. Laetsch, D. R. & Blaxter, M. L. BlobTools: Interrogation of genome assemblies. *F1000Research* **6**, 1287 (2017).

109. Li, W. & Godzik, A. Cd-hit: A fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

110. Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

111. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: Quality assessment tool for genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

112. Okonechnikov, K., Conesa, A. & García-Alcalde, F. Qualimap 2: Advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292–294 (2015).

113. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

114. Bernt, M. *et al.* MITOS: Improved de novo metazoan mitochondrial genome annotation. *Mol. Phylogenet. Evol.* **69**, 313–319 (2013).

115. Dierckxsens, N., Mardulyn, P. & Smits, G. NOVOPlasty: de novo assembly of organelle genomes from whole genome data. *Nucleic Acids Res.* **45**, e18 (2017).

116. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

117. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

118. Kearse, M. *et al.* Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649 (2012).

119. Price, M. N., Dehal, P. S. & Arkin, A. P. Fasttree: Computing large minimum evolution trees with profiles instead of a distance matrix. *Mol. Biol. Evol.* **26**, 1641–1650 (2009).

120. Lartillot, N., Lepage, T. & Blanquart, S. PhyloBayes 3: A Bayesian software package for phylogenetic reconstruction and molecular dating. *Bioinformatics* **25**, 2286–2288 (2009).

121. Minh, B. Q., Nguyen, M. A. T. & Von Haeseler, A. Ultrafast approximation for
phylogenetic bootstrap. *Mol. Biol. Evol.* **30**, 1188–1195 (2013).

122. Anisimova, M., Gil, M., Dufayard, J. F., Dessimoz, C. & Gascuel, O. Survey of branch
support methods demonstrates accuracy, power, and robustness of fast likelihood-based
approximation schemes. *Syst. Biol.* **60**, 685–699 (2011).

123. Rambaut, A. FigTree v1.4.3. *http://tree.bio.ed.ac.uk/software/figtree/* (2016).

124. Mesquite Project Team. Mesquite: A modular system for evolutionary analysis. *Available*
from: *http://*   *mesquiteproject.wikispaces.com/home*   (2014)
doi:10.1017/CBO9781107415324.004.

125. Felsenstein, J. PHYLIP - Phylogenetic Inference Package (Version 3.5). *Cladistics* **5,** 164–
166 (1989) doi:10.2987/10-6090.1.

126. R Development Core Team, R. *R: A Language and Environment for Statistical Computing.*
R Foundation for Statistical Computing, Vienna, Austria. Available online at
https://www.R-project.org/ (2018). doi:10.1007/978-3-540-74686-7.

127. Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG Tools
for Functional Characterization of Genome and Metagenome Sequences. *J. Mol. Biol.* **428**,
726–731 (2016).

128. Carvalho-Santos, Z., Azimzadeh, J., Pereira-Leal, J. B. & Bettencourt-Dias, M. Tracing the
origins of centrioles, cilia, and flagella. *Journal of Cell Biology* **194,** 165–175 (2011)
doi:10.1083/jcb.201011152.

129. Wickstead, B. & Gull, K. Evolutionary biology of dyneins. in *Dyneins* 88–121 (2012).
doi:10.1016/B978-0-12-382004-4.10002-0.

130. Van Dam, T. J. P. *et al.* Evolution of modular intraflagellar transport from a coatomer-like
progenitor. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 6943–6948 (2013).

131. Kubicek, C. P., Starr, T. L. & Glass, N. L. Plant Cell Wall - Degrading Enzymes and Their
Secretion in Plant-Pathogenic Fungi. *Annu Rev Phytopathol* **52**, 427–51 (2014).

132. Sista Kameshwar, A. K. & Qin, W. Systematic review of publicly available non-Dikarya
fungal proteomes for understanding their plant biomass-degrading and bioremediation
potentials. *Bioresources and Bioprocessing* **6,** 30 (2019) doi:10.1186/s40643-019-0264-6.

133. Nguyen, L. T., Schmidt, H. A., Von Haeseler, A. & Minh, B. Q. IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol.*

987   *Evol.* **32**, 268–274 (2015).

988 134. Csurös, M. Count: Evolutionary analysis of phylogenetic profiles with parsimony and

989   likelihood. *Bioinformatics* **26**, 1910–1912 (2010).

992 **Table 1.** Statistics of sanchytrid genome assemblies before and after decontamination and
993 comparison with related zoosporic lineages.

	Amoeboradix gromovi		Sanchytrium tribonematis		Allomyces macrogynus	Catenaria anguillulae	Rhizoclosmatium globosum	Spizellomyces punctatus	Anaeromyces robustus
	Before	After	Before	After					
Genome size (Mb)	48.7	10.5	37.1	11.2	52.62	41.34	57.02	24.1	71.69
GC%	51.06	36.27	42.9	34.64	61.6	56	44.9	47.6	16.3
Number of contigs	8,420	1,167	8,015	1,960	8,973	509	437	329	1,035
N50	27,236	13,376	17,350	11,874	35,497	217,825	292,246	155,888	141,798
Predicted proteins	87,868	7,220	50,780	9,368	19,447	12,763	16,987	9,422	12,083
Mt genome size (bp)	27,055		24,749		57,473	---	---	126,474	---
Mt genome GC%	30.69		25.86		39.5	---	---	29.88	---

Fig. 1. Light microscopy images of life cycle stages of *Sanchytrium tribonematis* (a-e) and *Amoeboradix gromovi* (f-j). a-d, f-i – amoeboid crawling zoospores with filopodia (f) and posterior pseudocilium (pc). **g-i** – zoospores with retracted pseudocilium. **e, j** – sporangium (sp) with one (**e**) or two (**j**) papillae (p) on the host (h) surface. Scale bars: **a-d, f-i** – 5 µm; **e, j** – 10 µm.

Fig. 2. **Phylogenomic analysis of Holomycota. a**, Bayesian inference (BI) phylogenomic tree based on the GBE dataset of 264 conserved proteins (93,743 amino acid positions). The tree was reconstructed using 59 species and the CAT-Poisson model and the PMSF approximation of the LG+R8+C20 model for maximum likelihood (ML). **b**, Evolution of IQ-TREE ML bootstrap support for Chytridiomycota sister to all other fungi (C+F), Blastocladiomycota+Sanchytriomycota sister to all other fungi (B+F), Sanchytriomycota within Blastocladiomycota (S+B), and the monophyly of Dikarya (Dikarya) as a function of the proportion of fast-evolving sites removed from the GBE59 dataset. Holomycota are highlighted in violet, sanchytrids in pink and outgroup taxa in other colors (green, orange and yellow). All phylogenomic trees can be seen in Supplementary Figs. 2a-e. **c**, Schematic BI phylogeny showing the results obtained with the BMC dataset of 53 conserved proteins (14,965 amino acid positions) using 59 species (Fungi: violet; Sanchytriomycota: red; outgroup taxa: black) and the CAT-GTR model (BI) and the PMSF approximation of the LG+R7+C60 model for maximum likelihood (ML). In both trees, branches with support values higher or equal to 0.99 BI posterior probability and 99% ML bootstrap are indicated by black dots.

Fig. 3. **Distribution patterns of primary metabolism genes in Holomycota. a**, Binary heat-map and **b**, principal coordinate analysis (PCoA) species clustering based on the presence/absence of 1,158 orthologous genes belonging to 8 primary metabolism Gene Ontology categories across 43 eukaryotic genomes and transcriptomes. Species are color- and shaped-coded according to their taxonomic affiliation as indicated in the legend. COG presence is depicted in blue and absence is depicted in white.

Fig. 4. **Comparison of flagellar protein distribution and structure in sanchytrids and other Holomycota. a**, Presence/absence heatmap of 61 flagellum-specific proteins in 43 eukaryotic

flagellated (purple; red: Sanchytriomycota) and non-flagellated (pink) lineages. The right column lists microtubular genes and flagellum-specific genes (orange). **b**, Illustration depicting the main structural elements present in a typical eukaryotic flagellum (left) and a reduced sanchytrid flagellum (right). Gene presence is depicted in blue and absence in white. **c**, Representation of the evolutionary relationships of holomycotan lineages and their patterns of presence/absence of key molecular components of the flagellar apparatus (full or half circles). Red crosses on branches indicate independent flagellum losses.

Fig. 5. Hyphae-related genes across Holomycota. a, Cladogram of Holomycota depicting the phylogenetic relationships according to the GBE59 phylogenomic reconstruction. Bubble size on the nodes and tips represents the total number of reconstructed ancestral and extant hyphal multicellularity-related proteins. CBZ (Chytridiomycota, Blastocladiomycota and Zoopagomycota) and NRA (Nucleariids, Rozellida-Microsporidia and Aphelida) nodes are indicated with letters within the corresponding bubbles. Unicellular lineages are highlighted in colors according to their taxonomic affiliation (bottom to top: Amoebozoa [yellow], Apusomonadida [light green], Holozoa [dark green], nucleariids [light blue], Rozellida-Microsporidia [purple], Aphelida [pink], Chytridiomycota [violet], Sanchytriomycota-Blastocladiomycota [red] and Zoopagomycota [orange]). **b**, Presence/absence heatmap of 619 hyphal morphogenesis proteins in 59 unicellular (taxa are color-highlighted as in Fig. 5a) and multicellular (not color-highlighted) eukaryotic proteomes. Gene presence is depicted in blue and absence in white. **c**, Heatmap clustered by similarity showing the correlation between unicellular (taxa are color-highlighted as in Fig. 5a) and multicellular (not color-highlighted) taxa according to the presence/absence of hyphal morphogenesis gene proteins.

Galindo et al. Fig. 1

a

Acanthamoeba castellanii — Amoebozoa
Dictyostelium discoideum
Thecamonas trahens — Apusomonadida
Monosiga brevicollis — Holozoa
Salpingoeca rosetta
Nuclearia thermophila — nucleariids
Parvularia atlantis
Fonticula alba
Rozella allomycis — Microsporidia + Rozellida
Paramicrosporidium saccamoebae
Mitosporidium daphniae
Paraphelidium tribonemae — Aphelida
Piromyces sp.
Anaeromyces robustus
Neocallimastix californiae
Pecoramyces ruminatium 0.51/100
Gonapodya prolifera
Synchytrium microbalum 1/76
Synchytrium taraxaci
Rhizoclosmatium globosum
Caulochytrium protostelioides
Spizellomyces punctatus 1/77
Blyttiomyces helicus
Batrachochytrium dendrobatidis
Homolaphlyctis polyrhiza — Chytridiomycota
Sanchytrium tribonematis — Sanchytriomycota
Amoeboradix gromovi
Allomyces macrogynus
Blastocladiella emersonii — Blastocladiomycota
Catenaria anguillulae
Basidiobolus meristosporus
Basidiobolus heterosporus
Conidiobolus coronatus 1/92
Entomophthora muscae
Massospora cicadina — Zoopagomycota
Piptocephalis cylindrospora 1/---
Linderina pennispora
Coemansia reversa 0.51/---
Capniomyces stellatus
Smittium culicis
Mortierella verticillata
Mortierella alpina
Umbelopsis isabellina
Phycomyces blakesleeanus — Mucoromycota
Rhizopus oryzae
Mucor circinelloides
Cokeromyces recurvatus
Glomus cerebriforme 1/75
Rhizophagus irregularis — Glomeromycota
Puccinia graminis
Ustilago maydis 1/---
Cryptococcus neoformans — Basidiomycota
Postia placenta
Laccaria bicolor
Saitoella complicata
Schizosaccharomyces pombe
Sclerotinia sclerotiorum — Ascomycota
Yarrowia lipolytica
Saccharomyces cerevisiae

"Zygomycota"
Dikarya
F u n g i

0.2

b

Bootstrap percentage

C+F   B+F   S+B   Dikarya

100 95 90 85 80 75 70 65 60 55 50 45 40 35 30 25 20 15 10 5
Percentage of data used for phylogenomic analyses

c

Outgroups
Chytridiomycota
Sanchytriomycota
Blastocladiomycota
Zoopagomycota
Glomeromycota
Mucoromycota
Dikarya

0.2

Galindo et al. Fig. 2

Galindo et al., Fig. 3

Galindo et al., Fig. 4

Galindo et al., Fig. 5