Microbial Metagenomic Approach Uncovers the First Rabbit Haemorrhagic Disease Virus genome in Sub-Saharan Africa

Anise N. Happi[1*], Olusola A. Ogunsanya[1], Judith U. Oguzie[2,3], Paul E. Oluniyi [2,3], Alhaji S. Olono[1], Jonathan L. Heeney[4], Christian T. Happi[2,3*]

1. Department of Veterinary Pathology, Faculty of Veterinary Medicine, University of Ibadan, Ibadan, Nigeria. Email: anisehappi@yahoo.com, shorlar7@gmail.com, halhaagy1@gmail.com

2. Department of Biological Sciences, Faculty of Natural Sciences, Redeemer's University, Ede, Nigeria.

3. African Centre of Excellence for Genomics of Infectious Diseases (ACEGID), Redeemer's University, Ede, Osun State, Nigeria. Email: oguziej@run.edu.ng, oluniyip@run.edu.ng, happic@run.edu.ng

4. Lab of Viral Zoonotics, Department of Veterinary medicine, University of Cambridge, Cambridge, UK. Email: jlh66@cam.ac.uk

* Corresponding authors: **Anise N. Happi**, Email: anisehappi@yahoo.com and **Christian T. Happi**, Email: happic@run.edu.ng

ABSTRACT

Rabbit Haemorrhagic Disease (RHD) causes high morbidity and mortality in rabbits and hares. Here, we report the first genomic characterization of Rabbit Hemorrhagic Disease Virus (RHDV) from sub-Saharan Africa. While suspected, only a single PCR finding without sequence confirmation or characterization has been reported. Here, we used a microbial metagenomic approach to confirm and characterize pathogens causing the suspected outbreak of RHD in Ibadan, Nigeria. On the 25th September 2020, the liver, spleen, and lung samples of five rabbits from an outbreak in 2 farms in Ibadan, Nigeria, were analyzed for the vp60 gene of RHDV by RT-PCR. Subsequently, Next Generation Sequencing on 1st of October   revealed one full and two partial RHDV2 genomes on both farms. Phylogenetic analysis showed close clustering with RHDV2 lineages from Europe, in particular, 98.6% similarity with RHDV2 in the Netherlands, and 99.1 to 100% identity with RHDV2 in Germany, suggesting potential importation from Europe. The detection of twelve unique mutations in RHDV2 sequences from the Ibadan outbreak compared to other RHDV2 sequences in the same clade suggests significant genetic diversity of the GI.2

strains in Nigeria. This highlights the need to further understand the genetic diversity of Lagoviruses to, inform vaccine development, and for accurate tracking, monitoring, and control of outbreaks in Africa.

Key words: Rabbit Haemorrhagic disease, Nigeria, RHDV2

Introduction

Rabbit haemorrhagic disease (RHD) is a highly infectious and deadly viral haemorrhagic disease of rabbits. The disease is caused by Rabbit haemorrhagic disease virus (RHDV) a *Lagovirus* of the *Caliciviridae* family[1]. The virus causes a high morbidity and mortality rates, killing more than 90% of infected adult animals in 2-3 days following infection[2,3]. This disease causes economic losses to the rabbit meat and fur industry and great negative ecological impact in wild rabbit populations[3, 4]. RHD is among the diseases notifiable to the World Organization for Animal Health (OIE).

Transmission is via nasal, oral, conjunctival routes with possible vector-borne transmission reported[5]. The virus is also shed through excretions from infected animals[6]. The lesions of RHD include petechial hemorrhages in multiple organs as a result of virus-induced hyper-coagulopathy. The most severe form of these lesions appears in the liver, trachea, and lungs[7]. RHDV also promotes fatal hepatitis in adult rabbits[8].

The RHDV genome is positive sense and single stranded RNA of approximately 7437 nucleotides in length[1]. There are two open reading frames (ORFs); ORF1 encoding the nonstructural proteins (RNA-dependent RNA polymerase and the major capsid protein VP60) and ORF2 encoding the VP10 protein a minor structural protein[9,10].

RHD was first reported in China in 1984[11]: where over 140 million rabbits were killed in the course of an outbreak[12]. Subsequent spread of the virus was reported in Europe and other continents[13-19]. In Africa, RHDV has been reported since the late 1980s[20], but molecular characterization and diversity of the circulating variants and their impact are yet to be understood.

There are six serogroups (G1-G6) RHDV[21] and one serotype[22] before 2010. These groups are known as "classical RHDV". A new strain RHDV2 (GI.2 or RHDVb) was discovered in France in 2010[23] and is now reported in many parts of Europe[22, 24-26] and other parts of the world[27, 28]. RHDV2 was responsible for major outbreaks causing deaths in previously vaccinated adult rabbits as well as younger rabbits known to be immune to classical RHD[22, 23, 29]. This strain is now reported in Europe, Australia, Canada and more recently the USA[27, 30, 31]. RHD has been reported in some African countries like Tunisia, Morocco, Cape Verde and Egypt[7, 32-34]. In sub-Saharan Africa, outbreaks of GI.2 were reported and confirmed by enzyme immunoassays (EIA) in domestic rabbits from Benin Republic[34], and PCR from Cote d'Ivoire[30]. However, to date there are no available virus genome data from any Sub-Saharan African country, rendering it difficult to trace their origin, evolution and genetic diversity.

Following devastating outbreaks of suspected cases of RHD affecting several farmers from the southwestern region of Nigeria, there was considerable loss of investment with no surviving rabbits in some of the establishments. During the decline in the number of cases in August 2020, two farms (Farm A and Farm B) reported high mortalities and

were visited to carry out post-mortem examinations and sampling for molecular
diagnosis. During the outbreak, farmers had observed symptoms similar to those caused
by RHDV. The fatalities affected exotic breeds of rabbits of all ages and both sexes. The
symptoms reported by farmers were anorexia (a day prior to death), lacrimal discharge,
lethargy, bleeding from the oral and nasal orifices and sudden death.
Here we report the use of microbial metagenomics sequencing to uncover the first
genomic characterization and whole genome sequencing of the RHDV2 responsible for
an outbreak of RHD in Sub-Saharan Africa.
90
91

Results
Gross Post mortem findings
The carcasses were found to be in good body condition and slightly autolyzed (5/5). The
oral and ocular mucous membranes were mildly pale (5/5). There were a few multifocal
widespread petechial haemorrhages on the ventral and dorsal abdominal subcutaneous
muscle (2/5), as well a few multifocal petechial haemorrhages on the pleura surface of
the lungs (2/5).
99

100 *RT-PCR*
101 Two samples (RT2 and RT4) out of five were positive by RT-PCR for RHDV targeting
102 the VP60 gene (Table 1). Bands were confirmed on electrophoresis at regions between
103 1500 and 2000bp (Results not shown).
104

Sequencing
We obtained one full genome (sample RT5) and two partial sequences (samples RT4 and
RT2) (Table 1). Metagenomics analysis identified the presence of Rabbit haemorrhagic
disease virus (RHDV) reads in three samples (RT2, RT4 and RT5) out of five samples
sequenced. Analysis of the sequence data revealed that all three of our sequences belong
to genotype RHDV2 also known as GI.2 or RHDVb. BLAST analysis revealed a sample
RT5 shared a 98.58% identity with a RHDV2 strain from the Netherlands, RT4 shared a
99.05% identity with a RHDV2 strain from while RT2 shared a 100% with the same
RHDV2 strain from Germany as RT4 (Figure 1). Phylogenetic analysis further confirmed
the results of our BLAST analysis, demonstrating that the sequences from these animals
belong to the RHDV2 genotype as they clustered together in the same clade with
previous RHDV2 sequences from Europe and other parts of Africa (Figure 1). Analysis
of Single nucleotide polymorphisms (SNPs) in the RHDV2 genome of sample RT5
(which was negative by RT-PCR) obtained from our study, revealed 11 unique mutations
resulting in amino acid changes. These are: Thr179Ile, Leu216Ser, Leu342Ser,
Arg827Lys, Gln928Arg, Thr1337Ile, Thr1742Ile, Leu1929Pro, Ser1978Phe,
Val2104Ala, Val2127Ala. Four of these mutations (Leu1929Pro, Ser1978Phe,
Val2104Ala, Val2127Ala) occur in the region of target for the RT-PCR primers. Sample
RT4 had a unique mutation Pro440Leu compared to other RHDV2 sequences they
clustered with on the tree.
125
126
127

128
129
130
131   Table 1: Sample Demographics
132

Sample ID	Sex	Age	RT-PCR	Genome Sequence
RT1	Female	8 weeks	Negative	No
RT2	Male	8 weeks	Positive	Yes (partial)
RT3	Male	5 weeks	Negative	No
RT4	Male	3 weeks	Positive	Yes (partial)
RT5	Male	Adult	Negative	Yes (full)

168
169
170 Figure 1: Phylogenetic relationship between the sequences from this study (coloured red)
171 and representative RHDV sequences obtained from the NCBI database
172

173
174
Discussion
Our initial suspicion of RHD based on case history, the extent of morbidity and mortality
and post mortem findings was confirmed by RT-PCR in two out of five tissue samples.
Further metagenomic sequence analysis resulted in three genome (one full and two
partial) assemblies.
To the best of our knowledge, this study represents the first full genome sequence of
Rabbit Hemorrhagic Disease Virus (RHDV) in sub-Saharan Africa. Bioinformatics
analysis of the virus genomes revealed that the three sequences from this study belong to
genotype *Lagovirus* europaeus/GI.2 also known as RDHV2 or RHDVb. This genotype
was first identified in 2010 as a novel pathogenic form of *Lagovirus* in France[8] after
which it spread rapidly through Europe and other parts of the world (Oceania, North
America and Africa)[27, 28, 30-33]. It has also been reported to replace the former circulating
GI.1 strain in Australia and Portugal[35, 36]. Phylogenetic analysis showed that our
sequences clustered closely with previous sequences from Europe. BLAST analysis
revealed that sample RT5 shared a 98.58% identity with a RHDV2 strain from the
Netherlands, RT4 shared a 99.05% identity with a RHDV2 strain from Germany while
RT2 shared a 100% with the same RHDV2 strain from Germany as RT4 suggesting that
the virus was most likely imported into the country from Europe.
The *Lagovirus* europaeus/GI.2 strain is known to dominate in most regions it has been
found and most recently in Morocco[37]. RHDV has been circulating in some African
countries and more recently, there have been reports of GI.2 outbreaks in a few countries
from North Africa (Tunisia, Egypt and Morocco)[37].
RHDV strains have yet to be characterized in Sub-Saharan African despite its devastating
effect on rabbit farms and the industry. Given the growing distribution and the
devastating effect, there is a need for heightened surveillance. In Australia, it spread to all
states and territories and rapidly became the dominant circulating strain within 18 months
of initial detection[36]. Ongoing surveillance and genomic sequencing are therefore needed
to understand the interaction of the various *Lagoviruses* in Nigeria and their impact on
host populations. Next generation sequencing of more samples is also needed in order to
determine the time of introduction of the virus into the country and how it is spreading
through the country.
Our phylogenetic analysis also revealed high genetic diversity of the GI.2 strain as shown
by the number of branching events within the clade. In addition, the high number of
unique mutations observed in the sequences from this study further emphasizes this
diversity, which is likely due in part to the rapid spread and evolution of this virus. Also,
it may explain why it affects rabbits of all ages more or less equally in our communities.
The fact that some of the mutations occurred in the region of target of the RT-PCR
primers may be the reason why some of the RT-PCR were negative (RT5), and yet
yielded a full genome after NGS and metagenomic analysis. This finding emphasizes the
need to develop primers for diagnosis based on RHDV sequences from the country of
outbreak in order to avoid false negative RT-PCR results. Compiled genomic data should
be carefully considered when developing diagnostics and vaccines.
The findings from this study is a significant landmark in the field, as it has revealed the
circulation of the RHDV2 in Nigeria and reports the first genomics characterization of

RHDV in sub-Saharan Africa. The close sequence homology suggests that the virus was
most likely imported from Europe. In addition, the high genetic diversity of the GI.2
strains found highlights the need for characterization of many more samples across sub-
Saharan Africa and in high incidence zones to guide the development of improved
diagnostics and vaccines.

The detection of RHDV2 on unbiased metagenomic sequencing, as shown in this study
illustrates the power of genomics in explaining a suspected outbreak. This ability to
rapidly identify and characterize an emerging virus (RHDV 2) – in an unusual cluster–
highlights the value of in-country genomics capacity. Serology using ELISA and RT-
PCR are the current methods of choice for RHDV diagnosis in Sub-Saharan Africa.
These diagnostic methods despite their limitations are done in very few selected
laboratories. The integration of genomics capacity into the established, but siloed,
pathogen-specific diagnostic platforms provides exciting opportunities for Veterinary
public health surveillance.

233
METHODS
Post Mortem and Sample collection
In August 2020, following reports of devastating outbreaks of suspected Rabbit
Haemorrhagic Disease (RHD) in Rabbit farms around, Ibadan, South-western region of
Nigeria, post mortem was carried out on four carcasses from farm A and one carcass
from farm B. Signalment from farm A consists of 1 female (8 weeks old) and 3 males (8
weeks, 5 weeks and 3weeks old). The breeds on the farm were Hyla. Farm B consists of
one adult male Hyla. Tissue sections were collected into RNAlater from the five (5)
rabbit carcasses suspected to have died from RHD. The samples were tagged RT1, RT2,
RT3, RT4 (farm A) and RT5 (farm B). Tissues (liver, spleen, lungs) of each animal were
pooled for RT1 - RT4 while only liver was collected for RT5. The samples were then
maintained in a cold chain and RNALater during transportation to the African Centre of
Excellence for the Genomics of Infectious Disease (ACEGID), Redeemer's University,
Ede, Nigeria for PCR confirmation and metagenomics sequencing analysis.

248
RNA Extraction and RT-PCR
Samples were first washed in PBS and thereafter homogenized and macerated. Total
RNA was extracted from tissues macerated in TRIzol using QIAamp Viral RNA
extraction kit (Qiagen, Hilden, Germany) according to manufacturer's instructions.
Extracted RNA was stored in -20^{o}C until RT-PCR and sequencing.
On the 16[th] of September, 2020, RT-PCR was performed on extracted RNA based on
modified established protocol[14] with primers targeting the VP60 gene. One-step
SuperScript™ III One-Step RT-PCR System with Platinum™ Taq DNA Polymerase
(Invitrogen, USA) was used for PCR amplification targeting the VP60 capsid protein
giving a 1740 base pair product[11]. The RT-PCR final reaction volume of 25ul was made
up of 12.5 µl of 2X reaction mix, 1.25µl of 20µM each of forward and reverse primers,
1µl SuperScript™ III RT/Platinum™ Taq Mix, 1µl RNA template and MgSo4
optimization to a final concentration of 2.5µM and nuclease free water to make up the
reaction volume. Our thermocycling condition includes; 55 °C for 30 min, 95 °C for 15
min, then 40 cycles at 95 °C for 1 min, 58 °C for 30 s and 72 °C for 1 min, and a final

extension step at 72 °C for 10 min. PCR products were viewed in 1% gel electrophoresis
(Results not shown).
266

Next Generation Sequencing and Bioinformatics Analysis
Upon RT-PCR confirmation on September 25, 2020, Nextera XT sequencing Libraries
were made based on established unbiased protocol[38]. Briefly, RNA was turbo DNase
treated to remove contaminating DNA, and converted to cDNA. Nextera XT sequencing
libraries were made from cDNA. Libraries were sequenced using the Miseq Illumina
platform in our sequencing platform at ACEGID, Redeemer's University, Ede, Osun
State, Nigeria.
Following sequencing, raw reads from the next-generation sequencing machine were
uploaded to our cloud-based platform (DNAnexus, www.dnanexus.com). Quality control
was carried out on the raw reads using fastqc
(https://www.bioinformatics.babraham.ac.uk/projects/fastqc). Metagenomics analysis
was carried out using Kraken2[39].
RHDV genomes were assembled using our publicly-available software viral-ngs v2.1.8
(https://github.com/broadinstitute/viral-ngs) implemented on DNAnexus.
Following BLASTn analysis, ninety representative RHDV sequences were obtained from
the NCBI database and aligned with our three sequences using MAFFT v7.388[40] with
further adjustment made manually as necessary in Geneious[41]. A neighbour-joining tree
was generated also in Geneious from a distance matrix corrected for nucleotide
substitutions by the Tamura-Nei model. The tree was viewed and manually edited using
FigTree (http://tree.bio.ed.ac.uk/software/figtree/). We aligned the three sequences from
this study with all sequences in the same RHDV2 clade on the tree to check for amino
acid mutations specific to our new sequences from Nigeria. We also mapped the RT-PCR
primers to our full genome obtained from this study to check if any mutation observed
occurred in the target regions of the primers.
291
292
293 Data Availability
294 All sequences from this study were submitted to the National Center for Biotechnology
295 Information (NCBI) database and the accession numbers (MW123059 - MW123061)
296 received on the 16 October, 2020.
297
298
.

References

1. Meyers, G., Wirblich, C., & Thiel, H. J. Rabbit hemorrhagic disease virus-
molecular cloning and nucleotide sequencing of a calicivirus genome. *Virology* **184**,
664–676. https://doi:10.1016/0042-6822(91)90436-F (1991).

2. Guittre, C. I., Baginski, G. & Gall, L. E. Detection of rabbit hemorrhagic disease
virus isolates and sequence comparison of the N-terminus of the capsid protein gene
by the polymerase chain reaction. *Res. Vet. Sci.* **58**, 128 - 132 (1995).

3. Abrantes, J., van der Loo, W., Le Pendu, J. & Esteves, P. J. Rabbit haemorrhagic
disease (RHD) and rabbit haemorrhagic disease virus (RHDV): a review. *Vet. Res.*
43, 12. https://doi.org/10.1186/1297-9716-43-12 (2012).

4. Mitro, S. & Krauss, H. Rabbit hemorrhagic disease: A review with special
reference to its epizootiology. *Eur. J. Epidemiol.* **9**, 70–78 (1993).

5. Asgari, S., Hardy, J. R., Sinclair, R. G. & Cooke B. D. Field evidence for
mechanical transmission of rabbit haemorrhagic disease virus (RHDV) by flies
(*Diptera Calliphoridae*) among wild rabbits in Australia. *Virus Res.* **54(**2) 123–132
(1998).

6. Ohlinger, V. F., Haas, B. & Thiel, H. J. Rabbit haemorrhagic disease (RHD):
Characterization of the causative calicivirus. *Vet. Res.* **24**(2):103–116 (1993).

7. World Organization for Animal Health (OIE) 2010. Terrestrial manual. In:Rabbit
Haemorrhagic Disease. 2010 Ch. 2.6.2. OIE Paris. France.
https://www.oie.int/fileadmin/Home/eng/Animal_Health_in_the_World/docs/pdf/Dis
ease_cards/RHD.pdf (2010).

8. Le Gall-Reculé, G. *et al.* Detection of a new variant of rabbit haemorrhagic disease
virus in France. *Vet. Rec.* **168**, 137–138 (2011)

9. Dalton, K. P. *et al.* Vaccine breaks: Outbreaks of myxomatosis on Spanish
commercial rabbit farms. *Vet. Microbiol.* **178**, 208–216 (2015).

10. Meyers, G., Wirblich, C., Thiel, H. J. & Thumfart, J. O. Rabbit hemorrhagic
disease virus: genome organization and polyprotein processing of a calicivirus studied
after transient expression of cDNA constructs. *Virology* **276**, 349–363 (2000).

11. Liu, S. J., Xue, H. P., Pu, B. Q., Qian, N. H. & Others. A new viral disease in
rabbits. *Zhongguo Xu Mu Shou Yi* **16**, 253–255 (1984).

12. Xu, W. Y. Viral haemorrhagic disease of rabbits in the People's Republic of
China: epidemiology and virus characterisation. *Rev. Sci. Tech.* **10**, 393–408 (1991).

13. Rahali, N., Sghaier, S., Kbaier, H., Zanati, A. & Bahloul, C. Genetic
characterization and phylogenetic analysis of rabbit hemorrhagic disease virus

isolated in Tunisia from 2015 to 2018. *Arch. Virol.* **164**, 2327–2332. https://doi:10.1007/s00705-019-04311-z (2019).

14. Ismail, M. M., Mohamed, M. H. A., El-Sabagh, I. M. & Al-Hammadi, M. A. Emergence of new virulent rabbit hemorrhagic disease virus strains in Saudi Arabia. *Trop. Anim. Health Prod.* **49**, 295–301. https://doi:10.1007/s11250-016-1192-5 (2017).

15. Gould, E. A. First case of rabbit haemorrhagic disease in Canada: contaminated flying insect, vs. long-term infection hypothesis: NEWS AND VIEWS: REPLY. *Mol. Ecol.* **21**, 1042–1047. https://doi:10.1111/j.1365-294X.2012.05462.x (2012).

16. Fitzner, A. & Niedbalski, W. Phylogenetic analysis of rabbit haemorrhagic disease virus (RHDV) strains isolated in Poland. *Arch. Virol.* **162**, 3197–3203. https://doi:10.1007/s00705-017-3476-0 (2017).

17. Elliott, S. & Saunders, R. Rabbit haemorrhagic disease in the UK. *Vet. Rec.* **181**, 516–516. https://doi:10.1136/vr.j5149 (2017).

18. Bergin, I. L. *et al.* Novel calicivirus identified in rabbits, Michigan, USA. *Emerg. Infect. Dis.* **15**, 1955–1962. https://doi:10.3201/eid1512.090839 (2009).

19. Abrantes, J. *et al.* New variant of rabbit hemorrhagic disease virus, Portugal, 2012-2013. *Emerg. Infect. Dis.* **19**, 1900–1902. https://doi:10.3201/eid1911.130908 (2013).

20. Morisse, J. P., Le Gall, G. & Boilletot, E. Hepatitis of viral origin in Leporidae: introduction and aetiological hypotheses. *Rev. Sci. Tech.* **10**, 269–310 (1991).

21. Le Gall-Reculé, G. *et al.* Phylogenetic analysis of rabbit haemorrhagic disease virus in France between 1993 and 2000, and the characterisation of RHDV antigenic variants. *Arch. Virol.* **148**, 65–81 (2003).

22. Dalton, K. P. *et al.* Variant rabbit hemorrhagic disease virus in young rabbits, Spain. *Emerg. Infect. Dis.* **18**, 2009–2012 (2012).

23. Carvalho, C. L. *et al.* Emergence of rabbit haemorrhagic disease virus 2 in the archipelago of Madeira, Portugal (2016-2017). *Virus Genes* **53**, 922–926. https://doi:10.1007/s11262-017-1483-6 (2017).

24. Baily, J. L., Dagleish, M. P., Graham, M., Maley, M. & Rocchi, M. S. RHDV variant 2 presence detected in Scotland. *Vet. Rec.* **174**, 411. doi: 10.1136/vr.g2781 (2014).

25. Westcott, D. G. & Choudhury, B. Rabbit haemorrhagic disease virus 2-like
variant in Great Britain. *Vet. Rec.* **176**, 74. https://doi:10.1136/vr.102830 (2015).

26. Hall, R. N. *et al.* Emerging Rabbit Hemorrhagic Disease Virus 2 (RHDVb),
Australia. *Emerg. Infect. Dis.* **21**, 2276–2278 (2015).

27. Martin-Alonso, A., Martin-Carrillo, N., Garcia-Livia, K., Valladares, B. &
Foronda, P. Emerging rabbit haemorrhagic disease virus 2 (RHDV2) at the gates of
the African continent. *Infect. Genet. Evol.* **44**, 46–50.
https://doi:10.1016/j.meegid.2016.06.034 (2016).

28. Dalton, K. P., Nicieza, I., Abrantes, J., Esteves, P. J. & Parra, F. Spread of new
variant RHDV in domestic rabbits on the Iberian Peninsula. *Vet. Microbiol.* **169**, 67–
73 (2014).

29. World Organization for Animal Health (OIE) 2016 Rabbit haemorrhagic disease
CoteD'Ivoirehttps://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?pa
ge_refer=MapFullEventReport&reportid=20852 (2016).

30. World Organization for Animal Health (OIE) 2020. Rabbit Haemorrhagic
Disease.   https://www.aphis.usda.gov/publications/animal_health/fs-rhdv2.pdf
(2020).

31. World Organization for Animal Health (OIE) 2005. World Animal Health
Publication and Handistatus II (data set for 2004). Paris, France:Office International
des Epizooties. https://www.oie.int/animal-health-in-the-world/the-world-animal-
health-information-system/data-before-2005-handistatus/ (2005).

32. World Organization for Animal Health (OIE) 2009. World Animal Health
Information Database - Version: 1.4. World Animal Health Information Database.
Paris, France: World Organisation for Animal Health. http://www.oie.int (2009).

33. World Organization for Animal Health (OIE) 2015. Rabbit haemorrhagic
disease,Benin.https://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?p
age_refer=MapEventSummary&reportid=19903 (2015).

34. Erfan, A. M. & Shalaby, A. G. Genotyping of rabbit hemorrhagic disease virus
detected in diseased rabbits in Egyptian Provinces by VP60 sequencing. *Vet World*
13, 1098–1107 (2020).

35. Lopes, A. M., Correia, J., Abrantes, J., Melo, P. & Ramada, M. Is the new variant
RHDV replacing genogroup 1 in Portuguese wild rabbit populations? *Viruses* **7**, 27-
36 (2015).

36. Mahar, J. E. *et al.* Rabbit Hemorrhagic Disease Virus 2 (RHDV2; GI.2) Is
Replacing Endemic Strains of RHDV in the Australian Landscape within 18 Months
of Its Arrival. *J. Virol.* **92**, 01374-17. https://doi.org/10.1128/JVI.01374-17 (2018).

37. Lopes, A. M., Rouco, C., Esteves, P. J. & Abrantes, J. GI.1b/GI.1b/GI.2
recombinant rabbit hemorrhagic disease virus 2 (Lagovirus europaeus/GI.2) in
Morocco, Africa. *Arch. Virol.* **164**, 279–283. https://doi.org/10.1007/s00705-018-
4052-y (2019).

38. Matranga, C. B. *et al.* Unbiased Deep Sequencing of RNA Viruses from Clinical
Samples. *J. Vis. Exp.* **113**, 54117. https://doi.org/10.3791/54117 (2016).

39. Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with
Kraken 2. *Genome Biol.* **20**, 257. https://doi.org/10.1186/s13059-019-1891-0 (2019).

40. Katoh, K., Misawa, K., Kuma, K.-I. & Miyata, T. MAFFT: a novel method for
rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids*
Res. **30**, 3059–3066 (2002).

41. Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–
1649 (2012).

Acknowledgements
This work is supported by the BBSRC-OVEL project BB/R020116/1 and WELLCOME
TRUST project 216619/Z/19/Z to CTH and JLH. We are grateful to Mrs. Philomena
Eromon (African Centre of Excellence for Genomics of Infectious Diseases (ACEGID),
Redeemer's University, Ede, Osun State, Nigeria) for advice during the RT-PCR trouble-
shooting.

Authors Contribution
A.N.H.- Initiated the work, performed the necropsy examination, advised on NGS
analysis and co wrote the first draft. O.A.O.-performed the necropsy examination and
sample collection, performed RT-PCR and analysis, and co-wrote the first draft. J.U.O. -
performed the RT-PCR and NGS experiments and co-wrote the first draft. P.E.O. –
performed Bioinformatics analysis of NGS data and discussed them. A.S.O.-participated
in the necropsy examination and sample collection, revised the first draft. J. L. H.-
Reviewed the manuscript. C.T.H.-Guided the NGS experiments, genomic data analysis
and discussions, review the manuscript. All authors contributed to the final version of the
manuscript.

Competing Interest
The authors declare no competing interest.
438

439
440   **Additional Information:** No additional information available
441
442