Title: Peroxisome import stress impairs ribosome biogenesis and induces endoplasmic

reticulum genes

Authors:

Kerui Huang [1†], Jinoh Kim [1†], Hua Bai [1*]

[†] Co-first authors

Affiliations:

1 Department of Genetics, Development, and Cell Biology, Iowa State University, Ames, IA

50011, USA

*Corresponding Author:

Hua Bai

Email: hbai@iastate.edu

Abstract

Peroxisome biogenesis diseases (PBDs) are characterized by global defects in peroxisomal

function and can result in severe brain, liver, kidney, and bone malfunctions. PBDs are due to

mutations in peroxisome biogenesis factors (PEX genes) that are responsible for peroxisome

assembly and function. There is increasing evidence suggesting that peroxisome import functions

decline during aging. The transcriptome profiling of peroxisome import defects is still lacking.

To identify conserved responses, we undertook a bioinformatic transcriptomic analysis on

Drosophila oenocyte specific Pex1, Pex12 and Pex5 knockdowns. In addition, we performed

analysis on human cells with induced peroxisome import stress. We uncovered that oenocyte-

specific Pex1, Pex12 and Pex5 have distinct transcriptional profiles with each other. Using gene

set enrichment analysis (GSEA), we identified protein processing in endoplasmic reticulum

pathway, specifically ER-associated protein degradation (ERAD) pathway is enriched and

induced in all PEX knockdowns in Drosophila. Moreover, we uncovered decreased expression in

ribosome biogenesis genes in flies and human cells. Indeed, we identified a stall at the 5'-ETS

cleavage sites during the ribosome biogenesis and impaired 40S small ribosomal export in both

flies and human. Our data indicates an unexpected link between peroxisome and ribosome

biogenesis. Our results suggest that reduced ribosome biogenesis could be conserved cellular

response to reduce peroxisome import stress.

Introduction

Peroxisomes are single-membrane organelles found in almost all eukaryotic cells. Peroxisomes are highly dynamic and versatile, as their composition and activity varies between organisms, different cell types and under different environmental stresses[1]. Generally peroxisomes are responsible for metabolizing branched-chain fatty acids (BCFA) or very long chain fatty acids (VLCFAs), purine catabolism, synthesis of plasmalogens, ether-lipids and bile acids, regulation of hydrogen peroxide and other reactive oxygen species (ROS) [1, 2]. In mammals, β-oxidation occurs largely in the mitochondria. Peroxisomes are normally required for catabolism of BCFA and VLCFAs, however it can perform β-oxidation on other fatty acids when mitochondria are compromised [3]. Peroxisomes have close interaction with many other organelles. Peroxisomes can physically interact with other organelles including the endoplasmic reticulum (ER), mitochondria, lipid droplets, lysosomes and chloroplasts, to perform specific functions [4].

Peroxisomes are formed by the action of 14 peroxisome assembly factors (Peroxins). The majority of which are involved in translocation of peroxisomal enzymes into the peroxisome matrix. Others are responsible for transporting peroxisome membrane proteins[5-8]. Unlike mitochondria, the peroxisome is composed of single membrane and it could import fully folded and oligomeric proteins into the matrix [9]. Peroxisomal matrix proteins contain one of two types of intrinsic peroxisomal targeting signals (PTSs), PTS1 or PTS2, which can direct import into the organelle [9]. Most matrix proteins are targeted through the PTS1 pathway. PTS1 consists of a tripeptide (-SKL or conserved variant) at the C-terminal and it can be recognized by a soluble receptor, Pex5 [9]. Pex7 will recognize PTS2 containing proteins near the N terminus. Pex5 and Pex7 bind their respective PTSs in the cytosol, and the receptor-cargo complexes will dock to the peroxisome surface, containing Pex13 and Pex14 docking complexes. Cargo is released from

Pex5 and imported into peroxisomes. Then Pex5 is mono-ubiquitylated by Pex4 and Pex12,

enables Pex5 recycling back to the cytosol for another round of import. Pex5 can also undergo

polyubiquitylation (by the ubiquitin-conjugating and ligase enzymes, Ubc4 and Pex2

respectively), which directs Pex5 to the proteasome for degradation (see excellent review in [10]).

Mutations in these peroxin genes and peroxisome resident enzymes can lead to a variety of

disorders, known as peroxisome biogenesis disorders (PBD) with involvement of kidney, brain,

bone and liver, and death in infants [11-14].

Recently there are increasing evidence placing peroxisome, especially peroxisomal

import function, as an important regulator for aging. Several studies suggest that peroxisomal

import function declines with age [15-17]. Consistently, our translatomic study shows that the

majority of peroxisome genes are downregulated in aged fly oenocytes [18]. Our previous study

identified that peroxisome import activities in fly oenocytes can non-autonomously regulate

cardiac arrhythmia during aging [19]. Several other studies have also implicated that peroxisome is

involved in longevity[20-22]. However, it remains elusive how impaired peroxisome import activity

contributes to cellular processes such as aging.

Ribosome biogenesis is one of the regulators of aging. The reduced level of ribosomal

genes, ribosome biogenesis factors or nutrient sensing pathways (such as TOR signaling), which

stimulate ribosome production, can increase the lifespan in multiple organisms, including *C.*

elegans, D. melanogaster, yeast, mice and human [23]. Because the rate of protein translation is

proportional to the rate of ribosome biogenesis [24, 25], it was suggested that reduced ribosome

biogenesis can reduce protein synthesis thus maintaining global proteostasis [23]. The biogenesis of

the 60S and 40S ribosomal subunits follows a complicated pathway in eukaryotic cells, which

requires the assembly of four rRNAs and ~80 ribosomal proteins (RPs) [26]. Ribosome biogenesis

occurs in the nucleolus and is initiated by transcription of a large precursor rRNA (pre-rRNA)

through RNA polymerase I (Pol I), from which the mature 18S, 5.8S, and 25S rRNAs are

generated [27]. The 5S pre-rRNA is transcribed by RNA polymerase III and incorporated into

nascent pre-ribosome in later step [28]. The pre-rRNA will assemble co-transcriptionally with

numerous trans-acting factors and early binding ribosomal proteins for the small subunit,

forming a macromolecular complex, named 90S pre-ribosome or small-subunit (SSU)

processome [29, 30]. In the firs 90S assembly intermediate, the pre-rRNA undergoes site-specific

base modifications at conserved sites and cleavage reactions. These early cleavages will produce

20S pre-rRNA (precursor to 18S), which becomes part of the early pre-40S particle. The pre-40S

particle will follow a simple maturation route to the cytoplasm where mature 40S is produced.

Ribosome biogenesis is tightly controlled by nutrient availability and extracellular conditions.

Myc and TOR pathways are known regulators of ribosome biogenesis [31, 32], however it remains

elusive whether there are additional regulators of this complex biological process.

Using human HEK293 and HepG2 cell lines, in combination of *Drosophila* tissue

specific RNAi, we sought to characterize conserved cellular responses under peroxisome import

stress. Our study discovered distinct transcriptional responses from knocking down different

components involved in peroxisome import pathways, suggesting PEXs proteins perform

additional activities besides facilitating Pex5. Genes participate in protein process and in

endoplasmic-reticulum-associated protein degradation (ERAD) are highly induced across all

PEXs knockdowns in flies, suggesting an induced ER stress under peroxisome defects. We also

identified repression of oxidative phosphorylation genes and induced inflammation pathways

across fly and humans, suggesting close interplay of peroxisome and mitochondria. Finally, for

the first time our study discovered a conserved reduction of ribosome biogenesis genes and

reduced ribosome biogenesis activity, under peroxisome import stress across fly and human.

Together, these findings shed lights on mechanisms of PBD pathology and aging.

Results

Transcriptomic analysis for peroxisomal stress response in Drosophila oenocytes and

human cells

Despite the emerging importance of peroxisome in the regulation of aging and

metabolism, little is known how cells mount defensive response to maintain cellular homeostasis.

Oenocytes are hepatocyte-like cells that are highly enriched with peroxisomes [19, 33]. In addition,

the oenocyte peroxisome was identified as an important regulator for age-related production of

inflammatory factor, unpaired 3 (upd3) [19], which could dampen cardiac function non-

autonomously. To study cellular responses induced by peroxisomal stress, we performed

transcriptomic analysis from *Drosophila* oenocytes and human cell cultures. We target on

peroxisome import stress by knocking down key factors (peroxins) involved in peroxisomal

import process (Pex5, Pex12 and Pex1) specifically in oenocytes, which were shown to

efficiently induce the ROS level and upd3 production [19]. Pex5 is one of the major peroxisome

protein import receptors, which can bind cargo proteins containing peroxisomal targeting signal

type 1 (PTS1) and deliver them to peroxisomal matrix through Pex13/Pex14 docking complex.

After releasing its cargo in peroxisomal matrix, Pex5 will be poly- or mono-ubiquitinated

through complex Pex2/Pex10/Pex12. Monoubiquitylated receptor is then released from the

membrane by the ATPases of the ATPases associated with diverse cellular activities (AAA)

family, Pex1 and Pex6 [34] (Fig. 1A). In the cytosol, the ubiquitin moiety is removed and Pex5

becomes available for another round of import. The polyubiquitylated Pex5 receptor is released

from peroxisome following the same way as the monoubiquitylated receptor, except that it is

directed for proteasomal degradation [34]. Knocking down these genes have been shown to greatly

impair peroxisome import in oenocytes and S2 cells [19, 35]. We utilized oenocyte-specific

GeneSwitch driver (*PromE^GS-Gal4*) [19] to transiently knock down peroxin genes in adult stage.

RU486 (mifepristone, or RU) was used to activate *PromE^GS-Gal4* (+RU), whereas control

genotype is the same, but with no RU feeding (-RU). After 5 days of RU activation, fly

oenocytes are dissected and polyA-tailed RNA was isolated for downstream RNA sequencing

and data analysis (Fig. 1A). To eliminate the transcriptional changes induced by RU feeding, we

included a control group (*PromE-GS-gal4>yw)* for RNA-seq analysis.

147   To identify conserved pathways involved in peroxisomal stress responses, we have

recently developed an inducible peroxisomal stress system in the mammalian cell culture. In this

system, we knocked in a Tet-ON 3G tetracycline-inducible expression construct into the AAVS1

safe harbor locus in human embryonic kidney-derived HEK293 cells (Fig. 2B). The expression

construct contained a FLAG-tagged full-length human PEX5 with a single amino acid

substitution at position 11 (cysteine to alanine, C11A), a conserved ubiquitination site involved

in PEX5 recycling [36]. Stable expression of PEX5^{C11A} mutant exerts dominant-negative effect on

wild type PEX5 recycling and efficiently blocks peroxisomal import in mammalian cell culture

[36]. PEX5^{C11A} protein level can be robustly induced by treating cells with doxycycline (Dox) at the

concentration of 1 μg/ml, but not in wild type (Fig. 2C). Induction of PEX5^{C11A} mutant was able

to block GFP-PTS1 import into peroxisomes (marked by PEX14), without significantly affecting

peroxisomal number (Fig. 1D-1F). To conduct RNA-seq analysis, PEX5^{C11A} cells were treated

with or without doxycycline for 72 hours prior for total RNA extraction. To directly compare

similarity with Pex5 knockdown in *Drosophila* oenocytes, we transfected PEX5-targeting siRNA

or non-targeting siRNA into HepG2 cells for 72 hours prior to total RNA isolation. PolyA-tailed

RNA was then isolated and RNA-seq libraries were constructed and pooled for the following

analysis (Fig. 1G-1H).

Distinct transcriptional profiling in Pex1, Pex12 and Pex5 knockdown in oenocytes

To understand how peroxisomal stress induced by different peroxins' knockdown affect

transcriptome, we conducted hierarchical cluster analysis across Pex1^{RNAi}, Pex12^{RNAi}, Pex5^{RNAi}

and yw control samples, using log2 fold change value (RU treatment / no RU treatment) (Fig.

2A). Intriguingly, we found three peroxin knockdowns produced distinct transcriptomic changes

with very little similarities, as reflected on the heatmap (Fig. 2A) and PCA analysis

(Supplementary Fig. 1A). The transcriptomic profile also varies between PEX5^{C11A} and PEX5-

siRNA in human cell culture (Supplementary Fig.1C). The little similarity between each gene

knockdowns is possibly due to different methods of inducing peroxisome stress and different cell

types used. Cluster 2 in Fig.2A represents genes that are highly induced by both Pex5 and Pex12

and contains GO terms such as response to ER stress, carbohydrate metabolic process, glycogen

biosynthetic process and lipid particle (Supplementary Fig 1C). Cluster 4 is enriched with genes

specifically induced by RU feeding, which are involved in oogenesis, transcription, histone

modification (Supplementary Fig.1D). These results are consistent with previous findings that

mifepristone can regulate genes involved in X-chromosome gene expression and oogenesis [37].

Cluster 9 and 11, which contains genes commonly induced by Pex5 and Pex1, are enriched with

GO terms including fatty acid beta-oxidation, mRNA splicing, protein folding, mitochondrial

translation, and peroxisome. Induction of peroxisomal genes suggests there is retrograde

signaling between peroxisome and nucleus. In cluster 10 and 12 contains genes induced by both

Pex1 and Pex12. These clusters contain genes involved in proteasome assembly, translation,

mitochondrial electron transport and ribosome (Supplementary Fig1f – 1E).

There are 248 differentially expressed genes (DEGs) in Pex12 knock down from

oenocytes ($|$ fold change $| \geq 1.2$, FDR ≤ 0.1), 254 DEGs induced by RU feeding. Both Pex5

knockdown and Pex1 knockdown induced great transcriptional changes: Pex1 knockdown

produced 1055 DEGs and Pex5 knockdown produced 1056 DEGs (Fig. 2B). Next, we ask what

genes are commonly regulated by all Pexs' knockdowns and what genes are specifically

controlled by individual Pexs. We compiled venn diagram using up-regulated DEGs across all

Pex-knockdown samples and yw control group. Consistent with the heatmap analysis (Fig.2A),

only 40 genes are commonly induced by all Pex knockdowns, but not in yw control group (Fig.

2C), confirming the finding that these Pex knockdowns generate different transcriptional

responses. To understand how each Pex genes specifically regulate transcriptions, we conducted

GO term analysis on genes specifically induced by individual Pex knock downs. Pex5

knockdown specifically induced 337 DEGs, which are enriched in glutathione, proteolysis, and

oxidation-reduction processes. Pex12 knockdown specifically induced genes involved in glucose

homeostasis and lipid metabolism. In contrast, Pex1 specifically induced genes involved in

protein homeostasis process, such as proteasome, ribosome, and translation (Fig. 2D).

Next, we sought to understand the function of all the commonly induced genes. When we

analyzed the function of these genes, we found 10% of them function in mitochondria (Fig. 2E),

7% of them function in ER, 5% in Golgi, whereas only 1 of them functions in peroxisome.

Localization predication was based on published datasets on peroxisome and mitochondria [38, 39],

predicated localization on Flybase and human orthologues predication. From those mitochondrial

genes, COX7A (cytochrome c oxidase subunit 7A) is a subunit for mitochondrial complex IV;

CG9512 encodes a glucose-methanol-choline oxidoreductase, its human orthologue CHDH

(choline dehydrogenase) localizes to the outer membrane of mitochondria in a potential-

dependent manner[40]. CHDH interacts with SQSTM1, a mitophagy adaptor molecule, and CHDH

is required for mitophagy [40]. This suggests mitochondrial abnormalities and possibly mitophagies

are induced under peroxisome import stress. Spidey, an oenocyte enriched gene essential for its

growth and lipid metabolism, is also induced by peroxisome stress. Its human orthologue (very-

long-chain 3-oxoacyl-CoA reductase, HSD17B12) is predicted to localize in the ER. Spidey is

reported to be a 3-ketoacyl-CoA reduction for elongation of long chain fatty acids into VLCFAs

is also induced by peroxisome stress[41] (Fig. 2E). Together, these data suggest that other

organelles' homeostasis are tightly controlled by peroxisomal and these changes can lead

collaborative response against disruption of homeostasis.

Comparing to commonly induced genes, Pex knockdowns only commonly repressed the

expression of 10 genes (Fig. 2F and Supplementary Fig. 1F). Among these 10 genes, 3 of them

encode transferases (CG1941, CG2781, CG14615), which can transfer acyl groups other than

amino acyl groups. CG2781 (ELOVL fatty acid elongase) is predicated to mediate elongation of

fatty acids into very-long-chain fatty acids. Eaat1 (excitatory amino acid transporter 1), which

encodes a transmembrane protein with glutamate-sodium symporter activity [42], is also commonly

repressed by all Pex knockdowns. Its function in oenocytes remain unknown. But glutamate

transport could indirectly regulates the synthesis of antioxidant glutathione [43], Eaat1 might also

regulate glutathione synthesis, thus regulating the redox level in oenocytes. We further analyzed

the genes specifically repressed by individual Pex knockdowns. Pex5 knockdown specifically

repressed genes involved in maturation of small subunit ribosomal ribonucleic acid (SSU-rRNA).

Pex12 knockdown specifically induced glutamine metabolic process and nucleolus genes. Pex1

knockdown repressed genes involved in mitotic cytokinesis, and defense response. Why Pex1

regulates genes in mitotic cytokinesis? It is possibly because they perform other functions in

oenocytes, such as cytoskeleton organization. Taken together, our Pex specific transcriptomic

analysis revealed common distinct changes elicited by peroxisome stress in adult oenocytes.

Peroxisomal stress induced endoplasmic reticulum changes in oenocytes

To further characterize peroxisome-stress induced signaling pathways, we performed

Gene Set Enrichment Analysis (GSEA) on three Pex knockdown samples, using a collection of

pre-defined gene sets retrieved from Kyoto Encyclopedia of Genes and Genomes (KEGG)

database on *Drosophila Melanogaster*. Interestingly, Pex12 knockdown affected many pathways

involved in RNA metabolism and processing, including RNA polymerase, RNA transport, RNA

degradation and spliceosome (Table 1). Spliceosome can carry intron removal to form the mature

mRNA. Spliceosome is a massive complex that includes hundreds of proteins and small nuclear

ribonucleoproteins (snRNPs), which are crucial in defining intron borders and facilitating

alternative splicing [44]. Many RNA binding proteins act as splicing regulators to facilitate or

inhibit splice site recognition by spliceosome[45]. Alternative splicing defects can arise when the

levels of spliceosome components are altered. Our GSEA analysis identified that small

ribonucleoprotein particle protein B (SmB), which is orthologous to human SNRPN that can

modulates functions of the splicing machinery and lead to alternative splicing [46]. In addition,

snRNP components pre-mRNA processing factor 8 (Prp8), U2 small nuclear riboprotein

auxiliary factor 38 (U2af38) and splicing factor 3b subunit 1 (sf3b1) are also repressed under

Pex12^{RNAi}. Alterations of these genes' orthologues in human are associated with broad changes

in the production of splice variant and diseases [47]. It has been reported that splicing changes can

regulate cellular responses under stressed condition such as aging [44]. Because majority of RNA

processing pathways are downregulated, it is possible that Pex12^{RNAi} induces cellular responses

primarily through regulation of alternative splicing.

GSEA results showed all three Pex knock down induced genes involved in DNA

replication, which might be attributable to disrupted reactive oxygen species (ROS) level,

because ROS are well-known mediators of DNA and mitochondrial DNA damage[48, 49]. We also

observed a decrease of ribosomal subunits, including mitochondrial ribosomes, in Pex5

knockdown. Consistent with this, down-regulation of Pex5 and Pex12 had a significant reduction

on ribosome biogenesis (FDR < 0.0001, Table 1), suggesting dampened ribosome biogenesis. In

addition, proteasome pathway has been also induced in Pex1 and Pex12 knockdowns (Table 1),

which might be possibly because of increasing needs to degrade poly-ubiquitinated Pex5 [50, 51].

Among all three different Pex knockdowns, we found protein processing in endoplasmic

reticulum pathway is up-regulated (Table 1). Density plot showed that comparing to total gene

expression in respective RNAi, Pex5[RNAi], Pex1[RNAi] and Pex12[RNAi] all showed a consistent, albeit

modest increase of the expression level involved in the ER pathway (Fig. 4A-4C). After

examining the pathway closely, we identified nine genes that are commonly induced by all Pex

RNAi (Fig. 4D). Some of these genes were not identified in previously venn analysis possibly

because they did not reach the cut-off (FC ≥ 1.2, FDR ≤ 0.1). Interestingly, CG30156, CG3061,

Ubc7, p47, Hsp70Ba and Hsp70Bb (highlighted in red) all participate in ER-associated

degradation (ERAD). Among them, p47, Hsp70Ba and Hsp70Bb are highly induced in all Pex

knockdowns, by which Hsp70Bb is induced by more than 6-fold in Pex1[RNAi]. p47 is predicted to

have ubiquitin binding activity and its human orthologue can interact with ubiquitinated

substrates during ERAD and are elevated upon ER stress [52]. Similarly, several ubiquitin ligase

complex components are also up regulated, such as sip3 (belongs to the HRD1 family) and Der-1.

The evidence suggests that there is higher level of ER stress which induced ERAD activity under

peroxisomal stress. Further investigations are needed to verify this possibility.

Peroxisome dysfunction represses oxidative phosphorylation and induced inflammation, metabolic pathways in human cells

To understand how peroxisome stress changes transcriptions in human cell cultures, we performed GO term analysis on DEGs ($|$fold change$| \geq 1.2$, FDR ≤ 0.1) from PEX5^{C11A} and PEX5-siRNA treated cells. PEX5-siRNA treatment produced 587 DEGs, whereas PEX5^{C11A} elicited dramatic transcriptional response, with 7393 DEGs. Interestingly, these two treatments share few common transcriptional changes: 26 genes commonly induced versus only 6 genes commonly repressed by two treatments (Fig. 3A). Among these 26 commonly up-regulated genes (Supplementary Fig.1G), majority of them regulates cell proliferation (CCNA2, FGFR3, CDCA5, PSMB9 and UBA7) and mitochondrial metabolism (PCK1, ACSM3, ALDH8A1 and BTG2). Then we looked at the specific genes induced by PEX5^{C11A} and PEX5-siRNA. PEX5^{C11A} specifically induced immune response, RNA metabolism, nitrogen metabolism (Fig. 3B), whereas PEX5-siRNA specifically induced peroxisome genes, cholesterol homeostasis, steroid metabolism, carboxylic acid metabolism and lipid metabolism (Fig. 3C). PEX5^{C11A} represses mitochondrial genes, translation, ribosome biogenesis. PEX5-siRNA represses negative regulation of protein phosphorylation and MAPK cascade, indicating an induction of MAPK signaling under PEX5 knockdown, which is consistent with previous findings in oenocytes [19].

To further characterize peroxisome-stress induced signaling pathways, we performed GSEA, using a collection of pre-defined gene sets retrieved from KEGG database on *Homo Sapiens*. PEX5^{C11A} significantly regulated 103 signaling pathways, whereas only 2 pathways are significant under PEX5-siRNA (FDR ≤ 0.05). Through GSEA analysis, we found oxidative phosphorylation pathway and valine leucine and isoleucine degradation pathway are downregulated under PEX5^{C11A} treatment (Table 2 and 3G). In oxidative phosphorylation

pathway, we found that key components of Complex I-IV of the respiratory chain are down-

regulated. For example, NADH dehydrogenase: NDUFV1, NDUFV2, NDUFS2, NDUFS3,

NDUFS4, NDUFS6, NDUFS7, NDUFS8 and NDUFSA11, which are components of Complex I

and deficiency of which is the most common cause of mitochondrial disease[53]. Cytochrome C1

(CYC1) which is part of the metal centers responsible for electron transfer in complex III, is also

downregulated. Almost all cytochrome c oxidase (COX, complex IV) subunits are repressed

(COX4l1, COX4l2, COX5A, COX5B, COX6A1, COX6B1, COX6C, COX7A1, COX7A2,

COX7A2L, COX7B, COX7C, COX8A and COX15) and many ATPase subunits (ATP6V1G2,

ATP6V0C, ATP6V1G1 etc.) are repressed as well. These results suggest that mitochondrial

electron transport activity is greatly impaired under peroxisomal stress, which is consistent with

previous publications[54-56]. Even though we did not observe decreased oxidative phosphorylation

genes in *Drosophila* oenocytes, we observed an altered mitochondria morphology with higher

fusion activity under Pex5 knockdown in oenocytes (unpublished observation). This indicates

peroxisome and mitochondria are dynamically connected organelles and the response to

peroxisome stress is conserved between flies and human.

It has been well established that defects in electron transport chain leads to elevated

reactive oxygen species (ROS) and can induce inflammatory pathways[57 58]. Interestingly, we also

observed activated MAPK signaling, type II diabetes mellitus under PEX5[C11A] treatment (Table

1). Pex5 knockdown was previously identified to activate MAPK/JNK signaling in *Drosophila*

41 Hawthorne St #1 oenocytes [19], this suggests a conserved cellular response from peroxisome

stress between flies and human. In addition, we also observed an upregulation of HIPPO

pathway, including key regulators such as MOB1B, LATS1/2, YAP1. HIPPO pathway which is

originally discovered for its regulation on organ size. However, recent evidence has discovered

that Hippo signaling is also involved in oncogenesis [59, 60] as well as apoptosis [61-67]. Peroxins

mutants exhibit enhanced cell apoptosis and cell death [68] (based on unpublished data) , our

evidence suggest that peroxisome stress induced by PEX5^{C11A} activates apoptosis through

HIPPO signaling activation.

PEX5-siRNA treatment produced less significant GSEA pathways comparing to

PEX5^{C11A}, with only spliceosome showing significant negative correlation with age (NES = 1.85,

FDR = 0.03). Spliceosome and RNA processing are also negatively correlated with age in

human[69]. which suggests that peroxisome defects are one of the causes for tissue aging.

According to GSEA analysis, under PEX5-siRNA induced genes involved in cholesterol

metabolism pathway (NES = -1.45, P-value = 0.019) (Fig. 3F). Among the upregulated genes in

the pathway, apolipoprotein genes are highly induced, including APOC1, APPOA1, APOA5 and

APOC3 (Fig. 3M). APOC3 had a significant 5-fold increase after PEX5 knockdown (Fig. 3M).

Apolipoproteins are crucial for lipoprotein metabolism. Not only can guide lipoprotein formation,

they also serve as activator or inhibitors of enzymes involved in lipoprotein metabolism [70].

Notably, single-nucleotide polymorphisms within the apoliprotein (APO) gene cluster, which

includes APOA1/C3/A4/5 genes, are strong risk alleles associated with hypertriglyceridemia and

increased coronary heart disease (CHD) risk in humans [71, 72]. APOA5 encodes a protein which is

secreted into plasma to control plasma triglyceride (TG) metabolism by acting as an activator of

lipoprotein lipase, thus promoting TG catabolism[73]. In contrast, APOC3 encodes a protein that

acts as an inhibitor of lipoprotein lipase and high expression of APOC3 have been associated

with increased plasma level of TG and hypertriglyceridemia [74]. Interestingly, APOC3 protein is

also a modulator for pro-inflammatory pathways, such as PKCβ and NF-κB in endothelial cells.

It is interesting that both APOC3 and APOA5, which have opposing effects on TG, all are

induced under PEX5-siRNA. Pex5 perturbation's effects on plasma TG require further

investigations. Many of pathway regulations are cell type or treatment specific. Despite of these

differences, we were able to identify several pathways, including MAPK, mitochondrial

oxidative phosphorylation, potentially PERK-mediated ER stress as conserved response to

peroxisome dysfunction. It is yet unknown whether there are additional conserved cellular

responses.

GSEA uncovers ribosome biogenesis declines as a conserved response in the peroxisomal

defects

To identify conserved response to peroxisomal stress, we compared the GSEA results in

Drosophila oenocytes and in human cells. GSEA showed a conserved downregulation of genes

functioning in the KEGG pathway ribosome biogenesis in Pex5^{RNAi}, Pex12^{RNAi} (*Drosophila*

oenocytes) and PEX5^{siRNA} in human cells (Fig. 5A and Supplementary Fig. 2A). As shown by

density plots (Fig. 5A), most of genes in the ribosome biogenesis gene set showed a consistent,

although modest, decrease in the expression level between knockdown and control, compared to

total gene expression (Fig. 4A-B and Supplementary Fig. 2B-C). A similar trend was observed

with ribosome gene set (Fig. 5B), with a decrease in ribosome genes in PEX5^{C11A} and Pex5^{RNAi},

including majority of genes encoding mitochondrial ribosomal proteins and ribosomal proteins.

Even though RU feeding also induced the ribosome pathway, however the heatmap showed RU

and Pex5^{RNAi} showed few similarities on the expression level (Supplementary Fig. 2B).

A closer look on ribosome biogenesis pathway revealed that multiple steps regulating the

process are repressed. Peroxisomal stress from three different knockdowns all target on 60S

ribosome processing and 90S pre-ribosome components (Fig. 5C). The eukaryotes contain two

ribosome subunits, the 40S (small) and the 60S (large), which contain four rRNA species (18S,

5.8S, 25S and 5S) and ribosomal proteins (RPs). Ribosome assembly starts in the nucleolus by

transcribing a large precursor rRNA (47S pre-rRNA in *Drosophila* and mammals) by RNA

polymerase I, from which the mature 18S, 5.8S, and 25S rRNAs are generated [27]. Our results

show multiple steps during ribosome processing are affected under Pex knockdown. For example,

NOP58, a core component of the box C/D small nucleolar ribonucleoprotein complex, which

controls specific pre-ribosomal RNA processing, is downregulated in PEX5[RNAi] in human cells.

Its fly orthologue, nop5, is also repressed under Pex12[RNAi] in oenocytes. Ribosome biogenesis

protein BMS1 homolog (BMS1), which is crucial for 40S maturation are repressed in human

PEX5[RNAi], same with its orthologue is flies. Additionally, elongation factor like GTPase 1 (EFL1)

can assist the release of eukaryotic translation initiation factor 6 (eIF6) release from 60S-

ribosome [75, 76]. Thus, it plays an important role in translational activation. CG33158 is the fly

orthologue of EFL1. Interestingly, we found reduced expression level of CG33158 in Pex5[RNAi]

and Pex12[RNAi] in oenocytes, suggesting the translation activity is dampened under peroxisome

stress.

During ribosome biogenesis, the 47S pre-rRNA assembles co-transcriptionally with

various *trans*-acting factors and early binding ribosomal proteins of the small subunit. Together a

huge macromolecular complex, named 90S pre-ribosome or small-subunit (SSU) processome

will form[26, 29, 30]. 90S pre-ribosome is composed of 60-70 non-ribosomal factors, with most being

called U three proteins (Utp) [26], and they can form structurally autonomous subcomplexes (UTP-

A, UTP-B, UTP-C, Mpp10-Imp3-Imp4, U3 snoRNP, and Bms1-Rcl1 modules). These modules

will associate with pre-rRNA in a sequential and hierarchical order [77-79]. UTP-A and UTP-B are

the earliest 90S modules that assemble on the 5'-external transcribed spacer (ETS) of the pre-

rRNA, whereas UTP-C and other complexes bind later. It has been reported that UTP-A complex

is critical for the initial cleavage of 5'ETS site on 47S pre-rRNA, thus regulating the ribosome

biogenesis [80]. We found multiple genes in SSU processome are downregulated under peroxisome

stress. UTP15 small subunit processome component (UTP15), nucleolar protein 6 (NOL6),

PWP2 small subunit processome (PWP2), are all repressed in PEX5^{RNAi} in human cells or their

fly orthologues are repressed in oenocytes. Because UTP-A, UTP-B and UTP-C components are

largely dampened under peroxisomal stress, we sought to verify whether peroxisomal stress also

caused pre-rRNA processing, especially at the 5'-ETS removal steps.

pre-rRNA processing starts with the removal of the 5' and 3'-ETS, and the internal

transcribed spacers 1 and 2 (ITS1 and ITS2, respectively) (Fig. 6A). To test whether pre-rRNA

processing is impaired under peroxisomal stress, we designed primers against the 5'-ETS region

and 3'-ETS regions in Pex5^{RNAi} oenocytes and PEX5^{C11A} induced human cells (Fig 6A). We

discovered strong accumulation of higher level of 5'-ETS and ITS1 could indicate a defect in

rRNA processing, or induced level of rRNA transcription (Fig. 6B-6C). Next, we measured

uncleaved ETS using primer 1 and 2 as shown in figure 6A, normalized to 18S (primer 3 and 4)

which represents total level of rRNA transcribed [81]. Higher relative expression value is indicative

of accumulated unprocessed rRNA. Pex5 knockdown specifically in oenocytes can increase the

fraction of 5'ETS-18S unprocessed rRNA (Fig. 6D), without significantly affecting the total

rRNA level (Fig.6E). We also found a consistent increase of unprocessed rRNA in human cells

after PEX5^{C11A} induction, but not 18S level (Fig. 6G-6H). To further verify whether ribosome

synthesis is impaired, we measured the localization of ribosomal protein S6 (Rps6) under

peroxisomal stress. The rationale for this assay is based on the observation that defects if

biogenesis pathway can cause ribosomal subunit export defects [82, 83]. Rps6 is an early assembling

40S subunit ribosomal protein [84-86], allowing us to score early defects in 40S synthesis. Indeed,

we found an accumulation of RPS6 in the nucleus in Pex5^{RNAi} oenocytes, after normalized to

total intensity (Fig. 6I – 6J). We also identified higher percentage of cells contain nucleus rpS6

signal in PEX5^{C11A} expressing human cells, comparing to control (Fig 6K-6L). These results

indicate a defect in rRNA processing, instead of increased rRNA transcription, under peroxisome

disruption.

Discussion

The peroxisome is the important metabolic site to perform β-oxidation of fatty acids and

the degradation of toxic hydrogen peroxide, yet it has received little respect until now. Although

peroxisomal dysfunction has been linked to severe diseases in man and aging [15, 17, 19, 87], whether

cells can mount defensive mechanism and how cells respond to peroxisome stress remain to be

determined. Here we characterized a variety of transcriptional alterations under multiple

peroxisomal stress in both *Drosophila* oenocytes and human cell cultures. We show that defects

in peroxisome import process can elicits distinct transcriptional response in *Drosophila*

oenocytes. Peroxisomes are highly interconnected with other organelles, such as ER and

mitochondria, so that peroxisome dysfunction can induce gene expression targeting to other

organelles. Specifically, peroxisome defects in oenocytes up-regulates genes involved in ER and

ERAD. PEX5^{C11A} induction induced genes involved in oxidative phosphorylation and altered

mitochondrial ribosome genes' expression. PEX5^{RNAi} specifically up-regulated genes involved in

cholesterol metabolism and transport. We also identified conserved decrease in ribosome

biogenesis and ribosome between *Drosophila* and human cells, highlighting it as a direct and

crucial mechanism to peroxisome stress.

Different importing peroxins control distinct transcriptional processes

Maintaining peroxisomal activity is a highly dynamic and complicated process,

facilitated by peroxisomal biogenesis factors (peroxins). In fact, most of the peroxins identified

so far are known to be directly involved in different stages of peroxisomal matrix protein import

[88]. Newly synthesized and folded peroxisome proteins from cytosol are targeted to the organelle

in a post-translational manner[88]. The protein import process into peroxisome is involved with

multiple steps, including cargo recognition, docking of the cargo/receptor- complex at the

peroxisome membrane, cargo translocation, cargo release into the peroxisomal matrix and

receptor recycling (Fig. 1A). Disrupting key regulators during these processes can compromise

the peroxisome import activity and lead to severe physiological consequences[19, 35], including

development of PBDs. Even though these Peroxin genes work cooperatively to maintain

peroxisome activity, it remains unknown whether their sole activity is to regulate peroxisome

biogenesis.

Interestingly, our transcriptomic analysis on multiple import-regulatory genes (*Pex5,*

Pex12 and *Pex1*) suggests that they may regulate other cellular processes. First, we found that

Pex1 and *Pex5* RNAi produced high level of transcriptional changes comparing to *Pex12* (1055

and 1056 DEGs versus 248 DEGs). This might be because the severity of peroxisome import is

different, because *Pex12* dsRNA in S2 human cells show less diffusion of cytosolic GFP-PTS

comparing to *Pex1* and *Pex5* [35]. However, both our GO ontology and GSEA analysis suggest that

even *Pex1* and *Pex5* knockdown induced different genetic pathways. *Pex5* knock down induced

GO terms related to oxidation-reduction and glutathione pathways, whereas *Pex1* dramatically

induced genes involved in translation, ribosome, and proteasome (Table 1). The discrepancies in

their transcriptomic profiles, despite sharing the same phenotype in peroxisome import, suggest

that they regulate different processes other than Pex5 recycling. Pex1 and Pex6 both belong to

the AAA ATPase family, which is known to be involved in protein quality control, including

folding, assembly, transport, and protein degradation [89]. This could explain the induction of

translation and proteasome in *Pex1*^{RNAi}. Pex1, Pex15 and Pex6 will form an exportomer and

deficiency of these components can trigger pexophagy in mammalian cells [90]. Another study

found that a missense mutation of Pex1 can ameliorated the defects of Pex6 mutant without

restoring PEX5 recycling, possibly through prevention of pexophagy, suggesting other functions

Pex1-Pex6 complex [91]. In support of this view, studies have unexpectedly identified *PEX6*

regulating mitochondrial inheritance in daughter cells in yeast. Seo et al. showed that PEX6

overexpression was able to rescue the deficit in Atp2p (the β-subunit of mitochondrial F_1, F_0-

ATPase [92]) mutant by improving its import into mitochondria. Studies suggest that Pex1 was

able to translocate between mitochondria and peroxisomes[93], where they showed that

mitochondria localized PEX6-PEX1-PEX26 complex can rescue PEX26 deficient cells.

Altogether, our findings provide new insights in peroxisome biogenesis disorders, peroxins

functions and their significance of other organelles' activities.

Peroxisome and inter-organelle communications

To optimize their activities, peroxisomes must coordinate with other organelles

frequently. No peroxisome is in isolation. A growing body of evidence showed that peroxisome

share physical membrane contact sites with other organelles (see an excellent review here [4]).

This suggests there is intimate interaction between peroxisomes with these organelles, which is

important for controlling processes such as metabolism and organelle proliferation. In agreement

of this concept, we identified changes of gene expression for other organelles, especially for ER

and mitochondria, under various Pex knockdowns.

Our results of PEX5^{RNAi} in HepG show significant up-regulation of cholesterol

metabolism (Fig. 3C and 3F). Previous study shows that cholesterol accumulates drastically in

animals and human patients with peroxisomal disorders, as a result of disrupted peroxisome-

lysosome contact sites [94]. The majority of cellular cholesterol (60% - 80%) is localized at the

plasma membrane (PM) [95]. Cholesterol is synthesized in the ER or it can be imported

exogenously and stored in the lysosome. Cholesterol can be transported from the ER and

lysosome to the PM. Recently it was shown that peroxisome mediates the transfer of cholesterol

from lysosome to PM. A peroxisome-lysosome contact site was shown to be required for this

transport [94]. An up-regulation of cholesterol metabolism genes could be a compensatory

mechanism to degrade excess cholesterol accumulated inside the cells.

Besides lysosome, peroxisomes also actively interact with ER and mitochondria. Our

results show that mitochondrial ribosomal subunits are repressed under Pex5 knockdown in

oenocytes (Table 1), as well as in PEX5^{C11A} mutant (Table 2). Consistently, genes involved in

oxidative phosphorylation are largely repressed in PEX5^{C11A}, which was also been observed in

Pex16 KO [96]. Our results suggest that mitochondria function is altered under peroxisome stress.

In fact, previous studies as well as our observation have found that mitochondria underwent

functional and morphological changes under Pex5 and Pex16 deficiencies [96, 97]. In these studies,

they consistently observed a more fused mitochondria structure, reduced level of the respiratory

chain activities. Park et. al identified that peroxisome-derived phospholipids are responsible

peroxisome dysfunction induced mitochondrial abnormalities [96]. Park et. al inhibited

plasmalogen synthesis by knocking down expression of GNPAT, an enzyme responsible for the

first step in plasmalogen production. GNPAT knockdown was able to recapitulate phenotypes

with Pex16 knockdown, including altered mitochondrial morphology and decreased mtDNA.

Interestingly, they also found dietary supplementation of plasmalogen precursors can rescue

thermogenesis in Pex16-KO mice. However, whether plasmalogen level is the sole contributor to

mitochondrial dysfunction and why there are tissue specific mitochondria abnormalities under

peroxisome deficiency, these questions still require investigation.

510   Our results suggest different Pex knockdowns commonly induce genes VLCFA

synthesis, whose protein products can localize on ER (e.g. *spidey* and *Cyp6g1*). Biosynthesis of

VLCFA takes place at the ER and can be metabolized in peroxisomes [98, 99]. Our findings suggest

two organelles work closely to regulate metabolism of VLCFA. EM images showed that two

organelles are adjacent to each other and that ER membrane can wrap around peroxisomes [100-103].

In fact, peroxisome-ER crosstalk is important for many lipid-related metabolic pathways,

including the biosynthesis of ether-phospholipids, production of polyunsaturated fatty acids, bile

acids, isoprenoids, and cholesterol.

518   Interestingly, we found peroxisome dysfunction in oenocyte cells can trigger protein

process pathway in the ER and ERAD (including CG30156, CG30161, Ubc7, p47, Hsp70Ba and

Hsp70Bb), which is possibly due to elevated ER stress (Fig. 3). Consistently, it has been

previously identified that peroxisome-deficient *Pex2* and *Pex5* knockout mice had elevated level

of ERAD gene expression, ER stress and ER abnormalities, prominently through PERK

activation [97, 104]. How does peroxisome dysfunction lead to the ER stress? Possible reasons

include perturbed flux of mevalonate metabolites, changes in fatty acid levels or composition and

increased oxidative stress [105]. Specific induction of PERK under peroxisome stress, not

activation of Xbp1 splicing, suggesting peroxisome stress induced a specific cellular response

distinct from protein misfolding induced ER stress. Genetic experiments and biochemical assays

should be utilized to discern the nature of peroxisome induced ER stress.

Peroxisome in ribosome biogenesis and protein homeostasis

531   Our results identified decreased transcription on ribosomal proteins and ribosome

biogenesis factors, as a conserved response to peroxisome stress between humans and

Drosophila. This is also the first study to verify the reduced ribosome biogenesis under Pex5^{RNAi}

in oenocytes and PEX5^{C11A} mutant in human cells. Our results show that the 5' cleavage process

is inhibited under Pex5 knockdown. Our result indicates that peroxisome defects may

specifically regulate ribosome biogenesis by altering ribosome biogenesis process, which might

be more efficient in dealing with transient stress. In addition, we also observed an accumulation

of RPS6 in the nucleus under Pex5 stress, which is often an indicator of ribosome biogenesis

defects, specifically involved with 40S maturation process [82]. However, it is highly possible that

60S maturation is also affected, judged by down-regulated expression of several crucial genes:

Midasin (MDN1)[106], GTPBP4 [107, 108] and EFL1[75, 76]. Overall, our results indicate a down-

regulation of ribosome availabilities, and possibly lower translational activities under

peroxisome stress.

544   Why multiple peroxisome defects converge to impair ribosome biogenesis? It is possible

that nucleolus is sensitive to peroxisomal or mitochondrial ROS level, which can hamper the

function of ribosome biogenesis proteins that primarily localize in the nucleolus. However, very

little is known on what stresses can alter nucleolus protein activity or composition. Further

research is needed to investigate this area.

549   It has been well established that reduced ribosome proteins and ribosome biogenesis

factors can increase longevity in model organisms (see [23] for review). The evidence also suggest

that reduced ribosome biogenesis is a protective mechanism to confer peroxisomal stresses.

Being a regulator of ribosome biogenesis, Pex5's effect on lifespan had been controversial. In

Δpex5 yeast cells, there was a strong reduction in chronological lifespan[20]. However, post

developmental knockdown of prx-5, a *C. elegans* homolog for mouse *Pex5*, increased the

worm's lifespan [21, 22]. The paradoxically increased lifespan by peroxisome disruption could be

attributed to decreased level of ribosome and translational activities. It will be interesting to see

whether post developmental knockdown of Pex5 in oenocytes can also prolong lifespan through

the regulation on ribosome biogenesis. In addition, how peroxisome import defect reduces

ribosome biogenesis will be an interesting avenue to investigate.

Methods

Plasmid Construction

Human Pex5 cDNA from the Mammalian Gene Collection (MGC) was purchased from

Dharmacon. The hPEX5 was amplified by PCR using the forward and reverse primers (5'-

CACTATAGGGAGACCCAAGCTTATCTAGACATGGCAATGCGGGAGCT-3' and 5'-

TCTTACTTGTCATCGTCGTCCTTGTAGTCGCCCTGGGGCAGGCC-3') and introduced

between XhoI and BamHI sites in c-Flag pcDNA3 (addgene #20011) to generate Flag-tagged

hPEX5 using NEBuilder® HiFi DNA assembly Master mix (New England Biolabs). Site-

directed mutagenesis for amino acid substitution was performed using the Q5® Site-directed

mutagenesis kit (New England Biolabs) according to the manufacture's instruction. The primers

for C11A mutant were 5'-GGAGGCCGAAgctGGGGGTGCCAACC-3' and 5'-

ACCAGCTCCCGCATTGCC-3'.

To generate Tetracycline inducible PEX5C11A plasmid, we modified pMK243 (Tet-

OsTIR1-PURO) from Masato Kanemaki (Addgene #72835). pMK243 was digested by BgIII and

MluI to remove OsTIR sequence. Flag-PEX5C11A was amplified by PCR using the forward and

reverse primers (5'- gattatgatcctctagacatatgctgcagattacttgtcatcgtcgtccttgtagt-3' and 5'-

tcctaccctcgtaaagaattcgcggccgcaatggcaatgcgggagctggt-3') and introduced between BgIII and

MluI sites in digested pMK243 plasmid to generate Tet-PEX5C11A-PURO plasmid. All

plasmids are confirmed by Sanger sequencing.

Generation of CRISPR Knock-in HEK293 cells expressing PEX5C11A

HEK293 cells were maintained in Dulbecco's Modified Eagle Medium containing 10%

fetal bovine serum (FBS), with penicillin and streptomycin. To generate stable cell line, we

followed the protocol described in Natsume et al. 1×10^6 HEK293 cells were plated in one well

of a 6-well plate. After 24hrs, 800ng of AAVS1 T2 CRISPR in pX330 (Addgene #72833) and

1ug of Tet-PEX5C11A-PURO were transfected using Effectene (Qiagen) according to the

manufacturer's instructions. After 48hrs, the cells were detached and diluted at 10 to 100 times

in 10ml of medium containing 1ug/ml of puromycin and transferred to 10 cm dishes. The

selection medium was exchanged every 3 to 4 days. After 8 to 10 days, colonies were marked

using a marker pen under a microscope, picked by pipetting with 10ul of trypsin-EDTA, and

subsequently transferred to a 96-well plate. 100ul of the selection medium was added. When the

cells were confluent, they were transferred to a 24-well plate. After 2-3 days, the cells were

transferred to a 6-well plate. After 2-3 days, cells were detached and half of the cells were

frozen, and the rest were used for genomic DNA preparation.

Genomic DNA isolation and PCR

To prepare genomic DNA, cells were lysed in buffer A solution (100mM Tris-HCl

[pH7.5], 100mM EDTA, 100mM NaCl, 0.5% SDS) followed by incubation at 65°C for 30min.

Buffer B (1.43M potassium acetate, 4.28M lithium chloride) was added and incubated on ice for

10min. After centrifuge at 12,000 rpm for 15 min, the supernatant was transferred to new

microtube and added isopropanol. After isopropanol precipitation, DNA pellets were washed in

70% ethanol and resuspended with DNAse-free Water.

To verify Tet-PEX5C11A-PURO insertion into AAVS1 locus, genomic PCR was

performed using Q5® High-Fidelity DNA polymerase (New England BioLabs) according to the

manufacturer's instruction. 5'-cgtttcttaggatggccttc-3' and 5'-agaaggatggagaaagagaa-3' were used

for WT cell validation. 5'-cgtttcttaggatggccttc-3' and 5'-ccgggtaaatctccagagga-3' were used for

Tet-PEX5C11A-PURO integration.

Fly husbandry and strains

The following RNAi lines were used in the KD experiments: y[1] v[1]; P{y[+t7.7]

v[+t1.8]=TRiP.HMJ21920}attP40 (BDSC # 58064, Pex5 RNAi), y[1] sc[*] v[1] sev[21];

P{y[+t7.7] v[+t1.8]=TRiP.HMC03536}attP2/TM3, Sb[1] (BDSC # 53308), y[1] v[1]; P{y[+t7.7]

v[+t1.8]=TRiP.HM05190}attP2 (BDSC # 28979). The control line used for the KD experiments

is yw, a gift from Rochele Lab. All fly lines are crossed to oenocyte specific gene-switch driver,

yw; PromE800-GS-gal4/Cyo;+, a gift from Heinrich Jasper.

Female flies were used in all experiments. Flies were maintained at 25°C, 60% relative

humidity, and 12-h light/dark cycle. Adults and larvae were reared on a standard cornmeal and

yeast-based diet, unless otherwise noted. The standard cornmeal diet consists of the following

materials: 0.8% cornmeal, 10% sugar, and 2.5% yeast. RU486 (mifepristone, Fisher Scientific)

was dissolved in 95% ethanol, and added to standard food at a final concentration of 100□µM

for all the experiments. Gene KD or overexpression was achieved by feeding flies on RU486

food for 5-6 days, before RNA isolation.

Fly oenocyte RNA isolation

623  Adult tissues oenocytes (20 females per replicate) were all dissected in cold 1□×□PBS

before RNA extraction. For oenocyte dissection, we first removed FB through liposuction and

then detached oenocytes from the cuticle using a small glass needle. Tissue lysis, RNA

extraction was performed using RNeasy Micro kit from QIAGEN (catalog number 74034) with

the following modifications. Samples were first collected in 1.7ml centrifuge tubes. Dissected

tissues were saved in 150ul of Buffer RLT (a component of RNeasy kits) with 143mM β-

mercaptoethanol on ice during the dissection. Samples were then put at room temperature for 3

minutes prior to lysis and centrifuged at 7.5xg for 3minutes. Additional 150ul buffer RLT was

added and pellet pestle grinder (Kimble pellet pestles, catalog number 749540-0000) was used to

lyse the tissue for 1 minutes. 20ng carrier RNA was added to the cell lysates. Freezed samples

was thawed in 37°C water bath for 1 minute. Total RNA was extracted using RNeasy Plus Micro

columns following company manual.

Drosophila oenocyte RNA-seq library construction and sequencing

636  RNA-seq libraries were constructed using 100 ng of total RNA and NEBNext Ultra II

RNA Lib Prep (NEB, Ipswich, MA, USA. Catalog number: E7770L). RNA concentrations were

measured using Qubit RNA BR Assay Kit (Thermo Fisher Scientific, catalog number: 10210).

Poly(A) mRNA was isolated using NEBNext Oligo d(T)25 beads and fragmented into 200□nt in

size. Purification of the ligation products are performed using Beckman Coulter AMPURE XP

(BECKMAN COULTER, catalog number: A63880). After cDNA synthesis, each cDNA library

was ligated with a NEBNext adaptor and barcoded with an adaptor-specific index (NEBNext®

Multiplex Oligos for Illumina, NEB, catalog number: E7335S). Twelve libraries were pooled in

equal concentration and sequenced using Illumina HiSeq 3000 platform (single end, 150bp reads

format).

Human cells library preparation

HEK293 expressing Tet-PEX5C11A-Puro cells were cultured in Dulbecco's Modified

Eagle Medium containing 10% fetal bovine serum (FBS), with penicillin and streptomycin.

HepG2 cells were cultured in Minimum Essential Medium containing 10% FBS, 1mM sodium

pyruvate, 1X MEM Non-Essential Amino Acids solution (ThermoFisher, #11140050) with

penicillin and streptomycin. Cells were incubated in a 37°C incubator in an atmosphere of 5%

CO_2 in air.

To prepare RNA-seq libraries, 4x10^5 HEK293 expressing Tet-PEX5C11A-Puro cells

were seeded in a 12 well plate. After 1 day, 1ug/ml of doxycycline was added to the cells to

induce PEX5C11A in the cells. After 72hrs, total RNA was isolated from the cells with or

without doxycycline treatment. 3x10^5 HepG2 cells were seeded in a 6-well plate. Transfection

of siRNA into HepG2 cells was performed with lipofectamine RNAiMAX (ThermoFisher)

according to the manufacturer's instructions. On-TARGET plus human PEX5 siRNA

(Dharmacon # L-015788-00-0005) was used for knockdown of PEX5 in HepG2 cells. For

negative control, On-TARGET plus non-targeting siRNA (Dharmacon # D-001810-02-05) was

used. After 72hrs, total RNA was isolated from the cells. All samples were treated TURBOTM

DNase (Thermofisher #AM1907) to remove all traces of DNA. RNA-seq libraries were

constructed using 1ug DNA-free total RNA with NEBNext Ultra II RNA Library Prep Kit for

Illumina (New England Biolabs, Ipswich, MA, USA. NEB#E7770). Poly (A) mRNA was

isolated using NEBNext Poly(A) mRNA Magnetic isolation module (New England Biolabs,

Ipswich, MA, USA. NEB#E7490). After first strand and second strand and second strand cDNA

synthesis, each cDNA library was ligated with a NEBNext adaptor and barcoded with an

adaptor-specific index. Twelve libraries were pooled in equal concentrations and sequenced

using Illumina HiSeq 3000 platform.

RNA-Seq data processing and differential expression analysis

FaastQC (v0.11.8) was first performed to check the sequencing read quality. Then

sequence alignment and mapping was performed using the Spliced Transcripts Alignment to a

Reference (STAR) software (v2.7.3a) [110]. The raw reads were mapped to *D. melanogaster*

genome (BDGP Release 6) or Genome Reference Consortium Human Build 38 (GRCh38).

Reads mapped were then counted with summarizedOverlaps function using "Union" mode in R.

Counts are then analyzed in DESeq2 (v1.26.0) [111] for batch control analysis and test for

differential expression.

Principal component analysis (PCA), heatmap and expression correlation plot

PCA graph was generated using plotPCA function of R package DESeq2[111]. Heatmaps

and hierarchy clustering analysis were generated using heatmap.2 function of R package gplots.

The density plots were plotted using R package ggplot.

Gene set enrichment analysis (GSEA)

For GSEA analysis, all pre-defined set of 132 KEGG pathways in Drosophila were

downloaded from KEGG. Text were trimmed and organized using Java script. Normalized

counts were used as input for parametric analysis and organized as suggested by GSEA tutorial

site (GSEA [112, 113]). Collapse dataset to gene symbols was set to false. Permutation type was set

to gene set; number of permutations was set to 1000; enrichment statistic used as weighted

analysis; metric for ranking genes was set to Signal to Noise.

Gene ontology and pathway analysis

Functional annotation analysis of differentially expressed genes was performed using

STRING[114] or DAVID[115, 116]. GO terms (Biological Process, Molecular Function, Cellular

Component), KEGG pathway, INTERPRO Protein Domains and Features, were retrieved from

the analysis.

Quantitative real-time polymerase chain reaction (qRT-PCR)

qRT-PCR was performed using Quantstudio 3 Real-Time PCR system and SYBR green

master mixture (Thermo Fisher Scientific, USA Catalog number: A25778). All gene expression

levels were normalized to Rpl32 (in *Drosophila*), GAPDH (in humans) by the method of

comparative Ct[117]. Mean and standard errors for each gene were obtained from the averages of

three biological replicates, with two technical repeats.

Immunostaining

Anti-RpS6 (Cell Signaling, catalog number 2317) at concentration of 1:100 was used to

stain oenocytes' RpS6[118]. Anti-RpS6 at concentration of 1:50 was used to stain human cells.

Secondary antibodies were obtained from Jackson ImmunoResearch.

Adult oenocyte tissues were dissected in PBS and fixed in 4% paraformaldehyde for

15 min at room temperature. Tissues were washed with 1x PBS with 0.3% Triton X-100

(PBST) for three times (~5 min each time), and blocked in PBST with 5% normal goat serum

for 30 min. Tissues were then incubated overnight at 4 °C with primary antibodies diluted in

PBST, followed by the incubation with secondary antibodies for 1 h at room temperature. After

washes, tissues were mounted using ProLong Gold antifade reagent (Thermo Fisher Scientific)

and imaged with an FV3000 Confocal Laser Scanning Microscope (Olympus). DAPI or Hoechst

33342 was used for nuclear staining.

1.2 x 10^5 HEK293-PEX5C11A cells were seeded in 24-well plates on coverslips

(Neuvitro #GG1215PLL). After 1day, cells were treated with or without doxycycline (1ug/ml)

for 3 days. The cells were rinsed with PBS, fixed in 4% paraformaldehyde for 10 min, and rinsed

with PBS again. Cells were permeabilized in 0.5% Triton X-100 in PBS for 10 min. Cells were

treated with PBS containing 1% bovine serum albumin (BSA) for 1 hr at room temperature. The

cells were incubated with antibody against RPS6 (1:50) diluted in PBS for overnight at 4°C.

Next day, cells were washed with 0.05% Triton X-100 in PBS for three times and incubated with

Alexa Flour® 488 donkey anti-mouse IgG (1:500) and Hoechst (1:1000) diluted in PBS for 1hr

at room temperature in the dark. After that, cells were washed with 0.05% Triton X-100 in PBS

for three times. Cells were briefly washed with PBS and mounted on glass slides using mounting

medium (Thermo Fisher Scientific, #P36961). Images were visualized by confocal microscope.

Image analysis and quantification

Confocal images were quantified using Olympus CellSense Dimension software

(Olympus, v 1.16) and ImageJ (v1.49). For oenocyte RpS6 signaling quantification, five

oenocyte nucleus were randomly selected from each image. Each single nucleus was set as a

region of interest (ROI). A background ROI was also selected. The intensity of RpS6 signal was

calculated for nucleus intensity and normalized to total intensity (total intensity = nucleus

intensity + background). For human RpS6 quantification, we counted the number of cells that

that contain RpS6 signal inside the nucleus versus total number of cells in one image.

Statistical analysis

GraphPad Prism (GraphPad Software, La Jolla, CA, USA, v6.07) or Deseq2 was used for

statistical analysis. To compare the mean value of treatment groups versus that of control,

student t-test was used. Log2 fold change and FDR values were calculated by Deseq2 using

Benjamini-Hochberg method for sequencing data.

Figure legends

Figure 1 Transcriptomic analysis for peroxisomal stress response in *Drosophila* oenocytes

and human cells. A Schematic diagram showing the key genes involved in peroxisomal import,

as well as oenocyte specific RNA-seq analysis. RNAi targets genes marked in red. **B** Schematic

diagram for generation of CRISPR Knock-in HEK293 cells expressing PEX5^{C11A}. **C** Western

blot shows inducible PEX5^{C11A} construct after treatment of Doxycycline. **D** Immunostaining

showing GPF-PTS1 importing activity in wild type cells (Dox-) and PEX5^{C11A} expressing cells

(Dox+). Scale bar represents 10μm. **E** Quantification of **D** on % cells with PTS1 import activity

in Dox- versus Dox+ cells. **F** Quantification in **D** showing the number of PEX14 positive

peroxisomes in Dox- versus Dox+ cells. **G – H** Schematic diagram showing RNA-seq analysis

using human cells.

Figure 2 Pex1, Pex12 knock-down show distinct transcriptional pattern from Pex5. A

Hierarchical clustering heatmap analysis, plotting log2 fold change (+RU / -RU) in yw, Pex1

RNAi, Pex12 RNAi, Pex5 RNAi oenocyte samples. **B** Number of differentially expressed genes

(DEGs) across all oenocyte samples. **C** Venn diagram analysis showing genes commonly up

regulated by yw, Pex1 RNAi, Pex12 RNAi and Pex5 RNAi (marked by red). **D** Gene Ontology

(GO) term analysis on commonly induced genes from **C. E** Organelle localization of proteins

produced commonly induced genes. **F** Venn diagram analysis showing genes commonly down

regulated by yw, Pex1 RNAi, Pex12 RNAi, and Pex5 RNAi (marked by red). **G** Gene Ontology

(GO) term analysis on commonly induced genes from **F.**

Table 1 GSEA pathway analysis results under Pex5, Pex12 and Pex1 RNAi. NES:

normalized enrichment score. "Up or down regulated" indicates whether genes within that

pathway are induced or reduced under RNAi treatment.

Figure 3 Peroxisomal stress induced endoplasmic reticulum changes in oenocytes. A-C Density

plot showing the fold change (Pex RNAi / Control) of ER pathway genes in oenocytes. **D** Plotting

of fold change in selected genes involved in ERAD pathway across Pex1 RNAi, Pex12 RNAi, and

Pex5 RNAi samples.

Figure 4: Peroxisome dysfunction up-regulates inflammation and cholesterol pathways in

human cells. A Venn diagram analysis showing commonly up-regulated genes by PEX5^{C11A}

induction and PEX5 RNAi. **B-C** Gene ontology analysis on up-regulated genes only in PEX5^{C11A}

or only in PEX5 RNAi cells. **D** Venn diagram analysis showing commonly down-regulated genes

by PEX5^{C11A} induction and PEX5 RNAi. **E-F** Gene ontology analysis on down-regulated genes

only in PEX5^{C11A} or only in PEX5 RNAi cells. **G** List of enriched pathways from GSEA analysis

on PEX5 RNAi cells. **H-K** GSEA enrichment profiles on PEX5^{C11A}: Oxidative phosphorylation,

MAPK signaling, regulation of lipolysis in adipocytes, HIPPO signaling. **L-M** GSEA enrichment

profiles on PEX5 RNAi: spliceosome, cholesterol metabolism. **N** Selected genes from cholesterol

metabolism pathways on PEX5 RNAi cells. Y-axis represents fold change (RNAi / Control).

Figure 5 Peroxisome dysfunction represses ribosomal genes in both flies and humans: A

Density plot showing the fold change (Treatment / Control) of ribosome biogenesis pathway

genes in oenocytes and PEX5 RNAi in human cells. **B** Density plot showing the fold change

(Treatment / Control) of ribosome pathway genes in oenocytes and $PEX5^{C11A}$ in human cells. **C**

Schematic diagram showing ribosome biogenesis pathway, the role of biogenesis factors and

their fold change in oenocytes (Pex5, Pex12 RNAi) and human cells (PEX5 RNAi). Genes

reduced the expression under treatment for more than 0.7-fold are noted on the diagram (red

arrow indicates genes reduced under Pex5 RNAi in oenocytes, yellow indicates Pex12 RNAi,

blue indicates PEX5 KD in human cells).

Figure 6 Impaired ribosome biogenesis genes in human and flies: A-B Overview of *D.*

melanogaster, and Mammalian rRNA biogenesis Pathways. rRNA biogenesis intermediates are

generally conserved among vertebrates; 5.8S rRNA is cleaved into 2S and a short 5.8S, and 28S

rRNAs are cleaved into 28Sa and 28Sb. Blue arrows indicate the locations of the primers for

ETS. Green label indicates primer locations for ITS1. Red arrow indicates primers for 18S. **C-F**

The relative amounts of the ETS, ITS1, unprocessed ETS/18S ratio, 18S were measured by

qPCR using RNA isolated from control oenocytes or Pex5 RNAi oenocytes. Data are presented

as the mean ± s.e.m (n = 2). **G-H** The relative amounts of the unprocessed ETS/18S ratio, 18S

were measured by qPCR using RNA isolated from control cells (-DOX) or $PEX5^{C11A}$ (+DOX)

cells. Data are presented as the mean ± s.e.m (n = 3). **I-L** Analysis of ribosome proteins (RPs)

nucleolar localization. **I** Depletion of Pex5 in oenocytes results in the accumulation of RpS6

nucleolus. Scale bar shows 3.3μm. **J** Quantification of RpS6 intensity in nucleolus normalized to

total intensity within the cell. **K** Expressing PEX^{C11A} (+Dox) results in the accumulation of RpS6

nucleolus in HEK293 cells (n=6). Scale bar shows 20μm. **L** Quantification of number of cells

containing RpS6 signal in nucleolus normalized to total number of cells, in control (-Dox) and

PEX5^{C11A} expressing cells (+Dox) (n = 6).

Supplementary figure 1: A PCA analysis on gene expression from oenocyte samples: Pex1,

Pex12, Pex5 and yw. 0RU represents the control group, 200RU represents the RNAi group. **B**

PCA analysis on gene expression from human cell culture samples: PEX5^{C11A} (-DOX) versus

PEX5^{C11A} (+DOX); control versus PEX5^{RNAi}. **C-D** GO analysis for clusters identified in Figure

2A. **E** Oenocyte genes that are commonly repressed by all Pex knockdowns but not affected by

RU feeding. **F** Genes that are commonly induced by PEX5 RNAi and PEX5^{C11A} in human cell

cultures.

812
Supplementary figure 2: A GSEA enrichment profiles on ribosome biogenesis pathway in Pex5

RNAi, Pex12 RNAi of oenocyte samples and PEX5 knockdown in human cell. **B** Heatmap

analysis plotting log2 (fold change) value in yw samples (200 RU/0RU) and Pex5 RNAi samples

in oenocytes. **C** GESA enrichment profiles in ribosome pathway in Pex5 RNAi in oenocyte

samples, PEX5 knockdown and PEX5^{C11A} samples in human cells. **D-E** Density plot on fold

change of all genes in PEX5^{C11A} group and PEX5 knockdown group. **F** Density plot on fold

change of ER pathway genes in PEX5^{C11A} group and PEX5 knockdown group. **G-I** GSEA

enrichment profiles on protein processing in endoplasmic reticulum in Pex5 RNAi, Pex12 RNAi

and Pex1 RNAi samples in oenocytes.

References

842 1. Pridie, C., Ueda, K. & Simmonds, A.J. Rosy Beginnings: Studying Peroxisomes in
843   Drosophila. *Frontiers in Cell and Developmental Biology* **8** (2020).
844

845 2. Fujiki, Y. *et al.* Recent insights into peroxisome biogenesis and associated diseases. *J
846   Cell Sci* **133** (2020).
847

848 3. Violante, S. *et al.* Peroxisomes can oxidize medium- and long-chain fatty acids through a
849   pathway involving ABCD3 and HSD17B4. *FASEB J* **33**, 4355-4364 (2019).
850

851 4. Shai, N., Schuldiner, M. & Zalckvar, E. No peroxisome is an island - Peroxisome contact
852   sites. *Biochim Biophys Acta* **1863**, 1061-1069 (2016).
853

854 5.   Fujiki, Y., Okumoto, K., Mukai, S., Honsho, M. & Tamura, S. Peroxisome biogenesis in
855   mammalian cells. *Front Physiol* **5**, 307 (2014).
856

857 6.   Steinberg, S.J. *et al.* Peroxisome biogenesis disorders. *Biochim Biophys Acta* **1763**, 1733-
858   1748 (2006).
859

860 7.   Subramani, S. Components involved in peroxisome import, biogenesis, proliferation,
861   turnover, and movement. *Physiol Rev* **78**, 171-188 (1998).
862

863 8.   Waterham, H.R. & Cregg, J.M. Peroxisome biogenesis. *Bioessays* **19**, 57-66 (1997).
864

865 9.   Smith, M.D. & Schnell, D.J. Peroxisomal protein import. the paradigm shifts. *Cell* **105**,
866   293-296 (2001).
867

868 10. Smith, J.J. & Aitchison, J.D. Peroxisomes take shape. *Nat Rev Mol Cell Biol* **14**, 803-817
869   (2013).
870

871 11. Wanders, R.J. Metabolic and molecular basis of peroxisomal disorders: a review. *Am J
872   Med Genet A* **126A**, 355-375 (2004).
873

874 12. Wanders, R.J. & Waterham, H.R. Peroxisomal disorders I: biochemistry and genetics of
875   peroxisome biogenesis disorders. *Clin Genet* **67**, 107-133 (2005).
876

877 13. Baumgartner, M.R. & Saudubray, J.M. Peroxisomal disorders. *Semin Neonatol* **7**, 85-94
878   (2002).
879

880 14. Gould, S.J. & Valle, D. Peroxisome biogenesis disorders: genetics and cell biology.
881   *Trends Genet* **16**, 340-345 (2000).
882

883 15. Narayan, V. *et al.* Deep Proteome Analysis Identifies Age-Related Processes in C.
884   elegans. *Cell Syst* **3**, 144-159 (2016).
885

886 16. Weir, H.J. *et al.* Dietary Restriction and AMPK Increase Lifespan via Mitochondrial
887   Network and Peroxisome Remodeling. *Cell Metab* **26**, 884-896 e885 (2017).
888

889 17. Legakis, J.E. *et al.* Peroxisome senescence in human fibroblasts. *Mol Biol Cell* **13**, 4243-
890   4255 (2002).
891

892 18. Huang, K. *et al.* RiboTag translatomic profiling of Drosophila oenocytes under aging and
893   induced oxidative stress. *BMC Genomics* **20**, 50 (2019).
894

895 19. Huang, K. *et al.* Impaired peroxisomal import in Drosophila oenocytes causes cardiac
896   dysfunction by inducing upd3 as a peroxikine. *Nat Commun* **11**, 2943 (2020).
897

20.   Lefevre, S.D., van Roermund, C.W., Wanders, R.J., Veenhuis, M. & van der Klei, I.J.
The significance of peroxisome function in chronological aging of Saccharomyces
cerevisiae. *Aging Cell* **12**, 784-793 (2013).

21.   Zhou, B. *et al.* Midlife gene expressions identify modulators of aging through dietary
interventions. *Proc Natl Acad Sci U S A* **109**, E1201-1209 (2012).

22.   Curran, S.P. & Ruvkun, G. Lifespan regulation by evolutionarily conserved genes
essential for viability. *PLoS Genet* **3**, e56 (2007).

23.   Turi, Z., Lacey, M., Mistrik, M. & Moudry, P. Impaired ribosome biogenesis:
mechanisms and relevance to cancer and aging. *Aging (Albany NY)* **11**, 2512-2540
(2019).

24.   Russell, J. & Zomerdijk, J.C. RNA-polymerase-I-directed rDNA transcription, life and
works. *Trends Biochem Sci* **30**, 87-96 (2005).

25.   Ruggero, D. & Pandolfi, P.P. Does the ribosome translate cancer? *Nat Rev Cancer* **3**,
179-192 (2003).

26.   Kornprobst, M. *et al.* Architecture of the 90S Pre-ribosome: A Structural View on the
Birth of the Eukaryotic Ribosome. *Cell* **166**, 380-393 (2016).

27.   Woolford, J.L., Jr. & Baserga, S.J. Ribosome biogenesis in the yeast Saccharomyces
cerevisiae. *Genetics* **195**, 643-681 (2013).

28.   Zhang, J. *et al.* Assembly factors Rpf2 and Rrs1 recruit 5S rRNA and ribosomal proteins
rpL5 and rpL11 into nascent ribosomes. *Genes Dev* **21**, 2580-2592 (2007).

29.   Dragon, F. *et al.* A large nucleolar U3 ribonucleoprotein required for 18S ribosomal RNA
biogenesis. *Nature* **417**, 967-970 (2002).

30.   Grandi, P. *et al.* 90S pre-ribosomes include the 35S pre-rRNA, the U3 snoRNP, and 40S
subunit processing factors but predominantly lack 60S synthesis factors. *Mol Cell* **10**,
105-115 (2002).

31.   Mayer, C. & Grummt, I. Ribosome biogenesis and cell growth: mTOR coordinates
transcription by all three classes of nuclear RNA polymerases. *Oncogene* **25**, 6384-6391
(2006).

32.   van Riggelen, J., Yetil, A. & Felsher, D.W. MYC as a regulator of ribosome biogenesis
and protein synthesis. *Nat Rev Cancer* **10**, 301-309 (2010).

33.   Gutierrez, E., Wiggins, D., Fielding, B. & Gould, A.P. Specialized hepatocyte-like cells
regulate Drosophila lipid metabolism. *Nature* **445**, 275-280 (2007).
943

944 34. Schliebs, W., Girzalsky, W. & Erdmann, R. Peroxisomal protein import and ERAD:
945   variations on a common theme. *Nat Rev Mol Cell Biol* **11**, 885-890 (2010).
946

947 35. Mast, F.D. *et al.* A Drosophila model for the Zellweger spectrum of peroxisome
948   biogenesis disorders. *Dis Model Mech* **4**, 659-672 (2011).
949

950 36. Okumoto, K. *et al.* Cysteine ubiquitination of PTS1 receptor Pex5p regulates Pex5p
951   recycling. *Traffic* **12**, 1067-1083 (2011).
952

953 37. Landis, G.N. *et al.* The progesterone antagonist mifepristone/RU486 blocks the negative
954   effect on life span caused by mating in female Drosophila. *Aging (Albany NY)* **7**, 53-69
955   (2015).
956

957 38. Chen, C.L. *et al.* Proteomic mapping in live Drosophila tissues using an engineered
958   ascorbate peroxidase. *Proc Natl Acad Sci U S A* **112**, 12093-12098 (2015).
959

960 39. Faust, J.E., Verma, A., Peng, C. & McNew, J.A. An inventory of peroxisomal proteins
961   and pathways in Drosophila melanogaster. *Traffic* **13**, 1378-1392 (2012).
962

963 40. Park, S., Choi, S.G., Yoo, S.M., Son, J.H. & Jung, Y.K. Choline dehydrogenase interacts
964   with SQSTM1/p62 to recruit LC3 and stimulate mitophagy. *Autophagy* **10**, 1906-1920
965   (2014).
966

967 41. Cinnamon, E. *et al.* Drosophila Spidey/Kar Regulates Oenocyte Growth via PI3-Kinase
968   Signaling. *PLoS Genet* **12**, e1006154 (2016).
969

970 42. Seal, R.P., Daniels, G.M., Wolfgang, W.J., Forte, M.A. & Amara, S.G. Identification and
971   characterization of a cDNA encoding a neuronal glutamate transporter from Drosophila
972   melanogaster. *Recept Channels* **6**, 51-64 (1998).
973

974 43. Gonzalez, E.A., Garg, A., Tang, J., Nazario-Toole, A.E. & Wu, L.P. A glutamate-
975   dependent redox system in blood cells is integral for phagocytosis in Drosophila
976   melanogaster. *Curr Biol* **23**, 2319-2324 (2013).
977

978 44. Deschenes, M. & Chabot, B. The emerging role of alternative splicing in senescence and
979   aging. *Aging Cell* **16**, 918-933 (2017).
980

981 45. Fu, X.D. & Ares, M., Jr. Context-dependent control of alternative splicing by RNA-
982   binding proteins. *Nat Rev Genet* **15**, 689-701 (2014).
983

984 46. Lee, M.S. *et al.* Modulation of alternative splicing by expression of small nuclear
985   ribonucleoprotein polypeptide N. *FEBS J* **281**, 5194-5207 (2014).
986

987 47. Chabot, B. & Shkreta, L. Defective control of pre-messenger RNA splicing in human
988   disease. *J Cell Biol* **212**, 13-27 (2016).
989

990   48.   Srinivas, U.S., Tan, B.W.Q., Vellayappan, B.A. & Jeyasekharan, A.D. ROS and the DNA
991   damage response in cancer. *Redox Biol* **25**, 101084 (2019).
992

993   49.   Shokolenko, I., Venediktova, N., Bochkareva, A., Wilson, G.L. & Alexeyev, M.F.
994   Oxidative stress induces degradation of mitochondrial DNA. *Nucleic Acids Res* **37**, 2539-
995   2548 (2009).
996

997   50.   Gonzalez, K.L. *et al.* Disparate peroxisome-related defects in Arabidopsis pex6 and
998   pex26 mutants link peroxisomal retrotranslocation and oil body utilization. *Plant J* **92**,
999   110-128 (2017).
1000

1001   51.   Kao, Y.T., Fleming, W.A., Ventura, M.J. & Bartel, B. Genetic Interactions between
1002   PEROXIN12 and Other Peroxisome-Associated Ubiquitination Components. *Plant*
1003   *Physiol* **172**, 1643-1656 (2016).
1004

1005   52.   Park, E.S., Yoo, Y.J. & Elangovan, M. The opposite role of two UBA-UBX containing
1006   proteins, p47 and SAKS1 in the degradation of a single ERAD substrate, alpha-TCR. *Mol*
1007   *Cell Biochem* **425**, 37-45 (2017).
1008

1009   53.   Fernandez-Vizarra, E., Tiranti, V. & Zeviani, M. Assembly of the oxidative
1010   phosphorylation system in humans: what we have learned by studying its defects.
1011   *Biochim Biophys Acta* **1793**, 200-211 (2009).
1012

1013   54.   Peeters, A. *et al.* Carbohydrate metabolism is perturbed in peroxisome-deficient
1014   hepatocytes due to mitochondrial dysfunction, AMP-activated protein kinase (AMPK)
1015   activation, and peroxisome proliferator-activated receptor gamma coactivator 1alpha
1016   (PGC-1alpha) suppression. *J Biol Chem* **286**, 42162-42179 (2011).
1017

1018   55.   Peeters, A. *et al.* Mitochondria in peroxisome-deficient hepatocytes exhibit impaired
1019   respiration, depleted DNA, and PGC-1alpha independent proliferation. *Biochim Biophys*
1020   *Acta* **1853**, 285-298 (2015).
1021

1022   56.   Shinde, A.B. *et al.* Mitochondrial disruption in peroxisome deficient cells is hepatocyte
1023   selective but is not mediated by common hepatic peroxisomal metabolites.
1024   *Mitochondrion* **39**, 51-59 (2018).
1025

1026   57.   Mittal, M., Siddiqui, M.R., Tran, K., Reddy, S.P. & Malik, A.B. Reactive oxygen species
1027   in inflammation and tissue injury. *Antioxid Redox Signal* **20**, 1126-1167 (2014).
1028

1029   58.   Murphy, M.P. How mitochondria produce reactive oxygen species. *Biochem J* **417**, 1-13
1030   (2009).
1031

1032   59.   Zhou, W. & Zhao, M. How Hippo Signaling Pathway Modulates Cardiovascular
1033   Development and Diseases. *J Immunol Res* **2018**, 3696914-3696914 (2018).
1034

1035   60.   Johnson, R. & Halder, G. The two faces of Hippo: targeting the Hippo pathway for
1036   regenerative medicine and cancer treatment. *Nat Rev Drug Discov* **13**, 63-79 (2014).
1037

1038   61.   Lapi, E. *et al.* PML, YAP, and p73 are components of a proapoptotic autoregulatory
1039   feedback loop.
1040

1041   62.   Bertini, E., Oka T Fau - Sudol, M., Sudol M Fau - Strano, S., Strano S Fau - Blandino, G.
1042   & Blandino, G. YAP: at the crossroad between transformation and tumor suppression.
1043

1044   63.   Strano, S. *et al.* The transcriptional coactivator Yes-associated protein drives p73 gene-
1045   target specificity in response to DNA Damage.
1046

1047   64.   Cottini, F., Anderson, K.C. & Tonon, G. Awakening the Hippo co-activator YAP1, a
1048   mercurial cancer gene, in hematologic cancers.
1049

1050   65.   Matallanas, D. *et al.* RASSF1A elicits apoptosis through an MST2 pathway directing
1051   proapoptotic transcription by the p73 tumor suppressor protein. *Mol Cell* **27**, 962-975
1052   (2007).
1053

1054   66.   Cottini, F. *et al.* Rescue of Hippo coactivator YAP1 triggers DNA damage-induced
1055   apoptosis in hematological cancers. *Nat Med* **20**, 599-606 (2014).
1056

1057   67.   Xiao, Q. *et al.* Depletion of CABYR-a/b sensitizes lung cancer cells to TRAIL-induced
1058   apoptosis through YAP/p73-mediated DR5 upregulation. *Oncotarget* **7**, 9513-9524
1059   (2016).
1060

1061   68.   Di Cara, F., Bulow, M.H., Simmonds, A.J. & Rachubinski, R.A. Dysfunctional
1062   peroxisomes compromise gut structure and host defense by increased cell death and Tor-
1063   dependent autophagy. *Mol Biol Cell* **29**, 2766-2783 (2018).
1064

1065   69.   Harries, L.W. *et al.* Human aging is characterized by focused changes in gene expression
1066   and deregulation of alternative splicing. *Aging Cell* **10**, 868-878 (2011).
1067

1068   70.   Feingold, K.R. & Grunfeld, C. Introduction to Lipids and Lipoproteins, in *Endotext*. (eds.
1069   K.R. Feingold *et al.*) (South Dartmouth (MA); 2000).
1070

1071   71.   Lai, C.Q., Parnell, L.D. & Ordovas, J.M. The APOA1/C3/A4/A5 gene cluster, lipid
1072   metabolism and cardiovascular disease risk. *Curr Opin Lipidol* **16**, 153-166 (2005).
1073

1074   72.   van Dijk, K.W., Rensen, P.C., Voshol, P.J. & Havekes, L.M. The role and mode of action
1075   of apolipoproteins CIII and AV: synergistic actors in triglyceride metabolism? *Curr Opin
1076   Lipidol* **15**, 239-246 (2004).
1077

1078   73.   Grosskopf, I. *et al.* Apolipoprotein A-V deficiency results in marked
1079   hypertriglyceridemia attributable to decreased lipolysis of triglyceride-rich lipoproteins
1080   and removal of their remnants. *Arterioscler Thromb Vasc Biol* **25**, 2573-2579 (2005).

1081
1082   74.   Kohan, A.B. Apolipoprotein C-III: a potent modulator of hypertriglyceridemia and
1083   cardiovascular disease. *Curr Opin Endocrinol Diabetes Obes* **22**, 119-125 (2015).
1084
1085   75.   Panse, V.G. & Johnson, A.W. Maturation of eukaryotic ribosomes: acquisition of
1086   functionality. *Trends Biochem Sci* **35**, 260-266 (2010).
1087
1088   76.   Saito, K. *et al.* Functional analysis of a novel glioma antigen, EFTUD1. *Neuro Oncol* **16**,
1089   1618-1629 (2014).
1090
1091   77.   Chaker-Margot, M., Hunziker, M., Barandun, J., Dill, B.D. & Klinge, S. Stage-specific
1092   assembly events of the 6-MDa small-subunit processome initiate eukaryotic ribosome
1093   biogenesis. *Nat Struct Mol Biol* **22**, 920-923 (2015).
1094
1095   78.   Perez-Fernandez, J., Martin-Marcos, P. & Dosil, M. Elucidation of the assembly events
1096   required for the recruitment of Utp20, Imp4 and Bms1 onto nascent pre-ribosomes.
1097   *Nucleic Acids Res* **39**, 8105-8121 (2011).
1098
1099   79.   Zhang, L., Wu, C., Cai, G., Chen, S. & Ye, K. Stepwise and dynamic assembly of the
1100   earliest precursors of small ribosomal subunits in yeast. *Genes Dev* **30**, 718-732 (2016).
1101
1102   80.   Sloan, K.E., Bohnsack, M.T., Schneider, C. & Watkins, N.J. The roles of SSU
1103   processome components and surveillance factors in the initial processing of human
1104   ribosomal RNA. *RNA* **20**, 540-550 (2014).
1105
1106   81.   Hofmann, E.R. *et al.* Identification and characterization of GRIM-1, a cell-death-
1107   associated gene product. *J Cell Sci* **123**, 2781-2791 (2010).
1108
1109   82.   Awad, D. *et al.* Inhibiting eukaryotic ribosome biogenesis. *BMC Biol* **17**, 46 (2019).
1110
1111   83.   Gadal, O. *et al.* A nuclear AAA-type ATPase (Rix7p) is required for biogenesis and
1112   nuclear export of 60S ribosomal subunits. *EMBO J* **20**, 3695-3704 (2001).
1113
1114   84.   Ferreira-Cerca, S., Poll, G., Gleizes, P.E., Tschochner, H. & Milkereit, P. Roles of
1115   eukaryotic ribosomal proteins in maturation and transport of pre-18S rRNA and ribosome
1116   function. *Mol Cell* **20**, 263-275 (2005).
1117
1118   85.   Kruger, T., Zentgraf, H. & Scheer, U. Intranucleolar sites of ribosome biogenesis defined
1119   by the localization of early binding ribosomal proteins. *J Cell Biol* **177**, 573-578 (2007).
1120
1121   86.   Zhang, D. *et al.* Aggregation of Ribosomal Protein S6 at Nucleolus Is Cell Cycle-
1122   Controlled and Its Function in Pre-rRNA Processing Is Phosphorylation Dependent. *J
1123   Cell Biochem* **117**, 1649-1657 (2016).
1124
1125   87.   Weller, S., Gould, S.J. & Valle, D. Peroxisome biogenesis disorders. *Annu Rev Genomics
1126   Hum Genet* **4**, 165-211 (2003).

88. Hasan, S., Platta, H. & Erdmann, R. Import of proteins into the peroxisomal matrix. *Frontiers in Physiology* **4** (2013).

89. Ogura, T. & Wilkinson, A.J. AAA+ superfamily ATPases: common structure--diverse function. *Genes Cells* **6**, 575-597 (2001).

90. Deosaran, E. *et al.* NBR1 acts as an autophagy receptor for peroxisomes. *J Cell Sci* **126**, 939-952 (2013).

91. Gonzalez, K.L. *et al.* A pex1 missense mutation improves peroxisome function in a subset of Arabidopsis pex6 mutants without restoring PEX5 recycling. *Proc Natl Acad Sci U S A* **115**, E3163-E3172 (2018).

92. Seo, J.G., Lai, C.Y., Miceli, M.V. & Jazwinski, S.M. A novel role of peroxin PEX6: suppression of aging defects in mitochondria. *Aging Cell* **6**, 405-413 (2007).

93. Weller, S. *et al.* Alternative splicing suggests extended function of PEX26 in peroxisome biogenesis. *Am J Hum Genet* **76**, 987-1007 (2005).

94. Chu, B.B. *et al.* Cholesterol transport through lysosome-peroxisome membrane contacts. *Cell* **161**, 291-306 (2015).

95. Maxfield, F.R. & Wustner, D. Intracellular cholesterol transport. *J Clin Invest* **110**, 891-898 (2002).

96. Park, H. *et al.* Peroxisome-derived lipids regulate adipose thermogenesis by mediating cold-induced mitochondrial fission. *J Clin Invest* **129**, 694-711 (2019).

97. Dirkx, R. *et al.* Absence of peroxisomes in mouse hepatocytes causes mitochondrial and ER abnormalities. *Hepatology* **41**, 868-878 (2005).

98. Islinger, M., Voelkl, A., Fahimi, H.D. & Schrader, M. The peroxisome: an update on mysteries 2.0. *Histochem Cell Biol* **150**, 443-471 (2018).

99. Jakobsson, A., Westerberg, R. & Jacobsson, A. Fatty acid elongases in mammals: their regulation and roles in metabolism. *Prog Lipid Res* **45**, 237-249 (2006).

100. Novikoff, P.M. & Novikoff, A.B. Peroxisomes in absorptive cells of mammalian small intestine. *J Cell Biol* **53**, 532-560 (1972).

101. Zaar, K., Volkl, A. & Fahimi, H.D. Association of isolated bovine kidney cortex peroxisomes with endoplasmic reticulum. *Biochim Biophys Acta* **897**, 135-142 (1987).

102. Grabenbauer, M., Satzler, K., Baumgart, E. & Fahimi, H.D. Three-dimensional ultrastructural analysis of peroxisomes in HepG2 cells. Absence of peroxisomal

1173   reticulum but evidence of close spatial association with the endoplasmic reticulum. *Cell*
1174   *Biochem Biophys* **32 Spring**, 37-49 (2000).
1175

1176   103.   Fahimi, H.D., Baumgart, E. & Volkl, A. Ultrastructural aspects of the biogenesis of
1177   peroxisomes in rat liver. *Biochimie* **75**, 201-208 (1993).
1178

1179   104.   Kovacs, W.J. *et al.* Peroxisome deficiency causes a complex phenotype because of
1180   hepatic SREBP/Insig dysregulation associated with endoplasmic reticulum stress. *J Biol*
1181   *Chem* **284**, 7232-7245 (2009).
1182

1183   105.   Faust, P.L. & Kovacs, W.J. Cholesterol biosynthesis and ER stress in peroxisome
1184   deficiency. *Biochimie* **98**, 75-85 (2014).
1185

1186   106.   Raman, N., Weir, E. & Muller, S. The AAA ATPase MDN1 Acts as a SUMO-Targeted
1187   Regulator in Mammalian Pre-ribosome Remodeling. *Mol Cell* **64**, 607-615 (2016).
1188

1189   107.   Kallstrom, G., Hedges, J. & Johnson, A. The putative GTPases Nog1p and Lsg1p are
1190   required for 60S ribosomal subunit biogenesis and are localized to the nucleus and
1191   cytoplasm, respectively. *Mol Cell Biol* **23**, 4344-4355 (2003).
1192

1193   108.   Saveanu, C. *et al.* Sequential protein association with nascent 60S ribosomal particles.
1194   *Mol Cell Biol* **23**, 4449-4460 (2003).
1195

1196   109.   Steffen, K.K. *et al.* Ribosome deficiency protects against ER stress in Saccharomyces
1197   cerevisiae. *Genetics* **191**, 107-118 (2012).
1198

1199   110.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
1200   (2013).
1201

1202   111.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
1203   for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
1204

1205   112.   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
1206   interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-
1207   15550 (2005).
1208

1209   113.   Mootha, V.K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation
1210   are coordinately downregulated in human diabetes. *Nat Genet* **34**, 267-273 (2003).
1211

1212   114.   Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased
1213   coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic*
1214   *Acids Res* **47**, D607-D613 (2019).
1215

1216   115.   Huang, D.W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis of
1217   large gene lists using DAVID bioinformatics resources. *Nature Protocols* **4**, 44-57
1218   (2009).

116. Huang da, W., Sherman, B.T. & Lempicki, R.A. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**, 1-13 (2009).

117. Schmittgen, T.D. & Livak, K.J. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* **3**, 1101-1108 (2008).

118. Martins, T. *et al.* TGFbeta/Activin signalling is required for ribosome biogenesis and cell growth in Drosophila salivary glands. *Open Biol* **7** (2017).

Figure 1 Transcriptomic analysis for peroxisomal stress response in *Drosophila* oenocytes and human cells.

Figure 2 Pex1, Pex12 knock-down show distinct transcriptional pattern from Pex5.

NAME	NES	FDR	Type of RNAi	Up or down regulated
Ribosome biogenesis in eukaryotes	2.44	<0.0001	Pex5	Down
Ribosome	1.79	0.021	Pex5	Down
DNA replication	1.68	0.057	Pex5	Down
Protein processing in endoplasmic reticulum	-1.77	0.052	Pex5	Up
Ribosome biogenesis in eukaryotes	2.89	<0.0001	Pex12	Down
DNA replication	2.11	<0.0001	Pex12	Down
RNA polymerase	1.91	0.001	Pex12	Down
RNA transport	1.85	0.002	Pex12	Down
Spliceosome	1.78	0.005	Pex12	Down
RNA degradation	1.67	0.002	Pex12	Down
Proteasome	-1.90	0.01	Pex12	Up
Protein processing in endoplasmic reticulum	-1.46	0.188	Pex12	Up
TOLL and IMD signaling pathway	1.74	0.041	Pex1	Down
DNA replication	1.70	0.034	Pex1	Down
Ribosome	-2.41	<0.0001	Pex1	Up
Proteasome	-1.84	0.012	Pex1	Up
Protein processing in endoplasmic reticulum	-1.54	0.190	Pex1	Up
Citrate cycle (TCA-cycle)	-1.43	0.295	Pex1	Up

Table 1 GSEA pathway analysis results under Pex5, Pex12 and Pex1 RNAi.

Figure 3 Peroxisomal stress induced endoplasmic reticulum changes in oenocytes.

Figure 4: Peroxisome dysfunction up-regulates inflammation and cholesterol pathways in human cells.

PEX5 C11A - DOWN

NAME	NES	FDR
Ribosome	2.35	<0.0001
Oxidative phosphorylation	1.97	0.024
Parkinson disease	1.72	0.013
Valine and isoleucine degradation	1.67	0.024

PEX5 C11A – UP (Top 20 pathways)

NAME	NES	FDR
Signaling pathways regulating pluripotency of stem cells	-2.06	<0.0001
Spnocerebellar ataxia	-2.00	<0.0001
MAPK signaling pathway	-1.94	0.001
MicroRNAs in cancer	-1.89	0.004
Osteoclast differentiation	-1.88	0.003
Neuroactive ligand-receptor interaction	-1.87	0.004
Regulation of lipolysis in adipocytes	-1.85	0.004
HIPPO signaling pathway	-1.84	0.004
Type II diabetes mellitus	-1.83	0.004
WNT signaling pathway	-1.83	0.004
Acute myeloid leukemia	-1.82	0.003
Breast cancer	-1.80	0.004
Cushing syndrome	-1.80	0.004
B cell receptor signaling pathway	-1.78	0.005
CAMP signaling pathway	-1.78	0.004
Aldosterone-regulated sodium reabsorption	-1.78	0.004
CGMP-PKG signaling pathway	-1.77	0.004
Proteoglycans in cancer	-1.76	0.005
Olfactory transduction	-1.76	0.004
TH1 and TH2 cell differentiation	-1.75	0.005

Table 2 Selected list of PEX5^{C11A} regulated pathways

Figure 5 Peroxisome dysfunction represses ribosomal genes in both flies and humans

Figure 6 Impaired ribosome biogenesis genes in human and flies