High-Throughput Sequencing can be used to Measure and Maximize Cultivated Diversity

Emily N. Junkins[a] and Bradley S. Stevenson[a]#

[a]Department of Microbiology and Plant Biology, University of Oklahoma, Norman, OK, USA

Running Head: Using sequencing to measure cultivated diversity

#Address correspondence to Bradley S. Stevenson, bradley.stevenson@ou.edu

Abstract word count: 359

Text (excluding Materials and Methods, references, tables, and figure legends) word count: 4129

Abstract

Molecular techniques continue to reveal a growing disparity between the immense diversity of microbial life and the small proportion that is in pure culture. The disparity, originally dubbed "the great plate count anomaly" by Staley and Konopka, has become even more vexing given our increased understanding of the importance of microbiomes to a host and the role of microorganisms in the vital biogeochemical functions of our biosphere. Searching for novel antimicrobial drug targets often focuses on screening a broad diversity of microorganisms. If diverse microorganisms are to be screened, they need to be cultivated. Recent innovative research has used molecular techniques to assess the efficacy of cultivation efforts, providing invaluable feedback to cultivation strategies for isolating targeted and/or novel microorganisms. Here, we aimed to determine the efficiency of cultivating representative microorganisms from a non-human, mammalian microbiome, identify those microorganisms, and determine the bioactivity of isolates. Molecular methods indicated that around 57% of the ASVs detected in the original inoculum were cultivated in our experiments, but nearly 53% of the total ASVs that were present in our cultivation experiments were *not* detected in the original inoculum. In light of our controls, our data suggests that when molecular tools were used to characterize our cultivation efforts, they provided a more complete, albeit more complex, understanding of which organisms were present compared to what was eventually cultivated. Lastly, about 3% of the isolates collected from our cultivation experiments showed inhibitory bioactivity against a multidrug-resistant pathogen panel, further highlighting the importance of informing and directing future cultivation efforts with molecular tools.

Importance

Cultivation is the definitive tool to understand a microorganism's physiology,

metabolism, and ecological role(s). Despite continuous efforts to hone this skill, researchers are

still observing yet-to-be cultivated organisms through high-throughput sequencing studies. Here,

we use the very same tool that highlights biodiversity to assess cultivation efficiency. When

applied to drug discovery, where screening a vast number of isolates for bioactive metabolites is

common, cultivating redundant organisms is a hindrance. However, we observed that cultivating

in combination with molecular tools can expand the observed diversity of an environment and its

community, potentially increasing the number of microorganisms to be screened for natural

products.

Introduction

Molecular techniques and sequencing technologies have helped to describe many

microbial communities and provide tools that allow us to study representative microorganisms

and their activities without isolation. Application of molecular approaches to the study of

microbiology have led to the discovery and characterization of rare biospheres of the most

extreme environments (1) and enabled researchers to decipher the metabolic potential and

activity of microorganisms from the human gut (2) to the bottom of the ocean (3). Now, changes

in microbial community structure and function can be monitored with molecular techniques

throughout the course of a disease (4) or with changing environmental conditions (5), such that

we may begin to predict the roles of microbial populations and outcomes over time (6, 7).

Isolating, or even just cultivating representative microorganisms is still considered the

best means towards understanding their physiology, metabolism, and potential ecological roles,

but this remains to be a major hurdle. The degree to which any microbial community is

represented in culture varies considerably. Since Staley and Konopka coined the term "great

plate count anomaly" (8), numerous studies have sought to quantify the culturable organisms

from various biomes (see (9) for a compiled list of most probable number (MPN) based

approaches). A recent study estimated that 81% of all microbial cells on Earth belong to

uncultured genera or higher classifications and about 25% of microbial cells are from uncultured

phyla (9). Specifically, 26% of marine, 31% of host-associated, 31% of eutrophic lakes, and 30%

of soil metagenomic sequences have cultured representatives at the family level (9). Not

surprisingly, human associated microbiomes have more cultured representatives, likely affected

by a focused effort (10). Sequencing technology has shed light onto the previously unknown

diversity of the microbial world, outpacing our ability to cultivate these organisms by an

estimated rate of 2.4- and 2.5-fold for bacteria and archaea, respectively (11).

Efforts to isolate yet uncultured microorganisms have gathered momentum as researchers

have begun to apply new technologies and innovations (12-14). Microbiologists routinely

manipulate the composition of growth media and treatments of the inocula in order to increase

the diversity of microorganisms represented in culture (e.g., (14-22)). As an example, treating an

inoculum with ethanol can select for spore-forming bacteria by killing vegetative cells and

leaving spores intact that would otherwise be overgrown and remain unisolated (23).

Furthermore, antioxidants in media can support the cultivation of both aerobic and anaerobic

organisms, by using compounds that alleviate oxidative stress (24, 25). With the goal of this

study being to cultivate many, different microorganisms, the approaches described above were of

particular interest. We hypothesized that they would select for gut-associated bacteria;

antioxidants would help mitigate oxygen stress for organisms best suited for microoxic or anoxic

environments, and ethanol treatment would reduce the abundant vegetative cells, allowing us to

better access the many spore-forming taxa that reside in the gut.

When cultivation approaches are coupled with molecular analyses, the total diversity and

identity of cultivated taxa can be determined (26-28), and specific, targeted taxa can be detected

(13). More recently, high throughput sequencing has made it possible to quantify exactly what

proportion of the original sample has been cultivated (26, 29, 30). Studies using this approach

showed that roughly 23% of microorganisms in bovine rumen fluid were cultivable in liquid

enrichments (29), while 96% of those in human bronchoalveolar lavage samples (30), and 60%

of those detected in/on American toads (26) were cultivated. A recent study characterized the

microbial diversity within the same inoculum spread onto both high- and low-nutrient media by

harvesting biomass on each plate and identifying all taxa present (26). Their approach built upon

a previous study that sought to isolate abundant but yet-uncultured taxa using defined, low

nutrient media and a PCR-based screening approach using group specific primers, termed plate-

wash PCR (PWPCR) (13). Medina et al. confirmed that low nutrient-media would allow for the

growth of a more diverse subset of bacteria from the original sample and that molecular

screening (i.e., 16S PWPCR) is a practical way to characterize the microbial community growing

on an entire agar plate (26). This data is invaluable in identifying medium composition, inoculum

treatments, and incubation conditions that can target certain taxa, or maximize the cultivated

diversity for large-scale isolation efforts (see 16, 18). However, these studies using solid agar

plates may have overestimated diversity for some taxa by focusing on the relative abundance of

taxa in their analyses (see Discussion).

Pure cultures have always been the "gold standard" for the in-depth study of

microorganisms. The need for isolation is especially important when screening for the

production of secondary metabolites for drug discovery. Historically, the process of isolating

microorganisms for drug discovery has relied mainly on high throughput cultivation efforts that

increase the number of isolates being screened (e.g., (31-33)), enrich for bioactive production

from known producers like *Streptomycetes* spp. (34), or focus on novel or rare taxa thought to

have uncharacterized biosynthetic capabilities (35-37). We aimed to do the same by sampling a

diverse, host-associated microbial community and cultivating them under conditions designed to

obtain as many different organisms as possible through the use of different media types.

Screening cultivated microorganisms with the broadest diversity is expected to provide the

greatest chance of uncovering natural products that are novel drug targets. Previous studies have

focused on extreme environments for novel natural products (38), but we expand on this idea to

include non-human mammals. We predicted that this would increase the diversity of

microorganisms and compounds screened for useful antimicrobial compounds and argue that the

shared life-history between a host and its host-associated microbial communities may produce

compounds that are relatively less cytotoxic to the host. This idea was originally supported by a

survey of microbiomes collected from roadkill mammals, which produced two new cyclic

lipodepsipeptides of pharmaceutical interest (33).

Here, we sought to use both general and selective media along with various treatments to

the inocula to collectively enrich for as many different microorganisms as possible from the oral

and gut microbiomes of a roadkill mammal (racoon). The goals for this study were to (1)

compare the diversity of cultured bacterial communities on multiple types of media combined

with various treatments of the inoculum, (2) determine which media, treatments, or combination

cultivated the highest richness of bacterial taxa, and (3) determine how many isolates produced

bioactive molecules against a panel of multi-drug resistant pathogenic microorganisms.

Materials and Methods

Sample Collection and Storage

A 3 mile stretch of Oklahoma State Highway 9 in Norman, OK between the University of

Oklahoma, Norman campus and Lake Thunderbird was used for our opportunistic sampling area.

This well-traveled highway runs through undeveloped prairie, scrub and forest, cattle ranches,

and rural residential land, providing many mammalian roadkill specimens. Sample collection

was conducted under Oklahoma Scientific Collector Permit #5250 (33). The oral cavity and

rectum of a racoon (*Procyon lotor*) deceased less than 8 h was sampled in triplicate (n=3) with

sterile, nylon swabs. To ensure equal biomass was collected for immediate sequencing and

subsequent cultivation, two swabs were used to sample each orifice simultaneously (to increase

amount of sample) and stored together in a sterile conical tube containing 3.0 mL 1X PBS, stored

on ice, and immediately transferred back to the laboratory (under 45 minutes). At the laboratory,

each set of swabs was vortexed to suspend cells. A 900 µL aliquot of the cell suspension

intended for molecular analysis was centrifuged at 10,000 x g for 1 min, the supernatant was

removed. The cell pellet was resuspended in 750 mL of BashingBead Buffer (Zymo Research

Corp.) and transferred to BashingBead Lysis Tubes (Zymo Research Corp., Irvine, CA) prior to

lysis through homogenization in a Mini-BeadBeater-8 (BioSpec Products Inc., Bartlesville, OK)

at maximum speed for 45 seconds. Homogenized samples were then stored at -20˚C until needed

for DNA extraction. For cultivation, the remaining 2.1 mL of the cell suspension for each sample

was serially diluted with PBS. Aliquots of 50 µL from the 10^{0}-10^{-6} dilutions were spread onto 13

different media/treatment combinations (see Table S1). This 2.1 mL volume represented the

exact amount needed for cultivation with no suspension left over so that the 10^{0} dilution was not

diluted, and represented the surface area of the swab. In all, each inoculum was spread onto the

agar media, in triplicate, for each media type and incubated for 6 days at 30˚C.

Culture-Dependent Bacterial Community Analysis

Cultivation was conducted with a combination of broad-spectrum media with decreased

nutrients and selective media in order to "cast a wide net" and maximize the cultivable diversity

of each inoculum (see quasi-factorial design in Table S1). The media used included 0.1X

strength tryptic soy broth (Bacto, USA) with 1.5% agar, ROXY and derivatives with hemin (0.1

g/L) and alpha-ketoglutarate (2.0 g/L) (25), 0.25X strength R2A medium (15), and blood agar

(per L: 10.0 g meat extract, 10.0 g peptone, 5.0 g sodium chloride, 15 g agar, 5% sheep's blood

(ThermoFisher, USA), pH 7). To date, this is the first instance using ROXY medium and plate-

wash PCR techniques. In addition to different media types, one variation included the

pretreatment of the inocula for 4 hours in 70% ethanol at 22-24°C (23). Catalase (40,000 U/L)

was added as a treatment to TSA, ROXY and R2A to remove peroxides produced during

autoclaving (19, 20). The last treatment included the addition of streptomycin at 50 mg/L as a

broad spectrum selection agent against Gram-negative and some Gram-positive species (39).

To test which combinations of media and treatments resulted in cultivated communities

that represented the diversity observed in the original sample, we grew each inoculum (mouth or

rectum) on each combination of media/treatment types and sequenced the resulting biomass.

Collection of Total Biomass from Agar Plates, DNA Extraction and PCR (PWPCR)

Biomass was harvested from each agar plate in order to identify the microorganisms

present and to compare this to culture-independent analyses of the inoculum. The biomass from

each agar plate was harvested by adding 2.0 mL of PBS to the surface of the plate and

suspending the colony biomass by agitation with a sterile spreader. The suspended colony

biomass was collected and transferred to a 2.0 mL microcentrifuge tube, pelleted at 10,000 x g

for 1 min, and resuspended in 750 uL of BashingBead Buffer (Zymo Research Corp.). Each

sample was transferred to a BashingBead Lysis Tube (Zymo Research Corp.) and homogenized

for 45 seconds at maximum speed using a Mini-BeadBeater-8 (BioSpec Products Inc.,

Bartlesville, OK). The lysed samples were stored at -20˚C until DNA was extracted.

DNA Extraction and Sequencing

Before DNA extraction, each sample was thawed on ice and homogenized for 30 seconds

at maximum speed using a Mini-BeadBeater-8 (BioSpec Products Inc., Bartlesville, OK). DNA

was extracted according to manufacturer specifications using the Zymo *Quick*-DNA

Fungal/Bacterial Miniprep kit (Cat# D6005, Zymo Research Corp., Irvine, CA). For community

characterization, a conserved region of the SSU rRNA gene of most bacteria, archaea, and

eukarya was amplified using primers 515F-Y and 926R (40) via the following PCR protocol:

initial denaturation at 94˚C for 2 min, followed by 30 cycles of denaturation at 94˚C for 45 s,

annealing at 50˚C for 45 s, and extension at 68˚C for 90 s, with a final extension at 68˚C for 5

min. These primers produced two amplicons, a ~400 bp fragment for bacteria and archaea, and a

600 bp fragment for eukaryotes. The forward primer 515F-Y (5'-<u>GTA AAA CGA CGG CCA</u>

<u>G</u> CCG TGY CAG CMG CCG CGG TAA-3') contains the M13 tag (underlined) fused to the 5'

end of the forward primer (41). The reverse primer 926R (5'-CCG YCA ATT YMT TTR AGT

TT-3') was unmodified from Parada et. al 2015. Each PCR contained 5 PRIME HOT master mix

(1X; 5 PRIME Inc., Gaithersburg, MD), 0.2 µM of each primer, and 3.0 µL of extracted DNA at

a final volume of 50 µL. The amplified fragments were purified using Sera-Mag magnetic beads

(GE) with the AmPureXP (Beckman Coulter) protocol at a final concentration of 1.8x v/v. After

purification, 3 μL of each PCR product, 1x 5 PRIME HOT master mix (Quantabio,

Massachusetts, USA), 0.2μM of the 926R primer, and 0.2 μM of a specific 12 bp oligonucleotide

was used in a separate barcoding PCR (6 cycles) in 50 μL reactions to attach a unique barcode to

amplicons of each library. The same thermocycler protocol was used as above but only run for 6

cycles. The now barcoded amplicons were purified using Sera-Mag (GE) beads with the

AmPureXP (Beckman Coulter) protocol to a final volume of 40 μL, quantified using the QuBit

HS DS DNA assay kit (Thermo Fisher Scientific Inc., Waltham, MA), and pooled in equimolar

amounts. The pooled, barcoded amplicon libraries were then concentrated to a final volume of 40

μL (209 ng/ μL) with an Amicon-Ultra filter (Millipore, Burlington, MA, USA) following

manufacturer's protocol. The combined amplicon libraries were denatured according to Miseq

library preparations protocol (Illumina, San Diego, CA, USA). The sample was loaded at a

concentration of 10 pM and sequenced using 2x250 paired-end strategy on the Miseq (Illumina

San Diego, CA, USA) platform for 251 cycles.

ASV Inference and Bacterial Community Characterization

Barcodes from raw SSU rRNA amplicon sequences were removed and demultiplexed

using QIIME v 1.9.1 (42). Demultiplexed reads were trimmed for adapters and quality filtered as

a part of the *dada2* pipeline (43) and amplicon sequence variants (ASVs) were inferred using the

forward reads (203 bp). Taxonomy was assigned using the SILVA database v32 (44, 45). The

final dataset consisted of ~1.7 million reads resulting in 463 ASVs with a median of 17,347

sequences per sample (min = 132, max= 266,678).

Data Availability: Sequence data has been deposited at NCBI's Sequence Read Archive (SRA)

database under accession number PRJNA675861.

Data Analysis

Community diversity was analyzed using the *phyloseq* (46) and *vegan* (47) packages in

R. Relative abundance was only used to directly assess the diversity of the original inocula.

Presence/absence and richness were used as the main metrics to compare cultivated samples

because of the variation in biomass (i.e. colony size) between microbial colonies on and between

agar plates. Also, because of the sampling procedure with duplicate swabs used in triplicate, the

data from the molecular control was pooled for analysis since the variability in swabbing would

be different than the variability of triplicates resulting from cultivation.

Alpha diversity was calculated using richness of ASVs, while beta diversity was

measured with non-metric multidimentional scaling (NMDS) using Jaccard distances. Statistical

comparisons across media types, treatment, and orifices where done using permanova with the

adonis2 function in the *vegan* package (47). Due to the nestedness of orifice and cultivation

conditions, orifice was used as a strata or fixed factor, and media and treatment were used as

predictor variables (~media*treatment). Communities were also compared without using orifice

as a fixed factor to capture any influence that orifice could have on the cultivable communities.

To fully understand the significance described in the permanova, a permdisp with the *betadisper*

function in *vegan* was used to describe any within sample variance (i.e. across replicate agar

plates) that could explain any significant differences detected in the permanova. Hypothesis

testing via anova with permutations (n=999) was used with permdisp to determine any

significant differences in variation within samples.

To identify certain ASV families more associated with an orifice, medium, or treatment, a

species indicator analysis was performed using the *indicspecies* package (permutations = 999)

(48, 49).

Bioassays and Isolate Identification

Colonies were selected for isolation based on morphology, with the intent of sampling as

many different colony morphologies as possible. From average of 8.1 x 10^{10} CFUs/mL cultivated

across all medium types and dilutions, a total 238 colonies were chosen for isolation based on

differential colony morphology from the mouth and rectum on a subset of the media types (0.1X

TSA, ROXY with alpha-ketoglutarate and hemin, and 0.25X R2A). To determine bioactivity,

soft agar overlays in 0.1X TSA (0.8% agar) were used to test for growth inhibition of a panel of

drug-resistant pathogens (*Klebsiella pneumonia* ATCC# 13883, *Enterococcus faecium* ATCC#

51559, *Pseudomonas aeruginosa* ATCC# 10145, and *Candida albicans* ATCC# 565304). To

prepare for the bioassay, pathogens were incubated overnight, shaking at 250 rpm at 30°C, in

0.1X tryptic soy broth (TSB) (Bacto, USA). A 400 μL aliquot of each pathogen was used to

inoculate 3.6 mL of molten TSA soft agar (at 55 ºC), mixed, and immediately poured onto a

0.1X TSA agar plate for a final overlay volume of 4 mL. Colony material from each isolate was

transferred to the bioassay plate with a sterile toothpick by etching an 'X' into the overlay. The

inoculated bioassay plates were incubated at 30 ºC for 24 hours. Inhibition of the pathogen in the

overlay was characterized by a zone of clearing surrounding the isolate (Figure S1). For

cryopreservation, each isolate was grown in 5 mL of 0.1X TSB in a 16 x 150 mm test tube

shaking at 250 rpm for 48 h. An aliquot of 800 uL of the culture was transferred to a 2 mL screw

cap tube, along with 200 uL of 80% glycerol, for a final concentration of 20% glycerol. After

mixing by vortex for 30 s, it was then stored at -80ºC.

Partial SSU rRNA sequences were used to identify each isolate. Genomic DNA was

extracted from each isolate using QuickExtract (Lucigen, Wisconsin, USA) according to

manufacturer's protocol from the above mentioned 0.1X TSB cultures. The SSU rRNA gene was

amplified using primers 8F (3'-ATGC-5') and 1492R (3'-ATGC-5') at a concentration of 0.2

μM, 1x 5 PRIME HOT master mix (Quantabio, Massachusetts, USA) , and 2.0 μL of DNA

template per 50 μL reaction via the following PCR protocol: initial denaturation at 94˚C for 2

min, followed by 30 cycles of denaturation at 94˚C for 30 s, annealing at 55˚C for 45 s, and

extension at 72˚C for 45 s, with a final extension at 72˚C for 10 min. Amplified fragments were

purified using Sera-Mag magnetic beads (GE) with the AmPureXP (Beckman Coulter) protocol

at a final concentration of 1.8x v/v. Purified amplicons (a total of 192 samples) were sent for

Sanger sequencing using the 8F primer (Genewiz, New Jersey, USA). Resulting sequences and

chromatograms were assessed for quality using IGV (50). Sequences with low quality (< Q20)

were removed, low quality ends were trimmed with MEGA X (51), and the final sequences were

identified using the SILVA v. 132 classifier online server (44). A total of 117 isolates had

suitable reads and met quality thresholds for taxonomic identification.

Results

Cultivated Microbial Diversity Differed Between Orifice and Cultivation Condition

As expected, a library of SSU rRNA genes from the original inocula showed that it had

higher richness (alpha diversity) than the cultivated organisms from both orifices (Figure 1). The

microbial community sampled from the rectum had higher overall richness (153 ASVs)

compared to that of the mouth (86 ASVs) (Figure 1). Overall, we were able to cultivate 57.3% of

the ASVs detected in the inocula (58.1% from the mouth and 57.5% from the rectum).

Interestingly, about half (52.7%) of the ASVs detected from the cultivation experiments were not

represented in either library from the inocula (i.e. molecular control) (Figure 2). The number of

ASVs detected on a certain medium or treatment differed between orifices. For instance,

cultivation on TSA medium treated with catalase resulted in only 29 ASVs from the rectum

inoculum compared to around 70 different ASVs from the mouth. The fewest number of ASVs

was detected on the ROXY medium amended with alpha-ketoglutarate and hemin from both the

mouth and rectum. The cultivated beta diversity differed based on orifice and cultivation

condition. Specifically, community structure differed based on orifice and treatments, more so

than base medium type (Figure 3). Differences in the cultivated community composition and

diversity between medium and treatment confirmed that, collectively, using different cultivation

techniques increased the number of organisms cultivated from each inoculum.

Selective Treatments Increased Cultivated Richness Compared to Media Type or Orifice

An indicator species analysis was used to determine which microbial families were

significantly associated with certain conditions. A total of ten families were identified to be

significantly ($p < 0.05$) associated with orifice, media type, and/or treatment. Only one family,

Aeromonadaceae was significantly associated with a particular orifice, the mouth. This family

was also the only family significantly ($p = 0.001$) associated with a certain base medium type,

ROXY (Table 1). Treatment of the medium and/or the inoculum selected for the most diverse set

of indicator species, with a total of 9 families significantly ($p < 0.05$) associated with a certain

treatment or treatments (Table 1). Certain bacterial families like Staphylococcaceae and

Paenibacillaceae were associated with the ethanol pretreatment of inocula, whereas the

Wohlfarhtimonodaceae and Metaschinikowiceae were more associated with the addition of

streptomycin to media (Table 1).

Despite some unique taxa being associated with certain conditions, much of the cultivable

microbial diversity was shared between the mouth and the rectum (Figure 5), indicated by the

lack of significant differences in community structure described by the permanova and permdisp

analyses (discussed below). Based on molecular analyses, 27 families where shared between the

mouth and rectum before cultivation, while 6 families were unique to the mouth and 6 families

were unique to the rectum. Negative cultivation controls (e.g., PBS) yielded no growth,

supporting the assumption that any biomass collected from plates originated from the inoculum.

Lastly, our cultivation experiments led to the detection of 18 families from the mouth and 8

families from the rectum that were not detected with direct molecular analysis of the inoculum.

This accounted for around half (52.7%) of the ASVs detected during cultivation (Figure 2). This

could be explained by differences in biomass and growth characteristics not being captured by

sequence data from the inocula (discussed below).

The Raccoon Microbiome Contained Cultivable Bioactive Bacteria

A total of 238 isolates were collected from a subset of media (0.25X R2A, 0.1X TSA,

and ROXY with alpha-ketoglutarate and hemin). Many of these isolates were expected to be

redundant, since the diversity of the initial inoculum was low (Figure 1). This was confirmed

with many isolates redundant at the genus level and the majority of the identifiable isolates

belonging to genera *Serratia* (34.2%) and *Klebsiella* (17.9%) (Figure S2). Each of the 238

isolates was assayed for antimicrobial activity against a panel of multi-drug resistant pathogens.

A total of 7 isolates showed antimicrobial activity, identified as a zone of clearing of one of the

antibiotic resistant panel organisms. This equated to a ~3% "hit" rate against already resistant

organisms. Six of these isolates were recovered from the mouth, while one, a *Bacillus* sp., was

recovered from the rectum. Three isolates were recovered from R2A, 3 from TSA and 1, an

Enterobacteriaceae sp., from ROXYakgh. Three showed activity against *Klebsiella pneumonia*,

three against *Enterococcus faecium*, and one, a *Pseudomonas* sp., against *Candida albicans.* Five

of the isolates were identified by partial 16S rRNA sequence identity (Table 2).

The Composition of Cultivated Taxa was Highly Variable

Due to the nestedness of this experiment (i.e. taxa observed during cultivation theoretically

would be observed in sequencing), the stratification of taxa based on sampled orifice was

corrected for hypothesis testing with permanova by defining orifice as a stratum in the *adonis2*

function in the *vegan* package (47). The majority of the explained variation and significant

differences in cultivated taxa were due to treatment of the inoculm or medium type (F= 2.44, R^2

= 0.12, p = 0.001), while base medium type alone explained only ~5% of the observed variation

(F= 1.54, R^2 = 0.05, p = 0.002). The interactions of base medium and treatment only explained

~6% of the variation (F= 1.52, R^2= 0.06, p= 0.208). An analysis of variances (permdisp) was

used to determine if any significance was driven by a dispersion effect rather than a location

effect for orifice, base medium type, treatments, and medium+treatments. Mean distances to

centroids did not differ significantly for orifice (F=0.69, p=0.41; Mouth 0.51; Rectum 0.53) or

medium+treatment (F= 1.38, p=0.20; TSA=0.47, TSAcatalase=0.52, TSAstrep=0.51,

TSAethanol=0.45, R2A=0.44, R2Acatalase=0.47, R2Aethanol=0.45, Blood=0.45,

Bloodethanol=0.50, Bloodstrep=0.53, ROXY=0.48, ROXYcatalase=0.49, ROXYgh=0.52),

meaning little inter-sample variation was detected between replicates. In contrast, the mean

distances did differ for treatment (F=3.3, p=0.02; alpha-ketoglutarate+hemin=0.52, no treatment

(base)=0.50, catalase=0.53, ethanol=0.49, streptomycin=0.56) and base medium type (F=4.4,

p=0.01; Blood=0.56, R2A=0.50, ROXY=0.53, TSA=0.55). The significance in dispersion

within the treatment and media could correspond to the significance in permanova results, where

much of the variation explained in the permanova may be due to the intrinsic variability and

stochasticity among replicates of the same inoculum.

Discussion

We used mouth and rectal samples from a roadkill mammal (raccoon, *Procyon lotor*) to

(1) compare the diversity of cultured bacterial communities on multiple media types and/or

treatments of the inoculum, (2) determine which media/treatments cultivated the highest richness

of bacterial taxa, and (3) determine if any isolated organisms produced bioactive molecules

against a drug resistant pathogen panel. The occurrence of antimicrobial resistance is increasing

at a faster rate than new therapies are entering the market (52, 53). Since nearly two-thirds of our

current antimicrobial drugs are linked to microorganisms (52, 54), one of our goals was to

cultivate a diverse array of bacteria from a non-human mammalian microbiome and screen these

isolates for the production of antimicrobial metabolites. Growing microbial populations in the

laboratory that are representative of the overall microbial diversity within an inoculum continues

to be a major limitation for the field of microbiology. Here, we sought to use a combination of

selective medium types and treatments of the inocula that might enrich for certain groups present

in the oral and gut microbiomes of a roadkill racoon. In order to capture a more comprehensive

sample of microbial diversity, we used high-throughput sequencing of 16S rRNA gene amplified

libraries to characterize the original inocula and the cultivated populations that were washed

from the surface of agar media.

We anticipated finding a significantly lower percentage of cultivable organisms relative

to the molecular-based measures of diversity from each orifice, which we described as the

cultivated percentage. We also anticipated that the identity of recovered taxa would be similar

between media types and even orifices. The differences between organisms cultivated on

different media, from different orifices was low. However, when observing the significance

described in the permanova and permdisp, the percentage of explained variation was low. This

suggested that the stoichiometry of diverse and variable inocula could play a large part in the

outcome of this and any similar cultivation experiment. Based on the low percentage of

explained variation, we also approached this data set from the perspective of the unobserved or

residual variation, such that some of this discussion will aim to pose questions about what makes

cultivation so difficult. Overall, we observed that cultivation yielded not just a fraction of what is

detected in sequence data, but rather expanded the total diversity observed in the original sample.

Cultivation can be optimized at large scales

Convergence of microbial communities occurred between the mouth and rectum for both

the molecular samples and the cultivated communities. This was expected since as an animal

undergoes decomposition, the microbial diversity decreases, especially in rectal communities

(55), and becomes dominated by only certain taxa, namely Proteobacteria (56). In our samples,

despite being a single time point roughly 8 hours after death, the mouth was dominated by

Proteobacteria while the rectum samples contained mostly Firmicutes, Proteobacteria, and

Bacteroides (Figure S3). Overall, about 87% of the microbial families detected in the molecular

controls were shared between the mouth and rectum. This increase in shared taxa was also

expected, as the proportion of cultivable organisms was expected to be lower (57) and the

differences in diversity would not be detected through the selection of cultivability. Because of

the small proportion of molecularly characterized microorganisms that are routinely recovered in

the laboratory, we employed multiple variations in cultivation, like medium type and inoculum

treatments to "cast a wider net."

Treatments selected for organisms that can withstand ethanol stress (e.g., spore-formers),

or were streptomycin resistant. These had a greater effect on the cultivable diversity compared to

changes in base medium alone, likely because the base media shared many of the same

oxidizable substrates (see media recipes in methods section). It was logical to hypothesize that

the inoculum exposed to selective treatments could have decreased cultivated richness compared

to the same inoculum on an untreated medium of the same type, since the selective agent would

inhibit the growth of a portion of the population. Instead, the cultivable richness and diversity

varied for the same treatments on different base medium types and between orifices (Figure 1

and Figure 4). This could be due to the treatments selecting for specific groups of taxa (i.e.

antibiotics) or creating an environment where certain taxa were able to out-compete on non-

selective medium (i.e. low nutrients, increased incubation and slow growers) (17). Specifically,

medium amended with catalase or streptomycin and inocula treated with ethanol resulted in

cultivated organisms not identified on other plates, or even the molecular controls in some cases

(Figure 5). One possible explanation would be that this was due to the inherit stochasticity

associated with cultivation (discussed below). Despite the variability, we were able to cultivate

roughly 57% of the species richness observed with molecular approaches. Previous studies

measuring cultivation success with high-throughput sequencing have reported similar levels of

cultivable richness from the skin of toads (~60%) (26) and rumen fluid (23%) (29). Medina et al.

(2017) reported greater variation in cultivability between inocula (different animals). Our data

showed that more variability in cultivated taxa was observed between selective additions to the

medium or treatments of the inocula (Figure 4), and not the inocula (different orifices) (Figure

3). Similar to Zehavi et al, (2018), different medium types and treatments increased the

cultivable proportion of the original sample.

Bioactivity and phylogenetic diversity of isolates

The research reported here was based on previous success of using roadkill mammals as a

source of antimicrobial-producing microorganisms (33). Our intent was to quantify and broaden

the diversity of cultivated bacteria by the addition of various media and treatments. This affect

was measured by using plate-wash PCR and SSU rRNA gene sequencing. Simultaneously,

colonies were also picked for isolation from replicate plates (n = 3), identified through SSU

rRNA sequencing, and screened for antimicrobial production. Overall, 7 of the 238 recovered

isolates showed bioactivity against the antibiotic-resistant pathogen panel, resulting in a ~3% hit

rate. A total of five bioactive isolates were successfully sequenced and met our quality control

standards described in the methods. Three belonged to the phylum Proteobacteria (class

Gammaproteobacteria) and two belonged to the phylum Firmicutes. Two isolates were identified

as a *Bacillus* sp. and a *Pseudomonas* sp., both genera known to contain species that produce

bioactive compounds (33, 58-61). Interestingly, the isolates identified as an *Enterococcus* sp.

and *Klebsiella* sp. showed inhibitory activity to a member of its own genus, *Enterococcus*

faecium and *Klebsiella pneumonia,* respectively. This finding was consistent with the hypothesis

that closely-related organisms often have a negative effect on each other due to resource overlap

(62), which has been demonstrated for these genera (63). In addition, some populations could be

making antimicrobials, like bacteriocins, that effect close relatives (64).

Cultivation Yields Complex Data

The cultivation and isolation of many organisms at once can be incredibly laborious. New

technologies and approaches can help increase throughput, which is necessary to overcome the

re-isolation of abundant, common, and easy to cultivate organisms. However, it is still important

to be able to assess the efficacy of any cultivation effort. Unfortunately, we uncovered an

intrinsic property of cultivation that is not readily realized unless assessed at a large scale,

variability. Here, we did not detect the same cells (e.g., colonies measured in PWPCR)

consistently growing on replicate plates (Figure S4).

Neither differences in cultivation strategy nor orifice sampled could explain the majority

of the observed variation in community structure (Figure 4). We can attribute unexplained

variance (i.e. residuals) observed between communities (i.e. beta diversity) to stochastic effects

of cultivation, but that is not fundamentally measurable in this case. We can speculate that

randomness would play a role in cultivation as samples are diluted (27, 29) or the spatial

heterogeneity of the inoculum is altered, as demonstrated on a much larger scale using leaf litter

(65). In these instances, we could expect different inocula resulting in different assemblages of

cultivated organisms. We also know that microorganisms immobilized on an agar surface are

still able to interact through motility or diffusion of metabolites, making certain competitive

adaptations more advantageous (64). In this experiment, the agar media allowed physical

separation of individual cells in the inocula and constrained microbial populations to form

colonies, potentially forcing interactions with neighboring colonies that could result in

competition and inhibition of growth through secondary metabolite production (64). The

randomness of which populations are in close proximity to one another could play a role in

which members will thrive from the same inoculum source, affecting the outcome of which taxa

are cultivated (66).

Cultivation on agar plates warrants special consideration when discussing diversity. The

relative distribution of populations on an agar surface affects their prevalence and the overall

composition of the cultivated community. Further, the physiology and colony morphology of

each population dictates their prevalence and that of the surrounding populations on an agar

surface. Colony size, especially surface area, is impacted by traits such as growth rate, surface

motility (i.e., gliding motility, swarming), presence of inhibitory metabolites, and the availability

of resources. These differences in biomass distribution were important when deciding how to

measure diversity, a critical metric for assessing cultivation on an agar surface. Unlike liquid

cultures, varying biomass (i.e. colony size) from different taxa that might produce an equal

number of colonies can over-estimate the relative abundance of that taxon, studies relying on

relative abundance to characterize diversity should consider this (26, 30). If a population (A) has

more proficient growth and forms a larger colony than population (B), DNA extracted from each

population would indicate that taxon A was more abundant in the original inoculum, incorrectly

representing the evenness of the original inoculum. This is especially true when relative

abundance is calculated based on equimolar DNA concentration during library prep. To mitigate

any biased evenness, species richness and Jaccard distances was used to measure alpha and beta

diversity, respectively.

In this data set, about half of the ASVs detected after cultivation were not detected in the

molecular controls. This same phenomenon was observed by Zehavi et al. 2018 when cultivated

rumen OTUs outnumbered the OTUs detected in the original rumen sample and its dilutions

(1,012 out of 1,698) (29). This discrepancy may speak to the power of cultivation relative to the

power of direct molecular analyses in describing the diversity of a community. However, both of

these approaches come with their own caveats. Cultivation makes assumptions about an

organism's ability to grow in the laboratory, while sequencing is dependent on methodology,

sequencing chemistry and sampling depth. In this study, rarefaction curves indicated that

sequencing efforts were sufficient (Figure S5). One potential explanation for the large proportion

of ASVs not detected in the controls could be that in combination with the different media and

treatments, we were able to give some of the less abundant microorganisms a growth advantage.

For example, two families, Wohlfarhtimonodaceae and Metaschinikowiceae, were more

associated with the addition of streptomycin to media (Table 1), and not detected in the

molecular controls (Figure 5). Streptomycin may have selected against the more abundant or

more competitive organisms, allowing taxa from these families to grow.

Cultivation is critical to answer broad challenges of microbial ecology like deciphering

microbial metabolisms or specific challenges like obtaining new isolates for antibiotic discovery.

The study of pure cultures remains the best approach to comprehensively describe an organism's

physiological and metabolic properties, and efforts to optimize cultivation strategy have proven

successful (13, 14, 16, 18, 21, 23, 28, 67-69). Our data adds to previous studies (26, 29) that also

sought to assess how medium type and treatment could increase cultivated diversity, but adds the

goal of "casting a wider net" to increase the diversity of bioactive isolates. We were able to add

to our library of bioactive isolates and show that microbiomes from roadkill mammals are a

useful source of bioactivity, as observed in Motley et al 2017. Lastly, we showed that through a

thoughtful cultivation approach driven by bulk molecular analysis of cultivated taxa, we could

cultivate a larger proportion of the diversity within an inoculum and even taxa not detected in the

molecular controls. This work adds to the growing assessment of cultivation strategies using

newer tools, like high-throughput sequencing, and shows how these methods can be applied to

drug discovery efforts. Cultivation is influenced by many factors, and this work highlights some

of those intricacies, like variability and stochasticity. Cultivation is complex and challenging but

cultivating in combination with molecular tools can expand the observed diversity of an

environment and its community.

Acknowledgments

The authors thank Dr. H. Nunn and the team of Mammalian Microbiome undergraduate

researchers (A. Byrd, J. Dewberry, B. Hansen, C. Hauger, J. Jacobs) for sample processing

during cultivation. The authors also thank Dr. A. K. Dunn for helpful comments during the

writing process. The isolates used for the pathogen panel were kindly provided by Dr. Robert

Cichewicz, Institute for Natural Products and Research Technologies (INPART), University of

Oklahoma.

Tables and Figures

Tables and Figures

Figure 1: Alpha diversity of the mouth and rectum. For the controls, richness, measured using

ASVs, was higher in the rectum than the mouth. For cultivated communities, richness varied

based on medium type and treatment. Points represent each medium or treatment type, indicated

by color (n=3). Molecular 16S/18S refers to the directly sequenced inoculum sample (n=3).

Figure 2: Distribution of ASVs between cultivation and molecular controls. The majority of

ASVs detected in the original inocula were shared between the mouth and rectum, while

approximately half of all ASVs detected through cultivation were not detected in the directly

sequenced samples.

Figure 3: Beta diversity measured in Jaccard distance and NMDS ordination. Differences in

microbial community structure were driven by orifice and treatment, specifically ethanol and

streptomycin. Shape corresponds to orifice sampled, while color differentiates between

treatments of the medium or inoculum.

Figure 4: Percent of variation described by permanova models based on (a) media and treatment

with orifice as a strata, (b) orifice and treatment, and (c) pairwise comparisons of treatments on

media types. Largely, the differences in community structure during cultivation could not be

explained by media, orifice, or treatment alone. In all graphs, residual variation (in grey), is the

proportion of variance not explained by the model.

Figure 5: Heatmaps describing presence absence of taxonomic families observed in the (a)

mouth and (b) rectum. Purple indicates that a family was observed while grey indicates that a

family was absent. The dendrograms highlight clusters of families observed during cultivation.

Table 1. Taxonomic associations with media type using Indicator Species Analysis

Grouping Factor	Condition	Indicator ASV Family	Index Value	p value
Orifice	Mouth	Aeromonadaceae	0.483	0.007
Media Type	ROXY	Aeromonadaceae	0.589	0.001
	Blood + TSA + R2A	Cladosporiaceae	0.604	0.026
Treatment	Ethanol	Staphylococcaceae	0.562	0.014
		Paenibacillaceae	0.408	0.041
	Streptomycin	Wohlfahrtiimonadaceae	0.408	0.034
		Metschnikowiaceae	0.408	0.030
	Ethanol + Streptomycin	Peptostreptococcaceae	0.669	0.003
	Catalase + Ethanol + Streptomycin	Aspergillaceae	0.692	0.004
		Cladosporiaceae	0.633	0.006
	Catalase + Ethanol + Streptomycin + No Treatment	Debaryomycetaceae	0.899	0.002
		Clostridiaceae 1	0.860	0.006

Table 2. Cultivated bioactive isolates

Isolate	SILVA Taxonomy	Orifice	Medium	Activity Against
RC2RCR2A30-212	*Bacillus* sp.	Rectum	R2A	*Klebsiella pneumoniae*
RC2MOR2A30-238	*Pseudomonas* sp.	Mouth	R2A	*Candida albicans*
RC2MORGH30-018	Enterobacteriaceae	Mouth	ROXYgh	*Enterococcus faecium*
RC2MOTSA30-050	*Enterococcus* sp.	Mouth	TSA	*Enterococcus faecium*
RC2MOTSA30-142	*Klebsiella* sp.	Mouth	TSA	*Klebsiella pneumoniae*

588 **Supplementary Material:**

589 Figure S1: Bioassay plate showing three isolates generating a zone of inhibition within pathogen

590 overlay.

592 Figure S2: Pie chart of 117 identified isolates at the genus level.

Table S1. A quasi-factorial approach using targeted media and treatments for gut microorganisms.

Treatment/ Selection agents	Media			
	0.1 TSA	ROXY	0.25 R2A	Blood Agar
None	X	X	X	X
Catalase	X	X	X	
Hemin and alpha-ketoglutarate		X		
70% ethanol	X		X	X
Streptomycin	X		X	X

Figure S3: Phylum-level characterization of molecular controls, triplicates shown for mouth

(RC2M) and rectum (RC2R).

Figure S4: Bar chart describing the variability of consistent recovery of ASVs, data subset. Y

axis indicates how many replicates of TSA plates in which the top 25 ASV were observed. Color

corresponds to orifice.

Figure S5: Rarefaction curves showing plateaued sequencing depth of (a) cultivation samples

and (b) molecular controls.

References:

1. Sogin ML, Morrison HG, Huber JA, Welch DM, Huse SM, Neal PR, Arrieta JM, Herndl GJ. 2006. Microbial diversity in the deep sea and the underexplored ''rare biosphere''. Proceedings of the National Academy of Sciences 103:12115–12120.

2. Donia MS, Cimermancic P, Schulze CJ, Brown LCW, Martin J, Mitreva M, Clardy J, Linington RG, Fischbach MA. 2014. A Systematic Analysis of Biosynthetic Gene Clusters in the Human Microbiome Reveals a Common Family of Antibiotics. Cell 158:1402–1414.

3. Tortorella E, Tedesco P, Palma Esposito F, January G, Fani R, Jaspars M, de Pascale D. 2018. Antibiotics from Deep-Sea Microorganisms: Current Discoveries and Perspectives. Marine Drugs 16:355–16.

4. Halfvarson J, Brislawn CJ, Lamendella R, Vázquez-Baeza Y, Walters WA, Bramer LM, D'Amato M, Bonfiglio F, McDonald D, González A, McClure EE, Dunklebarger MF, Knight R, Jansson JK. 2017. Dynamics of the human gut microbiome in inflammatory bowel disease. Nature Microbiology 2:1–7.

5. Arora-Williams K, Olesen SW, Scandella BP, Delwiche K, Spencer SJ, Myers EM, Abraham S, Sooklal A, Preheim SP. 2018. Dynamics of microbial populations mediating biogeochemical cycling in a freshwater lake. Microbiome 6:165–16.

6. Gilbert JA, Quinn RA, Debelius J, Xu ZZ, Morton J, Garg N, Jansson JK, Dorrestein PC, Knight R. 2016. Microbiome-wide association studies link dynamic microbial consortia to disease. Nature 535:94–103.

7. Singh BK, Bardgett RD, Smith P, Reay DS. 2010. Microorganisms and climate change: terrestrial feedbacks and mitigation options. Nature Reviews 8:779–790.

8. Staley JT, Konopka A. 1985. Meaurement of In Situ Activities of Nonphotosynthetic Microorganisms in Aquatic and Terrestrial Habitats. Annu Rev Microbiol 39:321–346.

9. Lloyd KG, Steen AD, Ladau J, Yin J, Crosby L. 2018. Phylogenetically Novel Uncultured Microbial Cells Dominate Earth Microbiomes. mSystems 3:431–12.

10. Eloe-Fadrosh EA, Ivanova NN, Woyke T, Kyrpides NC. 2016. Metagenomics uncovers gaps in amplicon-based detection of microbial diversity. Nature Microbiology 1:15032–4.

11. Schloss PD, Girard RA, Martin T, Edwards J, Thrash JC. 2016. Status of the Archaeal and Bacterial Census: an Update. mBio 7:995–10.

12. Nichols D, Cahoon N, Trakhtenberg EM, Pham L, Mehta A, Belanger A, Kanigan T,
Lewis K, Epstein SS. 2010. Use of ichip for high-throughput in situ cultivation of
"uncultivable" microbial species. Appl Environ Microbiol 76:2445–2450.

13. Stevenson BS, Eichorst SA, Wertz JT, Schmidt TM, Breznak JA. 2004. New strategies for
cultivation and detection of previously uncultured microbes. Appl Environ Microbiol
70:4748–4755.

14. Connon SA, Giovannoni SJ. 2002. High-Throughput Methods for Culturing
Microorganisms in Very-Low-Nutrient Media Yield Diverse New Marine Isolates. Appl
Environ Microbiol 68:3878–3885.

15. Reasoner DJ, Geldreich EE. 1985. A New Medium for the Enumeration and Subculture of
Bacteria from Potable Water. Appl Environ Microbiol 49:1–7.

16. Bartelme RP, Custer JM, Dupont CL, Espinoza JL, Torralba M, Khalili B, Carini P. 2020.
Influence of Substrate Concentration on the Culturability of Heterotrophic Soil Microbes
Isolated by High-Throughput Dilution-to-Extinction Cultivation. mSphere 5:489–15.

17. Davis KER, Joseph SJ, Janssen PH. 2005. Effects of Growth Medium, Inoculum Size, and
Incubation Time on Culturability and Isolation of Soil Bacteria. Appl Environ Microbiol
71:826–834.

18. Henson MW, Pitre DM, Weckhorst JL, Lanclos VC, Webber AT, Thrash JC. 2016.
Artificial Seawater Media Facilitate Cultivating Members of the Microbial Majority From
the Gulf of Mexico. mSphere 1:e00028–16.

19. Kawasaki K, Kamagata Y. 2017. Phosphate-Catalyzed Hydrogen Peroxide Formation
from Agar, Gellan, and κ-Carrageenan and Recovery of Microbial Cultivability via
Catalase and Pyruvate. Appl Environ Microbiol 83:e01366–17.

20. Kim S, Kang I, Seo J-H, Cho J-C. 2019. Culturing the ubiquitous freshwater
actinobacterial acI lineage by supplying a biochemical "helper" catalase. The ISME
Journal 13:2252–2263.

21. Oberhardt MA, Zarecki R, Gronow S, Lang E, Klenk H-P, Gophna U, Ruppin E. 2015.
Harnessing the landscape of microbial culture media to predict new organism-media
pairings. Nature Communications 6:1–14.

22. Kopke B, Wilms R, Engelen B, Cypionka H, Sass H. 2005. Microbial Diversity in Coastal
Subsurface Sediments: a Cultivation Approach Using Various Electron Acceptors and
Substrate Gradients. Appl Environ Microbiol 71:7819–7830.

23. Browne HP, Forster SC, Anonye BO, Kumar N, Neville BA, Stares MD, Goulding D,
Lawley TD. 2016. Culturing of "unculturable" human microbiota reveals novel taxa and
extensive sporulation. Nature 533:543–546.

24.   La Scola B, Khelaifia S, Lagier JC, Raoult D. 2014. Aerobic culture of anaerobic bacteria
using antioxidants: a preliminary report. Eur J Clin Microbiol Infect Dis 33:1781–1783.

25.   Dione N, Khelaifia S, La Scola B, Lagier JC, Raoult D. 2016. A quasi-universal medium
to break the aerobic/anaerobic bacterial culture dichotomy in clinical microbiology.
Clinical Microbiology and Infection 22:53–58.

26.   Medina D, Walke JB, Gajewski Z, Becker MH, Swartwout MC, Belden LK. 2017. Culture
Media and Individual Hosts Affect the Recovery of Culturable Bacterial Diversity from
Amphibian Skin. Front Microbiol 8:32–15.

27.   Justice NB, Sczesnak A, Hazen TC, Arkin AP. 2017. Environmental Selection, Dispersal,
and Organism Interactions Shape Community Assembly in High-Throughput Enrichment
Culturing. Appl Environ Microbiol 83:e01253–17–16.

28.   Yang S-J, Kang I, Cho J-C. 2015. Expansion of Cultured Bacterial Diversity by Large-
Scale Dilution-to-Extinction Culturing from a Single Seawater Sample. Microb Ecol
71:29–43.

29.   Zehavi T, Probst M, Mizrahi I. 2018. Insights Into Culturomics of the Rumen
Microbiome. Front Microbiol 9:D46–10.

30.   Cummings LA, Hoogestraat DR, Rassoulian-Barrett SL, Rosenthal CA, Salipante SJ,
Cookson BT, Hoffman NG. 2020. Comprehensive evaluation of complex polymicrobial
specimens using next generation sequencing and standard microbiological culture. Sci
Rep 10:1–12.

31.   Hoffmann T, Krug D, Bozkurt N, Duddela S, Jansen R, Garcia R, Gerth K, Steinmetz H,
Muller R. 2018. Correlating chemical diversity with taxonomic distance for discovery of
natural products in myxobacteria. Nature Communications 9:1–10.

32.   Helfrich EJN, Vogel CM, Ueoka R, Schafer M, Ryffel F, Muller DB, Probst S, Kreuzer
M, Piel J, Vorholt JA. 2018. Bipartite interactions, antibiotic production and biosynthetic
potential of the *Arabidopsis* leaf microbiome. Nature Microbiology 3:909–919.

33.   Motley JL, Stamps BW, Mitchell CA, Thompson AT, Cross J, You J, Powell DR,
Stevenson BS, Cichewicz RH. 2017. Opportunistic Sampling of Roadkill as an Entry
Point to Accessing Natural Products Assembled by Bacteria Associated with Non-
anthropoidal Mammalian Microbiomes. J Nat Prod 80:598–608.

34.   van der Meij A, Worsley SF, Hutchings MI, van Wezel GP. 2017. Chemical ecology of
antibiotic production by actinomycetes. FEMS Microbiology Reviews 41:392–416.

35.   Stierle AA, Stierle DB, Kelley K. 2006. Berkelic Acid, A Novel Spiroketal with Selective
Anticancer Activity from an Acid Mine Waste Fungal Extremophile. Journal of Organic
Chemistry 71:5357–5360.

36. Ding Z-G, Li M-G, Zhao J-Y, Ren J, Huang R, Xie M-J, Cui X-L, Zhu H-J, Wen M-L.
2010. Naphthospironone A: An Unprecedented and Highly Functionalized Polycyclic
Metabolite from an Alkaline Mine Waste Extremophile. Chem Eur J 16:3902–3905.

37. Rangseekaew P, Pathom-aree W. 2019. Cave Actinobacteria as Producers of Bioactive
Metabolites. Front Microbiol 10:381–11.

38. Smanski MJ, Schlatter DC, Kinkel LL. 2015. Leveraging ecological theory to guide
natural product discovery. Journal of Industrial Microbiology & Biotechnology 43:115–
128.

39. Schatz A, Bugle E, Waksman SA. 1944. Streptomycin, a substance exhibiting antibiotic
activity against Gram-positive and Gram-negative bacteria. Experimental Biology and
Medicine 55:66–69.

40. Parada AE, Needham DM, Fuhrman JA. 2015. Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. Environ Microbiol 18:1403–1414.

41. Kraus EA, Beeler SR, Mors RA, Floyd JG, GeoBiology 2016, Stamps BW, Nunn HS,
Stevenson BS, Johnson HA, Shapiro RS, Loyd SJ, Spear JR, Corsetti FA. 2018.
Microscale Biosignatures and Abiotic Mineral Authigenesis in Little Hot Creek,
California. Front Microbiol 9:544–13.

42. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer
N, Peña AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley
RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR,
Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010.
QIIME allows analysis of high-throughput community sequencing data. Nat Methods
7:335–336.

43. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.
DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods
13:581–583.

44. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.
2013. The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. Nucleic Acids Research 41:D590–6.

45. Yilmaz, P, Parfrey, LW, Yarza, P, Gerken, J, Pruesse, E, Quast, C, Schweer, T, Peplies, J,
Ludwig, W, Glöckner, FO. 2014. Nucleic Acids Research 42:D643–8.

46. McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS ONE 8:e61217.

47. Dixon P. 2003. VEGAN, a package of R functions for community ecology. Journal of
Vegetation Science 14:927–930.

48. De Cáceres M, Legendre P. 2009. Associations between species and groups of sites: indices and statistical inference. Ecology 90:3566–3574.

49. De Cáceres M, Sol D, Lapiedra O, Legendre P. 2011. A framework for estimating niche metrics using the resemblance between qualitative resources. Oikos 120:1341–1350.

50. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011. Integrative genomics viewer. Nature Biotechnology 29:24–26.

51. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Molecular Biology and Evolution 35:1547–1549.

52. Newman DJ, Cragg GM. 2016. Natural Products as Sources of New Drugs from 1981 to 2014. J Nat Prod 79:629–661.

53. 2019. Antibiotic Resistance and Threats in the United States 2019. Center for Disease Control (CDC).

54. Clardy J, Fischbach MA, Walsh CT. 2006. New antibiotics from bacterial natural products. Nat Biotech 24:1541–1550.

55. Guo J, Fu X, Liao H, Hu Z, Long L, Yan W, Ding Y, Zha L, Guo Y, Yan J, Chang Y, Cai J. 2016. Potential use of bacterial community succession for estimating post-mortem interval as revealed by high-throughput sequencing. Sci Rep 6:1–11.

56. Metcalf JL, Wegener Parfrey L, González A, Lauber CL, Knights D, Ackermann G, Humphrey GC, Gebert MJ, Van Treuren W, Berg-Lyons D, Keepers K, Guo Y, Bullard J, Fierer N, Carter DO, Knight R. 2013. A microbial clock provides an accurate estimate of the postmortem interval in a mouse model system. eLife 2:403–19.

57. Jensen, V. 1968. The plate count technique. In TRG Gray & D Parkinson (Eds.), *The ecology of soil bacteria* (pp. 158–170). Liverpool University Press.

58. Hamdache A, Lamarti A, Aleu J, Collado IG. 2011. Non-peptide Metabolites from the Genus Bacillus. J Nat Prod 74:893–899.

59. Sumi CD, Yang BW, Yeo I-C, Hahm YT. 2015. Antimicrobial peptides of the genus Bacillus: a new era for antibiotics. Can J Microbiol 61:93–103.

60. Raaijmakers JM, Weller DM, Thomashow LS. 1997. Frequency of Antibiotic-Producing *Pseudomonas* spp. in Natural Environments. Appl Environ Microbiol 63:881–887.

61. Fuller AT, Mellows G, Woolford M, Banks GT, Barrow KD, Chain EB. 1971. Pseudomonic Acid: an Antibiotic produced by *Pseudomonas fluorescens*. Nature 234:416–417.

779   62.   Hibbing ME, Fuqua C, Parsek MR, Peterson SB. 2009. Bacterial competition: surviving
780   and thriving in the microbial jungle. Nature Reviews Microbiology 8:15–25.

781   63.   de Vos MGJ, Zagorski M, McNally A, Bollenbach T. 2017. Interaction networks,
782   ecological stability, and collective antibiotic tolerance in polymicrobial infections.
783   Proceedings of the National Academy of Sciences 114:10666–10671.

784   64.   Chao L, Levin BR. 1981. Structured habitats and the evolution of anticompetitor toxins in
785   bacteria. Proceedings of the National Academy of Sciences 78:6324–6328.

786   65.   Albright MBN, Chase AB, Martiny JBH. 2019. Experimental Evidence that Stochasticity
787   Contributes to Bacterial Composition and Functioning in a Decomposer Community.
788   mBio 10:247–13.

789   66.   Traxler MF, Watrous JD, Alexandrov T, Dorrestein PC, Kolter R. 2013. Interspecies
790   Interactions Stimulate Diversification of the *Streptomyces coelicolor* Secreted
791   Metabolome. mBio 4:e00459–13.

792   67.   Marmann A, Aly A, Lin W, Wang B, Proksch P. 2014. Co-Cultivation—A Powerful
793   Emerging Tool for Enhancing the Chemical Diversity of Microorganisms. Marine Drugs
794   12:1043–1065.

795   68.   Lagier J-C, Hugon P, Khelaifia S, Fournier P-E, La Scola B, Raoult D. 2015. The rebirth
796   of culture in microbiology through the example of culturomics to study human gut
797   microbiota. Clin Microbiol Rev 28:237–264.

798   69.   Berdy B, Spoering AL, Ling LL, Epstein SS. 2017. In situ cultivation of previously
799   uncultivable microorganisms using the ichip. Nat Protoc 12:2232–2242.

Table 1. Taxonomic associations with media type using Indicator Species Analysis

Grouping Factor	Condition	Indicator ASV Family	Index Value	p value
Orifice	Mouth	Aeromonadaceae	0.483	0.007
Media Type	ROXY	Aeromonadaceae	0.589	0.001
	Blood + TSA + R2A	Cladosporiaceae	0.604	0.026
Treatment	Ethanol	Staphylococcaceae	0.562	0.014
		Paenibacillaceae	0.408	0.041
	Streptomycin	Wohlfahrtiimonadaceae	0.408	0.034
		Metschnikowiaceae	0.408	0.030
	Ethanol + Streptomycin	Peptostreptococcaceae	0.669	0.003
	Catalase + Ethanol + Streptomycin	Aspergillaceae	0.692	0.004
		Cladosporiaceae	0.633	0.006
	Catalase + Ethanol + Streptomycin + No Treatment	Debaryomycetaceae	0.899	0.002
		Clostridiaceae 1	0.860	0.006

Table 2. Cultivated bioactive isolates

Isolate	SILVA Taxonomy	Orifice	Medium	Activity Against
RC2RCR2A30-212	*Bacillus* sp.	Rectum	R2A	*Klebsiella pneumoniae*
RC2MOR2A30-238	*Pseudomonas* sp.	Mouth	R2A	*Candida albicans*
RC2MORGH30-018	Enterobacteriaceae	Mouth	ROXYgh	*Enterococcus faecium*
RC2MOTSA30-050	*Enterococcus* sp.	Mouth	TSA	*Enterococcus faecium*
RC2MOTSA30-142	*Klebsiella* sp.	Mouth	TSA	*Klebsiella pneumoniae*

(a) **Permanova**

(b) **Permanova**

% Variation Explained

Residual
OrificeTreatment
Treatment
Orifice

~Orifice*treatment, no strata

Pairwise Permanova ~Treatment

(c)

% Variation Explained

Treatment

Residual
Ethanol vs Streptomycin
Base vs Streptomycin
Base vs Ethanol
AKGH vs Streptomycin
AKGH vs Ethanol
AKGH vs Base
Catalase vs Streptomycin
Catalase vs Ethanol
Catalase vs Base
Catalase vs AKGH