Spatio-temporal Control of ERK Pulse Frequency Coordinates Fate Decisions

during Mammary Acinar Morphogenesis

Pascal Ender [1,2], Paolo Armando Gagliardi [1], Maciej Dobrzyński [1], Agne Frismantiene

[1], Coralie Dessauges [1], Thomas Höhener [1], Marc-Antoine Jacques [1,3], Andrew R.

Cohen [4], and Olivier Pertz [1]*

1 Institute of Cell Biology, University of Bern, Baltzerstrasse 4, 3012 Bern, Switzerland

2 present address: Department of Autonomous Matter, AMOLF, Science Park 104,

1098 XG Amsterdam, The Netherlands

3 present addresses: Cancer Research UK Cambridge Institute, University of

Cambridge, Li Ka Shing Centre, Robinson Way, Cambridge, CB2 0RE, United

Kingdom; and, Wellcome Genome Campus, Hinxton, Cambridgeshire, CB10 1SD,

United Kingdom

4 Department of Electrical and Computer Engineering, Drexel University, 3120-40

Market Street, Suite 313, Philadelphia, PA 19104, USA

*Correspondence and requests for materials should be addressed to O.P. (email:

olivier.pertz@unibe.ch).

Abstract

The signaling events controlling proliferation, survival, and apoptosis during mammary epithelial acinar morphogenesis remain poorly characterized. By imaging single-cell ERK activity dynamics in MCF10A acini, we find that these fates depend on the average frequency of non-periodic ERK pulses. High pulse frequency is observed during initial acinus growth, correlating with rapid cell motility. Subsequent decrease in motility correlates with lower ERK pulse frequency and quiescence. Later, during lumen formation, coordinated ERK waves emerge across multiple cells of an acinus, correlating with high and low ERK pulse frequency in outer surviving and inner dying cells respectively. Optogenetic entrainment of ERK pulses causally connects high ERK pulse frequency with inner cell survival. Acini harboring the PIK3CA H1047R mutation, commonly observed in breast cancer, display increased ERK pulse frequency, inner cell survival and loss of lumen formation. Thus, fate decisions during acinar morphogenesis are fine-tuned by different spatio-temporal coordination modalities of ERK pulse frequency.

Introduction

Mammary organogenesis involves formation of a rudimentary gland during embryogenesis, followed by proliferation and branching invasion led by multi-layered terminal end buds (TEBs) during puberty. Cells in the inner TEB layers then undergo apoptosis to form the ductal lumen. During pregnancy, secretory alveoli are then formed at the ends of the ductal tree (Inman et al., 2015; Paine and Lewis, 2017). Morphogenesis of this complex structure requires spatial and temporal control of cell fates such as proliferation, survival, migration and death. However, the spatio-temporal signaling events that regulate such fate decisions remain poorly explored. The epidermal growth factor (EGF) receptor (EGFR) – mitogen activated protein kinase (MAPK) signaling cascade, that ultimately leads to activation of the extracellular regulated kinase (ERK) is a key pathway involved in mammary gland development. EGFR-ERK signaling results in the upregulation of gene products involved in a wide variety of processes such as proliferation, survival, migration and differentiation (Lavoie et al., 2020). EGFR activity is required for mammary gland morphogenesis in mice (Sebastian et al., 1998). Paracrine amphiregulin release by the matrix metallo-protease (MMP) ADAM-17 and its binding to EGFR mediates the effects of estrogen receptor α to promote mammary gland development and growth (Ciarloni et al., 2007; Sternlicht et al., 2005). EGFR-dependent ERK activity is enriched at the front of elongating tubes and coordinates cell migration (Huebner et al., 2016). In 3D mammary acini, oncogenic ERK activation suppresses apoptosis and thus lumen formation (Reginato et al., 2005). In primary mammary cell culture, ERK activity is also crucial for survival (Finlay et al., 2000).

Recent single-cell measurements of ERK activity dynamics in a variety of 2D epithelial monolayer cultures have revealed the existence of non-periodic ERK pulses of constant amplitude and duration (Aikin et al., 2020; Albeck et al., 2013; Aoki et al., 2013; Gagliardi et al., 2021; Goglia et al., 2020; Hino et al., 2020; Hiratsuka et al., 2015). These ERK pulses originate from MAPK network properties such as ultrasensitivity (leading to steep ERK activation at a threshold EGFR input), and negative feedback (leading to ERK adaptation) (Kholodenko et al., 2010; Sparta et al., 2015). An emerging theme is that the average frequency of these non-periodic ERK pulses, referred to as ERK frequency from now on, control apoptosis (low frequency), survival (medium frequency) or proliferation (high frequency) (Albeck et al., 2013; Aoki et al., 2013; Gagliardi et al., 2021). These ERK pulses can either exhibit a stochastic

spatially uncorrelated behavior (Albeck et al., 2013; Goglia et al., 2020), or can be organized as wave patterns that regulate collective cell migration in a wound (Aoki et al., 2017; Hino et al., 2020; Hiratsuka et al., 2015), cell survival around sites of apoptotic cell extrusion (Gagliardi et al., 2021; Valon et al., 2021), or extrusion of cancer cells (Aikin et al., 2020). These ERK signaling patterns consist of trigger waves that involve sequential activation of ERK pulses in adjacent epithelial cells through paracrine signaling involving MMP-mediated cleavage of pro-EGF ligands.

In this work, we explore single cell ERK dynamics in 3D mammary acini. Culturing mammary MCF10A cells in a basement membrane matrix (Matrigel) produces acini that retain key features of in vivo breast alveoli. Based on previous work (Anderson et al., 2010; Debnath et al., 2003), we subdivided this process into four stages as depicted in Figure S1A. Stage 1 is characterized by high proliferation rates and basement membrane deposition; stage 2 consists of a quiescent state and presence of an outer cell layer with clear basolateral polarity; stage 3 consists of apoptosis of inner cells that allows formation of a hollow lumen in stage 4. The whole process takes approximately two weeks.

We document different spatio-temporal modalities of ERK signaling during different developmental stages of MCF10A acini formation. Stage 1 is characterized by high ERK frequency, robust proliferation and rapid collective motility. This is followed by a transition to lower ERK frequency and slower collective motility. During stage 2, formation of ERK wave patterns correlates with domains of different ERK frequencies: outer/inner cells display medium/low ERK frequencies, respectively controlling survival and apoptosis. Optogenetic control of ERK signaling shows that ERK pulses control collective migration during stage 1, and that a critical ERK frequency is necessary for survival during stage 2. We characterize a crosstalk between phospho-inositide-3 kinase (PI3K) and MAPK/ERK signaling that feeds into the regulation of ERK frequency. This provides insight into how oncogenic PI3K signaling crosstalks with ERK to contribute to loss of lumen formation. Our work reveals how spatio-temporal control of ERK frequency organizes mammary acinar morphogenesis.

Results

Stage 1 proliferative acini exhibit non-periodic MMP/EGFR-dependent ERK pulses whose frequency correlates with collective cell migration speed.

To explore single-cell ERK dynamics during acinar morphogenesis, we created MCF10A reporter lines expressing the nuclear marker histone 2B (H2B) fused to miRFP703 with either the ERK activity biosensor ERK-KTR fused to mTurquoise2, or a mCherry-geminin S/G2/M cell cycle marker (Sakaue-Sawano et al., 2017). ERK-KTR reports on ERK activity through reversible nucleus/cytosol shuttling after its phosphorylation by active ERK (Regot et al., 2014) (Figure 1A). These lines were then used to grow acini according to a modified version of a previously described protocol (Debnath et al., 2003). After three days, in which EGF, serum and insulin were required for initial acinar growth, these growth factors (GFs) were removed to study ERK signaling dynamics intrinsic to acinar morphogenesis. Using the geminin marker and a fluorogenic caspase substrate, we evaluated if our protocol recapitulated the proliferation, quiescence and apoptosis fates previously documented during acinar morphogenesis (Figure S1B-E) (Debnath et al., 2002; Liu et al., 2012). Stage 1 acini (4 days post-seeding) displayed elevated levels of proliferation and absence of apoptosis. Stage 2 and 3 acini (7 and 11 days post-seeding) revealed quiescent cells and increased apoptosis leading to lumen formation. Stage 4 acini (14 days post-seeding) exhibited a mature lumen, abundant apoptotic debris, and a small increase in proliferation.

We then imaged single-cell ERK dynamics in stage 1 acini using confocal spinning disk microscopy of both the H2B and ERK-KTR channels with time resolutions of 3 - 5 minutes, until acini started to suffer from phototoxicity (observed after 10 - 23 hours). Cells in stage 1 acini displayed asynchronous, non-periodic ERK pulses (Figure 1B) as observed in 2D culture (Aikin et al., 2020; Albeck et al., 2013; Gagliardi et al., 2021). To extract single-cell ERK activity trajectories, we used a customized version of the open-source LEVERJS software (Wait et al., 2014; Winter et al., 2016) that segments and tracks nuclei based on H2B signal; and calculates ERK activity as a ratio of ERK-KTR fluorescence intensities in cytosolic and nuclear voxel masks (Figure 1C). Detrending of the ERK trajectories and normalization of the values to [0,1] generated a reliable input for automated detection of ERK pulses (Figure 1C). Single cell ERK trajectories revealed spontaneous ERK pulses with slightly different amplitudes (Figure 1D). Trametinib-mediated MEK, gefitinib-mediated EGFR, as well as

Batimastat-mediated MMP inhibition abolished ERK pulses (Figure 1E). MEK or EGFR

inhibition for multiple days led to massive cell death and disintegration of the acini

(Figure S2A), suggesting that ERK provides a pro-survival signal. Acute stimulation

with 1 ng/ml EGF induced ERK pulses of amplitudes similar to those of spontaneous

pulses, while 100 ng/ml EGF induced ERK pulses of higher amplitudes (Figure 1F,G).

These results document spontaneous, asynchronous EGFR- and MMP-depedent

ERK pulses in stage 1 acini.

As previously described (Wang et al., 2013), stage 1 acini displayed collective cell

motility correlating with a rotational movement of 360 degrees of whole acini over

multiple hours. Later during stage 1, we observed a transition to a state of slower

motility (Figure 2A, Movie S1). This was shown to correlate with deposition of

basement membrane around the acinus (Wang et al., 2013). The reduction in

migration speed correlated with decreased ERK frequency (Figure 2B-D and S3A,

Movie S2), without having a significant effect on ERK pulse amplitude and duration

(Figure S3B,C).

Figure 1 **Spontaneous EGFR/MMP-dependent ERK pulses in acini.** (A) Schematic representation of the ERK activity - dependent subcellular localization of ERK-KTR. (B) Time-series of the equatorial optical section of an acinus (red plain line) expressing fluorescent H2B and ERK-KTR. Highlighted is a cell (green dotted line) that displays spontaneous ERK activity pulses (arrowheads) resulting in the nuclear to cytoplasmic translocation of ERK-KTR. Scale bar = 10 µm. (C) Image analysis pipeline to extract ERK trajectories from 3D time lapse datasets. Nuclei are segmented and tracked by LEVERJS based on the H2B signal. Single-cell ERK activity levels are the ratio of the median ERK-KTR signal pixel intensities in the voxel mask around the nucleus and the one of the segmented nuclear volume. ERK pulses are detected on detrended ERK trajectories normalized to [0,1]. (D) Representative single-cell ERK trajectories from one acinus. (E) Overlayed ERK trajectories from control and drug-treated acini. Control trajectories correspond to the same acinus as in (D). (F) ERK trajectories from acini treated with 1 or 100 ng/ml EGF at the indicated time points. (G) ERK pulse amplitudes in cells from control and EGF-treated acini. Wilcoxon tests (n.s., P > 0.05; ***, P < 0.001).

Figure 2 **Collective cell migration and ERK pulsing in stage 1 acini** (A) Time series renderings of the cross section of an acinus transitioning from the rapid motility to the slow motility stage. Nuclei and motility tracks color-coded by instantaneous velocity are shown. Scale bar = 10 μm. (B) Analysis of ERK activity in the acinus from (A). Heatmap shows detrended and normalized single cell ERK activity levels over time. Gray areas correspond to time points when a cell was not within the imaged volume. Asterisks indicate individual ERK pulses. (C) Analysis of motility and ERK activity in the acinus from (A). Graph shows mean binarized ERK activity and mean instantaneous velocity with 95% confidence intervals of all imaged cells over time and their Pearson correlation coefficient. Mean binarized ERK activity is used as a measure for the fraction of the cell population in a state of active ERK. (D) ERK pulse frequency from trajectories at different developmental timepoints. Trajectories pooled from 7 (stage 1 rotation), 5 (stage 1 no rotation) and 11 (stage 2) acini. Wilcoxon tests (***, P < 0.001).

Stage 2 quiescent acini exhibit different ERK frequencies in inner and outer cells, which emerges from collective waves of ERK pulses.

We then evaluated ERK dynamics in larger, stage 2 quiescent acini that are characterized by an outer layer of polarized cells, and a less organized inner cell mass destined for apoptosis for future lumen formation (Debnath et al., 2002). When comparing low-motility stage 1 and stage 2 acini, we observed a further reduction in median ERK frequency (Figure 2D), while ERK pulse amplitudes and durations remained almost identical (Figure S3B,C). This change in ERK frequency resulted from a bimodal distribution in which a part of the cell population did not display any ERK pulses (Figure 2D). Because ERK pulse frequency can regulate proliferation, survival and apoptosis fates in MCF10A cells (Aikin et al., 2020; Albeck et al., 2013; Gagliardi et al., 2021), and because inner cells in stage 2 acini are destined to undergo apoptosis, we evaluated ERK pulse frequencies in inner versus outer cells. Outer cells exhibited a significantly higher ERK frequency than inner cells, with the latter often not exhibiting ERK pulses at all (Figure 3A,B). Similar ERK activity amplitude and duration were found in inner/outer cells (Figure S3D,E). Together with our characterization of fate decisions (Figure S1), these results suggest a spatio-temporal mechanism that controls survival versus apoptosis fates through regulation of ERK frequency.

A striking feature of stage 2 acini was that they exhibited spatially correlated ERK pulses in the form of waves spreading across multiple cells (Figure 3B,C, Movie S3). We devised computational methods (Figure S3F-H, Materials and Methods) to detect, track, and extract features that describe ERK waves. These ERK waves were observed in all of the stage 2 acini that we imaged (N=11), and exhibited different geometries (Figure 3D). While some ERK pulses remained restricted to single cells, most of the ERK pulses occurred within collective waves (Figure 3E). ERK waves typically involved a median of 6 cells for a median duration of 54 minutes (Figure 3F,G). However, a large variance was observed with some ERK waves involving as little as 2 and as many as 39 cells (almost the whole acinus). ERK waves, as well as isolated ERK pulses, were predominantly initiated in the outer cell layer (Figure 3H). Further, ERK waves that originated in the outer layer displayed a higher bias to remain at that location than those originating in inner cells (Figure 3I). We have previously shown that in 2D MCF10A cultures, as well as in acini, apoptotic cells trigger ERK waves in their neighboring cells (Gagliardi et al., 2021). However, the ERK waves we

observed here only rarely coincided with apoptotic events, suggesting that they
originate through a different mechanism. Our results strongly suggest that ERK waves
contribute to spatially position different ERK pulse frequencies in inner and outer
acinar cells.

Figure 3 **Different ERK pulse frequencies in inner and outer acinar cell layers in stage**
2, and collective waves of ERK pulses. (A) ERK pulse frequency from trajectories of cells
located in inner versus outer acini layers. Trajectories pooled from 11 acini. Wilcoxon test (***,
P < 0.001). (B) Left: heatmap of detrended/normalized single-cell ERK trajectories in outer
and inner cells of a representative acinus. Right: detection of individual ERK activity waves in
the same acinus. (C) Representative time-series micrographs of ERK wave ID 10 in (B). Nuclei
color-coded by ERK-KTR ratios. Arrows depict wave directionality. The arrowhead indicates
the initiator cell. Dashed line indicates the acinus border. Scale bar = 10 μm. (D) 2D projection
representations of isolated ERK waves from four different acini. Cells that participate in the
wave are color-coded by their relative time of activation. Size represents the relative Z position
of the cell and shape if they belong to the inner or outer cell population. Top and bottom panels

depict two isolated ERK waves for each acinus. (E) Percentage of initiator events that remain
restricted to a single cell vs those that lead to collective events. (F) Total numbers of unique
cells involved in individual collective events. (G) Durations of individual collective events. (H)
Average frequency of single and collective events per acinus, normalized by the number of
cells in the acinus. (I) Penetration of collective events across acini. Calculated as the time-
averaged fraction of localization of a collective event between the inner (0) and outer (1) cell
layer.

Optogenetic control of ERK frequency regulates collective motility, survival and apoptosis fates

To explore the role of ERK pulse frequency during stage 1 collective motility, as well as stage 2 apoptosis and survival fates required for lumen morphogenesis, we used two optogenetic actuators to evoke different ERK pulse frequencies in acini (Figure 4A). Optogenetic fibroblast growth factor receptor (optoFGFR) is a Cry2-based light-activatable receptor tyrosine kinase that activates ERK, Akt and calcium signaling (Kim et al., 2014). OptoRaf is a CIBN/Cry2-based system in which a catalytic Raf domain is recruited to a plasma membrane targeted anchor in a light-dependent fashion to specifically activate ERK (Aoki et al., 2017). We generated stable lines expressing any of the two optogenetic constructs, a spectrally compatible ERK-KTR-mRuby, and H2B-miRFP. Application of a single blue light pulse evoked a discrete optoFGFR- or optoRaf-mediated ERK pulse with similar shape and duration as spontaneous ones (Figure 4B).

Since ERK pulse wave patterns can coordinate collective cell migration through regulation of myosin activity (Aoki et al., 2017; Hino et al., 2020), we hypothesized that the asynchronous ERK pulses we observed might coordinate the collective motility pattern in stage 1 acini. We therefore sought to disrupt this process by synchronizing ERK pulses across all the cells of an acinus. We imaged rotating stage 1 acini for 4 hours in the absence of blue light, and observed asynchronous ERK pulses (Figure 4C). We then applied high frequency light pulses at 30 minute intervals, which synchronized ERK pulses across cells (Figure 4C, Movie S4). This switch to synchronous high-frequency ERK pulsing in all cells immediately resulted in decreased cell migration speed and persistence (Figure 4D-F). This suggest that the asynchronous high frequency ERK pulses organize collective cell migration in stage 1 acini.

Next, we tested the hypothesis that the different ERK frequencies observed between outer and inner cells in stage 2 acini regulate survival vs apoptosis cell fates. Using optoFGFR and optoRaf, we evoked frequency-modulated, population-synchronous ERK dynamics in all the cells of an acinus (Figure 5A, Movie S5). Endogenous collective ERK pulses were however still occurring. Because the regulation of apoptosis/survival fates required for lumen formation spans over one week, we could not use our live cell imaging platform to study this process. We therefore used LITOS (LED Illumination Tool for Optogenetic Stimulation) (Höhener et al., 2022) to evoke

frequency-modulated ERK pulse regimes in multiwell plates in a tissue culture
incubator (Figure S4A). This system evoked similar ERK pulses as observed in our
live cell imaging system (Figure S4B). We then stimulated stage 2 acini with light
pulses delivered at different frequencies for 7 days and scored the distribution of acini
that exhibited filled lumen, partially cleared lumen or cleared lumen (Figure 5B). Using
both optoFGFR and optoRaf, we observed that ERK pulses induced every 0.5, 1, 2,
3, 4 but not 10 hours led to survival of inner cells, increasing the number of acini with
filled or partially cleared lumina (Figure 5C). These results further suggest that survival
and apoptosis fates are regulated by a frequency encoded ERK signal. The optoRaf
experiments indicate that high frequency ERK pulsing alone is sufficient to induce
survival independently of PI3K/Akt signaling.
Because endogenous ERK pulses occur on top of optogenetically-induced ones, we
used optoFGFR to evoke different ERK pulse frequencies in EGFR-inhibited acini that
do not exhibit spontaneous ERK pulses. Here, unstimulated acini and those stimulated
every 10 hours displayed cell death. In marked contrast, ERK pulses applied at 0.5, 1,
2, 3 or 4 hours led to cell survival (Figure 5D). These results show that cells in acini
must experience at least one ERK pulse every 4 hours to survive.

Figure 4 **Optogenetic stimulation of acini controls 3D migration properties in stage 1 acini** (A) Cartoon of the optoFGFR and optoRAF systems. OptoFGFR consists of the intracellular domain of FGFR1 linked to the plasma membrane and a Cry2 PHR domain which dimerizes upon blue light stimulation, leading to receptor autophosphorylation and activation of downstream cascades. OptoRAF consists of a Cry2 linked to cRaf and a membrane-linked CIBN domain. CIBN and Cry2 dimerize upon blue light stimulation which recruits cRaf to the plasma membrane where it phosphorylates MEK. (B) Average ERK trajectories from isolated spontaneous and optogenetically induced ERK pulses with 95% confidence interval. Time = 0 corresponds to maximal amplitude of peaks. (C) Overlayed detrended ERK activity trajectories from a stage 1 rotating acinus expressing optoFGFR and stimulated every 30 min starting from 4 hours. Vertical blue lines indicate the blue light stimulation. (D) Six single-cell migration trajectories from the same example organoid color coded according to migration speed. Spheres represent the nuclei positions at the end of the trajectory. The micrographs were taken at 4 and 8 hours of the experiment, each one with the migration trajectories of the past 4 hours. Dashed line indicates the acinar border. (E) Speed and (F) persistence of single-cell migration in the same acinus. Population average and 90% confidence interval are shown.

Figure 5 **Controlling survival/apoptosis decisions with optogenetic actuators.** (A) Overlayed detrended ERK activity trajectories from optoFGFR or optoRAF expressing acinar cells stimulated with different blue light pulse frequencies. (B) Colored: Equatorial optical sections of acini displaying cleared, partially cleared or filled luminal space or underwent complete cell death at stage 4. Black and white: maximal intensity projections of equatorial Z planes spanning 12 µm. Acini borders (red lines), luminal space (dashed red lines), and the border between the cleared and filled part of the luminal space (dashed green lines) are indicated. Scale bar = 20 µm. (C) Percentages of acini displaying cleared, partially cleared or filled luminal space at day 14, after 7 days on an LED plate that emitted blue light pulses at defined intervals. N = 36 - 72 acini per condition from 2 independent replicates. Pearson's chi-squared test. optoFGFR: X^2 (12 degrees of freedom, N = 384) = 78, P < 0.001. optoRAF: X^2 (12 degrees of freedom, N = 326) = 32, P < 0.005 (D) Percentages of acini displaying cleared,

partially cleared or filled luminal space or complete cell death at day 14, after 7 days in the
presence of gefitinib on an LED plate that emitted blue light pulses at defined intervals. N =
24 - 46 acini per condition. X^2 (15 degrees of freedom, N = 201) = 79, P < 0.001. (E) MCF10A
cells in monolayer culture were stimulated every 3 hours with blue light pulses or every 9 hours
with the same AUC, achieved with 20 consecutive blue light pulses, and compared with
unstimulated cells. Randomly selected trajectories (left) and whole population average (right).
(F) Distribution of apoptotic rates in 5 biological replicates each one normalized on the
experiment mean. t-test (**, P < 0.01; ***, P < 0.001). (G) Percentages of optoFGFR-
expressing acini that displayed cleared, partially cleared or filled luminal space at day 14, after
7 days on an LED plate that emitted either a single blue light pulse every 3 hours, 20
subsequent blue light pulses every 9 hours or no blue light pulses. N = 27-39 acini per
condition. X^2 (4 degrees of freedom, N = 104) = 23, P < .001.

ERK frequency but not integrated ERK activity regulates the survival fate Next,
we explored if the frequency of ERK pulses or the total integrated ERK activity over
time controls survival. To test this, we used optoFGFR to induce synthetic ERK
pulses of different widths, that when applied at different frequencies can evoke the
same integrated ERK activity over a specific time period. Since ERK pulses display an
identical shape and promote survival for 4 hours both in serum-deprived 2D
monolayers (Gagliardi et al., 2021) and 3D acini, we first identified light stimulation
schemes capable of inducing ERK pulses of different widths in monolayers. We
applied a series of 1 to 86 successive blue light pulses delivered at 2-minute intervals,
and recorded the resulting ERK dynamics (Figure S4C). We have previously shown
that light stimulation applied at this frequency leads to sustained optoFGFR activity
(Dessauges et al., 2021). The integrated ERK activity (area under the curve = AUC)
displayed a linear relationship with the number of light pulses (Figure S4D). 20 blue
light pulses delivered every 2 minutes led to a single ERK pulse with a three-fold
increase in integrated ERK activity. We then applied 2 distinct optoFGFR stimulation
schemes consisting of 1 AUC equivalent of ERK activity evoked every 3 hours versus
3 AUC equivalents of ERK activity evoked every 9 hours, resulting in the identical
integrated ERK activity over a period of 9 hours (Figure 5E). We found that the first
but not second optogenetic stimulation scheme induced cell survival in serum-
deprived 2D cultures (Figure 5F). We then performed the identical experiment in stage
2 acini by applying the two optogenetic stimulation schemes using LITOS for 7 days.
Quantification of lumen formation efficiency at day 14 revealed increased inner cell
survival when the first, but not the second, optogenetic stimulation scheme was
applied (Figure 5G). These results show that the frequency of ERK pulses rather than
the integrated ERK activity regulates survival.

Finally, we also used our optogenetic toolkit to explore a number of scenarios of how
different ERK frequencies in inner/outer cells might be regulated in stage 2 acini.
These results are discussed in the Supplemental Text and Figure S5.

**Oncogenic PI3K signaling increases ERK frequency leading to loss of acinar
lumen formation**

We then explored ERK dynamics in the context of a pathological alteration of acinar
morphogenesis induced by the H1047R mutation in the alpha subunit of PI3K
(PIK3CA), which is frequently mutated in breast cancer (Cancer Genome Atlas

Network, 2012). The H1047R PIK3CA mutation leads to absence of lumen in MCF10A
knockin acini (Berglund et al., 2013; Chakrabarty et al., 2010; Chen et al., 2013, Isakoff
et al., 2005; Lauring et al., 2010), as well as ductal hyperplasia in a transgenic mouse
model (Tikoo et al., 2012). H1047R PIK3CA MCF10A knock-in cells have been shown
to display elevated ERK activity using western blot (Gustin et al., 2009), strongly
suggesting the existence of a crosstalk between oncogenic PI3K and MAPK/ERK
signaling. Consistently, we found that H1047R PIK3CA MCF10A knock-in cells
cultured as monolayers displayed higher median ERK frequency than their wild-type
(WT) counterparts (Figure 6A-C), while maintaining similar ERK pulse shape, duration
and amplitude (Figure 6D, S6A,B). The finding of a similar ERK pulse shape in WT
and mutant cells indicates that the PI3K to MAPK crosstalk must occur upstream of
Ras because the MAPK network structure that shapes ERK dynamics is maintained
(Kholodenko et al., 2010). The increased ERK frequency suggests the involvement of
receptor   level   interactions   (Sparta   et   al.,   2015).
To better understand the effect of the PIK3CA H1047R on acinar morphogenesis, we
evaluated proliferation and apoptosis during different stages (Figure 6E-I). Stage 1
mutant acini displayed increased proliferation compared to their WT counterparts, as
evidenced by augmented cell numbers, and geminin quantification (Figure 6E-G).
While remaining slightly higher than in WT acini, proliferation also diminished during
stages 2 - 4, but displayed a small upshoot during stage 4 (Figure 6E,G). In contrast
to the steep apoptosis rise observed starting on day 7 in WT acini, apoptotic rates
remained lower in PIK3CA mutant acini at all stages (Figure 6E,H). Increased
proliferation at stage 1, and decreased apoptosis at stages 3 and 4, thus mostly
contribute to increased cell number and absence of lumen formation in PIK3CA-
mutant acini (Figure 6I).
We then evaluated ERK dynamics during the pathological acinar morphogenesis
induced by PIK3CA H1047R. As in monolayers, PIK3CA-mutant acini displayed ERK
pulses like those of WT acini (Figure 7A). Stage 1 rotating PIK3CA-mutant acini
displayed ERK frequencies as high as those observed in their WT counterparts (Figure
7B). However, stage 1 non-rotating PIK3CA-mutant acini did not display the decreased
ERK frequency observed in WT (Figure 7B). Stage 2 PIK3CA-mutant acini displayed
increased ERK frequency compared to WT, as well as prominent ERK waves (Figure
7B-D, Movie S6). However, most likely due to their heterogeneity, we could not
pinpoint a specific feature of ERK waves associated with the increased ERK frequency

observed in PIK3CA-mutant versus WT acini. PIK3CA-mutant stage 2 displayed
higher ERK frequencies than WT acini, leading the inner cells of PIK3CA-mutant to
exhibit a similar ERK frequency than the outer cells of WT acini (Figure 7B,C).
Amplitude and duration of ERK pulses were similar for both WT and PIK3CA H1047R
acinar cells at all stages (Figure S6C,D).

Pictilisib-mediated PI3K or AZD5363-mediated Akt inhibition decreased ERK
frequency in both WT and mutant acini (Figure 7E). Stimulation of WT acini with
insulin-like growth factor (IGF1), that primarily activates PI3K-Akt signaling (Myers et
al., 1993), also resulted in increased ERK pulse frequency (Figure S7A,B). Gefitinib-
mediated EGFR inhibition abolished ERK pulses in PIK3CA-mutant acini, suggesting
an EGFR-dependent mechanism (Figure S7C). Batimastat-mediated MMP inhibition
in PIK3CA H1047R cells led to reduction of ERK phosphorylation to levels
observed in WT cells, without affecting Akt signaling (Figure S7D). PIK3CA
H1047R MCF10A cells have been shown to exhibit increased expression of
the EGFR-ligand amphiregulin in comparison to WT cells (Young et al., 2015),
possibly explaining the increase in EGFR-dependent ERK frequency. Consistently,
we observed increased amphiregulin expression at mRNA level in PI3K-mutant when
compared with WT cells. Amphiregulin expression levels were were decreased
upon pictilisib-mediated inhibition of PI3K activity in PIK3CA mutant cells (Figure
S7E). Together, these results strongly suggest that the PI3K to MAPK/ERK crosstalk
that regulates ERK frequency involves amphiregulin/MMP-activation of EGFR.
Increased survival in PIK3CA H1047R acini might therefore involve increased
ERK frequency through this crosstalk mechanism, in addition to PI3K-Akt signaling.

Figure 6 **Increased 2D monolayer ERK frequency and altered acinar morphogenesis of PIK3CA H1047R cells.** (A) Micrographs of WT and PIK3CA H1047R MCF10A 2D monolayers expressing fluorescent H2B (left) and ERK-KTR (middle). Right: nuclei of the same cells color-coded by ERK-KTR ratio. Scale bar = 100 μm. (B) Heatmap of single-cell ERK trajectories in WT and PIK3CA H1047R monolayers. (C) ERK frequencies in WT and PIK3CA H1047R monolayer cells. (D) Average ERK trajectories from isolated pulses in WT and PIK3CA H1047R cells within monolayers. 95% confidence intervals are shown. Time = 0 corresponds to maximal amplitude of peaks. (E) Micrographs and 3D reconstructions of H2B, caspase 3/7 fluorogenic substrate and geminin signals in PIK3CA H1047R acini at different stages. Micrographs show maximal intensity projections of equatorial Z planes spanning 12 μm. Plain lines mark the borders of the acini, dashed lines mark the outer cell layer. Scale bar = 10 μm. (F) Cell numbers per acinus at the different stages. N = 54 - 60 PIK3CA H1047R acini each. (G) Fraction of Geminin positive cells per acinus at different stages. (H) Number of Caspase

3/7 apoptotic debris divided by the acinar cell number at different days. (I) Percentages of
acini that either displayed a cleared, partially cleared or filled luminal space at day 14.
Pearson's chi-squared test: X^2 (2 degrees of freedom, N = 82) = 50, P < 0.001. Representative
examples used for classification are shown (maximal intensity projections of equatorial Z
planes spanning 12 µm). Acini borders (red lines), luminal space (dashed red lines), and the
border between the cleared and filled part of the luminal space (dashed green lines) are
indicated. Scale bar = 10 µm. (F-I) Measurements taken on the same acini. WT acini are the
same as in figure S1, for comparison. (G, H) Error bars represent standard error of the mean.
(C, F, G and H) Wilcoxon tests (n.s., P > 0.05; *, P < 0.05; **, P < 0.01; ***, P < 0.001).

Figure 7 **Increased ERK frequency in PIK3CA H1047R mutant acini.** (A) Average ERK trajectories from isolated pulses in WT and PIK3CA H1047R cells within acini. 95% confidence intervals are shown. Time = 0 corresponds to maximal amplitude of peaks. (B) ERK frequencies of WT and PIK3CA H1047R cells at different stages and locations within the acinus. Mutant trajectories pooled from 7 (stage 1 rotation), 2 (stage 1 no rotation) and 6 (stage 2) acini. (C) Heatmap of detrended/normalized single-cell ERK trajectories in outer and inner cells of a representative stage 2 PIK3CA H1047R acinus. (D) Representative time-series of an ERK wave in a stage 2 PIK3CA H1047R acinus (dashed line) cross section. Nuclei are color-coded by ERK activity levels. Arrows show directionality of activation. Arrowhead indicates the initiator cell. Scale bar = 10 µm. (E) ERK frequencies of WT and PIK3CA H1047R cells from control acini and acini treated with 250 nM pictilisib or 5 µM AZD5363. (B and E) wt control data is the same as in figure 2 and 3, and shown again for comparison. Wilcoxon tests (n.s., P > 0.05; *, P < 0.05, **, P < 0.01; ***, P < 0.001).

Discussion

Recent work in epithelial monolayers have revealed the existence of non-periodic single-cell ERK pulses, whose frequency controls apoptosis, survival or proliferation fates (Albeck et al., 2013; Aoki et al., 2013, Gagliardi et al., 2021; Valon et al., 2021). At the level of a cell population, these ERK pulses can be stochastic when cells are stimulated with EGF (Albeck et al., 2013), or can be organized as collective ERK waves during collective cell migration (Aoki et al., 2017; Hino et al., 2020), cancer cell extrusion (Aikin et al., 2020), or spatial regulation of survival in response to stress (Gagliardi et al., 2021; Valon et al., 2021). Here we show that similar ERK pulses/waves coordinate fate decisions during mammary acinar morphogenesis. As in the cell systems mentioned above, ERK pulses are triggered by MMP-mediated cleavage of pro-EGF ligands and subsequent activation of EGFR. Downstream of EGFR, MAPK network properties such as ultrasensitivity and negative feedback (Huang and Ferrell, 1996, Kholodenko et al., 2010) might allow to translate minute amounts of MMP-cleaved pro-EGF into clear cut digital ERK pulses. Consistently, the slightly lower amplitude of spontaneous ERK pulses versus those induced by acute EGF stimulation suggests that the EGFR/MAPK system functions at the threshold input to generate digital ERK pulses (Figure 1G). This might allow the MMP/EGFR/MAPK signaling network to translate small variations in the EGFR input into frequency-modulated regimes of ERK pulses that can specify proliferation, survival and apoptosis. Note that exogenous addition of EGF impedes apoptosis-mediated lumen formation (Gaiko-Shcherbak et al., 2015), further suggesting that small amounts of EGFR ligands synthesized by the acinus itself are necessary for self-organisation of its morphogenesis. This MMP/EGFR/MAPK network has also been shown to produce ERK trigger waves in which activated cells sequentially switch on ERK pulses in adjacent cells (Aoki et al., 2017; Boocock et al., 2020; Hino et al., 2020). Our results strongly suggest that the ERK waves observed in stage 2 acini rely on this mechanism. Thus, a relatively simple signaling network might allow to produce conserved and sophisticated ERK behaviors in epithelial collectives. This excitable MMP/EGFR/MAPK network that generates pulsatile ERK activity strongly contrasts with the oscillatory ERK behavior observed in the segmentation clock in vertebrate embryos that is regulated on slower timescales by rhythmic transcriptional regulation of MAPK phosphatases (Hubaud and Pourquié, 2014). We speculate that the pulsatile MAPK network observed in epithelia provides an opportunity to constantly sense and

react to environmental inputs such as growth factors and insults to warrant epithelial homeostasis during acinar development. This illustrates how the MAPK network can be differently wired to produce distinct ERK dynamics at different timescales depending on the developmental context.

ERK dynamics regulate collective migration and proliferation in stage 1 acini

We found that stage 1 acini displayed high frequency, asynchronous ERK pulses during rapid collective cell migration that leads to a global rotation behavior of the acinus (Figure 2). This rotation behavior has previously been implicated in the morphogenesis of spherical tissue buds during mammary organogenesis (Fernández et al., 2021). During migration of 2D epithelial sheets, ERK waves co-ordinate myosin activity necessary for collective motility, and are shaped by a mechanochemical feedback from myosin to ERK (Aoki et al., 2017; Boocock et al., 2020; Hino et al., 2020). Further, ERK also has been shown to control myosin activity in MCF10A acini single-cell motility (Pearson and Hunter, 2007). We therefore propose that asynchronous ERK pulses spatially coordinate myosin contractility necessary for this collective motility behavior. This is consistent with our result that optogenetic synchronization of ERK pulses immediately leads to decreased collective motility (Figure 4C-F). The transition to a state of slower motility in late stage 1 acini has been shown to result from assembly of an endogenous laminin-rich basement membrane (Wang et al., 2013). We speculate that assembly of this basement membrane might modify the myosin-ERK mechanochemical feedback loop mentioned above, leading to decreased ERK frequency and motility, allowing to regulate the transition from proliferation to quiescence. Assembly of the basement membrane might therefore act as a checkpoint coordinating ERK frequency-dependent regulation of motility and transition from proliferation to quiescence. Future experimental/modeling studies will be necessary to further refine this hypothesis.

ERK waves spatially regulate apoptosis and survival fates during stage 2 acinar lumen morphogenesis

Our experiments in stage 2 acini suggest that spatial control of different ERK frequencies regulates survival in outer versus apoptosis fates in inner acinar cells. Outer cells display a median ERK frequency of one pulse every 4 hours, while inner cells display lower ERK frequencies (Figure 3A). This is consistent with the ability of

one ERK pulse to provide about 4 hours of survival in MCF10A monolayers (Gagliardi
et al., 2021). Using optogenetic control of ERK pulses, we excluded mechanisms such
as differential growth factor receptor sensitivity or refractory time of inner and outer
cells for regulation of different ERK frequencies (Supplementary Text, Figure S5A,B).
Instead, our results suggest a role for collective ERK waves to define the outer and
inner spatial domains of high and low ERK frequency, that respectively specify survival
and apoptosis fates (Figure 3B-I). ERK wave properties such as that they are initiated
mostly in the outer layer, and propagate more efficiently in the outer versus the inner
layer, might dynamically specify the two domains of ERK frequencies on timescale of
hours throughout the 7 days of the acinus cavitation process. While apoptosis is
predominantly responsible for the clearance of luminal cells in acini (Debnath et al.,
2002), we cannot exclude that an alternative mechanism such as autophagy resulting
from metabolic defects in inner cells, which is regulated by EGFR - PI3K signaling
might also contribute to this process (Schafer et al., 2009).

ERK pulse frequency but not integrated activity regulates survival versus
apoptosis fates
To formally test if ERK frequency regulates survival/apoptosis fates, we used
optogenetic stimulation of all the cells of acini to causally link ERK frequency with
lumen formation. When used to evoke ERK pulses for up to at least every 4 hours,
both optoFGFR and optoRaf led to loss of lumen formation (Figure 5C). ERK pulses
evoked every 10 hours were not sufficient to rescue apoptosis. These experiments
also functioned when EGFR was completely inhibited (Figure 5D). By optogenetically
varying the ERK pulse width, we also showed that ERK frequency rather than the
integrated ERK activity over time regulates survival versus apoptosis fates, both in
monolayers and in stage 2 acinus lumen formation (Figure 5E-G). Our data therefore
suggests that short ERK pulses are the signaling unit that allows cells in acini to
commit to survival for about 4 hours. This might allow cells to integrate signaling inputs
that fluctuate on timescales of minutes/hours, to dynamically control a morphogenetic
program on timescales of days. Interpretation of a specific ERK frequency into
survival/apoptosis fates might involve the ERK substrate Bcl-2-like protein 11 (Harada
et al., 2004) or ERK-dependent transcriptional control of immediate early genes (IEGs)
(Avraham and Yarden, 2011). IEGs produce transcripts with lifetimes of around 30
minutes, that encode proteins with lifetimes of 1-3 hours. Notably, IEGs include Jun

and Fos transcription factors which are important regulators of cell survival (Shaulian
and Karin, 2001). IEG's short lifetime is compatible with the ERK frequency required
for survival. Higher ERK frequencies as observed in stage 1 acini might control
proliferation by regulation of IEGs such as Fra-1 (Gillies et al., 2017).

Oncogenic PI3K signaling modulates ERK frequency contributing to aberrant
acinar morphogenesis.

Our results show that the breast cancer relevant PIK3CA H1047R mutation increases
ERK frequency, which might contribute at least in part to increased proliferation and
survival leading to larger acini without lumen formation, that phenocopies ductal
hyperplasia observed in mouse models (Tikoo et al., 2012). We show that this
crosstalk from PI3K to ERK signaling feeds into the control of ERK frequency both in
WT and PIK3CA H1047R monolayers and acini (Figure 7, S6-7). This crosstalk also
functions downstream of IGF1, that primarily activates PI3K-Akt signaling (Myers et
al., 1993) (Figure S7A,B), suggesting that it is a conserved feature downstream of
multiple receptor tyrosine kinases. During stage 1, PIK3CA H1047R acini display
similarly high ERK frequencies as in WT acini (Figure 7B). During stage 2, PIK3CA
H1047R acini then still display a substantial decrease in ERK frequency as in WT acini
compared to stage 1 (Figure 7B), which correlates with decreased proliferation (Figure
6E-H). However, ERK frequencies remain slightly higher both in outer and inner cells
in PIK3CA H1047R versus WT acini, correlating with the strong survival phenotype in
inner cells, and absence of lumen formation. These results suggest that the control of
ERK frequency is still subject to some degree of regulation even when the crosstalk
from PIK3CA H1047R is constitutively switched on. Because the ERK pulse shape is
identical in WT and PIK3CA mutant cells, this crosstalk must occur upstream of the
core Raf/MEK/ERK circuit. Consistently, ERK activation is sensitive to both EGFR and
MMP inhibition in PIK3CA H1047R cells (Figure S7C,D). As reported previously
(Young et al., 2015), our data strongly suggests that constitutive PI3K activity in
PIK3CA H1047R cells leads to increased expression of the EGFR ligand amphiregulin
that in turn might increase ERK frequency (Sternlicht et al., 2005) (Figure S7E).
PIK3CA H1047R also has been found to decrease expression of the protein tyrosine
phosphatase receptor type F (PTPRF) (Young et al., 2015), which might further
augment EGFR excitability and thus ERK frequency. Further work is required to
elucidate the specific contributions of PI3K and ERK signaling to control the

survival/apoptosis fate decisions and how the PI3K/ERK crosstalk might spatially fine
tune ERK frequency. Our result strongly suggests that oncogenic PI3K signaling-
induced aberrant spatial regulation of ERK frequency contributes to pathological
acinar morphogenesis.
585

Conclusion
We provide an initial characterization into how single-cell ERK dynamics control fate
decisions in space and time during the morphogenesis of a simple prototype organ
structure. Future studies are required to mechanistically understand how the different
dynamic signaling states are encoded and spatially organized, and if they provide
robustness against environmental perturbations occurring during development. This
will require the ever-expanding arsenal of optogenetic tools to manipulate specific cells
and evaluate how the cell collective responds. Further questions include how
additional signaling pathways might fine tune this morphogenetic process, and how
the ERK frequency is decoded into transcriptional programs that actuate the different
fates that shape acinus morphogenesis.

Material and methods

2D cell culture

MCF10A cells were cultured in DMEM/F12 supplemented with 5% horse serum, 20 ng/ml recombinant human EGF (Peprotech), 0.5 µg/ml hydrocortisone (Sigma-Aldrich/Merck), 10 µg/ml insulin (Sigma-Aldrich/Merck), 200 U/ml penicillin and 200 µg/ml streptomycin. The PIK3CA H1047R knockin cell line (Gustin et al., 2009) was a gift of Ben Ho Park. We regularly verified the presence of the mutation by sequencing the corresponding genomic locus. To generate stable cell lines, cells were transfected with FuGene (Promega) according to the manufacturer's protocol and clones were selected by antibiotic resistance and image-based screening.

3D cell culture

For acinus formation, single MCF10A cell suspensions were mixed with 4 volumes of growth factor-reduced Matrigel (Corning) at 4 °C and spread evenly on the surface of glass bottom cell culture plates at a concentration of 1.5 x 10^4 cells/cm^2. Acini were cultured in DMEM/F12 supplemented with 2% horse serum, 20 ng/ml recombinant human EGF, 0.5 µg/ml hydrocortisone, 10 µg/ml insulin, 200 U/ml penicillin and 200 µg/ml streptomycin. Horse serum, insulin and EGF were removed after 3 days of culture. For live imaging, 25 mM Hepes was added to the medium prior to mounting on the microscope. CellEvent Caspase 3/7 Green Detection Reagent was obtained from Thermo Fisher Scientific and used according to the manufacturer's protocol.

Plasmids

ERK-KTR-mTurquoise2 and ERK-KTR-mRuby2 were generated by fusion of the coding region of ERK-KTR (Regot et al., 2014) with that of mTurquoise2 (Goedhart et al., 2012) or mRuby2 (Lam et al., 2012). H2B-miRFP703 was generated by fusion of the coding region of human H2B clustered histone 11 (H2BC11) with that of miRFP703 (Shcherbakova et al., 2016). Geminin-mCherry was generated by fusion of the ubiquitylation domain of human Geminin (Sakaue-Sawano et al., 2017) to mCherry. The above mentioned fusion proteins were cloned in the piggyBac vectors pMP-PB, pSB-HPB (Balasubramanian et al., 2016) (gift of David Hacker, EPFL), or pPB3.0.Blast, an improved version of pPB generated in our lab.

pPB3.0-PuroCRY2-cRAF-mCitrine-P2A-CIBN-KrasCT, referred to in the manuscript

as OptoRAF, was generated in the following way: The CRY2-cRaf sequence was
excised from pCX4puro-CRY2-cRAF (gift from Kazuhiro Aoki, (Aoki et al., 2017))
using EcoRI and NotI. mCitrine was PCR amplified from the optoFGFR plasmid, while
adding NotI and XhoI sites, and digested. Both sequences were ligated into pPB3.0-
Puro, previously digested with EcoRI and XhoI. The GSGP2A-CIBN-KRasCT
sequence (synthesized by GENEWIZ) was digested with BsrGI and AflIII and ligated
into pPB3.0-Puro-CRY2-cRAF-mCitrine.
The piggyBac plasmids were co-transfected with a helper plasmid expressing a
hyperactive piggyBac transposase (Yusa et al., 2011).
Lyn-cytoFGFR1-PHR-mCit, expressing myristoylated FGFR1 cytoplasmic region
fused with the PHR domain of cryptochrome2 and mCitrine (gift from Won Do Heo
(Addgene plasmid # 59776), (Kim et al., 2014)), referred to in the manuscript as
OptoFGFR, was subcloned in a lentiviral backbone for stable cell line generation.
644

Imaging
All acini images were acquired on an epifluorescence Eclipse Ti2 inverted
fluorescence microscope (Nikon) equipped with a CSU-W1 spinning disk confocal
system (Yokogawa) and a Plan Apo VC 60X water immersion objective (NA = 1.2).
For time lapse imaging, laser-based autofocus was used. Images were acquired with
a Prime 95B or a Prime BSI sCMOS camera (both Teledyne Photometrics) at 16-bit
depth. Temperature, CO_2 and humidity were controlled throughout live imaging with a
temperature control system and gas mixer (both Life Imaging Services).
All monolayer cell images were acquired on an epifluorescence Eclipse Ti inverted
fluorescence microscope (Nikon) with a Plan Apo 20x air objective (NA = 0.8) or a
Plan Apo 40X air objective (NA = 0.9). Laser-based autofocus was used throughout
imaging. Images were acquired with an Andor Zyla 4.2 plus camera at 16-bit depth.
Both microscopes were controlled by NIS elements (Nikon).
658

Inhibitors and growth factors
Gefitinib was obtained from Sigma-Aldrich/Merck and used at a concentration of 10
µM. Trametinib was obtained from Selleck Chemicals and used at a concentration of
5 µM. Batimastat was obtained from MedChem Express and used at a concentration
of 30 µM. Pictilisib was obtained from Selleck Chemicals and used at a concentration
of 250 nM. AZD5363 was obtained from Selleck Chemicals and used at a

concentration of 5 μM. IGF1 was obtained from Peprotech and used at a concentration

of 100 nM.

Optogenetic experiments

For short term optogenetics experiments performed directly on the microscope, acini

were illuminated with wide field blue light (470 nm LED) at defined time points during

spinning disc time lapse imaging. Acini expressing optoFGF were illuminated for 100

ms at 50% LED intensity. Acini expressing optoRAF were illuminated for 100 ms at

60% LED intensity. The NIS elements JOBS module was used to program the imaging

and stimulation patterns.

The 2D monolayer optogenetic experiments were executed by culturing MCF10A

expressing optoFGFR, ERK-KTR-mRuby2 and H2B-miRFP703 as confluent

monolayer on 24 well plates with glass bottom. Optogenetic stimulation was done with

488 nm LED light at 100% light intensity for 100ms and using a 20x air objective. To

generate larger ERK activity pulses, cells were stimulated with trains of 488 nm light

pulses with a 2 min interval. The area under the curve (AUC) of ERK activity was

calculated using the ERK activity levels before and after the ERK pulses to set the

baseline. The correlation function between the number of 488 nm light pulses and

AUC was obtained by linear regression. The effects of different frequencies with same

AUC on apoptosis was measured after 24 hours of time-lapse acquisition and manual

annotation of apoptotic events on the base of morphological alterations of cell nucleus.

For long term optogenetic stimulation with the LITOS system, glass bottom 96-well cell

culture plates with 7-day old acini were fitted on a 32 x 64 RGB LED matrix (Boxtec)

inside a cell culture incubator. The matrix was connected to a custom printed circuit

board with an ESP32 microcontroller. This system was programmed to emit 1-minute

blue light pulses at maximal intensity at defined intervals for 7 days, after which lumen

formation efficiency was assessed. Acini were fixed with 4% paraformaldehyde prior

to imaging.

Image analysis of 3D acini

The open source LEVERJS software (Cohen, 2014; Wait et al., 2014; Winter et al.,

2016) was used to analyze the 3D time lapse movies. The LEVERJS software was

updated to include improved processing and visualization capabilities. The processing

pipeline began with a GPU-accelerated 3D non-local means denoising algorithm (Wait

et al., 2019). After denoising, a new ensemble-based segmentation algorithm was
applied. This ensemble segmentation combined an adaptive thresholding into
foreground/background regions with an anisotropic 3D Laplacian of Gaussian filter
targeted to a specific cell radius to separate touching cells (Winter et al., 2019). The
base segmentation was run at different cell radii and the results were combined using
unsupervised learning techniques from the field of algorithmic information theory
(Cohen et al., 2009). Here, the radii evaluated ranged from 2.5 µm to 4 µm in 0.5 µm
steps. These values were set empirically based on expected cell size ranges.
Following segmentation, the cells were tracked using Multitemporal Association
Tracking (Winter et al., 2011); (Winter et al., 2012).

Following segmentation and tracking of the image sequences, the ERK-KTR signal
was extracted and processed to a detrended and normalized signal. To extract the
ERK-KTR signal, distance transforms were computed for each segmented image. The
interior distance transform assigned each cell interior voxel a numeric value indicating
its distance starting at the cell boundary and increasing to the centroid. The exterior
distance transform assigned each boundary voxel a numeric value indicating its
distance to the nearest cell-assigned voxel. The exterior distance transform also
provided the identity of the nearest cell-assigned voxel for each background voxel.
The ERK-KTR signal was computed as the ratio of the image values around the center
of the cell to the image values around the boundary of the cell. The center region of
the cell included voxels in the 95[th] percentile of the interior distance transform. For the
boundary region of the cell we included interior voxels within one unit of the boundary
and exterior voxels within three units of the boundary. The resulting ERK-KTR signal
was computed as the ratio of the median voxel value in the outer region to the median
voxel value in the inner region. The extracted ERK-KTR signals for each cell had
different base intensities and showed different amounts of fluctuation. To normalize
this and to allow for quantitative comparison and visual representation with different
cell ERK-KTR expression levels we computed a detrended and normalized ERK-KTR
signal as follows. The signal S was first detrended by subtracting the median filter
signal, $S_d = S - \mathrm{median\_filter}(S)$. The signal was then normalized to the range [0,1]
using $S_{nd} = S_d / \max(S_d)$, unless the signal range in the detrended trajectory was below
an empirically set threshold. Inner and outer cells in stage 2 acini were identified
visually based on the 3D reconstructions in LEVERJS.

For segmentation and quantification of steady state Z-stacks, CellProfiler 3.1.8

(McQuin et al., 2018) with 3D functionalities was used. Nuclei were identified based

on H2B-miRFP signals. Apoptotic debris were identified using Caspase 3/7 Green

Detection kit signals using adaptive thresholding and watershed segmentation.

Geminin-mCherry intensities were measured within nuclei voxel masks. ERK-KTR

cytosolic/nuclear intensity ratios were generated by measuring median ERK-KTR

intensities in the nuclear area and in a spherical voxel mask 1 pixel around the nuclear

objects. Imaris software (Bitplane) was used for 3D rendering of confocal stacks and

to track and measure motility parameters in Figures 2 and 4.

Image analysis of 2D cell cultures

Nuclei of monolayer cells were segmented using Ilastik (Berg et al., 2019) and

CellProfiler 3.0. Ilastik was used for training a random forest classifier on different pixel

features of the H2B-miRFP channel and background pixels. The training data was

generated by manual annotation of 20 - 50 cells. The resulting probability map was

imported into CellProfiler for thresholding and segmentation. Cytosolic ERK-KTR

fluorescence intensities were measured by expansion of the nuclear objects. Cells

were tracked using μ-track 2.2.1 (Jaqaman et al., 2008).

Data analysis

All analysis and visualization of ERK activity peaks and time series was performed

with custom R/Matlab/Python code.

ERK activity pulse detection

ERK activity peaks were identified and counted by the following steps. 1. Application

of a short median filter to smooth the time series. 2. Application of a long median filter

to produce a bias estimate which was subtracted from the smoothed time series. 3.

Detrended time-series with real peaks were then identified by selecting those with an

activity difference above an empirical threshold. Those were rescaled to [0,1] and a

local maxima detection algorithm was used to identify peaks above an amplitude of

0.1.

Identification of collective events

To identify waves of collective ERK activation we developed a custom code and

implemented it in R. The algorithm works on a binarised signal that is calculated by

detrending and normalising ERK-KTR cytosolic/nuclear intensity ratio time series as
described above (Figure S3F,G). The algorithm searches for the first occurrence of
cells with ERK switched on (Figure S3H). If several such cells exist and they are
located within a threshold radius, they initiate the first collective event. A single active
cell can also become a collective event. In the next time point, the algorithm repeats
the search for active cells and compares their distances to cells in the previous frame.
If new active cells are located within the threshold distance to active cells at the
previous time point, they are added to respective collective events. If new active cells
are located outside of the threshold distance, they form new collective events. This
process of growing clusters of collective activity is repeated for all remaining time
points. The resulting statistics include the total number of cells involved in a collective
event, the duration and the average size of an event and the location (inner or outer
layer) of the cells that initiates the event.
778

Immunoblotting and qPCR

Cells were plated into 6-well dishes (2 x 10^5 cells/well) and cultured for 48h. The
resulting cell monolayers were washed twice with room temperature PBS, then
starving medium was added. For immunoblotting 1uL/mL DMSO or 10ug/mL
Batimastat was added together with the starving medium and cells were further
cultured for 72h. Media was removed, monolayers were washed twice with ice-cold
PBS, whole cell lysates were prepared and analyzed by immunoblotting as described
before (Dessauges et al., 2021). Primary antibodies against the following
proteins/epitopes were used: phospho-AKT^{Ser473} (cat. # 4058), AKT (cat. # 9272),
phospho-p44/42 MAPK (Erk1/2)$^{Thr202/Tyr204}$ (cat. # 4370), p44/42 MAPK (Erk1/2) (cat.
4695), all from Cell Signaling Technologies, BioConcept Ltd. Secondary
IRDye680LT- or IRDye800CW-conjugated anti-species specific IgGs were from LI-
COR. For qPCR cells were first starved for 24h, then fresh starving medium containing
1uL/mL DMSO or 2uM and 10uM pictilisib was added and cells were cultured for 24h.
Media was removed, monolayers were washed twice with ice-cold PBS, RNA was
isolated using TRIzol reagent. Reverse transcription was done with the ProtoScript II
reverse transcriptase kit (Bioconcept, M0368L). Real-time qPCR reactions were run
using the MESA Green qPCR MasterMix Plus for SYBR Green assay (Eurogenetec,
RT-SY2X-03+WOU) on the Rotor-Gen Q device (Qiagen). Each sample was tested in
triplicate. Expression of the gene of interest was calculated using the $2^{-\Delta\Delta Ct}$ method.

The sequences of the primer sets used are as follows: AREG, 5' -ACA TTT CCA TTC

TCT TGT CG- 3' (forward), and 5' - ACA TTT CCA TTC TCT TGT CG- 3' (reverse);

FLJ22101, 5' -TTC CCT GTG GCA CTT GAC ATT- 3' (forward), and 5'-CTT TTG CCT

CTG GCA GTA CTC A-3' (reverse).

Quantification and statistical analysis

All graphs were assembled and statistics were performed using R or Excel. Box plots

depict the median and the 25th and 75th percentiles; whiskers correspond to minimum

and maximum non-outlier values in Figures 1G, 2D, 3A, 3H, 3I, 5F, 6C, 6F, 7B, 7E,

S1C, S3B, S3C, S3D, S3E, S4B, S6A, S6B, S6C, S6D and S7A. Dot plots show

distribution of 50 randomly selected data points per condition, or all data points if there

are less than 50. Red lines in ERK activity trajectories represent the population

average. The statistical tests used and the significance thresholds are indicated in

each respective legend.

Data and code availability

The open-source code for LEVERJS is available at https://leverjs.net/git. All 3D

datasets that were analyzed with LEVERJS for this publication can be browsed

interactively at https://leverjs.net/mcf10a_3d.

Acknowledgments

The authors are grateful to Ben Ho Park for providing the MCF10A PIK3CA H1047R

knockin MCF10A cells, to Kazuhiro Aoki for the optoRaf construct, and to Won Do Heo

for the optoFGFR construct. This work was supported by grants from the Swiss

National Science Foundation and the Swiss Cancer Research Foundation to Olivier

Pertz and from a Human Frontiers Science Program grant to Olivier Pertz and Andrew

Cohen. We acknowledge support of the Microscopy Imaging Center of the University

of Bern (https://www.mic.unibe.ch/).

Author contribution

PE, PAG and OP designed the study. PE, PAG and AF performed experiments and analyzed

data. MD and M-AJ analyzed data. CD provided expertise with the optogenetic tools. TH

provided expertise with LITOS. ARC performed image analysis using LEVERJS. PE, PAG and
OP wrote the paper.

Conflict of interest

The authors declare that they have no conflict of interest.

References

Aikin, T.J., Peterson, A.F., Pokrass, M.J., Clark, H.R., and Regot, S. (2020). MAPK activity
dynamics regulate non-cell autonomous effects of oncogene expression. ELife *9*.

Albeck, J.G., Mills, G.B., and Brugge, J.S. (2013). Frequency-modulated pulses of ERK
activity transmit quantitative proliferation signals. Mol. Cell *49*, 249–261.

Anderson, L.R., Sutherland, R.L., and Butt, A.J. (2010). BAG-1 overexpression attenuates
luminal apoptosis in MCF-10A mammary epithelial cells through enhanced RAF-1 activation.
Oncogene *29*, 527–538.

Aoki, K., Kumagai, Y., Sakurai, A., Komatsu, N., Fujita, Y., Shionyu, C., and Matsuda, M.
(2013). Stochastic ERK activation induced by noise and cell-to-cell propagation regulates cell
density-dependent proliferation. Mol. Cell *52*, 529–540.

Aoki, K., Kondo, Y., Naoki, H., Hiratsuka, T., Itoh, R.E., and Matsuda, M. (2017). Propagating
wave of ERK activation orients collective cell migration. Dev. Cell *43*, 305-317.e5.

Avraham, R., and Yarden, Y. (2011). Feedback regulation of EGFR signalling: decision
making by early and delayed loops. Nat. Rev. Mol. Cell Biol. *12*, 104–117.

Balasubramanian, S., Wurm, F.M., and Hacker, D.L. (2016). Multigene expression in stable
CHO cell pools generated with the piggyBac transposon system. Biotechnol. Prog. *32*, 1308–
1317.

Berglund, F.M., Weerasinghe, N.R., Davidson, L., Lim, J.C., Eickholt, B.J., and Leslie, N.R.
(2013). Disruption of epithelial architecture caused by loss of PTEN or by oncogenic mutant
p110α/PIK3CA but not by HER2 or mutant AKT1. Oncogene *32*, 4417–4426.

Berg, S., Kutra, D., Kroeger, T., Straehle, C.N., Kausler, B.X., Haubold, C., Schiegg, M., Ales,
J., Beier, T., Rudy, M., et al. (2019). ilastik: interactive machine learning for (bio)image
analysis. Nat. Methods *16*, 1226–1232.

Boocock, D., Hino, N., Ruzickova, N., Hirashima, T., and Hannezo, E. (2020). Theory of
mechanochemical patterning and optimal migration in cell monolayers. Nat. Phys.

Cancer Genome Atlas Network (2012). Comprehensive molecular portraits of human breast
tumours. Nature *490*, 61–70.

Chakrabarty, A., Rexer, B.N., Wang, S.E., Cook, R.S., Engelman, J.A., and Arteaga, C.L.
(2010). H1047R phosphatidylinositol 3-kinase mutant enhances HER2-mediated
transformation by heregulin production and activation of HER3. Oncogene *29*, 5193–5203.

Chen, N., Eritja, N., Lock, R., and Debnath, J. (2013). Autophagy restricts proliferation driven
by oncogenic phosphatidylinositol 3-kinase in three-dimensional culture. Oncogene *32*, 2543–
2554.

Ciarloni, L., Mallepell, S., and Brisken, C. (2007). Amphiregulin is an essential mediator of
estrogen receptor alpha function in mammary gland development. Proc Natl Acad Sci USA
104, 5455–5460.

Cohen, A.R. (2014). Extracting meaning from biological imaging data. Mol. Biol. Cell *25*, 3470–
3473.

Cohen, A.R., Bjornsson, C.S., Temple, S., Banker, G., and Roysam, B. (2009). Automatic
summarization of changes in biological image sequences using algorithmic information theory.
IEEE Trans. Pattern Anal. Mach. Intell. *31*, 1386–1403.

Debnath, J., Mills, K.R., Collins, N.L., Reginato, M.J., Muthuswamy, S.K., and Brugge, J.S.
(2002). The role of apoptosis in creating and maintaining luminal space within normal and
oncogene-expressing mammary acini. Cell *111*, 29–40.

Debnath, J., Muthuswamy, S.K., and Brugge, J.S. (2003). Morphogenesis and oncogenesis
of MCF-10A mammary epithelial acini grown in three-dimensional basement membrane
cultures. Methods *30*, 256–268.

Dessauges, C., Mikelson, J., Dobrzynski, M., Jacques, M.-A., Frismantiene, A., Gagliardi,
P.A., Khammash, M., and Pertz, O. (2021). Optogenetic actuator/biosensor circuits for large-
scale interrogation of ERK dynamics identify sources of MAPK signaling robustness. BioRxiv.

Du, W.W., Fang, L., Li, M., Yang, X., Liang, Y., Peng, C., Qian, W., O'Malley, Y.Q., Askeland,
R.W., Sugg, S.L., et al. (2013). MicroRNA miR-24 enhances tumor invasion and metastasis
by targeting PTPN9 and PTPRF to promote EGF signaling. J. Cell Sci. *126*, 1440–1453.

Fernández, P.A., Buchmann, B., Goychuk, A., Engelbrecht, L.K., Raich, M.K., Scheel, C.H.,
Frey, E., and Bausch, A.R. (2021). Surface-tension-induced budding drives alveologenesis in
human mammary gland organoids. Nat. Phys. *17*, 1130–1136.

Finlay, D., Healy, V., Furlong, F., O'Connell, F.C., Keon, N.K., and Martin, F. (2000). MAP
kinase pathway signalling is essential for extracellular matrix determined mammary epithelial
cell survival. Cell Death Differ. *7*, 302–313.

Gagliardi, P.A., Dobrzyński, M., Jacques, M.-A., Dessauges, C., Ender, P., Blum, Y., Hughes,
R.M., Cohen, A.R., and Pertz, O. (2021). Collective ERK/Akt activity waves orchestrate
epithelial homeostasis by driving apoptosis-induced survival. Dev. Cell *56*, 1712-1726.e6.

Gaiko-Shcherbak, A., Fabris, G., Dreissen, G., Merkel, R., Hoffmann, B., and Noetzel, E.
(2015). The acinar cage: basement membranes determine molecule exchange and
mechanical stability of human breast cell acini. PLoS ONE *10*, e0145174.

Gillies, T.E., Pargett, M., Minguet, M., Davies, A.E., and Albeck, J.G. (2017). Linear Integration
of ERK Activity Predominates over Persistence Detection in Fra-1 Regulation. Cell Syst. *5*,
549-563.e5.

Goedhart, J., von Stetten, D., Noirclerc-Savoye, M., Lelimousin, M., Joosen, L., Hink, M.A.,
van Weeren, L., Gadella, T.W.J., and Royant, A. (2012). Structure-guided evolution of cyan
fluorescent proteins towards a quantum yield of 93%. Nat. Commun. *3*, 751.

Goglia, A.G., Wilson, M.Z., Jena, S.G., Silbert, J., Basta, L.P., Devenport, D., and Toettcher,
J.E. (2020). A Live-Cell Screen for Altered Erk Dynamics Reveals Principles of Proliferative
Control. Cell Syst. *10*, 240-253.e6.

Gustin, J.P., Karakas, B., Weiss, M.B., Abukhdeir, A.M., Lauring, J., Garay, J.P., Cosgrove,
D., Tamaki, A., Konishi, H., Konishi, Y., et al. (2009). Knockin of mutant PIK3CA activates
multiple oncogenic pathways. Proc Natl Acad Sci USA *106*, 2835–2840.

Harada, H., Quearry, B., Ruiz-Vela, A., and Korsmeyer, S.J. (2004). Survival factor-induced
extracellular signal-regulated kinase phosphorylates BIM, inhibiting its association with BAX
and proapoptotic activity. Proc Natl Acad Sci USA *101*, 15313–15317.

Hino, N., Rossetti, L., Marín-Llauradó, A., Aoki, K., Trepat, X., Matsuda, M., and Hirashima,
T. (2020). ERK-Mediated Mechanochemical Waves Direct Collective Cell Polarization. Dev.
Cell *53*, 646-660.e8.

Hiratsuka, T., Fujita, Y., Naoki, H., Aoki, K., Kamioka, Y., and Matsuda, M. (2015). Intercellular
propagation of extracellular signal-regulated kinase activation revealed by in vivo imaging of
mouse skin. ELife *4*, e05178.

Höhener, T.C., Landolt, A., Dessauges, C., Gagliardi, P.A., and Pertz, O. (2022). LITOS - a
versatile LED illumination tool for optogenetic stimulation. BioRxiv.

Huang, C.Y., and Ferrell, J.E. (1996). Ultrasensitivity in the mitogen-activated protein kinase
cascade. Proc Natl Acad Sci USA *93*, 10078–10083.

Hubaud, A., and Pourquié, O. (2014). Signalling dynamics in vertebrate segmentation. Nat.
Rev. Mol. Cell Biol. *15*, 709–721.

Huebner, R.J., Neumann, N.M., and Ewald, A.J. (2016). Mammary epithelial tubes elongate
through MAPK-dependent coordination of cell migration. Development *143*, 983–993.

Inman, J.L., Robertson, C., Mott, J.D., and Bissell, M.J. (2015). Mammary gland development:
cell fate specification, stem cells and the microenvironment. Development *142*, 1028–1042.

Isakoff, S.J., Engelman, J.A., Irie, H.Y., Luo, J., Brachmann, S.M., Pearline, R.V., Cantley,
L.C., and Brugge, J.S. (2005). Breast cancer-associated PIK3CA mutations are oncogenic in
mammary epithelial cells. Cancer Res. *65*, 10992–11000.

Jaqaman, K., Loerke, D., Mettlen, M., Kuwata, H., Grinstein, S., Schmid, S.L., and Danuser,
G. (2008). Robust single-particle tracking in live-cell time-lapse sequences. Nat. Methods 5,
695–702.

Kholodenko, B.N., Hancock, J.F., and Kolch, W. (2010). Signalling ballet in space and time.
Nat. Rev. Mol. Cell Biol. *11*, 414–426.

Kim, N., Kim, J.M., Lee, M., Kim, C.Y., Chang, K.-Y., and Heo, W.D. (2014). Spatiotemporal
control of fibroblast growth factor receptor signals by blue light. Chem. Biol. *21*, 903–912.

Lam, A.J., St-Pierre, F., Gong, Y., Marshall, J.D., Cranfill, P.J., Baird, M.A., McKeown, M.R.,
Wiedenmann, J., Davidson, M.W., Schnitzer, M.J., et al. (2012). Improving FRET dynamic
range with bright green and red fluorescent proteins. Nat. Methods *9*, 1005–1012.

Lauring, J., Cosgrove, D.P., Fontana, S., Gustin, J.P., Konishi, H., Abukhdeir, A.M., Garay,
J.P., Mohseni, M., Wang, G.M., Higgins, M.J., et al. (2010). Knock in of the AKT1 E17K
mutation in human breast epithelial cells does not recapitulate oncogenic PIK3CA mutations.
Oncogene *29*, 2337–2345.

Lavoie, H., Gagnon, J., and Therrien, M. (2020). ERK signalling: a master regulator of cell
behaviour, life and fate. Nat. Rev. Mol. Cell Biol. *21*, 607–632.

Liu, J.S., Farlow, J.T., Paulson, A.K., Labarge, M.A., and Gartner, Z.J. (2012). Programmed
cell-to-cell variability in Ras activity triggers emergent behaviors during mammary epithelial
morphogenesis. Cell Rep. *2*, 1461–1470.

Mailleux, A.A., Overholtzer, M., Schmelzle, T., Bouillet, P., Strasser, A., and Brugge, J.S.
(2007). BIM regulates apoptosis during mammary ductal morphogenesis, and its absence
reveals alternative cell death mechanisms. Dev. Cell *12*, 221–234.

McQuin, C., Goodman, A., Chernyshev, V., Kamentsky, L., Cimini, B.A., Karhohs, K.W., Doan,
M., Ding, L., Rafelski, S.M., Thirstrup, D., et al. (2018). CellProfiler 3.0: Next-generation image

processing for biology. PLoS Biol. *16*, e2005970.

Myers, M.G., Sun, X.J., Cheatham, B., Jachna, B.R., Glasheen, E.M., Backer, J.M., and White,
M.F. (1993). IRS-1 is a common element in insulin and insulin-like growth factor-I signaling to
the phosphatidylinositol 3'-kinase. Endocrinology *132*, 1421–1430.

Paine, I.S., and Lewis, M.T. (2017). The Terminal End Bud: the Little Engine that Could. J.
Mammary Gland Biol. Neoplasia *22*, 93–108.

Patterson, K.I., Brummer, T., O'Brien, P.M., and Daly, R.J. (2009). Dual-specificity
phosphatases: critical regulators with diverse cellular targets. Biochem. J. *418*, 475–489.

Pearson, G.W., and Hunter, T. (2007). Real-time imaging reveals that noninvasive mammary
epithelial acini can contain motile cells. J. Cell Biol. *179*, 1555–1567.

Reginato, M.J., Mills, K.R., Becker, E.B.E., Lynch, D.K., Bonni, A., Muthuswamy, S.K., and
Brugge, J.S. (2005). Bim regulation of lumen formation in cultured mammary epithelial acini is
targeted by oncogenes. Mol. Cell. Biol. *25*, 4591–4601.

Regot, S., Hughey, J.J., Bajar, B.T., Carrasco, S., and Covert, M.W. (2014). High-sensitivity
measurements of multiple kinase activities in live single cells. Cell *157*, 1724–1734.

Sakaue-Sawano, A., Yo, M., Komatsu, N., Hiratsuka, T., Kogure, T., Hoshida, T., Goshima,
N., Matsuda, M., Miyoshi, H., and Miyawaki, A. (2017). Genetically encoded tools for optical
dissection of the mammalian cell cycle. Mol. Cell *68*, 626-640.e5.

Schafer, Z.T., Grassian, A.R., Song, L., Jiang, Z., Gerhart-Hines, Z., Irie, H.Y., Gao, S.,
Puigserver, P., and Brugge, J.S. (2009). Antioxidant and oncogene rescue of metabolic
defects caused by loss of matrix attachment. Nature *461*, 109–113.

Sebastian, J., Richards, R.G., Walker, M.P., Wiesen, J.F., Werb, Z., Derynck, R., Hom, Y.K.,
Cunha, G.R., and DiAugustine, R.P. (1998). Activation and function of the epidermal growth
factor receptor and erbB-2 during mammary gland morphogenesis. Cell Growth Differ. *9*, 777–
785.

Shaulian, E., and Karin, M. (2001). AP-1 in cell proliferation and survival. Oncogene *20*, 2390–
2400.

Shcherbakova, D.M., Baloban, M., Emelyanov, A.V., Brenowitz, M., Guo, P., and Verkhusha,
V.V. (2016). Bright monomeric near-infrared fluorescent proteins as tags and biosensors for
multiscale imaging. Nat. Commun. *7*, 12405.

Sparta, B., Pargett, M., Minguet, M., Distor, K., Bell, G., and Albeck, J.G. (2015). Receptor
Level Mechanisms Are Required for Epidermal Growth Factor (EGF)-stimulated Extracellular
Signal-regulated Kinase (ERK) Activity Pulses. J. Biol. Chem. *290*, 24784–24792.

Sternlicht, M.D., Sunnarborg, S.W., Kouros-Mehr, H., Yu, Y., Lee, D.C., and Werb, Z. (2005). Mammary ductal morphogenesis requires paracrine activation of stromal EGFR via ADAM17-dependent shedding of epithelial amphiregulin. Development *132*, 3923–3933.

Tikoo, A., Roh, V., Montgomery, K.G., Ivetac, I., Waring, P., Pelzer, R., Hare, L., Shackleton, M., Humbert, P., and Phillips, W.A. (2012). Physiological levels of Pik3ca(H1047R) mutation in the mouse mammary gland results in ductal hyperplasia and formation of ERα-positive tumors. PLoS ONE *7*, e36924.

Valon, L., Davidović, A., Levillayer, F., Villars, A., Chouly, M., Cerqueira-Campos, F., and Levayer, R. (2021). Robustness of epithelial sealing is an emerging property of local ERK feedback driven by cell elimination. Dev. Cell *56*, 1700-1711.e8.

Wait, E., Winter, M., Bjornsson, C., Kokovay, E., Wang, Y., Goderie, S., Temple, S., and Cohen, A.R. (2014). Visualization and correction of automated segmentation, tracking and lineaging from 5-D stem cell image sequences. BMC Bioinformatics *15*, 328.

Wait, E., Winter, M., and Cohen, A.R. (2019). Hydra image processor: 5-D GPU image analysis library with MATLAB and python wrappers. Bioinformatics *35*, 5393–5395.

Wang, H., Lacoche, S., Huang, L., Xue, B., and Muthuswamy, S.K. (2013). Rotational motion during three-dimensional morphogenesis of mammary epithelial acini relates to laminin matrix assembly. Proc Natl Acad Sci USA *110*, 163–168.

Winter, M.R., Fang, C., Banker, G., Roysam, B., and Cohen, A.R. (2012). Axonal transport analysis using Multitemporal Association Tracking. Int. J. Comput. Biol. Drug Des. *5*, 35–48.

Winter, M., Wait, E., Roysam, B., Goderie, S.K., Ali, R.A.N., Kokovay, E., Temple, S., and Cohen, A.R. (2011). Vertebrate neural stem cell segmentation, tracking and lineaging with validation and editing. Nat. Protoc. *6*, 1942–1952.

Winter, M., Mankowski, W., Wait, E., Temple, S., and Cohen, A.R. (2016). LEVER: software tools for segmentation, tracking and lineaging of proliferating cells. Bioinformatics *32*, 3530–3531.

Winter, M., Mankowski, W., Wait, E., De La Hoz, E.C., Aguinaldo, A., and Cohen, A.R. (2019). Separating touching cells using pixel replicated elliptical shape models. IEEE Trans. Med. Imaging *38*, 883–893.

Young, C.D., Zimmerman, L.J., Hoshino, D., Formisano, L., Hanker, A.B., Gatza, M.L., Morrison, M.M., Moore, P.D., Whitwell, C.A., Dave, B., et al. (2015). Activating PIK3CA Mutations Induce an Epidermal Growth Factor Receptor (EGFR)/Extracellular Signal-regulated Kinase (ERK) Paracrine Signaling Axis in Basal-like Breast Cancer. Mol. Cell. Proteomics *14*, 1959–1976.

1031   Yusa, K., Zhou, L., Li, M.A., Bradley, A., and Craig, N.L. (2011). A hyperactive piggyBac
1032   transposase for mammalian applications. Proc Natl Acad Sci USA *108*, 1531–1536.