Chromosomal neighbourhoods allow identification of organ specific changes in gene expression

Rishi Das Roy[1*], Outi Hallikas[1], Mona M. Christensen[1], Elodie Renvoisé[1,2], Jukka Jernvall[1,3*]

[1]Institute of Biotechnology, University of Helsinki, P.O. Box 56, FI-00014 Helsinki, Finland.
[2]5 Rue de Portland, 72100 Le Mans, France.
[3]Department of Geosciences and Geography, University of Helsinki, P.O. Box 64, FI-00014 Helsinki, Finland.

[*]To whom correspondence should be addressed. Email: rishi.dasroy@helsinki.fi and jernvall@fastmail.fm

ABSTRACT

Although most genes share their chromosomal neighbourhood with other genes, distribution of genes has not been explored in the context of individual organ development; the common focus of developmental biology studies. Because developmental processes are often associated with initially subtle changes in gene expression, here we explored whether neighbouring genes are informative in the identification of differentially expressed genes. First, we quantified the chromosomal neighbourhood patterns of genes having related functional roles in the mammalian genome. Although the majority of protein coding genes have at least five neighbours within 1 Mb window around each gene, very few of these neighbours regulate development of the same organ. Analyses of transcriptomes of developing mouse molar teeth revealed that whereas expression of genes regulating tooth development changes, their neighbouring genes show no marked changes, irrespective of their level of expression. Finally, we test whether inclusion of gene neighbourhood in the analyses of differential expression could provide additional benefits. For the analyses, we developed an algorithm, called DELocal that identifies differentially expressed genes by comparing their expression changes to changes in adjacent genes in their chromosomal regions. Our results show that DELocal removes detection bias towards large changes in expression, thereby allowing identification of even subtle changes in development. Future studies, including the detection of differential expression, may benefit from, and further characterize the significance of gene-gene neighbour relationships.

Author Summary

Development of organs is typically associated with small and hard to detect changes in gene expression. Here we examined how often genes regulating specific organs are neighbours to each other in the genome, and whether this gene neighbourhood helps in the detection of changes in gene expression. We found that genes regulating individual organ development are very rarely close to each other in the mouse and human genomes. We built an algorithm, called DELocal, to detect changes in gene expression that incorporates information about neighbouring genes. Using transcriptomes of developing mouse molar teeth containing gene expression profiles of thousands of genes, we show how genes regulating tooth development are ranked high by DELocal even if their expression level changes are subtle. We propose that developmental biology studies can benefit from gene neighbourhood analyses in the detection of differential expression and identification of organ specific genes.

INTRODUCTION

Temporal and spatial regulation of gene expression is important in development and differentiation. During development, genes are expressed in a highly dynamic manner, and perturbations of the expression dynamics underlie many diseases and developmental defects. Regulation of gene expression through developmental time can be examined by quantifying gene expression levels at two or more time points. However, because development is typically a gradual process, detecting differential gene expression can be challenging due to the changes being initially subtle. This challenge underscores the continuing need for different strategies to identify biologically meaningful changes in gene expression.

The relationship between gene expression and genome organization is one aspect that has attracted considerable research interest. In prokaryotes, the genome structure and the regulation of gene expression are operationally linked. This is because the genes involved in the same process are typically located in the same operons, forming multigene clusters that were originally read as single transcripts [1]. In eukaryotes, operons controlling multiple genes are rare although nearly 15% of the 20,000 genes in *C. elegans* genome are located in operons [2, 3]. Nevertheless, most eukaryotic genes are scattered throughout the genome without apparent order related to expression or function [4]. Even though there does not appear to be any simple logic in the distribution of eukaryotic genes, it is not entirely random as most genes cannot be moved around in the genome without seriously disturbing their functionality [5]. Some non-random distribution is found both at the small scale, relating to a limited number of genes, and at the large scale, concerning large chromosomal regions [5, 6]. Clustering has been detected among co-expressed genes, such as the housekeeping genes, as also within functional groups, such as interacting pairs of proteins or biological processes containing thousands of genes [7, 8]. Overall, factors contributing to the genome organization include at least evolutionary history and current evolutionary forces, mechanisms of rearrangement, mechanisms of regulation of gene expression and control of chromatin composition. The main source of non-random gene order in mammalian genomes has been suggested to be tandem duplications [9].

An additional factor in the eukaryote nucleus is the three-dimensional organization of the genome [10, 11]. During metazoan development, enhancers regulate the gene activity in different cell and tissue types through dynamic three-dimensional hubs, in which enhancers and promoters assemble in physical proximity to activate gene expression. Genome-wide analyses of the spatial organization of chromatin have uncovered multiple topologically associating domains (TADs) that form largely self-interacting regions of regulatory elements and promoters [12, 13]. Although there are cases where gene expression is co-expressed within TADs,[14, 15] current evidence

points towards a more complex relationship between gene co-regulation and TADs [16,

17]. In addition, TADs appear to have multiple hierarchies [16], and questions remain on

the degree of TAD-conservation across species[18].

In this study we examine genome organization from the point of individual organ

development; the common focus of developmental biology and regeneration studies. We

first review the spatial distribution of protein coding genes in the mammalian genome and

quantify neighbouring genes involved in development of the same organs. Next, our

specific organ system for in-depth analyses of gene neighbourhoods and gene

expression is the mammalian tooth. Especially developing mouse molar teeth provide

many well characterized genes to examine links between gene neighbourhood and

expression [[19] and references therein]. Our starting point is 1 Mb chromosomal

neighbourhood around each organ specific gene as these encompass the majority of

both regulatory regions and constrained architecture seen in the conservation of large or

very orderly gene clusters. The choice of 1Mb window is obviously somewhat arbitrary,

but it should provide a readily applicable framework to diverse taxa and data. This

definition of neighbourhood should be fairly compatible with TADs as they happen to be

approximately 1Mb in the mouse (Dixon et al. Nature, 485, 2012; they say 880kb)[12].

Nevertheless, we also test our inferences using TADs calculated for each

neighbourhood. The 1Mb windows include well known gene clusters such as Hox genes

and immunoglobulin genes, whose spatial organization on the chromosome is crucial to

their regulation and function. Still, this strict co-localization of co-regulated genes is not

the general pattern in mammalian genomes [9], and the patterns of gene clustering

related to gene function are delicate and complex.

Finally, for detection of differentially expressed genes in organ development, we

developed a new method, called DELocal. This method does not depend solely on the

read counts from transcriptome sequencing, but it includes in the analysis the expression

data of the neighbouring genes. This approach removes the emphasis on fold-changes

that is common to other methods and allows DELocal to identify relatively subtle gene

expression changes during development. Using embryonic mouse dental transcriptomes

obtained with both microarray and RNAseq, we show how DELocal provides an

alternative ranking of differentially expressed (DE) genes.

RESULTS

Majority of protein coding genes have neighbouring genes within 1 Mb window

As our focus of interest is developmental regulation, we first obtained an overall pattern

of distribution of developmental genes by tabulating how closely genes are located in the

genome. For example, the human genome (GRCh38.p13) is 4.5*10-9 base pairs long
and has 20,449 protein coding genes, which means, on average, one gene for every
220,060 bases (Ensembl release 99). Similarly, for the mouse genome, this number is
154,290 (GRCm38.p6). Consequently, on average, 5 to 6 genes should reside in each 1
Mb window in the mouse genome. To express these statistics as a neighbourhood, we
can state that each gene has 4 to 5 neighbouring genes within a 1 Mb window. This
observation is obviously a broad generalization, but it does indicate that genes tend to
have some neighbours within 1 Mb. To examine the neighbourhood patterns in more
detail, we examined the 1 Mb neighbourhoods of protein coding genes of the mouse
genome. For every gene, number of neighbouring genes within 1 Mb window was
counted (Figure 1A, Supplementary Figure S1). This simple calculation shows that, at the
level of all protein coding genes, majority of genes in mouse genome have more than 4
neighbours, the median number of neighbours being 15 (Supplementary Figure S2). This
tabulation indicates that there is some level of clustering of genes in the eukaryotic
genome, a pattern well established in the literature [20].

Majority of protein coding genes lack neighbours regulating the same organ

To contrast the genome-wide pattern of gene neighbourhoods with that of a single organ
system, we first tabulated gene neighbourhoods associated with the development of the
mouse tooth. This specific organ focus basically illustrates whether genes participating in
the regulation of the same organ are also located close to each other (which is not
expected). Here mouse molar development provides a good example because its gene
regulation is relatively well understood, and because tooth development itself is a
relatively autonomous process [19, 21]. We analyzed the 1 Mb neighbourhoods of the
tooth developmental genes (Supplementary Table S1) and the number of tooth genes
that are sharing the same neighbourhood. The results show that tooth genes are mostly
located far from each other, suggesting that genes regulating this specific developmental
process are not co-localized (Figure 1B), a pattern that is in agreement with the
organization of eukaryotic genomes [10, 11]. Taken together, although majority of protein
coding genes are to some extent clustered in the mouse genome, tooth genes tend to be
located far from each other (Figure 1).

Figure 1. Although protein-coding genes typically have neighbours, tooth genes do not have other tooth gene neighbours within 1 Mb windows around each gene. (**A**) The number of neighbouring genes within 1 Mb window around each gene tabulated from the mouse genome for all protein coding genes and (**B**) genes involved in tooth development. Majority of 21,971 protein coding genes have at least five neighbours whereas most of 302 tooth developmental genes do not have tooth genes as neighbours. This pattern suggests that genes with specific functions are sparsely distributed.

Next, we examined how representative the developing mouse tooth (Figure 1B) is of

other developmental systems. We used the gene ontology (GO) terms to provide an

approximation of the adjacency of genes involved in the same developmental process.

Here we focused in the genes belonging to mouse GO slims (a concise list of terms) of

"biological process" [22, 23]. For every gene belonging to these terms, its 1 Mb

neighbourhood was investigated to tabulate genes belonging to the same term. The

tabulations show that most genes belonging to a certain GO term are sparsely distributed

in the chromosomes (Figure 2) which is in accordance with previous studies [5, 24].

There are only few genes from broader, or high level, GO terms that are densely located

(more than 3 genes from the same GO term within 1 Mb neighbourhood). However, these

few GO terms represent very broad descriptions of biological functions. With more

precise GO terms (Figure 2, bottom row), genes tend to have no neighbours belonging to

the same GO term. For generality, we examined these patterns in human genome and

they remained largely the same (Supplementary Figure S3).

Figure 2. The more specific the GO term is the fewer neighbours its genes have from the same GO term within 1 Mb. Each pie represents the genes of one GO term under the root GO term 'biological processes'. The top three rows represent GO slim terms. The GO terms in the box are children of the GO term 'system development'. The color-coding indicates the number of neighbours that a gene has from the same GO term. The number of genes in the GO term is indicated under the GO term. Analysis was done for the mouse genome. See Supplementary Table S2 for GO IDs.

The scarcity of neighbouring genes in the precise GO term categories is not surprising considering the limited number of genes in each category. To test if the patterns are a simple function of group size, we performed simulations where genes were assigned randomly to artificial GO terms containing different numbers of genes. For every group size, 10,000 simulations were performed and the density of genes was calculated for each artificial GO term in the 1 Mb neighbourhood. A plot showing the percentage of genes lacking neighbours shows the expected decrease as the number of genes increases in the artificial GO terms (Figure 3). The empirical patterns largely follow the randomizations, although some real GO terms with small number of genes (300 and less) show slightly lower percentages, indicating that there is a slight tendency for higher spatial clustering of genes belonging to the same GO terms than in the simulations. This may be partly due to GO terms having paralogous genes that are sometimes located near each other in the genome. Nevertheless, up to GO term categories containing 1000 genes, 80 % of genes have no neighbours belonging to the same GO terms. These

patterns indicate that the general expectation for the genes sharing the same genomic

neighbourhood (Figure 1A) is that they participate in the regulation of different systems

and organs. To the extent that this is the general rule of genomic organization, it provides

opportunities for new computational approaches. Below we develop a method to detect

differential expression using additional information from the neighbouring genes and test

it using our data on the developing mouse tooth.

Figure 3. Real GO terms with lower gene numbers are slightly more clustered in
the genome than artificial GO terms with the same gene numbers. Artificial GO
terms with different numbers of genes were made with randomly selected genes and
their distribution across the genome was measured. For each group size, 10,000
simulations were executed and for each simulation the percentage of genes with zero
within-1Mb-neighbours with the same artificial GO term were counted (black dots). Real
GO terms are marked with blue dots. See Supplementary Table S2 for GO IDs. The blue
line is median, and the shadow shows the observations between 1st and 3rd quartile.

Neighbouring genes do not show organ-specific changes in expression

Because genes linked to tooth development do not appear to be co-localized in the

genome (Figure 1B), we checked whether these genes differ in their expression

dynamics compared to the other genes in their neighbourhood. We examined differential

expression of genes at the onset of tooth crown formation, between mouse embryonic

day 13.5 (E13) and 14.5 (E14) when many of the tooth genes are known to be

upregulated [19, 25]. For example, in visual inspection *Ctnna1*, *Shh*, *Foxi3* and *Sostdc1*,

all genes required for normal tooth development [26], show upregulation between E13

and E14 (Figure 4). In contrast, the other genes in their neighbourhoods show no marked

changes between E13 and E14, irrespective of their expression levels (Figure 4).

Therefore, the distribution of genes involved in the regulation of the tooth (Figure 1 and 4)

is manifested also at the level of differential expression. Building from this observation,

we developed a new algorithm, DELocal, to identify DE genes based on their neighbours'

expression dynamics (Methods). To evaluate its potential performance, we used gene

expression data from embryonic mouse dental tissues generated by both microarray and

RNAseq.

Figure 4. Only tooth developmental genes are differentially expressed within 1 Mb windows in the developing mouse molar. Median expression levels of the tooth developmental genes *Ctnna1*, *Shh, Foxi3* and *Sostdc1* and their neighbouring genes at developmental stages E13 and E14. Regardless of their expression level, the surrounding genes show little change between the two stages whereas the tooth genes are upregulated. *Egr1* in the 1 Mb window of *Ctnna1* is also a tooth developmental gene.

The number of neighbours is not critical for detecting differential expression

Our hypothesis of neighbouring genes being informative in the detection of differential

expression is dependent on the definition of 'neighbourhood'. Therefore, it is important to

determine the right number of neighbours to include in the analysis by the DELocal

algorithm. To define the optimal number of neighbours we tested different numbers (1-

14) of closest genes within a fixed window (1 Mb) surrounding the gene of interest. We

evaluated the performance of DELocal with different numbers of closest neighbours in

identifying the genes involved in tooth development. Again we contrasted the expression

levels between E13 to E14 molar teeth (for 302 tooth genes), or so-called bud stage to

cap stage transition in tooth development, when many tooth genes are known to be

active [19]. The Matthews correlation coefficient (MCC) scores were measured for

different numbers of neighbours to examine the strength of DELocal to identify tooth

genes (true positive; TP) as well as non- tooth genes (true negatives; TN). The MCC

score was chosen to optimize the model due to very few TPs, or imbalanced dataset.

The results show that DELocal produces similar and stable MCC scores on both

microarray and RNAseq datasets, even though RNAseq data produces slightly higher

MCC scores than microarray (Figure 5). We note that only one nearest gene is enough to

obtain close to the highest MCC score. However, for RNAseq the best MCC score

corresponds to 5 nearest neighbours. Because there are fewer genes available in

microarray analyses compared to RNAseq, in the rest of the analyses we used DELocal

with 5 neighbours both for the RNAseq and microarray data. It should be also noted that

the distances of these 5 neighbours are different for each gene and therefore the final

neighbourhood windows are not constant in size. Additionally to the 1 Mb window,

median TAD boundaries were also used to define the neighbourhoods and to evaluate

DELocal. Although, TAD boundaries are different for each gene there was no marked

difference in the result compared to 1 Mb neighbourhood (Supplementary Figure S4).

Figure 5. DELocal performance is not strongly dependent on the number of gene
neighbours used in the analysis. Every gene is evaluated in relation to its neighbouring
genes. In the absence of any "gold standard" for the number of neighbours, different
numbers of nearest genes (within 1 Mb window) were used to identify the differentially
expressed genes. The overall performances were measured using MCC. The
performance of DELocal using RNAseq data was slightly better than with microarray
data.

DELocal shows increased precision compared to other methods

Microarray is one of the earliest successful high throughput technologies to measure a

large number of gene expressions, and consequently there are a good number of

statistical methods to identify differentially expressed genes from datasets generated by

this platform. Hence, the performance of DELocal can be evaluated by comparisons to

these methods using a microarray dataset. However, microarray is limited to only the

genes that have been targeted by microarray probes. Therefore, the expression of all the

genes cannot be accessed, resulting in fewer neighbouring genes being sampled. To

obtain a more comprehensive readout of differentially expressed genes, RNAseq was

also used to evaluate DELocal performance. The performance of all these methods was

measured by their ability to identify differentially expressed tooth genes.

For analysis of performance, we used the receiver operating curve (ROC) [27]

depicting the true positive rate against the false positive rate of differentially expressed

genes. The analyses show that DELocal outperforms other methods in identifying tooth

genes using both microarray and RNAseq (Figure 6). The performance is most similar to

Limma/DEMI (microarray) and Limma/DESeq (RNAseq). We used also other metrics like

specificity, recall (sensitivity), precision and MCC to evaluate and compare the different

methods. DELocal shows high specificity and accuracy compared to other methods for

microarray data (Supplementary Figure S5).

Figure 6. Compared to other methods, DELocal is at least as powerful in detecting
differentially expressed genes. Receiver operating characteristic (ROC) curves and
areas under the curves (within the parenthesis) show that DELocal outperforms other
methods on both microarray and RNAseq data.

In RNAseq data, DELocal outperforms other methods except in recall (sensitivity)

(Figure 7). The MCC scores remain equivalent to each other. The tooth gene dataset is

imbalanced due to the large number of non-tooth genes (true negatives), which hinders

the evaluation of accuracy, but does not affect F1 or MCC. Considering that the objective

of many experiments is to find true positives, the F1 score, which is a compound-term of

precision and recall, is an important metric. The F1 score ranges from zero (bad) to one

(good), and the F1 scores of all of the methods remain suboptimal. Nevertheless,

DELocal using RNAseq outperforms other methods also in F1 score. Below we discuss

the results from RNAseq only.

Figure 7. Comparison of DELocal with earlier methods to identify differential
expression. Evaluation matrices show that, except for recall (sensitivity), DELocal equals
or outperforms earlier methods. However due to large number of true negatives, the
significance scores of precision, F1 and MCC remained negligible. The evaluation
matrices are explained in materials and methods section. The analysis was done using
RNAseq data. For microarray data see Supplementary Figure S5.

DELocal provides an alternative ranking of genes by removing bias towards large
changes in expression

Because DELocal quantifies expression changes in the context of neigbouring genes,

genes in the chromosomal regions having generally large changes in expression should

not be as highly ranked as they would be in other methods. We examined the rankings

and expression changes of tooth genes among predicted differentially expressed genes

by all the methods. DELocal predicted tooth genes are enriched in top rank positions

compared to the other three methods (Supplementary Figure S6). In experimental

research, gene rankings are typically the first values to be examined in validating the

results biologically, and in practice high ranking is very influential when selecting

candidate genes for new downstream analyses. The results show that whereas DEseq,

edgeR and limma emphasize in their rankings genes that show large changes in their

expression, this bias is not present in the DELocal ranking (Figure 8A). These results

also indicate that the identity of highly ranked tooth genes is different between DELocal

and the other methods (Figure 8B). Thus, DELocal differs from the other methods more

substantially than is apparent based on the different performance measures alone

(Figure 6, 7).

Figure 8. DELocal provides an alternative ranking of differentially expressed

genes. (A). Unlike the other methods, DELocal rankings of differential expression do not

bias towards genes with large log-fold changes (1 denotes the highest rank). (B) Principal

component analyses of the rank orders of tooth developmental genes shows the ordering

of the DELocal to be distinct from the ones provided by the other three methods (using

correlation matrix of top 500 tooth genes).

Differentially expressed genes missed by DELocal tend to have local paralogues

The DELocal algorithm appears to efficiently identify the differentially expressed genes

(having high precision) as well as to filter out non tooth genes (having high specificity).

Still, DELocal missed 60 differentially expressed tooth genes which are identified by all

the other methods in RNAseq dataset. DELocal is built on the hypothesis that every true

differentially expressed gene should have neighbours and none of them should be

differentially expressed. Consequently, DELocal may fail to identify those differentially
expressed genes whose neighbours are also differentially expressed. For instance, 2 out
of 5 nearest neighbours of *Bmp7* were differentially expressed genes which could be the
reason of failure of DELocal to detect *Bmp7* (Figure 9). Additionally, the presence of
paralogous genes in the neighbourhood may complicate the detection of differential
expression. Most notably, *Dlx1* and *Dlx2* in chromosome 11 (Figure 9), *Dlx5* and *Dlx6* in
chromosomes 6, and *Dlx1* and *Dlx2* in chromosome 2, are co-regulated as pairs and can
compensate for the deletion of one another [28-31]. Additionally, the other neighbouring
paralogs in our set of tooth genes are *Cyp26c1* with *Cyp26a1*, and *Cdh1* with *Cdh3* [32].
Therefore, at least some of the genes missed by DELocal should be possible to detect by
incorporating information about gene paralogues into the algorithm.

Figure 9. Differentially expressed genes in the same neighbourhood interfere with
the detection of differentially expressed genes by DELocal. DELocal failed to identify
Bmp7 due to differential expression patterns of its neighbouring genes (top). *Dlx1* and
*Dlx2 are p*aralogous tooth genes which are in the same neighbourhood (bottom). Only
tooth developmental genes are labelled here.

DISCUSSION

Only seven protein-coding genes in Ensembl mouse annotation lack neighbours
altogether within their 1 Mb windows (Figure 1A). Hence, the overwhelming majority of
mouse genes have neighbours, and this pattern of organization is likely to apply to all
mammals, if not beyond. In the genomic scale of thousands of genes, neighbouring
genes can be co-expressed in eukaryotes [33]. Our analyses, focused on individual
organ development, suggest that the neighbouring genes are very rarely involved in the
regulation of the same organ or tissue (Figure 2), and that this is also reflected in the lack
of co-expression of neighbours (Figure 4). The lack of co-expressed neighbours is partly
explained by the limited number of genes that are associated with individual organ
regulation (Figures 2 and 3). In contrast, more general processes, such as cell

proliferation or cell death, are linked to many thousands of genes (Figure 2), increasing

the likelihood of neighbourhood co-expression. Indeed, an analysis using high-level GO

terms containing thousands of genes found that functionally related genes are often

located close to each other [8]. It is plausible that evolution of organs such as teeth may

have co-opted genes from multiple high-level clusters, perhaps further explaining the

rarity of co-expressed neighbours in organ regulation.

A distinct advantage of the rarity of co-expressed neighbours in organ development is

that this information can be used to detect even subtle changes in gene expression. To

this end, here we developed the DELocal algorithm to identify differentially expressed

genes by incorporating expression information from chromosomal neighbourhood.

DELocal provided highly precise detection of differentially expressed genes in mouse

tooth development (Figures 6 and 7), suggesting that this method can provide additional

benefits in the analysis of developmental systems. Specifically, DELocal is distinct from

the previous methods of determining differentially expressed genes by not relying on

large fold-changes in gene expression (Figure 8A). The ability of DELocal to identify

biologically significant but more subtle changes is based on its special feature: it can

track a change in gene expression that is unique among the surrounding genes in the

chromosomal location. Thus, DELocal can pinpoint genes which are being actively and

specifically regulated between two developmental stages, regardless of whether the

change of expression is large or small. Taken together, DELocal can complement other

methods by providing alternative rank listings of potential genes of interests (Figure 8B).

More technically, the developed DELocal algorithm is based on linear models that

have been successfully implemented and used in Limma, DeSeq2 and many other

methods to identify differentially expressed genes [34, 35]. With linear models, gene

expression can be modelled in two or more biological conditions and thereafter

differentially expressed genes of different contrasts of interest can be found. Linear

models are advantageous compared to other methods in that they can model complex

experimental conditions with multiple factors. Here we used DELocal to determine the

genes that are differentially expressed between bud and cap stage in the developing

mouse tooth. The extensive list of genes active in tooth development [19, 21] allowed us

to optimize and evaluate the performance of DELocal. Considering that the in vivo bud

and cap stage differences in gene expression are relatively subtle, the high specificity

and accuracy of DELocal is promising. Obviously, the optimization requires a list of genes

of interest. However, DELocal can also be run without any prior knowledge of genes that

are active in a particular developmental process. Figure 3 shows that in relation to the

number of neighbours, the performance of DELocal is very stable and even only one

nearest neighbour could be sufficient to build the models. The implication of this is that

DELocal could be used with only a single neighbour contrast in the absence of any

reference/training gene set. In this context it will be interesting to test whether the

optimum number of neighbours is the same for different organs. Another future direction

is to examine different TAD classifications using DELocal.

It remains to be explored whether the rarity of neighbouring genes regulating the

same process is related, for example, to the requirements of spacing required for folding

of the DNA, or whether it is the result of historical processes of genome rearrangements

or co-option of genes in the evolution of organs. Regardless, our results on the

developing mouse tooth suggest that the lack of organ specific neighbours is also

manifested at the level of differential expression. We further show that it is possible to

develop computational approaches to detect even subtle changes in gene expression

using the additional information from the neighbouring genes.

MATERIALS AND METHODS

Ethics statement

All mouse studies were approved and carried out in accordance with the guidelines of the

Finnish national animal experimentation board under licenses KEK16-021,

ESAVI/2984/04.10.07/2014 and ESAV/2363/04.10.07/2017.

Gene sets and gene neighbourhood analyses

The start position, chromosome name and gene biotype (protein coding, lncRNA,

ncRNA, pseudogene) of every gene of mouse and human genomes were downloaded

from Ensembl database using R package biomaRT [36] in February, 2020. Further

analysis was limited to protein coding genes. 302 genes linked to tooth development

were marked as tooth developmental genes (Supplementary Table S1 and refs [19, 37]).

Throughout the study, the gene start coordinates are used as the position of the gene in

the chromosome. The same coordinates are used to measure the distances between

genes. The GO slim terms were acquired from AmiGO [38] and they are listed in

Supplementary Table S2. For each GO term, the corresponding genes were downloaded

again from Ensembl using biomaRT. The artificial GO terms of different sizes for

simulation were created by randomly sampling protein coding genes from genome using

custom R scripts. For every group size, 10,000 simulations were performed and for each

simulation density of genes from the same artificial GO term in neighbourhood was

calculated.

For most of the analyses, we used the 1 Mb windows as a neighbourhood measure for

each focal gene as these are relatively straightforward to determine for different genomes

and genes. Additionally, we analyzed neighbourhoods using TADs. Although TADs are

considered to be conserved across different mammalian cell types, multiple TADs

surrounding a gene with different boundaries can be delineated [39]. For a robust

consensus of TAD boundaries, we downloaded four different mouse cell-lines and three

different algorithms (with 10 kb and 50 kb resolutions, June 2021 [40]). A consensus TAD

boundary for each gene was defined by selecting a median start and end co-ordinates

from all TADs surrounding that gene. The median size of these consensus TAD

boundaries was 990kb (within the range of 185kb to 6Mb, standard deviation = 547813.8)

which is roughly similar to previous report of 880kb[41]. TAD boundaries for more than

1500 genes (mostly from Y chromosome) were not determined.

RNAseq library preparation

Developing mouse molar teeth from embryonic days 13.5 (E13) and 14.5 (E14) were

dissected from wild type C57BL/Ola embryos. For RNAseq, five biological replicates were

used. The samples were stored in RNAlater (Qiagen GmbH, Hilden, Germany) in -75°C.

RNA was extracted first twice with guanidinium thiocyanate-phenol-chloroform extraction

and then further purified using RNeasy Plus micro kit (Qiagen GmbH, Hilden, Germany)

according to manufacturer's instructions. RNA quality of representative samples was

assessed with 2100 Bioanalyzer (Agilent, Santa Clara, CA) and the RIN values were 9 or

higher. The RNA concentrations were determined by Qubit RNA HS Assay kit (Thermo

Fisher Scientific, Waltham, MA). The cDNA libraries were prepared with Ovation RNAseq

System V2 (Nugene, Irvine, CA), and sequenced with NextSeq500 (Illumina, San Diego,

CA).

Microarray library preparation

Mouse E13 and E14 teeth were dissected from wild type NMRI embryos. Five biological

replicates were used. The amount of RNA available in each sample was measured with

2100 Bioanalyzer (Agilent, Santa Clara, CA). Only the samples showing a RIN value

above 9 were used for the microarray analysis.

Gene Expression analysis

Gene expression was measured both in microarray (platform: GPL6096, Affymetrix

Mouse Exon Array 1.0) and RNAseq (platforms GPL19057, Illumina NextSeq 500). The

microarray gene signals were normalized with aroma.affymetrix [42] package using

Brainarray custom CDF (Version 19, released on Nov 13, 2014) [43]. The RNAseq reads

(84 bp) were evaluated and bad reads were filtered out using FastQC [44], AfterQC [45]

and Trimmomatic [46]. This resulted in on average 63 million reads per sample. Then

good reads were aligned with STAR [47] to

Mus_musculus.GRCm38.dna.primary_assembly.fa genome and counts for each gene

was performed by HTSeq [48] tool using Mus_musculus.GRCm38.90.gtf annotation. On

average 89% reads were uniquely mapped to Mus musculus genome. Additionally,

RNAseq count values were normalized using DESeq2 [34]. All the transcriptomic data

are available in NCBI Gene Expression Omnibus under the accession number

GSE142201.

DELocal

In DELocal, it is hypothesized that differentially expressed genes have different

expression dynamics compared to their neighbouring genes. Specifically, we want to find

genes whose expression changes, while that of the surrounding genes do not. To

calculate this, we used a similar logic to ESLiM [49], an algorithm that detects changes in

exon usage. Whereas ESLiM is suitable for detection of alternative splicing (exon usage),

DELocal is applicable for identifying differentially expressed genes in the chromosomal

neighbourhood. Note that in principle one could choose the neighbouring gene or genes

randomly across the genome, but this would result in different results in each

randomization.

In this algorithm, gene's expression is modelled as a linear relationship with median

expression of neighbourhood genes, such as,

$\hat{g}_{ij} = s_i \times N\widetilde{g}w_{ij} + b_i$ .... (i),

Where $\hat{g}_{ij}$ is expected expression of i-th gene in j-th sample, $N\widehat{g}w_{ij}$ is median

expression of N nearest neighbouring genes within 1 Mb window of the i-th gene from j-th

sample and b_i is base line expression level of gene g_i. The slope s_i of every gene g_i

depends on its neighbouring genes. Therefore, the difference between expected and

observed values or residual,

$r_{ij} = g_{ij} - \hat{g}_{ij}$ ...... (ii)

where g_{ij} is observed value. For differentially expressed genes, these residual values

will be significantly different in different biological conditions.

Furthermore, with the aid of the residuals r_{ij} observed g_{ij} could be formulated as follow,

$g_{ij} = s_i \times N\widetilde{g}w_{ij} + b_i + r_{ij}$.... (iii)

Noticeably, this relationship Eq. (iii) is independent of experimental condition and only

dependent on neighbouring gene. Therefore, differentially expressed genes are detected

through significantly deviated residual values between the desired contrasts using

Empirical Bayes statistics, available from Limma package [35]. We tested the

performance of DELocal using from 1 to 14 neighbouring genes (N in Eq. (iii)). The log-

normalized and normalized count values were used in DELocal respectively for

microarray and RNAseq data. There are 334 protein coding genes in mouse genome

which do not have any other protein coding gene in their 1Mb neighbourhood. Therefore,

we also used available non-protein coding genes from the neighbourhood in DELocal

analysis. However, after the inclusion of non-protein coding genes there are still 17

protein coding genes without any neighbours within 1 Mb.

Performance measures

DELocal was compared with different publicly available tools applicable both for

microarray or RNAseq : RankProd[50], SAM[51], DEMI[52], Limma[35], edgeR[53] and

DESeq2 [34]. All these programs were executed with default parameters. Genes reported

with p-value <= 0.05 by these tools were marked as differentially expressed and used to

evaluate the performance of each tool using the following metrics and receiver operating

characteristic (ROC) curves.

• Sensitivity (Recall), true positive rate TPR = TP / (TP + FN)

• Specificity, true negative rate SPC = TN / (TN + FP)

• Precision, positive predictive value PPV = TP / (TP + FP)

• Accuracy, ACC = (TP + TN) / (TP + FP + FN + TN)

• F1 score, F1 = 2TP / (2TP + FP + FN)

• Mathews correlation coefficient (MCC) $= \dfrac{TP \times TN - FP \times FN}{\sqrt{[(TP+FP)(TP+FN)(TN+FP)(TN+FN)]}}$

where TP, true positive; FP, false positive; TN, true negative; FN, false negative

Tooth developmental genes were used to find the true and false positive rate for the

analyses (Supplementary Table S1 and refs [19, 37]). Rest of the genes (in thousands)

were considered as true negatives. Due to the large number of these, very high accuracy

can be achieved by predicting all genes as non-differentially expressed. Therefore,

because our objective was to find true tooth genes, we focused on Recall, Precision, F1,

MCC and area under the ROC curves rather than only Accuracy for evaluating the

methods. The areas under the ROC curves were calculated with ROCR [27] package.

Additionally, we compared differences in rank and log-fold changes in the genes ranked

high by the different methods (Figure 8).

FUNDING

558 Financial support was provided by the Academy of Finland, Jane and Aatos Erkko

559 Foundation, John Templeton Foundation, and Sigrid Jusélius Foundation.

560 *Conflict of interest statement*. None declared.

DATA AVAILABILITY

563 The DELocal algorithm is freely available here https://github.com/dasroy/DELocal as an

564 R package. The microarray and RNAseq data are deposited in the NCBI Gene

565 Expression Omnibus with accession codes GSE141907 and GSE142199.

ACKNOWLEDGEMENTS

567 We thank P. Rastas and P. Auvinen for discussions. We thank P. Auvinen, L. Paulin and

568 P. Laamanen at DNA Sequencing and Genomics Laboratory for bulk RNA sequencing.

REFERENCES

1.   Jacob F, Monod J. Genetic regulatory mechanisms in the synthesis of proteins.
J Mol Biol. 1961;3:318-56.
2.   Blumenthal T, Davis P, Garrido-Lecca A. Operon and non-operon gene clusters
in the C. elegans genome. WormBook. 2015:1-20.
3.   Girard LR, Fiedler TJ, Harris TW, Carvalho F, Antoshechkin I, Han M, et al.
WormBook: the online review of Caenorhabditis elegans biology. Nucleic Acids Res.
2007;35(Database issue):D472-5.
4.   Zabidi MA, Stark A. Regulatory Enhancer-Core-Promoter Communication via
Transcription Factors and Cofactors. Trends in genetics : TIG. 2016;32(12):801-14.
5.   Lercher MJ, Urrutia AO, Hurst LD. Clustering of housekeeping genes provides a
unified model of gene order in the human genome. Nature genetics. 2002;31(2):180-3.
6.   Vogel JH, von Heydebreck A, Purmann A, Sperling SJBB. Chromosomal
clustering of a human transcriptome reveals regulatory background. 2005;6(1):230.
7.   Thévenin A, Ein-Dor L, Ozery-Flato M, Shamir R. Functional gene groups are
concentrated within chromosomes, among chromosomes and in the nuclear space of
the human genome. Nucleic Acids Research. 2014;42(15):9854-61.
8.   Tiirikka T, Siermala M, Vihinen M. Clustering of gene ontology terms in
genomes. Gene. 2014;550(2):155-64.
9.   Lemay DG, Martin WF, Hinrichs AS, Rijnkels M, German JB, Korf I, et al. G-
NEST: a gene neighborhood scoring tool to identify co-conserved, co-expressed
genes. 2012;13(1):253.
10.   Furlong EEM, Levine M. Developmental enhancers and chromosome topology.
Science. 2018;361(6409):1341-5.

11.   Robson MI, Ringel AR, Mundlos S. Regulatory Landscaping: How Enhancer-
Promoter Communication Is Sculpted in 3D. Mol Cell. 2019;74(6):1110-22.
12.   Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, et al. Topological domains in
mammalian genomes identified by analysis of chromatin interactions. Nature.
2012;485(7398):376-80.
13.   Nora EP, Lajoie BR, Schulz EG, Giorgetti L, Okamoto I, Servant N, et al. Spatial
partitioning of the regulatory landscape of the X-inactivation centre. Nature.
2012;485(7398):381-5.
14.   Flavahan WA, Drier Y, Liau BB, Gillespie SM, Venteicher AS, Stemmer-
Rachamimov AO, et al. Insulator dysfunction and oncogene activation in IDH mutant
gliomas. Nature. 2016;529(7584):110-4.
15.   Symmons O, Uslu VV, Tsujimura T, Ruf S, Nassari S, Schwarzer W, et al.
Functional and topological characteristics of mammalian regulatory domains. Genome
Res. 2014;24(3):390-400.
16.   Sikorska N, Sexton T. Defining Functionally Relevant Spatial Chromatin
Domains: It is a TAD Complicated. J Mol Biol. 2020;432(3):653-64.
17.   Soler-Oliva ME, Guerrero-Martinez JA, Bachetti V, Reyes JC. Analysis of the
relationship between coexpression domains and chromatin 3D organization. PLoS
Comput Biol. 2017;13(9):e1005708.
18.   Eres IE, Gilad Y. A TAD Skeptic: Is 3D Genome Topology Conserved? Trends
in genetics : TIG. 2021;37(3):216-23.
19.   Hallikas O, Das Roy R, Christensen M, Renvoisé E, Sulic A-M, Jernvall J.
System-level analyses of keystone genes required for mammalian tooth development.
J Exp Zool Part B Mol Dev Evol [in press]. 2020.
20.   Lane N, Martin W. The energetics of genome complexity. Nature.
2010;467(7318):929-34.
21.   Jernvall J, Thesleff I. Tooth shape formation and tooth renewal: evolving with the
same signals. Development. 2012;139(19):3487-97.
22.   The Gene Ontology C. The Gene Ontology Resource: 20 years and still GOing
strong. Nucleic Acids Res. 2019;47(D1):D330-D8.
23.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene
ontology: tool for the unification of biology. The Gene Ontology Consortium. Nature
genetics. 2000;25(1):25-9.
24.   de Laat W, Grosveld F. Spatial organization of gene expression: the active
chromatin hub. Chromosome Research. 2003;11(5):447-59.
25.   Gene expression in tooth (WWW database), Developmental Biology Programme
of the University of Helsinki, [Internet]. University of Helsinki. 1996-2007. Available
from: http://bite-it.helsinki.fi.
26.   Li CY, Hu J, Lu H, Lan J, Du W, Galicia N, et al. alphaE-catenin inhibits
YAP/TAZ activity to regulate signalling centre formation during tooth development.
Nature communications. 2016;7:12133.
27.   Sing T, Sander O, Beerenwinkel N, Lengauer T. ROCR: visualizing classifier
performance in R. Bioinformatics (Oxford, England). 2005;21(20):3940-1.
28.   Thomas BL, Tucker AS, Qui M, Ferguson CA, Hardcastle Z, Rubenstein JL, et
al. Role of Dlx-1 and Dlx-2 genes in patterning of the murine dentition. Development.
1997;124(23):4811-8.

29.   Qiu M, Bulfone A, Ghattas I, Meneses JJ, Christensen L, Sharpe PT, et al. Role
of the Dlx homeobox genes in proximodistal patterning of the branchial arches:
mutations of Dlx-1, Dlx-2, and Dlx-1 and -2 alter morphogenesis of proximal skeletal
and soft tissue structures derived from the first and second arches. Dev Biol.
1997;185(2):165-84.
30.   Beverdam A, Merlo GR, Paleari L, Mantero S, Genova F, Barbieri O, et al. Jaw
transformation with gain of symmetry after Dlx5/Dlx6 inactivation: mirror of the past?
Genesis. 2002;34(4):221-7.
31.   Debiais-Thibaud M, Metcalfe CJ, Pollack J, Germon I, Ekker M, Depew M, et al.
Heterogeneous conservation of Dlx paralog co-expression in jawed vertebrates. PloS
one. 2013;8(6):e68182.
32.   Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al.
eggNOG 4.5: a hierarchical orthology framework with improved functional annotations
for eukaryotic, prokaryotic and viral sequences. Nucleic Acids Research.
2015;44(D1):D286-D93.
33.   Fukuoka Y, Inaoka H, Kohane IS. Inter-species differences of co-expression of
neighboring genes in eukaryotic genomes. BMC genomics. 2004;5(1):4.
34.   Love MI, Huber W, Anders SJGB. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. 2014;15(12):550.
35.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Research. 2015;43(7):e47-e.
36.   Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. Nature
protocols. 2009;4(8):1184-91.
37.   Nieminen P, Pekkanen M, Aberg T, Thesleff I. A graphical WWW-database on
gene expression in tooth. Eur J Oral Sci. 1998;106 Suppl 1:7-11.
38.   Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S, et al. AmiGO:
online access to ontology and annotation data. Bioinformatics (Oxford, England).
2009;25(2):288-9.
39.   Zufferey M, Tavernari D, Oricchio E, Ciriello G. Comparison of computational
methods for the identification of topologically associating domains. Genome Biol.
2018;19(1):217.
40.   Liu T, Porter J, Zhao C, Zhu H, Wang N, Sun Z, et al. TADKB: Family
classification and a knowledge base of topologically associating domains. BMC
genomics. 2019;20(1):217.
41.   Yu M, Ren B. The Three-Dimensional Organization of Mammalian Genomes.
Annu Rev Cell Dev Biol. 2017;33:265-89.
42.   Bengtsson H, Irizarry R, Carvalho B, Speed TP. Estimation and assessment of
raw copy numbers at the single locus level. Bioinformatics (Oxford, England).
2008;24(6):759-67.
43.   Dai M, Wang P, Boyd AD, Kostov G, Athey B, Jones EG, et al. Evolving
gene/transcript definitions significantly alter the interpretation of GeneChip data.
Nucleic Acids Res. 2005;33(20):e175.
44.   Andrews S. FastQC: A quality control tool for high throughput sequence data.
2010 [Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

45.   Chen S, Huang T, Zhou Y, Han Y, Xu M, Gu JJBB. AfterQC: automatic filtering,
trimming, error removing and quality control for fastq data. 2017;18(3):80.
46.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics (Oxford, England). 2014;30(15):2114-20.
47.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics (Oxford, England). 2013;29(1):15-
21.
48.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics (Oxford, England). 2015;31(2):166-9.
49.   Risueno A, Roson-Burgo B, Dolnik A, Hernandez-Rivas JM, Bullinger L, De Las
Rivas J. A robust estimation of exon expression to identify alternative spliced genes
applied to human tissues and cancer samples. BMC genomics. 2014;15:879.
50.   Del Carratore F, Jankevics A, Eisinga R, Heskes T, Hong F, Breitling R.
RankProd 2.0: a refactored bioconductor package for detecting differentially expressed
features in molecular profiling datasets. Bioinformatics (Oxford, England).
2017;33(17):2774-5.
51.   Tusher VG, Tibshirani R, Chu G. Significance analysis of microarrays applied to
the ionizing radiation response. Proceedings of the National Academy of Sciences of
the United States of America. 2001;98(9):5116-21.
52.   Ilmjärv S, Hundahl CA, Reimets R, Niitsoo M, Kolde R, Vilo J, et al. Estimating
differential expression from multiple indicators. Nucleic Acids Research.
2014;42(8):e72-e.
53.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics (Oxford,
England). 2009;26(1):139-40.
713
1.   Jacob F, Monod J. Genetic regulatory mechanisms in the synthesis of proteins.
J Mol Biol. 1961;3:318-56. Epub 1961/06/01. doi: 10.1016/s0022-2836(61)80072-7.
PubMed PMID: 13718526.
2.   Blumenthal T, Davis P, Garrido-Lecca A. Operon and non-operon gene clusters
in the C. elegans genome. WormBook. 2015:1-20. Epub 2015/05/06. doi:
10.1895/wormbook.1.175.1. PubMed PMID: 25936768; PubMed Central PMCID:
PMCPMC4781206.
3.   Girard LR, Fiedler TJ, Harris TW, Carvalho F, Antoshechkin I, Han M, et al.
WormBook: the online review of Caenorhabditis elegans biology. Nucleic Acids Res.
2007;35(Database issue):D472-5. Epub 2006/11/14. doi: 10.1093/nar/gkl894. PubMed
PMID: 17099225; PubMed Central PMCID: PMCPMC1669767.
4.   Zabidi MA, Stark A. Regulatory Enhancer-Core-Promoter Communication via
Transcription Factors and Cofactors. Trends in genetics : TIG. 2016;32(12):801-14.
Epub 2016/11/07. doi: 10.1016/j.tig.2016.10.003. PubMed PMID: 27816209.
5.   Lercher MJ, Urrutia AO, Hurst LD. Clustering of housekeeping genes provides a
unified model of gene order in the human genome. Nature genetics. 2002;31(2):180-3.
doi: 10.1038/ng887.
6.   Vogel JH, von Heydebreck A, Purmann A, Sperling SJBB. Chromosomal
clustering of a human transcriptome reveals regulatory background. 2005;6(1):230. doi:
10.1186/1471-2105-6-230.

7. Thévenin A, Ein-Dor L, Ozery-Flato M, Shamir R. Functional gene groups are
concentrated within chromosomes, among chromosomes and in the nuclear space of
the human genome. Nucleic Acids Research. 2014;42(15):9854-61. doi:
10.1093/nar/gku667 %J Nucleic Acids Research.
8. Tiirikka T, Siermala M, Vihinen M. Clustering of gene ontology terms in
genomes. Gene. 2014;550(2):155-64. Epub 2014/07/06. doi:
10.1016/j.gene.2014.06.060. PubMed PMID: 24995610.
9. Lemay DG, Martin WF, Hinrichs AS, Rijnkels M, German JB, Korf I, et al. G-
NEST: a gene neighborhood scoring tool to identify co-conserved, co-expressed
genes. 2012;13(1):253. doi: 10.1186/1471-2105-13-253.
10. Furlong EEM, Levine M. Developmental enhancers and chromosome topology.
Science. 2018;361(6409):1341-5. Epub 2018/09/29. doi: 10.1126/science.aau0320.
PubMed PMID: 30262496; PubMed Central PMCID: PMCPMC6986801.
11. Robson MI, Ringel AR, Mundlos S. Regulatory Landscaping: How Enhancer-
Promoter Communication Is Sculpted in 3D. Mol Cell. 2019;74(6):1110-22. Epub
2019/06/22. doi: 10.1016/j.molcel.2019.05.032. PubMed PMID: 31226276.
12. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, et al. Topological domains in
mammalian genomes identified by analysis of chromatin interactions. Nature.
2012;485(7398):376-80. Epub 2012/04/13. doi: 10.1038/nature11082. PubMed PMID:
22495300; PubMed Central PMCID: PMCPMC3356448.
13. Nora EP, Lajoie BR, Schulz EG, Giorgetti L, Okamoto I, Servant N, et al. Spatial
partitioning of the regulatory landscape of the X-inactivation centre. Nature.
2012;485(7398):381-5. Epub 2012/04/13. doi: 10.1038/nature11049. PubMed PMID:
22495304; PubMed Central PMCID: PMCPMC3555144.
14. Flavahan WA, Drier Y, Liau BB, Gillespie SM, Venteicher AS, Stemmer-
Rachamimov AO, et al. Insulator dysfunction and oncogene activation in IDH mutant
gliomas. Nature. 2016;529(7584):110-4. Epub 2015/12/25. doi: 10.1038/nature16490.
PubMed PMID: 26700815; PubMed Central PMCID: PMCPMC4831574.
15. Symmons O, Uslu VV, Tsujimura T, Ruf S, Nassari S, Schwarzer W, et al.
Functional and topological characteristics of mammalian regulatory domains. Genome
Res. 2014;24(3):390-400. Epub 2014/01/09. doi: 10.1101/gr.163519.113. PubMed
PMID: 24398455; PubMed Central PMCID: PMCPMC3941104.
16. Sikorska N, Sexton T. Defining Functionally Relevant Spatial Chromatin
Domains: It is a TAD Complicated. J Mol Biol. 2020;432(3):653-64. Epub 2019/12/22.
doi: 10.1016/j.jmb.2019.12.006. PubMed PMID: 31863747.
17. Soler-Oliva ME, Guerrero-Martinez JA, Bachetti V, Reyes JC. Analysis of the
relationship between coexpression domains and chromatin 3D organization. PLoS
Comput Biol. 2017;13(9):e1005708. Epub 2017/09/14. doi:
10.1371/journal.pcbi.1005708. PubMed PMID: 28902867; PubMed Central PMCID:
PMCPMC5612749.
18. Eres IE, Gilad Y. A TAD Skeptic: Is 3D Genome Topology Conserved? Trends
in genetics : TIG. 2021;37(3):216-23. Epub 2020/11/19. doi: 10.1016/j.tig.2020.10.009.
PubMed PMID: 33203573; PubMed Central PMCID: PMCPMC8120795.
19. Hallikas O, Das Roy R, Christensen M, Renvoisé E, Sulic A-M, Jernvall J.
System-level analyses of keystone genes required for mammalian tooth development.
J Exp Zool Part B Mol Dev Evol [in press]. 2020.

20.   Lane N, Martin W. The energetics of genome complexity. Nature.
2010;467(7318):929-34. doi: 10.1038/nature09486.
21.   Jernvall J, Thesleff I. Tooth shape formation and tooth renewal: evolving with the
same signals. Development. 2012;139(19):3487-97. Epub 2012/09/06. doi:
10.1242/dev.085084. PubMed PMID: 22949612.
22.   The Gene Ontology C. The Gene Ontology Resource: 20 years and still GOing
strong. Nucleic Acids Res. 2019;47(D1):D330-D8. Epub 2018/11/06. doi:
10.1093/nar/gky1055. PubMed PMID: 30395331; PubMed Central PMCID:
PMCPMC6323945.
23.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene
ontology: tool for the unification of biology. The Gene Ontology Consortium. Nature
genetics. 2000;25(1):25-9. Epub 2000/05/10. doi: 10.1038/75556. PubMed PMID:
10802651; PubMed Central PMCID: PMCPMC3037419.
24.   de Laat W, Grosveld F. Spatial organization of gene expression: the active
chromatin hub. Chromosome Research. 2003;11(5):447-59. doi:
10.1023/A:1024922626726.
25.   Gene expression in tooth (WWW database), Developmental Biology Programme
of the University of Helsinki, [Internet]. University of Helsinki. 1996-2007. Available
from: http://bite-it.helsinki.fi.
26.   Li CY, Hu J, Lu H, Lan J, Du W, Galicia N, et al. alphaE-catenin inhibits
YAP/TAZ activity to regulate signalling centre formation during tooth development.
Nature communications. 2016;7:12133. Epub 2016/07/14. doi: 10.1038/ncomms12133.
PubMed PMID: 27405641; PubMed Central PMCID: PMCPMC4947177.
27.   Sing T, Sander O, Beerenwinkel N, Lengauer T. ROCR: visualizing classifier
performance in R. Bioinformatics (Oxford, England). 2005;21(20):3940-1. doi:
10.1093/bioinformatics/bti623 %J Bioinformatics.
28.   Debiais-Thibaud M, Metcalfe CJ, Pollack J, Germon I, Ekker M, Depew M, et al.
Heterogeneous conservation of Dlx paralog co-expression in jawed vertebrates. PloS
one. 2013;8(6):e68182. Epub 2013/07/11. doi: 10.1371/journal.pone.0068182. PubMed
PMID: 23840829; PubMed Central PMCID: PMCPMC3695995.
29.   Beverdam A, Merlo GR, Paleari L, Mantero S, Genova F, Barbieri O, et al. Jaw
transformation with gain of symmetry after Dlx5/Dlx6 inactivation: mirror of the past?
Genesis. 2002;34(4):221-7. Epub 2002/11/16. doi: 10.1002/gene.10156. PubMed
PMID: 12434331.
30.   Thomas BL, Tucker AS, Qui M, Ferguson CA, Hardcastle Z, Rubenstein JL, et
al. Role of Dlx-1 and Dlx-2 genes in patterning of the murine dentition. Development.
1997;124(23):4811-8. Epub 1998/01/15. PubMed PMID: 9428417.
31.   Qiu M, Bulfone A, Ghattas I, Meneses JJ, Christensen L, Sharpe PT, et al. Role
of the Dlx homeobox genes in proximodistal patterning of the branchial arches:
mutations of Dlx-1, Dlx-2, and Dlx-1 and -2 alter morphogenesis of proximal skeletal
and soft tissue structures derived from the first and second arches. Dev Biol.
1997;185(2):165-84. Epub 1997/05/15. doi: 10.1006/dbio.1997.8556. PubMed PMID:
9187081.
32.   Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al.
eggNOG 4.5: a hierarchical orthology framework with improved functional annotations
for eukaryotic, prokaryotic and viral sequences. Nucleic Acids Research.
2015;44(D1):D286-D93. doi: 10.1093/nar/gkv1248 %J Nucleic Acids Research.

33.   Fukuoka Y, Inaoka H, Kohane IS. Inter-species differences of co-expression of
neighboring genes in eukaryotic genomes. BMC genomics. 2004;5(1):4. doi:
10.1186/1471-2164-5-4.
34.   Love MI, Huber W, Anders SJGB. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. 2014;15(12):550. doi: 10.1186/s13059-014-
0550-8.
35.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Research. 2015;43(7):e47-e. doi: 10.1093/nar/gkv007 %J Nucleic Acids
Research.
36.   Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. Nature
protocols. 2009;4(8):1184-91. Epub 2009/07/21. doi: 10.1038/nprot.2009.97. PubMed
PMID: 19617889; PubMed Central PMCID: PMCPMC3159387.
37.   Nieminen P, Pekkanen M, Aberg T, Thesleff I. A graphical WWW-database on
gene expression in tooth. Eur J Oral Sci. 1998;106 Suppl 1:7-11. Epub 1998/04/16. doi:
10.1111/j.1600-0722.1998.tb02146.x. PubMed PMID: 9541196.
38.   Carbon S, Ireland A, Mungall CJ, Shu S, Marshall B, Lewis S, et al. AmiGO:
online access to ontology and annotation data. Bioinformatics (Oxford, England).
2009;25(2):288-9. Epub 2008/11/27. doi: 10.1093/bioinformatics/btn615. PubMed
PMID: 19033274; PubMed Central PMCID: PMCPMC2639003.
39.   Zufferey M, Tavernari D, Oricchio E, Ciriello G. Comparison of computational
methods for the identification of topologically associating domains. Genome Biol.
2018;19(1):217. Epub 2018/12/12. doi: 10.1186/s13059-018-1596-9. PubMed PMID:
30526631; PubMed Central PMCID: PMCPMC6288901.
40.   Liu T, Porter J, Zhao C, Zhu H, Wang N, Sun Z, et al. TADKB: Family
classification and a knowledge base of topologically associating domains. BMC
genomics. 2019;20(1):217. Epub 2019/03/16. doi: 10.1186/s12864-019-5551-2.
PubMed PMID: 30871473; PubMed Central PMCID: PMCPMC6419456.
41.   Yu M, Ren B. The Three-Dimensional Organization of Mammalian Genomes.
Annu Rev Cell Dev Biol. 2017;33:265-89. Epub 2017/08/09. doi: 10.1146/annurev-
cellbio-100616-060531. PubMed PMID: 28783961; PubMed Central PMCID:
PMCPMC5837811.
42.   Bengtsson H, Irizarry R, Carvalho B, Speed TP. Estimation and assessment of
raw copy numbers at the single locus level. Bioinformatics (Oxford, England).
2008;24(6):759-67. Epub 2008/01/22. doi: 10.1093/bioinformatics/btn016. PubMed
PMID: 18204055.
43.   Dai M, Wang P, Boyd AD, Kostov G, Athey B, Jones EG, et al. Evolving
gene/transcript definitions significantly alter the interpretation of GeneChip data.
Nucleic Acids Res. 2005;33(20):e175. doi: 10.1093/nar/gni179. PubMed PMID:
16284200; PubMed Central PMCID: PMCPMC1283542.
44.   Andrews S. FastQC: A quality control tool for high throughput sequence data.
2010 [cited 2019]. Available from:
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.
45.   Chen S, Huang T, Zhou Y, Han Y, Xu M, Gu JJBB. AfterQC: automatic filtering,
trimming, error removing and quality control for fastq data. 2017;18(3):80. doi:
10.1186/s12859-017-1469-3.

46.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics (Oxford, England). 2014;30(15):2114-20. Epub
2014/04/01. doi: 10.1093/bioinformatics/btu170. PubMed PMID: 24695404.
47.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics (Oxford, England). 2013;29(1):15-
21. Epub 2012/10/25. doi: 10.1093/bioinformatics/bts635. PubMed PMID: 23104886.
48.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics (Oxford, England). 2015;31(2):166-9. Epub
2014/09/25. doi: 10.1093/bioinformatics/btu638. PubMed PMID: 25260700.
49.   Risueno A, Roson-Burgo B, Dolnik A, Hernandez-Rivas JM, Bullinger L, De Las
Rivas J. A robust estimation of exon expression to identify alternative spliced genes
applied to human tissues and cancer samples. BMC genomics. 2014;15:879. Epub
2014/10/10. doi: 10.1186/1471-2164-15-879. PubMed PMID: 25297679; PubMed
Central PMCID: PMCPMC4298068.
50.   Del Carratore F, Jankevics A, Eisinga R, Heskes T, Hong F, Breitling R.
RankProd 2.0: a refactored bioconductor package for detecting differentially expressed
features in molecular profiling datasets. Bioinformatics (Oxford, England).
2017;33(17):2774-5. Epub 2017/05/10. doi: 10.1093/bioinformatics/btx292. PubMed
PMID: 28481966; PubMed Central PMCID: PMCPMC5860065.
51.   Tusher VG, Tibshirani R, Chu G. Significance analysis of microarrays applied to
the ionizing radiation response. Proceedings of the National Academy of Sciences of
the United States of America. 2001;98(9):5116-21. Epub 2001/04/20. doi:
10.1073/pnas.091062498. PubMed PMID: 11309499; PubMed Central PMCID:
PMCPMC33173.
52.   Ilmjärv S, Hundahl CA, Reimets R, Niitsoo M, Kolde R, Vilo J, et al. Estimating
differential expression from multiple indicators. Nucleic Acids Research.
2014;42(8):e72-e. doi: 10.1093/nar/gku158 %J Nucleic Acids Research.
53.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics (Oxford,
England). 2009;26(1):139-40. doi: 10.1093/bioinformatics/btp616 %J Bioinformatics.
904