Structural dynamics of the functional nonameric Type III translocase export gate

Biao Yuan[1,2,3,4], Athina G. Portaliou[1], Rinky Parakra[1], Jochem H. Smit[1,5], Jiri Wald[2,3,4], Yichen Li[5], Bindu Srinivasu[1], Maria S. Loos[1], Harveer Singh Dhupar[6], Dirk Fahrenkamp[2,3,4], Charalampos G. Kalodimos[7], Franck Duong van Hoa[6], Thorben Cordes[5,8], Spyridoula Karamanou[1], Thomas C. Marlovits[2,3,4,9] and Anastassios Economou[1,9,10]

[1] KU Leuven, Department of Microbiology and Immunology, Rega Institute for Medical Research, Laboratory of Molecular Bacteriology, B-3000 Leuven, Belgium

[2] Centre for Structural Systems Biology (CSSB), Notkestrasse 85, D-22607 Hamburg, Germany

[3] University Medical Center Hamburg-Eppendorf (UKE), Institute for Structural and Systems Biology, Notkestrasse 85, D-22607 Hamburg, Germany

[4] German Electron Synchrotron Centre (DESY), Notkestrasse 85, D-22607 Hamburg, Germany

[5] Molecular Microscopy Research Group, Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

[6] Department of Biochemistry and Molecular Biology, Faculty of Medicine, Life Sciences Institute, University of British Columbia, Vancouver, Canada

[7] Department of Structural Biology, St. Jude Children's Research Hospital, 263 Danny Thomas Place, Memphis, Tennessee 38105

[8] Physical and Synthetic Biology, Faculty of Biology, Ludwig Maximilians-Universität München, Großhadernerstr. 2-4, 82152 Planegg-Martinsried, Germany

[9]For correspondence: Anastassios Economou (tassos.economou@kuleuven.be)

Thomas C. Marlovits (thomas.marlovits@cssb-hamburg.de)

[10] Lead contact

Character count incl. spaces: 44,236

Keywords: EPEC/ export apparatus/ Type III secretion/ SctV-C structure/ protein dynamics

Abstract

Type III protein secretion is widespread in Gram-negative pathogens. It comprises the injectisome with a surface-exposed needle and an inner membrane translocase. The translocase contains the SctRSTU export channel enveloped by the export gate subunit SctV that binds chaperone/exported clients and forms a putative ante-chamber. We probed the assembly, function, structure and dynamics of SctV from enteropathogenic *E.coli* (EPEC). In both EPEC and *E.coli* lab strains, SctV forms peripheral oligomeric clusters that are detergent-extracted as homo-nonamers. Membrane-embedded SctV$_9$ is necessary and sufficient to act as a receptor for different chaperone/exported protein pairs with distinct C-domain binding sites that are essential for secretion. Negative staining electron microscopy revealed that peptidisc-reconstituted His-SctV$_9$ forms a tripartite particle of ~22 nm with a N-terminal domain connected by a short linker to a C-domain ring structure with a ~5 nm-wide inner opening. The isolated C-domain ring was resolved with cryo-EM at 3.1 Å and structurally compared to other SctV homologues. Its four sub-domains undergo a three-stage "pinching" motion. Hydrogen-deuterium exchange mass spectrometry revealed this to involve dynamic and rigid hinges and a hyper-flexible sub-domain that flips out of the ring periphery and binds chaperones on and between adjacent protomers. These motions are coincident with pore surface and ring entry mouth local conformational changes that are also modulated by the ATPase inner stalk. We propose a model that the intrinsic dynamics of the SctV protomer are modulated by chaperones and the ATPase and could affect allosterically the other subunits of the nonameric ring during secretion.

Word count: 250

Authors e-mail addresses:

Biao Yuan: biao.yuan@cssb-hamburg.de

Athina G. Portaliou: athina.portaliou@kuleuven.be

Rinky Parakra: rinky.parakra@kuleuven.be

Jochem H. Smit: jochem.smit@kuleuven.be ; ORCID: 0000-0002-3597-9429

Jiri Wald: jiri.wald@cssb-hamburg.de

Yichen Li: yichen.li@rug.nl

Bindu Srinivasu: bindu.srinivasu@kuleuven.be ; ORCID: 0000-0003-0875-2680

Maria S. Loos: ms.mariam.ms@gmail.com

Harveer Singh Dhupar: harvirsinghdhupar@yahoo.com

Dirk Fahrenkamp: dirk.fahrenkamp@cssb-hamburg.de

Charalampos G. Kalodimos: babis.kalodimos@stjude.org

Franck Duong van Hoa: fduong@mail.ubc.ca

Thorben Cordes: cordes@bio.lmu.de

Spyridoula Karamanou: lily.karamanou@kuleuven.be ; ORCID: 0000-0002-8803-1404

Thomas C. Marlovits: thomas.marlovits@cssb-hamburg.de

Anastassios Economou: tassos.economou@kuleuven.be ; ORCID: 0000-0002-1770-507X
75

INTRODUCTION

The type III protein secretion system (T3SS) is widely used by many Gram-negative

pathogenic or symbiotic bacteria to inject proteins directly into the eukaryotic host cytoplasm

(1, 2). T3SS comprises the "injectisome", a nano-syringe bridging the bacterial and

eukaryotic cytoplasm. Injectisomes comprise four parts (Fig. 1A): a ~32-36 nm-long trans-

envelope "basal body" composed of stacked rings; an inner membrane-embedded "export

apparatus", located at the basal body base; a circular cytoplasmic sorting platform, including

the ATPase complex, that protrudes ~28 nm intro the cytoplasm and has a diameter of ~32

nm and finally, a filamentous needle lying at the top of the injectisome, that protrudes from

within the basal body to the extracellular milieu, and carries a "translocon" at its tip (2-5).

Translocons form pores in host plasma membranes through which effectors are delivered

(6).

The unifying nomenclature Sct (Secretion and cellular translocation) is used for the

conserved components of the pathogenic/symbiotic injectisomes across bacteria (7,

8)(www.stepdb.eu). Species/genus suffixes in subscript are added here to denote Sct

components from other bacteria (e.g. $_{Sf}$Sct for *Shigella flexneri*).

The export apparatus located centrally inside the injectisome is thought to be the

translocase that mediates chaperone/exported client targeting, switching from one client

class to another, client engagement and translocation through the secretion tunnel (2, 9,

10). Five components (SctRSTU and V) of the translocase contain many hydrophobic

transmembrane-like domains, which challenge structural characterization and functional

definition. This is further complicated by the peripheral nature of the association of SctV

with SctRST seen from cryo-ET analysis (3, 5, 11). SctV and U also contain large C-terminal

cytoplasmic domains. The SctV export gate is a nonamer and its C-domain forms a large

cytoplasmic ring with a ~5 nm vestibule connected to the N-terminal membrane part with

narrow stalks (12-15)(Fig. 1A). The isolated cytoplasmic domain of *Shigella flexneri* $_{Sf}$SctV-

C crystalized as a ring structure with an external diameter of 11-17 nm and a height from

the membrane surface of ~5.5 nm (12, 16, 17). Similarly, the flagellar SctV(FlhA)-C forms

rings with an external diameter of 10.1 ± 0.6 nm and a height of 5.1 ± 0.4 nm (18-21). The

C-terminal domain of $_{EPEC}$SctV has been structurally resolved at 4.6 Å resolution using cryo-

EM revealing that nonameric rings form via electrostatic interactions between subunits (14).

The nonameric $_{St}$SctV of *Salmonella typhimurium* has been determined using a high-

throughput cryo-ET pipeline (16, 22). Such a cryo-ET model still lacks sufficient resolution

to determine function and the assembly of its transmembrane channel formed around other

subunits of the export apparatus by 72 transmembrane helices. Notably, the ATPase is

located at a significant distance from SctV [~5 nm; (3, 5, 11)], but may connect to the SctV

cytoplasmic ring indirectly through the elongated SctO inner stalk (23-27). Additional

peripheral bridges formed by other components of the cytoplasmic sorting platform keep it

in place juxtaposed to SctV even when the stalk is missing (3). The dynamics of

ATPase/stalk/SctV C-domain during the translocation cycle remain unknown.

The monomeric cytoplasmic domains of SctV/FlhAs(SctV-C/FlhA-C) expressed as

individual polypeptides interact with chaperone/exported protein complexes in solution (19,

28, 29) with a low dissociation constant (K_d) of ~20 μM (29). Chaperone/exported protein

complexes also interact with the hexameric ATPase of the system in NMR-identified

complexes (30) and with cytoplasmic platform components in pull-down experiments (26,

31, 32). Although the order of chaperone/exported protein interactions with machinery

components during secretion remains debatable, such interactions presumably attract

exported substrates to SctV (9, 29). Then exported substrates dissociate from chaperones,

diffuse to the SctV vestibule and to the inner membrane SctRSTU for export, via a putative

channel running longitudinally through SctRST (10, 13, 33, 34). Recently, the first structure

of an active export apparatus engaged with an exported protein in two functional states was

reported, revealing the complete 800 Å-long secretion conduit and unravelling its critical role

in T3 secretion (13). However, export protein binding and trapping steps are poorly

understood and the precise order and roles of the translocase components are unknown.

cryo-ET revealed that within the assembled injectisome, $_{St}$SctRST sits on top of SctV

in an inverted cone formation that is largely periplasmically exposed and only partially buried

in the inner membrane plane (3, 13, 16). It is kept in this periplasmically protruding state

apparently through contacts with the SctJ lipoprotein and lipids (3). A similar arrangement

was seen in the flagellar T3S (33). The intricate formation of the injectisome rings would

imply possible architectural dependence of one sub-structure on the other and perhaps

suggest a temporal order of addition of each element. Nevertheless, deletion of *sctV* does

not prevent the correct assembly of the remaining injectisome including that of the

cytoplasmic sorting platform, the SctRST inverted cone, that should be internal to SctV ring,

and the SctD that surrounds SctV externally (16, 22). Moreover, the injectisome can be

purified with non-ionic detergents in stable structures in the apparent absence of SctV (13,

15, 35, 36). These data suggested that assembly of SctV may retain a significant autonomy

from that of the other architectural components.

Using the Gram-negative pathogen enteropathogenic *Escherichia coli* (EPEC), a

serious diarrhea threat to children in developing countries (37), we recently reconstituted

T3SS substrate membrane targeting process *in vitro* using inverted inner membrane

vesicles (IMVs) harboring EPEC injectisomes. We revealed a novel mechanism that

governs both membrane targeting of exported clients and switching from translocator

(middle client) to effector (late client) export (9). We now focus on the assembly, structure,

structural dynamics and receptor function of the export gate. We report that nonamers of

$_{EPEC}$SctV and of its C-domain self-assemble *in vivo* in the absence of any other T3S

components, with the C-domain having a major contribution to nonamerization. Membrane-

embedded SctV$_9$ is necessary and sufficient to act as a functional receptor for

chaperone/exported protein complexes in the absence of ongoing secretion and/or other

injectisome components. Specific patches on its C-domain are important for chaperone/

exported protein complexes to bind as demonstrated using immobilized peptide arrays and

confirmed with mutational analysis. The structure of the isolated C-domain polypeptide was

resolved at 3.1 Å resolution by cryo-EM. Negative staining EM analysis indicates that

peptidisc-reconstituted full-length His-SctV$_9$ is a two-domain particle connected by an

equatorial constriction. Both peptidisc-reconstituted His-SctV$_9$ and cryo-EM-resolved C-

domains showed nonameric rings with an internal pore of a ~5 nm diameter. Important

contacts between the protomers are essential for C-domain nonamerization. The C-domain

architecture revealed four sub-domains connected by linkers and compared to other

homologue structures suggest that they undergo an "open" to "closed" "pinching motion.

Hydrogen-deuterium exchange mass spectrometry revealed extensive local intrinsic

dynamics in parts of the structure that would allow rigid body motions and rotations including

those of the hyper-flexible sub-domain 2 (SD2) that is mainly responsible for the pinching

motion. SD2 is a non-conserved structure that overlooks a deep groove that separates two

adjacent protomers and forms a highly negatively charged ridge on either side of the groove.

This landscape defines a wide, multi-valent chaperone trap with multiple binding patches

that are essential for secretion. Direct contacts of the ATPase complex inner stalk SctO

would modulate both local and domain dynamics.

These findings lay the foundations for mechanistic dissection of how structural

dynamic states are coupled to T3S translocase-mediated secretion.

RESULTS

SctV forms multimeric clusters *in vivo*

To analyze SctV membrane assembly *in vivo*, we fused SctV to the C-terminus of eYFP and placed it under tetracycline promoter-control (anhydrotetracycline; AHT)(9). EPECΔ*sctV,* a non-polar deletion strain, is similarly complemented *in vivo* by vector-borne *sctV* (Fig. S1A, lane 3), *eyfp-sctV* (lanes 5-6) and *his-sctV* (Fig S1B, lanes 6-8). eYFP-SctV assembles in distinct focal clusters in the cellular periphery detected by live-cell fluorescence microscopy (Fig. 1B), while eYFP alone shows widespread diffusion with no clusters (Fig. S1C).

To test if eYFP-SctV clustered assembly *in vivo* requires any other T3S components or if it self-assembles, we imaged the *E.coli* lab strain C41, that is devoid of an injectisome, carrying *eyfp-sctV*. C41/*eyfp-sctV* cells synthesized a similar amount of eYFP-SctV as EPECΔ*sctV/eyfp-sctV* (Fig. S2A) that became visible after ~20 min (Fig. 1C; Fig. S2B). This presumably also encompasses the maturation time for the fluorophore (38). eYFP-SctV synthesized in C41, as in EPECΔ*sctV* (Fig. 1B), also showed a distinct peripheral punctate staining (Fig. 1C). This indicated that the apparent eYFP-SctV clustering *in vivo* is an inherent property of SctV.

The punctate staining pattern of eYFP-SctV was characteristic and distinct from that of other fluorescent fusion protein markers that showed either diffuse staining all over the cytoplasm (eYFP) or peripheral staining but of similar dispersed intensity [periplasmic TorA (39); outer membrane BamE (R. Ieva, unpublished) (Fig. S1C-E)]. eYFP-SctV foci fluoresced more strongly than the monomeric inner membrane protein LacY-eYFP (Fig. S1F)(40), suggesting that SctV self-assembles in higher order quaternary states in the inner membrane *in vivo* (see below). Analysis of 10,000-14,000 individual cells showing distinct clusters (Fig. S2B and C), revealed that commonly ~1-4 eYFP-SctV clusters are formed per EPECΔ*sctV*/C41 cell (Fig. S2D) and are distributed along the cellular periphery (Fig. S2E).

SctV natively extracted from membranes is nonameric

To study the oligomeric state and assembly of SctV, we isolated EPEC membranes,

treated them with various concentrations of the non-ionic detergent dodecyl-maltoside

(DDM), separated the resulting complexes by blue native-polyacrylamide gel

electrophoresis (BN-PAGE) and visualized them by immuno-staining with an antibody

against the C-domain of SctV (Fig. 1D).

DDM concentrations ≤0.7% [w/v; 13.2 mM; ~80-fold above the critical micellar

concentration (CMC_{H2O} = 0.17 mM)] did not solubilize discrete SctV-containing complexes

(Fig. 1E, lane 2). At DDM concentrations ≥1.0%, the SctV-containing species migrated as

a tighter band, with an apparent mass of ~700 kDa, consistent with the mass of a stable

nonamer (Fig. 1E, lanes 3-5).

When membranes were solubilized with 0.8% DDM, higher order SctV-containing

complexes with apparent sizes of ~1 MDa immuno-stained with α-SctV-C (Fig. 1E, lane 6,

bracket). These might be sub-assemblies of the injectisome [mass >3 MDa; at least 18

different proteins (8, 35)]. When 0.8% DDM extracts were treated with additional DDM to a

final concentration of 1.8%, a sharp band of $SctV_9$ was obtained (Fig. 1E, lane 7),

suggesting that the assembled nonamer was a pre-existing component of the ~1 MDa

species. Elevated amounts of DDM likely dissociated all other partner subunits but left the

$SctV_9$ intact.

These data suggested that full-length SctV in EPEC membranes assembles in stable,

detergent-extractable, nonameric assemblies that easily dissociate from other injectisome

components depending on detergent concentration.

Self-assembly of SctV nonamers in membranes is injectisome-independent

SctV might nonamerize on demand by enveloping existing membrane-embedded

cores of SctRSTU or other injectisome components (22, 41-43). However, EPEC-derived

SctV$_9$ is stable at up to 1.8% DDM (Fig. 1E) and SctV$_9$ forms clusters *in vivo* even in C41

that is devoid of any injectisome components (Fig. 1C). This raised the possibility that SctV

might self-assemble into nonamers in membranes.

To test this, we over-expressed the *sctV* gene behind a T7 promoter in C41 cells

grown in LB medium (Fig. S3A, lane 3). The membranes were isolated and the membrane-

embedded SctV complexes were characterized by BN-PAGE as above. At low DDM

concentrations (0.4% DDM), C41 extracts contained negligible amounts of DDM-solubilized

SctV (Fig. 1F, lane 2), while at DDM concentrations above 0.6%, discrete nonameric species

of SctV of ~700 kDa were detected (Fig. 1F, lanes 3-7), similar in size to those seen in the

EPEC-derived extracts (Fig. 1E, lanes 3-5). One visible difference though was that SctV$_9$

extraction from C41 membranes initiated at almost half the DDM concentration needed to

extract it from EPEC membranes (Fig. 1F) suggesting that in EPEC, additional components

may further stabilize membrane-embedded and assembled SctV$_9$. The SctV$_9$ species can

also be extracted with other non-ionic detergents (Fig. S3B), implying a stable, pre-existing,

detergent-independent, oligomeric state. SctV$_9$ in C41 membranes was destabilized and

aggregated by heat treatment (Fig. S3C) or once purified from membranes, dissociated into

smaller complexes when treated with an ionic detergent (Fig. S3D), but was very stable at

high concentrations of different non-ionic detergents (Fig. S3E and F).

SctV$_9$ may interact with additional injectisome components (16, 33). We tested

directly whether the export apparatus components SctRSTU form complexes with SctV$_9$.

His-SctRSTU synthesized in C41, gives rise to a complex that is best extracted with 3.6%

digitonin and has a mass of ~240kDa when analyzed by BN-PAGE and immuno-stained

with α-SctU C-domain antibody (Fig. 1G, lane 1). This apparent mass would be consistent

with the R$_5$:S$_4$:T$_1$:U$_1$ stoichiometry anticipated from cryo-EM and MS experiments (3, 10, 16,

33, 34). When co-synthesized with SctV in the same cells, a higher order species of >1MDa

is observed at the low DDM concentration of 0.8% used (Fig. 1H, lane 4). This species runs

as a diffuse band suggesting poor stability and increased dynamics. Once sequentially

treated with 1.8% DDM (w/v) the nonameric species of SctV of ~700kDa appears as a

prominent sharp band, indicating dissociation from other components (Fig. 1I, lane 2).

Collectively, these data suggested that $SctV_9$ stably self-nonamerizes in the absence

of a scaffold provided by other T3S components but weakly associates with pre-assembled

SctRSTU once the latter is present in the membrane.

Contributions of the N- and C-terminal domains of SctV to its nonamerization

SctV comprises a 34.8 kDa N-terminal domain with 8 predicted transmembrane

helices connected by a short ~3 kDa linker to a 37.1 kDa C-terminal cytoplasmic domain.

To determine which SctV segment is important for nonamerization, His-tagged N-terminal

and C-terminal SctV fragments (Fig. 2A, top) were expressed in C41 and their

oligomerization properties were compared to those of SctV after cytoplasmic and membrane

extracts were fractionated by ultra-centrifugation.

Full-length SctV and SctV-N were mainly distributed in the membrane fraction, while

most of SctV-C was detected in the cytosolic fraction (Fig. 2B, lanes 1-3). To exclude co-

sedimentation with membranes due to aggregation and inclusion body formation, the

membrane fraction was further washed with 8M Urea in two steps. Full-length SctV and

SctV-N remained largely in the urea-insoluble membrane fraction (Fig. 2B, top and middle,

lane 7), while SctV-C was extracted by urea completely (Fig. 2B, bottom, lane 7). We

concluded that SctV-N can become membrane integrated in the presence or absence of the

soluble C-domain. Membrane-embedded His-SctV-N extracted with detergent migrates

diffusely at apparent masses of 200-500 kDa in BN-PAGE gels (Fig. 2C, lane 2, bracket),

that are discrete from those of SctV$_9$ (lane 1). This might imply formation of dynamic but

unstable oligomers.

Next, we purified His-SctV-C by metal affinity chromatography and sucrose gradient

ultra-centrifugation (Fig. S4A and B). His SctV-C had a mass of 384.3 (± 3.5) kDa and a

hydrodynamic (Stokes) diameter of 15.14 (± 0.82) nm (shown by analytical gel permeation

chromatography coupled to multi-angle and quasi-elastic light scattering detectors (GPC-

MALS/QELS) (Fig. S4C) or 379.2 (± 2.07) kDa (by native mass spectrometry; Fig. S4D).

Both in good agreement to the theoretical mass of His-SctV-C$_9$ (374.9 kDa) indicating that

the C-domain of SctV inherently self-nonamerizes.

To test the nonamer-promoting role of the C-domain in the context of the full-length

SctV, we introduced different mutations in residues of SctV-C that lie at the protomer-

protomer interface [(12, 14); see below]. In all cases, the mutations resulted in loss of the

nonameric states of both the SctV-C (Fig. 2D; lane 2, one example shown; all mutants in

Fig. S5A) and the SctV full length (Fig. 2E; lane 2, one example shown; all mutants in Fig.

S5B). All oligomerization mutants were functionally defective in the secretion of either middle

(Fig. 2F, lane 3, one example shown; all mutants in Fig. S5C) or late (lane 7 one example

shown; all mutants in Fig. S5C) clients, indicating that self-nonamerization is essential for

secretion and that several residues on the oligomerization surface have similar essential

contributions to oligomerization.

We concluded that SctV-C has a strong tendency to self-nonamerize. This leads to

nonamerization of the full-length SctV, perhaps aided by weak SctV-N oligomeric

assemblies in the membrane.

SctV$_9$ is a necessary and sufficient receptor for chaperone/exported client complexes

SctV interacts with T3S chaperones with or without exported clients and/or the

gatekeeper SctW (9, 29, 44). Nonamerization may be required for high affinity binding since

chaperone/exported substrate complexes bind with low affinity to monomeric flagellar SctV

(29). To determine whether self-assembled $SctV_9$ is functional as an exported protein

receptor, we probed its binding to the dimeric chaperone CesT alone or in complex with its

cognate client Tir, using an *in vitro* affinity determination assay developed previously (9).

IMVs were prepared from *E. coli* C41, harbouring either *sctV* on a plasmid or the empty

vector and were urea-treated to strip away peripheral and non-specifically associated

proteins. The external face of the IMVs corresponds to that of the cytoplasm in the cell; their

lumen represents the periplasmic face. $SctV_9$-containing, C41-derived IMVs were

functionally competent for high affinity saturable binding of CesT or the CesT/Tir complex

(Fig. 2G; Fig. S6A), with K_d values equivalent to those measured in IMVs from EPEC cells

in the presence of the full complement of injectisome components [Fig. 2G; (9)]. In contrast,

IMVs from C41 devoid of SctV showed only a linear, non-specific binding component that

did not yield a measurable K_d (Fig. 2G; Fig. S6A). The receptor function of SctV was further

corroborated using a second chaperone and chaperone/effector complex CesF/EspF (45)

that also bound with high affinity to SctV-containing C41-derived IMVs, but not to C41-

derived IMVs not containing SctV (Fig. 2G; Fig. S6B).

Chaperone/exported protein pairs are likely to share common receptor docking

regions on the $SctV_9$ receptor (29). Attesting to this, incubation of excess of CesF chaperone

(at concentrations of 10-fold K_d) prevents the binding of CesT on IMVs prepared from

C41/*sctV* and *vice versa* (Fig. 2G).

Chaperone/exported middle client complexes and the SctW gatekeeper bind to six

different patches on the SctV C-domain determined using arrays of immobilized peptides

(9). Using the same approach CesT, CesT/Tir, CesF, and CesF/EspF were shown to bind

to patches $\alpha-\varepsilon$ (Fig. S6C-E) that may be used in multi-valent binding (see below). IMVs

prepared from C41 cells expressing *sctVγ* were used as an example. $SctVγ^-$ formed

nonamers as did the wild type (Fig. S5D, lanes 1 and 2) but showed a 3-4-fold affinity

reduction for CesT/Tir and CesF/EspF (Fig. 2G). All patches are functionally important

beyond their *sensu stricto* receptor capacity for *in vivo* secretion of middle and late clients

(9)(Fig. 2F, lanes 4 and 8, the γ derivative is completely defective).

We concluded that $SctV_9$ embedded in C41 membranes is necessary and sufficient

to act as a high affinity receptor for T3S chaperone/exported client complexes *via* its

cytoplasmic domain. The important mechanistic implication of this observation is that the

receptor function of SctV neither depends on the presence of any other T3S components

nor on ongoing translocation.

Purified full length His-SctV$_9$ forms a ring structure

To determine the ultrastructure of full-length $SctV_9$, we synthesized His-SctV in C41.

His-SctV forms nonamers like SctV (Fig. S1G, lanes 1 and 2), was extracted from

membranes with 1.8% Triton X-100, and separated from higher order aggregates on a

sucrose gradient (Fig. 3A and B; Fig. S4E). Fractions enriched in His-SctV$_9$ (determined by

BN-PAGE; Fig. 3C) were collected, pooled and incubated with Ni^{2+}-nitrilotriacetic acid (Ni^{2+}-

NTA) resin. Resin-bound His-SctV$_9$ assemblies were on-bead reconstituted into peptidiscs

using 30% fluorescently labelled peptides (46). To probe if the peptidisc particles were

indeed reconstituted in soluble states in the absence of detergent, we analysed them by

clear native PAGE (CN-PAGE), which separates molecules on the basis of charge, mass

and shape. Silver-stained CN-PAGE gels of His-SctV$_9$-PR revealed a band of ~700-800 kDa

(Fig. 3D) that was also fluorescent, indicating the presence of the bound solubilizing

peptidisc peptides (Fig. 3E). GPC-MALS/QELS revealed that His-SctV$_9$-PR has a mass of

762.8 ± 6.9 kDa (Fig. 3F, red) consistent with a nonameric species bound to ~17 solubilizing

peptides of 4.5 kDa and a hydrodynamic diameter of 22.0 ± 0.98 nm (Fig. 3F, blue) close to

the anticipated dimensions derived from cryo-ET (16).

The His-SctV$_9$-PR was analysed by negative staining EM (Fig. 3G, overview). Class

averaged side views of individual His-SctV$_9$-PR particles (Fig. 3G, bottom left) reveal that

they comprise two distinct domains likely representing the membrane-embedded N-domain

and cytoplasmic C-domain parts of SctV separated by a narrow constriction (9, 12, 16).

Class averaged bottom views reveal a multimeric ring with an inner opening (Fig. 3G, bottom

right).

High resolution structure of the SctV C-terminal domain nonameric ring

Repeated attempts to obtain a high-resolution structure of full length SctV failed to

yield sufficient detail in the membrane part of the ring. We therefore focused our attention

on the cytoplasmic domain. His-tagged SctV-C was purified and its structure determined

using cryo-EM. Both single- and double-ringed species were observed in an approximately

4:1 ratio and were sorted out by using a 3D classification in Relion 3.1 (Fig. S7A; Table S1).

The reconstructed maps of the single-ringed particles with imposed C9 symmetry and

double-ringed particles with imposed D9 symmetry were refined to 3.6 Å and 3.4 Å resolution,

separately (Fig. S7A). The map of the single-rings (map1) is well aligned with parts of the

double ringed map (map2) (Fig. S7B). Therefore, the double-ringed particles were

subtracted into two single-ringed particles and combined with the single-ringed particle

population. Thus, a final 3.1 Å map (map3) using 105,670 particles from 3739 micrographs

was achieved (Fig. S7C-D). Local resolution regions of the EM map are consistent with

predicted disordered regions (Fig. S6E). The model of the regions of 345-415, 462-583, and

626-670 of SctV-C was obtained *de novo* from map3. While, the initial model of $_{sf}$SctV-C$_9$

(PDB: 4a5p, 2x49) (12, 20) was used to guide the model building process for $_{EPEC}$SctV-C

(Fig. 4A).

The SctV-C$_9$ ring contains a charged pore lined with positively charged residues as

are the two outer rims, close to the entry and exit mouths of the pore, at the cytoplasmic-

facing bottom and the membrane-facing top of the structure, respectively (Fig. S8A). The

outer periphery of the ring is characterized by separated upper and bottom negatively

charged bands. Significant conservation is seen in the residues that build the nonamer and

line the inner ring pore (Fig. S8B). In contrast, outwardly facing peripheral surfaces of the

ring are not conserved.

The SctV-C protomer is connected to the transmembrane N-domain by a linker (Fig.

2A; 4B; S6E) and is built of four sub-domains (SD) connected by hinges. SD1 is

discontinuous (aa353-415 and 463-483); SD2 (416-463) is inserted in SD1; SD3 (488-570

and SD4 (571-676) [Fig. 4B; Fig. S6E and S8D (47)]. SD3 is the only domain that lines the

narrowest constriction of the pore and nonamerizes the protomer and is the most conserved

domain, as are some regions of SD1 and SD4 (Fig. S8B).

The nonamer assembles using an interprotomeric interface that is not extensively

hydrophobic and includes multiple intermolecular electrostatic bridges that connect one sub-

domain in one protomer A to a sub-domain in the adjacent protomer B, primarily through

conserved residues in SD3-SD3 and SD3-SD1 contacts. These connect $linker_A$-$SD3_B$

(S347-R532), $SD1_A$-$SD3_B$ (E483/R564) and $SD3_A$-$SD3_B$ (Q488-E503/Y501 and E489-R535)

interfaces (Fig. S8C-G). Moreover, the SD1-SD3 hinge, is positioned between two

protomers and may contribute to ring formation and conformational changes (47). All of

these residues, except E503, were shown by mutagenesis to be crucial for nonamerization

of both SctV-C and full length SctV (Fig. 2D and E; S5A and B; S8F and G) and for secretion

(Fig. 2F; S5C). The oligomerization interface, provided primarily by SD3 residues (Fig. S8D),

is the most conserved part of the protein (Fig. S8B). In contrast, the regions that are away

from the ring pore formed by the conserved SD3, including the outwardly facing SD1 and 2

and parts of SD4 are the least conserved.

The ring pore is built exclusively of SD3 contributed from the 9 subunits (Fig. 4C),

with the highly conserved $K506_{SD3}$, $R510_{SD3}$, and $K549_{SD3}$ residues lining it (Fig. S8E) and

has a diameter of ~5nm. The cytoplasmic entry to the pore is built exclusively of SD4 that

lines the pore entry with its 2β-tip (residues 594-604; β11/12) that forms a hook-like

protrusion (Fig. 4C). SD4 is recessed outwards relative to SD3, thus providing a wider,

funnel-like entry mouth to the pore (Fig. 4C, entry funnel) with a diameter of ~8nm between

hooks. On the other end, the exit mouth of the pore facing the membrane, widens out away

over SD3, is built exclusively by SD1 and together with the top of SD3 form a wide exit mouth

"ledge" (Fig. 4C, exit funnel). SD2 is unrelated to the pore and faces outwards. When a

protomer is viewed head on in a side view of the ring, SD3 and SD4 have a near vertical

placement, like barrel staves, while SD1/2 tilt leftward towards the adjacent subunit by ~30°

from the vertical axis (Fig. 4D). This positioning creates large elongated grooves between

the staves (Fig. 4D, dashed line) including the conserved residues $R532_{SD3}$, $E407_{SD1}$ and

$R564_{SD3}$ (Fig. S8B, middle) and also on either side of the ring "castle embrasures", between

adjacent SD4 domains on the entry funnel at the bottom and between SD1 domains at the

exit funnel at the top(asterisks).

Local structural dynamics of SctV-C$_9$ determined by HDX-MS

Given that the same nonameric SctV ring changes its affinities for clients on the basis

of associated protein regulators (9), we hypothesized that the intrinsic local dynamics of

SctV may underlie its function. To test this, we used a hydrogen deuterium exchange mass

spectrometry pipeline (HDX-MS; Fig. S9) which non-invasively monitors loss or gain of

backbone H-bonds, commonly participating in secondary structure. The labeling reaction is

at low micromolar concentrations and at physiological buffers (48-50). SctV-C was diluted

to ~5 μM into D_2O buffer for various time-points. Samples were acid-quenched, protease-

digested and D-uptake was determined. 233 peptides with good signal/noise ratio yielded

~99.7% primary sequence coverage (Table S2). The D-uptake for a tandem array of

peptides that cover all of the sequence was expressed as a percentage of each one's fully

deuterated control (taken as 100%) (Fig. 4E). Dynamics data are subsequently used to

derive Gibbs free energy of exchange (ΔG_{ex}, kJ mol^{-1}) per residue of a protein's sequence

using our in-house software PyHDX (Fig. 4F)(51, 52). ΔG_{ex} is inversely correlated to the

degree of dynamics in the protein backbone, therefore lower ΔG_{ex} values represent higher

backbone flexibility.

SctV-C displays significant dynamics including regions yielding lower resolution in

the cryo-EM structure and predicted to contain disordered sequence (Fig. S6E; S7D). The

highest level of dynamics is observed in the periphery of the ring (Fig. 4E and F, left and

middle) and the hook at the entry funnel (right). In contrast, the pore and the exit funnel are

rather rigid (Fig. 4E and F, left) as is a region next to the hook (Fig. 4E, middle and right).

The observed elevated dynamics of the ring align along the ridges that flank the side groove,

in which one elevated dynamics region of each protomer apposes the elevated dynamics

region of the adjacent protomer (Fig. 4G).

Intrinsic local dynamics of the SctV-C protomer

To gain insight in the regions of elevated dynamics and rigidity in the four sub-

domains of the SctV protomer we mapped the HDX-MS data (Fig. 4E) onto a protomer

structure derived from cryo-EM in ribbon representation (Fig. 5A). Highly dynamic residues

are in red hues, while the less dynamic ones are in blue hues. SD2 displays the most

elevated and SD3 the more reduced dynamics in the structure. The only regions of SD3 that

display elevated dynamics are its connecting linker to SD4 (Fig. 5A.I, V572-T575) and a

short connection between α5 and α6 (G513-I517) that faces the inner ring surface (Fig.

5A.II). On the other hand, the hinge that connects SD3 to SD1 shows reduced dynamics

(III). In addition to its intrinsic flexibility, SD2 is connected to SD1, into which it is rooted,

using highly flexible loops connecting to the highly flexible ends of two secondary structure

elements (β4 and α3; IV) and this would greatly facilitate rigid body motions of SD2 around

this two-element hinge. The highly flexible β4 of SD1 participates in a 4-stranded sheet,

displaying additional flexible regions (V), that face the side groove of the ring (Fig. 4G). SD4

is rather rigid with three main internal regions of high dynamics: the hook (β10-β11; VI), the

loop$_{N604-R609}$ (VII) and the extreme C-terminal region (I662-A676; VIII). These overall

features were largely retained in the SctV monomer, obtained by analysing the monomeric

SctV(R535A) derivative. Some regions of the monomer displayed more enhanced (e.g.

V477-L504 in SD3 that face the pore) and two more reduced (N345-L357 and D550-L554)

dynamics (Fig. 5B). These differences were visualized in a differential uptake map were

purple hues indicated enhanced rigidity and green hues enhanced dynamics (Fig. 5C).

This pattern of intrinsic dynamics of SctV-C yields several regions of the protein with

residues that have very low ΔG thresholds between open and closed states, explaining their

flexibility (Fig. 5D, coloured circles). This pattern of flexibility was also corroborated by an

orthogonal approach, normal mode analysis, in which low frequency vibrational states for

the atoms of a protein can be derived using coarse grain models derived from PDB

structures and calculating their possible displacement (Fig. 5D, magenta line) (53). In the

context of the nonamer-embedded protomer, flexible regions are located exclusively in

SD1,2 and 4 and are practically absent from SD3.

Sub-domain motions in SctV-C

Local dynamics in critical parts of a structure, as detected in Fig. 5A, B and D for

SctV-C, are commonly accompanied by and permit domain motions (14, 27, 52). All

available structures of SctV-C have been solved in slightly different domain arrangements

revealing a repertoire of domain conformational states that can be acquired by the four sub-

domains of the C-domain. These can be seen to broadly ranging from an "open" to a "closed"

state (12, 14, 18, 27, 29), primarily determined by the motion of the hyper-flexible SD2. The

SctV-C structure resolved here approaches the closed state but its SD2 is closer to its SD1

compared to other structures and its SD2 is more detached from SD4 than in the "closed"

models. To visualize the inherent dynamics of the C-domain sub-domains in the context of

the protomer and the nonamer, we modeled the potential transition motions from an open

to a closed state of SctV-C homologues into a movie (Fig. 5A; Movie S1).

The main rigid body motions modelled are those of the SD2 which makes use of its

hinge connecting it to SD1 (linkers between β4 and β5 and β8 and α3; Fig. 5A.IV; E, left;

S6E) to extend outwards from the ring periphery to yield an "open" state and then retracts

to an "intermediate" position, resting its 2β-tip$_{SD2}$ (residues 433-444; β6/7) against SD1 and

a flexible loop (residues 423-429; IX) against the C-terminus of SD4. This retraction of SD2

is coincident to a rigid body rotation of SD4 around its hinge with SD3 by ~30° (Fig. 5E,

middle). This brings the protruding hook/2β-tip$_{SD4}$ (β10/11; VI) to move towards SD3 and the

inside of the pore. In a second step, SD1 undergoes a rigid body motion along its hinge with

SD3 (Fig. 5E, right, III) and moves towards SD4. This displaces the 2β-tip$_{SD2}$ which now

moves towards the C-terminus of SD4, while the 423-429 loop of SD2 (IX) moves away from

SD4. Concomitantly, the hook reverses its previous motion (Fig. 5E, right; Movie S1).

During the "pinching" motion of SD2 against SD4 in the ring periphery to yield the

closed state, the inner ring surface and the diameter of the pore remain largely unchanged,

except for individual charged and bulky residues of the inner ring side of SD3 (aa K500,

N502, E503, K506, F543, K549; Fig. S8E).

Collectively, this modelling suggested possible rigid body domain transitions derived

from the high-resolution structural snapshots of end states. Our HDX-MS data would now

rationally explain these motions as resulting by the local dynamics of StcV-C and involving

the hinges and the hyper-flexible SD2 (Fig. 5A; Movie S2). Rather than distinct stable

"states", the intrinsic dynamics analysis raises the more possibility of a conformational

ensemble of states.

SctV-C sub-domain dynamics are related to association of interactors

Two main groups of SctV interactors have been analysed to date: chaperones with

or without clients (24, 29, 47, 54) and the inner stalk subunit SctO of the ATPase complex

(27). Binding of chaperone/client complexes occurs mainly in the periphery of the ring

extending to the exit funnel. Binding sites determined by NMR and peptide arrays are

located on either side of the side grooves of the ring (Fig. 6A; S6E; Movie S3). A prominent

chaperone binding site (patches α and β; S6E) is the 4-stranded beta sheet of SD1 that

forms a continuous surface with SD2 residues, with residues like (V393$_{SD1}$, K411$_{SD1}$,

T413$_{SD1}$, K436$_{SD2}$, L437$_{SD2}$ in protomer B and apposed to H455$_{SD2}$, I456$_{SD2}$ in protomer A;

Fig. 6B, top) overlooking the side groove (Fig. 4F). The non-conserved and highly dynamic

SD2 is a major site of chaperone binding on a single protomer but can also contribute binding

interfaces for multivalent inter protomeric binding. Sub-domain motions affect the degree of

exposure of the binding sites (Movie S3). Most patches and NMR binding sites also coincide

wholly or partly with islands of high or very high flexibility, particularly on SD2 and SD1 (Fig.

S6E). Mutagenesis of surface-exposed residues in several of these regions revealed that

they all contribute to chaperone binding affinity and are all essential for secretion [patch γ as

an example; Fig. 2F, lanes 4 and 8 and 2G;(9)].

SctO binds exclusively to SD4, almost vertically against the entry funnel of the ring

[Fig. 6A;(27)] approaching it from the cytoplasm and inserting in sockets formed by the

vertical walls of two adjacent SD4 domains and a flat roof provided by the bottom side of

SD3 (Fig. 6 C; Movie S4). SctO binds on one side directly the SD4 hook of protomer A, and

on the other SD4 residues of protomer B.

DISCUSSION

How injectisomes function remains elusive. With the development of *in vitro*

reconstitution systems (9, 55), guidance from cryo-EM-derived structures and structural

dynamics introduced here for the first time, this issue can now be systematically addressed,

and assembly processes can be dissected away from functional ones (i.e. secretory client

docking and translocation).

The assembly of the SctRSTU translocase and its SctV export gate are key to the

initiation of the downstream export events (22, 42). A major mechanistic challenge for the

assembly of all these subunits is that although they enter the membrane using the universal

membrane-protein integration machineries (*e.g.* Sec/SRP, YidC), SctV needs to

subsequently multimerize and be "plugged" by SctRSTU. While oligomerization in the

presence of other injectisome components *in vivo* has been implied for $_{Ye}$SctV-eGFP (43),

our experiments demonstrate that SctV self-assembles into functional nonamers, driven

largely by its C-domain (Fig. 2C-D; S5A; S8), *in vivo* and *in vitro* even in cells that have no

injectisome components. Therefore, nonamerization is an inherent SctV property. In the

absence of SctV, the rest of the injectisome still assembles and SctRSTU acquires its

periplasmically-exposed location inside the SctD/J rings (16, 22). Thus, SctV, perhaps pre-

assembled, may arrive late to half-pre-assembled injectisomes into which it can integrate to

meet SctRST (16). In EPEC, this may be simply a kinetically controlled stochastic event,

through tight regulation of transcription and/or translation, given that SctV is synthesized in

cells much later than SctRSTU (56).

SctV-C/FlhA-C form nonameric rings in solution via electrostatic interactions (14, 34,

57). Here with an improved map at 3.1 Å resolution we confirmed that multiple interfaces

(Fig. S8C-G) are indeed involved in cytoplasmic ring assembly. Mutational analysis revealed

that these contacts are all essential in forming the nonameric ring both in the isolated C-

domain (Fig. 2D; S5A) and in the full-length protein (Fig. 2E; S5B), as is the interprotomeric

SD1-SD3 hinge involved in conformational changes (47)(Fig. S8C-G). Formation of the C-

domain ring also leads to nonamerization and function of the full-length SctV, with a minor

possible contribution from the transmembrane N-domain. The latter domain cannot be

resolved well by cryo-EM in isolated SctV$_9$ (17, 21) presumably due to the absence of other

T3SS components and/or lipids.

A significant body of data had implied that SctV possesses significant dynamics: the

loss of nonamerization by single residue C-domain mutations (Fig. S5A-B), the reduced

resolution in regions of the cryo-EM map, alternate domain states in the different SctV

structures (12, 14, 47), the significant effects of patch mutations in both the affinity and the

secretory function of SctV (Fig. 2F, G; S5C), the isolation of point mutants in SctV that mimic

the effects of bound interactors (58), the ability of the ring to switch affinities (9) and be

influenced by pH (17) and Ca^{2+} ions (14). The local HDX-MS analysis performed here

provided for the first-time residue level structural understanding of these dynamics (Fig. 4E;

5A). It revealed the coexistence of a very rigid pore-building/nonamerizing sub-domain

attached to two moderately dynamic domains that build the entry and the exit funnels of the

pore. Moreover, it revealed a hyper-dynamic, non-conserved, outward facing sub-domain

that controls chaperone access to the ring periphery. Flexible and inflexible hinges between

the sub-domains and flexible elements like the entry funnel hook and the inner pore exposed

dynamic region of SD3 (Fig. 5A) suggest possibilities for motions and functional roles. In

common with multiple other examples (59-61), the dynamics described here are intrinsic,

and reflect an inherent property of the monomeric polypeptide (Fig. 5B). As such, their

exploitation by binding ligands does not require high energy input.

Our data corroborated and extended the major role of SctV as a chaperone/exported

client receptor (9, 29), via its C-domain. T3SS chaperones are structurally distinct and

flexible and atomistic detail of their binding in the context of the SctV nonamer is missing.

Nevertheless, localization of binding interfaces on the SctV-C$_9$, hints to important functional

regions made available between adjacent protomers, in the context of the nonamer (Fig. 6A

and B, light green; Movie S3). Several residues in these regions are important for

chaperone/gatekeeper binding in structures solved by NMR (29) and in genetic analyses

(23, 24, 62-64)(Fig. S6E; green; Table S3). Using different T3SS-chaperones we

demonstrated that binding sites on SctV-C are universal and shared (Fig. 2G). The absence

of conservation, charged surfaces with excessive dynamics would allow the hyper-flexible

SD2 to co-evolve to specifically bind the multiple different T3SS chaperones of a particular

organism. The rather flat nature of this receptor landscape and its dynamics make it

structurally feasible to accommodate very large chaperone/client complexes, while dimeric

chaperones and/or clients may additionally secure docking on multiple regions of adjacent

protomers. NMR structures suggested that chaperone complexes insert their binding

elements inside both the side groove of the ring but also in the crevice between SD2-SD4

uncovered when SD2 moves outward to the open state [(29); Fig. 6B, red arrow]. SD2

interactions with chaperones may also control their release or accessibility since chaperone

binding sites are hidden when SD2 retracts (Movie S3, green) (14, 27).

The identified chaperone binding regions on SctV-C have important mechanistic

implications as they demonstrate that while targeting and subsequent secretion are

structurally and biochemically separable, these processes must also be somehow coupled

since when the receptor site γ is mutated alone, it leads to a modest reduction in affinity but

completely abrogates secretion [Fig. 2F-G; (9)].

Multiple limited area binding sites on receptors, dynamic non-folded polypeptides and

excessive dynamics are common themes in chaperone-non-folded client recognition and

delivery systems (52, 65, 66). Multi-valent binding secures synergistic low nM affinities, from

binding sites that individually possess only μM binding strength and the non-folded client

does not face a severe entropic penalty when bound as it retains most of its sequence

unstructured. A powerful mechanistic reason for these evolutionary choices is their facile

loss of the nM binding for the chaperone and/or client and the release of the latter by minor

conformational modulation of the receptor. This "disruptive" role is anticipated here to be

played by the SctO inner stalk of the ATPase since it penetrates between two adjacent SctV-

C subunits at SD4 and binds the hook [Fig. 6C; (27)] and will undoubtedly affect the intrinsic

dynamics of both subunits and the wider ring. Moreover, SctO binding might stabilize the

SD2 open state [MovieS4; (27)]. As the machinery has nine binding sites for SctO and

chaperone/clients but only one SctO subunit attached to one hexameric ATPase, a rotary

mechanism would disrupt consecutively bound multiple chaperone/client complexes.

Because the receptor is intrinsically dynamic, the ATP expenditure for client release may be

minor.

Future analysis will be necessary to dissect the structural basis of these events and

define the path that the exported clients follow upon their dissociation from their chaperones.

As SD2 retraction coincides with SD4 hook rotation towards the central pore, this motion of

and those of highly conserved pore-lining residues $K506_{SD3}$, $R510_{SD3}$ and $K549_{SD3}$, might

also guide and translocate clients into the pore (Fig. S8 D and E; Movie S1). Both the hook

and the conserved pore-lining residues affect substrate switching (17). The development

here of tools like the peptidisc-reconstituted $SctV_9$ pave the way for functional reconstitution

of the inner membrane translocase to decipher translocase mechanism and dynamics.

Materials and Methods

For the complete list of strains, plasmids, mutants, primers, buffers, antibodies, see

the supplementary material.

Cell growth, induction of gene expression, *in vivo* secretion

EPEC strains were grown in the optimized M9 medium (67) (37°C; 6 h). Plasmid gene

expression was induced from the tetracycline promoter (OD_{600}=0.3; AHT; 2.5 ng/mL; 3 h or

as indicated). Cells were harvested (5,000 x g; 20 min; 4°C; Sigma 3-16KL; rotor 11180);

the spent medium was TCA-precipitated (20% w/v) and resuspended with 50 mM Tris-HCl

(pH 8.0) in volumes adjusted according to OD_{600}. An equal number of cells or supernatant

volumes derived from an equal number of bacterial cells, were analyzed by SDS-PAGE, and

immuno- or Coomasie Blue- stained.

Live-cell imaging

Cells were imaged on inverted fluorescence microscope (Olympus IX-83) with a 1.49

NA oil-immersion objective (Olympus UAPON 100x) and ET442/514/561 Laser triple band

set filtercube (69904, Chroma). The excitation light was provided by a 514 nm laser

(Sapphire LP, Coherent) and the fluorescence emission was collected on an EM-CCD

Camera (Hamamatsu C9100-13). Camera frames were collected using CellSens software

(Olympus) for multi-position acquisition. Brightfield images were flattened by subtracting

dark offset and dividing by an empty brightfield image. Fluorescence image were corrected

for illumination beam profile and dark count. The brightfield images are segmented to identify

cell positions by convolutional neural networks with the U-Net architecture (68-70). Next,

individual cells were analyzed with the software ColiCoords (71). Binary objects from

segmented images were filtered based on morphological features (size, ellipse axis) to

select only single planktonic *E. coli* cells. Next, the obtained cell objects were further filtered

in consecutive steps 1) Shape of the binary image; 2) radius of the cell measured from the

brightfield image (C41; 280 nm $\leqslant$ r $\leqslant$ 437 nm, EPEC; 467 nm $\leqslant$ r $\leqslant$ 568 nm); 3) shape of

the brightfield image radial distribution; 4) fluorescence intensity, which resulted in removal

of out-of-focus cells and overexpressing cells. To localize fluorescent foci, first the cytosolic

background was obtained by filtering the images with a spatial median filter (72) (kernel size

5 pixels). Foci were identified in the background-subtracted images by a local maximum

filter (73). To remove aggregates and inclusion bodies from the identified peaks, the Zernike

moments (74) for each peak were determined (R_2^0), and by thresholding only Gaussian-

shaped peaks were selected.

Membrane solubilization and BN-PAGE analysis

Cell pellets were resuspended in Buffer A supplemented with 1 mM MgCl$_2$, 2.5 mM

PMSF, and 50 mg/mL DNase and lysed by using French press (1,000 psi; 6 passes). The

unbroken cells were removed by centrifugation (3,000 x g; 4°C; 5 min; Sigma 3-16KL; rotor

11180). The membrane fraction was pelleted down by high-speed centrifugation (100,000 x

g; 4°C; 30 min; 45Ti rotor; Optima XPN-80, Beckman Coulter). The membrane pellet was

resuspended in solubilization buffer, homogenized with a Dounce homogenizer to a final

protein concentration of ~40 mg/mL. Membrane proteins were extracted by incubating with

non-ionic detergent as indicated (4°C; 1 h). Following ultra-centrifugation (100,000 x g; 4°C;

15 min; rotor TL-100; OptimaMax-XP, Beckman-Coulter), solubilized membrane protein

samples (20µl extract with BN-PAGE loading buffer, 10% glycerol) were loaded on 3-12%

gradient Bis-Tris NativePAGE^{TM} precast protein gels (Invitrogen) and subjected to BN-

PAGE electrophoresis as described previously (75) with little modification. BN-PAGE

electrophoresis was performed with anode buffer and cathode buffer 1 using XCell^{TM}

SureLock^{TM} Mini-Cell (approximately 2.5 hours, 4°C, 100V) until the blue running front has

moved about one-third of gel. Then cathode buffer 1 was replaced with cathode buffer 2 for

electrophoresis (15 hours, 4°C, 70V).

Reconstitution of His-SctV$_9$ in peptidiscs

The peptidisc reconstitution of His-SctV was conducted 'on-bead' as described

previously (46) with some modifications. Crude membrane suspensions were derived from

C41 cells overexpressing His-*sctV* by French press (6 passes; 1,000 psi; pre-cooled with

ice). A total of 10 mL suspension (200 mg/mL) was mixed with 200 μl 100 x protease inhibitor

cocktail solution (Sigma Aldrich, Cat. S8830), solubilized in Buffer F (supplemented with 1.8%

Triton X-100) and incubated (1 h; 4°C). Insoluble aggregates were removed by

centrifugation (20,000 x g, 30 min, 4°C; Sigma 1-16K). Solubilized membrane proteins (1

mL; total of 400 mg) were loaded on a 30 mL 15-35% w/v linear sucrose gradient prepared

as previously described (76). Sucrose solutions were made in Buffer G and 25 x 89 mm 38.5

mL open-up polyallomer tubes were used (Beckman Coutler). Ultra-centrifugation was

performed (160,000 x g; 16 hours; 4°C; SW 32 Ti rotor; Optima XPN-80, Beckman Coulter)

and 1mL fractions were collected and analyzed by BN-PAGE. Fractions containing SctV$_9$

without higher order aggregates of SctV were selected and incubated with 1.5 mL of Ni^{2+}-

NTA resin (Qiagen) pre-equilibrated in Buffer F (6 h; 4°C). The Ni^{2+}-NTA beads were

transferred to a gravity column, washed with 100 column volumes (CV) of Buffer F

supplemented with 0.9% Triton X-100 and 10 CV of Buffer H supplemented with 0.02%

DDM, sequentially. Post-washing, 1 CV of Assembly Buffer conta

ining 0.8 mg/mL NSPr mix (1 fluorescent: 2 non-fluorescent NSPr; Peptidisc, an

amphipathic bi-helical peptide; PEPTIDISC BIOTECH) in 20 mM Tris-HCl pH 8.0) was

added to the beads and incubated (2 h; 4°C). Following washing with buffer H (10 CV), His-

SctV$_9$-PR was eluted in 0.5 mL fractions (5 mL of Buffer H + 300 mM imidazole). The elution

fractions were loaded onto a Superose™6 10/300 GL column previously equilibrated with

buffer L using an ÄKTA Pure system (GE Healthcare). The purified His-SctV$_9$-PR Peptidisc

complex was subsequently analyzed by negative-stain electron microscopy.

Purification of His-SctV-C$_9$

BL21(DE3) cells overexpressing *his-sctV-C* were resuspended in buffer A and lysed

by French press 1,000 psi; 5-6 passes; pre-cooled with ice). Insoluble material was removed

by centrifugation (100,000 x g; 30 min; 4°C; 45 Ti rotor; Optima XPN-80; Beckman Coulter).

The cell lysate supernatant was passed through a Ni^{+2}-NTA resin column (Qiagen), and

protein was purified following the manufacturer's instructions. Eluted His-SctV-C was loaded

on a 13 mL 15-35% linear sucrose gradient and centrifuged (as described above). Fractions

were loaded on 3-12% gradient Bis-Tris NativePAGE^{TM} precast protein gels (Invitrogen) and

subjected to CN-PAGE electrophoresis. CN-PAGE electrophoresis was performed using

XCell^{TM} *SureLock*^{TM} Mini-Cell (approximately 15 hours, 4°C, 15 mA) and gels stained with

Coomasie Blue. Fractions containing mainly His-SctV-C$_9$ were further purified on a

Superdex200 10/300 GL column previously equilibrated with buffer M using an ÄKTA Pure

system (GE Healthcare). His-SctV-C$_9$ complexes were subsequently analyzed by cryo-EM.

Negative Staining and EM

Sample preparations with the concentration of 20-100 ng/μl were applied to glow-

discharged, carbon-coated copper grids, and stained with a 2% (wt/vol) solution of uranyl

acetate, and examined under a FEI Talos L120C (120 KV) microscope equipped with a 4k

× 4k Ceta^{TM} 16M camera. The micrographs were acquired at 92,000X magnification with 2

seconds exposure. Particle picking and 2D classification were performed using Relion

3.1(77)

Single-Particle cryo-EM and Data processing

SctV-C was vitrified on Quantifoil 1.2/1.3. Briefly, 4 μl of the sample was applied onto

glow discharged grids and allowed to disperse for 0.5-2min. The grids were blotted for the

4-7s set at 100% humidity and plunge-frozen in liquid propane/ethane mixture cooled with

liquid nitrogen to about minus 180 -190 °C by using a Vitrobot Mark V. Vitrified specimen

were imaged on a FEI Titan Krios operating at 300 kV and equipped with a field emission

gun (XFEG) and a Gatan Bioquantum energy filter. Movies consisting of 25 frames were

automatically recorded using FEI EPU software and the K2 Summit camera resulting in 0.55

Å per physical pixel. For individual frames, an electron dose of 1.65 e$^-$/Å^2 was used,

corresponding to a cumulative electron dose of 41.25 e$^-$/Å^2 equally distributed over a 5-sec

movie. Movies were recorded at 0-4 µm defocus. Samples for diameter measurements were

recorded with LEGINON13 on a FEI Polara (300 kV) equipped with a field emission gun

(FEG) and a Gatan CCD Camera (UHS 4000). Single-particle reconstructions were

performed using Relion 3.1 (77). 3739 Movies were motion-corrected, dose-weighted, and

binned by 2. The CTF estimation of the resulting micrographs was determined using

CTFFIND4 (78). Particles were picked from the motion-corrected micrographs using

crYOLO (79) trained with a sub-set of manually picked particles. Particles were extracted

into 256 x 256 boxes and subsequently binned by 2 for several rounds of 2D classification.

A cleaned dataset was obtained by re-extraction and aligned to a rotationally averaged

structure. 3D classification with C1 symmetry was performed to sort out the single-ringed

particles and double-ringed particles. Focused refinements with applying C9 for single-

ringed particles and D9 symmetry for double-ringed particles were performed. After

converged refinements, per-particle CTF and Bayesian polishing were used to generate new

data sets for another round of focused refinements. Overall gold-standard resolution: Fourier

shell correlation (FSC=0.143) and local resolution were calculated with Relion3.1. The

double-ringed particles were subtracted into two single-ringed particles and combined with

single-ringed particles for the refinements for generating a better single-ringed map.

Model building, refinement and validation

Ab initio model building was performed with Coot (v0.9-beta)(80). Interactive refinement

against the cryo-EM map density was performed with ISOLDE (v.1.1.0)(81), a molecular

dynamics-guided structure refinement tool within in ChimeraX (v.1.1)(82). The resulting

coordinate file was further refined with Phenix.real_space_refine (v.1.18-6831)(83) using

reference model restraints, strict rotamer matching and disabled grid search. Model

validation was carried out using MolProbity server (84) and EMRinger (85) within the Phenix

software package. More details can be found in Table S1.

SctV sub-domain motions analysis

EPECSctV-C was compared to its homologs with existing PDB files using ChimeraX 1.0

(https://www.rbvi.ucsf.edu/chimerax/). The conformational states were then classified into

different classes according to the distance between SD2 and SD4. Two models were

generated after structural alignment using chimeraX 1.0 that had apparently the most

extreme distances in the conformational states of SD2 and SD4. To build the open and close

state of the EPECSctV-C structural model we used SWISS-MODEL

(https://swissmodel.expasy.org/) and PDB: 6wa6 *Chlamydia pneumoniae* and PDB: 2x49

Salmonella typhimurium were chosen as the template for open and close state respectively.

Then the open and close state of SctV-C, and the intermediate state of SctV-C obtained

from cryo-EM here were morphed using ChimeraX to see how the sub-domains might move.

The movie was recorded in ChimeraX and further processed using iMOVIE

(https://www.apple.com/imovie/).

HDX-MS experimental procedure, data collection and analysis

D exchange reaction and Quenching

The purified SctV-C_9 protein was diluted to 5 µM for mass spectrometry experiments using

H_2O Buffer (25 mM Tris pH 8; 25 mM KCl). For HDX-MS experiments, 4 µl of SctV-C_9 in

H_2O buffer were mixed with 46 µl of the D_2O buffer (25 mM Tris pD 7.6; 25 mM KCl) at 30°C

and incubated at various time intervals (10s, 30 sec, 1min, 5 min, 10 min, 30 min, 100 min)

before being added to 50 µl of quench solution (8M Urea; 0.1% DDM; 5 mM TCEP pH 2.5).

A 100-µL Hamilton syringe was used for sample injection (100 µl). The mobile-phase flow

paths were held at 0.5 °C using a three-valve unit (Trio Valve, Leap Technologies)

constructed with a custom thermal chamber such that on-line protein-digestion, peptide

desalting and reversed-phase HPLC separation are performed prior to infusion into the ESI

ion source of the mass spectrometer. Loading of sample, digestion, and desalting (3 min)

was driven with an isocratic HPLC pump (IPro-500, IRIS Technologies, Lawrence, KS) at a

flow rate of 100 µL min-1 through the 50-µL sample loop, the immobilized nepenthesin

column (Affipro 2.1 mm ID x 20 mm length, Cat.No.: AP-PC-004), across a VanGuard C18

Pre-column, (130 Å, 1.7 mm, 2.1 x 5 mm, Waters), and out to waste. After isolation of the

enzyme column from the flow path, gradient elution from the trap across a C18 analytical

column (130 A°, 1.7 mm, 1 x 100 mm, Waters) was carried out with a capillary-scale HPLC

pump (Agilent 1100, Palo Alto, CA). The flow rate was held constant at 40 µL min-1 while

the composition of the mobile phase was increased from 0% ACN containing 0.23% (FA) to

40% ACN containing 0.23% FA over 12 min gradient. Following gradient elution, mobile-

phase composition was increased over a 1-min period to 90% ACN, 0.1% FA and held at

that composition for 6 min prior to reduction over 1 min to 100% H_2O, 0.1% FA. All HPLC

connections were made with 1/16 in. × 0.05 in. PEEK tubing.

Peptide identification and HDX data analysis

To map the peptides of SctV-C_9, identification MS runs were performed, where 5 µM of

protein in H_2O buffer was used. The sample was quenched as described above and

analysed in the MS^E acquisition mode in a SYNAPT G2 ESI-Q-TOF mass spectrometer over

the m/z range 100-2,000 Da with the collision energy was ramped from 15 to 35 V.

HDX-MS experiments were performed on SYNAPT G2 ESI-Q-TOF mass spectrometer

using a capillary voltage 3.0 kV, sampling cone voltage 20 V, extraction cone voltage 3.6 V,

source temperature 80°C, desolvation gas flow 500 L/h at 150°C. The primary sequence of

SctV-C$_9$ as a search template was used to identify SctV-C$_9$ peptides identification using

ProteinLynx Global Server (PLGS v3.0.1, Waters, UK). HDX experiments were then

analyzed in Dynamx 3.0 (Waters, Milford MA) software. All the other parameters were as

previously described (52). Full deuteration controls were obtained by incubating SctV-C$_9$ in

D$_2$O buffer overnight at 30°C. D-uptake (%) was calculated using the full deuteration control

D-uptake values. The data has not been corrected for back exchange and is represented

either as absolute D values or as a percent of the full deuteration control (86).

Gibbs free energy and normal mode analysis

Gibbs free energy of local unfolding (ΔG_{ex}, kJ mol^{-1}) was determined assuming the

Linderstrøm-Lang model of H/D exchange (87), using the PyHDX software (51). Normal

modes of SctV were calculated using the WebNM@ web server (53), using the SctV cryo-

EM structure as input (PDB XXX). Per-residue displacement was averaged across all protein

chains and the first 6 non-trivial normal modes were summed with equal weights to obtain

the final NMA displacement.

Miscellaneous

In vivo protein secretion, IMVs preparation, peptide arrays, SEC-MALS- QELS analysis and

determination of equilibrium dissociation constants (K_d) were as described (9).The details of

peptide array and Kd determination can be also found in supplementary methods. Ni^{2+}-NTA

purification was according to the manufacturer's instructions (Qiagen). Purification of

plasmids, PCR, digestion, and DNA fragments were done using Wizard® DNA Clean-Up

System from Promega. DNA fragment ligation was done by using T4 DNA ligase as per the

manufacturer's protocol (Promega). Sequencing of the genetic constructs was performed

by Macrogen Europe. Oligos were synthesized by Eurogentec.

ACCESSION NUMBERS:

PDB: 7OSL

EMD-13054

Authors' contributions:

B.Y. performed *in vivo* and *in vitro* assays, molecular cloning, protein purification, GPC-

MALS-QELS, reconstituted SctV in peptidiscs and performed negative staining, and cryo-

EM data processing; A.G.P. performed molecular cloning, membrane binding assays,

peptide array analysis, GPC-MALS, *in vivo* and *in vitro* assays; R.P. and S.K. performed

HDX-MS experiments and analysis; Y.L. and J.S. performed live-cell imaging; J.S.

performed and supervised free energy calculations of the HDX-MS data and developed

analysis software; J.W. performed negative staining and cryo-EM data collection and

processing, and analyzed data; D.F. built the atomic model; H.S.D. set up peptidisc

reconstitution; B.S. performed native MS; M.S.L contributed in molecular cloning, antibody

preparation, gene knockouts. S.K. trained, designed and supervised protein purification,

detergent extraction and reconstitution, biochemical and biophysical experiments. T.C.M.

supervised electron microscopy and structure determination experiments and analysis,

C.G.K and T.C. contributed to data analysis, A.E., A.G.P., and B.Y. wrote the first draft of

the paper with feedback from all the authors. A.E., T.C.M. and S.K. conceived and

supervised the project.

Acknowledgements

We thank K. C. Tsolis for preliminary SctV C-domain characterization and S.Krishnamurthy

for help with analysis of HDX-MS data; C. Robinson (TorA-GFP), and R. Ieva (pDR*bamE-*

gfp) for constructs. Work in our labs was supported by grants (to AE and TCM): T3RecS

[#G002516N; Fonds Wetenschappelijk Onderzoek (FWO); https://www.fwo.be/ and "I 2408-

B22" from the Austrian Science Fund (FWF); https://www.fwf.ac.at/en/]; (to AE): DIP-BiD

(#AKUL/15/40-G0H2116N; Hercules/FWO); RUN (#RUN/16/001 KU Leuven); the

FWO/F.R.S.-FNRS "Excellence of Science - EOS" programme grant #30550343);

PROFOUND (Protein folding/un-folding and dynamics; W002421N; WoG/FWO) and (to AE

and SK); C1 (FOscil; KU Leuven). This project was supported by funds through the Behörde

für Wissenschaft, Forschung und Gleichstellung of the city of Hamburg available to TCM.

B.Y. and Y.L. were Chinese Scholarship Council doctoral fellows. JHS is a PDM/KU Leuven

fellow. The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript. Image data processing has been performed at the

German Electron Synchrotron Centre (DESY) using the High-Performance Computing

Cluster.

Data Availability

All relevant data are within the manuscript and its Supporting Information files and are fully

available without restriction. Coordinate and associated volume and metadata have been

deposited in the PDB and EMDB respectively. SctV-C EMD-xxx / PDB xxx

Competing interests

The authors declare they have no competing financial interests or other conflicts of interest

References

1.   Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, et al. Assembly, structure,
function and regulation of type III secretion systems. Nature reviews Microbiology. 2017.
2.   Galán JE, Lara-Tejero M, Marlovits TC, Wagner SJArom. Bacterial type III secretion systems:
specialized nanomachines for protein delivery into target cells. 2014;68:415-38.
3.   Hu J, Worrall LJ, Vuckovic M, Hong C, Deng W, Atkinson CE, et al. T3S injectisome needle complex
structures in four distinct states reveal the basis of membrane coupling and assembly. Nature microbiology.
2019.
4.   Lara-Tejero M, Galán JEJEP. The Injectisome, a Complex Nanomachine for Protein Injection into
Mammalian Cells. 2019;8(2).
5.   Butan C, Lara-Tejero M, Li W, Liu J, Galan JE. High-resolution view of the type III secretion export
apparatus in situ reveals membrane remodeling and a secretion pathway. Proceedings of the National
Academy of Sciences of the United States of America. 2019;116(49):24786-95.
6.   Park D, Lara-Tejero M, Waxham MN, Li W, Hu B, Galán JE, et al. Visualization of the type III secretion
mediated Salmonella–host cell interface using cryo-electron tomography. 2018;7:e39514.
7.   Hueck CJ. Type III protein secretion systems in bacterial pathogens of animals and plants.
Microbiology and molecular biology reviews. 1998;62(2):379-433.
8.   Portaliou AG, Tsolis KC, Loos MS, Zorzini V, Economou A. Type III secretion: building and operating
a remarkable nanomachine. Trends in biochemical sciences. 2016;41(2):175-89.
9.   Portaliou AG, Tsolis KC, Loos MS, Balabanidou V, Rayo J, Tsirigotaki A, et al. Hierarchical protein
targeting and secretion is controlled by an affinity switch in the type III secretion system of enteropathogenic
Escherichia coli. The EMBO journal. 2017;36(23):3517-31.
10.   Miletic S, Goessweiner-Mohr N, Marlovits TC. The Structure of the Type III Secretion System Needle
Complex. Current topics in microbiology and immunology. 2020;427:67-90.
11.   Berger C, Ravelli RBG, Lopez-Iglesias C, Kudryashev M, Diepold A, Peters PJ. Structure of the
Yersinia injectisome in intracellular host cell phagosomes revealed by cryo FIB electron tomography. Journal
of structural biology. 2021;213(1):107701.
12.   Abrusci P, Vergara-Irigaray M, Johnson S, Beeby MD, Hendrixson DR, Roversi P, et al. Architecture
of the major component of the type III secretion system export apparatus. Nature structural & molecular
biology. 2013;20(1):99-104.
13.   Sean Miletic DF, Nikolaus Goessweiner-Mohr, Jiri Wald, Maurice Pantel, Oliver Vesper, Vadim Kotov,
Thomas C. Marlovits. Substrate-engaged type III secretion system structures reveal gating mechanism for
unfolded protein translocation. BioRxiV. 2020.
14.   Majewski DD, Lyons BJE, Atkinson CE, Strynadka NCJ. Cryo-EM analysis of the SctV cytosolic
domain from the enteropathogenic E. coli T3SS injectisome. Journal of structural biology. 2020;212(3):107660.
15.   Worrall LJ, Hu J, Strynadka NCJ. Aligning the Symmetry of the Type III Secretion System Needle
Complex. Journal of chemical information and modeling. 2020;60(5):2430-5.
16.   Butan C, Lara-Tejero M, Li W, Liu J, Galán JE. High-resolution view of the type III secretion export
apparatus in situ reveals membrane remodeling and a secretion pathway. 2019:709592.
17.   Matthews-Palmer TRS, Gonzalez-Rodriguez N, Calcraft T, Lagercrantz S, Zachs T, Yu XJ, et al.
Structure of the cytoplasmic domain of SctV (SsaV) from the Salmonella SPI-2 injectisome and implications
for a pH sensing mechanism. Journal of structural biology. 2021;213(2):107729.
18.   Terahara N, Inoue Y, Kodera N, Morimoto YV, Uchihashi T, Imada K, et al. Insight into structural
remodeling of the FlhA ring responsible for bacterial flagellar type III protein export. Science advances.
2018;4(4):eaao7054.
19.   Saijo‐Hamano Y, Imada K, Minamino T, Kihara M, Shimada M, Kitao A, et al. Structure of the
cytoplasmic domain of FlhA and implication for flagellar type III protein export. Molecular microbiology.
2010;76(1):260-8.
20.   Worrall LJ, Vuckovic M, Strynadka NC. Crystal structure of the C‐terminal domain of the Salmonella
type III secretion system export apparatus protein InvA. Protein Science. 2010;19(5):1091-6.
21.   Lucas Kuhlen SJ, Jerry Cao, Justin C. Deme, Susan M. Lea. Nonameric structures of the cytoplasmic
domain of FlhA and SctV in the context of the full-length protein. bioRxiv. 2020.
22.   van der Velde JHM, Oelerich J, Huang J, Smit JH, Jazi AA, Galiani S, et al. Corrigendum: A simple
and versatile design concept for fluorophore derivatives with intramolecular photostabilization. Nature
communications. 2017;8:16173.
23.   Hara N, Morimoto YV, Kawamoto A, Namba K, Minamino TJJob. Interaction of the extreme N-terminal
region of FliH with FlhA is required for efficient bacterial flagellar protein export. 2012;194(19):5353-60.
24.   Ibuki T, Uchida Y, Hironaka Y, Namba K, Imada K, Minamino TJJob. Interaction between FliJ and
FlhA, components of the bacterial flagellar type III export apparatus. 2013;195(3):466-73.
25.   Lee P-C, Zmina SE, Stopford CM, Toska J, Rietsch AJPotNAoS. Control of type III secretion activity
and substrate specificity by the cytoplasmic regulator PcrG. 2014;111(19):E2027-E36.

26.   Majewski DD, Worrall LJ, Hong C, Atkinson CE, Vuckovic M, Watanabe N, et al. Cryo-EM structure of
the homohexameric T3SS ATPase-central stalk complex reveals rotary ATPase-like asymmetry.
2019;10(1):626.
27.   Jensen JL, Yamini S, Rietsch A, Spiller BW. "The structure of the Type III secretion system export
gate with CdsO, an ATPase lever arm". PLoS pathogens. 2020;16(10):e1008923.
28.   Bange G, Kümmerer N, Engel C, Bozkurt G, Wild K, Sinning I. FlhA provides the adaptor for
coordinated delivery of late flagella building blocks to the type III secretion system. Proceedings of the National
Academy of Sciences. 2010;107(25):11295-300.
29.   Xing Q, Shi K, Portaliou A, Rossi P, Economou A, Kalodimos CG. Structures of chaperone-substrate
complexes docked onto the export gate in a type III secretion system. Nature communications. 2018;9(1):1773.
30.   Stefl S, Nishi H, Petukh M, Panchenko AR, Alexov E. Molecular mechanisms of disease-causing
missense mutations. Journal of molecular biology. 2013;425(21):3919-36.
31.   Romo-Castillo M, Andrade A, Espinosa N, Monjaras Feria J, Soto E, Diaz-Guerrero M, et al. EscO,
a functional and structural analog of the flagellar FliJ protein, is a positive regulator of EscN ATPase activity of
the enteropathogenic Escherichia coli injectisome. Journal of bacteriology. 2014;196(12):2227-41.
32.   Blaylock B, Riordan KE, Missiakas DM, Schneewind O. Characterization of the Yersinia enterocolitica
type III secretion ATPase YscN and its regulator, YscL. Journal of bacteriology. 2006;188(10):3525-34.
33.   Kuhlen L, Abrusci P, Johnson S, Gault J, Deme J, Caesar J, et al. Structure of the core of the type III
secretion system export apparatus. Nature structural & molecular biology. 2018;25(7):583-90.
34.   Johnson S, Kuhlen L, Deme JC, Abrusci P, Lea SMJm. The Structure of an Injectisome Export Gate
Demonstrates Conservation of Architecture in the Core Export Gate between Flagellar and Virulence Type III
Secretion Systems. 2019;10(3):e00818-19.
35.   Radics J, Königsmaier L, Marlovits TC. Structure of a pathogenic type 3 secretion system in action.
Nature Structural and Molecular Biology. 2014;21(1):82.
36.   Marlovits TC, Kubori T, Sukhan A, Thomas DR, Galan JE, Unger VM. Structural insights into the
assembly of the type III secretion needle complex. Science. 2004;306(5698):1040-2.
37.   Goosney DL, Gruenheid S, Finlay BBJAroc, biology d. Gut feelings: enteropathogenic E. coli (EPEC)
interactions with the host. 2000;16(1):173-89.
38.   Macdonald PJ, Chen Y, Mueller JD. Chromophore maturation and fluorescence fluctuation
spectroscopy of fluorescent proteins in a cell-free expression system. Analytical biochemistry.
2012;421(1):291-8.
39.   Barrett CM, Ray N, Thomas JD, Robinson C, Bolhuis A. Quantitative export of a reporter protein, GFP,
by the twin-arginine translocation pathway in Escherichia coli. Biochemical and biophysical research
communications. 2003;304(2):279-84.
40.   Robinson A, McDonald JP, Caldas VE, Patel M, Wood EA, Punter CM, et al. Regulation of mutagenic
DNA polymerase V activation in space and time. 2015;11(8):e1005482.
41.   Dietsche T, Mebrhatu MT, Brunner MJ, Abrusci P, Yan J, Franz-Wachtel M, et al. Structural and
functional characterization of the bacterial type III secretion export apparatus. PLoS pathogens.
2016;12(12):e1006071.
42.   Wagner S, Königsmaier L, Lara-Tejero M, Lefebre M, Marlovits TC, Galán JE. Organization and
coordinated assembly of the type III secretion export apparatus. Proceedings of the National Academy of
Sciences. 2010;107(41):17745-50.
43.   Diepold A, Wiesand U, Cornelis GR. The assembly of the export apparatus (YscR, S, T, U, V) of the
Yersinia type III secretion apparatus occurs independently of other structural components and involves the
formation of an YscV oligomer. Molecular microbiology. 2011;82(2):502-14.
44.   Gaytán MO, Monjarás Feria J, Soto E, Espinosa N, Benítez JM, Georgellis D, et al. Novel insights into
the mechanism of SepL‐mediated control of effector secretion in enteropathogenic Escherichia coli.
MicrobiologyOpen. 2018;7(3):e00571.
45.   Elliott SJ, O'Connell CB, Koutsouris A, Brinkley C, Donnenberg MS, Hecht G, et al. A gene from the
locus of enterocyte effacement that is required for enteropathogenic Escherichia coli to increase tight-junction
permeability encodes a chaperone for EspF. Infection and immunity. 2002;70(5):2271-7.
46.   Carlson ML, Young JW, Zhao Z, Fabre L, Jun D, Li J, et al. The Peptidisc, a simple method for
stabilizing membrane proteins in detergent-free solution. eLife. 2018;7:e34085.
47.   Inoue Y, Ogawa Y, Kinoshita M, Terahara N, Shimada M, Kodera N, et al. Structural Insights into the
Substrate Specificity Switch Mechanism of the Type III Protein Export Apparatus. Structure. 2019;27(6):965-
76 e6.
48.   Hu W, Walters BT, Kan ZY, Mayne L, Rosen LE, Marqusee S, et al. Stepwise protein folding at near
amino acid resolution by hydrogen exchange and mass spectrometry. Proceedings of the National Academy
of Sciences of the United States of America. 2013;110(19):7684-9.
49.   Brown KA, Wilson DJ. Bottom-up hydrogen deuterium exchange mass spectrometry: data analysis
and interpretation. Analyst. 2017;142(16):2874-86.
50.   Skinner JJ, Lim WK, Bedard S, Black BE, Englander SW. Protein hydrogen exchange: testing current
models. Protein Sci. 2012;21(7):987-95.

51. Smit J.H. KS, Srinivasu B.,Y., Karamanou S., Economou A. PyHDX: Derivation and visualization of
protection factors from Hydrogen-Deuterium Exchange Mass Spectrometry at near residue resolution.
bioRxiV. 2020.
52. Krishnamurthy S, Eleftheriadis, N., Karathanou, K., Smit, J., Portaliou, A., Chatzi, K., Karamanou,S.,
Bondar, N., Gouridis, G., and Economou, A. . A nexus of intrinsic dynamics underlies protein translocase
priming. Structure. 2021;in press.
53. Tiwari SP, Fuglebakk E, Hollup SM, Skjærven L, Cragnolini T, Grindhaug SH, et al. WEBnm@ v2.0:
Web server and services for comparing protein flexibility. BMC Bioinformatics. 2014;15(1):427.
54. Portaliou AG, Tsolis KC, Loos MS, Balabanidou V, Rayo J, Tsirigotaki A, et al. Hierarchical protein
targeting and secretion is controlled by an affinity switch in the type III secretion system of enteropathogenic
Escherichia coli. EMBO J. 2017;36(23):3517-31.
55. Terashima H, Kawamoto A, Tatsumi C, Namba K, Minamino T, Imada K. In vitro reconstitution of
functional type III protein export and insights into flagellar assembly. MBio. 2018;9(3):e00988-18.
56. Yerushalmi G, Litvak Y, Gur-Arie L, Rosenshine I. Dynamics of expression and maturation of the type
III secretion system of enteropathogenic Escherichia coli. Journal of bacteriology. 2014;196(15):2798-806.
57. Kuhlen L, Johnson S, Zeitler A, Baurle S, Deme JC, Caesar JJE, et al. The substrate specificity switch
FlhB assembles onto the export gate to regulate type three secretion. Nature communications.
2020;11(1):1296.
58. Yu XJ, Grabe GJ, Liu M, Mota LJ, Holden DW. SsaV Interacts with SsaL to Control the Translocon-
to-Effector Switch in the Salmonella SPI-2 Type Three Secretion System. mBio. 2018;9(5).
59. Bhabha G, Biel JT, Fraser JS. Keep on moving: discovering and perturbing the conformational
dynamics of enzymes. Accounts of chemical research. 2015;48(2):423-30.
60. Loutchko D, Flechsig H. Allosteric communication in molecular machines via information exchange:
what can be learned from dynamical modeling. Biophysical reviews. 2020;12(2):443-52.
61. Zhang Y, Doruker P, Kaynak B, Zhang S, Krieger J, Li H, et al. Intrinsic dynamics is evolutionarily
optimized to enable allosteric behavior. Current opinion in structural biology. 2020;62:14-21.
62. Hara N, Namba K, Minamino T. Genetic characterization of conserved charged residues in the
bacterial flagellar type III export protein FlhA. PloS one. 2011;6(7):e22417.
63. Kinoshita M, Hara N, Imada K, Namba K, Minamino T. Interactions of bacterial flagellar chaperone-
substrate complexes with FlhA contribute to co-ordinating assembly of the flagellar filament. Mol Microbiol.
2013;90(6):1249-61.
64. Minamino T, Kinoshita M, Hara N, Takeuchi S, Hida A, Koya S, et al. Interaction of a bacterial flagellar
chaperone FlgN with FlhA is required for efficient export of its cognate substrates. Mol Microbiol.
2012;83(4):775-88.
65. Chatzi KE, Sardis MF, Tsirigotaki A, Koukaki M, Sostaric N, Konijnenberg A, et al. Preprotein mature
domains contain translocase targeting signals that are essential for secretion. The Journal of cell biology.
2017;216(5):1357-69.
66. Saio T, Guan X, Rossi P, Economou A, Kalodimos CG. Structural basis for protein antiaggregation
activity of the trigger factor chaperone. Science. 2014;344(6184):1250494.
67. Yuan B, Economou A, Karamanou SJFML. Optimization of Type 3 protein secretion in
enteropathogenic E. coli. 2018.
68. Long J, Shelhamer E, Darrell T, editors. Fully convolutional networks for semantic segmentation.
Proceedings of the IEEE conference on computer vision and pattern recognition; 2015.
69. Martín Abadi AA, Paul Barham, Eugene Brevdo, Zhifeng Chen, Craig Citro, Greg S. Corrado, Andy
Davis, Jeffrey Dean, Matthieu Devin, Sanjay Ghemawat, Ian Goodfellow, Andrew Harp, Geoffrey Irving,
Michael Isard, Yangqing Jia, Rafal Jozefowicz, Lukasz Kaiser, Manjunath Kudlur, Josh Levenberg, Dan Mane,
Rajat Monga, Sherry Moore, Derek Murray, Chris Olah, Mike Schuster, Jonathon Shlens, Benoit Steiner, Ilya
Sutskever, Kunal Talwar, Paul Tucker, Vincent Vanhoucke, Vijay Vasudevan, Fernanda Viegas, Oriol Vinyals,
Pete Warden, Martin Wattenberg, Martin Wicke, Yuan Yu, Xiaoqiang Zheng. TensorFlow: Large-Scale
Machine Learning on Heterogeneous Distributed Systems. arXiv:160304467. 2016.
70. Chollet F. https://github.com/fchollet/keras. Keras. 2015.
71. Smit JH, Li Y, Warszawik EM, Herrmann A, Cordes T. ColiCoords: A Python package for the analysis
of bacterial fluorescence microscopy data. PLoS One. 2019;14(6):e0217524.
72. Pauli Virtanen RG, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni
Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua
Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, CJ
Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert
Cimrman, Ian Henriksen, E.A. Quintero, Charles R Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian
Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0–Fundamental Algorithms for Scientific
Computing in Python. arXiv:190710121. 2019.
73. Stéfan van der Walt JLS, Juan Nunez-Iglesias, François Boulogne, Joshua D. Warner, Neil Yager,
Emmanuelle Gouillart, Tony Yu and the scikit-image contributors. scikit-image: Image processing in Python.
Peer Journal. 2014.

74.   Coelho LP. Mahotas: Open source software for scriptable computer vision. Journal of Open Research
Software. 2013;1(1)(e3).
75.   Wittig I, Braun H-P, Schägger HJNp. Blue native PAGE. 2006;1(1):418.
76.   Stone AB. A simplified method for preparing sucrose gradients. Biochemical Journal. 1974;137(1):117.
77.   Zivanov J, Nakane T, Forsberg BO, Kimanius D, Hagen WJ, Lindahl E, et al. New tools for automated
high-resolution cryo-EM structure determination in RELION-3. Elife. 2018;7.
78.   Rohou A, Grigorieff N. CTFFIND4: Fast and accurate defocus estimation from electron micrographs.
Journal of structural biology. 2015;192(2):216-21.
79.   Wagner T, Merino F, Stabrin M, Moriya T, Antoni C, Apelbaum A, et al. SPHIRE-crYOLO is a fast and
accurate fully automated particle picker for cryo-EM. Communications biology. 2019;2:218.
80.   Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. Acta
crystallographica Section D, Biological crystallography. 2010;66(Pt 4):486-501.
81.   Croll TI. ISOLDE: a physically realistic environment for model building into low-resolution electron-
density maps. Acta crystallographica Section D, Structural biology. 2018;74(Pt 6):519-30.
82.   Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al. UCSF ChimeraX:
Structure visualization for researchers, educators, and developers. Protein Sci. 2021;30(1):70-82.
83.   Liebschner D, Afonine PV, Baker ML, Bunkoczi G, Chen VB, Croll TI, et al. Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments in Phenix. Acta crystallographica
Section D, Structural biology. 2019;75(Pt 10):861-77.
84.   Williams CJ, Headd JJ, Moriarty NW, Prisant MG, Videau LL, Deis LN, et al. MolProbity: More and
better reference data for improved all-atom structure validation. Protein Sci. 2018;27(1):293-315.
85.   Barad BA, Echols N, Wang RY, Cheng Y, DiMaio F, Adams PD, et al. EMRinger: side chain-directed
model and map validation for 3D cryo-electron microscopy. Nature methods. 2015;12(10):943-6.
86.   Wales TE, Eggertson MJ, Engen JR. Considerations in the analysis of hydrogen exchange mass
spectrometry data. Methods in molecular biology. 2013;1007:263-88.
87.   Linderstrøm-Lang K. Deuterium exchange between peptides and water. Chem Soc Spec Publ.
1955;2:1-20.
1104
1105

1106

Figure legends

Fig. 1. Subcellular localization of SctV and characterization of SctV nonamers

A. Cartoon representation of the EPEC injectisome. SctV$_9$ is shown in red.

B and **C.** eYFP-SctV forms distinct foci in EPECΔ*sctV* (**B**) and C41(**C**) cell peripheral. Scale

bar: 5 μm; *eYFP-sctV* expression was induced with AHT: 20 ng/mL Representative images

from YFP and brightfield channel are shown; *n=3* biological replicates.

D: Pipeline for SctV membrane complexes analysis by BN-PAGE. Following French press,

lysed cells were removed and membrane fractions were harvested by ultra-centrifugation.

Non-ionic detergents DDM was used to extract membrane proteins for further analysis by

BN-PAGE and immuno-staining with α-SctV-C.

E. BN-PAGE analysis of EPEC-derived, DDM-solubilized membrane complexes containing

SctV. Membrane proteins were extracted with different concentrations of DDM, analyzed by

BN-PAGE and α-SctV-C immuno-staining (lanes 1-6). ~1 MDa species containing SctV

component was seen when extracted with 0.8% DDM (lane 6, bracket). A predominant

~700kDa species also appeared when extra DDM was added into 0.8% DDM sample (lane

7) migrating on the gel with the similar mass size as extractions with ~1.0% DDM (lanes 3-

5). Representative images are shown; *n=3*

F. As in E, DDM-solubilized membrane complexes derived from C41/ *sctV* cells. Membrane

proteins were extracted with different concentrations of DDM, analyzed by BN-PAGE and

α-SctV-C immuno-staining. Representative images are shown; *n=3*

G SctRSTU complex formation in C41 cells. BN-PAGE analysis of C41 or C41/*sctRSTU*,

Digitonin-solubilized membrane complexes and α-SctU-C immuno-staining. A

representative image is shown; *n=3*

H-I. SctRSTU-V complex formation in C41 cells. Complexes derived from C41cells carrying

compatible plasmids with *sctV* and *sctRSTU* (expressed as indicated) extracted with 0.8%

DDM **(H)** or further treated with 1% DDM **(I)** and analyzed by BN-PAGE and α-SctV-C

immuno-staining. Representative images are shown; *n=3*

1134

Fig. 2. Analysis of SctV$_9$ as a functional receptor for chaperone/exported protein

complexes

A. Schematic representation of SctV transmembrane, linker and cytoplasmic domains (20)

B. Sub-cellular localization of full length SctV and its sub-domains in C41 cells expressing

plasmid-borne SctV, SctV-N or SctV-C (Bottom panel). Following fractionation, equal

fraction volume was analyzed by 12% SDS-PAGE and immuno-staining. Tot: Total cell

lysate; Cyt: cytoplasmic fraction; Mem: Membrane fraction; Sol: Solubilized membrane

protein fraction. Representative images are shown; *n=3.*

C. BN-PAGE analysis of His-SctV-N in C41. The membrane fraction of C41 cells expressing

plasmid-borne SctV or His-SctV-N was extracted with 1.8% DDM. Samples were analyzed

by BN-PAGE and immuno-staining with an α-SctV-C or α-His antibody (lanes 1 and 2,

respectively). Representative image is shown; *n=3*

D. The oligomerization states of SctV-C and its derivative. Purified SctV-C and one of its

interface disrupting mutants were analyzed by CN-PAGE and stained with Coomasie. SctV-

C mutant ERR: E503A-R535A-R564A; Representative images are shown; *n=3*

E. The oligomerization states of full-length SctV and its derivative. As in C, proteins from

membrane fractions derived from C41/*sctV* wt or mutated were extracted with 1.8% of DDM,

analyzed by BN-PAGE, and immuno-staining with *α*-SctV-C. SctV mutant ERR: E503A-

R535A-R564A; Representative images are shown; *n=6.*

F. Secretion of EspA and Tir from EPEC Δ*sctV* cells carrying *sctV* wt or derivatives (as

indicated). Secreted protein amounts were quantified and secretion derived from EPEC

Δ*sctV/ sctV* wt was considered 100%. SctV mutant ERR: E503A-R535A-R564A; SctV Patch

γ-: DKITFLLKKL➤GSIAVLLASS; All other values are expressed as % of this. n=3.

G. Equilibrium dissociation constants (K_d) of protein ligands for IMVs derived from C41 cells

with or without SctV or with SctV mutant (as indicated). The K_ds of CesT and CesF were

also determined for IMVs that had been pre-incubated with excess of CesF or CesT,

respectively (bottom); SctV Patch γ-: DKITFLLKKL►GSIAVLLASS; mean values (± SEM)

are shown; *n=6.*

1163

Fig. 3. Full-length SctV purification, peptidiscs reconstitution, and characterization

A. Pipeline of removal of SctV higher order aggregates and on-bead reconstitution into

peptidiscs (His-SctV$_9$-PR) and purification. Crude membrane fraction of C41/*his-sctV$_9$* cells,

following French press disruption and ultra-centrifugation, was treated with 1.8% v/v Triton

X-100 and loaded onto the 15%-35% sucrose gradient to separate His-SctV$_9$ from soluble

higher order aggregates.

B. Sucrose gradient fractions containing His-SctV$_9$ without higher order aggregates were

analyzed by BN-PAGE and stained with Coomassie blue (representative image shown).

These fractions were used for further peptidisc reconstitution. Representative image is

shown; *n=3*

C-E. Validation of His-SctV$_9$-PR reconstitution. His-SctV$_9$-PR was analyzed by BN-PAGE

followed by Coomasie blue staining (**C**) and CN-PAGE followed by silver staining (**D**) and

fluorescence peptidisc detection **(E).** Representative images are shown; *n=3.*

F. GPC-MALS/QELS analysis of His-SctV$_9$-PR using Superose^{TM} 6 (GE Healthcare). UV (in

black), mass traces (in red) and hydrodynamic diameter (in blue) of are shown. *n=3.*

G. Negative staining EM analysis of His-SctV$_9$-PR. Representative "overview" (upper panel;

scale bar: 50 nm) and "side" and "bottom views" of 2D class averages from His-SctV$_9$-PR

are shown (lower panel; scale bar: 10 nm).

1182

Fig. 4 Structure and structural dynamics of SctV-C$_9$ using Cryo-EM and HDX-MS

A. Overviews of the nonameric SctV-C model. Different views of SctV-C$_9$ model were shown

with each protomer being differentially colored.

B. Ribbon diagram indicating the sub-domains of SctV-C (SD; SD1: 353-415 and 463-483;

SD2:416-463; SD3: 488-570; SD4: 570-676). The protomer is shown in side view orientated

as the orange protomer in 4A, side view.

C. Central cross-section showing the two-funnel-like transport channel of SctV-C. Two cross

sectional protomers were shown in ribbon and surface mode, and coloured as in panel B.

The entry/exit funnels, the central pore, and the hook-shape region of SctV-C are indicated

with green arrows.

D. Side groove formed by two adjacent protomers of SctV-C. The groove of one protomer

(shown in ribbon and colored as in panel B) is indicated with a green line. Asterisks indicate

the entry and exit site.

E. Local intrinsic dynamics of SctV-C$_9$ assessed by HDX-MS. SctV-C$_9$ indicated views in

surface representation, colored according to the %D uptake. Differences (D) in %D at 5 min

are visualized on the structure of SctV-C$_9$ using a color gradient that correlates with the %

of D uptake relative to the fully deuterated control (as indicated). Side groove is indicated

with a green line. Asterisks indicate the entry and exit site. Blue hues: rigid regions; Red

hues: disordered regions; grey: unidentified. Biological replicates *n=2,* Technical replicates

n=3.

F. ΔG_{ex} values (in kJ/mol) were calculated for each residue of SctV-C by PyHDX (51) from

time-course D-exchange HDX-MS experiments and visualized on the nonameric SctV-C

protein structure. Residues are colored on a linear scale from grey (33 kJ/mol, rigid) to red

(19 kJ/mol, dynamic). Highly rigid residues (ΔG_{ex} >33 kJ/mol, transparent grey).

G. As in E, SctV-C$_9$ side view in surface representation, colored according to the %D uptake

is shown, highlighting flexible regions on one protomer. Charged residues of those regions

that are important for interactors binding are indicated, colored according to the subdomains

that they belong to.

1211

Fig. 5 Local and domain dynamics in the SctV-C protomer

A. Conformational dynamics of the SctV-C protomer assessed by HDX-MS and colored in a

ribbon representation (as in Fig. 4E). Tracing lines indicate the respective subdomains,

colored as in Fig. 4B; Latin numerals indicate important dynamic motions; Secondary

elements are represented for clarification.

B. Conformational dynamics of the SctV-C R535A mutant assessed by HDX-MS and colored

in a ribbon representation (as in Fig. 4E). %D uptake is shown.

C. Differences (D) in %D uptake on mutating residue R535A (which forms monomer) at 5

min are visualized on the structure of SctV-C9 using a color gradient that correlates with

the % of D uptake relative to the fully deuterated control (as indicated). Highly flexible

regions are displayed in green and rigid regions are shown in purple on mutating R535A.

D. ΔG_{ex} values were calculated for each residue of SctV-C using PyHDX software (as in Fig.

4F). ΔG_{ex} values are shown per residue of SctV-C-domain (residues 334-675) and colored

on a linear scale from grey (33 kJmol^{-1}; rigid), to orange (21 kJmol^{-1}, flexible) to red (19

kJmol^{-1}; dynamic). Normal modes are calculated using the WebNM@ web serve.

Displacements per normal mode are then summed with equal weights for the first 6 non-

trivial normal modes. Normal mode flexibility is derived from normal mode eigenvalues as

described previously.

E. Comparison of SctV-C sub-domains in the three indicated states derived from analysis of

the 2 PDB structures integrated into a movie (Supplementary methods; Movie S1). The

EPECSctV-C structure determined here is that of the "intermediate" state. Important elements

and arrows showing domain movements are indicated.

1234

Fig. 6 Chaperone and ATPase inner stalk binding sites on the SctV-C protomer

A. Chaperone and SctO binding domains in the indicated views of the SctV-C$_9$ surface

structure at the open (left) and closed (right) state. Chaperone binding sites determined from

peptide arrays (Fig. S6E, α,β and γ) and NMR analysis (9, 29) are coloured light green in

two adjacent protomers (dark grey) of the nonamer model. SctO binding sites determined

from cryoEM analysis (27) are colored cyan. For detailed residues and mutations see Fig.

S6E and Table S3. Red arrow: crevice between SD2-SD4 revealed by SD2 opening.

B. Chaperone binding sites (light green) are shown in the open (top) and closed (bottom)

state SctV-C of two adjacent protomers (coloured as in Fig. 4B) in a surface

representation. Red arrow: crevice between SD2-SD4 revealed by SD2 opening.

C. SctO binding sites (cyan) from (27) are shown on open (top) and closed (bottom) state

SctV-C of two adjacent protomers (coloured as in Fig.4B) in a surface representation.

Fig.1 Yuan et al,

Fig. 2, Yuan et al

Fig. 3, Yuan et al

Fig. 4 Yuan et al.

Fig. 5 Yuan et al.

A

Open   Closed

top

180°

bottom

90°

side

A B   A B

Binding sites
Chaperones *SctO*

B   C

Open

Closed

A B   A B

Fig. 6 Yuan et al.