A putative *erm* gene is present in Δ*ermB Clostridioides difficile* isolates showing high levels of resistance to clindamycin

Viviana Ramírez-Hernández[1], Gabriel Ramírez-Vargas[2]

[1]Hospital "Dr. Rafael ÁngelCalderón Guardia", San José, Costa Rica

[2]Hospital Nacional de Niños"Dr. Carlos Sáenz Herrera", San José, Costa Rica

[2]Corresponding author: garava291@hotmail.es

Abstract:

Clostridioides difficile has become the leading cause of hospital-acquired diarrhea. Clindamycin isa member of the macrolide-lincosamide-streptogramin B family of protein synthesis inhibitors. Been the ribosomal methylation the most widespread mechanism of resistance to these antibiotics in *C difficile erm* genes usually produce high levels of resistance to these drugs. In this short report we present evidence about the presence of an unreported putative *erm* gene in *C difficile* isolates that despite of presenting a Δ*ermB* genotype maintain high levels of clindamycin resistance.

Introduction:

Clostridiodes difficile is a Gram-positive, sporulating and anaerobic bacterium that has become the leading cause of hospital-acquired diarrhea(Banawas, 2018;Guery *et al*, 2019). *C difficile* infection (CDI) is associated with the exposure of the normal intestinal microbiota to antibiotics. Theresulting disruption of this normal intestinal microflora allows the multiplication and establishment of *C difficile*in the large intestine causing the pathology (Banawas, 2018;Guery *et al*, 2019).

Antibiotic resistance is crucial in spreading CDI among hospitalized patients, particularly in the older ones(Banawas, 2018). Antibiotics used for treatment of any type of infection could potentially promote (CDI) and resistant to multiple agents may have a selective advantage for the bacteria(Spigaglia, 2016). Most of the antibiotics associated with CDI appearance, including clindamycin,

remaining to be associated with the highest risk for CDI(Spigaglia, 2016; Banawas, 2018).

Clindamycin and erythromycin are members of the macrolide-lincosamide-streptogramin B (MLSB) family of protein synthesis inhibitors. Ribosomal methylation is the most widespread mechanism of resistance to these antibiotics in *C difficile*. *erm* genes usuallyproduce high levels of resistance to these drugs this bymodifying the ribosomal 23S rRNA(Spigaglia, 2016; Banawas, 2018).

The *ermB* gene can be located in the transposon Tn5398, a mobilizable nonconjugative element of 9.6 kb found in *C difficile* genomes andcontains two copies of *ermB*(Spigaglia, 2016; Banawas, 2018). Isolates whit a truncated version of Tn5398, containing only one copy of *ermB*, are called Δ*ermB*isolates and fail in inducing the MLS_Bresistance phenotype (Hussain *et al*, 2005).

In this short report we present evidence about the presence ofan unreported putative *erm* gen in *C difficile* isolates that despite

presenting a Δ*ermB* genotype maintain high levels of clindamycinresistance.

Results:

Figure 1. Alignmentof *erm* genes. Marked in red boxesare the amino acids associated whit essential ribosome methylation activities.

The gene product analyzed showed a 96% of coverage and 26% identitywhen compared to ermB from Tn5398. In addition compared to ermB this putativeerm protein presents 4 of6 essential amino acids necessary for ribosome

methylation activityalready described(Farrow *et al*, 2002).

Table 1. Features of analyzed C difficile isolates.

Isolate	*cfrC*	Δ*ermB*	CLI MIC (μg/ml) *
LIBA-5707	Yes	Yes	>256
LIBA-5701	No	Yes	>256

*CLI MIC: clindamycin minimum inhibitory concentration from Ramírez-Vargas*et al*, 2017.

Lacking *cfrC* gene (another ribosome methylating enzyme) and having Δ*ermB* genotype, LIBA-5701is reported to be resistant to clindamycin(MIC>256μg/ml)according to CLSI guides (Ramirez-Vargas *et al*, 2017;Stojković *et al*, 2019).

Discussion:

Here we report the presence of a putative *erm* gen in isolates from Costa Rican Hospitals recovered from CDI patients in the 2000s. The isolates showed similarly pathologic capabilities than other known*C difficile*epidemic isolates like NAP1 (Quesada-Gómez *et al*, 2015;López-Ureña *et al*, 2016).

This putative *erm* genpresents the key amino acidsthat has been associated to the ribosome methylation activity essential for the high levels of resistanceto for MLS$_B$ agents such as clindamycinin other erm proteins (Farrow *et al*, 2002). Mutagenesis showed that changes in these amino acids result in lowerminimum inhibitory concentration for erythromycin, another agent belonging to the MLS$_B$group (Farrow *et al*, 2002).

In Δ*ermBC difficile* isolates the MLS$_B$resistancephenotypeis lost (Hussain *et al*, 2005). So, other resistance mechanism should explain theresistance to clindamycin LIBA-5701. As reported before (Ramirez-Vargas *et al*, 2017;Stojković *et al*, 2019) lack of the element containing *cfrC* did not result in changes in the clindamycin resistance. Using the CARD database(Alcock *et al*, 2020) no other resistance genes associated to MLS$_B$resistance genotype were found (data not shown). At this point we cannot exclude the possibility of other new resistancemechanismsnot detected here and themethylationactivity of

the putative erm gen product must be probe.

Materials and methods:

Raw Illumina SRA files for LIBA-5701 and LIBA-5700used in this work can be found under the accession numbersERR467550andERR467555 respectively at the SRA database. The genomes were assembly using EDENA(Hernandez *et al*, 2008) and annotated using the RAST platform (Aziz, 2008). The predicted erm protein (supplementary material 1) was aligned against ermB from Tn5398 in *C difficle* 630 genome (accessionnumber NZ_CP019870.1) using Genius platform (Kearse *et al*, 2012).Similarly, they were compared using pBLAST (https://blast.ncbi.nlm.nih.gov/Blas t.cgi?PAGE=Proteins) for determination of identity and coverage between the sequences.

Authors declare no conflict of interest.

References:

Alcock, B. P., Raphenya, A. R., Lau, T., Tsang, K. K., Bouchard, M., Edalatmand, A., Huynh, W., Nguyen, A. V., Cheng, A. A., Liu, S., Min, S. Y., Miroshnichenko, A., Tran, H. K., Werfalli, R. E., Nasir, J. A., Oloni, M., Speicher, D. J., Florescu, A., Singh, B., Faltyn, M., … McArthur, A. G. (2020). CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic acids research*, *48*(D1), D517–D525. https://doi.org/10.1093/nar/gkz93 5

Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., Formsma, K., Gerdes, S., Glass, E. M., Kubal, M., Meyer, F., Olsen, G. J., Olson, R., Osterman, A. L., Overbeek, R. A., McNeil, L. K., Paarmann, D., Paczian, T., Parrello, B., Pusch, G. D., … Zagnitko, O. (2008). The RAST Server: rapid annotations using subsystems technology. *BMC genomics*, *9*, 75. https://doi.org/10.1186/1471-2164-9-75

Banawas S. S. (2018). *Clostridium difficile* Infections: A Global Overview of Drug Sensitivity and Resistance Mechanisms. *BioMed research international*, *2018*, 8414257. https://doi.org/10.1155/2018/841 4257

Farrow, K. A., Lyras, D., Polekhina, G., Koutsis, K., Parker, M. W., & Rood, J. I. (2002). Identification of essential residues in the Erm(B) rRNA methyltransferase of

Clostridium perfringens. *Antimicrobial agents and chemotherapy*, *46*(5), 1253–1261. https://doi.org/10.1128/aac.46.5.1253-1261.2002

Guery, B., Galperine, T., & Barbut, F. (2019). *Clostridioides difficile*: diagnosis and treatments. *BMJ (Clinical research ed.)*, *366*, l4609. https://doi.org/10.1136/bmj.l4609

Hernandez, D., François, P., Farinelli, L., Osterås, M., & Schrenzel, J. (2008). De novo bacterial genome sequencing: millions of very short reads assembled on a desktop computer. *Genome research*, *18*(5), 802–809. https://doi.org/10.1101/gr.072033.107

Hussain, H. A., Roberts, A. P., & Mullany, P. (2005). Generation of an erythromycin-sensitive derivative of Clostridium difficile strain 630 (630Deltaerm) and demonstration that the conjugative transposon Tn916DeltaE enters the genome of this strain at multiple sites. *Journal of medical microbiology*, *54*(Pt 2), 137–141. https://doi.org/10.1099/jmm.0.45790-0

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., &

Drummond, A. (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)*, *28*(12), 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., & Drummond, A. (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)*, *28*(12), 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

López-Ureña, D., Quesada-Gómez, C., Montoya-Ramírez, M., del Mar Gamboa-Coronado, M., Somogyi, T., Rodríguez, C., & Rodríguez-Cavallini, E. (2016). Predominance and high antibiotic resistance of the emerging Clostridium difficile genotypes NAPCR1 and NAP9 in a Costa Rican hospital over a 2-year period without outbreaks. *Emerging microbes & infections*, *5*(5), e42. https://doi.org/10.1038/emi.2016.38

Quesada-Gómez, C., López-Ureña, D., Acuña-Amador, L., Villalobos-Zúñiga, M., Du, T., Freire, R., Guzmán-Verri, C., del Mar Gamboa-Coronado, M., Lawley, T. D., Moreno, E., Mulvey, M. R., de Castro Brito, G. A., Rodríguez-Cavallini, E., Rodríguez, C., & Chaves-Olarte, E. (2015). Emergence of an outbreak-associated Clostridium difficile variant with increased virulence. *Journal of clinical microbiology*, *53*(4), 1216–1226. https://doi.org/10.1128/JCM.0305 8-14

Ramírez-Vargas, G., Quesada-Gómez, C., Acuña-Amador, L., López-Ureña, D., Murillo, T., Del Mar Gamboa-Coronado, M., Chaves-Olarte, E., Thomson, N., Rodríguez-Cavallini, E., & Rodríguez, C. (2017). A Clostridium difficile Lineage Endemic to Costa Rican Hospitals Is Multidrug Resistant by Acquisition of Chromosomal Mutations and Novel Mobile Genetic Elements. *Antimicrobial agents and chemotherapy*, *61*(4), e02054-16. https://doi.org/10.1128/AAC.0205 4-16

Spigaglia P. (2016). Recent advances in the understanding of antibiotic resistance in Clostridium difficile infection. *Therapeutic advances in infectious disease*, *3*(1),

23–42. https://doi.org/10.1177/20499361 15622891

Stojković, V., Ulate, M. F., Hidalgo-Villeda, F., Aguilar, E., Monge-Cascante, C., Pizarro-Guajardo, M., Tsai, K., Tzoc, E., Camorlinga, M., Paredes-Sabja, D., Quesada-Gómez, C., Fujimori, D. G., & Rodríguez, C. (2019). *cfr*(B), *cfr*(C), and a New *cfr*-Like Gene, *cfr*(E), in Clostridium difficile Strains Recovered across Latin America. *Antimicrobial agents and chemotherapy*, *64*(1), e01074-19. https://doi.org/10.1128/AAC.0107 4-19

Suplemnetary material 1. Putative erm protein sequence in this study

MKNLGYSQNFLVNKKLVERLISKSNID
VTDYVIEIGPGKGIITDVLSQHAGEVVA
VEYDQELYNNLVRYHSHDNVTYIFGDF
LKYKLPLNRRYKIFSNIPFQITADIIRKLT
DDVNPPSDINIIIQREAAKKNCGIPLQK
YEGFRAAIIKAQYKVEITHSFKRSDFFPS
PNVDTVMLHMQLWDDRLSGDDLQN
YKDLVAFFYTNIKGETAKERLSILFSNEQ
IKRLGKANRISLSNSYTLITAEQWMNIY
YYSKIGLTDEKKKKFQDAYKKLQKMNK
KLKKQNRTSLRKSSSNKKCRTRIDTSN
GTKNR