Molecular signatures of resource competition: clonal interference
drives the emergence of ecotypes

Massimo Amicone[+, *] and Isabel Gordo[+, *]

[+]Instituto Gulbenkian de Ciência (IGC)

[*]corresponding authors: mamicone@igc.gulbenkian.pt, igordo@igc.gulbenkian.pt

Short running title: Clonal interactions drive ecotypes formation.

Keywords: Eco-evolutionary dynamics, rapid adaptation, clonal interference, resource
competition, competitive exclusion, diversification, community assembly.

Type of article: Letters.

Content (number of words/items): Abstract (141), Main text (4990), References (70),
Figure legends (6).

Author contributions

MA and IG designed the study, all authors wrote the manuscript and provided final
approval for publication.

Conflict of interest

The authors declare no conflict of interest.

Data accessibility

Should the manuscript be accepted, the data supporting the results will be archived in
an appropriate public repository (Figshare) and the data DOI will be included at the end
of the article.

Abstract

Microbial ecosystems harbor an astonishing diversity that can persist for long times. To understand how such diversity is generated and maintained, ecological and evolutionary processes need to be integrated at similar timescales, but this remains a difficult challenge. Here, we extend an ecological model of resource competition to allow for evolution via *de novo* mutation, focusing on large and rapidly adapting asexual populations. Through numerical and analytical approaches, we characterize adaptation and diversity at different levels and show how clonal interference – the interaction between simultaneously emerging lineages – shapes the eco-evolutionary dynamics. We find that large mutational inputs can foster diversification under sympatry, increasing the probability that phenotypically and genetically distinct clusters arise and stably coexist, constituting an initial form of community. Our findings have implications beyond microbial populations, providing novel insights about the interplay between ecology and evolution in clonal populations.

Introduction

Understanding the mechanisms behind the evolution of biodiversity and the formation of communities remains a difficult challenge. One must integrate ecology and evolution over similar timescales, as taken together, they can give rise to phenomena that could not be explained by either alone (Schoener, 2011). The competitive exclusion principle (Hardin, 1960) theoretically bounds the number of species by the number of limiting resources. This principle generated an apparent contradiction between theoretical expectations and observations, as ecosystems can be replete of diversity even in limiting environments, both in nature (Hutchinson, 1961; Tilman, 1982; Huston, 1994) and in more controlled laboratory conditions (Maharjan *et al.*, 2006; Gresham *et al.*, 2008; Kinnersley, Holben and Rosenzweig, 2009; Herron and Doebeli, 2013). Much theoretical work has been done to resolve such controversy, often referred to as the "paradox of the plankton" (Hutchinson, 1961). Different ecological mechanisms were proposed to maintain diversity, including trade-offs on the species' traits (Posfai, Taillefumier and Wingreen, 2017), heterogeneity in space (Abrams, 1988) and time (Litchman and Klausmeier, 2001) or gene regulation (Pacciani-Mori *et al.*, 2020). However, solutions often rely on specific conditions, where a small change in the parameters could cause a collapse of the species richness. What amount of diversity can be maintained if the species' traits are not constant but rather evolve? is a critical question that remains to be answered.

Ecological theories have been challenged by evolutionary processes (Geritz *et al.*, 1998). In eco-evolutionary frameworks, mutations generate new genetic variants whose fate depends on the state of the ecosystem and, in turn, their increase in frequency can alter the populations. A common outcome of such eco-evolutionary feedbacks is that

evolution limits diversity by reducing the effectiveness of coexistence mechanisms

(Edwards *et al.*, 2018). The diversity that could be possible by ecological principles

alone, is reduced by selection of the fittest and competitive exclusion. Although several

studies have produced novel understanding on the evolution of diversity (Dieckmann

and Doebeli, 1999; Shoresh, Hegreness and Kishony, 2008; Doebeli, 2011; Kremer and

Klausmeier, 2017), the majority of them rely on a timescale separation between

ecological and evolutionary processes, i.e. on the strong-selection-weak-mutation

assumption. The emergence of beneficial mutations is assumed to be much slower than

the ecological dynamics, thus, before a new lineage arises, the population has already

reached ecological equilibrium. While allowing for analytical tractability this

assumption comes at a cost: it neglects the overlap between multiple evolving lineages –

clonal interference – a phenomena that has been extensively observed in microbial

communities *in vitro* and *in vivo* (Desai, Fisher and Murray, 2007; Barroso-Batista *et al.*,

2014). Population genetics models incorporating clonal interference have generated

predictions for the adaptation rate, fixation probabilities and genetic diversity of a

population (Gerrish and Lenski, 1998; Park and Krug, 2007; Good *et al.*, 2012; de Sousa

et al., 2016), yet excluding ecological interactions. When the population size (N) and/or

the mutation rate (U) are not small ($NU >> 1$), multiple lineages can emerge

simultaneously, ecologically interact with each other and evolve in non-trivial ways.

Although these processes are inevitably intertwined (Lawrence *et al.*, 2012; Barroso-

Batista *et al.*, 2014, 2020; Garud *et al.*, 2019), little was done to investigate how they act

in chorus.

Here, we study how clonal interference affects the interplay between ecological and

evolutionary processes and structures communities. Following a model of competition

for resources (Posfai, Taillefumier and Wingreen, 2017), we build an eco-evolutionary

framework, where no restrictive assumption on the timescales is made, as common in

other models (Geritz *et al.*, 1998; Shoresh, Hegreness and Kishony, 2008; Good, Martis

and Hallatschek, 2018). We follow an initially isogenic population throughout time and

show how different patterns of adaptation and diversity emerge. Every generation,

multiple mutations arise simultaneously in the population and interactions between

clones, which occur via competition, affect both the sign and the strength of selection.

To our knowledge this is the first study to investigate adaptation under competition for

resources across phenotypic and genetic diversity and to characterize the probability of

diversification under different molecular evolution parameters. We describe how a

large amount of variation can be maintained via a balance between selection and

mutation, and how ecotypes - functionally distinct clusters of genotypes - can emerge,

coexist and establish a first form of community.

Model and Methods

Eco-Evolutionary model

We model the dynamics of a single clonal lineage evolving to consume a set of different

substitutable resources, constantly replenished in a well-mixed environment.

Individuals mutate at a rate U and the fate of the emerging mutations depends on their

phenotypic effects, on the resource concentration, on the other individuals present in

the environment and on drift.

The underlying dynamics are determined by the MacArthur's consumer resource model

(Mac Arthur, 1969), recently formalized by *Posfai et al.* to explain high levels of diversity

in a purely ecological context, *i.e.* in the absence of mutations (Posfai, Taillefumier and

Wingreen, 2017). Briefly, let M be the number of types present at time t with densities

(*#cells/V*) n_i, *(i=1…M)* and R the number of substitutable resources with input

concentrations r_j, *(j=1…R)*. The expected density dynamics of each type are:

$$113 \qquad \frac{dn_i}{dt} = n_i(t) \left(\sum_{j=1}^{R} \frac{\alpha_j^{(i)} r_j}{\sum_{k=1}^{M(t)} n_k(t) \cdot \alpha_j^{(k)}} - \delta \right) \qquad (1)$$

where $\alpha_j^{(i)}$ represents the consumption rate of resource j by type i and δ is the death

rate. The resource amounts are constant in this model since, as *Posfai et al.*, we assume

that metabolic reactions occur much faster than cell division (Posfai, Taillefumier and

Wingreen, 2017).

We also assume a finite amount of energy available for each cell and limit their ability of

consuming resources by an energetic constraint (E):

$$120 \qquad 0 \leq \sum_{j=1}^{R} \alpha_j^{(i)} \leq E \,, \; \forall i = 1, \ldots, M \qquad (2)$$

A trade-off has been considered in previous studies as a fixed energy budget (Posfai,

Taillefumier and Wingreen, 2017; de Oliveira, Amado and Campos, 2018; Amado and

Campos, 2019), but in our model it acts as an upper limit. Assuming equally supplied

resources ($r_j = r \; \forall j$) and unitary energy, volume and death rate ($E, V, \delta=1$), the

population size is $N=Rr$.

We model an initial isogenic population (M(t_0) =1) with given traits $\vec{\alpha}^{(1)}$ and allow for

mutations that change the heritable traits and give rise to new genotypes. Every

generation, each genotype i *(i=1…M(t))* will generate a Poisson number of mutants with

expected value

$$E[\#mut_i] = n_i(t) \cdot U$$

130   with U being the per-genome, per-generation rate of non-lethal mutations. Assuming an

131   infinite site model, a mutation on genotype i will result into a new individual with

132   unique genotype (i') whose phenotypes differ from the parental traits as:

$$\vec{\alpha}^{(i')} = \vec{\alpha}^{(i)} + \vec{\Delta}, \qquad \vec{\Delta} \in \mathbb{R}^R$$

133   For simplicity, we assume the Δ_j to be normally distributed. The mutation effects are

134   defined as follows:

$$\vec{\Delta}: \begin{cases} \Delta_j \sim N(0, \sigma), j \text{ sampled from } \{1, \dots, R\} \\ \Delta_z \sim N(0, \rho \cdot \sigma), \text{for all } z \neq j \end{cases}$$

If ρ =1, a mutation changes all the traits with equal probability. If 0< ρ <1, mutations

target one trait (randomly sampled with probability $1/R$), but also alter partially the

others. If ρ =0, a mutation only changes one trait. The parameter ρ modulates different

degrees of trait interdependence or equivalently the pleiotropic effect of mutations,

while the parameter σ modulates the magnitude of the mutation effects (see Fig. 1B).

In order to respect the boundary condition (2), we assumed that: i) mutations leading to

negative values of α_j are loss of function and thus assigned α_j =0; ii) mutations that do

not respect the energy constraint cannot exist, therefore Δ_j are drawn until the upper

limit of (2) is satisfied.

In the limit of discrete time steps, we define the selection acting on genotype i at time t,

$s_i(t)$, as the expected increase in abundance in the absence of drift, such that:

$$E[n_i(t+1)] = n_i(t)(1 + s_i(t))$$

146   and from (1):

$$s_i(t) = \sum_{j=1}^{R} \frac{\alpha_j^{(i)} r_j}{\sum_{k=1}^{M(t)} n_k(t) \cdot \alpha_j^{(k)}} - \delta \, . \tag{3}$$

Thus, the fate of each genotype depends on its ability to consume each of the resources

and on the ecosystem's ecology.

To simulate drift, we draw the final abundances via multinomial sampling with

probabilities $\frac{E[n_i(t+1)]}{n_{tot}}$ $\forall i=1...M(t)$. Every generation, the number of genotypes $M(t+1)$ is

updated together with their abundances and traits. Along the simulations, we record the

genotypic and phenotypic composition of the population, from which classical statistics

used in population genetics are calculated as discussed below.

Phenotypic and genotypic diversity

In the two-resources case, within a population, each type i is characterized by a vector

of its consumption traits $(\alpha_1^{(i)}, \alpha_2^{(i)})$ and a vector of the mutations that gave rise to it,

each with a unique identifier (e.g. the vector [1,2,7,10] represents genotype 10 whose

ancestors are, in order, genotypes 7,2 and 1, and 1 is the ancestor common to every

type). From this implementation we can reconstruct the entire phylogenetic tree of a

population at any time point and map it on the phenotypic space. To measure the

genetic diversity of a population we quantify the average pairwise genetic distance π_G in

a sample of m individuals: $\pi_G(m) = \frac{\sum_{(i,j)} d_G(i,j)}{\binom{m}{2}}$, where $m=100$ and $d_G(i,j)$ is the number of

mutations that separate genotype i from j. From π_G and the number of segregating sites

in the sample, we further compute another population genetics statistic: Tajima's D,

whose expectation under the simplest population genetics neutral model is zero

(Tajima, 1989). At the functional level, we compute the average pairwise phenotypic

distance π_P, defined as: $\pi_P(m) = \frac{\sum_{(i,j)} d_E(i,j)}{\binom{m}{2}} \cdot \frac{1}{\sigma}$, where d_E is the classical Euclidean

distance.

Neutral mutation model

To understand the dynamics in the absence of selection, we run simulations with only

drift. Genotypes acquire mutations with the same trait effects and probability as

described before, but their growth probabilities are equal and do not depend on the

phenotypes. Modelling the explicit α_j under neutrality, instead of assuming that the

mutations have no effect (i.e. $\Delta_j=0$), allows for a better comparison with the model of

selection. These neutral simulations were run for the same time as the selection case.

Even though this time is much smaller than that needed to accumulate neutral changes

in a classical Wright-Fisher neutral model (Wright, 1930; Fisher, 1958), the outcome

informs on how drift contributes to the patterns of molecular and phenotypic evolution

measured here.

Parameters of the numerical simulations

In each simulation, an initially maladapted ($\alpha_1^{(1)} = \alpha_2^{(1)} = 0.05$) monomorphic

population undergoes the eco-evolutionary process described above for 10000

generations. Its genotypic and phenotypic compositions are analyzed along time, for the

following set of parameters: $N=10^7$, U: $\{10^{-8}, 5 \cdot 10^{-8}, 10^{-7}, 5 \cdot 10^{-7}, 10^{-6}, 5 \cdot 10^{-6}, 10^{-5}, 5 \cdot 10^{-5}\}$, σ:

$\{0.0125, 0.025, 0.05\}$, ρ: $\{0, 0.5, 1\}$. The parameter combinations are studied under

selection or neutrality, with the exception of fixing σ=0.05 and ρ=0 in the latter, as by

changing these values, the outcome would be equivalent. Of these, each set was

simulated in 100 independent replicates to obtain the statistics of diversity. In order to

disentangle the role of the energetic constraint assumption, we simulated adaptation

under selection but without any boundary condition (i.e. E=+∞ in (2)), with parameters:

$N=10^7$, U: $\{10^{-8}, 10^{-7}, 10^{-6}, 10^{-5}\}$, σ: $\{0.0125, 0.025, 0.05\}$, ρ=0. When we tested additional

parameters, this is specified in the text. The algorithm was written in *R* (*version 3.6.1*)

and the results analyzed in *RStudio*.

Results

Competition-driven diminishing return and the rate of adaptation

The initially monomorphic population, which is poorly adapted, is expected to acquire

mutations that improve the ability to consume the available resources and to advance in

the phenotypic space towards better adapted states. Our aim is to identify what

influences the speed of this adaptive process.

We first obtain some analytical approximations on how the selective pressure changes

in time and with the genetic composition of the population, under the competition for

resources set by (1). Selection acting on an emerging genotype i is given by (3) and

depends on the population investment on each resource j: $e_j^{(M(t))} := \sum_{i=1}^{M(t)} n_i(t) \cdot \alpha_j^{(i)}$.

Let us first simplify the problem by considering a monomorphic ($M=1$) population

whose phenotype ($\vec{\alpha}$) mirrors the resource input proportions: $\frac{\alpha_j}{\sum_k^R \alpha_k} = \frac{r_j}{\sum_k^R r_k}, \forall j = 1, \dots, R$

which consists of the local optimal strategy for a given energetic investment ($\sum_k^R \alpha_k$). In

the absence of mutations, such population at equilibrium satisfies:

$$e_j^{(1)} = r_j \cdot \sum_{k=1}^R \alpha_k \quad \forall j = 1, \dots, R. \tag{4}$$

Now consider a mutant that emerges from this population with phenotypes

$\vec{\alpha}_{mut} = \vec{\alpha} + \vec{\Delta}$. From (3) and (4) it follows that the selection acting on such mutant is:

$$s_{mut}(\vec{\alpha}, \vec{\Delta}) = \frac{\sum_j^R \Delta_j}{\sum_j^R \alpha_j}. \tag{5}$$

While bigger steps result in stronger selection, equation (5) also implies that the same

mutations are subject to weaker selection when emerging on a better adapted

background. Thus, this system exhibits diminishing returns epistasis that emerges from

the competition dynamics. Because we sample the phenotypic changes from a normal

distribution (see *Methods*), their additive effect will also follow a normal distribution

$\sum_j^R \Delta_j \sim N(0,\sigma^*)$ where $\sigma^* = \sigma\sqrt{(R-1)\cdot\rho^2 + 1}$; therefore, stronger pleiotropic effects

lead to stronger selection. From the continuous univariate distribution theory (Johnson

et al., 1994) we can retrieve the expected beneficial mutation effect $(E[\sum_j^R \Delta_j])$ and,

from (5), compute the corresponding expected selection coefficients $(E[s^+])$ for varying

values of $\sum_j^R \alpha_j$ (see Appendix S1 in Supporting Information). Figure 2A shows how the

selection strength decreases for better adapted genetic backgrounds across different σ

and ρ conditions. It is important to note that the diminishing return epistasis in this

model is not due to the energy constraint; nonetheless, such boundary condition

truncates the distribution of mutation effects (see Fig. S1A and Appendix S1) and

further slows down the rate of adaptation of well adapted populations (inset in Fig. 2A).

Following (5), the proportion of beneficial mutations, computed as

229
$$P(s^+) = \frac{\int_0^{(1-\sum_j^R \alpha_j)} N(0,\sigma*)}{\int_{-\infty}^{(1-\sum_j^R \alpha_j)} N(0,\sigma*)}$$

drops rapidly in well adapted populations as the energetic trade-off makes the

deleterious mutations more common (Fig. S1). Next, we tested how well the analytical

approximations – obtained by assuming monomorphic populations at consecutive

equilibria – predict the regimes of extensive clonal interference, where the adapting

populations are polymorphic and out of equilibrium. To do so, after running simulations

with *NU=100* and two resources, we calculate the average population trait sum

$\hat{\alpha}(t) := \frac{\sum_i^{M(t)}\left(\alpha_1^{(i)}+\alpha_2^{(i)}\right)\cdot n_i(t)}{N}$, compute the expected beneficial selection coefficient as

$E[s^+|\hat{\alpha}] = \frac{E[\Delta_1 + \Delta_2|\hat{\alpha}]}{\hat{\alpha}}$ and compare it with the mean beneficial selection observed in the

simulations, during the first 300 generations. The strength of selection acting on

polymorphic populations follows the predicted diminishing return pattern, but is often

underestimated (Fig. 2B). In fact, it decreases with the mean population phenotype $\hat{a}$,

but it increases with the population phenotype variance (Fig. S2).

Numerical simulations further allow us to link the strength of selection with the speed

of phenotypic adaptation: larger phenotypic changes imply stronger selection, resulting

in faster adaptation (Fig. 2C). We find that, when time is scaled by $\int_{\hat{a}(t0)}^{1} E[s^+|\hat{a}]d\hat{a} \cdot \sigma^*$,

the populations' mean phenotype moves with similar velocity, demonstrating that both

the complex form of selection and the mutation type mediate the speed of adaptation

(see Fig. 2D and Appendix S1). Finally, the simulations show how larger mutation rates

further accelerate adaptation. Specifically, populations under intense clonal

interference ($NU\square1$) can approach the phenotypic optimum rapidly (Fig. 2E).

In summary, these results show how the availability and the effect of beneficial

mutations, together with the complex form of selection, dictate the rate of adaptation

that slows down over time and drives the populations from an initially strong to a

finally weak selection regime.

Number of coexisting genotypes

During adaptation of an initially monomorphic population, *de novo* mutations can

generate polymorphism but at the same time purifying selection tends to reduce such

diversity. How many genotypes are generated and maintained under competition for

resources? The simulations show that, after a first burst of diversity, the mean number

of genotypes first declines and later plateaus (Fig. 3A). This reduction in the mean

number of genotypes is due to the energetic constraint; In fact, populations evolving

under neutrality or without such boundary do not suffer any decline (Fig. S3). In

contrast, when the populations' phenotypes approach the energetic constraint,

beneficial mutations become rarer and selection reduces the number of coexisting

genotypes (Fig. S3). Despite the more abundant deleterious mutations, the populations

can maintain a dynamic balance between the lineages that are purged and the newly

emerging ones (inset of Fig. 3A). The long-term number of genotypes, *M**, averaged

across the last 1000 generations, can be fitted by a linear function of the population

mutation rate whose slope decreases for stronger selection: *M*≈aNU+1*, with an

inferred *a= {8.25±0.06, 6.83±0.02, 5.48±0.02}* for *σ= {0.0125,0.025,0.05}*, respectively

(Fig. 3B). Additional simulations, run over longer times (50000 generations), confirmed

the long-term plateau of the number of genotypes (*M*), which can be fitted to a power

law decay *M(t)=ct⁻ᵝ+d*, with asymptote *d≈1* (see Fig. S4, for parameter values *N=10⁷*,

U=10⁻⁵, σ=0.05, ρ=0).

The number of genotypes, *M**, deviates from *NU/σ* - the expected mean number of

deleterious mutations under mutation-selection balance in a simple model of constant

negative selection (Haigh, 1978) (see Fig. S5).

Population diversification into ecotypes

In this model, adapting populations consist of a cloud of many genotypes (Fig. 3) and we

now characterize the phenotypic structures of these clouds. In trait space, the

population optimum Ω (represented by the star in Fig. 1B) has $\sum_j^R \alpha_j^{(\Omega)} = 1$ and all

individuals' phenotypes mirroring the resource supply proportions $\left(\alpha_j^{(\Omega)} = \frac{r_j}{r_{tot}}, \forall j = \right.$

$\left. 1, ..., R \right)$. Such state cannot be invaded by any mutant, thus excluding diversity.

However, *Posfai et al.* have shown that, in the absence of *de novo* mutations, large

collections of phenotypes can stably coexist when they are distributed around the

optimum, if polymorphism and metabolic trade-off are the initial conditions (Posfai,

Taillefumier and Wingreen, 2017). Thus, we now ask: if mutation is the only source of

variation, will an initially maladapted isogenic population evolve towards a single

optimal state or towards distinct states?

The simulations show that, due to the stochastic nature of mutation, populations

adapting under exactly the same conditions can evolve either one or multiple states

(Fig. 4A). Remarkably, the same ancestral genotype can give rise to a better adapted

population with many genotypes, all functionally similar (Fig. 4A, left panel), or

diversify into different ecotypes: clusters of genotypes with distinct metabolic

preferences, capable of coexisting indefinitely (Fig. 4A, right panel).

We now investigate the conditions favoring such ecotype diversification, in the regime

of strong selection ($Ns \square 1$). Using the mean shift clustering algorithm (Cheng, 1995),

which tests for multimodality, one can group each adapted population into functional

clusters (see details in Appendix S2). We find that the proportion of populations that

evolved into 1, 2 or more clusters, changes dramatically with NU. Under regimes of

more intense clonal interference, distinct ecotypes more probably emerge and coexist.

Importantly, when NU is very large, the number of ecotypes may exceed the number of

limiting resources (Fig. 4B). The probability of diversification (P) – computed as the

proportion of populations that evolved more than one cluster – is close to zero when

$NU<1$ but significantly increases when $NU \geq 1$. Such increase can be fitted by a logistic

function: $P = \frac{1}{1+e^{a(\ln(NU)-b)}}$ (see Fig. 4C and Tables S1-2). In contrast, the populations do

not diversify into clusters under neutrality (Fig. 4C).

Both the rate and the type of mutations influence the diversification process. Under

intense clonal interference, larger mutation effects (σ) and/or smaller pleiotropic

effects (ρ) promote the formation of multiple ecotypes (Fig. 4C, Fig S6 and S7).

310 In summary, during the process of adaptation to improve consumption of the available

311 resources, different outcomes can evolve: i) if the input of new beneficial mutations is

312 low, the recurrently fittest takes over as a cloud of genotypes until the population forms

313 a single generalist ecotype (e.g. Video S1); ii) if NU is large, the availability of many

314 beneficial mutations provides the potential for several genotypes to coevolve and

315 distinct ecotypes to stably coexist (e.g. Video S2).

316 **Phenotypic and genotypic diversity within populations**

317 Can we predict ecotypes from the genotypes in the population? Populations composed

318 by a single functional cluster can have considerable genetic diversity if multiple lineages

319 have converged to similar phenotypes (see examples in Fig. S8). We characterized the

320 adapting populations by calculating their average pairwise genetic (π_G) and phenotypic

321 (π_P) distances. Both measures of diversity increase with NU and always exceed the

322 neutral case (Fig. 5A). The increase of π_P, but not of π_G, is negatively affected by larger

323 pleiotropic effect (ρ) (ANOVA p-value = 3.2×10^{-10} and 0.09, respectively). Correlating π_P

324 and π_G in the evolved populations, we confirmed that more intense clonal interference

325 fosters functional diversification but stronger pleiotropy constrains it (Fig. S9A-B).

326 We next attempted to predict the existence of multiple ecotypes using the genotypic

327 diversity. The populations that evolved more than one cluster have on average larger

328 π_G. However, it is difficult to establish a threshold above which multiple ecotypes can be

329 predicted (Fig. 5B). The Receiver Operating Characteristic (ROC) curves – obtained via

330 varying thresholds – show the predictive power of π_G and confirm that this improves for

331 those populations that evolved under weaker pleiotropy (Fig. 5C). For example, the

332 thresholds needed to predict ecotypes with 95% specificity are: $\pi_G> \{2.7, 3.6, 4.5\}$ for $\rho=$

333 $\{0, 0.5, 1\}$, respectively (as shown by the dotted lines in Fig. 5B and the dots in Fig. 5C).

A common statistic used to infer population genetic structure is the Tajima's D (Tajima,

1989), which compares the number of segregating mutations with the average pairwise

genetic distance. The expected value of D is zero under an equilibrium model of

mutation and drift, without selection. We note that the time simulated (10^4 generations)

is much shorter than that required to achieve a neutral equilibrium at these population

sizes ($N=10^7$), so as expected our results in the absence of selection show a negative

Tajima's D, whose mean decreases with NU due to larger numbers of newly emerging

segregating sites. Under selection, populations with the same NU may present different

genetic structures (see three examples in Fig. S8) and Tajima's D ranges from very

negative to very positive values (Fig. 6). Interestingly, the mean Tajima's D is maximal

for intermediate NU and it can be close to zero under both low or very large NU

(compare the black and the gold distributions in Fig. 6). These patterns suggest that

other statistics beyond Tajima's D are needed to understand the structure of non-

equilibrium populations undergoing strong selection.

Discussion

Microbial communities are vital for humans and many other host species (Nicholson *et*

al., 2012; Sunagawa *et al.*, 2015). Emerging observations of evolution in such

ecosystems (Barroso-Batista *et al.*, 2014; Garud *et al.*, 2019; Zhao *et al.*, 2019) motivate

new theories where the mechanisms that generate diversity involve complex forms of

selection and clonal interference (Gordo, 2019). We propose that an eco-evolutionary

model of resource competition, describing the mechanisms behind ecological

divergence, can help understand diversification within ecosystems. Our framework can

be generalized to incorporate other evolutionary mechanisms, including transmission

and horizontal gene transfer, and can serve as a bridge between ecology and population

genetics.

Population genetics models of clonal interference have greatly advanced our

understanding on adaptation under this condition (Gerrish and Lenski, 1998; Park and

Krug, 2007; Good *et al.*, 2012; de Sousa *et al.*, 2016). However, clonal interference is

rarely considered in theoretical studies of ecosystems (Farahpour *et al.*, 2018), even

though it greatly impacts the evolution of microbes within communities (Barroso-

Batista *et al.*, 2014) and may be relevant in key ecosystems such as the human

microbiota (Zhao *et al.*, 2019). Commensal species in the gut have large population sizes

$\sim 10^8$ cells/g; If each bacterium mutates in the gut as it does in the laboratory (Drake,

1991), then each gram of material will host around 10^5 new mutant cells every

generation. Even if only 0.1% brings up a benefit (Perfeito *et al.*, 2007), clonal

interference still extensively affects the gut microbiota dynamics.

Here we have studied an ecological model where clones do not compete for fixation but

for resources. Modeling competition explicitly allows to make testable predictions

about diversity as both the traits and genomes can now easily be studied. We show that

clonal interactions can drive an initial monomorphic population to polymorphism with

distinct ecotypes, deviating from the simple expectation of adapting to a single optimum

phenotype (Fig. 4).

Taking the MacArthur model, *Posfai et al.* have demonstrated that metabolic trade-offs

promote coexistence of more species than resource types, thus overcoming the

competitive exclusion principle in the absence of evolution (Posfai, Taillefumier and

Wingreen, 2017). Extensions of this framework already demonstrated its power in

recapitulating experimental results from studies of soil, plant (Goldford *et al.*, 2018) or

mammalian gut microbiotas (Leónidas Cardoso *et al.*, 2020). In addition, a similar

framework provided analytical descriptions of how populations adapt under

competition for resources and demonstrated that directional selection can limit

ecological diversification (Good, Martis and Hallatschek, 2018). This pattern is also

observed in our simulations; In fact, stronger pleiotropy, which causes stronger

directional selection, limits the emergence of clusters (Fig. 4C). Our work differs from

the latter by focusing on clonal interference and by introducing an energetic trade-off.

We show that under clonal interference, the outcome of phenotypic adaptation is

probabilistic, whereby the populations can evolve paths that would be impossible to

observe through equilibrium assumptions.

Trade-offs are commonly assumed and expected to affect evolutionary trajectories

(Farahpour *et al.*, 2018; Amado and Campos, 2019), but this is not always observed.

While many empirical results have confirmed the role of trade-offs during adaptation

(Bell and Reboud, 1997; Bull, Badgett and Wichman, 2000; Turner and Elena, 2000;

Dykhuizen and Dean, 2004; Greene *et al.*, 2005; Duffy, Turner and Burch, 2006; Coffey *et*

al., 2008; Ward, Perron and MacLean, 2009; Bailey and Kassen, 2012; Li, Petrov and

Sherlock, 2019), others did not find evidence for any (Reboud and Bell, 1997; Kassen

and Bell, 1998; Turner and Elena, 2000; Trindade *et al.*, 2009; Bedhomme, Lafforgue

and Elena, 2012). Here we assumed a trade-off in the form of an energetic constraint,

which only affects well adapted genotypes. Thus, in our model the observation of a

trade-off depends on the time at which it is measured. It would be interesting to test for

trade-offs at different times during adaptation, as this could explain some of the

contrasting findings outlined above. Compatible with this hypothesis, a trade-off in

Escherichia coli ability to grow in the presence of both glucose and lactose was found,

but it only emerged after a period of constraint-free adaptation (Satterwhite and

Cooper, 2015). We find that the metabolic trade-off in the resource consumption is not

required for the formation of distinct ecotypes but it promotes their stable coexistence

(Fig. 4, S8 and Video S2).

We introduced mutation in a model of competition for resources to couple ecology and

evolution. This framework allowed some analytical approximations of the selection that

underlies adaptation. Consistent with empirical data, competition-driven selection is

characterized by diminishing return epistasis - the benefit decline in populations with

higher mean fitness (Chou *et al.*, 2011; Kryazhimskiy *et al.*, 2014; Schoustra *et al.*, 2016;

Wünsche *et al.*, 2017).

In many ecosystems coexisting types seem to outnumber the limiting resources, and,

solving this contention has motivated numerous studies. Previous eco-evolutionary

analysis suggest that adding evolutionary changes confirms (Edwards *et al.*, 2018) or

even exacerbates (Shoresh, Hegreness and Kishony, 2008) this paradox. Perhaps

surprisingly, our simulations show that large mutational inputs maintain a dynamically

stable number of types that overcome the competitive exclusion (Fig. 3). And at the

functional level, we show that diversity generally respects the exclusion principle

(number of types ≤ number of resources) but with exceptions: in a regime of strong

clonal interference, the number of extant ecotypes can even exceed the number of

limiting resources (Fig. 4B). The process leading to such ecological diversification is

strongly influenced by the underlying molecular parameters: regimes of low clonal

interference (NU <1) lead to the evolution of a single generalist population, but large

mutational inputs ($NU{\geq}1$) can lead to the formation of two or more differentially

specialized ecotypes (Fig. 4 and Video S1-S2).

Our analysis demonstrates that both large mutation effects and weak pleiotropy foster

ecological diversification (Fig. 4C). Different pleiotropic effects are meant to represent

different interactions between the traits under selection. If the available resources are

similar (e.g. chemical composition) and/or the metabolic processes involved in their

consumption share many genes, this could increase the chances that a mutation affects

the two traits simultaneously leading to large pleiotropy. Contrarily, less related

resources could involve more independent effects leading to smaller pleiotropy,

promoting diversification (Fig. 4C). This interpretation could explain why adaptive

diversification occurred in some experimental evolution setups (Friesen *et al.*, 2004;

Sandberg *et al.* 2017) but was not observed in others (Satterwhite and Cooper, 2015;

Sandberg *et al.* 2017). In agreement with this hypothesis, Sandberg and colleagues

showed that evolving on less metabolically related resources promoted ecological

diversification (Sandberg *et al.* 2017).

Sympatric diversification can be observed experimentally and predicted by theoretical

models (Friesen *et al.*, 2004). The framework of adaptive dynamics has been extensively

used in this context as it describes evolution on fitness landscapes that change

dynamically due to frequency-dependent interactions (Geritz *et al.*, 1998; Doebeli,

2011). While very useful to understand diversification, adaptive dynamics are based on

equilibrium assumptions: the populations first evolve to an equilibrium state before

diversification occurs, as explained by the concept of evolutionary branching points.

Our approach in contrast follows adaptation through an individual-based model that

undergoes strong non-equilibrium dynamics. Previous studies (Rosindell, Harmon and

Etienne, 2015; Ispolatov, Madhok and Doebeli, 2016) highlighted how considering

evolution at the individual level is necessary to fully understand the adaptation process.

We find that the genetic diversity of a population can be used to predict the underlying

phenotypic structure, but with limitations. The accuracy of such prediction decreases if

mutations are not constrained and can affect equally all the traits under selection (Fig.

5). Other statistics based on genetic data (such as the Tajima's D) can strongly deviate

from the expected values if the populations are not yet at (evolutionary) equilibrium

and can be hard to interpret if the underlying populations' parameters are unknown.

The model studied here can be considered a first step of integration. It predicts that the

typical high mutational input of bacterial species and cancer cells, coupled with an

energetic constraint, is a mechanism capable of generating functionally diverse clonal

communities. Future frameworks addressing microbial ecology and evolution will need

to address how space, migration and/or fluctuating conditions affect the patterns of

diversity observed here. In addition, cooperation between genotypes (such as cross-

feeding or use of costly public goods) have the potential to shape diversity levels and

should be investigated in future studies.

Acknowledgements

The authors acknowledge C Bank and the members of the Evolutionary Dynamics and

Evolutionary Biology labs of the IGC for their assistance throughout the development of

this work; PR Campos, T Paixão, S Miller, G Sgarlata and R Ramiro for their comments

on the manuscript.

This work was supported by Portuguese Science Foundation (FCT) Grant

PTDC/BIA-EVL/31528/2017; Deutsche Forschungs-gemeinschaft (DFG) Grant

SFB 1310; and a cooperation agreement between University of

Cologne and Gulbenkian Institute. MA was supported by FCT Grant

PD/BD/138735/2018.

References

Abrams, P. A. (1988) 'How should resources be counted?', *Theoretical Population Biology*. doi:
10.1016/0040-5809(88)90014-7.

Amado, A. and Campos, P. R. A. (2019) 'Ecological specialization under multidimensional trade-offs',
Evolutionary Ecology. Springer International Publishing, 33(6), pp. 769–789. doi: 10.1007/s10682-019-
10013-4.

Mac Arthur, R. (1969) 'Species Packing, and What Competition Minimizes', *Proceedings of the National*
Academy of Sciences, 64(4), pp. 1369–1371. doi: 10.1073/pnas.64.4.1369.

Bailey, S. F. and Kassen, R. (2012) 'Spatial structure of ecological opportunity drives adaptation in a
bacterium', *American Naturalist*. doi: 10.1086/666609.

Barroso-Batista, J. *et al.* (2014) 'The First Steps of Adaptation of Escherichia coli to the Gut Are Dominated
by Soft Sweeps', *PLoS Genetics*. doi: 10.1371/journal.pgen.1004182.

Barroso-Batista, J. *et al.* (2020) 'Specific Eco-evolutionary Contexts in the Mouse Gut Reveal Escherichia
coli Metabolic Versatility', *Current Biology*. doi: 10.1016/j.cub.2020.01.050.

Bedhomme, S., Lafforgue, G. and Elena, S. F. (2012) 'Multihost experimental evolution of a plant RNA virus
reveals local adaptation and host-specific mutations', *Molecular Biology and Evolution*. doi:
10.1093/molbev/msr314.

Bell, G. and Reboud, X. (1997) 'Experimental evolution in Chlamydomonas II. Genetic variation in strongly
contrasted environments', *Heredity*. doi: 10.1038/hdy.1997.78.

Bull, J. J., Badgett, M. R. and Wichman, H. A. (2000) 'Big-benefit mutations in a bacteriophage inhibited
with heat', *Molecular Biology and Evolution*. doi: 10.1093/oxfordjournals.molbev.a026375.

Cheng, Y. (1995) 'Mean Shift, Mode Seeking, and Clustering', *IEEE Transactions on Pattern Analysis and*
Machine Intelligence. doi: 10.1109/34.400568.

Chou, H. H. *et al.* (2011) 'Diminishing returns epistasis among beneficial mutations decelerates
adaptation', *Science*, 332(6034), pp. 1190–1192. doi: 10.1126/science.1203799.

Coffey, L. L. *et al.* (2008) 'Arbovirus evolution in vivo is constrained by host alternation', *Proceedings of*
the National Academy of Sciences of the United States of America. doi: 10.1073/pnas.0712130105.

Desai, M. M., Fisher, D. S. and Murray, A. W. (2007) 'The Speed of Evolution and Maintenance of Variation
in Asexual Populations', *Current Biology*. doi: 10.1016/j.cub.2007.01.072.

Dieckmann, U. and Doebeli, M. (1999) 'On the origin of species by sympatric speciation', *Science*, 400(July),
pp. 0–3. doi: 10.1038/22521.

Doebeli, M. (2011) *Adaptive diversification (MPB-48)*, *Adaptive Diversification (MPB-48)*. doi:
10.23943/princeton/9780691128931.001.0001.

Drake, J. W. (1991) 'A constant rate of spontaneous mutation in DNA-based microbes', *Proceedings of the*
National Academy of Sciences of the United States of America. doi: 10.1073/pnas.88.16.7160.

Duffy, S., Turner, P. E. and Burch, C. L. (2006) 'Pleiotropic costs of niche expansion in the RNA
bacteriophage Φ6', *Genetics*. doi: 10.1534/genetics.105.051136.

Dykhuizen, D. E. and Dean, A. M. (2004) 'Evolution of specialists in an experimental microcosm', *Genetics*. doi: 10.1534/genetics.103.025205.

Edwards, K. F. *et al.* (2018) 'Evolutionarily stable communities: a framework for understanding the role of trait evolution in the maintenance of diversity', *Ecology Letters*, 21(12), pp. 1853–1868. doi: 10.1111/ele.13142.

Farahpour, F. *et al.* (2018) 'Trade-off shapes diversity in ecoevolutionary dynamics', *eLife*, 7. doi: 10.7554/eLife.36273.

Fisher, R. A. (1958) 'The genetical theory of natural selection, 2nd edn.', *Dover Publication, New York*. doi: 10.1111/jeb.12566.

Friesen, M. L. *et al.* (2004) 'Experimental evidence for sympatric ecological diversification due to frequency-dependent competition in Escherichia coli', *Evolution*, 58(2), pp. 245–260. doi: 10.1111/ele.12430.

Garud, N. R. *et al.* (2019) 'Evolutionary dynamics of bacteria in the gut microbiome within and across hosts', *PLoS biology*, 17(1), p. e3000102. doi: 10.1371/journal.pbio.3000102.

Geritz, S. A. H. *et al.* (1998) 'Evolutionary singular strategies and the adaptive growth and branching of the evolutionary tree', *Evolutionary Ecology*, 12, pp. 35–57.

Goldford, J. E. *et al.* (2018) 'Emergent simplicity in microbial community assembly', *Science*. doi: 10.1126/science.aat1168.

Good, B. H. *et al.* (2012) 'Distribution of fixed beneficial mutations and the rate of adaptation in asexual populations', *Proceedings of the National Academy of Sciences of the United States of America*, 109(13), pp. 4950–4955. doi: 10.1073/pnas.1119910109.

Good, B. H., Martis, S. and Hallatschek, O. (2018) 'Adaptation limits ecological diversification and promotes ecological tinkering during the competition for substitutable resources', *Proceedings of the National Academy of Sciences of the United States of America*, 115(44), pp. E10407–E10416. doi: 10.1073/pnas.1807530115.

Gordo, I. (2019) 'Evolutionary change in the human gut microbiome: From a static to a dynamic view', *PLoS Biology*, 17(2), pp. 1–5. doi: 10.1371/journal.pbio.3000126.

Greene, I. P. *et al.* (2005) 'Effect of Alternating Passage on Adaptation of Sindbis Virus to Vertebrate and Invertebrate Cells', *Journal of Virology*. doi: 10.1128/jvi.79.22.14253-14260.2005.

Gresham, D. *et al.* (2008) 'The repertoire and dynamics of evolutionary adaptations to controlled nutrient-limited environments in yeast', *PLoS Genetics*. doi: 10.1371/journal.pgen.1000303.

Haigh, J. (1978) 'The accumulation of deleterious genes in a population-Muller's Ratchet', *Theoretical Population Biology*, 14(2), pp. 251–267. doi: 10.1016/0040-5809(78)90027-8.

Hardin, G. (1960) 'The Competitive Exclusion Principle', *Science*, 131(November).

Herron, M. D. and Doebeli, M. (2013) 'Parallel Evolutionary Dynamics of Adaptive Diversification in Escherichia coli', *PLoS Biology*. doi: 10.1371/journal.pbio.1001490.

Huston, M. A. (1994) 'Biological diversity: the coexistence of species on changing landscapes', *Biological diversity: the coexistence of species on changing landscapes*. doi: 10.2307/2404662.

Hutchinson, G. E. (1961) 'THE PARADOX OF THE PLANKTON * New Haven , Connecticut Osborn Zoological Laboratory , The problem that I wish to discuss in the present contribution is raised by the very paradoxical situation of the plankton , particularly the phyto- plankton , of relativ', XCV(882), pp. 137–145.

Ispolatov, I., Madhok, V. and Doebeli, M. (2016) 'Individual-based models for adaptive diversification in high-dimensional phenotype spaces', *Journal of Theoretical Biology*. Elsevier, 390, pp. 97–105. doi: 10.1016/j.jtbi.2015.10.009.

Johnson, Norman L *et al.* (1994) 'Continuous Univariate Distributions', *Wiley*, 1, p. Section 10.1.

Kassen, R. and Bell, G. (1998) 'Experimental evolution in Chlamydomonas. IV. Selection in environments
that vary through time at different scales', *Heredity*. doi: 10.1046/j.1365-2540.1998.00329.x.

Kinnersley, M. A., Holben, W. E. and Rosenzweig, F. (2009) 'E unibus plurum: Genomic analysis of an
experimentally evolved polymorphism in Escherichia coli', *PLoS Genetics*. doi:
10.1371/journal.pgen.1000713.

Kremer, C. T. and Klausmeier, C. A. (2017) 'Species packing in eco-evolutionary models of seasonally
fluctuating environments', *Ecology Letters*. doi: 10.1111/ele.12813.

Kryazhimskiy, S. *et al.* (2014) 'Global epistasis makes adaptation predictable despite sequence-level
stochasticity', *Science*, 344(6191), pp. 1519–1522. doi: 10.1126/science.1250939.

Lawrence, D. *et al.* (2012) 'Species interactions alter evolutionary responses to a novel environment',
PLoS Biology. doi: 10.1371/journal.pbio.1001330.

Leónidas Cardoso, L, Durão, P, Amicone, M and Gordo, I (2020) 'Dysbiosis individualizes the fitness effect
of antibiotic resistance in the mammalian gut', *Nature Ecology and Evolution*. Springer US, 4(9), pp. 1268–
1278. doi: 10.1038/s41559-020-1235-1.

Li, Y., Petrov, D. A. and Sherlock, G. (2019) 'Single nucleotide mapping of trait space reveals Pareto fronts
that constrain adaptation', *Nature Ecology and Evolution*. Springer US, 3(11), pp. 1539–1551. doi:
10.1038/s41559-019-0993-0.

Litchman, E. and Klausmeier, C. A. (2001) 'Competition of phytoplankton under fluctuating light',
American Naturalist. doi: 10.1086/318628.

Maharjan, R. *et al.* (2006) 'Clonal adaptive radiation in a constant environment', *Science*. doi:
10.1126/science.1129865.

Nicholson, J. K. *et al.* (2012) 'Host-gut microbiota metabolic interactions', *Science*. doi:
10.1126/science.1223813.

de Oliveira, V. M., Amado, A. and Campos, P. R. A. (2018) 'The interplay of tradeoffs within the framework
of a resource-based modelling', *Ecological Modelling*. Elsevier, 384(June), pp. 249–260. doi:
10.1016/j.ecolmodel.2018.06.026.

Pacciani-Mori, L. *et al.* (2020) 'Dynamic metabolic adaptation can promote species coexistence in
competitive microbial communities', *PLoS Computational Biology*, 16(5), pp. 1–18. doi:
10.1371/journal.pcbi.1007896.

Park, S. C. and Krug, J. (2007) 'Clonal interference in large populations', *Proceedings of the National
Academy of Sciences of the United States of America*, 104(46), pp. 18135–18140. doi:
10.1073/pnas.0705778104.

Perfeito, L. *et al.* (2007) 'Adaptive mutations in bacteria: High rate and small effects', *Science*, 317(5839),
pp. 813–815. doi: 10.1126/science.1142284.

Philip J. Gerrish & Richard E. Lenski (1998) 'The fate of competing beneficial mutations in an asexual
population', *Genetica*, 102/103, pp. 127–144,.

Posfai, A., Taillefumier, T. and Wingreen, N. S. (2017) 'Metabolic Trade-Offs Promote Diversity in a Model
Ecosystem', *Physical Review Letters*, 118(2), pp. 1–5. doi: 10.1103/PhysRevLett.118.028103.

Reboud, X. and Bell, G. (1997) 'Experimental evolution in Chlamydomonas III. Evolution of specialist and
generalist types in environments that vary in space and time', *Heredity*. doi: 10.1038/hdy.1997.79.

Rosindell, J., Harmon, L. J. and Etienne, R. S. (2015) 'Unifying ecology and macroevolution with individual-
based theory', *Ecology Letters*, 18(5), pp. 472–482. doi: 10.1111/ele.12430.

Satterwhite, R. S. and Cooper, T. F. (2015) 'Constraints on adaptation of Escherichia coli to mixed-

resource environments increase over time', *Evolution*, 69(8), pp. 2067–2078. doi: 10.1111/evo.12710.

Schoener, T. W. (2011) 'The Newest Synthesis⬚: Understanding Ecological Dynamics', *Science*,
331(January), pp. 426–429. doi: 10.1126/science.1193954.

Schoustra, S. *et al.* (2016) 'Diminishing-returns epistasis among random beneficial mutations in a
multicellular fungus', *Proceedings of the Royal Society B: Biological Sciences*, 283(1837). doi:
10.1098/rspb.2016.1376.

Shoresh, N., Hegreness, M. and Kishony, R. (2008) 'Evolution exacerbates the paradox of the plankton',
Proceedings of the National Academy of Sciences of the United States of America, 105(34), pp. 12365–
12369. doi: 10.1073/pnas.0803032105.

de Sousa, J. A. M. *et al.* (2016) 'Competition and fixation of cohorts of adaptive mutations under Fisher
geometrical model', *PeerJ*, 2016(2), pp. 1–17. doi: 10.7717/PEERJ.2256.

Sunagawa, S. *et al.* (2015) 'Structure and function of the global ocean microbiome', *Science*. doi:
10.1126/science.1261359.

T. E. Sandberg, C. J. Lloyd, B..O. Palsson, A. M. F. (2017) 'Laboratory Evolution to Alternating Substrate
Environments Yields Distinct Phenotypic and Genetic Adaptive Strategies', 83(13), pp. 1–15.

Tajima, F. (1989) 'Statistical method for testing the neutral mutation hypothesis by DNA polymorphism',
Genetics.

Tilman, D. (1982) 'Resource competition and community structure.', *Monographs in population biology*.
doi: 10.2307/4549.

Trindade, S. *et al.* (2009) 'Positive epistasis drives the acquisition of multidrug resistance', *PLoS Genetics*.
doi: 10.1371/journal.pgen.1000578.

Turner, P. E. and Elena, S. F. (2000) 'Cost of host radiation in an RNA virus', *Genetics*.

Ward, H., Perron, G. G. and MacLean, R. C. (2009) 'The cost of multiple drug resistance in Pseudomonas
aeruginosa', *Journal of Evolutionary Biology*. doi: 10.1111/j.1420-9101.2009.01712.x.

Wright, S. (1930) 'The genetical theory of natural selection: A review', *Journal of Heredity*. doi:
10.1093/oxfordjournals.jhered.a103361.

Wünsche, A. *et al.* (2017) 'Diminishing-returns epistasis decreases adaptability along an evolutionary
trajectory', *Nature Ecology and Evolution*, 1(4), pp. 1–6. doi: 10.1038/s41559-016-0061.

Zhao, S. *et al.* (2019) 'Adaptive Evolution within Gut Microbiomes of Healthy People', *Cell Host and
Microbe*. Elsevier Inc., 25(5), pp. 656-667.e8. doi: 10.1016/j.chom.2019.03.007.

Figure legends

Figure 1. **Ecological dynamics and individual-based evolutionary processes**. **A)** Illustration of the

eco-evolutionary dynamics. Bacterial genotypes, represented by circles of different colors, grow

according to the resources' concentration (squares and semicircles) and their own phenotypic traits,

represented by the enzyme-like structures on the cell wall. Mutation events (red arrow) can generate

new types whose fate will depend on drift and selection. **B)** Phenotypic space and mutation process. An

initially maladapted monomorphic population (green circle) can acquire de-novo mutations according to

the given assumptions, as explained in the inset. **C)** Examples of different clonal interference regimes, NU

<< 0.1 above, NU >>1 below. Each color represents a different mutant type.

Figure 2. Diminishing return epistasis and the adaptation rate. A) Analytical predictions of the

diminishing return epistasis in monomorphic populations. The inset shows the effect of the energy

constraint. **B)** Dots represent the mean of the observed positive selection during the first 300 generations

of adaptation. The dotted line represents the analytical expectation at each generation t, given the

average population trait sum $\left(\hat{\alpha}(t) := \frac{\sum_i^{M(t)} n_i(t)(\alpha_1^{(i)}+\alpha_2^{(i)})}{N} \right)$. Each color represents an independent

simulated population, which adapted with $\sigma = 0.05$, $\rho = 0.5$, $N = 10^7$ and U$= 10^{-5}$. **C)** The population

average trait sum $\hat{\alpha}(t)$ is shown as proxy of adaptation under different σ and ρ conditions. Other

parameters: $R = 2$, $N = 10^7$, $U = 10^{-8}$. Lines are the averages over 100 simulations and the shaded

areas represent the confidence interval. **D)** Same dynamics as in C, but on a different scale: Generations ·

I · σ*, where I and σ* are defined in the figure. **E)** Adaptation across different mutational inputs (color

coded) and the expected beneficial effect of mutations at the end of the adaptation process (generation

10000). The inset panel highlights the dynamics over the last 100 generations. Points in the right panel

are represented by the mean +/- one standard deviation in *log-log* scale. Other parameters for panel E:

$N = 10^7$, $\sigma = 0.05$, $\rho = 0.5$.

Figure 3. Genotypes' dynamics and the mutation-selection balance under competition for two

resources. A) Number of genotypes present in the environment over time, under neutrality (diamonds)

or under selection (lines). Lines represent the average across 100 populations and the shaded area their

confidence interval. The populations eventually plateau and dynamically maintain large number of

genotypes (M^*). Other parameters: $\sigma = 0.05$, $\rho = 0$ and $N = 10^7$. **B)** Long-lasting number of genotypes,

computed as the average over the last 1000 generations (e.g. dotted line in the right panel of A). The lines

represent the linear regressions with model: $M^* = aNU + 1$ and NU: $\{0.1, 0.5, 1, 5, 10, 50, 100, 500\}$ where

$a = \{8.25 \pm 0.06, 6.83 \pm 0.02, 5.48 \pm 0.02\}$ for $\sigma = \{0.0125, 0.025, 0.05\}$, respectively. Both axes are

represented in *log* scale with ticks every $\{1, ..., 9\} \cdot 10^x$. Other parameters: $\rho = 0$ and $N = 10^7$.

Figure 4. Ecological diversification under competition for two resources. A) Two example populations evolving under the same conditions ($N = 10^7, U = 10^{-5}, \rho = 0.5, \sigma = 0.05, \text{R} = 2$). The phenotypes and the preference distributions show one population that has evolved into a single optimal cluster (squares) and another population that gave rise to a stable diverse community composed by two clusters (circles). Lines connecting the shapes represent mutations. **B)** Counts of populations that evolved into 1,2 or more phenotypic clusters. Here, $N = 10^7, \sigma = 0.05, \text{R} = 2, \rho = 0$. **C)** Populations diversify with a probability that increases with $\ln(NU)$ and σ but decreases with ρ. The lines represent the fit of the data to the logistic function: $P = \frac{1}{1+e^{a(\ln(NU)-b)}}$, $NU: \{0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 1, 5, 10, 50, 100, 500\}$. The inferred parameters a and *b* are reported in Tables S1-2 and the full set of data is shown in Fig. S6. In the left plot: $\rho = 0$, while in the right one: $\sigma = 0.05$. The probabilities were computed as proportions out of 100 independent populations and their 95% confidence interval by normal approximation: $P \pm z\sqrt{\frac{P(1-P)}{100}}$, $z = 1.96$.

Figure 5. Diversity within and between populations and ecotypes' prediction. A) Average pairwise genotypic (π_G) and phenotypic (π_P) diversity were measured within each population, as defined in the *Methods*. 100 independent populations were simulated under the conditions specified on the x axis and by the colors. Other parameters: σ = 0.05, R=2 and $N = 10^7$. **B)** π_G distributions of the populations that evolved in more than one cluster (dark blue) or in a single cluster (light blue). Data form panel A with different NU were pulled together for a total of 800 populations per condition (ρ). The violin plots in A and B show the distribution of the data and their median. The dotted lines represent the π_G threshold that would ensure 95% specificity in ecotype prediction (i.e. 5% false positive) and correspond to the circles in panel C. **C)** π_G was used to predict whether each population is composed by one or multiple ecotypes. The ROC curves represent the sensitivity over one minus specificity of the prediction outcomes, for varying thresholds. The dotted line and the circles represent the thresholds that ensure 5% false positive rate and these are: $\pi_G > 2.7, 3.6$ or 4.5 for ρ=0, 0.5 or 1, respectively. The area under the curve (AUC) is reported on the figure. * or *** indicate p-value <0.01 or <0.0001, respectively.

Figure 6. **Tajima's D within and between populations.** The Tajima's D statistics was computed for each population at the end of the adaptation (generation 10000) from a sample of $m = 100$ genotypes. Samples under selection (empty circles) are compared with samples under neutrality (full diamonds). Each data point is an independent population. Those populations whose sample did not present polymorphism, are not represented (e.g. black diamonds are missing) because the Tajima's D is not defined as the number of segregating sites would be zero. The corresponding count of monomorphic samples is given in the lower panel. Other parameters: $N = 10^7, \rho = 0, \sigma = 0.05, R = 2$. The violin plots show the distributions of the data and their median.

A

Resources: r_1 ⬗ r_2 ◻

B

Consumption rate of r_2 (α_2)

Consumption rate of r_1 (α_1)

Pleiotropic effect

$\rho = 0$ ... $\rho = 1$

$\sim \sigma$

★ Optimal phenotype

C

Genotypes' abundance

NU << 1

N: population size
U: mutation rate

NU >> 1

Genotypes' abundance

Generations